

**ON THE POLYHEDRAL HOMOTOPY METHOD FOR SOLVING
GENERALIZED NASH EQUILIBRIUM PROBLEMS OF
POLYNOMIALS**

KISUN LEE AND XINDONG TANG

ABSTRACT. The *generalized Nash equilibrium problem* (GNEP) is a kind of games to find strategies for a group of players such that each player's objective function is optimized. Solutions for GNEPs are called *generalized Nash equilibria* (GNEs). In this paper, we propose a numerical method for finding GNEs of GNEPs of polynomials based on the polyhedral homotopy continuation and the Moment-SOS hierarchy of semidefinite relaxations. We show that our method can find all GNEs if they exist, or detect the nonexistence of GNEs, under some genericity assumptions. Some numerical experiments are made to demonstrate the efficiency of our method.

1. INTRODUCTION

Suppose there are N players and the i th player's strategy is a vector $x_i \in \mathbb{R}^{n_i}$ (the n_i -dimensional real Euclidean space). We write that

$$x_i := (x_{i,1}, \dots, x_{i,n_i}), \quad x := (x_1, \dots, x_N).$$

The total dimension of all strategies is $n := n_1 + \dots + n_N$. When the i th player's strategy is considered, we use x_{-i} to denote the subvector of all players' strategies except the i th one, i.e.,

$$x_{-i} := (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_N),$$

and write $x = (x_i, x_{-i})$ accordingly. The *generalized Nash equilibrium problem* (GNEP) is to find a tuple of strategies $u = (u_1, \dots, u_N)$ such that each u_i is a minimizer of the i th player's optimization

$$(1.1) \quad F_i(u_{-i}) \begin{cases} \min_{x_i \in \mathbb{R}^{n_i}} & f_i(x_i, u_{-i}) \\ \text{s.t.} & g_{i,j}(x_i, u_{-i}) = 0, (j \in \mathcal{E}_i), \\ & g_{i,j}(x_i, u_{-i}) \geq 0, (j \in \mathcal{I}_i). \end{cases}$$

In the above, the \mathcal{E}_i and \mathcal{I}_i are disjoint labeling sets (possibly empty), the f_i and $g_{i,j}$ are continuous functions in x and we suppose $\mathcal{E}_i \cup \mathcal{I}_i = \{1, \dots, m_i\}$ for each $i \in \{1, \dots, N\}$. A solution to the GNEP is called a *generalized Nash equilibrium* (GNE). If defining functions f_i and $g_{i,j}$ are polynomials in x for all $i \in \{1, \dots, N\}$ and $j \in \{1, \dots, m_i\}$, then we say the GNEP a *generalized Nash equilibrium problem of polynomials*. Besides that, we let X_i be the point-to-set map such that

$$X_i(x_{-i}) := \left\{ x_i \in \mathbb{R}^{n_i} \mid \begin{array}{l} g_{i,j}(x_i, x_{-i}) = 0, (j \in \mathcal{E}_i), \\ g_{i,j}(x_i, x_{-i}) \geq 0, (j \in \mathcal{I}_i) \end{array} \right\},$$

and

$$X := \{x \in \mathbb{R}^n \mid x_i \in X_i(x_{-i}) \text{ for all } i = 1, \dots, N\}.$$

A point $x \in \mathbb{R}^n$ is said to be *feasible* for this GNEP if $x \in X$. Moreover, if for every $i \in \{1, \dots, N\}$ and $j \in \{1, \dots, m_i\}$, the constraining function $g_{i,j}$ only has variables x_i , i.e., the i th player's feasible strategy set is independent with other players' strategies, then the GNEP is called a *(standard) Nash equilibrium problem* (NEP). The GNEP is said to be *convex* if for each i and all x_{-i} such that $X_i(x_{-i}) \neq \emptyset$, the $F_i(x_{-i})$ is a convex optimization, i.e., $f_i(x_i, x_{-i})$ is convex in x_i , $g_{i,j}$ ($j \in \mathcal{E}_i$) is linear in x_i , and $g_{i,j}$ ($j \in \mathcal{I}_i$) is convex in x_i .

GNEPs originated from economics in [2, 11], and have been widely used in many other areas, such as telecommunications [1], supply chain [42] and machine learning [33]. There are plenty of interesting models formulated as GNEP of polynomials, and we refer to [14, 15, 38, 39, 42] for them.

For recent studies on GNEPs, one primary task is to develop efficient methods for finding GNEs. Indeed, solving GNEPs may easily be out of reach, especially when convexity assumptions are not given. For NEPs, some methods are studied in [18, 44]. When the NEP is defined by polynomials, a method using the Moment-SOS semidefinite relaxation on the KKT system is introduced in [38]. There are some numerical algorithms to find GNEs, especially for convex GNEPs, such as the penalty method [3, 15], the augmented Lagrangian method [24], the variational and quasi-variational inequality approach [13, 16, 19], the Nikaido-Isoda function approach [48, 49], and the interior-point method on solving the KKT system [12]. Moreover, for convex GNEP of polynomials, a semidefinite relaxation method is introduced in [39], and it is extended to nonconvex rational GNEPs in [41]. The Gauss-Seidel method is studied in [40] for nonconvex GNEPs of polynomials. We refer to [14, 16] for surveys on GNEPs.

In this paper, we study the GNEPs of polynomials. The problems without convexity assumptions are mainly considered. We propose a method for finding GNEs based on the polyhedral homotopy continuation and the Moment-SOS semidefinite relaxations, and investigate its properties. Our main contributions are:

- We propose a numerical algorithm for solving GNEPs of polynomials. The polyhedral homotopy continuation is exploited for solving the complex KKT system of the GNEP, and we select GNEs from the set of complex KKT points with the help of Moment-SOS semidefinite relaxations.
- We show that when the GNEP is given by dense polynomials whose coefficients are generic, the mixed volume for the complex KKT system is identical with its algebraic degree. In this case, the polyhedral homotopy continuation can obtain all complex KKT points, and our algorithm finds all GNEs if they exist, or detect the nonexistence of them.
- Even when the number of complex KKT points obtained by the polyhedral homotopy is less than the mixed volume, or there exist infinitely many complex KKT points, our algorithm may still find one or more GNEs.
- Numerical experiments are presented to show the effectiveness of our algorithm.

The paper is organized as follows. In Section 2, we introduce some basics in optimality conditions for GNEPs, polyhedral homotopy and polynomial optimization. The algorithm for solving GNEPs of polynomials is proposed in Section 3. We show the polyhedral homotopy continuation is optimal for GNEPs of polynomials with generic coefficients in Section 4. Numerical experiments are presented in Section 5.

2. PRELIMINARIES

In this section, preliminary concepts for GNEPs, polyhedral homotopy continuation and polynomial optimization problems are reviewed. We introduce the optimality conditions for GNEPs to derive a system of polynomials whose collection of solutions contains GNEs. To give an upper bound of the number of solutions for a system of polynomials, we remind Bernstein's theorem. Finally, the polyhedral homotopy continuation and the Moment-SOS hierarchy of semidefinite relaxations are suggested as two main tools to solve GNEPs.

2.1. Optimality conditions for GNEPs. Under some suitable constraint qualifications (for example, the linear constraint qualification condition (LICQ), or Slater's Condition for convex problems; see [5]), if $u \in X$ is a GNE, then there exist Lagrange multiplier vectors $\lambda_1, \dots, \lambda_N$ such that

$$(2.1) \quad \begin{cases} \nabla_{x_i} f_i(x) - \sum_{j=1}^{m_i} \lambda_{i,j} \nabla_{x_i} g_{i,j}(x) = 0, & (i \in \{1, \dots, N\}) \\ \lambda_i \perp g_i(x), g_{i,j}(x) = 0, & (i \in \{1, \dots, N\}, j \in \mathcal{E}_i) \\ \lambda_{i,j} \geq 0, g_{i,j}(x) \geq 0, & (i \in \{1, \dots, N\}, j \in \mathcal{I}_i). \end{cases}$$

where $\nabla_{x_i} f_i(x)$ is the gradient of f_i with respect to x_i and $\lambda_i \perp g_i(x)$ implies that $\lambda_i^\top g_i(x) = 0$. The polynomial system (2.1) is called the *KKT system* of this GNEP. The solutions $(x, \lambda_1, \dots, \lambda_N)$ of the KKT system are called *KKT tuples* and the first block of coordinates x are called *KKT points*. For the GNEP of polynomials, the LICQ of $F_i(u_{-i})$ hold at every GNE [37], under some genericity conditions. Moreover, consider the system consists of all equations in (2.1), i.e.,

$$(2.2) \quad \begin{cases} \nabla_{x_i} f_i(x) - \sum_{j=1}^{m_i} \lambda_{i,j} \nabla_{x_i} g_{i,j}(x) = 0, & (i \in \{1, \dots, N\}) \\ \lambda_i \perp g_i(x), g_{i,j}(x) = 0, & (i \in \{1, \dots, N\}, j \in \mathcal{E}_i). \end{cases}$$

Then, the $(x, \lambda_1, \dots, \lambda_N)$ satisfying (2.2) is called a *complex KKT tuple*, and similarly, the first coordinate x is called a *complex KKT point*. For generic GNEPs of polynomials, there are finitely many solutions to (2.2) [37]. Under this case, the number of complex KKT points is called the *algebraic degree of the GNEP*.

2.2. Mixed volumes and Bernstein's theorem. Let $\mathbb{C}[z]$ be the set of all complex coefficient polynomials in the variable $z := (z_1, \dots, z_k)$. For a polynomial $p \in \mathbb{C}[z]$, suppose

$$p = \sum_{a \in \mathbb{N}^k} c_a z^a$$

where $z^a = z_1^{a_1} \cdots z_k^{a_k}$ for $a = (a_1, \dots, a_k)$. Then the *support* of p , denoted by A_p , is the set of exponent vectors for monomials such that

$$a \in A_p \quad \text{if and only if} \quad c_a \neq 0.$$

The convex hull Q_p of the support A_p is called the *Newton polytope* of p . For an integer vector $w \in \mathbb{Z}^k$, we define a map $h_w : \mathbb{Z}^k \rightarrow \mathbb{Z}$ such that

$$h_w(a) = \langle w, a \rangle \quad \text{for all} \quad a \in \mathbb{Z}^k.$$

Given a finite integer lattice of points $A \subset \mathbb{Z}^k$, the minimum value of h_w on A is denoted by $h_w^*(A)$. When it is clear from the context, we may omit the index w for h . Moreover, we let $A^w := \{a \in A \mid h_w(a) = h_w^*(A)\}$. Then, we define p^w be the

polynomial consists of terms of p supported on A^w , i.e., for each $p = \sum_{a \in A} c_a z^a \in \mathbb{C}[z]$, we have

$$p^w = \sum_{a \in A^w} c_a z^a.$$

For an m -tuple of polynomials $\mathcal{P} = (p_1, \dots, p_m)$, we denote $\mathcal{P}^w := (p_1^w, \dots, p_m^w)$. The m -tuple \mathcal{P}^w is called the *facial system* of \mathcal{P} with respect to w . The term ‘facial’ comes from the fact that A^w is a face of A exposed by a vector w .

Let Q_1, \dots, Q_m be polytopes in \mathbb{R}^k , and $\alpha_1, \dots, \alpha_m$ be nonnegative real scalars. The *Minkowski sum* of polytopes is

$$\alpha_1 Q_1 + \dots + \alpha_m Q_m := \{\alpha_1 v_1 + \dots + \alpha_m v_m \mid v_i \in Q_i\}.$$

The volume of the Minkowski sum $\alpha_1 Q_1 + \dots + \alpha_m Q_m$ can be understood as a homogeneous polynomial in variables of $\alpha_1, \dots, \alpha_m$. In particular, the coefficient for the term $\alpha_1 \alpha_2 \dots \alpha_m$ in the volume of $\alpha_1 Q_1 + \dots + \alpha_m Q_m$ is called the *mixed volume* of Q_1, \dots, Q_m , which is denoted by $MV(Q_1, \dots, Q_m)$.

In [4], it was proved that for a square polynomial system in $\mathbb{C}[z]$, the mixed volume of the system is an upper bound for the number of isolated roots in the complex torus $(\mathbb{C} \setminus \{\mathbf{0}\})^k$, which is called *Bernstein’s theorem*. Moreover, it states when the mixed volume bound is tight.

Theorem 2.1 (Bernstein’s theorem). [4, Theorem A and B] Let \mathcal{P} be a system consists of polynomials p_1, \dots, p_k in $\mathbb{C}[z]$. For each Newton polytope Q_{p_i} of p_i , we have

$$(2.3) \quad (\text{the number of isolated roots for } \mathcal{P} \text{ in } (\mathbb{C} \setminus \{\mathbf{0}\})^k) \leq MV(Q_{p_1}, \dots, Q_{p_k}).$$

The equality holds if and only if the facial system \mathcal{P}^w has no root in $(\mathbb{C} \setminus \{\mathbf{0}\})^k$ for any nonzero $w \in \mathbb{Z}^k$.

It is worth to note that Bernstein’s theorem concerns about roots in the torus $(\mathbb{C} \setminus \{\mathbf{0}\})^k$ because it allows to consider Laurent polynomials which are possible to have negative exponents. A system satisfies the equality in the above theorem is called *Bernstein generic*.

2.3. Polyhedral homotopy continuation. The *homotopy continuation* is an algorithmic method to find numerical approximations of roots for a system of polynomial equations. Consider a square system of polynomial equations $\mathcal{P} := \{p_1, \dots, p_k\} \subset \mathbb{C}[z_1, \dots, z_k]$ with k equations and k variables. We are interested in solving the system \mathcal{P} , i.e., computing a zero set

$$\mathbf{V}(\mathcal{P}) := \{z \in \mathbb{C}^k \mid p_1(z) = \dots = p_k(z) = 0\}.$$

The main idea for the homotopy continuation is solving \mathcal{P} by tracking a homotopy path from the known roots of a system \mathcal{Q} called a *start system* to that of the *target system* \mathcal{P} . Given the start system $\mathcal{Q} = \{q_1, \dots, q_k\} \subset \mathbb{C}[z_1, \dots, z_k]$ with the same number of variables and equations of \mathcal{P} , we construct a homotopy $\mathcal{H}(z, t)$ such that $\mathcal{H}(z, 0) = \mathcal{Q}$ and $\mathcal{H}(z, 1) = \mathcal{P}$. For tracking the homotopy from $t = 0$ to $t = 1$, numerical ODE solving techniques for *Dauidenko equations* and Newton’s iteration are applied; see [45] for more details.

Choosing a proper start system is an important task for the homotopy continuation as it determines the number of paths to track. In this paper, the *polyhedral homotopy continuation* established by Huber and Sturmfels [23] is considered. For each polynomial p_i in \mathcal{P} with its Newton polytope Q_{p_i} , the polyhedral homotopy

continuation constructs a start system \mathcal{Q} with the mixed volume $MV(Q_{p_1}, \dots, Q_{p_k})$ many solutions. We briefly introduce the framework of the polyhedral homotopy continuation. For a polynomial $p \in \mathbb{C}[z_1, \dots, z_k]$ and its support set A_p , we know that

$$p(z) = \sum_{a \in A_p} c_a z^a.$$

Let $\ell_p : A_p \rightarrow \mathbb{R}$ be a function defined on every lattice point in A_p , we define

$$(2.4) \quad \bar{p}(z, t) = \sum_{a \in A_p} c_a z^a t^{\ell_p(a)}$$

which is called a *lifted polynomial* of p by the lifting function ℓ_p . Lifting all polynomials p_1, \dots, p_k in \mathcal{P} gives a lifted system $\bar{\mathcal{P}}(z, t)$. Note that $\bar{\mathcal{P}}(z, 1) = \mathcal{P}$. A solution of $\bar{\mathcal{P}}$ can be expressed by a Puiseux series $z(t) = (z_1(t), \dots, z_k(t))$ where

$$z_i(t) = t^{\alpha_i} y_i + (\text{higher order terms})$$

for some rational number α_i and a nonzero constant y_i . As $z(t)$ is a solution for the lifted system, plugging $z(t)$ into each p_j gives

$$\bar{p}_j(z(t), t) = \sum_{a \in A_{p_j}} c_a y_i^a t^{(a, \alpha) + \ell_{p_j}(a)} + (\text{higher order terms}).$$

Dividing by $t^{(a, \alpha) + \ell_{p_j}(a)}$ and letting $t = 0$, we have a start system \mathcal{Q} . The solutions for \mathcal{Q} can be obtained from the branches of the algebraic function $z(t)$ near $t = 0$. The homotopy $\mathcal{H}(z, t)$ joining \mathcal{P} and \mathcal{Q} has $MV(Q_{p_1}, \dots, Q_{p_k})$ many paths as t varies from 0 to 1. It is motivated from Theorem 2.1 that a polynomial system supported on A_{p_1}, \dots, A_{p_k} has at most $MV(Q_{p_1}, \dots, Q_{p_k})$ many isolated solutions in the torus $(\mathbb{C} \setminus \{\mathbf{0}\})^k$. Polyhedral homotopy continuation is implemented robustly in many software HOM4PS2 [30], PHCPack [47] and HomotopyContinuation.jl [9].

Remark 2.2. Note that the polyhedral homotopy continuation algorithm may find all complex solutions even in the case that the number of solutions is smaller than the mixed volume. Also, the polyhedral homotopy continuation only finds roots in the torus $(\mathbb{C} \setminus \{\mathbf{0}\})^k$. Indeed, there may exist complex KKT tuples outside of the torus for GNEPs. For example, when there are KKT points with inactive constraints, Lagrange multipliers according to these constraints are 0; see also in Example 3.3. However, actual implementations are designed to find roots outside the torus by adding a small perturbation on the constant term. See [32] for details.

An algorithm for the polyhedral homotopy continuation for solving a polynomial system is given in the following:

Algorithm 2.3. For a system of polynomial equations $\mathcal{P} = \{p_1, \dots, p_k\}$, do the following:

- Step 1 For each $i = 1, \dots, k$, choose a function $\ell_{p_i} : A_{p_i} \rightarrow \mathbb{R}$. Then, construct the lifted polynomial $\bar{p}_i(z, t)$ as in (2.4), and define $\bar{\mathcal{P}}(z, t) := \{\bar{p}_1(z, t), \dots, \bar{p}_k(z, t)\}$.
- Step 2 Construct a start system \mathcal{Q} from the lifted system $\bar{\mathcal{P}}$ by trimming some powers of t and letting $t = 0$.
- Step 3 Starting from \mathcal{Q} , track $MV(Q_{p_1}, \dots, Q_{p_k})$ many paths from $t = 0$ to $t = 1$ with Puiseux series solutions $x(t)$ obtained near $t = 0$.

As the polyhedral homotopy continuation approximates roots of a system numerically, posteriori certifications are usually applied to verify the output obtained by numerical solvers, such as the Smale's α -theory [6, Chapter 8] and interval arithmetic [34, Chapter 8]. There are multiple known implementations for these methods. For α -theory certification, one can use `alphaCertified` [20] or `NumericalCertification` [29]. For certification using interval arithmetic, software `NumericalCertification` and `HomotopyContinuation.jl` [7] are available.

2.4. Basic concepts in polynomial optimization. We review basic concepts in polynomial optimization. For the set of real polynomials $\mathcal{H} = \{h_1, \dots, h_s\}$ in variables of $z := (z_1, \dots, z_k)$, the ideal generated by \mathcal{H} is

$$\text{Ideal}[\mathcal{H}] := h_1 \cdot \mathbb{R}[z] + \dots + h_s \cdot \mathbb{R}[z].$$

For a nonnegative integer d , the d -truncation of $\text{Ideal}[\mathcal{H}]$ is

$$\text{Ideal}[\mathcal{H}]_d := \text{Ideal}[\mathcal{H}] \cap \mathbb{R}[z]_d.$$

A polynomial $\sigma \in \mathbb{R}[z]$ is said to be a *sum of squares (SOS)* if $\sigma = \sigma_1^2 + \dots + \sigma_l^2$ for some polynomials $\sigma_i \in \mathbb{R}[z]$. The set of all SOS polynomials in z is denoted as $\Sigma[z]$. For a degree d , we denote the truncation

$$\Sigma[z]_d := \Sigma[z] \cap \mathbb{R}[z]_d.$$

For a set $\mathcal{Q} = \{q_1, \dots, q_t\}$ of polynomials in z , its quadratic module is the set

$$\text{Qmod}[\mathcal{Q}] := \Sigma[z] + q_1 \cdot \Sigma[z] + \dots + q_t \cdot \Sigma[z].$$

Similarly, we denote the truncation of $\text{Qmod}[\mathcal{Q}]$

$$\text{Qmod}[\mathcal{Q}]_{2d} := \Sigma[z]_{2d} + q_1 \cdot \Sigma[z]_{2d - \deg(q_1)} + \dots + q_t \cdot \Sigma[z]_{2d - \deg(q_t)}.$$

The tuple \mathcal{Q} determines the basic closed semi-algebraic set

$$\mathcal{S}(\mathcal{Q}) := \{z \in \mathbb{R}^k \mid q_1(z) \geq 0, \dots, q_t(z) \geq 0\}.$$

Moreover, for $\mathcal{H} = \{h_1, \dots, h_s\}$, its real zero set is

$$\mathbf{V}_{\mathbb{R}}(\mathcal{H}) := \mathbf{V}(\mathcal{H}) \cap \mathbb{R}^k = \{z \in \mathbb{R}^k \mid h_1(z) = \dots = h_s(z) = 0\}.$$

The set $\text{Ideal}[\mathcal{H}] + \text{Qmod}[\mathcal{Q}]$ is said to be *archimedean* if there exists $\rho \in \text{Ideal}[\mathcal{H}] + \text{Qmod}[\mathcal{Q}]$ such that the set $\mathcal{S}(\rho)$ is compact. If $\text{Ideal}[\mathcal{H}] + \text{Qmod}[\mathcal{Q}]$ is archimedean, then $\mathbf{V}_{\mathbb{R}}(\mathcal{H}) \cap \mathcal{S}(\mathcal{Q})$ must be compact. Conversely, if $\mathbf{V}_{\mathbb{R}}(\mathcal{H}) \cap \mathcal{S}(\mathcal{Q})$ is compact, say, $\mathbf{V}_{\mathbb{R}}(\mathcal{H}) \cap \mathcal{S}(\mathcal{Q})$ is contained in the ball $R - \|z\|^2 \geq 0$, then $\text{Ideal}[\mathcal{H}] + \text{Qmod}[\mathcal{Q} \cup \{R - \|z\|^2\}]$ is archimedean and $\mathbf{V}_{\mathbb{R}}(\mathcal{H}) \cap \mathcal{S}(\mathcal{Q}) = \mathbf{V}_{\mathbb{R}}(\mathcal{H}) \cap \mathcal{S}(\mathcal{Q} \cup \{R - \|z\|^2\})$. Clearly, if $f \in \text{Ideal}[\mathcal{H}] + \text{Qmod}[\mathcal{Q}]$, then $f \geq 0$ on $\mathbf{V}_{\mathbb{R}}(\mathcal{H}) \cap \mathcal{S}(\mathcal{Q})$. The reverse is not necessarily true. However, when $\text{Ideal}[\mathcal{H}] + \text{Qmod}[\mathcal{Q}]$ is archimedean, if the polynomial $f > 0$ on $\mathbf{V}_{\mathbb{R}}(\mathcal{H}) \cap \mathcal{S}(\mathcal{Q})$, then $f \in \text{Ideal}[\mathcal{H}] + \text{Qmod}[\mathcal{Q}]$. This conclusion is referenced as Putinar's Positivstellensatz [43]. Interestingly, if $f \geq 0$ on $\mathbf{V}_{\mathbb{R}}(\mathcal{H}) \cap \mathcal{S}(\mathcal{Q})$, we also have $f \in \text{Ideal}[\mathcal{H}] + \text{Qmod}[\mathcal{Q}]$, under some standard optimality conditions [36].

3. AN ALGORITHM FOR FINDING GNEs

In this section, we study an algorithm of finding GNEs based on the polyhedral homotopy continuation and the Moment-SOS semidefinite relaxations. First, we propose a framework for solving GNEPs. Then, we discuss the polyhedral homotopy continuation for solving the complex KKT systems for GNEPs of polynomials, and the Moment-SOS relaxations for selecting GNEs from the set of KKT points.

For the GNEP of polynomials, we consider its complex KKT system (2.2). Let $m := m_1 + \dots + m_N$ and define $\mathcal{K}_{\mathbb{C}} \subseteq \mathbb{C}^n \times \mathbb{C}^m$ as a finite set of complex KKT tuples, i.e., every point in $\mathcal{K}_{\mathbb{C}}$ solves the system (2.2). We further define

$$(3.1) \quad \mathcal{K} := \left\{ (x, \lambda) \in \mathcal{K}_{\mathbb{C}} \cap (\mathbb{R}^n \times \mathbb{R}^m) \mid \begin{array}{l} \lambda_{i,j} \geq 0, g_{i,j}(x) \geq 0 \\ \forall i \in \{1, \dots, N\}, j \in \mathcal{I}_i \end{array} \right\},$$

$$\mathcal{P} := \{x \in \mathbb{R}^n \mid \text{there is } \lambda \in \mathbb{R}^m \text{ such that } (x, \lambda) \in \mathcal{K}\}.$$

Then, the \mathcal{K} and \mathcal{P} are sets of KKT tuples and KKT points respectively. When the GNEP is convex, all points in \mathcal{P} are GNEs. Furthermore, when $\mathcal{K}_{\mathbb{C}}$ is the set of all complex KKT tuples and some constraint qualifications hold at every GNE, the \mathcal{P} is the set of all GNEs if it is nonempty, or the nonexistence of GNEs can be certified by the emptiness of \mathcal{P} . However, when there is no convexity assumed for the GNEP, the KKT conditions are usually not sufficient for $x \in \mathcal{P}$ being a GNE.

Suppose the GNEP is not convex. Let $u = (u_i, u_{-i}) \in \mathcal{P}$. For each $i \in \{1, \dots, N\}$, we consider the following optimization problem

$$(3.2) \quad \begin{cases} \delta_i := \min_{x_i \in \mathbb{R}^{n_i}} f_i(x_i, u_{-i}) - f_i(u_i, u_{-i}) \\ \text{s.t.} \quad g_{i,j}(x_i, u_{-i}) = 0 \quad (j \in \mathcal{E}_i), \\ \quad \quad g_{i,j}(x_i, u_{-i}) \geq 0 \quad (j \in \mathcal{I}_i). \end{cases}$$

Then, u is a GNE if and only if every $\delta_i \geq 0$, i.e., u_i minimizes (3.2) for each i . If $\delta_i < 0$ for some i , then u is not a GNE. In this case, assume (3.2) has a minimizer v_i . Then we know

$$f_i(v_i, u_{-i}) - f_i(u_i, u_{-i}) = \delta_i < 0.$$

So, if $x^* \in \mathcal{P}$ is a GNE and $v_i \in X_i(x_{-i}^*)$, then the following must hold at x^*

$$(3.3) \quad f_i(v_i, x_{-i}) - f_i(x_i, x_{-i}) \geq 0.$$

Therefore, we check the inequality (3.3) for all $x \in \mathcal{P}$ such that $v_i \in X_i(x_{-i})$. We remark that for NEPs, the $v_i \in X_i(x_{-i}^*)$ hold for all NEs. In summary, we propose the following algorithm for finding GNEs

Algorithm 3.1. For the GNEP of polynomials, do the following:

- Step 0 Let $S := \emptyset$ and $V_i := \emptyset$ for all $i \in \{1, \dots, N\}$.
- Step 1 Solve the complex KKT system (2.2) for a set of complex solutions $\mathcal{K}_{\mathbb{C}}$. Let \mathcal{P} be the set given as in (3.1).
- Step 2 If $\mathcal{P} \neq \emptyset$, then select $u \in \mathcal{P}$, let $\mathcal{P} := \mathcal{P} \setminus \{u\}$, and proceed to the next step. Otherwise, output the set S (possibly empty) of GNEs and stop.
- Step 3 If $f_i(v_i, u_{-i}) - f_i(u_i, u_{-i}) \geq 0$ for all $i \in \{1, \dots, N\}$ and for all $v_i \in V_i \cap X_i(u_{-i})$, then go to the next step. Otherwise, go back to Step 2.
- Step 4 For each $i \in \{1, \dots, N\}$, solve the polynomial optimization problem (3.2) for a minimizer v_i . If there exists $i \in \{1, \dots, N\}$ such that $\delta_i < 0$, let $V_i := V_i \cup \{v_i\}$ for all such i . Otherwise, u is a GNE and let $S := S \cup \{u\}$. Then, go back to Step 2.

In Section 3.1, we show how to find the set of complex solutions for the system (2.2) using the polyhedral homotopy continuation. Since the polyhedral homotopy tracks mixed volume many paths, the \mathcal{P} must be a finite set (possibly empty). Therefore, Algorithm 3.1 must terminate within finitely many loops. Moreover, if $\mathcal{K}_{\mathbb{C}}$ is the set of all complex KKT tuples, i.e., Algorithm 2.3 finds all complex solutions for (2.2), then Algorithm 3.1 will either find all GNEs, or detect the

nonexistence of GNEs. This is the case when Algorithm 2.3 finds the mixed volume many complex solutions for (2.2). In Section 4, we show that when the GNEP is defined by generic dense polynomials, Algorithm 2.3 can always find the mixed volume many solutions for (2.2). The following result is straightforward:

Theorem 3.2. For the GNEP, suppose $|\mathcal{K}_{\mathbb{C}}|$ equals the mixed volume of (2.2). If S produced by Algorithm 3.1 is nonempty, then S is the set of all GNEs. Otherwise, the GNEP does not have any GNE.

3.1. The polyhedral homotopy method for finding KKT tuples. In this subsection, we explain how the polyhedral homotopy continuation is applied to find complex solutions for the system (2.2). For each $i \in \{1, \dots, N\}$, denote the tuple of polynomials in variables of x and λ_i

$$F_i(x, \lambda_i) := \left\{ \nabla_{x_i} f_i(x) - \sum_{j=1}^{m_i} \lambda_{i,j} \nabla_{x_i} g_{i,j}(x) \right\} \\ \cup \{ \lambda_{i,j} g_{i,j}(x) \mid j \in \mathcal{I}_i \} \cup \{ g_{i,j}(x) \mid j \in \mathcal{E}_i \}.$$

We define a system

$$(3.4) \quad F(x, \lambda) := \bigcup_{i=1}^N F_i(x, \lambda_i).$$

Then, $F(x, \lambda)$ is a system with $n+m$ polynomial equations and $n+m$ variables, and we use Algorithm 2.3 to solve $F(x, \lambda) = 0$ by letting $z := (x, \lambda)$ and $\mathcal{P}(z) := F(z)$.

The example below shows detail of applying the homotopy method for finding KKT tuples from an actual NEP problem.

Example 3.3. Consider the two-players unconstrained NEP

$$\min_{x_1 \in \mathbb{R}^1} \frac{1}{2} x_1^2 x_2^3 - x_1 x_2^2 - 2x_1 x_2 \quad \Bigg| \quad \min_{x_2 \in \mathbb{R}^1} \frac{1}{2} x_1^3 x_2^2 - x_1^2 x_2 - 2x_1 x_2.$$

For such the problem, the complex KKT system reduces to vanishing the gradients $\nabla_{x_1} f_1$ and $\nabla_{x_2} f_2$, i.e., we have

$$F = \{ \nabla_{x_1} f_1, \nabla_{x_2} f_2 \} = \{ x_1 x_2^3 - x_2^2 - 2x_2, x_1^3 x_2 - x_1^2 - 2x_1 \}.$$

Considering a lifted system with generic lifting functions ℓ_{f_1} and ℓ_{f_2} , we have

$$\overline{F}(x, t) = (x_1 x_2^3 t^{\ell_{f_1}(1,3)} - x_2^2 t^{\ell_{f_1}(0,2)} - 2x_2 t^{\ell_{f_1}(0,1)}, \\ x_1^3 x_2 t^{\ell_{f_2}(3,1)} - x_1^2 t^{\ell_{f_2}(2,0)} - 2x_1 t^{\ell_{f_2}(1,0)})$$

such that $\overline{F}(x, 1) = F$ and a start system $\overline{F}(x, 0)$ after trimming some powers of t . The system F has the mixed volume equals to 8. Therefore, the polyhedral homotopy continuation provides 8 paths to track and 6 numerical solutions

$$(x_1, x_2) := \begin{cases} (1.52138, 1.52138), (-0.76069 + 0.857874i, -0.76069 + 0.857874i), \\ (-0.76069 - 0.857874i, -0.76069 - 0.857874i), (0, -2), \\ (-2, 0), (0, 0) \end{cases}$$

of F are found. Indeed, using the software `Macaulay2` [17], we find the system $F(x) = 0$ has exactly 6 complex solutions.

Remark 3.4. (1) Note that the system above has the number of solutions equals to 6, which is strictly less than its mixed volume. The homotopy continuation algorithm may find all complex solutions, even if the number of solutions is smaller than the mixed volume.

- (2) As presented in the Example 3.3, the polyhedral homotopy continuation may find solutions outside the torus.

In Section 4, we show that under genericity assumption, the polyhedral homotopy continuation provides the optimal number of paths for finding complex KKT points. In this case, the polyhedral homotopy continuation guarantees to find all complex KKT points, hence the complete collection of GNEs can be obtained by Algorithm 3.1.

3.2. The Moment-SOS relaxation for selecting GNEs. In this sequel, we discuss how to solve the polynomial optimization (3.2). For each i , denote

$$\begin{aligned}\theta_i(x_i) &:= f_i(x_i, u_{-i}) - f_i(u_i, u_{-i}), \\ \Phi_i(x_i) &:= \{g_{i,j}(x_i, u_{-i}) \mid j \in \mathcal{E}_i\}, \\ \Psi_i(x_i) &:= \{g_{i,j}(x_i, u_{-i}) \mid j \in \mathcal{I}_i\}.\end{aligned}$$

Denote the degree

$$(3.5) \quad d_i := \max\{\lceil \deg(\theta_i)/2 \rceil, \lceil \deg(\Phi_i(x_i))/2 \rceil, \lceil \deg(\Psi_i(x_i))/2 \rceil\},$$

where $\deg(\Phi_i(x_i)) := \max\{\deg(g_{i,j}(x_i, u_{-i})) \mid j \in \mathcal{E}_i\}$, and $\deg(\Psi_i(x_i))$ is similarly defined. For $d \geq d_i$, we denote by $\mathbb{N}_{2d}^{n_i}$ the set of all tuples of n_i nonnegative integers whose sum is not greater than d , and let $y \in \mathbb{R}^{\mathbb{N}_{2d}^{n_i}}$ be the variable that is labeled by tuples in $\mathbb{N}_{2d}^{n_i}$. Then the d th moment relaxation for (3.2) is

$$(3.6) \quad \begin{cases} \vartheta_i^{(d)} := \min_{y \in \mathbb{R}^{\mathbb{N}_{2d}^{n_i}}} \langle \theta_i, y \rangle \\ \text{s.t.} \quad L_p^{(d)}[y] \succeq 0 (p \in \Phi_i), \quad L_q^{(d)}[y] = 0 (q \in \Psi_i) \\ y_0 = 1, \quad M_d[y] \succeq 0, \end{cases}$$

Its dual optimization problem is the d th SOS relaxation

$$(3.7) \quad \begin{cases} \max & \gamma \\ \text{s.t.} & \theta_i - \gamma \in \text{Ideal}(\Psi_i)_{2d} + \text{Qmod}(\Phi_i)_{2d}. \end{cases}$$

By solving the relaxations (3.6)-(3.7) for $d = d_0, d_0 + 1, \dots$, we get the Moment-SOS hierarchy for solving (3.2). The following is the algorithm.

Algorithm 3.5. For the given $u \in \mathcal{P}$ and the i th player's optimization (3.2). Initialize $d := d_i$.

Step 1 Solve the moment relaxation (3.6) for the minimum value $\vartheta_i^{(d)}$ and a minimizer y^* . If $\vartheta_i^{(d)} \geq 0$, then $\delta_i = 0$ and stop; otherwise, go to the next step.

Step 2 Let $t := d_i$ as in (3.5). If y^* satisfies the rank condition

$$(3.8) \quad \text{rank } M_t[y^*] = \text{rank } M_{t-d_i}[y^*],$$

then extract a set U_i of $r := \text{rank } M_t(y^*)$ minimizers for (3.2) and stop.

Step 3 If (3.8) fails to hold and $t < d$, let $t := t + 1$ and then go to Step 2; otherwise, let $d := d + 1$ and go to Step 1.

The rank condition (3.8) is called *flat truncation* in the literature [35]. It is a sufficient (and almost necessary) condition for checking convergence of the Moment-SOS hierarchy. Indeed, the Moment-SOS hierarchy has finite convergence if and only if the flat truncation is satisfied for some relaxation order, under some generic conditions [35]. When (3.8) holds, the method in [21] can be used to extract r

minimizers for (3.2). The method is implemented in the software `GloptiPoly3` [22]. We refer to [21, 35] and [26, Chapter 6] for more details.

The convergence properties of Algorithm 3.5 are as follows. By solving the hierarchy of relaxations (3.6) and (3.7), we get a monotonically increasing sequence of lower bounds $\{\vartheta_d\}_{d=d_0}^{\infty}$ for the minimum value ϑ_{\min} , i.e.,

$$\vartheta_{d_0} \leq \vartheta_{d_0+1} \leq \cdots \leq \vartheta_{\min}.$$

When $\text{Ideal}(\Psi_i)_{2d} + \text{Qmod}(\Phi_i)_{2d}$ is archimedean, we have $\vartheta_d \rightarrow \vartheta_{\min}$ as $d \rightarrow \infty$, as shown in [25]. If $\vartheta_d = \vartheta_{\min}$ for some d , the relaxation (3.6) is said to be exact for solving (3.2). For such a case, the Moment-SOS hierarchy is said to have finite convergence. The Moment-SOS hierarchy has finite convergence when the archimedean and some optimality conditions hold [36]. Although there exist special polynomials such that the Moment-SOS hierarchy fails to have finite convergence, such special problems belong to a set of measure zero in the space of input polynomials [36]. We refer to [26–28, 36] for more work on polynomial and moment optimization.

4. THE MIXED VOLUME OF GNEPS

For a polynomial system, if the set of its complex solutions is zero-dimensional, then the algebraic degree of the polynomial system counts the number of complex solutions for the system. In this section, we prove that under some genericity assumptions on the GNEP, the mixed volume of the complex KKT system (2.2) equals to its algebraic degree. Throughout this section, we have a GNEP of polynomials consists of dense polynomials of certain degrees. Without loss of generality, we assume $\mathcal{I}_i = \emptyset$ for all $i \in \{1, \dots, N\}$, i.e., all players only have equality constraints, for convenience of our discussion. Note that if there exist inequality constraints, then all following results still hold by enumerating the active constraints. For a tuple $d := (d_1, \dots, d_N)$ of nonnegative integers, the $\mathbb{C}[x]_d$ represents the space of polynomials whose degree in x_i is not greater than d_i .

Recall that we say a system of polynomials is Bernstein generic if the number of isolated solutions in the complex torus equals its mixed volume. The main result of this section is the following:

Theorem 4.1. Consider the GNEP of polynomials given as in (1.1). For each i , let $d_{i,0}, \dots, d_{i,m_i} \in \mathbb{N}^N$ be tuples of nonnegative integers. Suppose all f_i and $g_{i,j}$ are generic dense polynomials in $\mathbb{C}[x]_{d_{i,0}}$ and $\mathbb{C}[x]_{d_{i,j}}$ respectively. Then, the complex KKT system (3.4) is Bernstein generic.

We first introduce some basic notation and useful lemmas, and then prove Theorem 4.1. We define a weight vector w for the complex KKT system (3.4). Let w_1, \dots, w_N be weight vectors such that

$$w_i = (\mathbf{0}, \dots, \mathbf{0}, (w_{i,1}, \dots, w_{i,n_i}, v_{i,1}, \dots, v_{i,m_i}), \mathbf{0}, \dots, \mathbf{0}),$$

where $w_{i,k}$ and $v_{i,j}$ are the weights for the variables $x_{i,k}$ and $\lambda_{i,j}$ respectively. Define $w := \sum_{i=1}^N w_i$. Each w_i is the weight vector for the system F applied only for x_i and λ_i variables.

The idea for the proof of the result is inspired by the paper [8]. We introduce the lemmas established from the paper that will be used for the proof of the Theorem 4.1.

Lemma 4.2. [8, Lemma 8] Let p be a polynomial in $\mathbb{C}[x]$ and $w \in \mathbb{Z}^n$ be an integer vector. If $\frac{\partial p^w}{\partial x_i} \neq 0$, then $\frac{\partial p^w}{\partial x_i} = \left(\frac{\partial p}{\partial x_i}\right)^w$ and $h^*(A_{\frac{\partial p}{\partial x_i}}) = h^*(A_p) - w_i$.

Lemma 4.3. Let p be a polynomial in $\mathbb{C}[x]_d$ and $w = \sum_{i=1}^N w_i$ be a weight vector in \mathbb{Z}^n . Then, we have

$$h_{w_i}^*(A_p) \cdot p^w = \sum_{k=1}^{n_i} w_{i,k} x_{i,k} \frac{\partial p^w}{\partial x_{i,k}}.$$

Proof. For a monomial x^a , note that $x_{i,k} \frac{\partial x^a}{\partial x_{i,k}} = a_{i,k} x^a$. Therefore, we have

$$\sum_{k=1}^{n_i} w_{i,k} x_{i,k} \frac{\partial x^a}{\partial x_{i,k}} = \sum_{k=1}^{n_i} w_{i,k} a_{i,k} x^a = \langle w_i, a \rangle x^a.$$

Summing over all monomials in p^w , we get

$$h_{w_i}^*(A_{p^w}) \cdot p^w = \sum_{k=1}^{n_i} w_{i,k} x_{i,k} \frac{\partial p^w}{\partial x_{i,k}}.$$

Noting the fact that $h_{w_i}^*(A_p) = h_{w_i}^*(A_{p^w})$, we get the desired result. \square

Note that Lemma 4.3 is a generalization of the Euler's formula for quasihomogeneous polynomials mentioned in [8, Lemma 9]. We now ready to prove Theorem 4.1.

Proof of Theorem 4.1. For each polynomial p and a weight vector w , let $\mathbf{I}_i^w(p)$ and $\mathbf{J}_i^w(p)$ be a partition set of indices $\{1, \dots, n_i\}$ for each i such that $\frac{\partial p^w}{\partial x_{i,j}} \neq 0$ if $j \in \mathbf{I}_i^w(p)$ and $\frac{\partial p^w}{\partial x_{i,j}} = 0$ if $j \in \mathbf{J}_i^w(p)$. Also, we let

$$\mathbf{I}_i^w = \mathbf{I}_i^w(f_i) \cup \left(\bigcup_{j=1}^{m_i} \mathbf{I}_i^w(g_{i,j}) \right), \quad \mathbf{I}^w = \bigcup_{i=1}^N \mathbf{I}_i^w, \quad \text{and} \quad \hat{n}_i = |\mathbf{I}_i^w|.$$

It is clear that if $j \in \mathbf{J}_i^w(p)$ and $a \in A_p^w$, then $a_{i,j} = 0$. Hence, we may consider p^w as a polynomial in $\mathbb{C}[\mathbf{I}^w] := \mathbb{C}[x_{i,j} \mid j \in \mathbf{I}_i^w(p), i = 1, \dots, N]$. Note that if p is a generic polynomial, then p^w can also be considered as a generic polynomial in $\mathbb{C}[\mathbf{I}^w]$.

In the following, for a fixed weight vector w , we show that there is no roots for the facial system F^w in the torus $(\mathbb{C} \setminus \{0\})^n$. Consider the facial system of F_i , say,

$$F_i^w = \left\{ (\nabla_{x_i} f_i - \sum_{j=1}^{m_i} \lambda_{i,j} \nabla_{x_i} g_{i,j})^w, g_{i,1}^w, \dots, g_{i,m_i}^w \right\}$$

where

$$\nabla_{x_i} f_i - \sum_{j=1}^{m_i} \lambda_{i,j} \nabla_{x_i} g_{i,j} = \left\{ \frac{\partial f_i}{\partial x_{i,1}} - \sum_{j=1}^{m_i} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,1}}, \dots, \frac{\partial f_i}{\partial x_{i,n_i}} - \sum_{j=1}^{m_i} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,n_i}} \right\}.$$

For each $k \in \{1, \dots, n_i\}$, let $A_{\partial_{i,k}}$ be the support of $\frac{\partial f_i}{\partial x_{i,k}} - \sum_{j=1}^{m_i} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,k}}$. Then, we have

$$h^*(A_{\partial_{i,k}}) = \min \left\{ h^* \left(A_{\frac{\partial f_i}{\partial x_{i,k}}} \right), \min_{j=1, \dots, m_i} \left\{ h^* \left(A_{\frac{\partial g_{i,j}}{\partial x_{i,k}}} \right) + v_{i,j} \right\} \right\}.$$

In the above, we recall that for a polynomial p , the $h^*(A_p)$ is the minimum of $h(a) := \langle w, a \rangle$ over $a \in A_p$; see Section 2.2 for more details. Depending on where the value of $h^*(A_{\partial_{i,k}})$ is attained, there are the following three cases:

$$\begin{aligned} \text{(a)} \quad & h^*(A_{\partial_{i,k}}) = \min_{j=1, \dots, m_i} \left\{ h^* \left(A_{\frac{\partial g_{i,j}}{\partial x_{i,k}}} \right) + v_{i,j} \right\}, \\ \text{(b)} \quad & h^*(A_{\partial_{i,k}}) = h^* \left(A_{\frac{\partial f_i}{\partial x_{i,k}}} \right), \\ \text{(c)} \quad & h^*(A_{\partial_{i,k}}) = h^* \left(A_{\frac{\partial f_i}{\partial x_{i,k}}} \right) = \min_{j=1, \dots, m_i} \left\{ h^* \left(A_{\frac{\partial g_{i,j}}{\partial x_{i,k}}} \right) + v_{i,j} \right\}. \end{aligned}$$

Let $M_i^w \subseteq \{1, \dots, m_i\}$ be the set of indices l such that

$$h^* \left(A_{\frac{\partial g_{i,l}}{\partial x_{i,k}}} \right) + v_{i,l} = \min_{j=1, \dots, m_i} \left\{ h^* \left(A_{\frac{\partial g_{i,j}}{\partial x_{i,k}}} \right) + v_{i,j} \right\}.$$

Then, for each $k = 1, \dots, n_i$, we have

$$\left(\frac{\partial f_i}{\partial x_{i,k}} - \sum_{i=1}^{m_i} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,k}} \right)^w = \begin{cases} \sum_{j \in M_i^w} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,k}}, & \text{Case (a)} \\ \frac{\partial f_i^w}{\partial x_{i,k}}, & \text{Case (b)} \\ \frac{\partial f_i^w}{\partial x_{i,k}} - \sum_{j \in M_i^w} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,k}}, & \text{Case (c)}. \end{cases}$$

Note that for a fixed i , if we consider a generic dense polynomial $p \in \mathbb{C}[x]$ with a fixed multidegree, then we have the same support $A_{\frac{\partial p}{\partial x_{i,k}}}$ for $\frac{\partial p}{\partial x_{i,k}}$, for any $k = 1, \dots, n_i$. Therefore, the values of $h^* \left(A_{\frac{\partial f_i}{\partial x_{i,k}}} \right)$ and $\min_{j=1, \dots, m_i} \left\{ h^* \left(A_{\frac{\partial g_{i,j}}{\partial x_{i,k}}} \right) + v_{i,j} \right\}$ do not depend on the choice of $k = 1, \dots, n_i$. It means that without loss of generality, if we have $h^* \left(A_{\frac{\partial f_i}{\partial x_{i,k}}} \right) > \min_{j=1, \dots, m_i} \left\{ h^* \left(A_{\frac{\partial g_{i,j}}{\partial x_{i,k}}} \right) + v_{i,j} \right\}$ for some $k \in \{1, \dots, n_i\}$, then so do all other indices in $\{1, \dots, n_i\}$.

For each index i , let

$$U_i^w := \mathbf{V}(g_{i,1}^w, \dots, g_{i,m_i}^w) \subset \mathbb{C}^{\hat{n}_1 + \dots + \hat{n}_N},$$

$$\text{Jac}_i^w := [\nabla_{x_i} f_i^w(x) \quad \nabla_{x_i} g_{i,1}^w(x) \quad \dots \quad \nabla_{x_i} g_{i,m_i}^w(x)], \text{ and}$$

$$W_i^w := \{x \in \mathbb{C}^{\hat{n}_1 + \dots + \hat{n}_N} \mid \text{rank}(\text{Jac}_i^w) \leq m_i\}.$$

Denote $V_{-i} = \bigcap_{k \neq i} (W_k^w \cap U_k^w)$. From the genericity of f_i^w and $g_{i,j}^w$, we have that $\dim V_{-i} \leq \hat{n}_i$. Also, note that if $x^* \in W_i^w$, then there exist $\lambda_{i,0}, \dots, \lambda_{i,m_i} \in \mathbb{C}$ such that

$$\lambda_{i,0} \nabla_{x_i} f_i^w(x^*) + \lambda_{i,1} \nabla_{x_i} g_{i,1}^w(x^*) + \dots + \lambda_{i,m_i} \nabla_{x_i} g_{i,m_i}^w(x^*) = 0.$$

It means that if $\left(\frac{\partial f_i}{\partial x_{i,k}} + \sum_{j=1}^{m_i} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,k}} \right)^w(x^*) = 0$, then $x^* \in W_i^w$. Indeed, all nonzero solutions to the facial system, if they exist, must be in $U_i^w \cap W_i^w$ for all $i = 1, \dots, N$. From now on, we prove the desired statement by considering each of the three cases mentioned above respectively.

Case (a). Suppose that there exists $i = 1, \dots, N$ such that

$$\left(\frac{\partial f_i}{\partial x_{i,k}} - \sum_{j=1}^{m_i} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,k}} \right)^w = \sum_{j \in M_i^w} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,k}}.$$

Without loss of generality, we assume $i = 1$. Note that if there is a root (x^*, λ^*) over $(\mathbb{C} \setminus \{\mathbf{0}\})$ of F^w , then

$$(4.1) \quad \sum_{j \in M_1^w} \lambda_{1,j}^* \frac{\partial g_{1,j}}{\partial x_{1,k}}(x^*) = 0.$$

Denote by $(\text{Jac}_i^w)^\circ$ the submatrix of the rightmost m_i columns of Jac_i^w , and

$$(W_1^w)^\circ := \{x \in \mathbb{C}^{\hat{n}_1 + \dots + \hat{n}_N} \mid \text{rank}(\text{Jac}_i^w)^\circ \leq m_i - 1\}.$$

Then the equation (4.1) implies that $x^* \in (W_1^w)^\circ \cap U_1^w$. By [37], $\text{codim}((W_1^w)^\circ \cap U_1^w) \geq n_1 + 1$ when $g_{i,j}$ are generic, thus we have

$$(W_1^w)^\circ \cap U_1^w \cap V_{-1} = \emptyset.$$

This contradicts to the fact that (x^*, λ^*) is a root of F^w .

Case (b). Suppose that there exists $i = 1, \dots, N$ such that

$$(4.2) \quad \left(\frac{\partial f_i}{\partial x_{i,k}} - \sum_{j=1}^{m_i} \lambda_{i,j} \frac{\partial g_{i,j}}{\partial x_{i,k}} \right)^w = \frac{\partial f_i^w}{\partial x_{i,k}}.$$

Without loss of generality assume that $i = 1$. In this case, we may further assume that $m_1 \neq 0$ because if $m_1 = 0$, then it can be considered as a special case of Case (c). For a root (x^*, λ^*) for the facial system, we have $\frac{\partial f_1^w}{\partial x_{1,k}}(x^*) = 0$ for all $k = 1, \dots, n_1$. Then, $\mathbf{V}(\frac{\partial f_1^w}{\partial x_{1,k}} \mid k = 1, \dots, n_1) \cap V_{-1}$ has the dimension at most zero due to the genericity of f_1^w . Hence, the genericity of $g_{1,j}^w$ concludes that there is no point in $\bigcap_{i=1}^N (W_i^w \cap U_i^w)$ satisfying (4.2).

Case (c). As the first two cases cannot be happened, we may assume that

$$h^* \left(A \frac{\partial f_i}{\partial x_{i,k}} \right) = \min_{j=1, \dots, m_i} \left\{ h^* \left(A \frac{\partial g_{i,j}}{\partial x_{i,k}} \right) + v_{i,j} \right\}$$

for all $i = 1, \dots, N$. We consider two subcases, the case that $w_{i,k} \geq 0$ for each indices $i = 1, \dots, N$ and $k \in \mathbf{I}_i^w$, and the case that there is $i \in 1, \dots, N$ such that $w_{i,k} < 0$ for some $k \in \mathbf{I}_i^w$.

First, we assume that $w_{i,k} \geq 0$ for each index i and $k \in \mathbf{I}_i^w$. Because we consider a dense polynomial f_i , its partial derivatives are also dense polynomials. Thus, we have $\mathbf{0}$ in the support A_p for each $p \in \{\frac{\partial f_i}{\partial x_{i,k}} \mid k \in \mathbf{I}_i^w\}$. Therefore, we have $0 \geq h^*(A_p)$. Since all $w_{i,k} \geq 0$, we also have $h^*(A_p) \geq 0$, and so we get $h^*(A_p) = 0$ for each p . It further concludes that $w_{i,k} = 0$ for all i and $k \in \mathbf{I}_i^w$. Also, since

$$h^* \left(A \frac{\partial f_i}{\partial x_{i,k}} \right) = \min_{j=1, \dots, m_i} \left\{ h^* \left(A \frac{\partial g_{i,j}}{\partial x_{i,k}} \right) + v_{i,j} \right\} = 0$$

for each i , we have $\min_{j=1, \dots, m_i} v_{i,j} = 0$. We assume that there is at least one index $i \in 1, \dots, N$ such that $v_{i,j} > 0$ for some $j \in \{1, \dots, m_i\}$. Otherwise, w can be considered as just a zero vector and there is nothing to prove. Recall that M_i^w is a subset of $\{1, \dots, m_i\}$ such that $v_l = \min_{j=1, \dots, m_i} v_{i,j} = 0$ for all $l \in M_i^w$. Then, we know that $M_i^w \subsetneq \{1, \dots, m_i\}$ for some $i = 1, \dots, N$; otherwise, F^w equals to F . Without

loss of generality, let $i = 1$ be such an index. Then, the size of M_1^w is exactly the number of variables $\lambda_{1,j}$ that appear in F_1^w (i.e., $\lambda_{1,j}$ variable appears in F_1^w if and only if $j \in M_1^w$). Without loss of generality, we further assume $M_1^w = \{1, \dots, \hat{m}_1\}$, and let $\widehat{\text{Jac}}_i^w$ be the submatrix of the left most $\hat{m}_1 + 1$ columns of Jac_i^w . If (x^*, λ^*) is a nonzero solution to the facial system, then

$$\text{rank}(\widehat{\text{Jac}}_i^w(x^*)) \leq \hat{m}_i.$$

Denote the determinantal variety by \widehat{W}_1^w . We have

$$\text{codim}(\widehat{W}_1^w \cap \mathbf{V}(g_{1,1}, \dots, g_{1,\hat{m}_1})) = \hat{n}_1.$$

Note that $g_{1,j}(x^*) = 0$ for all $j = 1, \dots, m_i$, and $\hat{m}_1 < m_1$, such x^* does not exist when f_1 and $g_{i,j}$ are generic.

Lastly, we deal with the subcase that there exists $i \in \{1, \dots, N\}$ such that $w_{i,k} < 0$ for some $k \in \mathbf{I}_i^w$. Without loss of generality, assume that $i = 1$ and suppose that $w_{1,\hat{k}} < 0$ for some $\hat{k} \in \mathbf{I}_1^w$. Since there is a negative entry $w_{1,\hat{k}}$, we have $h_1^*(A_{g_{1,t}}) < 0$ for some $t \in \{1, \dots, m_1\}$. Furthermore, suppose that we have a root (x^*, λ^*) of F^w . Note that $g_{1,j}^w(x^*) = 0$ for all $j = 1, \dots, m_i$. Let $t \in M_1^w$ be the index such that $h^*(A_{g_{1,t}}) < 0$. Then, by Lemma 4.3, we have

$$\begin{aligned} 0 &= h^*(A_{g_{1,t}}) \lambda_{1,t} g_{1,t}^w(x^*) = \sum_{i=1}^N \sum_{k \in \mathbf{I}_i^w} w_{i,k} x_{i,k} \lambda_{1,t} \frac{\partial g_{1,t}^w}{\partial x_{i,k}}(x^*) \\ &= \sum_{k \in \mathbf{I}_1^w} w_{1,k} x_{1,k} \left(\frac{\partial f_1^w}{\partial x_{1,k}}(x^*) - \sum_{j \in M_1^w \setminus \{t\}} \lambda_{1,j} \frac{\partial g_{1,j}^w}{\partial x_{1,k}}(x^*) \right) \\ &\quad + \sum_{i=2}^N \sum_{k \in \mathbf{I}_i^w} w_{i,k} x_{i,k} \lambda_{1,t} \frac{\partial g_{1,t}^w}{\partial x_{i,k}}(x^*) \\ &= h_{w_1}^*(A_{f_1}) f_1^w(x^*) - \sum_{j \in M_1^w \setminus \{t\}} \lambda_{1,j} h_{w_1}^*(A_{g_{1,j}}) g_{1,j}^w(x^*) \\ &\quad + \sum_{i=2}^N h_{w_i}^*(A_{g_{1,t}}) \lambda_{1,t} g_{1,t}^w(x^*). \end{aligned}$$

In the above, the third equality holds due to the fact that

$$\frac{\partial f_1^w}{\partial x_{1,k}}(x^*) - \sum_{j \in M_1^w} \lambda_{1,j} \frac{\partial g_{1,j}^w}{\partial x_{1,k}}(x^*) = 0,$$

and the last equality is obtained by applying Lemma 4.3. Let

$$q = h_{w_1}^* \cdot (A_{f_1}) f_1^w - \sum_{j \in M_1^w \setminus \{t\}} \lambda_{1,j} h_{w_1}^* \cdot (A_{g_{1,j}}) g_{1,j}^w + \sum_{i=2}^N h_{w_i}^* \cdot (A_{g_{1,t}}) \lambda_{1,t} g_{1,t}^w$$

be the polynomial obtained from the last equality. We know that a point x^* lies in $\mathbf{V}(q)$. It means that $q(x^*) = -h_{w_1}^*(A_{f_1}) \cdot f_1^w(x^*) = 0$ since $x^* \in U_1^w$. However, it contradicts to the genericity of f_1 . \square

Remark 4.4. (1) For the GNEP, if the defining functions are generic polynomials, then the set of complex KKT tuples is finite, and all KKT tuples lie in the torus. This is implied by [37, Theorem 3.1]. In this case, Bernstein

genericity implies that the mixed volume agrees with the algebraic degree. The explicit formula for the algebraic degree of generic GNEPs is studied in the recent paper [37].

- (2) Even when defining functions for the GNEP are not generic, the mixed volume still is an upper bound for the number of isolated solutions in the torus by Theorem 2.1. In this case, we may still find all complex KKT tuples using the homotopy continuation. However, it is still open in general that how to justify the completeness of solutions of a system found by the homotopy continuation. For partial results on the test of checking completeness, see [10, 31].

5. NUMERICAL EXAMPLES

In this section, we present some numerical experiments of solving GNEPs of polynomials using the polyhedral homotopy continuation. We apply the software `HomotopyContinuation.jl` to find complex KKT points of GNEPs by the polyhedral homotopy continuation, and apply `Gloptipoly3` to implement the Moment-SOS hierarchy of semidefinite relaxations for verifying GNEs. The computation is executed in a Macbook pro, 2 GHz Quad-Core Intel Core i5, 32 GB RAM.

When the GNEP is convex, if the complex KKT tuple $(x, \lambda_1, \dots, \lambda_N)$ satisfying

$$g_{i,j}(x) \geq 0, \lambda_{i,j} \geq 0 \quad \text{for all } i \in \{1, \dots, N\}, j \in \mathcal{I}_i,$$

i.e., if x is a KKT point, then it is a GNE. For nonconvex GNEPs, the tuple of strategies x is a GNE if and only if the

$$\delta := \min_{i=1, \dots, N} \delta_i \geq 0,$$

where δ_i is given by (3.2). The δ is called the *accuracy parameter* for x . In practical computation, one may not get $\min\{g_{i,j}(x), \lambda_{i,j}, \delta\} \geq 0$ exactly, due to rounding-off errors. In this section, we regard x being a GNE if $\min\{g_{i,j}(x), \lambda_{i,j}, \delta\} \geq -10^{-6}$.

Example 5.1. (i) Consider a 2-player NEP in [38]

$$\begin{aligned} \text{1st player: } & \begin{cases} \min_{x_1 \in \mathbb{R}^3} & \sum_{j=1}^3 x_{1,j}(x_{1,j} - j \cdot x_{2,j}) \\ \text{s.t.} & 1 - x_{1,1}x_{1,2} \geq 0, 1 - x_{1,2}x_{1,3} \geq 0, x_{1,1} \geq 0, \end{cases} \\ \text{2nd player: } & \begin{cases} \min_{x_2 \in \mathbb{R}^3} & \prod_{j=1}^3 x_{2,j} + \sum_{\substack{1 \leq i < j \leq 3 \\ 1 \leq k \leq 3}} x_{1,i}x_{1,j}x_{2,k} + \sum_{\substack{1 \leq i \leq 3 \\ 1 \leq j < k \leq 3}} x_{1,i}x_{2,j}x_{2,k} \\ \text{s.t.} & 1 - (x_{2,1})^2 - (x_{2,2})^2 - (x_{2,3})^2 = 0. \end{cases} \end{aligned}$$

This is a nonconvex NEP, since both players' optimization problems are nonconvex. Moreover, the feasible set for the first player's optimization problem is unbounded. By implementing the polyhedral homotopy continuation on the complex KKT system, we got 252 complex KKT tuples, and 8 of them satisfy the KKT system (2.1). Since this is a nonconvex problem, we ran Algorithm 3.1 for selecting NEs. We obtained four NEs $u = (u_1, u_2)$ with

$$\begin{aligned} u_1 &= (0.3198, 0.6396, -0.6396), & u_2 &= (0.6396, 0.6396, -0.4264); \\ u_1 &= (0.0000, 0.3895, 0.5842), & u_2 &= (-0.8346, 0.3895, 0.3895); \\ u_1 &= (0.2934, -0.5578, 0.8803), & u_2 &= (0.5869, -0.5578, 0.5869); \\ u_1 &= (0.0000, -0.5774, -0.8660), & u_2 &= (-0.5774, -0.5774, -0.5774). \end{aligned}$$

Their accuracy parameters are respectively

$$-5.2324 \cdot 10^{-11}, 1.7619 \cdot 10^{-9}, -4.8633 \cdot 10^{-9}, -7.1933 \cdot 10^{-9}.$$

Note that for this NEP, the mixed volume of the complex KKT system equals 252. The polyhedral homotopy found all complex KKT tuples, so all NEs are obtained by our method. It took about 7.81 seconds to find all NEs including 4 seconds to find all complex KKT tuples, and about 3.81 seconds to verify NEs.

(ii) If the second player's objective becomes

$$-\prod_{j=1}^3 x_{2,j} + \sum_{\substack{1 \leq i \leq 3 \\ 1 \leq j < k \leq 3}} x_{1,i} x_{2,j} x_{2,k} - \sum_{\substack{1 \leq i < j \leq 3 \\ 1 \leq k \leq 3}} x_{1,i} x_{1,j} x_{2,k},$$

then the polyhedral homotopy continuation found 252 complex KKT tuples, and there are 3 of them satisfying the KKT system (2.1). However, none of these KKT points are NEs, by Algorithm 3.1. Indeed, since the mixed volume for the complex KKT system equals 252, all complex KKT tuples were found by homotopy continuation. Therefore, we detected that this NEP does not have any NE. It took around 3 seconds to solve the complex KKT system, and 1.09 seconds to detect nonexistence of NEs.

Example 5.2. Consider a GNEP variation of the problem in Example 5.1(i).

$$\begin{aligned} \text{1st player: } & \left\{ \begin{array}{l} \min_{x_1 \in \mathbb{R}^3} \quad \sum_{j=1}^3 x_{1,j} (x_{1,j} - j \cdot x_{2,j}) \\ \text{s.t.} \quad x_{2,3} - x_{1,1} x_{1,2} \geq 0, \quad x_{2,1} - x_{1,2} x_{1,3} \geq 0, \quad x_{1,1} - x_{2,2} \geq 0, \end{array} \right. \\ \text{2nd player: } & \left\{ \begin{array}{l} \min_{x_2 \in \mathbb{R}^3} \quad \prod_{j=1}^3 x_{2,j} + \sum_{\substack{1 \leq i < j \leq 3 \\ 1 \leq k \leq 3}} x_{1,i} x_{1,j} x_{2,k} + \sum_{\substack{1 \leq i \leq 3 \\ 1 \leq j < k \leq 3}} x_{1,i} x_{2,j} x_{2,k} \\ \text{s.t.} \quad 1 - (x_{1,1} x_{2,1})^2 - (x_{2,2})^2 - (x_{2,3})^2 = 0. \end{array} \right. \end{aligned}$$

Similar to the problem in Example 5.1(i), this is a nonconvex GNEP, and the first player has an unbounded feasible strategy set. By implementing the polyhedral homotopy continuation on the complex KKT system, we computed the mixed volume 512 and found 484 complex KKT tuples, and 11 of them satisfy the KKT system (2.1). Since this is a nonconvex problem, we ran Algorithm 3.1 for selecting GNEs. We obtained two GNEs $u = (u_1, u_2)$ with

$$\begin{aligned} u_1 &= (0.8188, -0.3213, -0.3947), & u_2 &= (0.8868, 0.6353, -0.2631); \\ u_1 &= (0.5873, -0.5993, 0.6091), & u_2 &= (1.1747, -0.5993, 0.4061). \end{aligned}$$

Their accuracy parameters are respectively

$$-4.0433 \cdot 10^{-9}, \quad -6.7675 \cdot 10^{-9}.$$

It took about 9.85 seconds to find all GNEs including 4 seconds to solve the complex KKT system, and about 5.85 seconds to verify GNEs.

Example 5.3. Consider a 2-player convex GNEP in [39]

$$\begin{aligned} \min_{x_1 \in \mathbb{R}^2} \quad & \sum_{j=1}^2 (x_{1,j} - 1)^2 + x_2 (x_{1,1} - x_{1,2}) \quad \left| \quad \min_{x_2 \in \mathbb{R}^1} \quad (x_2)^3 - x_{1,1} x_{1,2} x_2 - x_2 \right. \\ \text{s.t.} \quad & 2 - x_1^\top x_1 - x_2 \geq 0; \quad \left. \text{s.t.} \quad 3x_2 - x_1^\top x_1 \geq 0, \quad 1 - x_2 \geq 0. \right. \end{aligned}$$

By implementing the polyhedral homotopy continuation on the complex KKT system, we knew the mixed volume is 23, and we got 17 complex KKT tuples. For

these KKT tuples, only one of them satisfies the KKT system (2.1). Because this is a convex GNEP, we got a GNE $u := (u_1, u_2)$ from this KKT tuple with

$$u_1 = (0.4897, 1.0259), \quad u_2 = (0.7077).$$

It took around 2 seconds to solve the complex KKT system.

Example 5.4. Consider a 2-player GNEP

$$\begin{aligned} \text{1st player:} & \begin{cases} \min_{x_1 \in \mathbb{R}^2} & 3x_{2,1}(x_{1,1})^3 + 5(x_{1,2})^3 - 2 \sum_{j=1}^2 x_{1,j} \cdot \sum_{j=1}^2 x_{2,j} \\ \text{s.t.} & 5x_{1,1} - 2x_{1,2} + 3x_{2,2} - 1 \geq 0, \quad 3 - x_{2,1} \cdot x_1^\top x_1 \geq 0, \\ & x_{1,1} \geq -2, \quad x_{1,2} \geq 1; \end{cases} \\ \text{2nd player:} & \begin{cases} \min_{x_2 \in \mathbb{R}^2} & (2x_{1,1} + 3x_{1,2})(x_{2,1})^3 - 3x_{2,1} + 7(x_{2,2})^2 + 5x_{1,1}x_{1,2}x_{2,2} \\ \text{s.t.} & 7x_{1,2} + 3x_{2,2} - 5x_{2,1}^2 + 3 \geq 0, \quad 2x_{2,1} \geq -1, \\ & 2 - x_{2,2} \geq 0, \quad 5 + x_{2,2} \geq 0. \end{cases} \end{aligned}$$

This is a nonconvex GNEP. By implementing the polyhedral homotopy continuation on the complex KKT system, we knew the mixed volume is 480 and found 480 complex KKT tuples. We ran Algorithm 3.1 and obtained the unique GNE $u := (u_1, u_2)$ with

$$u_1 = (0.7636, 1.0000), \quad u_2 = (0.4700, -0.2727), \quad \delta = 1.0220 \cdot 10^{-8}.$$

Note that for this GNEP, the mixed volume of the complex KKT system coincides with the number of complex KKT tuples we found. The polyhedral homotopy found all complex KKT tuples, so all GNEs are obtained by our method. It took around 5.75 seconds to find all GNEs including 4 seconds to solve the complex KKT system, and 1.75 seconds to select the GNE.

Example 5.5. Consider a GNEP whose optimization problems are

$$\begin{aligned} \text{1st player:} & \begin{cases} \min_{x_1 \in \mathbb{R}^2} & 2(x_{1,1})^2 + 7(x_{1,2})^2 + 3x_{1,1} + 5x_{1,2} \\ \text{s.t.} & 1 - 2(x_{1,1})^2 - (x_{1,2})^2 - 3(x_{2,1})^2 - 5(x_{2,2})^2 \geq 0, \\ & 1 - x_{1,1} \geq 0, \quad \frac{1}{2} - x_{1,2} \geq 0; \end{cases} \\ \text{2nd player:} & \begin{cases} \min_{x_2 \in \mathbb{R}^2} & 3(x_{2,2})^2 - 4x_{2,1}x_{2,2} \\ \text{s.t.} & 3(x_{1,1})^2 + (x_{1,2})^2 + \frac{7}{10}(x_{2,1})^2 + 6(x_{2,2})^2 - 1 \geq 0, \\ & 7 - x_{2,1} \geq 0, \quad x_{2,2} - \frac{3}{10} \geq 0, \quad \frac{8}{10} - x_{2,2} \geq 0. \end{cases} \end{aligned}$$

This is a nonconvex GNEP. By implementing the polyhedral homotopy continuation on the complex KKT system, we computed the mixed volume is 168 and found 168 complex KKT tuples. However, none of them are GNEs. It took around 3 seconds to solve this problem, including 3 seconds to solve the complex KKT system, and 0.001 seconds to detect the nonexistence of GNEs.

Example 5.6. Consider a convex GNEP of 3 players. For $i = 1, 2, 3$, the i th player aims to minimize the quadratic function

$$f_i(x) = \frac{1}{2} x_i^\top A_i x_i + x_i^\top (B_i x_{-i} + b_i).$$

All variables have box constraints $-10 \leq x_{i,j} \leq 10$, for all i, j . In addition to them, the first player has linear constraints $x_{1,1} + x_{1,2} + x_{1,3} \leq 20$, $x_{1,1} + x_{1,2} - x_{1,3} \leq x_{2,1} - x_{3,2} + 5$; the second player has $x_{2,1} - x_{2,2} \leq x_{1,2} + x_{1,3} - x_{3,1} + 7$; and the third player has $x_{3,2} \leq x_{1,1} + x_{1,3} - x_{2,1} + 4$.

(i) Consider the case that the values of parameters are set as in [15, Example A.3]:

$$A_1 = \begin{bmatrix} 20 & 5 & 3 \\ 5 & 5 & -5 \\ 3 & -5 & 15 \end{bmatrix}, A_2 = \begin{bmatrix} 11 & -1 \\ -1 & 9 \end{bmatrix}, A_3 = \begin{bmatrix} 48 & 39 \\ 39 & 53 \end{bmatrix},$$

$$B_1 = \begin{bmatrix} -6 & 10 & 11 & 20 \\ 10 & -4 & -17 & 9 \\ 15 & 8 & -22 & 21 \end{bmatrix}, B_2 = \begin{bmatrix} 20 & 1 & -3 & 12 & 1 \\ 10 & -4 & 8 & 16 & 21 \end{bmatrix},$$

$$B_3 = \begin{bmatrix} 10 & -2 & 22 & 12 & 16 \\ 9 & 19 & 21 & -4 & 20 \end{bmatrix}, b_1 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, b_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, b_3 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}.$$

This is a convex GNEP, since for all $i \in \{1, 2, 3\}$, the A_i is positive semidefinite and all constraints are linear. By implementing the polyhedral homotopy continuation on the complex KKT system, we got the mixed volume 12096, and found 11631 complex KKT tuples. There are 5 GNEs obtained by Algorithm 3.1, which are presented in the following table.

	u_1	u_2	u_3
1	(-0.3805,-0.1227,-0.9932)	(0.3903,1.1638)	(0.0504,0.0176)
2	(-0.9018,-4.4017,-2.1791)	(-2.0034,-2.4541)	(-0.0316,2.9225)
3	(-0.8039,-0.3062,-2.3541)	(0.9701, 3.1228)	(0.0751,-0.1281)
4	(1.9630,-1.3944, 5.1888)	(-3.1329,-10.0000)	(-0.0398,1.6392)
5	(0.6269,10.0000,9.3731)	(1.8689,10.0000)	(0.3353,-10.0000)

It took around 177 seconds to solve the complex KKT system. We would like to remark that in [15] and [39], only the first GNE was found, and the second to the fourth GNEs are new solutions found by our algorithm.

(ii) If we let

$$A_1 = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & -5 \\ 3 & -5 & 15 \end{bmatrix},$$

and all other parameters be given as in (i), then this GNEP is nonconvex. By implementing the polyhedral homotopy continuation on the complex KKT system, the mixed volume equals 12096 and we got 11620 complex KKT tuples, and five of them satisfy the KKT condition (2.1). Since this is a nonconvex problem, we ran Algorithm 3.1 for selecting GNEs, and obtained one GNE $u = (u_1, u_2)$ with

$$u_1 = (0.9968, 10.0000, 9.0032), u_2 = (0.6668, 10.0000), u_3 = (0.7283, -10.0000).$$

It took around 209.68 seconds to find all GNEs including 207 seconds to solve the complex KKT system, and 2.68 seconds to select the GNE.

5.1. Comparison with existing methods. In this subsection, we compare the performance of Algorithm 3.1 with some existing methods for solving GNEPs, such as the augmented Lagrangian method (ALM) in [24], Gauss-Sedel method (GSM) in [40], the interior point method (IPM) in [12], and the semidefinite relaxation method (KKT-SDP) in [39]. We tested these methods on all GNEPs of polynomials in Examples 5.1-5.6.

Given a computed tuple $u = (u_1, \dots, u_N)$ for an N -player game, the quantity

$$\kappa := \max \left\{ \max_{\substack{i=1, \dots, N \\ j \in \mathcal{I}_1^{(i)} \cup \mathcal{I}_2^{(i)}}} \{-g_{i,j}(u)\}, \max_{\substack{i=1, \dots, N \\ j \in \mathcal{I}_0^{(i)}}} \{|g_{i,j}(u)|\} \right\}$$

is exploited to measure the feasibility violation. It is clear that u is feasible for the GNEP if and only if $\kappa \leq 0$. Besides that, let δ_i be the minimum for the optimization problem (3.2), and $\delta := \min_{i=1, \dots, N} \delta_i$. Then, u is a GNE if and only if $\kappa \leq 0$ and $\delta \geq 0$. For the KKT-SDP method, the $\kappa \leq 0$ for all computed tuple, and we say the method find a GNE successfully if $\delta \geq -10^{-6}$, due to a numerical error. For other earlier algorithms mentioned above, since they are iterative methods, the stopping criterion is given as the following: For the computed tuple u , when $\kappa \leq 10^{-6}$, we solve (3.2) for each i and compute δ . If we further have $\delta \geq -10^{-6}$, then we stop the iteration and report that the method found a GNE successfully.

For the ALM, GSM and IPM, the same parameters are applied as in [12, 24, 40]. In the augmented-Lagrangian method, the full penalization is used, and a Levenberg-Marquardt type method (see [24, Algorithm 24]) is implemented to solve penalized subproblems. For Gauss-Seidel method, the normalization parameters are updated as (4.3) in [40], and the Moment-SOS relaxations are used to solve normalized subproblems. We let 1000 be the maximum number of iterations for the ALM and IPM, and at most 100 iterations are allowed in the GSM. For initial points, we use $(0, 0, 0, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})$ for Examples 5.1(i-ii), $(0, 0, 0, 0, -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ for Example 5.2, $(0, 0, 1, 1)$ for Example 5.4, $(0, 0, 0, \frac{1}{\sqrt{5}})$ for Example 5.5, and the zero vector for all other problems. For the one-shot KKT-SDP method, randomly generated positive semidefinite matrices are exploited to formulated polynomial optimization. Note that the ALM, IPM and KKT-SDP are designed for finding a KKT point of the GNEP. When the GNEP is convex (e.g., Examples 5.3 and 5.6(i)), the limit point is guaranteed to be a GNE, if these methods produce a convergent sequence. However, since we in general do not make any convexity assumption, it is possible that these methods converge to a KKT point which is not a GNE. For the ALM and IPM, the produced sequence is considered convergent to a KKT point if the last iterate satisfies the KKT conditions up to a small round-off error (say, 10^{-6}). If the iterations is convergent but the stopping criterion is not met, we still solve (3.2) to check if the latest iterating point is a GNE or not.

The numerical results are shown in Table 1. In general, the method can be regarded to solve the GNEP successfully if the error is small (e.g., less than 10^{-6}). For Algorithm 3.1, when there are more than one GNE obtained, we present the largest error among these GNEs.

The comparison is summarized as follows:

- (1) The augmented Lagrangian method converges to a KKT point that is not a GNE for Examples 5.1(i-ii) and 5.6(ii). For Example 5.2, the iteration cannot proceed because the maximum penalty parameter were reached at the 14th iteration. For Examples 5.4 and 5.5, it fails to converge because the penalized subproblem cannot be solved accurately.
- (2) The interior point method converges to a KKT point that is not a GNE for Examples 5.1(ii), 5.4 and 5.6(ii). For Examples 5.2 and 5.5, the algorithm does not converge. In this problem, the Newton type directions usually do not satisfy the sufficient descent conditions.

Example		ALM	IPM	GSM	KKT-SDP	Algorithm 3.1
5.1(i)	time	27.52	13.34	Fail	2.95	12.6
	error	4.67	4.67		1.48	$< 8 \cdot 10^{-9}$ (4 GNEs)
5.1(ii)	time	32.03	8.04	Fail	2.92	7.8
	error	1.11	1.11		0.19	no GNE
5.2	time	Fail	Fail	Fail	4.65	9.9
	error				0.66	$< 7 \cdot 10^{-9}$ (2 GNEs)
5.3	time	0.72	3.14	4.45	1.51	2.0
	error	$2 \cdot 10^{-7}$	$2 \cdot 10^{-7}$	$2 \cdot 10^{-7}$	$8 \cdot 10^{-9}$	$1 \cdot 10^{-8}$
5.4	time	Fail	1.69	11.47	17.89	5.75
	error		$2 \cdot 10^{-7}$	$4 \cdot 10^{-7}$	$1 \cdot 10^{-6}$	$2 \cdot 10^{-8}$
5.5	time	Fail	Fail	Fail	1.51	3
	error				no GNE	no GNE
5.6(i)	time	1.50	3.12	Fail	11.55	177
	error	$1 \cdot 10^{-7}$	$2 \cdot 10^{-7}$		$2 \cdot 10^{-7}$	$< 1 \cdot 10^{-6}$ (5 GNEs)
5.6(ii)	time	59.93	16.19	Fail	11.29	210
	error	123.22	123.22		123.22	$1 \cdot 10^{-7}$

TABLE 1. Comparison with other methods. The “time” gives the consumed time (in seconds) for finding a GNE or KKT points, and the “error” measures the quantity $\max\{|\delta|, \kappa\}$ of computed GNE candidates.

- (3) For Examples 5.1(i-ii) and 5.6(i), the Gauss-Seidel method failed to converge and it alternated between several points. For Examples 5.2, 5.5 and 5.6(ii), the iteration cannot proceed at some stages since global minimizers for normalized subproblems cannot be obtained. Usually, this is because the normalized subproblem is infeasible or unbounded.
- (4) The semidefinite relaxation method obtained a KKT point that is not a GNE for Examples 5.1(i-ii), 5.2 and 5.6(ii).
- (5) Algorithm 3.1 detected nonexistence of GNEs for Examples 5.1(ii) and 5.5. We would like to remark that if there exists KKT points that are not GNEs, then the semidefinite relaxation method may not detect nonexistence of GNEs. For all other GNEPs, Algorithm 3.1 found at least one GNE. Moreover, for Examples 5.1(i), 5.2 and 5.6, Algorithm 3.1 found more than one GNE, and the completeness of GNEs are guaranteed for Examples 5.1(i), 5.2 and 5.4.

5.2. GNEPs of polynomials with randomly generated coefficients. We present numerical results of Algorithm 3.1 on GNEPs defined by polynomials whose coefficients are randomly generated. For the GNEP with N players, we assume that all players have the same number of strategies, i.e., $n_1 = n_2 = \dots = n_N$. The i th

player's optimization problem is given by

$$(5.1) \quad \begin{cases} \min_{x_i \in \mathbb{R}^{n_i}} & f_i(x_i, x_{-i}) \\ s.t. & -x_i^\top A_i x_i + x_{-i}^\top B_i x_i + c_i^\top x \geq d_i. \end{cases}$$

In the above, we have $A_i = R_i^\top R_i$ with a randomly generated matrix $R_i \in \mathbb{R}^{n_i \times n_i}$. Also, $B_i \in \mathbb{R}^{n_i \times (n-n_i)}$, $c_i \in \mathbb{R}^n$, $d_i \in \mathbb{R}$ are randomly generated real matrices or vectors. Under this setting, the constraining function of (5.1) is a convex polynomial in x_i , and the $X_i(x_{-i})$ is compact, for all $x_{-i} \in \mathbb{R}^{n-n_i}$.

For the objective function f_i , we consider two cases. First, we let

$$f_i := x_i^\top \Sigma_i x_i + x_{-i}^\top \Lambda_i x_i + c_i^\top x,$$

where $\Sigma_i = \Theta_i^\top \Theta_i$ with a randomly generated matrix $\Theta_i \in \mathbb{R}^{n_i \times n_i}$, and Λ_i (resp., c_i) is a randomly generated matrix (resp., vector) in $\mathbb{R}^{n_i \times (n-n_i)}$ (resp., in \mathbb{R}^n). In this case, the GNEP given by (5.1) is convex, and all KKT points are GNEs. The second case is for GNEPs without convexity settings. Therefore, we consider a degree d dense polynomial with randomly generated real coefficients, i.e.,

$$f_i := \zeta^\top [x]_d,$$

where ζ is a randomly generated real vector of the proper size. To choose real matrices, vectors and coefficients randomly, we use the `Matlab` function `unifrnd` that chosen real numbers follow the uniform distribution.

The numerical results are presented in Table 2. By Theorem 3.2, if the mixed volume many complex KKT points are obtained, then Algorithm 3.1 can find all GNEs or detect nonexistence of GNEs. Because it considers random examples, the homotopy method mostly finds all mixed volume many KKT points. As the systems bigger, there are some cases that the homotopy method can't find all mixed volume many KKT points due to a numerical issue. Also, note that the homotopy continuation requires more time than selecting GNEs when the size of systems get bigger.

6. CONCLUSIONS AND DISCUSSIONS

This paper studies a new approach for solving GNEPs of polynomials using the polyhedral homotopy continuation and the Moment-SOS relaxations. We show that under some generic assumptions, the mixed volume and the algebraic degree for the complex KKT system are identical, and our method can find all GNEs or detect nonexistence of GNEs. Some numerical experiments are presented to show the effectivity of our method.

For future work, it is interesting to find local GNEs, i.e., find $x = (x_1, \dots, x_N)$ such that each x_i is local minimizer for $F_i(x_{-i})$. Note that every local GNE satisfies the KKT condition. However, it is difficult to select local GNEs from KKT points, especially when *the second-order sufficient optimality conditions* are not satisfied. Moreover, when the $|\mathcal{K}_{\mathbb{C}}|$ is strictly less than the mixed volume for (2.2), how do we know whether our method finds all GNEs or detects nonexistence of GNEs or not? These questions are mostly open, to the best of the authors' knowledge.

(a) Degree 2 convex GNEPs

N	n_i	mixed volume	rate of success	average time
2	2	25	100%	0.0575
2	3	49	100%	0.1721
2	4	81	100%	0.9539
3	2	125	100%	0.9118
3	3	343	100%	3.4150

(b) Degree d nonconvex GNEPs

d	N	n_i	mixed volume	rate of success	average time (seconds)
2	2	2	25	100%	0.0563 + 1.1330
	2	3	49	100%	0.1802 + 1.5098
	2	4	81	100%	0.8819 + 1.9762
	3	2	125	100%	0.8473 + 3.1890
	3	3	343	100%	3.3804 + 6.9738
3	2	2	100	100%	0.1893 + 2.5667
	2	3	484	100%	2.18 + 5.7500
	2	4	2116	98%	21.483 + 17.3477
	3	2	1000	97%	5.255 + 14.4360
4	2	2	289	100%	0.8270 + 4.4256
	2	3	2809	95%	24.533 + 21.9054
	3	2	4913	95%	44.0899 + 40.6792

TABLE 2. Numerical results for random GNEPs. N and n indicates the number of players and the number of variables for each player respectively. The “rate of success” indicates the percentage of GNEPs such that the homotopy continuation finds the mixed volume many KKT points. The “average time” represents the average of the elapsed times for (KKT points computation) + (GNE selection) in seconds. For convex problems, the elapsed time is only measured for the KKT points computation.

ACKNOWLEDGEMENTS

The authors would like to thank Jiawang Nie for motivating this paper and helpful comments. Xindong Tang is partially supported by the Start-up Fund P0038976/BD7L from The Hong Kong Polytechnic University.

REFERENCES

- [1] D. Ardagna, M. Ciavotta, and M. Passacantando. Generalized Nash equilibria for the service provisioning problem in multi-cloud systems. *IEEE Transactions on Services Computing*, 10(3):381–395, 2015.
- [2] K. J. Arrow and G. Debreu. Existence of an equilibrium for a competitive economy. *Econometrica: Journal of the Econometric Society*, pages 265–290, 1954.

- [3] Q. Ba and J.-S. Pang. Exact penalization of generalized Nash equilibrium problems. *Operations Research*, 2020.
- [4] D. N. Bernshtein. The number of roots of a system of equations. *Functional Analysis and its applications*, 9(3):183–185, 1975.
- [5] D. Bertsekas, *Nonlinear Programming*, third edition, Athena Scientific, 2016.
- [6] L. Blum, F. Cucker, M. Shub, and S. Smale. *Complexity and real computation*. Springer Science & Business Media, 2012.
- [7] P. Breiding, K. Rose, and S. Timme. Certifying zeros of polynomial systems using interval arithmetic. *arXiv preprint arXiv:2011.05000*, 2020.
- [8] P. Breiding, F. Sottile, and J. Woodcock. Euclidean distance degree and mixed volume. *Foundations of Computational Mathematics*, pages 1–23, 2021.
- [9] P. Breiding and S. Timme. `HomotopyContinuation.jl`: A package for homotopy continuation in Julia. In *International Congress on Mathematical Software*, pages 458–465. Springer, 2018.
- [10] T. Brysiewicz and M. Burr. Sparse trace tests. *arXiv preprint arXiv:2201.04268*, 2022.
- [11] G. Debreu. A social equilibrium existence theorem. *Proceedings of the National Academy of Sciences*, 38(10):886–893, 1952.
- [12] A. Dreves, F. Facchinei, C. Kanzow, and S. Sagratella. On the solution of the KKT conditions of generalized Nash equilibrium problems. *SIAM Journal on Optimization*, 21(3):1082–1108, 2011.
- [13] F. Facchinei, A. Fischer, and V. Piccialli. On generalized Nash games and variational inequalities. *Operations Research Letters*, 35(2):159–164, 2007.
- [14] F. Facchinei and C. Kanzow. Generalized Nash equilibrium problems. *Annals of Operations Research*, 175(1):177–211, 2010.
- [15] F. Facchinei and C. Kanzow. Penalty methods for the solution of generalized Nash equilibrium problems. *SIAM Journal on Optimization*, 20(5):2228–2253, 2010.
- [16] F. Facchinei and J.-S. Pang. 12 Nash equilibria: the variational approach. *Convex optimization in signal processing and communications*, page 443, 2010.
- [17] D. Grayson and M. Stillman, Macaulay2, a software system for research in algebraic geometry, <http://www.math.uiuc.edu/Macaulay2/>.
- [18] G. Gürkan and J.-S. Pang. Approximations of Nash equilibria. *Mathematical Programming*, 117(1):223–253, 2009.
- [19] P. T. Harker. Generalized Nash games and quasi-variational inequalities. *European journal of Operational research*, 54(1):81–94, 1991.
- [20] J. D. Hauenstein and F. Sottile. `alphaCertified`: Software for certifying numerical solutions to polynomial equations. Available at math.tamu.edu/~sottile/research/stories/alphaCertified, 2011.
- [21] D. Henrion and J. Lasserre. Detecting global optimality and extracting solutions in `GloptiPoly`. Positive polynomials in control, 293310. *Lecture Notes in Control and Inform. Sci.* 312, 2005.
- [22] D. Henrion, J.-B. Lasserre, and J. Löfberg. `GloptiPoly3`: moments, optimization and semi-definite programming. *Optimization Methods & Software*, 24(4-5):761–779, 2009.
- [23] B. Huber and B. Sturmfels. A polyhedral method for solving sparse polynomial systems. *Mathematics of computation*, 64(212):1541–1555, 1995.
- [24] C. Kanzow and D. Steck. Augmented Lagrangian methods for the solution of generalized Nash equilibrium problems. *SIAM Journal on Optimization*, 26(4):2034–2058, 2016.
- [25] J. B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM Journal on optimization*, 11(3):796–817, 2001.
- [26] J. B. Lasserre. *An introduction to polynomial and semi-algebraic optimization*, volume 52. Cambridge University Press, 2015.
- [27] J. Lasserre, *The Moment-SOS Hierarchy*, Proceedings of the International Congress of Mathematicians (ICM 2018), vol. 3, B. Sirakov, P. Ney de Souza and M. Viana (Eds.), 3761–3784, World Scientific, 2019.
- [28] M. Laurent, Optimization over polynomials: Selected topics, *Proceedings of the International Congress of Mathematicians*, S. Jang, Y. Kim, D-W. Lee, and I. Yie (eds.), ICM 2014, 843–869, 2014.
- [29] K. Lee. The NumericalCertification package in Macaulay2. *arXiv preprint arXiv:2208.01784*, 2022.

- [30] T.-L. Lee, T.-Y. Li, and C.-H. Tsai. **HOM4PS-2.0**: a software package for solving polynomial systems by the polyhedral homotopy continuation method. *Computing*, 83(2):109–133, 2008.
- [31] A. Leykin, J. I. Rodriguez, and F. Sottile. Trace test. *Arnold Mathematical Journal*, 4(1):113–125, 2018.
- [32] T.-Y. Li and X. Wang. The BKK root count in \mathbb{C}^n . *Mathematics of computation*, pages 1477–1484, 1996.
- [33] M. Liu, and O. Tuzel, Coupled generative adversarial networks. *Advances in neural information processing systems*, 29:469–477, 2016.
- [34] R. E. Moore, R. B. Kearfott, and M. J. Cloud. *Introduction to interval analysis*, volume 110. Siam, 2009.
- [35] J. Nie. Certifying convergence of Lasserre’s hierarchy via flat truncation. *Mathematical Programming*, 142(1):485–510, 2013.
- [36] J. Nie. Optimality conditions and finite convergence of Lasserre’s hierarchy. *Math. Program.*, 146(1-2):97–121, 2014.
- [37] J. Nie, K. Ranestad, X. Tang. Algebraic degree of generalized Nash equilibrium problems of polynomials. *arXiv preprint arXiv:2208.00357*, 2022.
- [38] J. Nie and X. Tang. Nash equilibrium problems of polynomials. *arXiv preprint arXiv:2006.09490*, 2020.
- [39] J. Nie and X. Tang. Convex generalized Nash equilibrium problems and polynomial optimization. *Mathematical Programming*, pages 1–34, 2021.
- [40] J. Nie, X. Tang, and L. Xu. The Gauss–Seidel method for generalized Nash equilibrium problems of polynomials. *Computational Optimization and Applications*, 78(2):529–557, 2021.
- [41] J. Nie, X. Tang, and S. Zhong. Rational generalized Nash equilibrium problems. *arXiv preprint arXiv:2110.12120*, 2021.
- [42] J.-S. Pang, G. Scutari, F. Facchinei, and C. Wang. Distributed power allocation with rate constraints in gaussian parallel interference channels. *IEEE Transactions on Information Theory*, 54(8):3471–3489, 2008.
- [43] M. Putinar. Positive polynomials on compact semi-algebraic sets. *Indiana University Mathematics Journal*, 42(3):969–984, 1993.
- [44] L. J. Ratliff, S. A. Burden, and S. S. Sastry. Characterization and computation of local Nash equilibria in continuous games. In *2013 51st Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 917–924. IEEE, 2013.
- [45] A. Sommese and C. Wampler. *The Numerical Solution of Systems of Polynomials Arising in Engineering and Science*. World Scientific, 2005.
- [46] B. Sturmfels. *Solving systems of polynomial equations*. Number 97 in CBMS Regional Conference Series in Mathematics. American Mathematical Soc., 2002.
- [47] J. Verschelde. Algorithm 795: **PHCpack**: A general-purpose solver for polynomial systems by homotopy continuation. *ACM Transactions on Mathematical Software (TOMS)*, 25(2):251–276, 1999.
- [48] A. Von Heusinger and C. Kanzow. Optimization reformulations of the generalized Nash equilibrium problem using Nikaido-Isoda-type functions. *Computational Optimization and Applications*, 43(3):353–377, 2009.
- [49] A. Von Heusinger and C. Kanzow. Relaxation methods for generalized Nash equilibrium problems with inexact line search. *Journal of Optimization Theory and Applications*, 143(1):159–183, 2009.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA SAN DIEGO, 9500 GILMAN DRIVE,
LA JOLLA, CA 92093

Email address: ki1004@ucsd.edu

URL: <https://klee669.github.io>

DEPARTMENT OF APPLIED MATHEMATICS, THE HONG KONG POLYTECHNIC UNIVERSITY, HUNG
HOM, KOWLOON, HONG KONG

Email address: xindong.tang@polyu.edu.hk

URL: <https://www.polyu.edu.hk/ama/profile/txd/>