

CYLINDRICAL STOCHASTIC INTEGRATION

JOHANNES ASSEFA AND PHILIPP HARMS

ABSTRACT. We show that any given integral can be turned into a cylindrical integral by the simple application of a Hom functor. In this way, we construct a theory of cylindrical stochastic integration and cylindrical stochastic evolution equations. As an application in mathematical finance, we develop cylindrically measure-valued forward rate models. These can be used to describe term structures with jumps in the underlying at predictable times, as observed in interest rate and energy markets.

CONTENTS

1. Introduction	1
2. Cylindrical vector integration	2
3. Cylindrical stochastic analysis	6
4. Cylindrically measure-valued term structure models	8
Appendix A. Measurable and predictable versions	15
References	16

1. INTRODUCTION

Cylindrical random variables came up in the 1960s and 1970s in the development of probability theory on Banach spaces and locally convex vector spaces [34]. Their raison d'être is that they are easy to construct, for instance by specifying their Fourier transform or by writing down a stochastic integral equation in weak form. However, it can be notoriously difficult to verify that a cylindrical random variable is non-cylindrical, i.e., a random variable in the usual sense. This is one of the main challenges in the construction of infinite-dimensional stochastic integrals, and the answer is negative beyond e.g. nuclear spaces [36] or UMD Banach spaces [31, 37].

We circumvent these difficulties by fully embracing the cylindrical approach. Thus, we give up on the hard problem of constructing non-cylindrical integrals and instead work with much simpler cylindrical integrals. Specifically, given any existing integral, stochastic or not, we define a new cylindrical integral by the simple application of a Hom functor; see Theorem 2.6. In important special cases, this yields the cylindrical Brownian or Lévy integrals of [3, 8, 30, 32]. Via the Hom functor, a-priori estimates for non-cylindrical stochastic evolution equations translate into a-priori estimates for cylindrical stochastic evolution equations, and we develop a corresponding theory in Theorem 3.4.

Date: August 15, 2022.

2020 Mathematics Subject Classification. 60H05 (Primary), 60H15, 91G30 (Secondary).

The authors thank Christa Cuchiero, Claudio Fontana, Johannes Müller, Thorsten Schmidt, Guillaume Szulda, and Josef Teichmann for helpful discussions. Philipp Harms gratefully acknowledges financial support by the National Research Foundation Singapore under the award NRF-NRFF13-2021-0012 and by Nanyang Technological University Singapore under the award NAP-SUG.

From a financial perspective, cylindrical processes can be seen as models of large financial markets. Indeed, a large financial market is simply a collection of tradeable assets, all of which are semimartingales, with no further structure imposed [17]. By the linearity of the stochastic integral, one may assume without loss of generality that linear combinations of tradeable assets are tradeable. Then, a large financial market is precisely a cylindrical semimartingale, i.e., a collection of semimartingales indexed by the elements of some linear space.

We exploit this connection to define cylindrical models for financial term structures with discontinuities. The discontinuities are caused by jumps in the underlying at predictable times and are empirically well documented in real-world markets. For example, interest rates are influenced by policy updates of the central bank, stock prices are influenced by earnings announcements of the company, energy prices are influenced by maintenance works in the power grid, and all of these events tend to happen at pre-scheduled times; cf. Figs. 1 and 2. The immediate modeling implication is that the forward rates cannot be continuous functions of the time to maturity but must be measure-valued [15, 27].

We develop a Heath–Jarrow–Morton (HJM) framework for measure-valued forward rates as in [27]. Actually, the forward rates are merely cylindrically measure-valued because measure-valued stochastic convolutions are ill-defined, as explained in Remark 4.1. Nevertheless, bond prices and a roll-over bank account are well-defined, as shown in Theorem 4.5, and the corresponding large financial market satisfies no asymptotic free lunch with vanishing risk [17] if a HJM drift condition is satisfied under an equivalent probability measure. Extensions to more general driving noise would be possible within the same framework, thanks to the universal construction of the cylindrical stochastic integral. Our modeling framework is versatile and can be applied to other financial markets, as well, including energy markets. In principle, it also lends itself to numerical discretization by similar methods as for non-cylindrical models [1, 12], but we have not explored this any further.

Markovian methods constitute an interesting complementary approach for constructing measure-valued processes. They can be used to treat non-Lipschitz coefficients as for instance in super Brownian motion [25] and more general affine or polynomial measure-valued processes [14, 16, 19]. Notably, these processes are non-cylindrical and take values in the cone of non-negative measures. There are wide-ranging financial applications in stochastic portfolio theory [18], rough volatility modeling [20], and recently also term structure modeling [15]. The connection to cylindrical stochastic analysis is via martingale problems with cylindrical test functionals, as explained in Remark 4.8.

The structure of the paper is as follows. Section 2 develops the cylindrical integral in full generality and explains connections to cylindrical vector measures. Section 3 considers the special case of stochastic integration and establishes well-posedness of cylindrical stochastic evolution equations. Section 4 develops applications to cylindrically measure-valued forward rate models. Appendix A contains some auxiliary results on measurable versions of stochastic processes.

2. CYLINDRICAL VECTOR INTEGRATION

This section describes both abstractly and by examples how any given integral can be turned into a cylindrical integral by the application of a Hom functor. Throughout this section, E , F , G , and H are Banach spaces, and there is a continuous bilinear function $F \times G \rightarrow H$, which is denoted by juxtaposition. Moreover, S is a set, \mathcal{A} is an algebra of subsets of S , $\mu : \mathcal{A} \rightarrow G$ is finitely additive, and Σ is the sigma algebra generated by \mathcal{A} . A function $f : S \rightarrow F$ is called elementary if




FIGURE 1. EONIA rates jump at predictable times due to pre-scheduled monetary policy meetings of the ECB and due to regulatory constraints coming into effect at pre-scheduled dates. Figure taken from [27].




FIGURE 2. Half hourly electricity prices in the UK jumped at predictable times in October 2016 due to pre-scheduled maintenance work on French nuclear power plants. Figure taken from [38].

it is of the form $f = \sum_{i=1}^n f_i \mathbb{1}_{A_i}$ for some $f_i \in F$, $A_i \in \mathcal{A}$, and $n \in \mathbb{N}$. The space of elementary functions with the supremum norm is denoted by $\mathcal{L}_{\text{elem}}^{\infty}(S, F)$. Limits of pointwise converging sequences of elementary functions are called strongly measurable. The space of bounded strongly measurable functions is denoted by $\mathcal{L}^{\infty}(S, F)$.

Definition 2.1 (Elementary integrals). For any $f = \sum_{i=1}^n f_i \mathbb{1}_{A_i} \in \mathcal{L}_{\text{elem}}^{\infty}(S, F)$ and $A \in \mathcal{A}$, we define

$$\int_A f \mu := \sum_i f_i \mu(A \cap A_i) \in H.$$

For fixed A and varying f , one obtains the elementary integral

$$\mathcal{L}_{\text{elem}}^{\infty}(S, F) \ni f \mapsto \int_A f \mu \in H.$$

We write $\|\mathbb{1}_A \mu\|_{F,G,H}$ for its operator norm, which is also called semivariation. For fixed f and varying A , this is a finitely additive function

$$f\mu : \mathcal{A} \ni A \mapsto \int_A f\mu \in H,$$

and the operator norm of its integral over scalar elementary integrands is $\|f\|_\mu := \|f\mu\|_{\mathbb{R},H,H}$.

A major challenge in building a good integration theory is to extend the elementary integral to a large set of integrands. We next list a selection of well-known examples.

Example 2.2 (Integral on bounded measurable functions). Assume that F is finite dimensional. Then $\mathcal{L}_{\text{elem}}^\infty(S, F)$ is dense in $\mathcal{L}^\infty(S, F)$. (Otherwise, it may be a strict subspace.) If $\|\mu\|_{F,G,H} < \infty$, then the elementary integral extends uniquely to a continuous linear operator

$$\mathcal{L}^\infty(S, F) \ni f \mapsto \int_S f\mu \in H.$$

A set $A \subset S$ is called a μ -null set if

$$\inf \left\{ \sum_{i \in \mathbb{N}} \|\mathbb{1}_{A_i} \mu\|_{F,G,H} : A_i \in \mathcal{A}, A \subseteq \bigcup_{i \in \mathbb{N}} A_i \right\} = 0.$$

The space $L^\infty(S, F)$ is defined as the quotient of $\mathcal{L}^\infty(S, F)$ modulo equality up to null sets, and the elementary integral factors through a continuous linear operator

$$L^\infty(S, F) \ni f \mapsto \int_A f\mu \in H.$$

Example 2.3 (Bartle integral). [5] Let $\mu : \Sigma \rightarrow G$ be countably additive on $\mathcal{A} = \Sigma$ with $\|\mu\|_{F,G,H} < \infty$ and suppose that there exists a countably additive function $\nu : \Sigma \rightarrow \mathbb{R}_{\geq 0}$ such that $\nu(A) \rightarrow 0$ is equivalent to $\|\mathbb{1}_A \mu\|_{F,G,H} \rightarrow 0$. Then the elementary integral extends uniquely to a continuous linear map

$$\mathcal{L}^\infty(S, F) \ni f \mapsto \int_S f\mu \in H,$$

where $\mathcal{L}^\infty(S, F)$ carries the structure¹ of boundedly pointwise convergence.

Example 2.4 (Lebesgue integral). Let $F = G = H = \mathbb{R}$ with bilinear form given by scalar multiplication, let $\mu : \Sigma \rightarrow \mathbb{R}_{\geq 0}$ be countably additive on $\mathcal{A} = \Sigma$, and let $L^1(S, \mathbb{R})$ be the completion of $\mathcal{L}_{\text{elem}}^\infty(S, \mathbb{R})$ with respect to the seminorm $\|\cdot\|_\mu$. Then $L^1(S, \mathbb{R})$ and can be identified with a set of measurable functions $S \rightarrow \mathbb{R}$ modulo equality up to null sets, and the elementary integral extends to a contraction

$$L^1(S, \mathbb{R}) \ni f \mapsto \int_S f\mu \in \mathbb{R}.$$

Example 2.5 (Bochner integral). [10] Let $F = H$ and $G = \mathbb{R}$ with bilinear form $F \times \mathbb{R} \rightarrow F$ given by scalar-times-vector multiplication, let $\mu : \Sigma \rightarrow \mathbb{R}_{\geq 0}$ be non-negative and countably additive on $\mathcal{A} = \Sigma$, and let $L^1(S, F)$ be the completion of $\mathcal{L}_{\text{elem}}^\infty(S, F)$ with respect to the norm $\|\cdot\|_\mu$. Then $L^1(S, F)$ and can be identified with the set of strongly measurable functions $f : S \rightarrow F$ such that $\|f\| \in L^1(S, \mathbb{R})$, modulo equality up to null sets. Moreover, the elementary integral extends to a contraction

$$L^1(S, H) \ni f \mapsto \int_S f\mu \in H.$$

¹Boundedly pointwise convergence is a non-topological notion of convergence; see e.g. [24].

Some further examples of integrals are presented in Section 3. Importantly, the goal of this work is *not* to develop a new integral in the sense of adding to this list of examples. Rather, we want to point out that any existing integral can be turned into a new *cylindrical integral* by a very simple construction, namely, by the application of a Hom functor.

Theorem 2.6 (Cylindrical integration). *Assume that the elementary integral in Definition 2.1 has a continuous extension to some Banach space I of integrands:*

$$I \ni f \mapsto \int_S f \mu \in H.$$

If E is non-trivial, then applying the functor $L(E, \cdot)$ yields a continuous linear operator

$$L(E, I) \ni f \mapsto \left(E \ni e \mapsto \int_S f(e) \mu \in H \right) \in L(E, H),$$

which has the same norm as the original integral, and which we call the cylindrical integral.

Proof. If E is non-trivial, then the functor $L(E, \cdot)$ is norm-preserving. \square

As an aside, the name cylindrical integral stems from stochastic integration: applying the Hom functor to the semimartingale integral produces an integral with values in a space of cylindrical random variables, as detailed in Section 3.

Remark 2.7 (Relation to vector measures). We next show that (a) cylindrical integrals in the sense of Theorem 2.6 are integrals with respect to cylindrical vector measures and conversely that (b) cylindrical vector measures give rise to a cylindrical integral similar to Theorem 2.6. Moreover, (c) cylindrical vector measures can be identified with operator-valued vector measures. These relations are clearest in the case $I = \mathcal{L}^\infty(S, F)$, which shall be assumed here.

- (a) The space $\mathcal{L}^\infty(S, L(E, F))$ of non-cylindrical integrands is continuously included in the space $L(E, \mathcal{L}^\infty(S, F))$ of cylindrical integrands. Therefore, the cylindrical integral restricts to a continuous linear operator $\mathcal{L}^\infty(S, L(E, F)) \rightarrow L(E, H)$ or, equivalently, a continuous linear operator $E \rightarrow L(\mathcal{L}^\infty(S, L(E, F)), H)$. In other words, for any $e \in E$, the function $e \otimes \mu : \mathcal{A} \rightarrow E \hat{\otimes} G$ belongs to the space ba of finitely additive functions $\mathcal{A} \rightarrow E \hat{\otimes} G$ with finite semivariation $\|\cdot\|_{L(E, F), E \hat{\otimes} G, H}$, where $\hat{\otimes}$ is the projective tensor product. Consequently, μ (or rather the linear map $e \mapsto e \otimes \mu$) belongs to $L(E, ba)$. In this sense, μ is a cylindrical vector measure, and the cylindrical integral is a continuous extension of its elementary integral.
- (b) Of course, not all elements of $L(E, ba)$ are of the simple form $e \mapsto e \otimes \mu$. To treat the general case, let $\mu \in L(E, ba)$, where ba now denotes the set of finitely additive functions $\mathcal{A} \rightarrow G$ with finite semivariation $\|\cdot\|_{F, G, H}$. Under the mild assumption that for all $e \in E$, the elementary integral with respect to $\mu(e)$ extends continuously to $\mathcal{L}^\infty(S, F)$, one obtains a continuous integral $\mathcal{L}^\infty(S, F) \rightarrow L(E, H)$. Thus, starting from an E -cylindrical vector measure, one obtains an integral with values in the space $L(E, H)$, similarly to Theorem 2.6.
- (c) Cylindrical vector measures $\mu : E \rightarrow (\mathcal{A} \rightarrow G)$ in the sense of (b) can be identified with operator-valued vector measures $\nu : \mathcal{A} \rightarrow (E \rightarrow G)$ by a flip of coordinates: $\nu(A)(e) = \mu(e)(A)$. This identification is a Banach isometry. To see this, recall that the semivariation norm of a vector measure is the operator norm of its elementary integral and use the fact that $L(E, L(F, G)) \cong L(F, L(E, G))$ for any Banach spaces E , F , and G . Similar results for countably additive vector measures have been obtained in [4, Theorem 4.2] building on [28].

3. CYLINDRICAL STOCHASTIC ANALYSIS

This section treats some stochastic integrals together with their cylindrical counterparts and establishes well-posedness of cylindrical stochastic evolution equations under Lipschitz conditions on the coefficients. Throughout this section, $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ is a filtered probability space satisfying the usual conditions, $\Omega_T = [0, T] \times \Omega$, \mathcal{R} is the semi-ring of predictable rectangles in Ω_T , i.e.,

$$\mathcal{R} = \{\{0\} \times A : A \in \mathcal{F}_0\} \cup \{(s, t] \times A : A \in \mathcal{F}_s, 0 \leq s < t \leq T\},$$

\mathcal{A} is the algebra generated by \mathcal{R} , and \mathcal{P} is the sigma algebra generated by \mathcal{A} . As any set in \mathcal{A} is a finite union of sets in \mathcal{R} , the elementary functions over \mathcal{R} and \mathcal{A} coincide and are denoted simply by $\mathcal{L}_{\text{elem}}^\infty(\Omega_T, \mathbb{R})$. Unless specified otherwise, Ω_T carries the predictable sigma algebra \mathcal{P} . Subscripts \mathbb{F} in names of functions spaces denote adaptedness to the filtration \mathbb{F} , examples being $C_{\mathbb{F}}([0, T], L^2(\Omega, H))$ or $L_{\mathbb{F}}^2(\Omega, L^2([0, T], H))$. The space of Hilbert-Schmidt operators between Hilbert spaces U and H is denoted by $L_2(U, H)$.

Example 3.1 (Semimartingale integral). [9, 23, 29, 30] Let $p \in [1, \infty)$, let $X : [0, T] \rightarrow L^p(\Omega, \mathbb{R})$ be a stochastic process which is adapted and right-continuous in probability, and let $dX : \mathcal{R} \rightarrow L^p(\Omega, \mathbb{R})$ be given by

$$\begin{aligned} dX(\{0\} \times A) &= X_0 \mathbb{1}_A, & A \in \mathcal{F}_0, \\ dX((s, t] \times A) &= (X_t - X_s) \mathbb{1}_A, & A \in \mathcal{F}_s, 0 \leq s < t \leq T. \end{aligned}$$

Then dX extends uniquely to an additive function $dX : \mathcal{A} \rightarrow L^p(\Omega, \mathbb{R})$ because any set \mathcal{A} is a finite union of sets in \mathcal{R} . Moreover, the following are equivalent:

- (a) dX extends (uniquely) to a countably additive function $dX : \mathcal{P} \rightarrow L^p(\Omega, \mathbb{R})$.
- (b) The elementary integral

$$\mathcal{L}_{\text{elem}}^\infty(\Omega_T, \mathbb{R}) \ni f \mapsto \int_{\Omega_T} f dX \in L^p(\Omega, \mathbb{R})$$

is continuous and therefore has a unique continuous extension

$$\mathcal{L}^\infty(\Omega_T, \mathbb{R}) \ni f \mapsto \int_{\Omega_T} f dX \in L^p(\Omega, \mathbb{R}),$$

whose operator norm $\|dX\|_{\mathbb{R}, L^p(\Omega), L^p(\Omega)}$ is called L^p -Emery metric.

- (c) Up to a modification, $X = M + A$ for a p -integrable martingale M and a process A of p -integrable variation.

For any Banach space E , the corresponding cylindrified integral is of the form

$$L(E, \mathcal{L}^\infty(\Omega_T, \mathbb{R})) \ni f \mapsto \int_S f dX \in L(E, L^p(\Omega, \mathbb{R})).$$

Example 3.2 (Itô integral). [13, 31] Let H and U be separable Hilbert spaces, and let W be U -cylindrical Brownian motion, i.e., $W : L^2([0, T], U) \rightarrow L^2(\Omega, \mathbb{R})$ is a linear isometry with values in centered Gaussian random variables. Then, the elementary integral

$$\mathcal{L}_{\text{elem}}^\infty(\Omega_T, L_2(U, H)) \ni f \mapsto \int_{\Omega_T} f dW \in L^2(\Omega, H)$$

is well defined and extends to an isometry

$$L^2(\Omega_T, L_2(U, H)) = L_{\mathbb{F}}^2(\Omega, L^2([0, T], L_2(U, H))) \ni f \mapsto \int_{\Omega_T} f dW \in L^2(\Omega, H).$$

The equality above is due to the fact that $[0, T]$ carries an absolutely continuous measure [33]. More generally, if H is merely UMD Banach and $p \in (1, \infty)$, the elementary integral extends to an isomorphism

$$L_{\mathbb{F}}^p(\Omega, \gamma(L^2([0, T], U)), H) \ni f \mapsto \int_{\Omega_T} f dW \in L^p(\Omega, H),$$

where γ stands for γ -radonifying linear operators. Even more generally, if H is merely an abstract L^1 space, one retains a continuous (but not necessarily isomorphic) integral of the above form. However, beyond these settings, there are Banach spaces H such that the above extension of the elementary integral does not exist. Nevertheless, it always exists in the following cylindrical form: if E is a dual or pre-dual of H , the cylindrical integral is a well-defined linear isomorphism

$$L(E, L_{\mathbb{F}}^p(\Omega, L^2([0, T], U))) \ni f \mapsto \int_{\Omega_T} f dW \in L(E, L^p(\Omega, \mathbb{R})),$$

which generalizes the above integrals in the sense that $L^p(\Omega, H)$ is continuously included in $L(E, L^p(\Omega, \mathbb{R}))$.

Example 3.3 (Cylindrical stochastic convolutions). Let H and U be separable Hilbert spaces, let W be U -cylindrical Brownian motion, let E be a Banach space, and let $S : \mathbb{R}_{\geq 0} \rightarrow L(E)$ be a strongly continuous semigroup. For any $f \in L(E, L^2(\Omega_T, L_2(U, H)))$, the function

$$E \times [0, T] \ni (e, t) \mapsto ((s, \omega) \mapsto \mathbf{1}_{[0, t]}(s) f(S_{t-s}e)(s, \omega) \in L^2(\Omega_T, L_2(U, H)))$$

is continuous. Applying the Itô integral of Example 3.2 yields a continuous function

$$E \times [0, T] \ni (e, t) \mapsto \int_0^t f(S_{t-s}e)(s) dW_s \in L^2(\Omega, H).$$

The integral is continuous and adapted in $t \in [0, T]$ and continuous and linear in $e \in E$. Thus, writing $f(S_{t-s}e)$ as a pull-back $S_{t-s}^* f(e)$, one obtains a continuous linear operator

$$L(E, L^2(\Omega_T, L_2(U, H))) \ni f \mapsto \int_0^{\cdot} S_{-\cdot}^* f_s dW_s \in L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H))),$$

which shall be called cylindrical stochastic convolution. Similarly, starting from the pathwise Bochner integral instead of the Itô integral, one obtains a convolution

$$L(E, L^2(\Omega_T, H)) \ni f \mapsto \int_0^{\cdot} S_{-\cdot}^* f_s ds \in L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H))).$$

Theorem 3.4 (Cylindrical stochastic evolution equations). *Let U and H be separable Hilbert spaces, let W be U -cylindrical Brownian motion, let E be a Banach space, let $S : \mathbb{R}_{\geq 0} \rightarrow L(E)$ be a strongly continuous semigroup, let $X_0 \in L(E, L^2((\Omega, \mathcal{F}_0, \mathbb{P}), H))$, and let F and G be Lipschitz functions*

$$F : L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H))) \rightarrow L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H))),$$

$$G : L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H))) \rightarrow L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, L_2(U, H)))).$$

Then, there exists a unique $X \in L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H)))$ which satisfies

$$X_t = S_t^* X_0 + \int_0^t S_{t-s}^* F(X)_s ds + \int_0^t S_{t-s}^* G(X)_s dW_s,$$

for all $t \in [0, T]$, where the integrals are E -cylindrical, and S_{t-s}^ denotes the pull-back along S_{t-s} .*

Proof. We first verify that the right-hand side of the integral equation is well-defined. For any $X \in L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H)))$, the integrands are given by

$$F(X) \in L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H))), \quad G(X) \in L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, L_2(U, H)))).$$

Any continuous adapted function with values in $L^2(\Omega, H)$ or $L^2(\Omega, L_2(U, H))$ has a predictable version by Lemma A.2, giving rise to continuous inclusions

$$\begin{aligned} C_{\mathbb{F}}([0, T], L^2(\Omega, H)) &\rightarrow L^2(\Omega_T, H), \\ C_{\mathbb{F}}([0, T], L^2(\Omega, L_2(U, H))) &\rightarrow L^2(\Omega_T, L_2(U, H)). \end{aligned}$$

For later use, we note that the operator norm of these inclusions is \sqrt{T} . Via these inclusions, the integrands are given by

$$F(X) \in L(E, L^2(\Omega_T, H)), \quad G(X) \in L(E, L^2(\Omega_T, L_2(U, H))).$$

By an application of the cylindrical stochastic convolutions in Example 3.3, one obtains that the right-hand side of the integral equation,

$$K(X) := S^* X_0 + \int_0^\cdot S_{\cdot-s}^* F(X)_s ds + \int_0^\cdot S_{\cdot-s}^* G(X)_s dW_s$$

belongs to $L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H)))$. The function

$$K : L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H))) \rightarrow L(E, C_{\mathbb{F}}([0, T], L^2(\Omega, H)))$$

is Lipschitz continuous as a composition of Lipschitz functions. The Lipschitz constant of K is bounded by a constant times \sqrt{T} . Thus, for sufficiently small T , the operator K is a contraction and has a unique fixed point by the Banach fixed point theorem. For arbitrary T , one obtains a unique fixed point by concatenation of solutions on small sub-intervals of $[0, T]$. \square

4. CYLINDRICALLY MEASURE-VALUED TERM STRUCTURE MODELS

This section is devoted to cylindrical random measures and applications to financial term structure models. Throughout this section, $T^* \in \mathbb{R}_{>0}$, $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{t \in [0, T^*]}, \mathbb{P})$ is a filtered probability space satisfying the usual conditions, $\Omega_{T^*} = [0, T^*] \times \Omega$ with the predictable sigma algebra \mathcal{P} , W is U -cylindrical Brownian motion on a separable Hilbert space U , and $\mathcal{B}(\mathbb{R}_{>0})$ is the Borel sigma algebra on the positive half-line $\mathbb{R}_{>0} = (0, \infty)$, and $C_0(\mathbb{R}_{>0})$ is the Banach space of continuous functions $f : \mathbb{R}_{>0} \rightarrow \mathbb{R}$ with $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow \infty} f(x) = 0$. By the Riesz–Markov–Kakutani representation theorem, $C_0(\mathbb{R}_{>0})^*$ is the space $\mathcal{M}(\mathbb{R}_{>0})$ of signed measures $\mu : \mathcal{B}(\mathbb{R}_{>0}) \rightarrow \mathbb{R}$ with finite total variation $\|\mu\|_{C_0(\mathbb{R}_{>0})^*} = \|\mu\|_{\mathbb{R}, \mathbb{R}, \mathbb{R}}$. Similarly, for any Banach space E , $\mathcal{M}(\mathbb{R}_{>0}, E)$ denotes the set of countably additive vector measures $\mu : \mathcal{B}(\mathbb{R}_{>0}) \rightarrow E$ with finite semivariation $\|\mu\|_{\mathbb{R}, E, E}$.

Remark 4.1 (Measure-valued stochastic integration). The development of measure-valued stochastic integrals meets several difficulties. As the space $\mathcal{M}(\mathbb{R}_{>0})$ is not UMD, there is no Itô isometry in the sense of Example 3.2. However, $\mathcal{M}(\mathbb{R}_{>0})$ is an abstract L^1 space, i.e., a Banach lattice such that $\|x + y\| = \|x\| + \|y\|$ for any non-negative elements x and y whose infimum $x \wedge y$ vanishes. Thus, it is lattice isometric to some L^1 space [2, Theorem 4.27]. Therefore, a decoupling inequality holds [11, Corollary 3.21], which implies that the elementary Itô integral extends to a continuous linear map [11, Theorem 3.29]

$$L_{\mathbb{F}}^2(\Omega, \gamma(L^2([0, T], U), \mathcal{M}(\mathbb{R}_{>0}))) \ni f \mapsto \int_0^T f dW \in L^2(\Omega, \mathcal{M}(\mathbb{R}_{>0})),$$

where γ denotes the space of gamma-radonifying operators. This integral can be used to build measure-valued term structure models which are parameterized by time-of-maturity. However, it is ill-suited for parameterization by time-to-maturity,

as in Heath–Jarrow–Morton models, because measure-valued stochastic convolutions involving the shift semigroup are ill-defined. For example, if W is scalar Brownian motion, $\delta_0 \in \mathcal{M}(\mathbb{R}_{>0})$ is the Dirac measure, and S is the semigroup of left-shifts of measures, then the integral $\int_0^t S_{t-s}\delta_0 dW_s$ is almost never a measure because it is the distributional derivative of a Brownian path: indeed, for any $f \in C_c^\infty((0, t))$,

$$\int_0^t S_{t-s}\delta_0(f)dW_s = \int_0^t f(t-s)dW_s = \int_0^t f'(t-s)W_s ds.$$

Abstractly, the problem is that the shift semigroup on $\mathcal{M}(\mathbb{R}_{>0})$ is not strongly measurable and not Rademacher-bounded. Similar problems occur when the norm topology on $\mathcal{M}(\mathbb{R}_{>0})$ is replaced by the weak-* topology because the above stochastic integral is not a measure-valued random variable despite the integrand $s \mapsto S_{t-s}\delta_0$ being bounded predictable. The integral is, however, a cylindrical measure-valued random variable, i.e., an element of $L(C_0(\mathbb{R}_{>0}), L^2(\Omega))$. This is detailed in the sequel.

The following lemma (a) describes cylindrical random measures as a generalization of random measures and (b) explains their relation to $L^p(\Omega)$ -valued vector measures. An important consequence of (b) is that cylindrical vector measures can be integrated not only against C_0 functions, as suggested by their definition, but also against bounded Borel-measurable functions.

Lemma 4.2 (Cylindrical random measures). *(a) For any $p \in [1, \infty]$, there is a continuous inclusion*

$$L^p(\Omega, \mathcal{M}(\mathbb{R}_{>0})) \ni \mu \mapsto \left(f \mapsto \int f\mu \right) \in L(C_0(\mathbb{R}_{>0}), L^p(\Omega)).$$

(b) For any $p \in (1, \infty)$, there is an isometric isomorphism

$$\mathcal{M}(\mathbb{R}_{>0}, L^p(\Omega)) \ni \mu \mapsto \left(f \mapsto \int f\mu \right) \in L(C_0(\mathbb{R}_{>0}), L^p(\Omega)).$$

Proof. (a) Let i denote the function in (a). To show that i is injective, let $\mu \in L^p(\Omega, \mathcal{M}(\mathbb{R}_{>0}))$ with $i(\mu) = 0$. Then the support of μ is $\{0\}$ because for any non-zero $\nu \in \mathcal{M}(\mathbb{R}_{>0})$, there is $f \in C_0(\mathbb{R}_{>0})$ with $\int f\nu \neq 0$, and then $U_f = \{\tilde{\nu} \in \mathcal{M}(\mathbb{R}_{>0}) : \int f\tilde{\nu} \neq 0\}$ is an open neighborhood of ν which satisfies $\mathbb{P}[\mu \in U_f] = \mathbb{P}[\int f\mu \neq 0] = 0$. Therefore, μ vanishes, and i is injective. Furthermore, i is continuous thanks to the following inequality, which holds for all $\mu \in L^p(\Omega, \mathcal{M}(\mathbb{R}_{>0}))$ and $f \in C_0(\mathbb{R}_{>0})$:

$$\left\| \int f\mu \right\|_{L^p(\Omega)} \leq \| \|f\|_{C_0(\mathbb{R}_{>0})} \| \mu \|_{\mathcal{M}(\mathbb{R}_{>0})} \|_{L^p(\Omega)} = \|f\|_{C_0(\mathbb{R}_{>0})} \| \mu \|_{L^p(\Omega, \mathcal{M}(\mathbb{R}_{>0}))}.$$

(b) Let i denote the function in (b). Then i is well defined because the elementary integral of any $\mu \in \mathcal{M}(\mathbb{R}_{>0}, L^p(\Omega))$ extends isometrically to a continuous linear operator $\mathcal{L}^\infty(\mathbb{R}_{>0}) \rightarrow L^p(\Omega)$, as explained in Example 2.2, and then restricts to a continuous linear operator $i(\mu) : C_0(\mathbb{R}_{>0}) \rightarrow L^p(\Omega)$. The semivariation of μ is the operator norm of the elementary integral of μ and coincides with the operator norms of the above extension and restriction. Thus, i is an isometry. The surjectivity of i follows from a version of the Riesz representation theorem; cf. [22, Section VI.2]. Thanks to the reflexivity of $L^p(\Omega)$, the bi-dual of any $T \in L(C_0(\mathbb{R}_{>0}), L^p(\Omega))$ is an extension $T^{**} : C_0(\mathbb{R}_{>0})^{**} \rightarrow L^p(\Omega)$ of T . Let $B(\mathbb{R}_{>0})$ be the closure of $\mathcal{L}_{\text{elem}}^\infty(\mathbb{R}_{>0})$ in the uniform norm. As $B(\mathbb{R}_{>0})$ is isometrically included in $C_0(\mathbb{R}_{>0})^{**}$ and contains all indicators of Borel sets, one can define $\mu : \mathcal{B}(\mathbb{R}_{>0}) \ni A \mapsto T^{**}\mathbf{1}_A \in L^p(\Omega)$. As T^{**} is weak-* to weak-* continuous, μ is weakly countably additive and, by the Orlicz–Pettis

theorem [22, Corollary I.4.4], countably additive. Moreover, $i(\mu) = T$ because the two operators coincide on the dense set of elementary integrands. Thus, i is surjective. \square

Remark 4.3 (Countable additivity). Let $p \in (1, \infty)$. Then any cylindrical random measure $\mu \in L(C_0(\mathbb{R}_{>0}), L^p(\Omega))$ is ‘almost’ countably additive in the following sense: for any sequence of disjoint sets $A_i \in \mathcal{B}(\mathbb{R}_{>0})$,

$$\mu\left(\bigcup_i A_i\right) = \sum_i \mu(A_i)$$

up to a null set, which may depend on the sequence A_i . If μ is non-cylindrical, i.e., $\mu \in L^p(\Omega, \mathcal{M}(\mathbb{R}_{>0}))$, the null set can be chosen independently of the sequence A_i . This difference is minor in financial applications.

Cylindrically measure-valued processes can be constructed by solving cylindrical stochastic evolution equations, as shown next. The corresponding theory has been developed in Theorem 3.4. For financial applications, the two most important cases are where the semigroup acts via right-shifts on $C_0(\mathbb{R}_{>0})$ and where the semigroup acts trivially as the identity on $C_0(\mathbb{R}_{>0})$. Note that right-shifts on $C_0(\mathbb{R}_{>0})$ correspond to left-shifts on the dual space $\mathcal{M}(\mathbb{R}_{>0})$, in line with the Heath–Jarrow–Morton framework.

Corollary 4.4 (Cylindrically measure-valued evolution equations). *Let $S : \mathbb{R}_{\geq 0} \rightarrow L(C_0(\mathbb{R}_{>0}))$ be a strongly continuous semigroup, let $X_0 \in L(C_0(\mathbb{R}_{>0}), L^2(\Omega, \mathcal{F}_0, \mathbb{P}))$, and let F and G be Lipschitz functions*

$$F : L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega))) \rightarrow L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega))),$$

$$G : L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega))) \rightarrow L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega, U))).$$

Then, there exists a unique $X \in L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega)))$ which satisfies for all $t \in [0, T]$ that

$$X_t = S_t^* X_0 + \int_0^t S_{t-s}^* F(X)_s ds + \int_0^t S_{t-s}^* G(X)_s dW_s.$$

Proof. This is a special case of Theorem 3.4. \square

We next describe financial term structure models where the collection of forward rates is given by a cylindrically measure-valued process, as given by Corollary 4.4 above. Then, the prices of bonds or futures are obtained as usual by integration of maturities and exponentiation, i.e., by the application of cylindrical functionals. This yields a large financial market, whose tradeable assets are cylindrical functionals of a cylindrical process. Besides the minimal assumption that linear combinations of tradeable assets are tradeable, there is little extra structure imposed. In particular, it is not assumed that the tradeable assets constitute a non-cylindrical process, and this assumption would be lacking an economic foundation.

As a proof of concept, we present in Theorem 4.5 below a term structure model with two components: one for interest rates, parameterized by time-to-maturity, and another one for energy prices, parameterized by time-of-maturity. The motivation is that both interest rate and energy markets exhibit jumps at predictable times, which force the forward rates to be measure-valued. Moreover, the parameterization by time-to-maturity is well suited for dynamics depending foremost on market structure, as in the case of interest rates, whereas the parameterization by time-of-maturity is well suited for dynamics depending strongly on seasonalities, as in the case of energy markets. The additive structure of energy forwards corresponds to advance settlement of future contracts and is broken for continuous

or terminal settlement unless the interest rate is constant [6, Section 4.1]. We do not know how to model continuously or terminally settled contracts with stochastic interest rates in the present generality; cf. [7].

Theorem 4.5 (Cylindrical term structure models). *Let $S : \mathbb{R}_{\geq 0} \rightarrow L(C_0(\mathbb{R}_{>0}))$ be the semigroup of right shifts, let $F, \tilde{F} \in L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T^*], L^2(\Omega)))$, let $G, \tilde{G} \in L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T^*], L^2(\Omega, U)))$, and let $X, \tilde{X} \in L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T^*], L^2(\Omega)))$ satisfy for all $t \in [0, T^*]$ that*

$$\begin{aligned} X_t &= S_t^* X_0 + \int_0^t S_{t-s}^* F_s ds + \int_0^t S_{t-s}^* G_s dW_s, \\ \tilde{X}_t &= \tilde{X}_0 + \int_0^t \tilde{F}_s ds + \int_0^t \tilde{G}_s dW_s. \end{aligned}$$

Then the following statements hold, with operators on $C_0(\mathbb{R}_{>0})$ viewed as vector measures using the Riesz representation in Lemma 4.2 (b):

(a) For every $T, T_1, T_2 \in [0, T^*]$, the bond price and energy futures processes

$$P(t, T) = \exp(-X_t(0, T-t)), \quad \tilde{P}(t, T_1, T_2) = \tilde{X}_t(T_1, T_2), \quad t \in [0, T^*],$$

have a predictable version, which is unique up to $dt \otimes \mathbb{P}$ -null sets.

(b) The roll-over bank account process

$$B_t = \lim_{n \rightarrow \infty} \prod_{i=1}^n P(t(i-1)/n, ti/n)^{-1}, \quad t \in [0, T^*],$$

exists, has a predictable version, and is given by

$$B_t = \exp\left(X_0(0, t] + \int_0^t F_s(0, t-s] ds + \int_0^t G_s(0, t-s] dW_s\right), \quad t \in [0, T^*].$$

(c) For every $T, T_1, T_2 \in [0, T^*]$, the discounted bond price $P(t, T)/B_t$ and the discounted energy future $\tilde{P}(t, T_1, T_2)/B_t$ are local martingales if and only if the following drift conditions hold:

$$F_t(0, T-t] = \frac{1}{2} \|G_t(0, T-t]\|_U^2, \quad \tilde{F}_t(T_1, T_2) = 0, \quad dt \otimes \mathbb{P}\text{-almost surely.}$$

(d) If the drift condition (c) is valid for all $T, T_1, T_2 \in [0, T^*]$, then there is no asymptotic free lunch with vanishing risk under any probability measure equivalent to \mathbb{P} in the large financial market whose primary traded assets are the discounted bond prices $P(t, T)/B_t$ together with the discounted energy futures $\tilde{P}(t, T_1, T_2)/B_t$, for $T, T_1, T_2 \in [0, T^*]$.

Proof. (a) Thanks to the continuous linear inclusion [35]

$$C_{\mathbb{F}}([0, T^*], L^2(\Omega)) \rightarrow L^2(\Omega_{T^*}),$$

the process X is a continuous linear operator

$$X : C_0(\mathbb{R}_{>0}) \rightarrow L^2(\Omega_{T^*}).$$

Then X has a Riesz representation as a countably additive vector measure

$$X : \mathcal{B}(\mathbb{R}_{>0}) \rightarrow L^2(\Omega_{T^*}),$$

by the same argument as in Lemma 4.2 (b). Thanks to the countable additivity, the function

$$\mathbb{R} \ni x \mapsto X(0, x] \in L^2(\Omega_{T^*})$$

is right-continuous with left limits. By Lemma A.1, it has a measurable version

$$\mathbb{R} \times \Omega_{T^*} \ni (x, t, \omega) \mapsto X(0, x](t, \omega) \in \mathbb{R}.$$

As for any $T \in [0, T^*]$, the function

$$\Omega_{T^*} \ni (t, \omega) \mapsto (T - t, t, \omega) \in \mathbb{R} \times \Omega_{T^*}$$

is measurable, one obtains the measurability of the function

$$\Omega_{T^*} \ni (t, \omega) \mapsto X(0, T - t](t, \omega) \in \mathbb{R}.$$

This function is uniquely determined up to $dt \otimes \mathbb{P}$ -null sets because, for any \tilde{X} which originates from another choice of measurable version above,

$$\int_0^{T^*} \mathbb{E} \left[(X(0, T - t](t, \omega) - \tilde{X}(0, T - t](t, \omega))^2 \right] dt = 0.$$

Thus, by exponentiation, one obtains that the bond price process $P(\cdot, T)$ has a predictable version, which is determined uniquely up to a $dt \otimes \mathbb{P}$ -null set. A similar reasoning applies to the energy futures process $\tilde{P}(\cdot, T_1, T_2)$.

(b) For the purpose of the proof, let B_t be defined for any $t \in [0, T^*]$ as

$$B_t = \exp \left(X_0(0, t] + \int_0^t F_s(0, t - s] ds + \int_0^t G_s(0, t - s] dW_s \right).$$

These integrals are well-defined because for any $t \in [0, T^*]$, the integrands have predictable versions

$$\Omega_{T^*} \ni (s, \omega) \mapsto F(0, t - s](s, \omega) \in \mathbb{R},$$

$$\Omega_{T^*} \ni (s, \omega) \mapsto G(0, t - s](s, \omega) \in U,$$

which are uniquely determined up to $ds \otimes \mathbb{P}$ -null sets and square-integrable, as can be seen similarly to (a). Let $t \in [0, T^*]$, $n \in \mathbb{N}$, $t_i = ti/n$, and

$$B_t^n = \prod_{i=1}^n P(t_{i-1}, t_i)^{-1} = \exp \left(\sum_{i=1}^n X_{t_{i-1}}(0, t_i - t_{i-1}] \right).$$

Then, by the evolution equation for X ,

$$\begin{aligned} \log B_t^n &= \sum_{i=1}^n S_{t_{i-1}}^* X_0(0, t_i - t_{i-1}] + \sum_{i=1}^n \int_0^{t_{i-1}} S_{t_{i-1}-s}^* F_s(0, t_i - t_{i-1}] ds \\ &\quad + \sum_{i=1}^n \int_0^{t_{i-1}} S_{t_{i-1}-s}^* G_s(0, t_i - t_{i-1}] dW_s \\ &= X_0(0, t_n] + \sum_{i=1}^n \int_0^{t_{i-1}} F_s(t_{i-1} - s, t_i - s] ds \\ &\quad + \sum_{i=1}^n \int_0^{t_{i-1}} G_s(t_{i-1} - s, t_i - s] dW_s \end{aligned}$$

In terms of the round-up-to-the-grid operator

$$[s]_n = \min\{t_i : i \in \mathbb{N}, t_i \geq s\}, \quad s \in [0, t],$$

this can be rewritten by purely algebraic manipulations as

$$\log B_t^n = X_0(0, t] + \int_0^t F_s([s]_n - s, t - s] ds + \int_0^t G_s([s]_n - s, t - s] dW_s.$$

For large n , $[s]_n - s$ tends to 0 from above. As F and G are countably additive vector measures on $\mathcal{B}(\mathbb{R}_{>0})$, it follows that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|F_s([s]_n - s, t - s] - F_s(0, t - s]\|_{L^2(\Omega)} &= 0, \\ \lim_{n \rightarrow \infty} \|G_s([s]_n - s, t - s] - G_s(0, t - s]\|_{L^2(\Omega, U)} &= 0. \end{aligned}$$

Moreover, one has the uniform bound

$$\begin{aligned}\|F_s(\lceil s \rceil_n - s, t - s)\|_{L^2(\Omega)} &\leq \|F\|_{L(C_0(\mathbb{R}_{>0}), C_T([0, T^*], L^2(\Omega)))}, \\ \|G_s(\lceil s \rceil_n - s, t - s)\|_{L^2(\Omega, U)} &\leq \|G\|_{L(C_0(\mathbb{R}_{>0}), C_T([0, T^*], L^2(\Omega, U)))}.\end{aligned}$$

Thus, by the dominated convergence theorem,

$$\begin{aligned}\lim_{n \rightarrow \infty} \int_0^t \|F_s(\lceil s \rceil_n - s, t - s) - F_s(0, t - s)\|_{L^2(\Omega)}^2 ds &= 0, \\ \lim_{n \rightarrow \infty} \int_0^t \|G_s(\lceil s \rceil_n - s, t - s) - G_s(0, t - s)\|_{L^2(\Omega, U)}^2 ds &= 0.\end{aligned}$$

This ensures convergence of the integrals in the above formula for $\log B_t^n$. Thus, we have shown that B_t^n converges to B_t as n tends to infinity.

- (c) This follows along the lines of [26, p. 61ff]. Fix $0 \leq t \leq T \leq T^*$. By the evolution equation for X , one has

$$\begin{aligned}-\log P(t, T) &= X_t(0, T - t] \\ &= S_t X_0(0, T - t] + \int_0^t S_{t-s} F_s(0, T - t] ds + \int_0^t S_{t-s} G_s(0, T - t] dW_s \\ &= X_0(t, T] + \int_0^t F_s(t - s, T - s] ds + \int_0^t G_s(t - s, T - s] dW_s.\end{aligned}$$

By the integral formula for B_t , the discounted bond price $Z(t, T) = P(t, T)/B_t$ satisfies

$$-\log Z(t, T) = X_0(0, T] + \int_0^t F_s(0, T - s] ds + \int_0^t G_s(0, T - s] dW_s.$$

This is a semimartingale, and Itô's formula gives

$$\begin{aligned}Z(t, T) &= Z(0, T) - \int_0^t Z(s, T) \left(F_s(0, T - s] - \frac{1}{2} \|G_s(0, T - s]\|_{\mathcal{U}}^2 \right) ds \\ &\quad - \int_0^t Z(s, T) G_s(0, T - s] dW_s.\end{aligned}$$

Thus, $Z(\cdot, T)$ is a local martingale if and only if the first drift condition in statement (c) is satisfied. A similar but simpler argument establishes the second drift condition.

- (d) Under the probability measure \mathbb{P} , the value process of any admissible trading strategy in finitely many discounted bonds and discounted energy futures is a supermartingale, thanks to the Ansel–Stricker lemma. Therefore, \mathbb{P} is a separating measure for the large financial market with discounted bonds and discounted energy futures as primary traded assets. Consequently, no asymptotic free lunch with vanishing risk holds under \mathbb{P} and also under any measure equivalent to it [17]. □

Remark 4.6 (Bank account process). The bank account process in Theorem 4.5 (b) coincides with the usual one for ‘regular’ Heath–Jarrow–Morton models. Here, regularity refers to requirement that the forward rate measures X_t are non-cylindrical and have continuous Lebesgue densities f_t , which are subject to some additional regularity and integrability conditions [26, Conditions H1–H3 and C1–C4]. Under these conditions, the bank account process may equivalently be defined by the well-known formula $B_t = \exp(\int_0^t r_s ds)$, where $r_s = f_s(0)$ is the short rate. In particular,

the bank account process is a semimartingale. Moreover, by no-arbitrage considerations, the discounted bond prices, and consequently also the undiscounted ones, are semimartingales. In contrast, beyond the regular case, the bank account process and the undiscounted bond price processes may not be semimartingales. For instance, this happens if $X_t = S_t^* X_0$ is deterministic and X_0 has infinite variation.

Example 4.7 (Cylindrical term structure models). It is easy to specify coefficients F, \tilde{F} and G, \tilde{G} which satisfy the assumptions and drift conditions of Theorems 3.4 and 4.5, as demonstrated in the following simple example. Let $e \in C_0(\mathbb{R}_{>0})^n$, and let $g, \tilde{g} : \mathbb{R}^n \rightarrow \mathcal{N}_2(U, \mathcal{M}(\mathbb{R}_{>0}))$ be bounded Lipschitz functions, where \mathcal{N}_2 denotes 2-nuclear operators. Then, for any $X \in L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega)))$, one has $X_t(e) \in L^2(\Omega, \mathbb{R}^n)$ via the Riesz isomorphism in Lemma 4.2 (b) and $g(X_t(e)) \in L^\infty(\Omega, \mathcal{N}_2(U, \mathcal{M}(\mathbb{R}_{>0})))$ thanks to the boundedness of g . Thus, one obtains Lipschitz functions

$$G, \tilde{G} : L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega))) \rightarrow L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega, U)))$$

by identifying U with U^* and setting

$$G(X)(f)(t) = g(X_t(e))(f), \quad \tilde{G}(X)(f)(t) = \tilde{g}(X_t(e))(f).$$

Moreover, one obtains Lipschitz functions

$$F, \tilde{F} : L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega))) \rightarrow L(C_0(\mathbb{R}_{>0}), C_{\mathbb{F}}([0, T], L^2(\Omega))),$$

which satisfy the drift conditions of Theorem 4.5, by the following definition, where the product of a measure $\mu \in \mathcal{M}(\mathbb{R}_{>0})$ with its distribution function $\int \mu \in \mathcal{L}^\infty(\mathbb{R}_{>0})$ is denoted by $\mu \int \mu \in \mathcal{M}(\mathbb{R}_{>0})$, and where Tr denotes the trace on U :

$$F(X)(f)(t) = \frac{1}{2} \text{Tr}(g(X_t(e)) \int g(X_t(e))) (f), \quad \tilde{F}(X) = 0.$$

Alternative specifications are possible. For instance, the coefficients could be time-dependent and could depend simultaneously on X and \tilde{X} . Note that G , as specified here, is non-cylindrically measure-valued. We have not been able to construct F subject to the drift condition when G is merely cylindrically measure-valued. Nevertheless, even for non-cylindrical G , a cylindrical setting is needed because the solution X will be merely cylindrical, as noted in Remark 4.1.

Remark 4.8 (Measure-valued processes via martingale problems). A well-trodden path for constructing measure-valued processes is via martingale problems. Recently, this approach has been used to define models for discontinuous term structures [15], similarly to the ones defined here via cylindrical stochastic analysis. The two approaches are closely related, as described next. Let X be a mild solution of the stochastic evolution equation in Corollary 4.4 with Markovian coefficients $F(X)_s = F(X_s)$ and $G(X)_s = G(X_s)$, i.e.,

$$X_t = S_t^* X_0 + \int_0^t S_{t-s}^* F(X_s) ds + \int_0^t S_{t-s}^* G(X_s) dW_s.$$

Then, thanks to the (square) integrability of the coefficients, X is also a weak solution of the stochastic evolution equation [21, Theorem 6.5], i.e., letting A denote the generator of the shift semigroup S , one has for all $e \in D(A)$ that

$$X_t(e) = X_0(e) + \int_0^t X_s(Ae) ds + \int_0^t F(X_s)(e) ds + \int_0^t G(X_s)(e) dW_s.$$

Thus, for any cylindrical functional $f(x) = \varphi(x(e_1), \dots, x(e_n))$, where $\varphi \in C_b^\infty(\mathbb{R}^n, \mathbb{R})$, $n \in \mathbb{N}$, and $e_1, \dots, e_n \in D(A)$, the process

$$f(X_t) - f(X_0) - \int_0^t Af'(X_s)X_s ds - \int_0^t f'(X_s)F(X_s) ds$$

$$-\frac{1}{2} \int_0^t f''(X_s)(G(X_s), G(X_s)) ds$$

is a local martingale, where Af' is interpreted in the following sense:

$$Af'(x)y = g'(x(e_1), \dots, x(e_n))(y(Ae_1), \dots, y(Ae_n)).$$

Assuming that F and G map non-cylindrical random measures to non-cylindrical random measures, as for instance in Example 4.7, it follows that X solves the martingale problem with extended generator

$$Lf(x) = Af'(x)x + f'(x)F(x) + \frac{1}{2}f''(x)(G(x), G(x)), \quad x \in \mathcal{M}(\mathbb{R}_{>0}).$$

Under suitable conditions, existence and uniqueness of solutions to such martingale problems can be shown directly by Markovian methods. Notably, this allows one to construct non-cylindrical processes with values in the cone of non-negative measures. Of course, coefficients which lead to genuinely cylindrical processes, as for example in Remark 4.1, are excluded. Solutions of the martingale problem are, in great generality, again weak and mild solutions as above, for some Brownian motion reconstructed from the solution on an extended probability space [21, Theorem 8.2].

APPENDIX A. MEASURABLE AND PREDICTABLE VERSIONS

Lemma A.1 (Measurable version). *Let $T \in \mathbb{R}_+$, let (S, Σ, μ) be a measure space, let E be a Banach space, and let $D([0, T], L^2(S, E))$ be the Banach space of càdlàg functions $[0, T] \rightarrow L^2(S, E)$ with the supremum norm. Then, any process $X \in D([0, T], L^2(S, E))$ has a strongly measurable version $Y : [0, T] \times S \rightarrow E$, which is uniquely defined up to $dt \otimes \mu$ -null sets, thereby giving rise to a linear embedding with operator norm bounded by \sqrt{T} :*

$$D([0, T], L^2(S, E)) \ni X \mapsto Y \in L^2([0, T] \times S, E).$$

Proof. We follow the lines of [21, Proposition 3.2] with continuity replaced by right-continuity with left limits. Let $X \in D([0, T], L^2(S, E))$. For any $n \in \mathbb{N}$, there exists a finite set $A_n \subset [0, T]$, which contains zero and satisfies

$$\sup_{t \in [0, T]} \|X_t - X_{\lfloor t \rfloor_n}\|_{L^2(\Omega, E)} \leq 2^{-n},$$

where $\lfloor t \rfloor_n = \max(A_n \cap [0, t])$ denotes rounding down to the grid A_n . For any $t \in [0, T]$, let N_t be the set of all $s \in S$ such that $X_{\lfloor t \rfloor_n}(s)$ does not converge to $X_t(s)$. By Borell–Cantelli, N_t is a null set. The set $N = \bigcup_{t \in [0, T]} \{t\} \times N_t$ is measurable and a $dt \otimes \mu$ -null set. Define $Y : [0, T] \times \Omega \rightarrow E$ as $\lim_{n \rightarrow \infty} X_t(\lfloor s \rfloor_n)$ for $(t, s) \notin N$ and as zero otherwise. Then, Y is a version of X and is strongly measurable as a pointwise limit of strongly measurable functions. The L^2 -norm of Y is bounded by \sqrt{T} times the supremum norm of X . In particular, for any other strongly measurable version \tilde{Y} , the L^2 norm of $Y - \tilde{Y}$ vanishes, which implies that Y is determined uniquely up to $dt \otimes \mathbb{P}$ -null sets. \square

Lemma A.2 (Predictable version). *Let $T \in \mathbb{R}_+$, let $(\Omega, \mathcal{F}, \mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ be a filtered probability space satisfying the usual conditions, let E be a Banach space, let $C_{\mathbb{F}}([0, T], L^2(\Omega, E))$ be the Banach space of adapted processes in $C([0, T], L^2(\Omega, E))$, and let $\Omega_T = [0, T] \times \Omega$ be endowed with the measure $dt \otimes \mathbb{P}$ restricted to the predictable sigma algebra. Then, any process $X \in C_{\mathbb{F}}([0, T], L^2(\Omega, E))$ has a predictable version $Y : \Omega_T \rightarrow E$, which is uniquely defined up to $dt \otimes \mathbb{P}$ -null sets, thereby giving rise to a continuous linear embedding of operator norm bounded by \sqrt{T} :*

$$C_{\mathbb{F}}([0, T], L^2(\Omega)) \ni X \mapsto Y \in L^2(\Omega_T).$$

Proof. See [21, Proposition 3.7] or [35, Lemma 2.4] for the existence of a predictable version and [35, Lemma 2.4 and Proposition 2.5] for the continuous linear embedding. \square

REFERENCES

- [1] P. Harms and M. Müller. “Weak convergence rates for stochastic evolution equations and applications to nonlinear stochastic wave, HJMM, stochastic Schrödinger and linearized stochastic Korteweg-de Vries equations”. In: *Zeitschrift für Angewandte Mathematik und Physik* 70.16 (2019).
- [2] C. D. Aliprantis and O. Burkinshaw. *Positive operators*. Vol. 119. Springer, 2006.
- [3] D. Applebaum and M. Riedle. “Cylindrical Lévy processes in Banach spaces”. In: *Proc. Lond. Math. Soc. (3)* 101.3 (2010), pp. 697–726.
- [4] J. Assefa. “Cylindrical stochastic integration”. MA thesis. University of Freiburg, 2022.
- [5] R. Bartle. “A general bilinear vector integral”. In: *Studia Mathematica* 15.3 (1956), pp. 337–352.
- [6] F. E. Benth, J. S. Benth, and S. Koekebakker. *Stochastic modelling of electricity and related markets*. Vol. 11. World Scientific, 2008.
- [7] F. E. Benth, M. Piccirilli, and T. Vargiolu. “Mean-reverting additive energy forward curves in a Heath–Jarrow–Morton framework”. In: *Mathematics and Financial Economics* 13.4 (2019), pp. 543–577.
- [8] N. Berman and W. L. Root. “A weak stochastic integral in Banach space with application to a linear stochastic differential equation”. In: *Applied Mathematics and Optimization* 10.1 (1983), pp. 97–125.
- [9] K. Bichteler. *Stochastic integration with jumps*. Vol. 89. Cambridge University Press, 2002.
- [10] S. Bochner. “Integration von Funktionen, deren Werte die Elemente eines Vektorraumes sind”. In: *Fundamenta Mathematicae* 20.1 (1933), pp. 262–176.
- [11] S. Cox. “Stochastic differential equations in Banach spaces: Decoupling, Delay Equations, and Approximations in Space and Time”. PhD thesis. Technical University of Delft, 2012.
- [12] S. Cox, A. Jentzen, and F. Lindner. “Weak convergence rates for temporal numerical approximations of stochastic wave equations with multiplicative noise”. In: (2019). arXiv: 1901.05535.
- [13] S. Cox and M. Veraar. “Vector-valued decoupling and the Burkholder-Davis-Gundy inequality”. In: *Illinois J. Math.* 55.1 (2011), 343–375 (2012).
- [14] C. Cuchiero. “Polynomial processes in stochastic portfolio theory”. In: *Stochastic processes and their applications* 129.5 (2019), pp. 1829–1872.
- [15] C. Cuchiero, F. Guida, and L. Di Persio. “Measure-valued processes for energy markets”. In preparation. 2022.
- [16] C. Cuchiero, F. Guida, L. di Persio, and S. Svaluto-Ferro. “Measure-valued affine and polynomial diffusions”. In: *Preprint arXiv:2112.15129* (2021).
- [17] C. Cuchiero, I. Klein, and J. Teichmann. “A new perspective on the fundamental theorem of asset pricing for large financial markets”. In: *Theory of Probability & Its Applications* 60.4 (2016), pp. 561–579.
- [18] C. Cuchiero, M. Larsson, and S. Svaluto-Ferro. “Probability measure-valued polynomial diffusions”. In: *Electronic Journal of Probability* 24 (2019), pp. 1–32.
- [19] C. Cuchiero and S. Svaluto-Ferro. “Infinite-dimensional polynomial processes”. In: *Finance and Stochastics* 25.2 (2021), pp. 383–426.

- [20] C. Cuchiero and J. Teichmann. “Generalized Feller processes and Markovian lifts of stochastic Volterra processes: the affine case”. In: *Journal of evolution equations* 20.4 (2020), pp. 1301–1348.
- [21] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*. Cambridge University Press, 2014.
- [22] J. Diestel and J. Uhl. *Vector measures*. Mathematical surveys and monographs. American Mathematical Society, 1977.
- [23] N. Dinculeanu. *Vector integration and stochastic integration in Banach spaces*. Vol. 48. John Wiley & Sons, 2000.
- [24] S. Dolecki. “An initiation into convergence theory”. In: *Beyond Topology, Contemporary Mathematics* 486 (2009), pp. 115–161.
- [25] A. Etheridge. *An introduction to superprocesses*. 20. American Mathematical Soc., 2000.
- [26] D. Filipović. *Consistency problems for Heath–Jarrow–Morton interest rate models*. Vol. 1760. Lecture Notes in Mathematics. Springer Verlag, Berlin Heidelberg New York, 2001.
- [27] C. Fontana, Z. Grbac, S. Gümbel, and T. Schmidt. “Term structure modelling for multiple curves with stochastic discontinuities”. In: *Finance and Stochastics* 24.2 (2020), pp. 465–511.
- [28] W. H. Graves. *On the theory of vector measures*. Vol. 195. American Mathematical Soc., 1977.
- [29] S. Kwapien and W. Wołczyński. *Random Series and Stochastic Integrals: Single and Multiple*. Birkhäuser, 1992.
- [30] M. Métivier and J. Pellaumail. *Stochastic integration*. Academic Press, 1980.
- [31] J. M. A. M. van Neerven, M. C. Veraar, and L. Weis. “Stochastic integration in UMD Banach spaces”. In: *Ann. Probab.* 35.4 (2007), pp. 1438–1478.
- [32] M. Ondreját. “Brownian representations of cylindrical local martingales, martingale problem and strong Markov property of weak solutions of SPDEs in Banach spaces”. In: *Czechoslovak Math. J.* 55(130).4 (2005), pp. 1003–1039.
- [33] B. Rüdiger and S. Tappe. “Isomorphism for Spaces of Predictable Processes and an Extension of the Ito Integral”. In: *Stochastic analysis and applications* 30.3 (2012), pp. 529–537.
- [34] L. Schwartz. *Radon measures on arbitrary topological spaces and cylindrical measures*. Oxford University Press, 1973.
- [35] S. Tappe. “A note on stochastic integrals as L^2 -curves”. In: *Statistics & probability letters* 80.13-14 (2010), pp. 1141–1145.
- [36] A. Ustunel. “Stochastic analysis on nuclear spaces and its applications”. In: *Stochastic Space—Time Models and Limit Theorems*. Springer, 1985, pp. 163–177.
- [37] M. Veraar and I. Yaroslavtsev. “Cylindrical continuous martingales and stochastic integration in infinite dimensions”. In: *Electronic Journal of Probability* 21 (2016).
- [38] C. Ward and H. Unwin. *What caused recent spikes in UK wholesale power prices?* 2017. URL: <https://myice.ice.org.uk/news-and-insight/the-civil-engineer/february-2017/what-caused-the-recent-spike-in-power-prices> (visited on 07/2022).

DEPARTMENT OF STOCHASTICS, UNIVERSITY OF FREIBURG
 Email address: johannes.assefa@uranus.uni-freiburg.de

DIVISION OF MATHEMATICAL SCIENCES, NANYANG TECHNOLOGICAL UNIVERSITY, SINGAPORE
 Email address: philipp.harms@ntu.edu.sg