

# CONSTRUCTION OF STRONG DERIVABLE MAPS VIA FUNCTIONAL CALCULUS OF UNBOUNDED SPECTRAL OPERATORS IN BANACH SPACES.

BENEDETTO SILVESTRI

**ABSTRACT.** We provide sufficient conditions for the existence of a strong derivable map and calculate its derivative by employing a result in our previous work on strong derivability of maps arising by functional calculus of unbounded scalar type spectral operators in Banach spaces and the extension to complete locally convex spaces of a classical result valid in the Banach space context.

**Notation 1.** If  $X, Y$  are topological spaces let  $\mathcal{C}(X, Y)$  be the set of continuous maps from  $X$  to  $Y$ , if  $Z$  is a uniform space, then  $\mathcal{C}_u(X, Z)$  and  $\mathcal{C}_s(X, Z)$  denote the uniform space of continuous maps from  $X$  to  $Z$  provided by the uniformity of uniform convergence and simple convergence respectively.

In what follows we use the notation in [4]. We let  $G$  be a complex Banach space,  $R$  be a possibly unbounded scalar type operator in  $G$  and  $E$  its resolution of the identity. We let  $L(G)$  be the linear space of linear endomorphisms of the linear space underlying  $G$ . Let  $U \in \mathcal{B}(\mathbb{C})$  thus,  $\mathcal{Q}_E^\infty(U)$  is always considered provided by the locally convex topology induced by the seminorm  $\|\cdot\|_\infty^E$ .

**Definition 2.** Let  $F$  denote the locally convex space whose underlying linear space equals  $\{T \upharpoonright \text{Dom}(R) \mid T \in L(G)\}$  and whose locally convex topology is generated by the following set of seminorms  $\{F \ni P \mapsto \|Pv\| \mid v \in \text{Dom}(R)\}$ , let  $j_{B(G)}^F : B(G) \rightarrow F$  be the map  $T \mapsto T \upharpoonright \text{Dom}(R)$ . Let  $\widehat{F}$  denote the Hausdorff completion of  $F$  and  $i_F$  the canonical map of  $F$  into  $\widehat{F}$ .

Notice that  $j_{B(G)}^F$  and  $i_F$  are linear and continuous maps so uniformly continuous.

**Theorem 3.** Let  $K \subseteq \mathbb{R}$  be a compact nondegenerate interval of  $\mathbb{R}$ , and  $U$  an open neighbourhood of  $\sigma(R)$  such that  $K \cdot U \subseteq U$ . Furthermore let  $X$  be a compact space,  $g \in \mathcal{C}(X \times K, \mathcal{Q}_E^\infty(U))$ ,  $\{x_n\}_{n \in \mathbb{N}}$  be a sequence in  $X$ ,  $t_0 \in K$  and  $\{f^n\}_{n \in \mathbb{N}}$  be a sequence of analytic maps defined on  $U$  with the following properties:  $\sup_{t \in K} \|\widetilde{f}_t^n\|_\infty^E < \infty$  for every  $n \in \mathbb{N}$ ,

$$(1) \quad (\forall v \in \text{Dom}(R))(\exists \lim_{n \in \mathbb{N}} f^n(t_0 R)v),$$

Date: August 9, 2022.

2010 Mathematics Subject Classification. 46G05, 47B40, 47A60.

Key words and phrases. strong operator derivability, unbounded spectral operators in Banach spaces, functional calculus.

and for every  $n \in \mathbb{N}$  and  $t \in K$  we have

$$(2) \quad g(x_n, t) = \left( \frac{df^n}{d\lambda} \right)_t.$$

Thus, there exist  $y \in X$  and a subsequence  $\{x_{n_k}\}_{k \in \mathbb{N}}$  of  $\{x_n\}_{n \in \mathbb{N}}$  converging to  $y$  such that for every open neighbourhood  $W$  of  $t_0$  for which  $W \subset K$  we have what follows: There exists a derivable map  $h \in \mathcal{C}(W, \widehat{F})$  whose derivative satisfies for every  $t \in W$

$$(Dh)(t) = i_F(g(y, t)(R)R),$$

while

$$h = \lim_{k \in \mathbb{N}} (W \ni t \mapsto (i_F \circ j_{B(G)}^F)(f^{n_k}(tR))) \text{ in } \mathcal{C}_u(W, \widehat{F}).$$

*Proof.* Let  $n \in \mathbb{N}$  and let  $W$  be an open interval of  $\mathbb{R}$  such that  $t_0 \in W$  and  $W \subset K$ . We have  $\sup_{t \in K} \|g(x_n, t)\|_\infty^E < \infty$  since  $\|\cdot\|_\infty^E \circ g$  is continuous and  $\{x_n\} \times K$  is compact. So we can apply [4, theorem 1.23] and state that for every  $v \in \text{Dom}(R)$  the map  $W \ni t \mapsto f^n(tR)v \in G$  is derivable and for every  $t \in W$

$$(3) \quad \frac{df^n(tR)v}{dt} = R \frac{df^n}{d\lambda}(tR)v.$$

Now by letting  $\iota : \mathbb{C} \rightarrow \mathbb{C}$  be the map  $\lambda \rightarrow \lambda$  clearly we have  $R = \iota(R)$  so

$$\begin{aligned} R \frac{df^n}{d\lambda}(tR) &= R \left( \frac{df^n}{d\lambda} \right)_t (R) \\ &= \iota(R) \left( \frac{df^n}{d\lambda} \right)_t (R) \\ &\subseteq \left( \iota \cdot \left( \frac{df^n}{d\lambda} \right)_t \right) (R) \\ &= \left( \left( \frac{df^n}{d\lambda} \right)_t \cdot \iota \right) (R); \end{aligned}$$

where the inclusion follows by the spectral theorem [3, 18.2.11(f)] which implies also that

$$\left( \frac{df^n}{d\lambda} \right)_t (R) \iota(R) \subseteq \left( \left( \frac{df^n}{d\lambda} \right)_t \cdot \iota \right) (R)$$

while clearly

$$\text{Dom} \left( \left( \frac{df^n}{d\lambda} \right)_t (R) \iota(R) \right) = \text{Dom}(R);$$

so, for every  $v \in \text{Dom}(R)$  we deduce that

$$R \frac{df^n}{d\lambda}(tR)v = \left( \frac{df^n}{d\lambda} \right)_t (R)Rv.$$

Therefore by (3) and hypothesis (2) we obtain for every  $v \in \text{Dom}(R)$  and  $t \in W$

$$(4) \quad \frac{df^n(tR)v}{dt} = g(x_n, t)(R)Rv,$$

which is equivalent to state that for every  $t \in W$

$$\frac{dj_{B(G)}^F(f^n(tR))}{dt} = g(x_n, t)(R)R.$$

Next  $g$  is uniformly continuous being a continuous map defined on a compact set and valued in a uniform space thus, by [1, X.13 Prp.2] we deduce for every  $z \in X$  that  $Q_z := \{g(z, \cdot)\} \cup \{g(x_n, \cdot) \mid n \in \mathbb{N}\}$  is an equicontinuous subset of  $\mathcal{C}(K, \mathcal{Q}_E^\infty(U))$  therefore, by [1, X.16 Thm.1] on  $Q_z$  the uniform structure inherited by that of  $\mathcal{C}_u(K, \mathcal{Q}_E^\infty(U))$  equals the uniform structure inherited by that of  $\mathcal{C}_s(K, \mathcal{Q}_E^\infty(U))$ . Now let  $\{x_{n_k}\}_{k \in \mathbb{N}}$  be a subsequence of  $\{x_n\}_{n \in \mathbb{N}}$  converging to a point  $y \in X$  which exists since  $X$  is compact. Thus, the sequence  $\{g(x_{n_k}, \cdot)\}_{k \in \mathbb{N}}$  converges to  $g(y, \cdot)$  in  $\mathcal{C}_s(K, \mathcal{Q}_E^\infty(U))$  due to the continuity of  $g$  therefore, for what above established applied to  $z = y$  we conclude that

$$(5) \quad g(y, \cdot) = \lim_{k \in \mathbb{N}} g(x_{n_k}, \cdot) \text{ in } \mathcal{C}_u(K, \mathcal{Q}_E^\infty(U)).$$

Next the functional calculus restricted at  $\mathcal{Q}_E^\infty(U)$  say  $I$  is a linear and continuous map at values in the Banach space  $B(G)$  since [3, 18.2.11(c)] so, uniformly continuous, moreover for every  $v \in \text{Dom}(R)$  the map  $\hat{v} : B(G) \rightarrow G$  defined by  $T \mapsto TRv$  is linear and continuous so uniformly continuous, therefore,  $\hat{v} \circ I : \mathcal{Q}_E^\infty(U) \rightarrow G$  is uniformly continuous being composition of uniformly continuous maps so, we deduce by (5) and by [1, X.5 Proposition 3(a)] applied to the uniformly continuous map  $\hat{v} \circ I$  that for every  $v \in \text{Dom}(R)$

$$g(y, \cdot)(R)Rv = \lim_{k \in \mathbb{N}} g(x_{n_k}, \cdot)(R)Rv \text{ in } \mathcal{C}_u(K, G),$$

and then (4) yields for every  $v \in \text{Dom}(R)$

$$\lim_{k \in \mathbb{N}} \left( W \ni t \mapsto \frac{df^{n_k}(tR)v}{dt} \right) = (W \ni t \mapsto g(y, t)(R)Rv) \text{ in } \mathcal{C}_u(W, G),$$

which is equivalent to state that

$$\lim_{k \in \mathbb{N}} \left( W \ni t \mapsto \frac{dj_{B(G)}^F(f^{n_k}(tR))}{dt} \right) = (W \ni t \mapsto g(y, t)(R)R) \text{ in } \mathcal{C}_u(W, F),$$

therefore in sequence by [1, X.5 Proposition 3(a)] and the fact that the derivative operator commutes with the operator  $f \mapsto u \circ f$  where  $u$  is a linear continuous operator between Hausdorff locally convex spaces, both applied to the map  $i_F$  we conclude that

$$(6) \quad \lim_{k \in \mathbb{N}} \left( W \ni t \mapsto \frac{d(i_F \circ j_{B(G)}^F)(f^{n_k}(tR))}{dt} \right) = (W \ni t \mapsto i_F(g(y, t)(R)R)) \text{ in } \mathcal{C}_u(W, \widehat{F}).$$

Next (1) is equivalent to state that  $\exists \lim_{n \in \mathbb{N}} j_{B(G)}^F(f^n(t_0 R))$  in  $F$  which then implies

$$(7) \quad \exists \lim_{n \in \mathbb{N}} (i_F \circ j_{B(G)}^F)(f^n(t_0 R)) \text{ in } \widehat{F}.$$

The statement then follows by (6), (7) and the extension of the result [2, Cor.8.6.4] to complete locally convex linear space valued maps defined on open sets of  $\mathbb{R}$ .  $\square$

#### REFERENCES

1. Bourbaki, N. *Topologie Generale*, CCLS, Ch 5-10 1974.
2. Dieudonne, J *Foundations of Modern Analysis* Academic Press 1969
3. Dunford; Schwartz *Linear Operators Part III* Wiley Interscience Publ. 1971
4. Silvestri, B. *Integral equalities for functions of unbounded spectral operators in Banach spaces* Dissertationes Math. 464 (2009), 60 pp.