

THE HOMOGENEOUS SPECTRUM OF A \mathbb{Z}_2 -GRADED COMMUTATIVE RING

MOHAMED AQALMOUN

ABSTRACT. Let $\mathbb{Z}_2 := \mathbb{Z}/2\mathbb{Z}$ be the additive group with two elements. In this article, we focus only on \mathbb{Z}_2 -graded commutative ring i.e commutative ring R such that $R = R_0 \oplus R_1$ as Abelian group and $R_i R_j \subseteq R_{i+j}$ for all $i, j \in \mathbb{Z}_2$. Our main goals is to establish a strong relation between \mathbb{Z}_2 -graded prime (maximal) ideals of R and prime (maximal) ideals of R_0 , for instance, it is showed that, the \mathbb{Z}_2 -graded spectrum of R is homeomorphic to the spectrum of R_0 with respect to the Zariski topologies.

Keywords: \mathbb{Z}_2 -graded ring, \mathbb{Z}_2 -graded prime ideal, \mathbb{Z}_2 -graded maximal ideal, \mathbb{Z}_2 -graded field.

MSC(2010): Primary: 13A02 ; Secondary: 13A99.

1. Introduction

Let G be an Abelian group with identity e and R be a commutative ring with unit. Then R is called a G -graded ring if there exist additive subgroups R_g of R indexed by elements $g \in G$ such that $R = \bigoplus_{g \in G} R_g$ and $R_g R_{g'} \subseteq R_{gg'}$ for all $g, g' \in G$, where $R_g R_{g'}$ consists of the finite sums of ring products ab with $a \in R_g$ and $b \in R_{g'}$. The elements of R_g are called homogeneous elements of R of degree g . The homogeneous elements of the ring R are denoted by $h(R)$, i.e. $h(R) = \cup_{g \in G} R_g$. If $a \in R$, then the element a can be written uniquely as $\sum_{g \in G} a_g$, where $a_g \in R_g$ is called the g -component of a in R_g .

Let R be a G -graded ring and I be an ideal of R . Then I is called a graded ideal of R if $I = \bigoplus_{g \in G} (I \cap R_g)$. If I is a G -graded ideal of R , then the quotient ring R/I is a G -graded ring. Indeed $R/I = \bigoplus_{g \in G} (R/I)_g$ where $(R/I)_g = (R_g + I)/I = \{x + I / x \in R_g\}$. Let $S \subseteq h(R)$ be a multiplicatively closed subset of R . Then the ring of fractions $S^{-1}R$ is a G -graded ring. Indeed $S^{-1}R = \bigoplus_{g \in G} (S^{-1}R)_g$ where $(S^{-1}R)_g = \{r/s / r \in h(R), s \in S \text{ and } \deg r = g \deg s\}$.

A G -graded ideal P of a G -graded ring R is called G -graded prime ideal of R if $P \neq R$ and if whenever r and s are homogeneous elements of R such that $rs \in P$, then either $r \in P$ or $s \in P$. The G -graded spectrum or homogeneous spectrum of R is the set of all G -graded

prime ideals of R , it is denoted by $G\text{Spec}R$. A graded ideal M of R is said to be graded maximal ideal of R if $M \neq R$ and if J is a graded ideal of R such that $M \subseteq J \subseteq R$, then $J = M$ or $J = R$, the set of all G -graded maximal ideals of R is denoted by $G\text{Max}R$.

For a commutative ring R and ideal I of R , the variety of I is the subset $V(I) := \{P \in \text{Spec}R \mid I \subseteq P\}$. Then the collection $\{V(I) \mid I \text{ ideal of } R\}$ satisfies the axioms for the closed sets of a topology on $\text{Spec}R$, called the Zariski topology. Note that for each ideal I , the set $\text{Spec}R - V(I)$ is an open subset, it is denoted $D(I)$. Recall that, the collection $D(f)$ where f runs through R forms a basis of opens. Similarly, let R be a G -graded commutative ring and I a G -graded ideal of R , the G -variety of I is the subset $V_G(I) := \{P \in G\text{Spec}R \mid I \subseteq P\}$. The collection $\{V_G(I) \mid I \text{ is a } G\text{-graded ideal of } R\}$ satisfies the axioms for the closed sets of a topology on $G\text{Spec}R$, called the Zariski topology, see [7], [10], [11]. Note that, the collection $D(r)$ where r runs through $h(R)$ forms a basis of opens for the Zariski topology.

Let $G = \mathbb{Z}_2 = \mathbb{Z}/2\mathbb{Z}$ and R be \mathbb{Z}_2 -graded commutative ring. Then $R = R_0 \oplus R_1$, where R_0 is a subring of R and R_1 is an R_0 -module such that $R_1R_1 \subseteq R_0$. Note that R_1R_1 is an ideal of R_0 , we denote it by R_1^2 . Similarly R_1^3 is an R_0 -submodule of R_1 . If P is a \mathbb{Z}_2 -graded prime ideal of R , then $P \cap R_0$ is a prime ideal of R_0 . This yields a natural map $\mathbb{Z}_2\text{Spec}R \rightarrow \text{Spec}R_0$.

Example 1.1. Let A be a commutative ring, then the polynomial ring $R = A[X]$ is \mathbb{Z}_2 -graded ring with the grading $R_0 = \bigoplus_n A(X^{2n})$ and $R_1 = \bigoplus_n A(X^{2n+1})$. In fact every \mathbb{N} -graded (respectively \mathbb{Z} -graded) commutative ring R has a natural \mathbb{Z}_2 -grading with $R'_0 = \bigoplus_n R_{2n}$ and $R'_1 = \bigoplus_n R_{2n+1}$.

Example 1.2. Let A be a commutative ring and $X = (X_\alpha)_\alpha$ be a set of indeterminate over A and $(a_{\alpha\beta})_{\alpha,\beta}$ be a family of elements of A . Consider the ideal I generated by all elements $X_\alpha X_\beta - a_{\alpha\beta}$. Then $R := \frac{A[X]}{I}$ is a \mathbb{Z}_2 -graded ring with the grading $R_0 = A$ and R_1 is the A -submodule of R generated by all $X_\alpha + I$.

Remark 1.3. Matrix ring representation of a \mathbb{Z}_2 -graded commutative ring. Let R be a \mathbb{Z}_2 -graded commutative ring. If $r = r_0 + r_1, s = s_0 + s_1 \in R = R_0 \oplus R_1$, then $rs = (r_0s_0 + r_1s_1) + (r_0s_1 + s_0r_1)$, that is $(rs)_0 = r_0s_0 + r_1s_1$ and $(rs)_1 = r_0s_1 + s_0r_1$. So, one can identifies R with the matrix ring

$$R' = \left\{ \begin{pmatrix} r_0 & r_1 \\ r_1 & r_0 \end{pmatrix} \mid r_0 \in R_0, r_1 \in R_1 \right\}$$

The purpose of this paper is to establish a strong relation between \mathbb{Z}_2 -graded prime ideals of a \mathbb{Z}_2 -graded commutative ring R and prime ideals of R_0 via the natural map $\mathbb{Z}_2\text{Spec}R \rightarrow \text{Spec}R_0$. It is proved that, this map is one to one, and in fact it is a homeomorphism with respect to the Zariski topologies. In fact, to fulfill this goal, we give the form of a \mathbb{Z}_2 -graded prime ideal P from its degree zero homogeneous part $P \cap R_0$, their expression lies on the fact that $P \cap R_0 \in V(R_1^2)$ or $P \cap R_0 \in D(R_1^2)$. Precisely, every \mathbb{Z}_2 -graded ideal can be discovered from its degree zero homogeneous part.

2. \mathbb{Z}_2 -Graded ideals

We start this section with some useful properties for \mathbb{Z}_2 -graded ideals of a \mathbb{Z}_2 -graded commutative ring. We will see that, when R is a strongly \mathbb{Z}_2 -graded commutative ring, those ideals are no thing but as ideals of R_0 .

Proposition 2.1. *Let R be a \mathbb{Z}_2 -graded ring and J be an ideal of R . The following statements are equivalent.*

- (1) J is a \mathbb{Z}_2 -graded ideal of R .
- (2) $J = I + R'$ where I is an ideal of R_0 and R' is an R_0 -submodule of R_1 with $IR_1 \subseteq R'$ and $R_1R' \subseteq I$.

Proof. Let J be a \mathbb{Z}_2 -graded ideal of R . Then $J = I + R'$ where $I := J \cap R_0$ is an ideal of R_0 and $R' := J \cap R_1$ is an R_0 -submodule of R_1 . If $x \in R_1$ and $y \in R'$, then $xy \in J$ is an element of degree 0, so $xy \in J \cap R_0 = I$. It follows that $R_1R' \subseteq I$. If $i \in I$ and $y \in R_1$, then $iy \in J$ is an element of degree 1, so $iy \in J \cap R_1 = R'$. Therefore $IR_1 \subseteq R'$.

Let $J = I + R'$ where I is an ideal of R_0 and R' is an R_0 -submodule of R_1 with $IR_1 \subseteq R'$ and $R_1R' \subseteq I$. Clearly J is a subgroup of R . Let $x = i + r' \in I + R'$ and $y = a + r \in R = R_0 + R_1$. Then $xy = (ia + r'r) + (ir + ar') \in I + R' = J$. \square

If R' is an R_0 -submodule of R_1 . Then the residual R' by R_1 is defined as follow

$$(R' :_{R_0} R_1) = \{a \in R_0 / aR_1 \subseteq R'\}$$

It is an ideal of R_0 . So, the condition $IR_1 \subseteq R'$ is equivalent to $I \subseteq (R' :_{R_0} R_1)$.

Remark 2.2. As a consequence, we have the two classes of \mathbb{Z}_2 -graded ideals, which will appear for our discussion.

- (1) If I is an ideal of R_0 , then $I + IR_1$ is a \mathbb{Z}_2 -graded ideal of R .
Indeed, $IR_1 \subseteq IR_1$ and $R_1(IR_1) = IR_1R_1 \subseteq IR_0 = I$.

- (2) If R' is a submodule of R_1 , then $(R' :_{R_0} R_1) + R'$ is a \mathbb{Z}_2 -graded ideal of R . Indeed; $(R' :_{R_0} R_1)R_1 \subseteq R'$ and $(R'R_1)R_1 \subseteq R'R_0 \subseteq R'$ that is $R'R_1 \subseteq (R' :_{R_0} R_1)$.

Definition 2.3. A G -graded commutative ring is called strongly graded if $R_g R_{g'} = R_{gg'}$ for all $g, g' \in G$.

Note that, if R is a \mathbb{Z}_2 -graded commutative ring, then R is strongly graded if and only if $R_1^2 = R_0$.

The following result establish a relation between graded ideals of a strongly \mathbb{Z}_2 -graded commutative ring R and ideals of R_0 .

Proposition 2.4. *Let R be a strongly \mathbb{Z}_2 -graded commutative ring. The following statements hold.*

- (1) R_1 is a finitely generated R_0 -module.
- (2) R_1 is a multiplication module.
- (3) If I and I' are ideals of R_0 such that $IR_1 \subseteq I'R_1$, then $I \subseteq I'$.
- (4) The \mathbb{Z}_2 -graded ideals of R are $I + IR_1$ where I is an ideal of R_0 .

Proof. (1) Since $R_1^2 = R_0$, there exist $a_i, b_i \in R_1$ such that $1 =$

$$\sum_{i=1}^n a_i b_i. \text{ Then } R_1 = (a_1, \dots, a_n). \text{ Indeed, if } x \in R_1, \text{ then}$$

$$x = \sum_{i=1}^n (x b_i) a_i. \text{ The elements } x b_i \text{ are of degree zero, so they are}$$

in R_0 . Therefore $R_1 = (a_1, \dots, a_n)$.

- (2) Let R' be an R_0 -submodule of R_1 . Clearly, $(R' :_{R_0} R_1)R_1 \subseteq R'$.

Let $x \in R'$, then $x = \sum_{i=1}^n (x b_i) a_i$. Each $x b_i$ is an element of

$(R' :_{R_0} R_1)$, in fact if $c \in R_1$, then $(x b_i) c = (b_i c) x \in R'$ since $b_i c \in R_0$ and $x \in R'$. Thus $R' = (R' :_{R_0} R_1)R_1$. Therefore R_1 is a multiplication module.

- (3) If $IR_1 \subseteq I'R_1$, then $IR_1^2 \subseteq I'R_1^2$, so $I \subseteq I'$.

- (4) Let J be a \mathbb{Z}_2 -graded ideal of R , then $J = I + R'$ where I is an ideal of R_0 and R' is an R_0 -submodule of R_1 such that $IR_1 \subseteq R'$ and $R'R_1 \subseteq I$. Since R_1 is a multiplication module, there is an ideal I' of R_0 such that $R' = I'R_1$. But $IR_1 \subseteq R' = I'R_1$, so $I \subseteq I'$. On other hand, $I' = I'R_1^2 = R'R_1 \subseteq I$. Thus $I = I'$ and $J = I + IR_1$. Conversely, if $J = I + IR_1$ where I is an ideal of R_0 , then J is a \mathbb{Z}_2 -graded ideal (see the Remark 2.2). □

Remark 2.5. Let R be a strongly \mathbb{Z}_2 -graded ring. The map $J \mapsto J \cap R_0$ realize a bijection between the set of \mathbb{Z}_2 -graded ideals of R and the set of ideals of R_0 .

One can show that, for a strongly \mathbb{Z}_2 -graded commutative ring, the \mathbb{Z}_2 -graded ideals of R are $(R' :_{R_0} R_1) + R'$ where R' is an R_0 -submodule of R_1 . It follows that the map $J \mapsto J \cap R_1$ realize a bijection between the set of \mathbb{Z}_2 -graded ideals of R and the set of R_0 -submodule R_1 .

3. \mathbb{Z}_2 -Graded prime ideals

Definition 3.1. Let R be a G -graded commutative ring and P be a G -graded ideal of R . Then P is called G -graded prime ideal (or homogeneous prime ideal) of R if $P \neq R$ and if whenever r and s are homogeneous elements of R such that $rs \in P$, then either $r \in P$ or $s \in P$

Remark 3.2. A G -graded prime ideal of a G -graded ring R is not necessarily a prime ideal. Let $R = \mathbb{Z} + i\mathbb{Z}$ (where $i^2 = -1$) as a \mathbb{Z}_2 -graded ring with $R_0 = \mathbb{Z}$ and $R_1 = i\mathbb{Z}$. It is easy to see that $P := 2\mathbb{Z} + i2\mathbb{Z}$ is a \mathbb{Z}_2 -graded prime ideal of R . But P is not a prime ideal of R , since $(3+i)(3-i) = 10 \in P$ and $3+i, 3-i \notin P$.

Recall from [3] that if A is a commutative ring and M is an A -module, then a prime submodule of M is a proper submodule N of M having the property that $am \in N$ implies that $a \in (N :_S M)$ or $m \in N$ for each $a \in S$ and $m \in M$.

The following Theorem describes the \mathbb{Z}_2 -graded prime ideals of R from prime ideals of R_0 and prime submodules of R_1 .

Theorem 3.3. *Let Q be a \mathbb{Z}_2 -graded ideal of R . Then Q is a \mathbb{Z}_2 -graded prime ideal of R if and only if*

- (1) *either $Q = p + R_1$ where $p \in V(R_0^2)$, or*
- (2) *$Q = (R' :_{R_0} R_1) + R'$ where R' is a prime submodule of R_1 such that $R_1^3 \not\subseteq R'$.*

Proof. (\Leftarrow): Let p be a prime ideal of R_0 containing R_1^2 . Then $p + R_1$ is \mathbb{Z}_2 -graded ideal of R and is a prime ideal of R since $R/(p + R_1) = R_0/p$ is an integral domain, so it is a \mathbb{Z}_2 -graded prime ideal of R .

Let R' be prime submodule of R_1 such that $R_1^3 \not\subseteq R'$. Then $(R' :_{R_0} R_1) + R'$ is a graded ideal of R . Let $a, b \in R_0$ such that $ab \in (R' :_{R_0} R_1) + R'$, then $ab \in (R' :_{R_0} R_1)$. Since R' is a prime submodule of R_1 , $(R' :_{R_0} R_1)$ is a prime ideal of R_0 . Thus $a \in (R' :_{R_0} R_1)$ or $b \in (R' :_{R_0} R_1)$. If $a \in R_0$ and $r \in R_1$ such that $ar \in (R' :_{R_0} R_1) + R'$, then $ar \in R'$. It follows that $a \in (R' :_{R_0} R_1)$ or $r \in R'$ since R' is a prime submodule

of R_1 . Now, let $r, r' \in R_1$ such that $rr' \in (R' :_{R_0} R_1) + R'$. Assume that $r, r' \notin R'$. We have $rr' \in (R' :_{R_0} R_1)$, so $rr'R_1^2 \subseteq (R' :_{R_0} R_1)$, that is $(rR_1)(r'R_1) \subseteq (R' :_{R_0} R_1)$. Since rR_1 and $r'R_1$ are ideals of R_0 and $(R' :_{R_0} R_1)$ is a prime ideal of R_0 , it follows that $rR_1 \subseteq (R' :_{R_0} R_1)$ or $r'R_1 \subseteq (R' :_{R_0} R_1)$, that is $rR_1^2 \subseteq R'$ or $r'R_1^2 \subseteq R'$. Since $r, r' \notin R'$ and R' is a prime submodule, we get $R_1^2 \subseteq (R' :_{R_0} R_1)$, so that $R_1^3 \subseteq R'$, a contradiction. Thus $r \in R'$ or $r' \in R'$.

(\Rightarrow): Let Q be a graded prime ideal of R . Then, by Proposition 2.1, $Q = p + R'$ where p is an ideal of R_0 and R' is a submodule of R_1 such that $pR_1 \subseteq R'$ and $R'R_1 \subseteq p$. It is easy to see that p is a prime ideal of R_0 . Indeed, if $a, b \in R_0$ such that $ab \in p$, then $ab \in Q$, so $a \in Q$ or $b \in Q$. That is $a \in p$ or $b \in p$.

First case, $R_1 \subseteq Q$: In this case $R_1 \subseteq R'$ that is $R' = R_1$. Therefore $Q = p + R_1$. Moreover $R_1^2 \subseteq Q \cap R_0 = p$. That is p is a prime ideal of R_0 containing R_1^2 .

Second case $R_1 \not\subseteq Q$: In this case $R' \neq R_1$. Let $a \in R_0$ and $r \in R_1$ such that $ar \in R'$. We see that $ar \in Q$, so $a \in Q$ or $r \in Q$ that is $a \in p \subseteq (R' :_{R_0} R_1)$ or $r \in R'$. It follows that R' is a prime submodule of R_1 . Clearly, $p \subseteq (R' :_{R_0} R_1)$. Let $a \in (R' :_{R_0} R_1)$. Since $R' \neq R_1$, there exist $b \in R_1$ such that $b \notin R'$. But $ab \in R'$, so that $ab \in Q$. Since Q is a \mathbb{Z}_2 -graded prime ideal of R and a, b are homogeneous elements, $a \in Q$ or $b \in Q$. Thus $a \in p = Q \cap R_0$ since $b \notin R'$. It follows that $p = (R' :_{R_0} R_1)$. Now, if $R_1^3 \subseteq R'$, then $R_1^2 \subseteq (R' :_{R_0} R_1) + R' = Q$. Pick $b \in R_1 - R'$, then $b^2 \in Q$, so $b \in Q \cap R_1 = R'$, a contradiction. Thus $R_1^3 \not\subseteq R'$. \square

In order to determine the homogeneous spectrum of a strongly \mathbb{Z}_2 -graded commutative ring, we need the following result.

Proposition 3.4. *Let R be a \mathbb{Z}_2 -graded commutative ring.*

- (1) *If p is a prime ideal of R_0 such that $R_1^2 \not\subseteq p$, then $(pR_1 :_{R_0} R_1) = p$.*
- (2) *Let Q be a \mathbb{Z}_2 -graded ideal of R such that $R_1^2 + Q \cap R_0 = R_0$. Then Q is a graded prime ideal if and only if $Q = p + pR_1$ where p is a prime ideal of R_0 .*

Proof. (1) Clearly, $p \subseteq (pR_1 :_{R_0} R_1)$. Let $a \in (pR_1 :_{R_0} R_1)$, then $aR_1 \subseteq pR_1$, so $aR_1^2 \subseteq pR_1^2 \subseteq p$, so that $a \in p$ since $R_1^2 \not\subseteq p$.

- (2) Since $R_1^2 + Q \cap R_0 = R_0$, $1 = \sum_{i=1}^n a_i b_i + \alpha$ where $a_i, b_i \in R_1$ and $\alpha \in Q \cap R_0$. If Q is a graded prime ideal of R , then $R_1 \not\subseteq Q$ since $R_1^2 \not\subseteq Q$. By Theorem 3.3, $Q = (R' :_{R_0} R_1) + R'$ where R' is a prime submodule of R_1 with $R_1^3 \not\subseteq R_1$. We see

that $p := (R' :_{R_0} R_1) = Q \cap R_0$ is a prime ideal and $pR_1 \subseteq R'$. Let $r \in R'$, then $r = \sum_{i=1}^n (b_i r) a_i + \alpha r$. If $r' \in R_1$, then $(b_i r) r' = (b_i r') r \in R'$. It follows that $b_i r \in (R' :_{R_0} R_1) = p$. So $r = \sum_{i=1}^n (b_i r) a_i + \alpha r \in pR_1$. Thus $R' = pR_1$, that is $Q = p + pR_1$.

Conversely, assume that $Q = p + pR_1$ where p is a prime ideal of R_0 , in fact $p = Q \cap R_0$. Since $R_1^2 + p = R_0$, $R_1^2 \not\subseteq p$. By the previous result $p = (pR_1 :_{R_0} R_1)$. Now, it is enough to show that pR_1 is a prime submodule of R_1 and $R_1^3 \not\subseteq pR_1$. Let $a \in R_0$ and $r \in R_1$ such that $ar \in pR_1$. We have $r = \sum_{i=1}^n (b_i r) a_i + \alpha r$,

so that $ar = \sum_{i=1}^n (ab_i r) a_i + \alpha ar$. We see that $a(b_i r) = (ar) b_i \in (pR_1)R_1 \subseteq p$, so $a \in p$ or $b_i r \in p$ for all i . If $a \in p$, then $a \in (pR_1, R_1)$. If $a \notin p$, then $r = \sum_{i=1}^n (b_i r) a_i + \alpha r \in pR_1$. On other hand $pR_1 \neq R_1$ since $(pR_1 : R_1) = p \neq R_0$. Thus pR_1 is a prime submodule of R_1 . If $R_1^3 \subseteq pR_1$, then $(R_1^2)^2 \subseteq pR_1^2 \subseteq p$, and in this case $R_1^2 \subseteq p$, which is not compatible with the hypothesis. It follows that $p + pR_1$ is a \mathbb{Z}_2 -graded prime ideal of R . □

Now, we have the following Corollary, which describes the homogeneous spectrum of a strongly \mathbb{Z}_2 -graded commutative ring.

Corollary 3.5. *Let R be a strongly \mathbb{Z}_2 -graded commutative ring. Then*

$$\mathbb{Z}_2 \text{Spec} R = \{p + pR_1 \mid p \in \text{Spec} R\}.$$

Proof. This follows from the previous Proposition and the fact that, for every prime ideal p of R_0 , $R_1^2 + p = R_0$. □

The following Corollary point out a class of \mathbb{Z}_2 -graded commutative ring for which prime ideals and \mathbb{Z}_2 -graded prime ideals coincide.

Corollary 3.6. *Let R be a \mathbb{Z}_2 -graded ring such that $R_1^2 \subseteq \sqrt{0}$. Then*

- (1) *The \mathbb{Z}_2 -graded prime of R are $p + R_1$ where p is a prime ideal of R_0 .*
- (2) *An ideal of R is \mathbb{Z}_2 -graded prime if and only if it is prime.*

Proof. (1) Immediate from 3.3.

- (2) If Q is a \mathbb{Z}_2 -graded prime ideal of R , then $R_1R_1 \subseteq Q$, so $R_1 \subseteq Q$. By Theorem 3.3, $Q = p + R_1$ where p is a prime ideal of R_0 . It is easy to see $R/Q = R_0/p$ is an integral domain, so Q is a prime ideal of R . Let P be any prime ideal of R and set $p = P \cap R_0$. Since $R_1^2 \subseteq \sqrt{0} \subseteq p \subseteq Q$, $R_1 \subseteq Q$. Therefore $p + R_1 \subseteq Q$. If $z = a + r \in Q$ where $a \in R_0$ and $r \in R_1$, then $a = z - r \in Q \cap R_0 = p$, so $z \in p + R_1$. It follows that $Q = p + R_1$ is a \mathbb{Z}_2 -graded prime ideal of R . \square

Example 3.7. Let R be a commutative ring and M be an R -module. Consider the trivial ring extension $S = R \times M$ as \mathbb{Z}_2 -graded ring with $S_0 = R$ and $S_1 = M$. Clearly, $S_1^2 = 0$. Therefore the \mathbb{Z}_2 -graded prime ideals and prime ideals coincides.

An immediate consequence of the Corollary 3.5 is as follow; if R is a strongly \mathbb{Z}_2 -graded commutative ring, then the homogeneous spectrum of R and the prime spectrum of R_0 are homeomorphic with respect to the Zariski topologies. The next Theorem extend this result by dropping the strongly graded hypothesis.

Theorem 3.8. *Let R be a \mathbb{Z}_2 -graded commutative ring. Then the map $\varphi : \mathbb{Z}_2\text{Spec}R \rightarrow \text{Spec}R_0$, $\varphi(P) = P \cap R_0$ is an homeomorphism of topological spaces with respect to the Zariski topologies.*

Proof. Let $P, Q \in \mathbb{Z}_2\text{Spec}R$ with $P \cap R_0 = Q \cap R_0$. If $x \in P \cap R_1$, then $x^2 \in P \cap R_0 = Q \cap R_0$, so $x^2 \in Q$, hence $x \in Q \cap R_1$. It follows that $P = P \cap R_0 + P \cap R_1 \subseteq Q \cap R_0 + Q \cap R_1 = Q$. By the same argument we have $Q \subseteq P$, thus $P = Q$.

Let $p \in \text{Spec}R_0$. Consider $R' = \{x \in R_1 \mid xR_1 \subseteq p\}$. Then R' is clearly an R_0 -submodule of R_1 .

First case, $R_1^2 \subseteq p$. By Theorem 3.3, $p + R_1$ is a \mathbb{Z}_2 -graded prime ideal of R and $\varphi(p + R_1) = p$.

Second case, $R_1^2 \not\subseteq p$. Note that using the Theorem 3.3, it enough to show that R' is a prime submodule of R_1 with $R_1^3 \not\subseteq R'$ and $p = (R' :_{R_0} R_1)$. If $R_1^3 \subseteq R'$, then $R_1^2R_1^2 = R_1^4 \subseteq p$, since p is prime, we get $R_1^2 \subseteq p$, a contradiction, so $R_1^3 \not\subseteq R'$. Let $a \in R_0$ and $x \in R_1$ such that $ax \in R'$. Then $axR_1 \subseteq p$. Since p is a prime ideal, $a \in p$ or $xR_1 \subseteq p$. If $xR_1 \subseteq p$, then $x \in R'$. If $a \in p$, then $(aR_1)R_1 \subseteq pR_1^2 \subseteq p$, so $aR_1 \subseteq R'$ that is $a \in (R' :_{R_0} R_1)$. Therefore R' is prime submodule of R_1 . Next, we show that $(R' :_{R_0} R_1) = p$. We have $(pR_1)R_1 = pR_1^2 \subseteq p$, so $pR_1 \subseteq R'$, that is $p \subseteq (R' :_{R_0} R_1)$. Now, let $a \in (R' :_{R_0} R_1)$. Then $aR_1 \subseteq R'$, that is $aR_1^2 = (aR_1)R_1 \subseteq p$. Since p is a prime ideal of R_0 and $R_1^2 \not\subseteq p$, we

get $a \in p$. Thus $(R' :_{R_0} R_1) = p$. Finally, $p + R'$ is a \mathbb{Z}_2 -graded prime ideal of R and $\varphi(p + R') = p$.

If $a \in R_0$, then $\varphi^{-1}(V(a)) = V(a)$. Thus φ is a continuous map.

If $r \in R$ is a homogeneous elements, then $\varphi(V(r)) = V(r^2)$. Thus φ^{-1} is also a continuous map. It follows that φ is an homeomorphism with respect to the Zariski topologies. \square

Next, we list a consequences of the previous Theorem. Recall that, a G -graded commutative ring has Noetherian homogeneous spectrum if $G\text{Spec}R$ is a Noetherian topological space with respect to the Zariski topology. The homogeneous dimension of R , denoted by $h \dim R$, is the maximum length n of a chain $P_0 \subset P_1 \subset \dots \subset P_n$ of homogeneous prime ideals of R .

Corollary 3.9. *Let R be \mathbb{Z}_2 -graded commutative ring. The following statements hold.*

- (1) *R has Noetherian homogeneous spectrum if and only if R_0 has Noetherian spectrum.*
- (2) *$h \dim R = \dim R_0$.*

Proof. (1) Immediate.

- (2) In $P_0 \subset P_1 \subset \dots \subset P_n$ is a chain of homogeneous ideals of R , then $\varphi(P_0) \subset \varphi(P_1) \subset \dots \subset \varphi(P_n)$ is a chain of prime ideals of R_0 . Conversely, if $p_0 \subset p_1 \subset \dots \subset p_n$ is a chain of prime ideals of R_0 . For each i , $p_i = \varphi(P_i)$ for some $P_i \in \mathbb{Z}_2\text{Spec}R$, we see that $P_0 \subset P_1 \subset \dots \subset P_n$. it follows that $h \dim R = \dim R_0$. \square

We close this section by a discuss of the homogeneous radical of a graded ideal. Recall that, if I is a G -graded ideal of a G -graded commutative ring R , then the graded radical of I , denoted by $\text{Grad}(I)$, is

$$\text{Grad}(I) = \{x = \sum_{g \in G} x_g / x_g \in \sqrt{I} \text{ for all } g \in G\}$$

Note that $\text{Grad}(I)$ is a G -graded ideal of R .

Lemma 3.10. *Let R be a G -graded commutative ring and I be a G -graded ideal of R . Then*

$$\text{Grad}(I) = \bigcap_P P$$

Where P runs through all homogeneous prime ideals of R containing I .

Proof. It is easy to see that $\text{Grad}(I) \subseteq \bigcap_P P$. Let $x \in \bigcap_P P$ and assume that $x \notin \text{Grad}(I)$. Then there exists x_g (g -component of x) for some

$g \in G$ such that $x_g \notin \sqrt{I}$, that is $x_g^k \notin I$ for all $k \in \mathbb{N}$. Consider the homogeneous multiplicative subset $S := \{x_g^k / k \in \mathbb{N}\}$. We see that $I \cap S = \emptyset$. By [Lemma 4.7,[9]], there exists a graded prime ideal P such that $I \subseteq P$ and $P \cap S = \emptyset$, that is $I \subseteq P$ and $x_g^k \notin P$ for all $k \in \mathbb{N}$, a contradiction. \square

For an ideal I_0 of R_0 , denote $R_1[I_0]$ the subset $R_1[I_0] = \{x \in R_1 / x^2 \in I_0\}$.

Lemma 3.11. *Let R be a \mathbb{Z}_2 -graded commutative ring and P be a \mathbb{Z}_2 -graded prime ideal of R . Then*

$$P = p + R_1[P]$$

Where $p = P \cap R_0$.

Proof. As in the proof of Theorem 3.8, we have $P = p + R'$ where $p = P \cap R_0$ and $R' = \{x \in R_1 / xR_1 \subseteq p\}$. So, it is enough to show that $R' = R_1[p]$. Clearly, $R' \subseteq R_1[p]$. Let $x \in R_1[p]$. Then $(xR_1)^2 = x^2R_1^2 \subseteq p$, hence $xR_1 \subseteq p$ since p is a prime ideal of R_0 and xR_1 is an ideal of R_0 . Thus $x \in R'$. \square

Theorem 3.12. *Let R be a \mathbb{Z}_2 -graded commutative ring and J be a \mathbb{Z}_2 -graded ideal of R . Then*

$$\text{Grad}(J) = \sqrt{J_0} + R_1[J_0]$$

Where $J_0 = J \cap R_0$.

Proof. Note that if P is a \mathbb{Z}_2 -graded ideal of R , then $J \subseteq P$ if and only if $J_0 \subseteq p$ where $p = P \cap R_0$. Thus

$$\begin{aligned} \text{Grad}(J) &= \bigcap_{J_0 \subseteq p} (p + R_1[p]) = \bigcap_{J_0 \subseteq p} p + \bigcap_{J_0 \subseteq p} R_1[p] \\ &= \sqrt{J_0} + \bigcap_{J_0 \subseteq p} R_1[p] \end{aligned}$$

On other hand,

$$\begin{aligned} \bigcap_{J_0 \subseteq p} R_1[p] &= \{x \in R_1 / x^2 \in p, \text{ for all } p \supseteq J_0\} \\ &= \{x \in R_1 / x^2 \in \sqrt{J_0}\} = R_1[J_0] \end{aligned}$$

It follows that $R' = R_1[p]$ as desired. \square

4. \mathbb{Z}_2 -Maximal graded ideals

Definition 4.1. Let R be a G -graded ring. A graded ideal M of R is said to be graded maximal ideal of R if $M \neq R$ and if J is a graded ideal of R such that $M \subseteq J \subseteq R$, then $J = M$ or $J = R$.

Example 4.2. Let $R = \mathbb{Z} + i\mathbb{Z}$ be a \mathbb{Z}_2 -graded ring with $R_0 = \mathbb{Z}$ and $R_1 = i\mathbb{Z}$. Let $M = 2\mathbb{Z} + i2\mathbb{Z}$. Then M is a \mathbb{Z}_2 -graded maximal ideal of R and it is not a maximal (prime) ideal of R .

Proof. Let J be a \mathbb{Z}_2 -graded ideal of R with $M \subsetneq J \subseteq R$. Let $x = a + ib \in J - M$ with $(a, b) \in \mathbb{Z}^2$. Then $a \notin 2\mathbb{Z}$ or $b \notin 2\mathbb{Z}$. If $a \notin 2\mathbb{Z}$, then $a = 2k + 1$ where $k \in \mathbb{Z}$. So $1 = a - (2k) \in J$, hence $J = R$. If $b \notin 2\mathbb{Z}$, then $b = (2l) + 1$ where $l \in \mathbb{Z}$, so $i = ib - 2il \in J$. Hence $-1 = i^2 \in J$. Thus $J = R$. Moreover M is not a prime ideal see the Remark 3.2, hence it is not a maximal ideal. \square

In the following Theorem we give a characterization of \mathbb{Z}_2 -graded maximal ideals of R .

Theorem 4.3. *Let R be a \mathbb{Z}_2 -graded commutative ring and Q be a \mathbb{Z}_2 -graded ideal of R . Then Q is a \mathbb{Z}_2 -graded maximal ideal of R if and only if*

- (1) *either $Q = p + R_1$ where $p \in V(R_1^2)$ is a maximal ideal of R_0 ,*
or
- (2) *$Q = (R' :_{R_0} R_1) + R'$ where R' is a maximal submodule of R_1
such that $R_1^3 \not\subseteq R'$.*

Proof. (\Rightarrow): Let Q be a \mathbb{Z}_2 -graded maximal ideal of R , in particular Q is a \mathbb{Z}_2 -graded prime ideal of R .

First case; $R_1 \subseteq Q$, from Theorem 3.3, $Q = p + R_1$ where $p \in V(R_1^2)$. Let p' be a maximal ideal containing p , then $Q \subseteq p' + R_1$ and $p' + R_1$ is a proper \mathbb{Z}_2 -graded ideal of R , by maximality, $Q = p' + R_1$, so $p = p'$ is a maximal ideal of R_0 containing R_1^2 .

Second case; $R_1 \not\subseteq Q$, by Theorem 3.3, $Q = (R' :_{R_0} R_1) + R'$ where R' is a prime R_0 -submodule of R_1 such that $R_1^3 \not\subseteq R'$. Let R'' be an R_0 -submodule of R_1 such that $R' \subseteq R''$ and consider the \mathbb{Z}_2 -graded ideal $J := (R'' :_{R_0} R_1) + R''$. Clearly $Q \subseteq J$, by maximality $J = Q$ or $J = R$, that is $R'' = R'$ or $R'' = R_1$. Thus R' is a maximal submodule of R_1 .

(\Leftarrow): Let p be a maximal ideal of R_0 containing R_1^2 and $Q := p + R_1$. Let J be a \mathbb{Z}_2 -graded ideal of R such that $Q \subseteq J$, then $R_1 = Q \cap R_1 \subseteq J \cap R_1$, so $J \cap R_1 = R_1$. Also $p = Q \cap R_0 \subseteq J \cap R_0$. Since p is a maximal ideal of R_0 , $J \cap R_0 = p$ or $J \cap R_0 = R_0$ that is $J = Q$ or $J = R$. Thus Q is a \mathbb{Z}_2 -graded maximal ideal of R .

Let R' be a maximal submodule of R_1 such that $R_1^3 \not\subseteq R'$ and $Q := (R' :_{R_0} R_1) + R'$. Clearly Q is a proper \mathbb{Z}_2 -graded ideal of R . Now, let P be any \mathbb{Z}_2 -graded maximal ideal of R containing Q . Denote $p' := P \cap R_0$. We have $(R' :_{R_0} R_1) = Q \cap R_0 \subseteq P \cap R_0$. Since R' is a maximal submodule of R_1 , $(R' :_{R_0} R_1)$ is a maximal ideal of R_0 ,

it follows that $p' = (R' :_{R_0} R_1)$ or $p' = R_0$. If $p' = R_0$, then $P = R$ since $1 \in P$, with is not compatible with the fact that P is a \mathbb{Z}_2 -graded maximal ideal. So $p' = (R' :_{R_0} R_1)$, hence $\varphi(Q) = \varphi(P)$ where φ is the map defined in Theorem 3.8, by the said Theorem, we have $P = Q$. Thus Q is a \mathbb{Z}_2 -graded maximal ideal of R . \square

Next, we establish a relation between \mathbb{Z}_2 -graded maximal ideal of R and maximal ideal of R_0 . Before, we establish the following Lemma.

Lemma 4.4. *Let R be a \mathbb{Z}_2 -graded ring and R' be an R_0 -submodule of R_1 such that $R_1^3 \not\subseteq R'$. The followings statements are equivalent.*

- (1) R' is a maximal submodule of R_1 .
- (2) $(R' :_{R_0} R_1)$ is maximal ideal of R_0 .

In this case $R' = (R' :_{R_0} R_1)R_1$.

Proof. If R' is a maximal submodule of R_1 , then clearly $(R' :_{R_0} R_1) \neq R_0$. Pike $\alpha \in R_0 - (R' :_{R_0} R_1)$, then $\alpha r \notin R'$ for some $r \in R_1$. It follows that $R' \subsetneq R' + R_0(\alpha r)$, so $R' + R_0(\alpha r) = R_1$. In particular, $r = \beta\alpha r + r'$ where $\beta \in R_0$ and $r' \in R'$. So that $(1 - \beta\alpha)r = r' \in R'$. Thus $1 - \beta\alpha \in (R' :_{R_0} R_1)$, that is $(R' :_{R_0} R_1) + (\alpha) = R_0$. Hence $(R' :_{R_0} R_1)$ is a maximal ideal of R_0 . For the converse, assume that $(R' :_{R_0} R_1)$ is a maximal ideal of R_0 . Let R'' be a submodule of R_1 with $R' \subseteq R''$. Then $(R' :_{R_0} R_1) \subseteq (R'' :_{R_0} R_1)$, hence $(R'' :_{R_0} R_1) = R_0$ or $(R'' :_{R_0} R_1) = (R' :_{R_0} R_1)$. If $(R'' :_{R_0} R_1) = R_0$, then clearly $R'' = R_1$. Assume that $(R'' :_{R_0} R_1) = (R' :_{R_0} R_1)$. Let $x \in R''$, then $xR_1 \subseteq (R'' :_{R_0} R_1) = (R' :_{R_0} R_1)$, so $xR_1^2 \subseteq R'$. Since R' is a prime submodule of R_1 and $R_1^2 R_1 = R_1^3 \not\subseteq R'$, it follows that $x \in R'$. Thus $R'' = R'$. Therefore R' is maximal submodule of R_1 .

We see that $R_1^2 \not\subseteq (R' :_{R_0} R_1)$, so that $(R' :_{R_0} R_1) + R_1^2 = R_0$ since $(R' :_{R_0} R_1)$ is a maximal ideal of R_0 . Thus $1 = \sum_{k=1}^n a_k b_k + \alpha$ where $a_k, b_k \in R_1$ and $\alpha \in (R' :_{R_0} R_1)$. Now, if $r \in R'$, then $r = \sum_{k=1}^n r a_k b_k + \alpha r \in (R' :_{R_0} R_1)R_1$ since $a_i r \in (R' :_{R_0} R_1)$ and $\alpha \in (R' :_{R_0} R_1)$. Therefore $R' = (R' :_{R_0} R_1)R_1$. \square

Now, we are able to establish a relation between \mathbb{Z}_2 -graded maximal ideals of R and maximal ideals of R_0 , as follow.

Theorem 4.5. *Let R be a \mathbb{Z}_2 -graded commutative ring. Then*

$$\mathbb{Z}_2\text{Max}R = \{p+R_1 / p \in \text{Max}R_0 \cap V(R_1^2)\} \cup \{p+pR_1 / p \in \text{Max}R_0 \cap D(R_1^2)\}$$

Proof. Let R' be a maximal submodule of R_1 such that $R_1^3 \not\subseteq R'$ and denote $p := (R' :_{R_0} R_1)$. By the previous Lemma, p is a maximal ideal of R_0 and $R' = pR_1$. Moreover $R_1^2 \not\subseteq p$ since $R_1^3 \not\subseteq R'$. Thus $(R' :_{R_0} R_1) + R' = p + pR_1$. Conversely, let p be a maximal ideal of R_0 such that $R_1^2 \not\subseteq p$ and denote $R' = pR_1$. By the Proposition 3.4, $(R' :_{R_0} R_1) = p$ is a maximal ideal of R_0 . Since $R_1^2 \not\subseteq p$, $R_1^3 \not\subseteq R'$. It follows from the previous Lemma that R' is a maximal submodule of R_1 . \square

An immediate consequence, we have the following Corollary.

Corollary 4.6. *Let R be a strongly \mathbb{Z}_2 -graded commutative ring. Then*

$$\mathbb{Z}_2\text{Max}R = \{p + pR_1 \mid p \in \text{Max}R_0\}$$

Proof. Follows immediately from the previous Proposition. \square

Corollary 4.7. *Let R be a \mathbb{Z}_2 -graded commutative ring. Then*

- (1) φ induce an homeomorphism between $\mathbb{Z}_2\text{Max}R$ and $\text{Max}R_0$.
- (2) R is a \mathbb{Z}_2 -graded local ring if and only if R_0 is a local ring.

Proof. (1) Due to the fact that, \mathbb{Z}_2 -graded maximal ideals of R (respectively, maximal ideals of R_0) are exactly the closed points of $\mathbb{Z}_2\text{Spec}R$ (respectively $\text{Spec}R_0$) with respect to the Zariski topology.

- (2) This is the case of one closed point. \square

5. \mathbb{Z}_2 -Graded field

Definition 5.1. Let R be a G -graded ring.

- (1) R is called a G -graded integral domain if whenever r and s are homogeneous elements of R such that $rs = 0$, then either $r = 0$ or $s = 0$.
- (2) R is called a G -graded field if every nonzero homogeneous element is a unit.

Remark 5.2. A G -graded commutative ring R is a G -graded domain if and only if the zero ideal is a G -graded prime ideal of R . Whereas, a G -graded commutative ring is a G -graded field if and only if the zero ideal is a G -graded maximal ideal of R .

Theorem 5.3. *Let R be a \mathbb{Z}_2 -graded commutative ring. Then R is a \mathbb{Z}_2 -graded integral domain if and only if $R_1 = 0$ and R_0 is an integral domain or 0 is a prime submodule of R_1 such that $(0 :_{R_0} R_1) = 0$ and $R_1^3 \neq 0$.*

Proof. (\Rightarrow) If R is a \mathbb{Z}_2 -graded integral domain, then 0 is a \mathbb{Z}_2 -graded prime ideal of R . By Theorem 3.3, $0 = p + R_1$ where $p \in V(R_1^2)$ or $0 = (R' :_{R_0} R_1) + R'$ where R' is a prime submodule of R_1 with $R_1^3 \not\subseteq R'$. If $0 = p + R_1$, then R_0 and $p = 0$ is a prime ideal of R_0 , that is $R_1 = 0$ and R_0 is an integral domain. If $0 = (R' :_{R_0} R_1) + R'$, then $R' = 0$ and $(R' :_{R_0} R_1) = 0$, as desired.

(\Leftarrow) Immediate. \square

Corollary 5.4. *Let R be a strongly \mathbb{Z}_2 -graded commutative ring. Then R is a \mathbb{Z}_2 -graded integral domain if and only if R_0 is an integral domain.*

Proof. By the Corollary 3.5. \square

Theorem 5.5. *Let R be a \mathbb{Z}_2 -graded commutative ring. Then R is a \mathbb{Z}_2 -graded field if and only if R_0 is a field and either $R_1 = 0$ or $R_1^2 \neq 0$.*

Proof. R is a \mathbb{Z}_2 -graded field if and only if 0 is a \mathbb{Z}_2 -graded maximal ideal of R . According to Theorem 4.5, $0 = p + R_1$ with p maximal ideal of R_0 containing R_1^2 , which is equivalent to $R_1 = 0$ and $0 = p$ is a maximal ideal of R_0 . Or $0 = p + pR_1$ with p is a maximal ideal of R_0 such that $R_1^2 \not\subseteq p$, which is equivalent to $0 = p$ is a maximal ideal of R_0 and $R_1^2 \neq 0$. \square

Corollary 5.6. *Let R be a \mathbb{Z}_2 -graded commutative ring. Then R is a \mathbb{Z}_2 -graded field if and only if R_0 is a field and either $R_1 = 0$ or $R_1 = R_0b$ where $b \in R_1$ such that $b^2 \neq 0$.*

Proof. We may assume that $R_1 \neq 0$. If R is a \mathbb{Z}_2 -graded field, then by the previous Theorem R_0 is a field and $R_1^2 \neq 0$. So $bb' \neq 0$ for some $b, b' \in R_1$ since R_1^2 is generated by all elements bb' where $b, b' \in R_1$. As $bb' \neq 0$ and R_0 is a field, bb' is a unit in R_0 . There exists $\alpha \in R_0$ such that $bb'\alpha = 1$. It follows that $r = \alpha(rb')b$ for all $r \in R_1$. Therefore $R_1 = R_0b$. Clearly $b^2 \neq 0$ since $R_1^2 \neq 0$. The converse is immediate, it is easy to see that every homogeneous elements in a unit. \square

For a commutative ring S we look at $S[X]$ as a \mathbb{Z}_2 -graded commutative ring see Example 1.1. We see that $(S[X])_0 = \{P(X^2) / P \in S[X]\}$ and $(S[X])_1 = \{XP(X^2) / P \in S[X]\}$. For $\alpha \in S$, $(X^2 - \alpha)$ is a \mathbb{Z}_2 -graded ideal of $S[X]$. Therefore $\frac{S[X]}{(X^2 - \alpha)}$ is a \mathbb{Z}_2 -graded commutative ring and clearly $\left(\frac{S[X]}{(X^2 - \alpha)}\right)_0 = S$ and $\left(\frac{S[X]}{(X^2 - \alpha)}\right)_1 = S\bar{X}$. Note that if S is a field and $\alpha \neq 0$, then $\frac{S[X]}{(X^2 - \alpha)}$ is a graded field.

Corollary 5.7. *Let R be a \mathbb{Z}_2 -graded commutative ring. If R is a \mathbb{Z}_2 -graded field, then R_0 is a field and either $R_1 = 0$ or R is isomorphic to $\frac{R_0[X]}{(X^2 - \alpha)}$ as \mathbb{Z}_2 -graded rings, where α is a nonzero element of R_0 .*

Proof. We may assume that $R_1 \neq 0$. By the previous Corollary R_0 is a field and $R_1 = R_0b$ where $b \in R_1$ such that $b^2 = 0$. Let $f : R_0[X] \rightarrow R$ the map defined by $f(P) = P(b)$. Clearly f is a rings morphism. We have $f(P(X^2)) = P(b^2) \in R_0$ since $b^2 \in R_0$, also $f(XP(X^2)) = bP(b^2) \in R_1$. It follows that f is a \mathbb{Z}_2 -graded morphism. For $r = r_0 + r_1 \in R$. We see that $r_1 = ab$ for some $a \in R_0$, so that $f(r_0 + aX) = r_0 + ab = r$. Now, let $\alpha = b^2 \in R_0$, then $(X^2 - \alpha) \subseteq \ker f$. Let $P \in \ker f$, by the euclidean division, $P = (X^2 - \alpha)Q + dX + c$ where $Q \in R_0[X]$ and $d, c \in R_0$. But $P(b) = 0$ implies that $db + c = 0$, so that $d = 0$ and $c = 0$ (if $d \neq 0$, then d is a unit, so $b = d^{-1}c \in R_0 \cap R_1$ which is not possible). It follows that $\ker f = (X^2 - \alpha)$. \square

6. \mathbb{Z}_2 -Graded integral domain and integral domain

For $x = a + r \in R$ where $(a, r) \in R_0 \times R_1$, define $N(x) := a^2 - r^2$. Then, $N(xy) = N(x)N(y)$ and $N(x) \in R_0$ for all $x, y \in R$. Also $N(0) = 0$ and $N(1) = 1$. Next we establish the relation between some properties of N and the integrity of R . To do, let $\mathcal{N} := \{x \in R / N(x) = 0\}$.

Theorem 6.1. *Let R be a \mathbb{Z}_2 -graded ring. The following statements are equivalent.*

- (1) R is an integral domain.
- (2) R is a \mathbb{Z}_2 -graded integral domain and $\mathcal{N} = 0$.

Proof. (1) \Rightarrow (2): Clearly, if R is an integral domain then R is a \mathbb{Z}_2 -graded integral domain. Let $x = a + r \in \mathcal{N}$, then $a^2 - r^2 = 0$, so that $(a+r)(a-r) = 0$, thus $a+r = 0$ or $a-r = 0$. Hence $x = 0$. It follows that $\mathcal{N} = 0$.

(2) \Rightarrow (1): Let $x, y \in R$ with $xy = 0$. Then $N(x)N(y) = 0$. Note that $N(x)$ and $N(y)$ are homogeneous, so that $N(x) = 0$ or $N(y) = 0$ that is $x \in \mathcal{N}$ or $y \in \mathcal{N}$. Thus $x = 0$ or $y = 0$. \square

REFERENCES

- [1] W. Heinzer and M. Roitman. *The homogeneous spectrum of a graded commutative ring*. Proc. Amer. Math. Soc. 130 (2002), 1573-1580
- [2] H. A. Khashan, *Graded rings in which every graded ideal is a product of Gr-primary ideals*, Int. J. Algebra, 2(13-16) (2008), 779-788.
- [3] CP. Lu *Prime submodules of modules*. Commentarii Mathematici Universitatis Sancti Pauli 1984; 33 (1): 61 – 69.
- [4] C. Nastasescu. *Strongly Graded Rings of Finite Groups*, Communications in Algebra, 11(10) (1983), 1033-1071.
- [5] C. Nastasescu and F. Van Oystaeyen, *Graded ring theory*, Mathematical Library 28, North Holland, Amesterdam, 1982.

- [6] J. Ohm and R. Pendleton. *Rings with Noetherian spectrum*. Duke Math. J.35:631-639, (1968).
- [7] M. Refai, M. Hailat and S. Obiedat, *Graded radicals on graded prime spectra*, FarEast J. of Math. Sci., part I (2000), 59-73.
- [8] A. Seidenberg. *A note on the dimension theory of rings*, Pac. J. Math., 3 (1953), 505-512.
- [9] X. Wu *Graded w -Noetherian modules over graded rings*. Bull. Korean Math. Soc. 2020;57:1319-1334. <https://doi.org/10.4134/BKMS.b191007>
- [10] A. Yousefian Darani. *Topologies on $\text{Spec}_g(M)$* , Buletinul Academiei de Ştiinţe a Moldovei. Matematica. 3(67), 2011, 45-53.
- [11] A. Yousefian Darania and S. Motmaen. *Zariski topology on the spectrum of graded classical prime submodules*. Appl. Gen. Topol. 14, no. 2 (2013), 159-169.

(Mohamed Aqalmoun) DEPARTMENT OF MATHEMATICS, HIGHER NORMAL SCHOOL,
SIDI MOHAMED BEN ABDELLAH UNIVERSITY, FEZ, MOROCCO.

Email address: maqalmoun@yahoo.fr