

Optimal scheduling of entropy regulariser for continuous-time linear-quadratic reinforcement learning

Lukasz Szpruch* Tanut Treetanthiploet† Yufei Zhang‡

Abstract. This work uses the entropy-regularised relaxed stochastic control perspective as a principled framework for designing reinforcement learning (RL) algorithms. Herein agent interacts with the environment by generating noisy controls distributed according to the optimal relaxed policy. The noisy policies on the one hand, explore the space and hence facilitate learning but, on the other hand, introduce bias by assigning a positive probability to non-optimal actions. This exploration-exploitation trade-off is determined by the strength of entropy regularisation. We study algorithms resulting from two entropy regularisation formulations: the exploratory control approach, where entropy is added to the cost objective, and the proximal policy update approach, where entropy penalises the divergence of policies between two consecutive episodes. We analyse the finite horizon continuous-time linear-quadratic (LQ) RL problem for which both algorithms yield a Gaussian relaxed policy. We quantify the precise difference between the value functions of a Gaussian policy and its noisy evaluation and show that the execution noise must be independent across time. By tuning the frequency of sampling from relaxed policies and the parameter governing the strength of entropy regularisation, we prove that the regret, for both learning algorithms, is of the order $\mathcal{O}(\sqrt{N})$ (up to a logarithmic factor) over N episodes, matching the best known result from the literature.

Key words. Continuous-time reinforcement learning, linear-quadratic, entropy regularisation, exploratory control, proximal policy update, regret analysis

AMS subject classifications. 62L05, 49N10, 93E35, 94A17

1 Introduction

Reinforcement learning (RL) is concerned with sequential decision-making in an uncertain environment and is a core topic in machine learning. Two key concepts in RL are *exploration*, which corresponds to learning via interactions with the environment, and *exploitation*, which corresponds to optimising the objective function given accumulated information (see [25]). They are at odds with each other, as learning the environment often requires acting suboptimally with respect to existing knowledge [18]. Thus it is crucial to develop effective exploration strategies and to optimally balance exploration and exploitation.

Randomisation and entropy-regularised relaxed control. A common exploration technique in RL is to adopt randomised actions. Randomisation generates stochastic policies that explore the environment, but simultaneously introduces additional biases by assigning positive probability to, potentially, non-optimal actions. The precise impact of randomisation has been studied extensively for discrete-time RL problems, including finite state-action models [7, 3] and linear-quadratic (LQ) models [20, 5, 24]. This is often quantified by the *regret* of learning, which measures the difference between the value function of designed policies and the optimal value function that could have been achieved had the agent known the environment. It is shown that to achieve the optimal regret, the level of randomisation has to decrease at a proper rate over the learning process.

For RL problems in continuous time and space, [28] proposes an entropy-regularised relaxed control framework to systematically design stochastic policies. Relaxed control formulation has been introduced in

*School of Mathematics, University of Edinburgh and Alan Turing Institute, L.Szpruch@ed.ac.uk

†Alan Turing Institute, ttreetanthiploet@turing.ac.uk

‡Department of Statistics, London School of Economics and Political Science, y.zhang389@lse.ac.uk

classical control theory for the existence of optimal controls (see e.g., [31, 21]), but its entropy regularised variants have recently attracted much attention for their wide applications in learning. Entropy regularisation ensures the optimal stochastic policy has a nondegenerate distribution over actions and hence explores the environment (see e.g., [28, 29, 4, 15, 9, 13, 14]). It also leads to a policy that is Lipschitz stable with respect to perturbations of the underlying model [22]. Such a stability property is crucial for the analysis of algorithm regret [12, 26]. These advantages make entropy-regularised relaxed stochastic control formulation a good choice for the principled design of efficient learning algorithms.

However, despite the recent influx of interest on entropy-regularised control problems, to the best of our knowledge there is no work on how to control the entropy regularising weight to achieve the best possible algorithm regret. A large degree of entropy regularisation often signifies a high level of exploration, which facilitates more efficient learning but also creates a larger exploration cost. Furthermore relaxed control formulation does not constitute a complete learning algorithm as the set of admissible policies for the agent are functions mapping time and space into actions and not into probability distributions over all actions.

Therefore, finding the optimal scheduling of entropy regularisation requires addressing the following two questions: A) How to follow a relaxed policy to interact with a continuous-time system? B) How do the policy execution and regularising weight affect the exploration-exploitation trade-off over the entire learning process?

Main Contributions. This work addresses the above questions for finite horizon continuous-time episodic LQ-RL problems.

- We prove that a Gaussian relaxed policy can be approximated with a single episode by piecewise constant independent Gaussian noises on a time grid. The error of this noisy execution is of half-order in both the mesh size of the grid and the variance of the relaxed policy in the high-probability sense, and is of first-order in both the mesh size and the variance under expectation (Theorem 3.3). This result should be contrasted with the heuristic implementation of a relaxed policy suggested in [28, 14], which involves independent agents interacting over a large number of episodes (see Example 2.1).
- We use entropy-regularised relaxed control problem formulation to design algorithms that achieve the best-known regret for episodic LQ-RL problems. At each episode, the algorithm executes a Gaussian relaxed policy on a time grid, estimates the unknown parameters by a Bayesian approach based on observed trajectories, and then designs a Gaussian policy by solving an entropy-regularised control problem. Two widely used entropy regularisation formulations are analysed: the exploratory control approach (i.e., Algorithm 1) where entropy serves as an exploration reward [28, 29, 9, 13, 14], and the proximal policy update approach (i.e., Algorithm 2) where entropy penalises the divergence of policies between two consecutive episodes [23, 11, 17]. By optimising the execution mesh size and regularising weight, both algorithms achieve an expected regret of the order $\mathcal{O}(\sqrt{N})$ (up to a logarithmic factor) over N episodes, matching the best possible results from the literature (see Theorems 2.2 and 2.3).

Related works. To the best of our knowledge, this is the first theoretical work on the execution of a relaxed policy, on the optimal scheduling of entropy regulariser for RL problems, and on regret bounds of learning algorithms with proximal policy updates.

The type of execution noise corresponds to a properly scaled discrete-time white noise process, which is related to the randomisation technique in [6] (see Section 2.4 for details). The scheduling of entropy regulariser relies on quantifying the precise impact of the regularising weight on the regret bound for continuous-time LQ-RL. Available theoretical results on regret bounds for continuous-time RL are very limited, including [10, 8] for infinite horizon problems with stationary policies, [12, 1] for finite horizon problems with pure exploitation policies, and [26] for finite horizon problems with explicit separation of exploitation and exploration phases. Regret analysis of relaxed policies is more involved than those in the existing literature. Indeed, the relaxed policies explore and exploit simultaneously, and one has to disentangle both effects in the analysis to optimise the regret. Moreover, the proximal policy update approach (i.e., Algorithm 2) results in relaxed policies depending on *all previously estimated parameters*. Such a memory dependence has to be carefully controlled via the regularising weights to optimise the regret.

Notation: For each $n \in \mathbb{N}$, we denote by \mathbb{S}^n (resp. \mathbb{S}_+^n) the space of $n \times n$ symmetric, (resp. symmetric positive definite) matrices, and by $\Lambda_{\min}(A)$ the smallest eigenvalues of a given matrix $A \in \mathbb{S}^n$. For each $T > 0$, complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and Euclidean space $(E, |\cdot|)$, we introduce the following spaces:

- $L^\infty([0, T]; E)$ (resp. $C([0, T]; E)$) is the space of measurable (resp. continuous) functions $\phi : [0, T] \rightarrow E$ satisfying $\|\phi\|_\infty = \text{ess sup}_{t \in [0, T]} |\phi_t| < \infty$.
- $L^\infty(\Omega \times [0, T]; E)$ is the space of measurable functions $\phi : \Omega \times [0, T] \rightarrow E$ satisfying $\|\phi\|_{L^\infty} = \text{ess sup}_{(\omega, t) \in \Omega \times [0, T]} |\phi_t(\omega)| < \infty$.

For notational simplicity, we write $\mathbb{R}_{>0} = (0, \infty)$, $\mathbb{R}_{\geq 0} = [0, \infty)$, and denote by $\mathcal{N}(m, s^2)$, $m \in \mathbb{R}$ and $s \geq 0$, the Gaussian measure on \mathbb{R} with mean m and standard deviation s .

2 Problem formulation and main results

This section studies a LQ-RL problem, where the state dynamics involves unknown drift coefficients. We prove that executing optimal relaxed policies of suitable entropy-regularised control problems leads to learning algorithms with optimal regrets. For the clarity of presentation, we assume all variables are one-dimensional, but the analysis and results can be naturally extended to a multidimensional setting.

2.1 LQ-RL with relaxed policies

LQ control. Let us first recall the standard LQ control problem over feedback controls. Let $T > 0$, $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space on which a Brownian motion $W = (W_t)_{t \in [0, T]}$ is defined, and $\mathcal{F}_0 \subset \mathcal{F}$ be a σ -algebra independent of W . For fixed $\theta^* = (A^*, B^*) \in \mathbb{R}^{1 \times 2}$, consider minimising the following cost functional:

$$J^{\theta^*}(\phi) := \mathbb{E} \left[\int_0^T f(t, X_t^{\theta^*, \phi}, \phi(t, X_t^{\theta^*, \phi})) dt + g(X_T^{\theta^*, \phi}) \right], \quad (2.1)$$

over all measurable functions $\phi : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ such that the controlled dynamics

$$dX_t = (A^* X_t + B^* \phi(t, X_t)) dt + \bar{\sigma}_t dW_t, \quad t \in [0, T]; \quad X_0 = x_0 \quad (2.2)$$

admits a unique square integrable solution $X^{\theta^*, \phi}$ satisfying $\mathbb{E} \left[\int_0^T |\phi(t, X_t^{\theta^*, \phi})|^2 dt \right] < \infty$, and f and g satisfy for all $(t, x, a) \in [0, T] \times \mathbb{R} \times \mathbb{R}$,

$$f(t, x, a) = Q_t x^2 + 2S_t x a + R_t a^2 + 2p_t x + 2q_t a, \quad g(x) = M x^2 + 2m x. \quad (2.3)$$

We impose the following conditions on the the coefficients.

H.1. $T > 0$, $x_0, A^*, B^* \in \mathbb{R}$, $\bar{\sigma}, Q, S, R, p, q \in C([0, T]; \mathbb{R})$, $M \geq 0$, $m \in \mathbb{R}$, and for all $t \in [0, T]$, $\bar{\sigma}_t, R_t > 0$ and $Q_t - S_t^2 R_t^{-1} \geq 0$.

Given θ^* , under (H.1), standard LQ control theory (see e.g., [30]) shows that (2.1)-(2.2) admits the optimal policy ϕ^{θ^*} , and for all $\theta = (A, B) \in \mathbb{R}^{1 \times 2}$ and $(t, x) \in [0, T] \times \mathbb{R}$

$$\phi^\theta(t, x) = k_t^\theta + K_t^\theta x, \quad \text{with } K_t^\theta := -(BP_t^\theta + S_t)R_t^{-1}, \quad k_t^\theta := -(B\eta_t^\theta + q_t)R_t^{-1}, \quad (2.4)$$

where $P^\theta \in C([0, T]; \mathbb{R}_{\geq 0})$ and $\eta^\theta \in C([0, T]; \mathbb{R})$ satisfy for all $t \in [0, T]$,

$$\frac{d}{dt} P_t^\theta + 2AP_t^\theta - (BP_t^\theta + S_t)^2 R_t^{-1} + Q_t = 0, \quad P_T^\theta = M; \quad (2.5a)$$

$$\frac{d}{dt} \eta_t^\theta + (A - (BP_t^\theta + S_t)R_t^{-1}B)\eta_t^\theta + p_t - (BP_t^\theta + S_t)R_t^{-1}q_t = 0, \quad \eta_T^\theta = m. \quad (2.5b)$$

Episodic LQ-RL and regularised relaxed control. Now assume that the parameter $\theta^* = (A^*, B^*)$ is unknown to the agent. Then the LQ control problem becomes an LQ-RL problem: the agent's objective is to control the system optimally while simultaneously learning the parameter θ^* . In an episodic RL framework, agent learns about θ^* by executing a sequence of policies and observing realisations of the corresponding controlled dynamics (see Section 2.3 for details).

Note that learning the dynamics often requires explicit exploration. Indeed, as shown in [26, 1], a greedy policy ϕ^θ , as defined in (2.4), based on the present estimate θ of θ^* in general does not guarantee exploration and consequently fails to converge to the optimal solution. As alluded to in Section 1, a common strategy in RL for encouraging exploration is to incorporate stochastic policy $\nu : [0, T] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$, which maps the current state to a probability measure over the action space \mathbb{R} . The associated cost is given by

$$\tilde{J}(\nu) := \mathbb{E} \left[\int_0^T \int_{\mathbb{R}} f(t, X_t^{\theta^*, \nu}, a) \nu(t, X_t^{\theta^*, \nu}; da) dt + g(X_T^{\theta^*, \nu}) \right], \quad (2.6)$$

where f and g are defined in (2.3), and $X^{\theta^*, \nu}$ is the solution to the following relaxed dynamics:

$$dX_t = \int_{\mathbb{R}} (A^* X_t + B^* a) \nu(t, X_t; da) dt + \bar{\sigma}_t dW_t, \quad t \in [0, T]; \quad X_0 = x_0. \quad (2.7)$$

Note that (2.6)-(2.7) incorporates (2.1)-(2.2) as a special case.

A principle approach to construct stochastic policies is to solve entropy-regularised relaxed control problems. Here we present two such formulations that are widely used in the literature. For notational simplicity, let $\theta = (A, B)$ be the current estimate of θ^* after the m -th episode, $\varrho > 0$ be a regularising weight, and for each measure $\bar{\nu}$ on \mathbb{R} , let $\mathcal{H}(\cdot \| \bar{\nu}) : \mathcal{P}(\mathbb{R}) \rightarrow \mathbb{R} \cup \{\infty\}$ be the relative entropy with respect to $\bar{\nu}$ such that for all $\nu \in \mathcal{P}(\mathbb{R})$,

$$\mathcal{H}(\nu \| \bar{\nu}) = \begin{cases} \int_{\mathbb{R}} \ln \left(\frac{\nu(da)}{\bar{\nu}(da)} \right) \nu(da), & \nu \text{ is absolutely continuous with respect to } \bar{\nu}, \\ \infty, & \text{otherwise.} \end{cases} \quad (2.8)$$

Approach 1 (Exploration reward). An exploratory policy can be chosen as the optimal policy of an exploratory control problem (see e.g., [28, 29, 22, 9, 13, 14]): consider minimising

$$\tilde{J}_\varrho^\theta(\nu) = \mathbb{E} \left[\int_0^T \left(\int_{\mathbb{R}} f(t, X_t^{\theta, \nu}, a) \nu(t, X_t^{\theta, \nu}; da) + \varrho \mathcal{H}(\nu(t, X_t^{\theta, \nu}) \| \nu_{\text{Leb}}) \right) dt + g(X_T^{\theta, \nu}) \right] \quad (2.9)$$

over all $\nu \in \mathcal{M}$, where \mathcal{M} consists of all measurable functions $\nu : [0, T] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$ such that the following controlled dynamics

$$dX_t = \int_{\mathbb{R}} (AX_t + Ba) \nu(t, X_t; da) dt + \bar{\sigma}_t dW_t, \quad t \in [0, T]; \quad X_0 = x_0 \quad (2.10)$$

admits a unique square integrable solution $X^{\theta, \nu}$ satisfying $\mathbb{E} \left[\int_0^T \int_{\mathbb{R}} |a|^2 \nu(t, X_t^{\theta, \nu}; da) dt \right] < \infty$, $\bar{\sigma}$, f and g are the same functions as in (2.1)-(2.2), and ν_{Leb} is the Lebesgue measure on \mathbb{R} .

The term $\varrho \mathcal{H}(\cdot \| \nu_{\text{Leb}})$ is an additional reward to encourage exploration. Under (H.1), standard verification arguments (see e.g., [28, 29]) show that the optimal policy of (2.9)-(2.10) is Gaussian:

$$\nu_\varrho^\theta(t, x) = \mathcal{N} \left(k_t^\theta + K_t^\theta x, \frac{\varrho}{2R_t} \right), \quad (t, x) \in [0, T] \times \mathbb{R}, \quad (2.11)$$

where K^θ and k^θ are defined as in (2.4). Note that $\varrho > 0$ ensures that $\nu_\varrho^\theta(t, x)$ executes each action with positive probability and hence explores the parameter space.

Approach 2 (Proximal policy update). An exploratory policy can also be determined by penalising the divergence of policies between two consecutive episodes. Recall that the greedy policy ϕ^θ in (2.4) satisfies the following pointwise minimisation condition (see e.g., [30, p. 317]):

$$\phi^\theta(t, x) = \arg \min_{a \in \mathbb{R}} H^\theta(t, x, a, 2(P_t^\theta x + \eta_t^\theta)), \quad (t, x) \in [0, T] \times \mathbb{R}, \quad (2.12)$$

where P^θ and η^θ are defined as in (2.5), and $H^\theta : [0, T] \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is the Hamiltonian of (2.1)-(2.2) with $\theta^* = \theta$:

$$H^\theta(t, x, a, y) := (Ax + Ba)y + f(t, x, a), \quad (t, x, a, y) \in [0, T] \times \mathbb{R} \times \mathbb{R} \times \mathbb{R}.$$

Instead of taking the greedy update (2.12), the agent determines a stochastic policy (for the $(m+1)$ -th episode) by minimising the following entropy-regularised Hamiltonian: for each $(t, x) \in [0, T] \times \mathbb{R}$,

$$\nu_\varrho^\theta(t, x) = \arg \min_{\nu \in \mathcal{P}(\mathbb{R})} \left(\int_{\mathbb{R}} H^\theta(t, x, a, 2(P_t^\theta x + \eta_t^\theta)) \nu(da) + \varrho \mathcal{H}(\nu \| \nu^m(t, x)) \right), \quad (2.13)$$

where ν^m is the relaxed policy for the m -th episode.

The relative entropy in (2.13) enforces the updated policy $\nu_\varrho^\theta(t, x)$ to stay close to $\nu^m(t, x)$. This prevents an excessively large policy update, and ensures that the updated policy ν_ϱ^θ explores the space if ν^m is an exploratory policy. Similar ideas have been used to design proximal policy gradient methods in [23, 17] and modified value iterations in [11]. The update rule (2.13) is analogous to the mirror descent algorithm for optimisation problems [2], with the Hamiltonian H^θ playing the role of the gradient in mirror descent.

The policy ν_ϱ^θ in (2.13) is Gaussian if the previous policy ν^m is Gaussian. Indeed, let $\nu^m(t, x) = \mathcal{N}(k_t^m + K_t^m x, (\lambda_t^m)^2)$ for some $(t, x) \in [0, T] \times \mathbb{R}$, then one can directly verify that $\nu_\varrho^\theta(t, x) = \mathcal{N}(k_t^{m+1} + K_t^{m+1} x, (\lambda_t^{m+1})^2)$, where

$$\begin{aligned} k_t^{m+1} &= h_t^m k_t^\theta + (1 - h_t^m) k_t^m, & K_t^{m+1} &= h_t^m K_t^\theta + (1 - h_t^m) K_t^m, \\ (\lambda_t^{m+1})^{-2} &= \frac{2R_t}{\varrho} + (\lambda_t^m)^{-2}, & h_t^m &= \frac{2R_t}{2R_t + \varrho(\lambda_t^m)^{-2}}, \end{aligned} \quad (2.14)$$

with K_t^θ and k_t^θ defined in (2.4).

Observe that in the present LQ setting, solving the relaxed control problems in Approaches 1 and 2 leads to Gaussian-measure-valued policies. The mean of these Gaussian measures is affine in the state and depends on the estimated parameter θ , while the variance of these Gaussian measures depends on the regularising weight ϱ . In the following, we demonstrate how to follow these Gaussian policies to interact with the dynamics (2.2), and how to choose the regularising weight for an efficient learning algorithm.

2.2 Execution of Gaussian relaxed policy

This section studies the execution of a Gaussian relaxed policy via randomisation. The precise definition of Gaussian policies is given as follows.

Definition 2.1. For each $k, K \in L^\infty([0, T]; \mathbb{R})$ and $\lambda \in L^\infty([0, T]; \mathbb{R}_{\geq 0})$, we define a Gaussian relaxed policy $\nu : [0, T] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$, denoted by $\nu \sim \mathcal{G}(k, K, \lambda)$, such that for all $(t, x) \in [0, T] \times \mathbb{R}$, $\nu(t, x) = \mathcal{N}(k_t + K_t x, \lambda_t^2)$.

To execute a Gaussian relaxed policy $\nu \sim \mathcal{G}(k, K, \lambda)$, at each given time and state, the agent will sample a control action based on the distribution $\nu(t, x) \sim \mathcal{N}(k_t + K_t x, \lambda_t^2)$. This suggests interacting with the dynamics (2.2) using a randomised policy $\varphi(t, x) = k_t + K_t x + \lambda_t \xi$, where $\xi \sim \mathcal{N}(0, 1)$ is an injected Gaussian noise independent of the Brownian motion W in (2.2). However, the following example shows that not all injected noises lead to the relaxed cost $\tilde{J}(\nu)$ (even in expectation).

Example 2.1. Let $\mu \in \mathbb{R}$ and $\lambda \geq 0$. Consider the relaxed policy $\nu : [0, 1] \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$ such that $\nu(t, x) = \mathcal{N}(\mu x, \lambda^2)$ for all $(t, x) \in [0, 1] \times \mathbb{R}$, and the cost

$$\tilde{J}(\nu) = \mathbb{E} \left[\int_0^1 \int_{\mathbb{R}} |a|^2 \nu(t, X_t^\nu; da) dt \right], \quad \text{with } X_t^\nu = \int_0^t \int_{\mathbb{R}} a \nu(s, X_s^\nu; da) ds, \quad \forall t \in [0, T].$$

Let $(\zeta_i)_{i \in \mathbb{N}}$ be \mathcal{F}_0 -measurable independent standard normal random variables, and for each $i \in \mathbb{N}$, let $\varphi^i : \Omega \times [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ satisfy $\varphi^i(\cdot, t, x) = \mu x + \lambda \zeta^i$ for all $(t, x) \in [0, 1] \times \mathbb{R}$, and

$$J(\varphi^i) := \mathbb{E} \left[\int_0^1 (\varphi^i(\cdot, t, X_t^i))^2 dt \middle| \mathcal{F}_0 \right], \quad \text{with } X_t^i = \int_0^t \varphi^i(\cdot, s, X_s^i) ds, \quad \forall t \in [0, T].$$

Then $\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N J(\varphi^i) = \tilde{J}(\nu) + \frac{\lambda^2}{2\mu} (e^{2\mu} - 2\mu - 1)$ a.s.

Example 2.1 follows from a direct computation and the strong law of large numbers (see also Lemma 3.1). It shows that repetitive sampling a relaxed policy among multiple episodes does not lead to the relaxed cost $\tilde{J}(\nu)$, and hence is not the correct execution of a relaxed policy.

Instead, we show that the relaxed cost (2.6) can be realised by independently sampling the target policy across different time points within a single episode. A technical difficulty of this approach is that there is no measurable process $\xi = (\xi_t)_{t \in [0, T]}$ such that $\mathbb{E}[\xi_t] = 0$ and $\mathbb{E}[\xi_t^2] = 1$ for all t , and ξ_t and ξ_s are independent for all $s \neq t$ (see [16, Example 1.2.5]). To address this technicality, we consider executing a relaxed policy on a time grid via piecewise constant independent Gaussian noises.

Definition 2.2. For each $\nu \sim \mathcal{G}(k, K, \lambda)$ and grid $\pi = \{0 = t_0 < \dots < t_N = T\}$ with mesh size $|\pi| = \max_{i=0, \dots, N-1} (t_{i+1} - t_i)$, we say $\varphi : \Omega \times [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ is a piecewise constant independent execution of ν on π , denoted by $\varphi \sim \mathcal{R}(k, K, \lambda, \xi^\pi)$, if there exist \mathcal{F}_0 -measurable independent standard normal random variables $(\zeta_i)_{i=0}^{N-1}$ such that for all $(\omega, t, x) \in \Omega \times [0, T] \times \mathbb{R}$, $\varphi(\omega, t, x) = k_t + K_t x + \lambda_t \xi_t^\pi(\omega)$, with $\xi_t^\pi(\omega) = \sum_{i=0}^{N-1} \zeta_i(\omega) \mathbf{1}_{t_i \leq t < t_{i+1}}$.

The process ξ^π models additional Gaussian noises used for executing a relaxed policy, and is assumed, without loss of generality, sampled at $t = 0$ (and hence independent of the Brownian motion W). For each $\varphi \sim \mathcal{R}(k, K, \lambda, \xi^\pi)$, the corresponding expected cost (with fixed realisation of ξ^π) is defined as

$$J(\varphi) := \mathbb{E} \left[\int_0^T f(t, X_t^\varphi, \varphi(\cdot, t, X_t^\varphi)) dt + g(X_T^\varphi) \mid \mathcal{F}_0 \right], \quad (2.15)$$

where f and g are given in (2.3), and X^φ satisfies the following controlled dynamics (cf. (2.2)):

$$dX_t = (A^* X_t + B^* \varphi(\cdot, t, X_t)) dt + \bar{\sigma}_t dW_t, \quad t \in [0, T]; \quad X_0 = x_0, \quad (2.16)$$

Note that $J(\varphi)$ is a random variable, as it depends on the realisations of the injected noise ξ^π .

The following theorem establishes the convergence of $J(\varphi) - \tilde{J}(\nu)$ with respect to the mesh size $|\pi|$ and the standard derivation λ , and consequently proves that φ is a proper execution of ν . The statement follows as a special case of Theorem 3.3 and Corollary 3.4, where the same result is proved for general LQ problems with controlled diffusion coefficients and nonconvex costs.

Theorem 2.1. *Suppose (H.1) holds and let $\nu \sim \mathcal{G}(k, K, \lambda)$. Then there exists a constant $C \geq 0$, depending only on the sup-norms of k, K and the coefficients in (H.1), such that*

(1) *for all grids π of $[0, T]$, all $\varphi \sim \mathcal{R}(k, K, \lambda, \xi^\pi)$ and all $\delta \geq 0$,*

$$\mathbb{P} \left(\left| J(\varphi) - \tilde{J}(\nu) \right| \geq C \sqrt{|\pi|} \|\lambda\|_\infty (1 + \|\lambda\|_\infty) \left(1 + \ln \left(\frac{1}{\delta} \right) \right) \right) \leq \delta,$$

(2) *if $\pi^n = \{i \frac{T}{n}\}_{i=0}^n$ and $\varphi^n \sim \mathcal{R}(k, K, \lambda, \xi^{\pi^n})$ for all $n \in \mathbb{N}$, then it holds a.s. that*

$$\left| J(\varphi^n) - \tilde{J}(\nu) \right| \leq C n^{-\frac{1}{2}} (\ln n) \|\lambda\|_\infty (1 + \|\lambda\|_\infty), \quad \text{for all sufficiently large } n \in \mathbb{N},$$

(3) *for all grids π of $[0, T]$ and $\varphi \sim \mathcal{R}(k, K, \lambda, \xi^\pi)$, $|\mathbb{E}[J(\varphi) - \tilde{J}(\nu)]| \leq C |\pi| \|\lambda\|_\infty^2$.*

Remark 2.1. Roughly speaking, Theorem 2.1 shows that if a randomised policy $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$ is driven by an external noise ξ with independent marginals across time, then its cost $J(\varphi)$ well represents the relaxed cost $\tilde{J}(\nu)$. Such a process ξ is closely related to the white noise in the distribution theory, and we refer the reader to Section 2.4 for a detailed discussion.

2.3 Regret analysis of episodic learning algorithms

Based on Theorem 2.1, this section analyses the episodic learning algorithms outlined in Section 2.1.

Probabilistic learning framework. We start by recalling the generic learning procedure for solving (2.1)-(2.2) with unknown θ^* : for each episode $m \in \mathbb{N}$, let $\nu^m = \mathcal{N}(k_t^m + K_t^m x, (\lambda_t^m)^2)$ be a given Gaussian relaxed policy. Here $k^m, K^m, \lambda^m : \Omega \times [0, T] \rightarrow \mathbb{R}$ are designed based on observations from previous episodes. The agent then chooses a grid π_m of $[0, T]$ and executes ν^m on π_m via a policy φ^m : for all $(\omega, t, x) \in \Omega \times [0, T] \times \mathbb{R}$,

$$\varphi^m(\omega, t, x) = k_t^m(\omega) + K_t^m(\omega)x + \lambda_t^m(\omega)\xi_t^{\pi_m}(\omega), \quad (2.17)$$

where ξ^{π_m} is a piecewise constant process on π_m generated by independent standard normal variables $(\zeta_i^m)_{i \in \mathbb{N}}$ (cf. Definition 2.2). The agent observes a trajectory of the corresponding state process X^m governed by the following dynamics (cf. (2.2)):

$$dX_t = \theta^* Z_t^m dt + \bar{\sigma}_t dW_t^m, \quad X_0 = x_0, \quad \text{with } Z_t^m = \begin{pmatrix} X_t^m \\ \varphi^m(\cdot, t, X_t^m) \end{pmatrix}, \quad (2.18)$$

where W^m is an independent Brownian motion corresponding to the m -th episode.¹ The agent then constructs an estimate of θ^* , denoted by θ_m , based on all observations $(X^n, \xi^{\pi_n})_{n=1}^m$, and solves an regularised relaxed control problem (see e.g., Approaches 1 and 2) based on θ_m and a properly chosen weight $\varrho_m > 0$. This determines the relaxed policy ν^{m+1} for the next episode.

Bayesian inference. To derive a concrete learning algorithm, we infer the parameter θ^* by a Bayesian approach as in [26]. Let $\mathcal{F}_m^{\text{ob}}$ be the observation σ -algebra generated by the state processes and the injected noises after the m -th episode:

$$\mathcal{F}_m^{\text{ob}} = \sigma\{X_t^n, \xi_t^{\pi_n} \mid t \in [0, T], n = 1, \dots, m\} \vee \mathcal{N}_0, \quad m \in \mathbb{N} \cup \{0\}, \quad (2.19)$$

where \mathcal{N}_0 is the σ -algebra generated by \mathbb{P} -null sets. By [19, Section 7.6.4], for each $m \in \mathbb{N}$, the likelihood function of θ^* with observations (X^m, ξ^{π_m}) is given by

$$\ell(\theta^* \mid X^m, \xi^{\pi_m}) \propto \exp\left(\int_0^T \frac{1}{\bar{\sigma}_s^2} \theta^* Z_s^m dX_s^m - \frac{1}{2} \int_0^T \frac{1}{\bar{\sigma}_s^2} (\theta^* Z_s^m)^2 ds\right), \quad (2.20)$$

where \propto stands for proportionality up to a constant independent of θ^* . Hence, given the initial belief that θ^* follows the prior distribution $\pi(\theta^* \mid \mathcal{F}_0^{\text{ob}}) = \mathcal{N}(\theta_0, V_0)$ for some $\theta \in \mathbb{R}^{1,2}$ and $V_0 \in \mathbb{S}_+^2$, the posterior distribution of $\pi(\theta^* \mid \mathcal{F}_m^{\text{ob}})$ after the m -th episode is given by

$$\pi(\theta^* \mid \mathcal{F}_m^{\text{ob}}) \propto \pi(\theta^* \mid \mathcal{F}_0^{\text{ob}}) \prod_{n=1}^m \ell(\theta^* \mid X^n, \xi^{\pi_n}) \propto \exp\left(-\frac{(\theta^* - \hat{\theta}_m)V_m^{-1}(\theta^* - \hat{\theta}_m)^\top}{2}\right), \quad (2.21)$$

where $\hat{\theta}_m$ and V_m are given by:

$$V_m := \left(V_0^{-1} + \sum_{n=1}^m \int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds\right)^{-1} \in \mathbb{S}_+^2, \quad (2.22a)$$

$$\hat{\theta}_m := \left(\theta_0 V_0^{-1} + \sum_{n=1}^m \left(\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n dX_s^n\right)^\top\right) V_m \in \mathbb{R}^{1 \times 2}. \quad (2.22b)$$

For simplicity, we assume that the agent has prior knowledge of the magnitude of θ^* :

H.2. Θ is a compact subset of $\mathbb{R}^{1 \times 2}$ such that $\theta^* \in \text{int}(\Theta)$.

Condition (H.2) suggests a truncated maximum a posteriori estimate θ_m as follows:

$$\theta_m := \Pi_\Theta(\hat{\theta}_m, V_m) \in \mathbb{R}^{1 \times 2}, \quad (2.23)$$

¹Without loss of generality, we assume that $(\xi^{\pi_m}, W^m)_{m \in \mathbb{N}}$ are defined on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, $(\xi^{\pi_m})_{m \in \mathbb{N}}$ are measurable with respect to the initial σ -algebra \mathcal{F}_0 , and all Brownian motions $(W^m)_{m \in \mathbb{N}}$ are independent.

where $\Pi_\Theta : \mathbb{R}^{1 \times 2} \times \mathbb{S}_+^2 \rightarrow \Theta$ is a measurable function such that $\Pi_\Theta(\theta, V) = \theta$ for all $\theta \in \Theta$ and $V \in \mathbb{S}_+^2$. Possible choices of Π_Θ include the orthogonal projection for closed and convex Θ , or a function that maximises the posterior distribution (2.21) over all $\theta^* \in \Theta$. We emphasise that the a-priori bound of θ^* is only imposed to simplify the analysis, which along with the projection step Π_Θ ensures the estimates $(\theta_m)_{m \in \mathbb{N}}$ are a-priori bounded. In the case without prior knowledge on θ^* , one can remove the projection step, and instead adopt a sufficiently long initial exploration step to guarantee a uniform bound of $(\hat{\theta}_m)_{m \in \mathbb{N}}$ with high probability (see e.g., [24, 1]).

Performance measure. In the above learning procedure, the agent solves suitable regularised relaxed control problems and executes the relaxed policies through randomisation. To measure the performance of an learning algorithm in this setting, we consider the regret of learning defined as follows (see e.g., [12, 26, 1]): for each $N \in \mathbb{N}$,

$$\text{Reg}(N) = \sum_{m=1}^N \left(J^{\theta^*}(\varphi^m) - J^{\theta^*}(\phi^{\theta^*}) \right), \quad (2.24)$$

where $J^{\theta^*}(\phi^{\theta^*})$ is the optimal cost that agent can achieve knowing the parameter θ^* (see (2.1)), and $J^{\theta^*}(\varphi^m)$ is the expected cost of the randomised policy φ^m for the m -th episode:

$$J^{\theta^*}(\varphi^m) := \mathbb{E} \left[\int_0^T f(t, X_t^m, \varphi^m(\cdot, t, X_t^m)) dt + g(X_T^m) \middle| \mathcal{F}_{m-1}^{\text{ob}} \vee \mathcal{Z}_m \right], \quad (2.25)$$

where $\mathcal{F}_{m-1}^{\text{ob}}$ is defined as in (2.19), and $\mathcal{Z}_m = \sigma\{\xi_t^{\pi_m} \mid t \in [0, T]\}$. The regret $\text{Reg}(N)$ characterises the cumulative loss from taking sub-optimal policies up to the N -th episode. Agent's aim is to construct policies whose regret grows sublinearly with respect to N .

Note that the expectation in (2.25) is only taken with respect to the Brownian motion W^m , and consequently, for each $m \in \mathbb{N}$, $J^{\theta^*}(\varphi^m)$ is a random variable depending on the realisations of the Brownian motions $(W^n)_{n=1}^{m-1}$ and the injected noises $(\xi^n)_{n=1}^m$.

Regret analysis. Given the parameter inference scheme (2.22)-(2.23), the performance of a learning algorithm now depends on the construction and execution of relaxed policies. In the sequel, we propose two algorithms based on regularised control problems in Approaches 1 and 2, and choose the regularising weights $(\varrho_m)_{m \in \mathbb{N}}$ and execution grids $(\pi_m)_{m \in \mathbb{N}}$ to minimise the algorithm regrets.

The first algorithm computes relaxed policies by solving the relaxed control problem (2.9) with exploration rewards. We summarise the algorithm as follows.

Algorithm 1: Learning with exploration reward

Input: $\theta_0 \in \mathbb{R}^{1 \times 2}$, $V_0 \in \mathbb{S}_+^2$, a truncation function Π_Θ , and a Gaussian relaxed policy ν^1 .

1 **for** $m = 1, 2, \dots$ **do**

2 Determine a grid π_m of $[0, T]$ and exercise ν^m on π_m via φ^m as in (2.17).

3 Obtain the updated estimate θ_m via (2.23).

4 Determine $\varrho_m > 0$ and compute ν^{m+1} by (2.11) with $\theta = \theta_m$ and $\varrho = \varrho_m$.

5 **end**

The following theorem chooses $(\varrho_m)_{m \in \mathbb{N}}$ and $(\pi_m)_{m \in \mathbb{N}}$ such that Algorithm 1 achieves a regret of the order $\mathcal{O}(\sqrt{N} \ln N)$ in expectation. This recovers the regret order of the phased-based algorithm with deterministic policies in [26]. Note that the additional exploration reward vanishes as the number of episodes tends to infinity.

Theorem 2.2. *Suppose (H.1) and (H.2) hold. Let $\theta_0 \in \mathbb{R}^{1 \times 2}$, $V_0 \in \mathbb{S}_+^2$, $\varrho_0 > 0$ and $\nu^1 \sim \mathcal{G}(k^{\theta_0}, K^{\theta_0}, \sqrt{\varrho_0/(2R)})$. Then there exists $c_0 > 0$ such that if one sets $\varrho_m = \varrho_0 m^{-\frac{1}{2}} \ln(m+1)$ and $|\pi_m| \leq c_0$ for all $m \in \mathbb{N}$, then there exists a constant $C \geq 0$ such that the regret of Algorithm 1 satisfies*

$$\mathbb{E}[\text{Reg}(N)] \leq C\sqrt{N} \ln N, \quad \forall N \in \mathbb{N} \cap [2, \infty).$$

Remark 2.2. The proof of Theorem 2.2 is given in Section 4.2, where the key steps are outlined at the end of this section for the reader's convenience. Note that for simplicity we choose ν^1 based on θ_0 , but the

same regret order can be achieved with $\nu^1 \sim \mathcal{G}(k^1, K^1, \lambda^1)$ for any deterministic k^1, K^1 and λ^1 . Moreover, we focus on optimising the regret order in expectation, and obtain a simple deterministic scheduling of the sampling frequency $(|\pi_m|)_{m \in \mathbb{N}}$ and regularising weights $(\varrho_m)_{m \in \mathbb{N}}$. The analysis can be extended to ensure that a similar regret bound holds with probability at least $1 - \delta$, for any given $\delta \in (0, 1)$, but $(\pi_m)_{m \in \mathbb{N}}$ and $(\varrho_m)_{m \in \mathbb{N}}$ have to be chosen depending explicitly on δ (see e.g., [24, 12, 1]). We leave a rigorous analysis of such a high-probability regret for future research.

The second algorithm computes relaxed policies with the proximal policy update (2.13). We summarise the algorithm as follows.

Algorithm 2: Learning with proximal policy update

- Input:** $\theta_0 \in \mathbb{R}^{1 \times 2}$, $V_0 \in \mathbb{S}_+^2$, a truncation function Π_Θ , and a Gaussian relaxed policy ν^1 .
- 1 **for** $m = 1, 2, \dots$ **do**
 - 2 Determine a grid π_m of $[0, T]$ and exercise ν^m on π_m via φ^m as in (2.17).
 - 3 Obtain the updated estimate θ_m via (2.23).
 - 4 Determine $\varrho_m > 0$ and compute ν^{m+1} by (2.14) with $\theta = \theta_m$ and $\varrho = \varrho_m$.
 - 5 **end**
-

The following theorem shows that Algorithm 2 achieves a similar sublinear regret as Algorithm 1. Note that unlike Algorithm 1, the regularising weight ϱ_m tends to infinity as the learning proceeds.

Theorem 2.3. *Suppose (H.1) and (H.2) hold. Let $\theta_0 \in \mathbb{R}^{1 \times 2}$, $V_0 \in \mathbb{S}_+^2$, $\varrho_0 > 0$ and $\nu^1 \sim \mathcal{G}(k^{\theta_0}, K^{\theta_0}, \varrho_0)$. Then there exists $c_0 > 0$ such that if one sets $\varrho_m = \varrho_0 m^{\frac{1}{2}} \ln(m+1)$ and $|\pi_m| \leq c_0$ for all $m \in \mathbb{N}$, then there exists a constant $C \geq 0$ such that the regret of Algorithm 2 satisfies*

$$\mathbb{E}[\text{Reg}(N)] \leq C(\sqrt{N}(\ln N) \ln(\ln N)), \quad \forall N \in \mathbb{N} \cap [3, \infty).$$

Remark 2.3. The proof of Theorem 2.3 is given in Section 4.3, whose key steps are outlined below. The argument is more involved than that of Theorem 2.2. This is due to the fact that ν^{m+1} , $m \in \mathbb{N}$, is a convex combination of all previous policies $(\nu^n)_{n=0}^m$ (cf. (2.14)), and hence the cost $J^{\theta^*}(\varphi^{m+1})$ depends explicitly on $(\theta_n)_{n=0}^m$. This memory dependence requires quantifying the precise dependence of the performance gap $J^{\theta^*}(\varphi^{m+1}) - J^{\theta^*}(\phi^{\theta^*})$ on the accuracy of $(\theta_n)_{n=0}^m$.

Sketched proofs of Theorems 2.2 and 2.3. The key idea of the regret analysis is to balance the exploration–exploitation trade-off for Algorithms 1 and 2. To this end, we decompose the regret into

$$\text{Reg}(N) = \sum_{m=1}^N \left(J^{\theta^*}(\varphi^m) - J^{\theta^*}(\phi^m) \right) + \sum_{m=1}^N \left(J^{\theta^*}(\phi^m) - J^{\theta^*}(\phi^{\theta^*}) \right), \quad (2.26)$$

where $\phi^m(t, x) = k_t^m + K_t^m x$ with (k^m, K^m) is defined in (2.17). The first term induces the exploration cost due to randomisation, and the second term describes the exploitation cost caused by the sub-optimal policies $(\phi^n)_{n=1}^N$. In particular, we prove that $\mathbb{E}[J^{\theta^*}(\varphi^m) - J^{\theta^*}(\phi^m) \mid \mathcal{F}_{m-1}^{\text{ob}}] = \mathcal{O}(\|\lambda^m\|_\infty^2 + \sqrt{|\pi_m|} \|\lambda^m\|_\infty^2)$ and $J^{\theta^*}(\phi^m) - J^{\theta^*}(\phi^{\theta^*}) = \mathcal{O}(\|K^m - K^{\theta^*}\|_\infty^2 + \|k^m - k^{\theta^*}\|_\infty^2)$. We then reduce the exploitation loss to the parameter estimate error by using Step 4 of Algorithms 1 and 2, and prove that $J^{\theta^*}(\phi^m) - J^{\theta^*}(\phi^{\theta^*}) = \mathcal{O}(|\theta_m - \theta^*|^2)$ for Algorithm 1 and $J^{\theta^*}(\phi^m) - J^{\theta^*}(\phi^{\theta^*}) = \mathcal{O}\left(\sum_{n=1}^m \frac{1}{n} \left\| \frac{\lambda_n^m}{\lambda^n} \right\|_\infty^2 |\theta_n - \theta^*|^2\right)$ for Algorithm 2 (see Proposition 4.10). We further quantify the estimation accuracy by $|\theta_m - \theta^*|^2 = \mathcal{O}\left(\frac{1}{\sum_{n=1}^m \min_{t \in [0, T]} (\lambda_t^n)^2}\right)$, provided that the mesh size $|\pi_m|$ is sufficiently small (see Section 4.1). These analyses show that the exploration cost in (2.26) increases in $(\lambda^n)_{n \in \mathbb{N}}$ and the exploitation loss in (2.26) decreases in $(\lambda^n)_{n \in \mathbb{N}}$.

Observe that for both Algorithms 1 and 2, the magnitude of λ^{m+1} is determined by the weights $(\varrho_n)_{n=1}^m$ (cf. (2.11) and (2.14)). Hence, by choosing the weights $(\varrho_m)_{m \in \mathbb{N}}$ as in Theorems 2.2 and 2.3, we can balance the exploration and exploitation and obtain algorithms with optimal regret orders (up to a logarithmic factor). \square

2.4 Discussions

Random execution and scaled white noise. The execution noise in Definition 2.2 is a properly scaled discrete-time white noise. Discretised white noise has been used in [6] for exploration. More precisely, let ϕ be a deterministic policy that is affine in the state, let \mathbb{W} be an independent Brownian motion, and for each $n \in \mathbb{N}$, let π^n be a uniform grid with mesh size $\delta_n = T/n$. Given $\varepsilon > 0$, [6] considers the randomised policy $\varphi_{n,\varepsilon} = \phi + \varepsilon \sum_{i=0}^{n-1} \frac{\mathbb{W}_{(i+1)\delta_n} - \mathbb{W}_{i\delta_n}}{\delta_n} \mathbf{1}_{[i\delta_n, (i+1)\delta_n)}$ for all large $n \in \mathbb{N}$, and the normalised cost $J(\varphi_{n,\varepsilon}) - Cn\varepsilon^2$ for some constant $C > 0$. As $\varphi_{n,\varepsilon}$ behaves similar to $\phi + \varepsilon \dot{\mathbb{W}}$ with $\dot{\mathbb{W}}$ being a white noise, the term $n\varepsilon^2$ is necessary for the finiteness of the normalised cost. Indeed, for a suitable choice of C , the normalised cost converges to $J(\phi)$ as $n \rightarrow \infty$.

Instead of fixing ε as in [6], the piecewise constant randomisation in Definition 2.2 adjusts ε according to the mesh size. To see it, let $\lambda > 0$ and for each $n \in \mathbb{N}$, let $\varepsilon_n = \lambda\sqrt{\delta_n}$. By Theorem 2.1 (or more generally Theorem 3.3), $J(\varphi_{n,\varepsilon_n})$ approximates the relaxed cost $J(\nu)$ with $\nu \sim \mathcal{N}(\phi, \lambda)$. The normalisation term $n\varepsilon_n^2 = \mathcal{O}(\lambda^2)$ corresponds to the exploration cost $J(\nu) - J(\phi)$ in (4.15) (see also [28]).

Random execution and chattering lemma. In the control theory, the chattering lemma (see e.g., [21]) ensures that there exists a sequence of strict control processes converging to a relaxed control process. However, the proof is nonconstructive and typically requires a compact action space. Also there is no explicit convergence rate.

Our work focuses on the LQ setting and provides an explicit approximation of Gaussian relaxed policies. We also quantify the precise approximation error in terms of the mesh size and the variance of the Gaussian relaxed policies.

LQ-RL with controlled diffusion. For general LQ-RL problems whose state dynamics involves controlled diffusions, the state process is no longer Gaussian. Moreover, as the noise of the state dynamics can degenerate, the Bayesian estimation (2.22) may not be well-defined. In this case, one has to derive learning algorithms in a problem dependent way.

For instance, let ϕ be a given policy and consider the state process X^ϕ satisfying the following one-dimensional dynamics:

$$dX_t = (AX_t + B\phi(t, X_t)) dt + (CX_t + D\phi(t, X_t) + \bar{\sigma}) dW_t, \quad t \in [0, T]; \quad X_0 = x_0. \quad (2.27)$$

Note that by observing X^ϕ continuously, the agent can infer the constants $(C, D, \bar{\sigma})$ using finite episodes. Indeed, the agent can compute the process $Y_t^\phi = CX_t^\phi + D\phi(t, X_t^\phi) + \bar{\sigma}$, $t \in [0, T]$, from the rate of increase of the quadratic variation of X^ϕ . Then the agent can recover D from the random variables $Y_0^{\phi^{(i)}}$, $i = 1, 2, 3$, with constant policies $\phi^{(1)} \equiv -1$, $\phi^{(2)} \equiv 0$, and $\phi^{(3)} \equiv 1$. The constants C and $\bar{\sigma}$ can further be inferred from observed paths of Y^ϕ , by executing a policy ϕ such that paths of X^ϕ are non-constant almost surely.

Given $(C, D, \bar{\sigma})$, the agent may eliminate the observation noise and simplify the learning process. To see it, suppose that $D, \bar{\sigma} \neq 0$. Then the agent will estimate (A, B) by executing the policy $\phi(t, x) = -D^{-1}(Cx + \bar{\sigma})$. This leads to a deterministic state dynamics: $dX_t^\phi = \theta Z_t^\phi dt$, with $\theta = (A, B)$ and $Z_t^\phi := \begin{pmatrix} X_t^\phi \\ \phi(t, X_t^\phi) \end{pmatrix}$. As $\bar{\sigma}$ is non-zero and X^ϕ is in general non-constant, X^ϕ and $\phi(\cdot, X^\phi)$ are linearly independent, and $\int_0^T Z_t^\phi (Z_t^\phi)^\top dt$ is invertible. This allows to identify θ using one episode. A complete study of LQ-RL with controlled diffusion is left to future research.

3 Execution gap analysis for Gaussian relaxed policies

This section analyses the execution of a Gaussian relaxed policy via piecewise constant randomisation as proposed in Section 2.2. The analysis is carried out in a general LQ setting, where both the drift and diffusion coefficients of the state dynamics are time-dependent and under controlled, and the cost functions are possibly nonconvex.

Let $b : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $\sigma : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $f : [0, T] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ and $g : \mathbb{R} \rightarrow \mathbb{R}$ be such that for all $(t, x, a) \in [0, T] \times \mathbb{R} \times \mathbb{R}$,

$$b(t, x, a) = A_t x + B_t a + \bar{b}_t, \quad \sigma(t, x, a) = C_t x + D_t a + \bar{\sigma}_t, \quad (3.1)$$

$$f(t, x, a) = Q_t x^2 + 2S_t x a + R_t a^2 + p_t x + q_t a, \quad g(x) = Mx^2 + mx. \quad (3.2)$$

The coefficients satisfy the following boundedness condition:

H.3. $T > 0$, $x_0 \in \mathbb{R}$, $A, B, \bar{b}, C, D, \bar{\sigma}, Q, S, R, p, q \in L^\infty([0, T]; \mathbb{R})$, and $M, m \in \mathbb{R}$.

For any Gaussian relaxed policy ν (cf. Definition 2.1), consider the controlled dynamics:

$$dX_t = \int_{\mathbb{R}} b(t, X_t, a) \nu(t, X_t; da) dt + \sqrt{\int_{\mathbb{R}} \sigma(t, X_t, a)^2 \nu(t, X_t; da)} dW_t, \quad t \in [0, T]; \quad X_0 = x_0, \quad (3.3)$$

where b and σ are defined in (3.1). By (3.1), for each $\nu \sim \mathcal{G}(k, K, \lambda)$ and $(t, x) \in [0, T] \times \mathbb{R}$,

$$\begin{aligned} \int_{\mathbb{R}} b(t, x, a) \nu(t, x; da) &= A_t x + B_t(k_t + K_t x) + \bar{b}_t, \\ \int_{\mathbb{R}} \sigma(t, x, a)^2 \nu(t, x; da) &= (C_t x + D_t(k_t + K_t x) + \bar{\sigma}_t)^2 + \lambda_t^2 D_t^2, \end{aligned}$$

which along with (H.3) shows that the coefficients of (3.3) are uniform Lipschitz continuous in x , and hence (3.3) admits a unique strong solution X^ν such that $\mathbb{E}[\sup_{t \in [0, T]} |X_t^\nu|^p] < \infty$ for any $p \geq 1$. The expected cost of ν is then defined by

$$\tilde{J}(\nu) := \mathbb{E} \left[\int_0^T \int_{\mathbb{R}} f(t, X_t^\nu, a) \nu(t, X_t^\nu; da) dt + g(X_T^\nu) \right], \quad (3.4)$$

with f and g defined in (3.2). Note that $\tilde{J}(\nu) < \infty$ due to the square-integrability of X^ν .

We then introduce a class of randomised policies with Gaussian noises, which includes as a special case the piecewise constant randomised policies defined in Definition 2.2.

Definition 3.1. For each $k, K \in L^\infty([0, T]; \mathbb{R})$, $\lambda \in L^\infty([0, T]; \mathbb{R}_{\geq 0})$, and measurable process $\xi : \Omega \times [0, T] \rightarrow \mathbb{R}$ satisfying for all $t \in [0, T]$, ξ_t is an \mathcal{F}_0 -measurable standard normal random variable, we define a randomised policy with Gaussian noise $\varphi : \Omega \times [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$, denoted by $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$, such that for all $(\omega, t, x) \in \Omega \times [0, T] \times \mathbb{R}$, $\varphi(\omega, t, x) = k_t + K_t x + \lambda_t \xi_t(\omega)$.

Compared with Definition 2.2, Definition 3.1 only imposes a Gaussianity of ξ_t for each $t \in [0, T]$, without requiring an independence of (ξ_t, ξ_s) for different $t, s \in [0, T]$. In the sequel, we simply refer to $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$ as a randomised policy if no confusion occurs.

For any randomised policy φ , consider the controlled dynamics:

$$dX_t = b(t, X_t, \varphi(\cdot, t, X_t)) dt + \sigma(t, X_t, \varphi(\cdot, t, X_t)) dW_t, \quad t \in [0, T]; \quad X_0 = x_0, \quad (3.5)$$

where b and σ are defined in (3.1), and $\varphi(\cdot, t, X_t)$ emphasises the additional randomness of the policy. By Definition 3.1 and (H.3), the coefficients of (3.5) are progressively measurable, uniformly Lipschitz continuous in x , and satisfies for all $p \geq 2$, $\mathbb{E}[\int_0^T |b(t, 0, \varphi(\cdot, t, 0))|^p dt] + \mathbb{E}[\int_0^T |\sigma(t, 0, \varphi(\cdot, t, 0))|^p dt] < \infty$. This implies that (3.3) admits a unique strong solution X^φ such that $\mathbb{E}[\sup_{t \in [0, T]} |X_t^\varphi|^p] < \infty$ for all $p \geq 2$ (see [32, Theorem 3.4.3]). The corresponding expected cost of $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$ (with fixed noise ξ) is then defined as

$$J(\varphi) := \mathbb{E} \left[\int_0^T f(t, X_t^\varphi, \varphi(\cdot, t, X_t^\varphi)) dt + g(X_T^\varphi) \middle| \mathcal{F}_0 \right], \quad (3.6)$$

where f and g are given in (3.2).

The following lemma quantifies the difference between $\tilde{J}(\nu)$ with $\nu \sim \mathcal{G}(k, K, \lambda)$ and $J(\varphi)$ with $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$. The proof follows from a straightforward but notationally cumbersome computation, whose detail is given in Appendix A.

Lemma 3.1. *Suppose (H.3) holds. For each $\nu \sim \mathcal{G}(k, K, \lambda)$ and $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$, let $\tilde{J}(\nu)$ and $J(\varphi)$ be defined by (3.4) and (3.6), respectively. Then for all $\nu \sim \mathcal{G}(k, K, \lambda)$ and $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$,*

$$J(\varphi) - \tilde{J}(\nu) = \int_0^T h_{1,t} \lambda_t \xi_t dt + \int_0^T h_{2,t} \lambda_t^2 (\xi_t^2 - 1) dt + \int_0^T \int_0^T h_{3,t,r} \lambda_r \lambda_t \xi_r \xi_t dr dt, \quad (3.7)$$

where $h_1, h_2 : [0, T] \rightarrow \mathbb{R}$ and $h_3 : [0, T]^2 \rightarrow \mathbb{R}$ are some bounded functions whose sup-norms depend only on the sup-norms of k, K and the coefficients in (H.3).

Now we are ready to quantify the performance of a piecewise constant independent policy in terms of the mesh size $|\pi|$. The proof relies on estimating (3.7) using concentration inequalities of independent normal random variables in the following lemma (see Appendix A for a proof).

Lemma 3.2. *There exists a constant $C \geq 0$ such that for all independent standard normal random variables $(\zeta_i)_{i \in \mathbb{N}}$, for all $N \in \mathbb{N}$, $(\rho_i)_{i=1}^N \subset \mathbb{R}$, $(\beta_{ij})_{i,j=1}^N \subset \mathbb{R}$ and $\delta > 0$,*

- (1) $\mathbb{P} \left(\left| \sum_{i=1}^N \rho_i \zeta_i \right| \geq C \|\rho\|_2 \left(1 + \sqrt{\ln \left(\frac{1}{\delta} \right)} \right) \right) \leq \delta$, with $\|\rho\|_2 = \sqrt{\sum_{i=1}^N \rho_i^2}$,
- (2) $\mathbb{P} \left(\left| \sum_{i=1}^N \rho_i (\zeta_i^2 - 1) \right| \geq C \|\rho\|_2 \left(1 + \ln \left(\frac{1}{\delta} \right) \right) \right) \leq \delta$, with $\|\rho\|_2 = \sqrt{\sum_{i=1}^N \rho_i^2}$,
- (3) $\mathbb{P} \left(\left| \sum_{i,j=1, i \neq j}^N \beta_{ij} \zeta_i \zeta_j \right| \geq C \|\beta\|_2 \left(1 + \ln \left(\frac{1}{\delta} \right) \right) \right) \leq \delta$, with $\|\beta\|_2 = \sqrt{\sum_{i,j=1, i \neq j}^N \beta_{ij}^2}$.

Theorem 3.3. *Suppose (H.3) holds and let $\nu \sim \mathcal{G}(k, K, \lambda)$. Then there exists a constant $C \geq 0$, depending only on the sup-norms of k, K and the coefficients in (H.3), such that for all grids π of $[0, T]$, all policies $\varphi \sim \mathcal{R}(k, K, \lambda, \xi^\pi)$ (cf. Definition 2.2) and all $\delta > 0$,*

$$\mathbb{P} \left(\left| J(\varphi) - \tilde{J}(\nu) \right| \geq C \sqrt{|\pi|} \|\lambda\|_\infty \left(1 + \|\lambda\|_\infty \left(1 + \ln \left(\frac{1}{\delta} \right) \right) \right) \right) \leq \delta, \quad (3.8)$$

and $|\mathbb{E}[J(\varphi) - \tilde{J}(\nu)]| \leq C |\pi| \|\lambda\|_\infty^2$.

Proof. Throughout this proof, let C be a generic constant independent of λ, ξ, δ and π , but may possibly depend on the sup-norms of the coefficients in (H.3) and the functions k and K . Observe that the functions h_i , $i = 1, \dots, 3$, in Lemma 3.1 are uniformly bounded by some constant $C \geq 0$. Let $\pi = \{0 = t_0 < \dots < t_N = T\}$ be a partition of $[0, T]$ with mesh size $|\pi|$, and let $\varphi \sim \mathcal{R}(\mu, \lambda, \xi^\pi)$ with $\xi_t^\pi = \sum_{i=0}^{N-1} \zeta_i \mathbf{1}_{t_i \leq t < t_{i+1}}$. We omit the superscript π in ξ^π for notational simplicity.

To prove (3.8), it suffices to upper bound all terms of (3.7) in high probability. The definition of ξ implies that $\int_0^T h_{1,t} \lambda_t \xi_t dt = \sum_{i=0}^{N-1} \zeta_i \left(\int_{t_i}^{t_{i+1}} h_{1,t} \lambda_t dt \right)$. By Lemma 3.2 Item (1), it holds with probability at least $1 - \delta$ that

$$\left| \int_0^T h_{1,t} \lambda_t \xi_t dt \right| \leq C \left(1 + \sqrt{\ln \left(\frac{1}{\delta} \right)} \right) \sqrt{\sum_{i=0}^{N-1} \left(\int_{t_i}^{t_{i+1}} h_{1,t} \lambda_t dt \right)^2} \leq C \sqrt{T} \sqrt{|\pi|} \|\lambda\|_\infty \left(1 + \sqrt{\ln \left(\frac{1}{\delta} \right)} \right). \quad (3.9)$$

Also applying Lemma 3.2 Item (2) to $\int_0^T h_{2,t} \lambda_t^2 (\xi_t^2 - 1) dt$ implies that with probability at least $1 - \delta$,

$$\left| \int_0^T h_{2,t} \lambda_t^2 (\xi_t^2 - 1) dt \right| \leq C \sqrt{T} \sqrt{|\pi|} \|\lambda\|_\infty \left(1 + \ln \left(\frac{1}{\delta} \right) \right). \quad (3.10)$$

It now remains to quantify the last term of (3.7). Observe that

$$\begin{aligned} & \int_0^T \int_0^T h_{3,t,r} \lambda_r \lambda_t \xi_r \xi_t dr dt \\ &= \sum_{i,j=0; i \neq j}^{N-1} \zeta_i \zeta_j \left(\int_{t_i}^{t_{i+1}} \int_{t_j}^{t_{j+1}} h_{3,t,r} \lambda_r \lambda_t dr dt \right) + \sum_{i=0}^{N-1} \zeta_i^2 \left(\int_{t_i}^{t_{i+1}} \int_{t_i}^{t_{i+1}} h_{3,t,r} \lambda_r \lambda_t dr dt \right). \end{aligned}$$

By Lemma 3.2 Item (3), it holds with probability at least $1 - \delta$,

$$\begin{aligned} & \left| \sum_{i,j=0;i \neq j}^{N-1} \zeta_i \zeta_j \left(\int_{t_i}^{t_{i+1}} \int_{t_j}^{t_{j+1}} h_{3,t,r} \lambda_r \lambda_t \, dr dt \right) \right| \\ & \leq C \|\lambda\|_\infty^2 \left(1 + \ln\left(\frac{1}{\delta}\right)\right) \sqrt{\sum_{i,j=0;i \neq j}^{N-1} (t_{i+1} - t_i)^2 (t_{j+1} - t_j)^2} \leq C \|\lambda\|_\infty^2 \left(1 + \ln\left(\frac{1}{\delta}\right)\right) T |\pi|, \end{aligned}$$

while by Lemma 3.2 Item (2), it holds with probability at least $1 - \delta$,

$$\begin{aligned} & \left| \sum_{i=0}^{N-1} \zeta_i^2 \left(\int_{t_i}^{t_{i+1}} \int_{t_i}^{t_{i+1}} h_{3,t,r} \lambda_r \lambda_t \, dr dt \right) \right| \\ & \leq \left| \sum_{i=0}^{N-1} (\zeta_i^2 - 1) \left(\int_{t_i}^{t_{i+1}} \int_{t_i}^{t_{i+1}} h_{3,t,r} \lambda_r \lambda_t \, dr dt \right) \right| + \left| \sum_{i=0}^{N-1} \left(\int_{t_i}^{t_{i+1}} \int_{t_i}^{t_{i+1}} h_{3,t,r} \lambda_r \lambda_t \, dr dt \right) \right| \\ & \leq C \|\lambda\|_\infty^2 \left(1 + \ln\left(\frac{1}{\delta}\right)\right) \sqrt{\sum_{i=0}^{N-1} (t_{i+1} - t_i)^4} + C \|\lambda\|_\infty^2 T |\pi| \leq C \|\lambda\|_\infty^2 \left(1 + \ln\left(\frac{1}{\delta}\right)\right) |\pi|. \end{aligned}$$

Summarising these inequalities shows that with probability at least $1 - \delta$,

$$\left| \int_0^T \int_0^T h_{3,t,r} \lambda_r \lambda_t \xi_r \xi_t \mathbf{1}_{r \leq t} \, dr dt \right| \leq C |\pi| \|\lambda\|_\infty^2 \left(1 + \ln\left(\frac{1}{\delta}\right)\right). \quad (3.11)$$

Combining (3.9), (3.10) and (3.11) with the inequality that $\sqrt{\ln\left(\frac{1}{\delta}\right)} \leq 1 + \ln\left(\frac{1}{\delta}\right)$ for all $\delta \geq 0$ leads to the estimate (3.8).

Furthermore, by Lemma 3.1 and the fact that $\xi_t \sim \mathcal{N}(0, 1)$ for all $t \in [0, T]$ and $|\mathbb{E}[\xi_t \xi_s]| \leq \mathbf{1}_{|t-s| \leq |\pi|}$, it follows from (3.7) that

$$\begin{aligned} |\mathbb{E}[J(\varphi) - \tilde{J}(\nu)]| &= \left| \mathbb{E} \left[\int_0^T \int_0^T h_{3,t,r} \lambda_r \lambda_t \xi_r \xi_t \, dr dt \right] \right| \leq C \|\lambda\|_\infty^2 \int_0^T \int_0^T |\mathbb{E}[\xi_r \xi_t]| \, dr dt \\ &= C \|\lambda\|_\infty^2 \int_0^T \int_{t-|\pi|}^{t+|\pi|} \, dr dt \leq C \|\lambda\|_\infty^2 \int_0^T 2|\pi| \, dt \leq C |\pi| \|\lambda\|_\infty^2. \end{aligned}$$

This proves the desired estimate of $|\mathbb{E}[J(\varphi) - \tilde{J}(\nu)]|$. \square

Theorem 3.3 implies the following almost sure convergence with uniform grids.

Corollary 3.4. *Suppose (H.3) holds and let $\nu \sim \mathcal{G}(k, K, \lambda)$. For each $n \in \mathbb{N}$, let $\pi^n = \{i \frac{T}{n}\}_{i=0}^n$, and $\varphi^n \sim \mathcal{R}(k, K, \lambda, \xi^{\pi^n})$ (cf. Definition 2.2). Then there exists a constant $C \geq 0$, depending only on the sup-norms of k, K and the coefficients in (H.3), such that for \mathbb{P} -a.s. $\omega \in \Omega$, there exists $N_\omega \in \mathbb{N}$ such that*

$$\left| J(\varphi^n) - \tilde{J}(\nu) \right| \leq C n^{-\frac{1}{2}} (\ln n) \|\lambda\|_\infty (1 + \|\lambda\|_\infty), \quad \forall n \geq N_\omega.$$

Proof. Let C be the same constant as that in Theorem 3.3, and for each $n \in \mathbb{N}$, let $\delta_n = \frac{1}{n^2}$, and $A_n = \left| J(\varphi^n) - \tilde{J}(\nu) \right| \geq C \sqrt{|\pi_n|} \|\lambda\|_\infty (1 + \|\lambda\|_\infty) \left(1 + \ln\left(\frac{1}{\delta_n}\right)\right)$. Theorem 3.3 implies $\sum_{n=1}^\infty \mathbb{P}(A_n) \leq \sum_{n=1}^\infty \delta_n < \infty$, which along with the Borel–Cantelli lemma shows that $\mathbb{P}(\limsup_{n \rightarrow \infty} A_n) = 0$. Consequently, for \mathbb{P} -a.s. $\omega \in \Omega$, there exists $N_\omega \in \mathbb{N}$, such that for all $n \geq N_\omega$,

$$\left| J(\varphi^n) - \tilde{J}(\nu) \right| \leq C \sqrt{|\pi_n|} \|\lambda\|_\infty (1 + \|\lambda\|_\infty) \left(1 + \ln\left(\frac{1}{\delta_n}\right)\right) \leq C n^{-\frac{1}{2}} (\ln n) \|\lambda\|_\infty (1 + \|\lambda\|_\infty). \quad \square$$

4 Regret analysis for episodic learning

4.1 Analysis of piecewise constant randomised policies

This section analyses a sequence of piecewise constant randomised policies that is adapted to the observation filtration. The precise definition of these policies are given in Setting 4.1, which includes as special cases the randomised policies generated by Algorithms 1 and 2.

Setting 4.1. Let $\theta_0 \in \mathbb{R}^{1 \times 2}$, $V_0 \in \mathbb{S}_+^2$, and for each $m \in \mathbb{N}$, let $K^m, k^m \in L^\infty(\Omega \times [0, T]; \mathbb{R})$, let $\lambda^m \in L^\infty([0, T]; \mathbb{R}_{\geq 0})$, let $\pi_m = (t_i^m)_{i=0}^{N_m}$ be a grid of $[0, T]$, and let $\varphi^m : \Omega \times [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be defined as in (2.17). Assume that $L := \sup_{m \in \mathbb{N}} (\|K^m\|_{L^\infty} + \|k^m\|_{L^\infty} + \|\lambda^m\|_\infty) < \infty$, and for all $m \in \mathbb{N}$, K^m and k^m are $\mathcal{F}_{m-1}^{\text{ob}} \otimes \mathcal{B}([0, T])$ -measurable, with $\mathcal{F}_{m-1}^{\text{ob}}$ defined by (2.19).

The following proposition establishes a subexponential concentration behaviour of $(V_m^{-1})_{m \in \mathbb{N}}$.

Proposition 4.1. *Suppose (H.1) and Setting 4.1 hold. Then there exists $C \geq 0$, depending only on the coefficients in (H.1) and L in Setting 4.1, such that for all $m \in \mathbb{N}$ and $\delta > 0$,*

$$\mathbb{P} \left(\left| V_m^{-1} - V_0^{-1} - \sum_{n=1}^m \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} Z_t^n (Z_t^n)^\top dt \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right| \geq C \max \left(\ln \left(\frac{2}{\delta} \right), \sqrt{m \ln \left(\frac{2}{\delta} \right)} \right) \right) \leq \delta,$$

Proof. Throughout this proof, let C be a generic constant depending only on the coefficients in (H.1) and L in Setting 4.1. For each $q \in [1, 2]$ and σ -algebra \mathcal{G} , let $\|\cdot\|_{q, \mathcal{G}} : \Omega \rightarrow [0, \infty]$ be the conditional Orlicz norm defined in [26, Definition 4.1], such that for all random variables $X : \Omega \rightarrow \mathbb{R}$,

$$\|X\|_{q, \mathcal{G}} := \mathcal{G}\text{-ess inf} \{ Y \in L^0(\mathcal{G}; (0, \infty)) \mid \mathbb{E} [\exp(|X|^q / Y^q) | \mathcal{G}] \leq 2 \},$$

where $L^0(\mathcal{G}; (0, \infty))$ is the set of all $\mathcal{G}/\mathcal{B}((0, \infty))$ -measurable functions $Y : \Omega \rightarrow (0, \infty)$.

We first prove that there exists a constant $C \geq 0$ such that $\|(\int_0^T |X_t^m|^2 dt)^{1/2}\|_{2, \mathcal{F}_{m-1}^{\text{ob}}} \leq C$ for all $m \in \mathbb{N}$. Observe from (2.18) that, for each $m \in \mathbb{N}$, $X^m = \mathcal{X}^m + \mathcal{Y}^m$, where

$$\begin{aligned} d\mathcal{X}_t^m &= (A^* \mathcal{X}_t^m + B^*(K_t^m \mathcal{X}_t^m + k_t^m)) dt + \bar{\sigma}_t dW_t^m, \quad t \in [0, T]; \quad \mathcal{X}_0^m = x_0, \\ d\mathcal{Y}_t^m &= (A^* \mathcal{Y}_t^m + B^*(K_t^m \mathcal{Y}_t^m + \lambda_t^m \xi_t^{\pi_m})) dt, \quad t \in [0, T]; \quad \mathcal{Y}_0^m = 0. \end{aligned} \quad (4.1)$$

By [26, Proposition 4.7], $\|(\int_0^T |\mathcal{X}_t^m|^2 dt)^{1/2}\|_{2, \mathcal{F}_{m-1}^{\text{ob}}} \leq C$ for all $m \in \mathbb{N}$. Moreover, let $\Phi^m : \Omega \times [0, T] \rightarrow \mathbb{R}$ be such that for all $(\omega, t) \in \Omega \times [0, T]$, $d\Phi_t^m(\omega) = (A^* + B^* K_t^m(\omega)) \Phi_t^m(\omega) dt$ and $\Phi_0^m(\omega) = 1$, then $\mathcal{Y}_t^m = \Phi_t^m \int_0^t (\Phi_s^m)^{-1} B^* \lambda_s^m \xi_s^{\pi_m} ds$ for all $t \in [0, T]$. Observe that Φ^m is invertible $d\mathbb{P} \otimes dt$ a.e., and by the uniform boundedness of $(K^m)_{m \in \mathbb{N}}$, $\|\Phi^m\|_{L^\infty} + \|(\Phi^m)^{-1}\|_{L^\infty} \leq C$ for all $m \in \mathbb{N}$. Hence, by Hölder's inequality,

$$\left(\int_0^T |\mathcal{Y}_t^m|^2 dt \right)^{\frac{1}{2}} \leq C \sup_{t \in [0, T]} |\mathcal{Y}_t^m| \leq C \sup_{t \in [0, T]} \int_0^t |\lambda_s^m \xi_s^{\pi_m}| ds \leq C \|\lambda^m\|_\infty \left(\int_0^T |\xi_s^{\pi_m}|^2 ds \right)^{\frac{1}{2}},$$

Assume without loss of generality that $\|\lambda^m\|_\infty > 0$. By the convexity of $x \mapsto e^x$ and Jensen's inequality, for a sufficiently large $\tilde{C} > 0$,

$$\begin{aligned} \mathbb{E} \left[\exp \left(\frac{1}{\tilde{C}^2 \|\lambda^m\|_\infty^2} \int_0^T |\mathcal{Y}_t^m|^2 dt \right) \middle| \mathcal{F}_{m-1}^{\text{ob}} \right] &\leq \mathbb{E} \left[\exp \left(\int_0^T \frac{C^2}{\tilde{C}^2} |\xi_t^{\pi_m}|^2 dt \right) \right] \\ &\leq \frac{1}{T} \mathbb{E} \left[\int_0^T \exp \left(\frac{TC^2}{\tilde{C}^2} |\xi_t^{\pi_m}|^2 \right) dt \right] \leq \mathbb{E} \left[\exp \left(\frac{TC^2}{\tilde{C}^2} |\xi_0^{\pi_m}|^2 \right) \right] \leq 2, \end{aligned}$$

where we used that $(\xi_t^{\pi_m})_{t \in [0, T]}$ are standard normal random variables independent of $\mathcal{F}_{m-1}^{\text{ob}}$. Consequently, $\|(\int_0^T |\mathcal{Y}_t^m|^2 dt)^{\frac{1}{2}}\|_{2, \mathcal{F}_{m-1}^{\text{ob}}} \leq \tilde{C}$, which implies that $\|(\int_0^T |X_t^m|^2 dt)^{1/2}\|_{2, \mathcal{F}_{m-1}^{\text{ob}}} \leq C$ for all $m \in \mathbb{N}$.

Observe that for all $m \in \mathbb{N}$ and $t \in [0, T]$, $\varphi^m(\cdot, t, X_t^m) = K_t^m X_t^m + k_t^m + \lambda_t^m \xi_t^{\pi^m}$. By the uniform boundedness of K^m and k^m , $\|(\int_0^T |\varphi^m(\cdot, t, X_t^m)|^2 dt)\|_{2, \mathcal{F}_{m-1}^{\text{ob}}} \leq C$ for all $m \in \mathbb{N}$, which along with $\bar{\sigma}^{-1} \in L^\infty([0, T]; \mathbb{R})$ and [26, Proposition 4.3] implies that

$$\begin{aligned} \left\| \int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^m (Z_s^m)^\top ds \right\|_{1, \mathcal{F}_{m-1}^{\text{ob}}} &\leq C \left\| \int_0^T \begin{pmatrix} X_s^m \\ \varphi^m(\cdot, s, X_s^m) \end{pmatrix} \begin{pmatrix} X_s^m \\ \varphi^m(\cdot, s, X_s^m) \end{pmatrix}^\top ds \right\|_{1, \mathcal{F}_{m-1}^{\text{ob}}} \\ &\leq C \left(1 + \left\| \left(\int_0^T |X_t^m|^2 dt \right) \right\|_{2, \mathcal{F}_{m-1}^{\text{ob}}}^2 + \left\| \left(\int_0^T |\varphi^m(\cdot, t, X_t^m)|^2 dt \right) \right\|_{2, \mathcal{F}_{m-1}^{\text{ob}}}^2 \right) \leq C. \end{aligned}$$

As $V_m^{-1} = V_0^{-1} + \sum_{n=1}^m \int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds$ and $\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^m (Z_s^m)^\top ds$ is $\mathcal{F}_m^{\text{ob}}$ -measurable, by the Bernstein inequality for martingale difference sequence in [26, Proposition 4.4], for all $m \in \mathbb{N}$ and $\varepsilon > 0$,

$$\mathbb{P} \left(\left| V_m^{-1} - V_0^{-1} - \sum_{n=1}^m \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right| \geq m\varepsilon \right) \leq 2 \exp(-mC \min(\varepsilon^2, \varepsilon)).$$

The desired result follows by substituting ε such that $2 \exp(-mC \min(\varepsilon^2, \varepsilon)) = \delta$. \square

We now establish upper and lower bounds of $\sum_{n=1}^m \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds \middle| \mathcal{F}_{n-1}^{\text{ob}} \right]$ in terms of m .

Proposition 4.2. *Suppose (H.1) and Setting 4.1 hold. Then there exists a constant $C > 0$, depending only on the coefficients in (H.1) and L in Setting 4.1, such that*

$$(1) \text{ for all } m \in \mathbb{N}, \sum_{n=1}^m \left\| \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} Z_t^n (Z_t^n)^\top dt \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right\| \leq Cm.$$

$$(2) \text{ if } |\pi^m| \|\lambda^m\|_\infty^2 \leq \frac{1}{C} \min_{t \in [0, T]} ((\lambda_t^m)^2 \wedge 1) \text{ for all } m \in \mathbb{N}, \text{ then}$$

$$\sum_{n=1}^m \Lambda_{\min} \left(\mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} Z_t^n (Z_t^n)^\top dt \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right) \geq \frac{1}{C} \sum_{n=1}^m \min_{t \in [0, T]} ((\lambda_t^n)^2 \wedge 1), \quad \forall m \in \mathbb{N},$$

where $a \wedge 1 = \min(a, 1)$ for all $a \in \mathbb{R}$.

Proof. For all $m \in \mathbb{N}$, by the definition of Z^m in (2.18), the measurability conditions of K^m and k^m and the fact that $\mathbb{E}[\xi_t^{\pi^m} | \mathcal{F}_{m-1}^{\text{ob}}] = 0$ for all $t \in [0, T]$ implies that for $d\mathbb{P}$ -a.s.,

$$\begin{aligned} \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} Z_t^m (Z_t^m)^\top dt \middle| \mathcal{F}_{m-1}^{\text{ob}} \right] (\omega) &= \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} \begin{pmatrix} X_t^m \\ K_t^m X_t^m + k_t^m + \lambda_t^m \xi_t^{\pi^m} \end{pmatrix} \begin{pmatrix} X_t^m \\ K_t^m X_t^m + k_t^m + \lambda_t^m \xi_t^{\pi^m} \end{pmatrix}^\top dt \middle| \mathcal{F}_{m-1}^{\text{ob}} \right] (\omega) \\ &= \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} \left(\begin{pmatrix} \tilde{X}_t^{m, \omega} & 0 \\ 0 & K_t^m(\omega) \tilde{X}_t^{m, \omega} \end{pmatrix} + \begin{pmatrix} 0 & \tilde{X}_t^{m, \omega} \lambda_t^m \xi_t^{\pi^m} \\ 0 & K_t^m(\omega) \tilde{X}_t^{m, \omega} \lambda_t^m \xi_t^{\pi^m} \end{pmatrix} \right) dt \right] \end{aligned} \quad (4.2)$$

$$+ \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} \left(\begin{pmatrix} \tilde{X}_t^{m, \omega} & 0 \\ K_t^m(\omega) \tilde{X}_t^{m, \omega} + k_t^m(\omega) \end{pmatrix} \begin{pmatrix} \tilde{X}_t^{m, \omega} & 0 \\ K_t^m(\omega) \tilde{X}_t^{m, \omega} + k_t^m(\omega) \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & (\lambda_t^m)^2 \end{pmatrix} \right) dt \right] \quad (4.3)$$

where $\tilde{X}^{m, \omega}$ is the state process X^m conditioned on $\mathcal{F}_{m-1}^{\text{ob}}$, such that \mathbb{P} -a.s. ω and for all $t \in [0, T]$,

$$\tilde{X}_t^{m, \omega}(\cdot) = \Phi_t^m(\omega) x_0 + \Phi_t^m(\omega) \int_0^t (\Phi_s^m(\omega))^{-1} \left(B^*(k_s^m(\omega) + \lambda_s^m \xi_s^{\pi^m}(\cdot)) \right) ds + \bar{\sigma}_s dW_s^m(\cdot), \quad (4.4)$$

with $d\Phi_t^m(\omega) = (A^* + B^* K_t^m(\omega)) \Phi_t^m(\omega) dt$ and $\Phi_0^m(\omega) = 1$. By the uniform boundedness of K^m , k^m and λ^m , $\mathbb{E} \left[\sup_{t \in [0, T]} |\tilde{X}_t^{m, \omega}|^2 \right] \leq C$ for all $m \in \mathbb{N}$. Hence, applying the Cauchy-Schwarz inequality to (4.2)-(4.3) and using the Gaussianity of $(\xi_t^{\pi^m})_{t \in [0, T]}$ yield Item (1).

To prove Item (2), we first establish a lower bound of (4.2). By (4.4) and the independence between ξ^{π_m} and W^m , for all $t \in [0, T]$,

$$\begin{aligned} \left| \mathbb{E}[\tilde{X}_t^{m,\omega} \lambda_t^m \xi_t^{\pi_m}] \right| &= \left| \lambda_t^m \Phi_t^m(\omega) \int_0^t (\Phi_s^m(\omega))^{-1} \lambda_s^m \mathbb{E}[\xi_s^{\pi_m} \xi_t^{\pi_m}] ds \right|. \\ \left| \mathbb{E}[K_t^m(\omega) \tilde{X}_t^{m,\omega} \lambda_t^m \xi_t^{\pi_m}] \right| &= \left| K_t^m(\omega) \lambda_t^m \Phi_t^m(\omega) \int_0^t (\Phi_s^m(\omega))^{-1} \lambda_s^m \mathbb{E}[\xi_s^{\pi_m} \xi_t^{\pi_m}] ds \right|. \end{aligned}$$

Note that $\mathbb{E}[\xi_s^{\pi_m} \xi_t^{\pi_m}]$ is 1 if there exists i_0 such that $s, t \in [t_{i_0}^m, t_{i_0+1}^m)$, and is zero otherwise. By the uniform boundedness of K^m , there exists $C \geq 0$ such that for all $m \in \mathbb{N}$,

$$\Lambda_{\min} \left(\mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} \left(\begin{pmatrix} \tilde{X}_t^{m,\omega} \lambda_t^m \xi_t^{\pi_m} & K_t^m(\omega) \tilde{X}_t^{m,\omega} \lambda_t^m \xi_t^{\pi_m} \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 & \tilde{X}_t^{m,\omega} \lambda_t^m \xi_t^{\pi_m} \\ 0 & K_t^m(\omega) \tilde{X}_t^{m,\omega} \lambda_t^m \xi_t^{\pi_m} \end{pmatrix} \right) dt \right] \right) \geq -C |\pi^m| \|\lambda^m\|_\infty^2. \quad (4.5)$$

We then prove that the minimum eigenvalue of (4.3) is bounded away from zero. Observe that for each $K \in \mathbb{R}$, a direct computation shows that all eigenvalues of the matrix $\Psi(K) := \begin{pmatrix} 1 \\ K \end{pmatrix} \begin{pmatrix} 1 \\ K \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$ are positive, which along with the continuity of the map $\mathbb{S}^2 \ni S \mapsto \Lambda_{\min}(S) \in \mathbb{R}$ implies that $c_0 := \inf_{|K| \leq L} \Lambda_{\min}(\Psi(K)) > 0$, which the constant L in Setting 4.1. Then for all $m \in \mathbb{N}$ and $t \in [0, T]$,

$$\begin{aligned} \Lambda_{\min} &\left(\mathbb{E} \left[\frac{1}{\bar{\sigma}_t^2} \left(\begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix} \begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & (\lambda_t^m)^2 \end{pmatrix} \right) \right] \right) \\ &\geq \frac{1}{\bar{\sigma}_t^2} \Lambda_{\min} \left(\text{Var} \left[\begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix} \right] + \begin{pmatrix} 0 & 0 \\ 0 & (\lambda_t^m)^2 \end{pmatrix} \right) \\ &\geq \frac{1}{\bar{\sigma}_t^2} \Lambda_{\min} \left(\begin{pmatrix} 1 \\ K_t^m(\omega) \end{pmatrix} \text{Var}[\tilde{X}_t^{m,\omega}] \begin{pmatrix} 1 \\ K_t^m(\omega) \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & (\lambda_t^m)^2 \end{pmatrix} \right) \\ &\geq \frac{1}{\bar{\sigma}_t^2} \left(\text{Var}[\tilde{X}_t^{m,\omega}] \wedge (\lambda_t^m)^2 \right) \Lambda_{\min} \left(\begin{pmatrix} 1 \\ K_t^m(\omega) \end{pmatrix} \begin{pmatrix} 1 \\ K_t^m(\omega) \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right) \geq c_0 \left(\text{Var}[\tilde{X}_t^{m,\omega}] \wedge (\lambda_t^m)^2 \right), \end{aligned} \quad (4.6)$$

as $|K_t^m(\omega)| \leq L$ for \mathbb{P} -a.s. Since $\inf_{t \in [0, T]} \bar{\sigma}_t > 0$, K^m is uniformly bounded and ξ^{π_m} and W^m are independent, by (4.4),

$$\begin{aligned} \inf_{m \in \mathbb{N}, t \in [T/2, T]} \text{Var}[\tilde{X}_t^{m,\omega}] &\geq \inf_{m \in \mathbb{N}, t \in [T/2, T]} \mathbb{E} \left[\left(\Phi_t^m(\omega) \int_0^t (\Phi_s^m(\omega))^{-1} \left(\lambda_s^m \xi_s^{\pi_m}(\cdot) \right) ds + \bar{\sigma}_s dW_s^m(\cdot) \right)^2 \right] \\ &\geq \inf_{m \in \mathbb{N}, t \in [T/2, T]} \left((\Phi_t^m(\omega))^2 \int_0^t ((\Phi_s^m(\omega))^{-1} \bar{\sigma}_s)^2 ds \right) > 0. \end{aligned} \quad (4.7)$$

Consequently, by using Fubini's theorem, the concavity of $\mathbb{S}^2 \ni S \mapsto \Lambda_{\min}(S) \in \mathbb{R}$ and Jensen's inequality,

$$\begin{aligned} \Lambda_{\min} &\left(\mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} \left(\begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix} \begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & (\lambda_t^m)^2 \end{pmatrix} \right) dt \right] \right) \\ &\geq \Lambda_{\min} \left(\mathbb{E} \left[\int_{T/2}^T \frac{1}{\bar{\sigma}_t^2} \left(\begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix} \begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & (\lambda_t^m)^2 \end{pmatrix} \right) dt \right] \right) \\ &\geq \int_{T/2}^T \Lambda_{\min} \left(\mathbb{E} \left[\frac{1}{\bar{\sigma}_t^2} \left(\begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix} \begin{pmatrix} \tilde{X}_t^{m,\omega} \\ K_t^m(\omega) \tilde{X}_t^{m,\omega} + k_t^m(\omega) \end{pmatrix}^\top + \begin{pmatrix} 0 & 0 \\ 0 & (\lambda_t^m)^2 \end{pmatrix} \right) \right] \right) dt \\ &\geq \tilde{c}_0 \min_{t \in [0, T]} \left(1 \wedge (\lambda_t^m)^2 \right), \end{aligned} \quad (4.8)$$

for some $\tilde{c}_0 > 0$, where the final inequality follows from (4.6) and (4.7). Hence setting $|\pi^m| \|\lambda^m\|_\infty^2 \leq \frac{\tilde{c}_0}{2C} \min_{t \in [0, T]} ((\lambda_t^m)^2 \wedge 1)$ with C given in (4.5) and using (4.2) (4.3), (4.5) and (4.8) prove Item (2). \square

Based on Propositions 4.1 and 4.2, the following lemma estimates the accuracy of $(\hat{\theta}_m)_{m \in \mathbb{N}}$ in terms of $(V_m^{-1})_{m \in \mathbb{N}}$, which extends [26, Proposition 4.6] to randomised policies.

Lemma 4.3. *Suppose (H.1) and Setting 4.1 hold. Then there exists a constant $C \geq 0$, depending only on the coefficients in (H.1) and θ_0, V_0, L in Setting 4.1, such that for all $m \in \mathbb{N}$,*

$$\mathbb{P}\left(|\hat{\theta}_m - \theta^*|^2 \leq C(\Lambda_{\min}(V_m^{-1}))^{-1}(1 + \ln m)\right) \geq 1 - 1/m. \quad (4.9)$$

Assume further that (H.2) holds. Then for all $m \in \mathbb{N}$,

$$\mathbb{E}[|\theta_m - \theta^*|^2] \leq C \left(\mathbb{E}[(\Lambda_{\min}(V_m^{-1}))^{-1}](1 + \ln m) + \frac{1}{m} + \mathbb{P}(\hat{\theta}_m \notin \Theta) \right). \quad (4.10)$$

Proof. Throughout this proof, let $C > 0$ be a generic constant which is independent of $\delta \in (0, 1)$ and $m \in \mathbb{N}$ and may take different values at each occurrence.

By the same arguments for Lemma 4.5 and Proposition 4.6 in [26] and the fact that $\inf_{t \in [0, T]} \bar{\sigma}_t > 0$, it follows that for all $\delta \in (0, 1)$ and $m \in \mathbb{N}$,

$$\mathbb{P}\left(\Lambda_{\min}(V_m^{-1})|\hat{\theta}_m - \theta^*|^2 \leq C(1 + \ln(\det V_m^{-1}) + \ln(\frac{1}{\delta}))\right) \geq 1 - \delta. \quad (4.11)$$

By Proposition 4.1 and Proposition 4.2 Item (1), it follows that for all $\delta \in (0, 1)$ and $m \in \mathbb{N}$, with probability at least $1 - \delta$,

$$\begin{aligned} \ln(\det V_m^{-1}) &\leq C \ln |V_m^{-1}| \leq C \ln \left(\left| V_0^{-1} + \sum_{n=1}^m \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} Z_t^n (Z_t^n)^\top dt \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right| + C \max \left(\ln \left(\frac{2}{\delta} \right), \sqrt{m \ln \left(\frac{2}{\delta} \right)} \right) \right) \\ &\leq C \ln \left(Cm + C \max \left(\ln \left(\frac{2}{\delta} \right), \sqrt{m \ln \left(\frac{2}{\delta} \right)} \right) \right) \leq C \left(1 + \ln m + \ln \left(\frac{1}{\delta} \right) \right). \end{aligned} \quad (4.12)$$

Substituting $\delta = 1/(2m)$ in (4.11) and (4.12) implies that

$$\mathbb{P}\left(|\hat{\theta}_m - \theta^*|^2 \leq C(\Lambda_{\min}(V_m^{-1}))^{-1}(1 + \ln m)\right) \geq 1 - 1/m.$$

Let A be the event such that the estimate $|\hat{\theta}_m - \theta^*|^2 \leq C(\Lambda_{\min}(V_m^{-1}))^{-1}(1 + \ln m)$ holds. Under (H.2), θ_m and θ^* are uniformly bounded (cf. (2.23)). Hence,

$$\begin{aligned} \mathbb{E}[|\theta_m - \theta^*|^2] &\leq \mathbb{E}[|\theta_m - \theta^*|^2 \mathbf{1}_{A \cap \{\hat{\theta}_m \in \Theta\}}] + C(\mathbb{P}(A^c) + \mathbb{P}(\hat{\theta}_m \notin \Theta)) \\ &\leq C \left(\mathbb{E}[(\Lambda_{\min}(V_m^{-1}))^{-1}](1 + \ln m) + \frac{1}{m} + \mathbb{P}(\hat{\theta}_m \notin \Theta) \right) \end{aligned}$$

where the last inequality uses the fact that $\theta_m = \hat{\theta}_m$ on the event $\{\hat{\theta}_m \in \Theta\}$. \square

The remaining part of this section is devoted to analyse the expected costs of $(\varphi_m)_{m \in \mathbb{N}}$. The following lemma proves stability of (2.5) and regularity of the cost function J^{θ^*} in (2.1). The proof adapts the arguments of Lemma 3.1 and Proposition 3.7 in [1] to the present setting, and is given in Appendix B.

Lemma 4.4. *Suppose (H.1) holds.*

- (1) *The map $\mathbb{R}^{1 \times 2} \ni \theta \mapsto P^\theta, \eta^\theta \in C^1([0, T]; \mathbb{R})$ is continuously differentiable, where $P^\theta \in C([0, T]; \mathbb{R}_{\geq 0})$ and $\eta^\theta \in C([0, T]; \mathbb{R})$ are the solution to (2.5).*
- (2) *For each $L \geq 0$, there exists a constant $C \geq 0$, depending only on L and the coefficients in (H.1), such that for all $K, k \in C([0, T]; \mathbb{R})$ with $\|K\|_\infty, \|k\|_\infty \in [0, L]$,*

$$J^{\theta^*}(\phi) - J^{\theta^*}(\phi^{\theta^*}) \leq C(\|K - K^{\theta^*}\|_\infty^2 + \|k - k^{\theta^*}\|_\infty^2),$$

where $\phi : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ satisfies $\phi(t, x) = K_t x + k_t$ for all $(t, x) \in [0, T] \times \mathbb{R}$, and ϕ^{θ^*} is the optimal policy of (2.1).

Based on Lemma 4.4, the next proposition quantifies the performance gaps between piecewise constant randomised policies $(\varphi_m)_{m \in \mathbb{N}}$ and the optimal policy ϕ^{θ^*} .

Proposition 4.5. *Suppose (H.1) and Setting 4.1 hold. Then there exists a constant $C \geq 0$, depending only on the coefficients in (H.1) and L in Setting 4.1, such that for all $m \in \mathbb{N}$,*

$$\mathbb{E}[J^{\theta^*}(\varphi^m) - J^{\theta^*}(\phi^{\theta^*})] \leq C \left((1 + |\pi_m|) \|\lambda^m\|_\infty^2 + \mathbb{E}[\|K^m - K^{\theta^*}\|_\infty^2] + \mathbb{E}[\|k^m - k^{\theta^*}\|_\infty^2] \right).$$

Proof. We first prove that there exists a constant $C \geq 0$ such that for all $K, k \in L^\infty([0, T]; \mathbb{R})$ with $\|K\|_\infty, \|k\|_\infty \in [0, L]$ and $\lambda \in L^\infty([0, T]; \mathbb{R}_{\geq 0})$, it holds with $\nu(t, x) := \mathcal{N}(K_t x + k_t, \lambda_t^2)$ for all $(t, x) \in [0, T] \times \mathbb{R}$,

$$|\tilde{J}_0^{\theta^*}(\nu) - J^{\theta^*}(\phi^{\theta^*})| \leq C \left(\|\lambda\|_\infty^2 + \|K - K^{\theta^*}\|_\infty^2 + \|k - k^{\theta^*}\|_\infty^2 \right), \quad (4.13)$$

with $\tilde{J}_0^{\theta^*}(\nu)$ defined as in (2.9). Indeed, let $\phi(t, x) := K_t x + k_t$ for all $(t, x) \in [0, T] \times \mathbb{R}$. Then

$$\tilde{J}_0^{\theta^*}(\nu) - J^{\theta^*}(\phi^{\theta^*}) = \left(\tilde{J}_0^{\theta^*}(\nu) - J^{\theta^*}(\phi) \right) + \left(J^{\theta^*}(\phi) - J^{\theta^*}(\phi^{\theta^*}) \right). \quad (4.14)$$

For the first term of (4.14), observe that $X^{\theta^*, \phi}$ and $X^{\theta^*, \nu}$ satisfy the same dynamics:

$$dX_t = (A^* X_t + B^*(K_t X_t + k_t)) dt + \bar{\sigma}_t dW_t, \quad t \in [0, T]; \quad X_0 = x_0,$$

and hence $X^{\theta^*, \phi} = X^{\theta^*, \nu}$ due to the uniqueness of solutions. Thus, by (2.9),

$$\tilde{J}_0^{\theta^*}(\nu) = \mathbb{E} \left[\int_0^T \left(f(t, X_t^{\theta^*, \phi}, K_t X_t^{\theta^*, \phi} + k_t) + R_t \lambda_t^2 \right) dt + g(X_T^{\theta^*, \phi}) \right] = J^{\theta^*}(\phi) + \int_0^T R_t \lambda_t^2 dt, \quad (4.15)$$

which along with $R \in L^\infty([0, T]; \mathbb{R})$ shows $|\tilde{J}_0^{\theta^*}(\nu) - J^{\theta^*}(\phi)| \leq C \|\lambda\|_\infty^2$. By Lemma 4.4, the second term of (4.14) can be estimated by $J^{\theta^*}(\phi) - J^{\theta^*}(\phi^{\theta^*}) \leq C(\|K - K^{\theta^*}\|_\infty^2 + \|k - k^{\theta^*}\|_\infty^2)$, which subsequently leads to (4.13).

Now for each $m \in \mathbb{N}$, consider

$$\tilde{J}_0^{\theta^*}(\nu^m) = \mathbb{E} \left[\int_0^T \left(\int_{\mathbb{R}} f(t, X_t^{\nu^m}, a) \nu^m(t, X_t^{\nu^m}; da) \right) dt + g(X_T^{\nu^m}) \Big| \mathcal{F}_{m-1}^{\text{ob}} \right],$$

where ν^m is defined in Setting 4.1, and X^{ν^m} satisfies for all $t \in [0, T]$,

$$dX_t^{\nu^m} = \int_{\mathbb{R}} (A^* X_t^{\nu^m} + B^* a) \nu^m(t, X_t^{\nu^m}; da) dt + \bar{\sigma}_t dW_t^m, \quad X_0^{\nu^m} = x_0.$$

The assumptions in Setting 4.1 ensures that $(K^m, k^m)_{m \in \mathbb{N} \cup \{0\}}$ are uniformly bounded. Then by (4.13), there exists $C \geq 0$ such that for all $m \in \mathbb{N}$,

$$|\tilde{J}_0^{\theta^*}(\nu^m) - J^{\theta^*}(\phi^{\theta^*})| \leq C \left(\|\lambda^m\|_\infty^2 + \|K^m - K^{\theta^*}\|_\infty^2 + \|k^m - k^{\theta^*}\|_\infty^2 \right), \quad d\mathbb{P}\text{-a.s.} \quad (4.16)$$

Moreover, Theorem 3.3 and the uniform boundedness of K^m, k^m, λ^m imply that

$$\mathbb{E} \left[J^{\theta^*}(\varphi^m) - \tilde{J}_0^{\theta^*}(\nu^m) \Big| \mathcal{F}_{m-1}^{\text{ob}} \right] \leq C |\pi_m| \|\lambda^m\|_\infty^2.$$

which along with the tower property of conditional expectations and (4.16) leads to the desired estimate. \square

4.2 Regret analysis of Algorithm 1

This section establishes the regret of Algorithm 1 (i.e., Theorem 2.2). For notational simplicity, we denote by $C > 0$ a generic constant which is independent of the episode number m and may take different values at each occurrence. The following lemma proves that the policies generated by Algorithm 1 satisfies Setting 4.1, and hence the results in Section 4.1 apply to these policies. The proof is given in Appendix B.

Lemma 4.6. *Suppose (H.1) and (H.2) hold and $\sup_{m \in \mathbb{N}} \varrho_m < \infty$. Let $k^1, K^1 \in C([0, T]; \mathbb{R})$, $\lambda^1 \in C([0, T]; \mathbb{R}_{\geq 0})$ and $\nu^1 \sim \mathcal{G}(k^1, K^1, \lambda^1)$. Then the policies $(\varphi^m)_{m \in \mathbb{N}}$ from Algorithm 1 satisfy Setting 4.1.*

We first establish the convergence rate of $(\theta_m)_{m \in \mathbb{N}}$ for suitable choices of $(\varrho_m)_{m \in \mathbb{N}}$ and $(\pi_m)_{m \in \mathbb{N}}$.

Proposition 4.7. *Suppose (H.1) and (H.2) hold. Let $\theta_0 \in \mathbb{R}^{1 \times 2}$, $V_0 \in \mathbb{S}_+^2$, $\varrho_0 > 0$ and $\nu^1 \sim \mathcal{G}(k^{\theta_0}, K^{\theta_0}, \sqrt{\frac{\varrho_0}{2R}})$. Then there exist constants $c_0, C > 0$ such that if one sets*

$$\varrho_m = \varrho_0 m^{-1/2} \ln(m+1) \quad \text{and} \quad |\pi_m| \varrho_{m-1} \leq c_0 (\varrho_{m-1} \wedge 1), \quad \forall m \in \mathbb{N},$$

then $\mathbb{E}[|\theta_m - \theta^*|^2] \leq C m^{-1/2}$ for all $m \in \mathbb{N}$.

Proof. We first establish a lower and upper bound of $(V_m^{-1})_{m \in \mathbb{N}}$ in terms of m . Note that by (2.11) and the choice of ν^1 , Algorithm 1 defines $K^m = K^{\theta_{m-1}}$, $k^m = k^{\theta_{m-1}}$ and $(\lambda^m)^2 = \frac{\varrho_{m-1}}{2R}$ for all $m \in \mathbb{N}$. As $0 < \inf_{t \in [0, T]} R_t \leq \sup_{t \in [0, T]} R_t < \infty$, by choosing a sufficiently small $c_0 > 0$ and setting $|\pi_m| \varrho_{m-1} \leq c_0 (\varrho_{m-1} \wedge 1)$ for all $m \in \mathbb{N}$, Proposition 4.2 Item (2) implies that for all $m \in \mathbb{N}$,

$$\sum_{n=1}^m \Lambda_{\min} \left(\mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right) \geq r \sum_{n=1}^m (\varrho_{n-1} \wedge 1), \quad (4.17)$$

with some constant $r > 0$ independent of m . By the definition of $(\varrho_m)_{m \in \mathbb{N}}$, there exists $\bar{N}_0 \in \mathbb{N}$ such that $\varrho_m \leq 1$ for all $m \geq \bar{N}_0$. In particular, for $m \geq 4\bar{N}_0$, it follows from (4.17) that

$$\begin{aligned} \sum_{n=1}^m \Lambda_{\min} \left(\mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right) &\geq r \sum_{n=\bar{N}_0}^m \varrho_0 (n-1)^{-1/2} \ln n \geq r \int_{\bar{N}_0}^m (x-1)^{-1/2} \ln x dx \\ &\geq r \int_{m/4}^m (x-1)^{-1/2} \ln x dx \geq r \ln \left(\frac{m}{4} \right) \int_{m/4}^m (x-1)^{-1/2} dx \geq r m^{1/2} \ln(m+1). \end{aligned} \quad (4.18)$$

By applying Proposition 4.1 with $\delta = 1/m$, for each $m \in \mathbb{N}$, with probability at least $1 - 1/m$,

$$\left| V_m^{-1} - V_0^{-1} - \sum_{n=1}^m \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} Z_t^n (Z_t^n)^\top dt \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right| \leq C \sqrt{m \ln(m+1)}. \quad (4.19)$$

Let A_m be the event that (4.19) holds. Without loss of generality, assume that \bar{N}_0 is sufficiently large such that $C \ln^{-1/2}(m+1) \leq r/2$ for all $m \geq 4\bar{N}_0$, with the constant C in (4.19). Then, for any $m \geq 4\bar{N}_0$, by (4.18) and (4.19), under the event A_m ,

$$\begin{aligned} &\Lambda_{\min}(V_m^{-1}) \\ &\geq \sum_{n=1}^m \Lambda_{\min} \left(V_0^{-1} + \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right) - \left| V_m^{-1} - V_0^{-1} - \sum_{n=1}^m \mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_t^2} Z_t^n (Z_t^n)^\top dt \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right| \\ &\geq r m^{\frac{1}{2}} \ln(m+1) - (C \ln^{-\frac{1}{2}}(m+1)) m^{\frac{1}{2}} \ln(m+1) \geq \frac{1}{2} r m^{\frac{1}{2}} \ln(m+1), \end{aligned}$$

which along with $(\Lambda_{\min}(V_m^{-1}))^{-1} \leq (\Lambda_{\min}(V_0^{-1}))^{-1}$ and $\mathbb{P}(A_m^c) \leq 1/m$ shows that for all $m \geq 4\bar{N}_0$,

$$\begin{aligned} \mathbb{E} \left[(\Lambda_{\min}(V_m^{-1}))^{-1} \right] &= \mathbb{E} \left[(\Lambda_{\min}(V_m^{-1}))^{-1} \mathbf{1}_{A_m} \right] + \mathbb{E} \left[(\Lambda_{\min}(V_m^{-1}))^{-1} \mathbf{1}_{A_m^c} \right] \\ &\leq C m^{-1/2} (\ln(m+1))^{-1} + \frac{1}{m} (\Lambda_{\min}(V_0^{-1}))^{-1} \leq C m^{-1/2} (\ln(m+1))^{-1}. \end{aligned} \quad (4.20)$$

Now consider the event such that the estimate (4.9) in Lemma 4.3 and (4.19) hold simultaneously. Then for all $m \geq 4\bar{N}_0$, with probability at least $1 - 2/m$,

$$|\hat{\theta}_m - \theta^*|^2 \leq C m^{-1/2} (\ln(m+1))^{-1} (1 + \ln m). \quad (4.21)$$

As $\theta^* \in \text{int}(\Theta)$ by (H.2), one can choose a sufficiently large $\bar{N}_0 \in \mathbb{N}$ such that for all $m \geq 4\bar{N}_0$, $C m^{-1/2} (\ln(m+1))^{-1} (1 + \ln m) \leq \inf_{\theta \notin \Theta} |\theta - \theta^*|^2$, with the constant C in (4.21). Hence, $\hat{\theta}_m \in \Theta$ on

the event that (4.21) holds, i.e. $\mathbb{P}(\hat{\theta}_m \notin \Theta) \leq 2/m$ for all $m \geq 4\bar{N}_0$. Combining this with the estimate (4.10) in Lemma 4.3 and (4.20), for all $m \geq 4\bar{N}_0$,

$$\mathbb{E}[|\theta_m - \theta^*|^2] \leq C \left(Cm^{-1/2}(\ln(m+1))^{-1}(1 + \ln m) + \frac{1}{m} + \frac{2}{m} \right) \leq Cm^{-1/2}.$$

By the uniform boundedness of $(\theta_m)_{m \in \mathbb{N}} \subset \Theta$, there exists a sufficiently large C such that the estimate holds for all $m \in \mathbb{N}$. \square

Proof of Theorem 2.2. By Lemma 4.4 Item (1), $\mathbb{R}^{1 \times 2} \ni \theta \mapsto (K^\theta, k^\theta) \in C([0, T]; \mathbb{R})^2$ is continuously differentiable and hence locally Lipschitz continuous. Then the fact that $(\theta_m)_{m \in \mathbb{N}} \subset \Theta$ and the boundedness of Θ (see (H.2)) imply that for all $m \in \mathbb{N}$ and \mathbb{P} -a.s.,

$$\|K^m - K^{\theta^*}\|_\infty^2 + \|k^m - k^{\theta^*}\|_\infty^2 = \|K^{\theta_{m-1}} - K^{\theta^*}\|_\infty^2 + \|k^{\theta_{m-1}} - k^{\theta^*}\|_\infty^2 \leq C|\theta_{m-1} - \theta^*|^2. \quad (4.22)$$

Hence, by Proposition 4.5 and the fact that $\|\lambda^m\|_\infty^2 \leq C\varrho_{m-1}$ for all $m \in \mathbb{N}$,

$$\mathbb{E}[|J^{\theta^*}(\varphi^m) - J^{\theta^*}(\phi^{\theta^*})|] \leq C \left((1 + |\pi_m|)\varrho_{m-1} + \mathbb{E}[|\theta_{m-1} - \theta^*|^2] \right), \quad \forall m \in \mathbb{N}.$$

Combining this with Proposition 4.7 shows that there exists $c_0 > 0$ such that if one sets

$$\varrho_m = \varrho_0 m^{-1/2} \ln(m+1), \quad |\pi_m| \varrho_{m-1} \leq c_0(\varrho_{m-1} \wedge 1), \quad \forall m \in \mathbb{N}, \quad (4.23)$$

then for all $N \in \mathbb{N}$,

$$\begin{aligned} \mathbb{E}[\text{Reg}(N)] &= \sum_{m=1}^N \mathbb{E}[|J^{\theta^*}(\varphi^m) - J^{\theta^*}(\phi^{\theta^*})|] \leq C \sum_{m=1}^N \left(|\pi_m| \varrho_{m-1} + \varrho_{m-1} + \mathbb{E}[|\theta_{m-1} - \theta^*|^2] \right) \\ &\leq C \sum_{m=1}^N \left(|\pi_m| \varrho_{m-1} + \varrho_{m-1} + m^{-1/2} \right). \end{aligned}$$

In particular, by choosing $|\pi_m| \leq \frac{c_0}{\max(\sup_{m \in \mathbb{N}} \varrho_{m-1}, 1)}$, the condition (4.23) holds and

$$\mathbb{E}[\text{Reg}(N)] \leq C \sum_{m=1}^N (\varrho_{m-1} + m^{-1/2}) \leq CN^{1/2} \ln(N+1). \quad \square$$

4.3 Regret analysis of Algorithm 2

This section establishes the regret of Algorithm 2 (i.e., Theorem 2.3). As in Section 4.2, $C > 0$ denotes a generic constant which is independent of the episode number m and may take different values at each occurrence. The following lemma proves an analogue of Lemma 4.6 for Algorithm 2. The proof is given in Appendix B.

Lemma 4.8. *Suppose (H.1) and (H.2) hold. Let $k^1, K^1 \in C([0, T]; \mathbb{R})$, $\lambda^1 \in C([0, T]; \mathbb{R}_{>0})$ and $\nu^1 \sim \mathcal{G}(k^1, K^1, \lambda^1)$. Then the policies $(\varphi^m)_{m \in \mathbb{N}}$ from Algorithm 2 satisfy Setting 4.1.*

The following lemma will be used to quantify the dependence of $\|K^m - K^{\theta^*}\|_\infty$ and $\|k^m - k^{\theta^*}\|_\infty$ on $(\theta_n - \theta^*)_{n=0}^m$.

Lemma 4.9. *Suppose (H.1) holds and for each $m \in \mathbb{N}$, let h^m be defined by (2.14). Then $h_t^m = 1 - \frac{(\lambda_t^{m+1})^2}{(\lambda_t^m)^2}$ for all $m \in \mathbb{N}$ and $t \in [0, T]$. Assume further that $(\varrho_m)_{m \in \mathbb{N}}$ is increasing, i.e., $\varrho_m \leq \varrho_{m+1}$ for all $m \in \mathbb{N}$. Then $h_t^m \leq \frac{1}{m}$ for all $m \in \mathbb{N}$ and $t \in [0, T]$.*

Proof. By (2.14), for all $m \in \mathbb{N}$,

$$1 - h_t^m = \frac{\varrho_m (\lambda_t^m)^{-2}}{2R_t + \varrho_m (\lambda_t^m)^{-2}} = \frac{\varrho_m (\lambda_t^m)^{-2}}{\varrho_m (\lambda_t^{m+1})^{-2}} = \frac{(\lambda_t^{m+1})^2}{(\lambda_t^m)^2}.$$

Assume further that $(\varrho_m)_{m \in \mathbb{N}}$ is increasing, by (2.14), for all $m \in \mathbb{N}$,

$$\frac{\varrho_m}{2R_t}(\lambda_t^m)^{-2} \geq \frac{\varrho_{m-1}}{2R_t}(\lambda_t^m)^{-2} = 1 + \frac{\varrho_{m-1}}{2R_t}(\lambda_t^{m-1})^{-2} \geq \dots \geq m-1 + \left(\frac{\varrho_1}{2R_t}\right)(\lambda_t^1)^{-2} \geq m-1,$$

which along with $h_t^m = \left(1 + \frac{\varrho_m}{2R_t}(\lambda_t^m)^{-2}\right)^{-1}$ leads to the desired estimate. \square

Based on Lemma 4.9, we quantify the accuracy of K^m and k^m in terms of $(\theta_n)_{n=0}^m$.

Proposition 4.10. *Suppose (H.1) and (H.2) hold. Let $\nu^1 \sim \mathcal{G}(k^1, K^1, \lambda^1)$ for some $k^1, K^1 \in C([0, T]; \mathbb{R})$ and $\lambda^1 \in C([0, T]; \mathbb{R}_{>0})$. Then there exists a constant $C \geq 0$ such that for all $\theta_0 \in \mathbb{R}^{1 \times 2}$, $V_0 \in \mathbb{S}_+^2$, $m \in \mathbb{N}$ and increasing sequences $(\varrho_m)_{m \in \mathbb{N}} \subset (0, \infty)$,*

$$\begin{aligned} & \|K^{m+1} - K^{\theta^*}\|_\infty^2 + \|k^{m+1} - k^{\theta^*}\|_\infty^2 \\ & \leq C \sum_{n=1}^m \frac{1}{n} \left\| \frac{\lambda^{m+1}}{\lambda^{n+1}} \right\|_\infty^2 |\theta_n - \theta^*|^2 + \left\| \frac{\lambda^{m+1}}{\lambda^2} \right\|_\infty^2 \left(\|K^1 - K^{\theta^*}\|_\infty^2 + \|k^1 - k^{\theta^*}\|_\infty^2 \right), \end{aligned}$$

where for each $m \in \mathbb{N}$, K^{m+1} , k^{m+1} and λ^m are defined by (2.14).

Proof. For all $m \in \mathbb{N}$ and $t \in [0, T]$, by (2.14), $h_t^m \in [0, 1]$ and

$$|K_t^{m+1} - K_t^{\theta^*}| \leq h_t^m |K_t^{\theta_m} - K_t^{\theta^*}| + (1 - h_t^m) |K_t^m - K_t^{\theta^*}|.$$

Then by using the convexity of $\mathbb{R} \ni x \mapsto x^2 \in \mathbb{R}$,

$$|K_t^{m+1} - K_t^{\theta^*}|^2 \leq h_t^m |K_t^{\theta_m} - K_t^{\theta^*}|^2 + (1 - h_t^m) |K_t^m - K_t^{\theta^*}|^2.$$

Similar to (4.22), one has $\|K^{\theta_m} - K^{\theta^*}\|_\infty^2 \leq C|\theta_m - \theta^*|^2$, which subsequently leads to

$$|K_t^{m+1} - K_t^{\theta^*}|^2 \leq Ch_t^m |\theta_m - \theta^*|^2 + (1 - h_t^m) |K_t^m - K_t^{\theta^*}|^2.$$

Applying the same argument to $|k_t^{m+1} - k_t^{\theta^*}|$ leads to the estimate:

$$|K_t^{m+1} - K_t^{\theta^*}|^2 + |k_t^{m+1} - k_t^{\theta^*}|^2 \leq Ch_t^m |\theta_m - \theta^*|^2 + (1 - h_t^m) (|K_t^m - K_t^{\theta^*}|^2 + |k_t^m - k_t^{\theta^*}|^2). \quad (4.24)$$

For each $m \in \mathbb{N}$ and $t \in [0, T]$, let $\kappa_{m,t} := |K_t^m - K_t^{\theta^*}|^2 + |k_t^m - k_t^{\theta^*}|^2$, $\Delta_m := |\theta_m - \theta^*|^2$ and $\eta_{m,t} := \prod_{k=1}^m (1 - h_t^k) = \frac{(\lambda_t^{m+1})^2}{(\lambda_t^1)^2}$ by Lemma 4.9. Then for all $m \in \mathbb{N}$ and $t \in [0, T]$, (4.24) and the fact that $h_t^{m+1} \leq \frac{1}{m}$ imply that

$$\begin{aligned} \eta_{m,t}^{-1} \kappa_{m+1,t} & \leq C \frac{\eta_{m,t}^{-1}}{m} \Delta_m + \eta_{m-1,t}^{-1} \kappa_{m-1,t} \leq C \frac{\eta_{m,t}^{-1}}{m} \Delta_m + C \frac{\eta_{m-1,t}^{-1}}{m-1} \Delta_{m-1} + \eta_{m-2,t}^{-1} \kappa_{m-2,t} \\ & \leq C \sum_{n=1}^m \frac{\eta_{n,t}^{-1}}{n} \Delta_n + \eta_1^{-1} \kappa_{1,t}. \end{aligned}$$

Multiplying both sides by η_m and using $\frac{\eta_{m,t}}{\eta_{n,t}} = \frac{(\lambda_t^{m+1})^2}{(\lambda_t^{n+1})^2}$ yield

$$\kappa_{m+1,t} \leq C \sum_{n=1}^m \frac{1}{n} \left(\frac{\lambda_t^{m+1}}{\lambda_t^{n+1}} \right)^2 \Delta_n + \left(\frac{\lambda_t^{m+1}}{\lambda_t^2} \right)^2 \kappa_{1,t}.$$

Taking the supremum over t leads to the desired inequality. \square

To simplify the estimate in Proposition 4.10, we quantify the behavior of $(\lambda^m)_{m \in \mathbb{N}}$ in terms of m .

Lemma 4.11. *Suppose (H.1) holds. Let $\varrho_0 > 0$ and $\inf_{t \in [0, T]} \lambda_t^1 > 0$. Then there exist constants $\bar{c} \geq \underline{c} > 0$ such that if one sets $\varrho_m = \varrho_0 m^{1/2} \ln(m+1)$ for all $m \in \mathbb{N}$, then*

$$\underline{c} m^{-1/2} \ln(m+1) \leq (\lambda_t^m)^2 \leq \bar{c} m^{-1/2} \ln(m+1), \quad \forall t \in [0, T], m \in \mathbb{N} \cap [2, \infty).$$

where λ^m , for $m \in \mathbb{N} \cap [2, \infty)$, is defined by (2.14).

Proof. Observe from (H.1) that $\inf_{t \in [0, T]} R_t > 0$. Then by (2.14) and the choice of $(\varrho_m)_{m \in \mathbb{N}}$, for all $m \geq 2$ and $t \in [0, T]$,

$$\begin{aligned} (\lambda_t^m)^{-2} &= 2R_t \varrho_{m-1}^{-1} + (\lambda_t^{m-1})^{-2} = 2R_t \sum_{k=1}^{m-1} \varrho_k^{-1} + (\lambda_t^1)^{-2} \geq C \sum_{k=1}^{m-1} k^{-1/2} (\ln(k+1))^{-1} \\ &\geq C (\ln(m+1))^{-1} \sum_{k=1}^{m-1} k^{-1/2} \geq C (\ln(m+1))^{-1} \int_1^m x^{-1/2} dx \geq Cm^{1/2} (\ln(m+1))^{-1}. \end{aligned}$$

Rearranging the terms leads to the desired upper bound.

On the other hand, observe that there exists $C > 0$ such that for all $x \geq 1$,

$$\frac{d}{dx} x^{1/2} (\ln(x+2))^{-1} = \frac{(x+2) \ln(x+2) - 2x}{2x^{1/2}(x+2) \ln^2(x+2)} \geq Cx^{-1/2} (\ln(x+1))^{-1}.$$

Then by the facts that $\sup_{t \in [0, T]} R_t < \infty$ and $(\lambda_t^1)^{-2} \leq C$, for all $m \geq 2$ and $t \in [0, T]$,

$$\begin{aligned} (\lambda_t^m)^{-2} &= 2R_t \sum_{k=1}^{m-1} \varrho_k^{-1} + (\lambda_t^1)^{-2} \leq C \sum_{k=1}^{m-1} k^{-1/2} (\ln(k+1))^{-1} \leq C \sum_{k=1}^{m-1} \frac{(k+2) \ln(k+2) - 2k}{2k^{1/2}(k+2) \ln^2(k+2)} \\ &\leq C \int_1^m \frac{(x+2) \ln(x+2) - 2x}{2x^{1/2}(x+2) \ln^2(x+2)} dx \leq Cm^{1/2} (\ln(m+2))^{-1} \leq Cm^{1/2} (\ln(m+1))^{-1}. \end{aligned}$$

Rearranging the terms yields to the desired lower bound. \square

Based on Lemma 4.11, the next proposition proves the convergence rate of $(\theta_m)_{m \in \mathbb{N}}$ in terms of m .

Proposition 4.12. *Suppose (H.1) and (H.2) hold. Let $\theta_0 \in \mathbb{R}^{1 \times 2}$, $V_0 \in \mathbb{S}_+^2$ and $\varrho_0 > 0$. Then there exist constants $c_0, C > 0$ such that if one sets $\nu^1 \sim \mathcal{G}(k^{\theta_0}, K^{\theta_0}, \varrho_0)$, $\varrho_m = \varrho_0 m^{1/2} \ln(m+1)$ and $|\pi_m| \leq c_0$ for all $m \in \mathbb{N}$, then $\mathbb{E}[|\theta_m - \theta^*|^2] \leq Cm^{-1/2}$ for all $m \in \mathbb{N}$.*

Proof. By Lemma 4.11, for all $m \geq 2$, $\|\lambda^m\|_\infty^2 \leq (\bar{c}/\underline{c}) \min_{t \in [0, T]} (\lambda_t^m)^2$ and $\sup_{m \in \mathbb{N}} \|\lambda^m\|_\infty^2 < \infty$. Let $c_0 := \frac{1}{\bar{c}} \min(\underline{c}/\bar{c}, 1/(\sup_{m \in \mathbb{N}} \|\lambda^m\|_\infty^2)) > 0$, with C being the constant in Proposition 4.2 Item (2). Hence, if $|\pi_m| \leq c_0$ for all $m \in \mathbb{N}$, then $|\pi^m| \|\lambda^m\|_\infty^2 \leq \frac{1}{\bar{c}} \min_{t \in [0, T]} ((\lambda_t^m)^2 \wedge 1)$ for all $m \in \mathbb{N}$, and

$$\sum_{n=1}^m \Lambda_{\min} \left(\mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right) \geq r \sum_{n=1}^m \min_{t \in [0, T]} ((\lambda_t^n)^2 \wedge 1), \quad (4.25)$$

with some constant $r > 0$ independent of m . Moreover, Lemma 4.11 implies that there exists $\bar{N}_0 \in \mathbb{N}$ such that for all $t \in [0, T]$ and $m \geq \bar{N}_0$, $(\lambda_t^m)^2 \leq 1$ for all $t \in [0, T]$. Hence, using (4.25) and following the same argument as in Proposition 4.7 yield (cf. (4.18)):

$$\sum_{n=1}^m \Lambda_{\min} \left(\mathbb{E} \left[\int_0^T \frac{1}{\bar{\sigma}_s^2} Z_s^n (Z_s^n)^\top ds \middle| \mathcal{F}_{n-1}^{\text{ob}} \right] \right) \geq rm^{1/2} \ln(m+1), \quad \forall m \geq 4\bar{N}_0.$$

Now proceeding along the lines of the proof of Proposition 4.7 leads to the desired estimate. \square

Proof of Theorem 2.3. Note that the choice of $(\varrho_m)_{m \in \mathbb{N}}$ ensures that $(\varrho_m)_{m \in \mathbb{N}}$ is increasing. By Propositions 4.10 and 4.12, for all $m \in \mathbb{N}$,

$$\begin{aligned} &\mathbb{E}[\|K^{m+1} - K^{\theta^*}\|_\infty^2] + \mathbb{E}[\|k^{m+1} - k^{\theta^*}\|_\infty^2] \\ &\leq C \sum_{n=1}^m \frac{1}{n} \left\| \frac{\lambda^{m+1}}{\lambda^{n+1}} \right\|_\infty^2 \mathbb{E}[|\theta_n - \theta^*|^2] + \left\| \frac{\lambda^{m+1}}{\lambda^2} \right\|_\infty^2 \left(\|K^{\theta_0} - K^{\theta^*}\|_\infty^2 + \|k^{\theta_0} - k^{\theta^*}\|_\infty^2 \right) \\ &\leq C \sum_{n=1}^m \frac{1}{n^{3/2}} \left\| \frac{\lambda^{m+1}}{\lambda^{n+1}} \right\|_\infty^2 + C \left\| \frac{\lambda^{m+1}}{\lambda^2} \right\|_\infty^2. \end{aligned}$$

By Lemma 4.11, for all $m \in \mathbb{N}$,

$$\begin{aligned} \mathbb{E}[\|K^{m+1} - K^{\theta^*}\|_\infty^2] + \mathbb{E}[\|k^{m+1} - k^{\theta^*}\|_\infty^2] &\leq Cm^{-\frac{1}{2}} \ln(m+1) \left(\sum_{n=1}^m \frac{1}{n \ln(n+1)} + 1 \right) \\ &\leq Cm^{-\frac{1}{2}} \ln(m+1) (\ln(\ln(m+1)) + 1). \end{aligned}$$

Summing over the index m then yields that for all $N \in \mathbb{N}$,

$$\sum_{m=1}^N \left(\mathbb{E}[\|K^{m+1} - K^{\theta^*}\|_\infty^2] + \mathbb{E}[\|k^{m+1} - k^{\theta^*}\|_\infty^2] \right) \leq CN^{\frac{1}{2}} \ln(N+1) (\ln(\ln(N+1)) + 1). \quad (4.26)$$

Note that the choices of ϱ_m and π_m and Lemma 4.11 imply that $\|\lambda^m\|_\infty^2 + |\pi_m| \|\lambda^m\|_\infty^2 \leq Cm^{-\frac{1}{2}} \ln(m+1)$, for all $m \in \mathbb{N} \cap [2, \infty)$, which along with (4.26) and Proposition 4.5 shows that for all $N \in \mathbb{N}$,

$$\begin{aligned} \mathbb{E}[\text{Reg}(N)] &= \mathbb{E}[J^{\theta^*}(\varphi^1) - J^{\theta^*}(\phi^{\theta^*})] + \sum_{m=2}^N \mathbb{E}[J^{\theta^*}(\varphi^m) - J^{\theta^*}(\phi^{\theta^*})] \\ &\leq C \left(\sum_{m=1}^N m^{-\frac{1}{2}} \ln(m+1) + N^{\frac{1}{2}} \ln(N+1) (\ln(\ln(N+1)) + 1) \right) \\ &\leq CN^{\frac{1}{2}} \ln(N+1) (\ln(\ln(N+1)) + 1). \end{aligned}$$

This proves the desired estimate. \square

Appendix A Proofs of technical results in Section 3

To prove Lemma 3.1, we first express the first and second moments of X^ν and X^φ in terms of the noise process ξ .

Lemma A.1. *Suppose (H.3) holds. For each $\nu \sim \mathcal{G}(k, K, \lambda)$ and $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$, let X^ν and X^φ be defined by (3.3) and (3.5), respectively. Then for all $\nu \sim \mathcal{G}(k, K, \lambda)$, $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$, and $t \in [0, T]$,*

$$\begin{aligned} \mathbb{E}[X_t^\varphi - X_t^\nu | \mathcal{F}_0] &= f_{1,t} \int_0^t (f_{1,s})^{-1} B_s \lambda_s \xi_s \, ds, \\ \mathbb{E}[(X_t^\varphi)^2 - (X_t^\nu)^2 | \mathcal{F}_0] &= f_{2,t} \int_0^t \int_0^t (f_{2,r})^{-1} f_{3,r} f_{1,r} (f_{1,s})^{-1} B_s \lambda_s \xi_s \mathbf{1}_{s \leq r} \, ds dr \\ &\quad + f_{2,t} \int_0^t \int_0^t (f_{2,r})^{-1} f_{4,r} f_{1,r} (f_{1,s})^{-1} B_s \lambda_r \lambda_s \xi_r \xi_s \mathbf{1}_{s \leq r} \, ds dr \\ &\quad + f_{2,t} \int_0^t (f_{2,r})^{-1} f_{5,r} \lambda_r \xi_r \, dr + f_{2,t} \int_0^t (f_{2,r})^{-1} D_r^2 \lambda_r^2 (\xi_r^2 - 1) \, dr, \end{aligned}$$

where $f_1, f_2, f_3, f_4, f_5 : [0, T] \rightarrow \mathbb{R}$ are defined by: for all $t \in [0, T]$,

$$\begin{aligned} f_{1,t} &= \exp\left(\int_0^t (A_s + B_s K_s) \, ds\right), & f_{2,t} &= \exp\left(\int_0^t \left(2(A_s + B_s K_s) + (C_s + D_s K_s)^2\right) \, ds\right), \\ f_{3,t} &= 2(B_t k_t + \bar{b}_t) + 2(C_t + D_t K_t)(\bar{\sigma}_t + D_t k_t), & f_{4,t} &= 2B_t + 2D_t(C_t + D_t K_t), \\ f_{5,t} &= 2D_t(\bar{\sigma}_t + D_t k_t) + 2(B_t + D_t(C_t + D_t K_t)) f_{1,t} \left(x_0 + \int_0^t (f_{1,s})^{-1} (B_s k_s + \bar{b}_s) \, ds\right). \end{aligned}$$

Proof. By (3.3) and (3.5),

$$d(X_t^\varphi - X_t^\nu) = (\eta_t(X_t^\varphi - X_t^\nu) + B_t \lambda_t \xi_t) dt + \mathcal{M}_t^1, \quad t \in [0, T], \quad (\text{A.1})$$

where $\eta_t := A_t + B_t K_t$ for all $t \in [0, T]$, and \mathcal{M}^1 is a local martingale. As X^ν and X^φ are square-integrable, the local martingale \mathcal{M}^1 is in fact a martingale. Let h be such that $h_t = \mathbb{E}[X_t^\varphi - X_t^\nu | \mathcal{F}_0]$ for all $t \in [0, T]$. By Fubini's theorem and (A.1), the process h is of finite variation and satisfies

$$dh_t = (\eta_t h_t + B_t \lambda_t \xi_t) dt, \quad t \in [0, T]; \quad h_0 = 0.$$

By introducing the integration factor $f_{1,t} = \exp\left(\int_0^t \eta_s ds\right)$ for all $t \in [0, T]$, the above equation can be solved analytically as:

$$\mathbb{E}[X_t^\varphi - X_t^\nu | \mathcal{F}_0] = h_t = f_{1,t} \int_0^t (f_{1,s})^{-1} B_s \lambda_s \xi_s ds, \quad t \in [0, T]. \quad (\text{A.2})$$

We then proceed to consider the difference of the second moments. By Itô's formula,

$$\begin{aligned} & d((X_t^\varphi)^2 - (X_t^\nu)^2) \\ &= \left(2X_t^\varphi dX_t^\varphi - 2X_t^\nu dX_t^\nu\right) + \left(d[X^\varphi]_t - d[X^\nu]_t\right) \\ &= \left(2X_t^\varphi (\eta_t X_t^\varphi + B_t k_t + \bar{b}_t + B_t \lambda_t \xi_t) - 2X_t^\nu (\eta_t X_t^\nu + B_t k_t + \bar{b}_t)\right) dt \\ & \quad + \left((\bar{\sigma}_t + \gamma_t X_t^\varphi + D_t k_t + D_t \lambda_t \xi_t)^2 - (\bar{\sigma}_t + \gamma_t X_t^\nu + D_t k_t)^2 - D_t^2 \lambda_t^2\right) dt + \mathcal{M}_t^2, \end{aligned} \quad (\text{A.3})$$

where $[X]$ is the quadratic variation of a process X , $\gamma_t := C_t + D_t K_t$ for all $t \in [0, T]$, and \mathcal{M}^2 is a local martingale. Let H and κ be such that for all $t \in [0, T]$, $H_t := \mathbb{E}[(X_t^\varphi)^2 - (X_t^\nu)^2 | \mathcal{F}_0]$ and

$$\kappa_t := \mathbb{E}[X_t^\nu | \mathcal{F}_0] = f_{1,t} \left(x_0 + \int_0^t (f_{1,s})^{-1} (B_s k_s + \bar{b}_s) ds \right). \quad (\text{A.4})$$

As $\mathbb{E}[\sup_{t \in [0, T]} |X_t^\nu|^4] < \infty$ and $\mathbb{E}[\sup_{t \in [0, T]} |X_t^\varphi|^4] < \infty$, the local martingale \mathcal{M}^2 in (A.3) is again a martingale. Taking conditional expectations on both sides of (A.3) and applying Fubini's theorem show that H satisfies the following SDE in terms of κ_t and h_t :

$$\begin{aligned} dH_t &= (2\eta_t + \gamma_t^2) H_t dt + \left(2(B_t k_t + \bar{b}_t) + 2\gamma_t(\bar{\sigma}_t + D_t k_t)\right) h_t dt \\ & \quad + \left(2B_t \mathbb{E}[X_t^\varphi | \mathcal{F}_0] + 2D_t(\bar{\sigma}_t + \gamma_t \mathbb{E}[X_t^\varphi | \mathcal{F}_0] + D_t k_t)\right) \lambda_t \xi_t dt + D_t^2 \lambda_t^2 (\xi_t^2 - 1) dt \\ &= (2\eta_t + \gamma_t^2) H_t dt + \left(2(B_t k_t + \bar{b}_t) + 2\gamma_t(\bar{\sigma}_t + D_t k_t)\right) h_t dt \\ & \quad + (2B_t + 2D_t \gamma_t) \lambda_t h_t \xi_t dt + \left(2\kappa_t B_t + 2D_t(\bar{\sigma}_t + \gamma_t \kappa_t + D_t k_t)\right) \lambda_t \xi_t dt + D_t^2 \lambda_t^2 (\xi_t^2 - 1) dt, \end{aligned}$$

where the last identity used $\mathbb{E}[X_t^\varphi | \mathcal{F}_0] = h_t + \kappa_t$ for all $t \in [0, T]$. Solving the linear differential equation using the integration factor f_2 and substituting the expressions of h_t and κ_t (i.e. (A.2) and (A.4)) into the solution lead to the desired result. \square

Based on Lemma A.1, we compute the term $J(\varphi) - \tilde{J}(\nu)$ for any $\varphi \sim \mathcal{R}(k, K, \lambda, \xi)$ and $\nu \sim \mathcal{G}(k, K, \lambda)$, and hence prove Lemma 3.1.

Proof of Lemma 3.1. Observe from (3.2) and the fact that X^ν is independent of \mathcal{F}_0 that

$$\tilde{J}(\nu) = \mathbb{E} \left[\int_0^T \left(f(t, X_t^\nu, k_t + K_t X_t^\nu) + R_t \lambda_t^2 \right) dt + g(X_T^\nu) \middle| \mathcal{F}_0 \right],$$

and

$$\begin{aligned} J(\varphi) &= \mathbb{E} \left[\int_0^T \left(f(t, X_t^\varphi, k_t + K_t X_t^\varphi) + R_t \lambda_t^2 \xi_t^2 \right) dt + g(X_T^\varphi) \middle| \mathcal{F}_0 \right] \\ & \quad + \mathbb{E} \left[\int_0^T \left(2S_t X_t^\varphi \lambda_t \xi_t + 2R_t (k_t + K_t X_t^\varphi) \lambda_t \xi_t + q_t \lambda_t \xi_t \right) dt \middle| \mathcal{F}_0 \right]. \end{aligned}$$

Hence, by Fubini's theorem and (3.2),

$$\begin{aligned}
J(\varphi) - \tilde{J}(\nu) &= \int_0^T \left(Q_t + 2S_t K_t + R_t (K_t)^2 \right) \mathbb{E}[(X_t^\varphi)^2 - (X_t^\nu)^2 | \mathcal{F}_0] dt \\
&\quad + \int_0^T \left(2S_t k_t + 2R_t K_t k_t + p_t + q_t K_t \right) \mathbb{E}[X_t^\varphi - X_t^\nu | \mathcal{F}_0] dt \\
&\quad + \int_0^T R_t \lambda_t^2 (\xi_t^2 - 1) dt + M \mathbb{E}[(X_T^\varphi)^2 - (X_T^\nu)^2 | \mathcal{F}_0] + m \mathbb{E}[X_T^\varphi - X_T^\nu | \mathcal{F}_0] \\
&\quad + \int_0^T (2S_t + 2R_t K_t) \mathbb{E}[X_t^\varphi | \mathcal{F}_0] \lambda_t \xi_t dt + \int_0^T (2R_t k_t + q_t) \lambda_t \xi_t dt.
\end{aligned} \tag{A.5}$$

By (A.2) and (A.4),

$$\mathbb{E}[X_t^\varphi | \mathcal{F}_0] = f_{1,t} \int_0^t (f_{1,s})^{-1} B_s \lambda_s \xi_s ds + f_{1,t} \left(x_0 + \int_0^t (f_{1,s})^{-1} (B_s k_s + \bar{b}_s) ds \right),$$

with $f_{1,t}$ defined in Lemma A.1. Substituting this into (A.5) and apply Lemma A.1 lead to the following identity:

$$\begin{aligned}
&J(\varphi) - \tilde{J}(\nu) \\
&= \int_0^T \tilde{h}_{1,t} \lambda_t \xi_t dt + \int_0^T \tilde{h}_{2,t} \lambda_t^2 (\xi_t^2 - 1) dt + \int_0^T \int_0^T \tilde{h}_{3,t,r} \lambda_r^2 (\xi_r^2 - 1) \mathbf{1}_{r \leq t} dr dt \\
&\quad + \int_0^T \int_0^T \tilde{h}_{4,t,r} \lambda_r \xi_r \mathbf{1}_{r \leq t} dr dt + \int_0^T \int_0^T \tilde{h}_{5,t,r} \lambda_r \lambda_t \xi_r \xi_t \mathbf{1}_{r \leq t} dr dt \\
&\quad + \int_0^T \int_0^T \int_0^T \tilde{h}_{6,t,r,s} \lambda_s \xi_s \mathbf{1}_{s \leq r \leq t} ds dr dt + \int_0^T \int_0^T \int_0^T \tilde{h}_{7,t,r,s} \lambda_r \lambda_s \xi_s \xi_r \mathbf{1}_{s \leq r \leq t} ds dr dt,
\end{aligned}$$

where $\tilde{h}_1, \tilde{h}_2 : [0, T] \rightarrow \mathbb{R}$, $\tilde{h}_3, \tilde{h}_4, \tilde{h}_5 : [0, T]^2 \rightarrow \mathbb{R}$, and $\tilde{h}_6, \tilde{h}_7 : [0, T]^3 \rightarrow \mathbb{R}$ satisfy for all $t, r, s \in [0, T]$,

$$\begin{aligned}
\tilde{h}_{1,t} &= m f_{1,T} (f_{1,t})^{-1} B_t + M f_{2,T} (f_{2,t})^{-1} f_{5,t} \\
&\quad + f_{1,t} \left(x_0 + \int_0^t (f_{1,s})^{-1} (B_s k_s + \bar{b}_s) ds \right) (2S_t + 2R_t K_t) + (2R_t k_t + q_t), \\
\tilde{h}_{2,t} &= R_t + M f_{2,T} (f_{2,t})^{-1} D_t^2, \quad \tilde{h}_{3,t,r} = (Q_t + 2S_t K_t + R_t (K_t)^2) f_{2,t} (f_{2,r})^{-1} D_r^2, \\
\tilde{h}_{4,t,r} &= (Q_t + 2S_t K_t + R_t (K_t)^2) f_{2,t} (f_{2,r})^{-1} f_{5,r} + (2S_t k_t + 2R_t K_t k_t + p_t + q_t K_t) f_{1,t} (f_{1,r})^{-1} B_r \\
&\quad + M f_{2,T} (f_{2,t})^{-1} f_{3,t} f_{1,t} (f_{1,r})^{-1} B_r, \\
\tilde{h}_{5,t,r} &= M f_{2,T} (f_{2,t})^{-1} f_{4,t} f_{1,t} (f_{1,r})^{-1} B_r + (2S_t + 2R_t K_t) f_{1,t} (f_{1,r})^{-1} B_r, \\
\tilde{h}_{6,t,r,s} &= (Q_t + 2S_t K_t + R_t (K_t)^2) f_{2,t} (f_{2,r})^{-1} f_{3,r} f_{1,r} (f_{1,s})^{-1} B_s, \\
\tilde{h}_{7,t,r,s} &= (Q_t + 2S_t K_t + R_t (K_t)^2) f_{2,t} (f_{2,r})^{-1} f_{4,r} f_{1,r} (f_{1,s})^{-1} B_s,
\end{aligned}$$

with the functions $f_1, f_2, f_3, f_4, f_5 : [0, T] \rightarrow \mathbb{R}$ given in Lemma A.1. Finally, by defining

$$\begin{aligned}
h_{1,t} &= \tilde{h}_{1,t} + \int_0^T \tilde{h}_{4,s,t} \mathbf{1}_{t \leq s} ds + \int_0^T \int_0^T \tilde{h}_{6,s,r,t} \mathbf{1}_{t \leq r \leq s} ds dr, \\
h_{2,t} &= \tilde{h}_{2,t} + \int_0^T \tilde{h}_{3,s,t} \mathbf{1}_{t \leq s} ds \quad \text{and} \quad h_{3,t,r} = \tilde{h}_{5,t,r} \mathbf{1}_{r \leq t} + \int_0^T \tilde{h}_{7,s,r,t} \mathbf{1}_{t \leq r \leq s} ds,
\end{aligned}$$

and applying Fubini's theorem lead to the required identity (3.7). \square

Proof of Lemma 3.2. Throughout this proof, let C be a generic constant independent of N , $(\rho_i)_{i=1}^N$, $(\beta_{ij})_{i,j=1}^N$ and δ . Item (1) follows directly from the general Hoeffding inequality [27, Theorem 2.6.3]. To prove Item

(2), observe that $(\zeta_i^2 - 1)_{i=1}^N$ are mean-zero sub-exponential random variables. Hence by Bernstein's inequality in [27, Theorem 2.8.2], there exists $C \geq 0$ such that for all $\delta \geq 0$, it holds with probability at least $1 - \delta$ that

$$\left| \sum_{i=1}^N \rho_i (\zeta_i^2 - 1) \right| \leq C \max \left(\|\rho\|_2 \sqrt{\ln \left(\frac{2}{\delta} \right)}, \|\rho\|_\infty \ln \left(\frac{2}{\delta} \right) \right) \leq C \|\rho\|_2 \left(1 + \ln \left(\frac{1}{\delta} \right) \right),$$

where the last inequality used the fact that $\|\rho\|_\infty \leq \|\rho\|_2$ and the Cauchy-Schwarz inequality.

It remains to prove Item (3). The independence of $(\zeta_i)_{i \in \mathbb{N}}$ implies that $\mathbb{E}[\sum_{i,j=1,i \neq j}^N \beta_{ij} \zeta_i \zeta_j] = 0$. Then, by considering $A \in \mathbb{R}^{N \times N}$ with diagonal entries $a_{ii} = 0$ for all $i = 1, \dots, N$, and off diagonal entries $a_{ij} = \rho_{ij}^N$ for all $i \neq j$, and by applying the Hanson–Wright inequality (see [27, Theorem 6.2.1]), there exists a constant $C \geq 0$ such that for all $s \geq 0$,

$$\mathbb{P} \left(\left| \sum_{i,j=1,i \neq j}^N \beta_{ij} \zeta_i \zeta_j \right| \geq s \right) \leq 2 \exp \left(-C \min \left(\frac{s^2}{\|A\|_F^2}, \frac{s}{\|A\|_F} \right) \right),$$

where $\|A\|_F$ is the Frobenius norm of A . Consequently for all $\delta > 0$, it holds with probability at least $1 - \delta$ that

$$\left| \sum_{i,j=1,i \neq j}^N \beta_{ij} \zeta_i \zeta_j \right| \leq C \|A\|_F \max \left(\sqrt{\ln \left(\frac{2}{\delta} \right)}, \ln \left(\frac{2}{\delta} \right) \right) \leq C \|A\|_F \left(1 + \ln \left(\frac{1}{\delta} \right) \right),$$

where the last inequality used the Cauchy-Schwarz inequality. \square

Appendix B Proofs of technical results in Section 4

Proof of Lemma 4.4. Item (1) has been proved in [1, Lemma 3.1] for $S = p = q = M = m = 0$. The argument there is based on the implicit function theorem and extends straightforwardly to the present setting with general convex quadratic cost functions (details omitted).

For Item (2), let C be a generic constant depending only on L and the coefficients in (H.1), and define $\|X\|_{\mathcal{S}^2} = \mathbb{E}[\sup_{t \in [0, T]} |X_t|^2]^{1/2}$ for any process $X : \Omega \times [0, T] \rightarrow \mathbb{R}$. Consider the control processes $U^{\phi^*} = \phi^{\theta^*}(\cdot, X^{\theta^*, \phi^*})$ and $U^\phi = \phi(\cdot, X^{\theta^*, \phi})$ $d\mathbb{P} \otimes dt$ a.e. As ϕ^{θ^*} is an optimal policy, a quadratic expansion of (2.1) at U^{ϕ^*} (see [1, Proposition 3.7]) implies that

$$\begin{aligned} J^{\theta^*}(\phi) - J^{\theta^*}(\phi^{\theta^*}) &= 2\mathbb{E} \left[\int_0^T \begin{pmatrix} X_t^{\theta^*, \phi} - X_t^{\theta^*, \phi^{\theta^*}} \\ U_t^\phi - U_t^{\phi^*} \end{pmatrix}^\top \begin{pmatrix} Q_t & S_t \\ S_t & R_t \end{pmatrix} \begin{pmatrix} X_t^{\theta^*, \phi} - X_t^{\theta^*, \phi^{\theta^*}} \\ U_t^\phi - U_t^{\phi^*} \end{pmatrix} dt \right] \\ &\quad + 2\mathbb{E} \left[M(X_T^{\theta^*, \phi} - X_T^{\theta^*, \phi^{\theta^*}})^2 \right] \\ &\leq C \left(\|X^{\theta^*, \phi} - X^{\theta^*, \phi^{\theta^*}}\|_{\mathcal{S}^2}^2 + \|\phi^\theta(\cdot, X^{\theta^*, \phi}) - \phi^{\theta^*}(\cdot, X^{\theta^*, \phi^{\theta^*}})\|_{\mathcal{S}^2}^2 \right). \end{aligned} \tag{B.1}$$

Recall that $X^{\theta^*, \phi}$ satisfies

$$dX_t^{\theta^*, \phi} = (A^* X_t^{\theta^*, \phi} + B^*(K_t X_t^{\theta^*, \phi} + k_t)) dt + \bar{\sigma}_t dW_t, \quad t \in [0, T]; \quad X_0^{\theta^*, \phi} = x_0. \tag{B.2}$$

The analytical solution to (B.2) and the assumption that $\|K\|_\infty, \|k\|_\infty \leq L$ yield $\|X^{\theta^*, \phi}\|_{\mathcal{S}^2} \leq C$. Observe that $X_0^{\theta^*, \phi} = X_0^{\theta^*, \phi^{\theta^*}}$ and for all $t \in [0, T]$,

$$\begin{aligned} d(X_t^{\theta^*, \phi} - X_t^{\theta^*, \phi^{\theta^*}}) &= \left(A^*(X_t^{\theta^*, \phi} - X_t^{\theta^*, \phi^{\theta^*}}) + B^*(K_t X_t^{\theta^*, \phi} + k_t - K_t^{\theta^*} X_t^{\theta^*, \phi^{\theta^*}} - k_t^{\theta^*}) \right) dt \\ &= \left((A^* + B^* K_t^{\theta^*})(X_t^{\theta^*, \phi} - X_t^{\theta^*, \phi^{\theta^*}}) + B^*(K_t - K_t^{\theta^*}) X_t^{\theta^*, \phi} + B^*(k_t - k_t^{\theta^*}) \right) dt. \end{aligned}$$

Let $\Phi \in C([0, T]; \mathbb{R})$ be the fundamental solution of $d\Phi_t^\theta = (A^* + B^*K_t^{\theta^*})\Phi_t^\theta dt$. Then

$$X_t^{\theta^*, \phi} - X_t^{\theta^*, \phi^{\theta^*}} = \Phi_t^\theta \int_0^t (\Phi_s^\theta)^{-1} \left(B^*(K_t - K_t^{\theta^*})X_t^{\theta^*, \phi} + B^*(k_t - k_t^{\theta^*}) \right) ds, \quad t \in [0, T],$$

which along with $\|K^{\theta^*}\|_\infty < \infty$ and $\|X^{\theta^*, \phi}\|_{S^2} \leq C$ shows $\|X^{\theta^*, \phi} - X^{\theta^*, \phi^{\theta^*}}\|_{S^2} \leq C(\|K - K^{\theta^*}\|_\infty + \|k - k^{\theta^*}\|_\infty)$. Furthermore, for all $t \in [0, T]$,

$$\begin{aligned} \phi^\theta(t, X_t^{\theta^*, \phi}) - \phi^{\theta^*}(t, X_t^{\theta^*, \phi^{\theta^*}}) &= K_t X_t^{\theta^*, \phi} + k_t - K_t^{\theta^*} X_t^{\theta^*, \phi^{\theta^*}} - k_t^{\theta^*} \\ &= K_t^{\theta^*} (X_t^{\theta^*, \phi} - X_t^{\theta^*, \phi^{\theta^*}}) + (K_t - K_t^{\theta^*}) X_t^{\theta^*, \phi} + k_t - k_t^{\theta^*}. \end{aligned}$$

Hence by $\|K^{\theta^*}\|_\infty < \infty$ and $\|X^{\theta^*, \phi}\|_{S^2} \leq C$, $\|\phi^\theta(\cdot, X^{\theta^*, \phi}) - \phi^{\theta^*}(\cdot, X^{\theta^*, \phi^{\theta^*}})\|_{S^2} \leq C(\|K - K^{\theta^*}\|_\infty + \|k - k^{\theta^*}\|_\infty)$, which along with (B.1) yields the desired estimate. \square

Proof of Lemma 4.6. Recall that for all $m \in \mathbb{N}$, Algorithm 1 defines $K^{m+1} = K^{\theta_m}$ and $k^{m+1} = k^{\theta_m}$. Lemma 4.4 Item (1) and (H.2) imply that $\sup_{\theta \in \Theta} (\|K^\theta\|_\infty + \|k^\theta\|_\infty) < \infty$. By the definition (2.23), $(\theta_m)_{m \in \mathbb{N}}$ takes values in a bounded set Θ , and hence $\sup_{m \in \mathbb{N}} (\|K^m\|_{L^\infty} + \|k^m\|_{L^\infty}) \leq L$. The uniform boundedness of $\|\lambda^m\|_\infty$ follows from (2.11), $\inf_{t \in [0, T]} R_t > 0$, and the condition that $\sup_{m \in \mathbb{N}} \varrho_m < \infty$. Finally, for each $m \in \mathbb{N}$, by (2.22) and (2.23), θ_m is $\mathcal{F}_m^{\text{ob}}$ measurable, which along with the continuity of $\mathbb{R}^{1 \times 2} \times [0, T] \ni (\theta, t) \mapsto (K_t^\theta, k_t^\theta) \in \mathbb{R}^2$ leads to the desired $\mathcal{F}_m^{\text{ob}} \otimes \mathcal{B}([0, T])$ -measurability of K^{m+1} and k^{m+1} . \square

Proof of Lemma 4.8. By similar arguments as those for Lemma 4.6, $L := \sup_{m \in \mathbb{N}} (\|K^{\theta_m}\|_{L^\infty} + \|k^{\theta_m}\|_{L^\infty}) < \infty$, and for all $m \in \mathbb{N}$, K^{θ_m} and k^{θ_m} are $\mathcal{F}_{m-1}^{\text{ob}} \otimes \mathcal{B}([0, T])$ -measurable. Observe that for each $m \in \mathbb{N}$ and $t \in [0, T]$, K_t^{m+1} (resp. k_t^{m+1}) is a convex combination of $(K_t^{\theta_n})_{n=0}^m$ (resp. $(k_t^{\theta_n})_{n=0}^m$) according to the weights $(h_t^n)_{n=0}^m$. Hence K^m and k^m are $\mathcal{F}_{m-1}^{\text{ob}} \otimes \mathcal{B}([0, T])$ -measurable, and $\sup_{m \in \mathbb{N}} (\|K^{m+1}\|_{L^\infty} + \|k^{m+1}\|_{L^\infty}) \leq L$, independent of $\theta_0, V_0, (\varrho_m)_{m \in \mathbb{N}}$ and $(\pi_m)_{m \in \mathbb{N}}$. Finally, by (2.14), $\varrho_m > 0$ and $R_t \geq 0$ for all $t \in [0, T]$, we have $\lambda_t^{m+1} \leq \lambda_t^m$ for all $t \in [0, T]$, which along with the fact that $\|\lambda^0\|_\infty < \infty$ implies the uniform boundedness of $(\lambda^m)_{m \in \mathbb{N}}$. \square

References

- [1] M. BASEI, X. GUO, A. HU, AND Y. ZHANG, *Logarithmic regret for episodic continuous-time linear-quadratic reinforcement learning over a finite-time horizon*, Journal of Machine Learning Research, 23 (2022), pp. 1–34.
- [2] A. BECK AND M. TEBoulLE, *Mirror descent and nonlinear projected subgradient methods for convex optimization*, Operations Research Letters, 31 (2003), pp. 167–175.
- [3] N. CESA-BIANCHI, C. GENTILE, G. LUGOSI, AND G. NEU, *Boltzmann exploration done right*, Advances in Neural Information Processing Systems, 30 (2017).
- [4] S. N. COHEN AND T. TREETANTHIpLOET, *Asymptotic randomised control with applications to bandits*, arXiv preprint arXiv:2010.07252, (2020).
- [5] S. DEAN, H. MANIA, N. MATNI, B. RECHT, AND S. TU, *On the sample complexity of the linear quadratic regulator*, Foundations of Computational Mathematics, 20 (2020), pp. 633–679.
- [6] F. DELARUE AND A. VASILEIADIS, *Exploration noise for learning linear-quadratic mean field games*, arXiv preprint arXiv:2107.00839, (2021).
- [7] E. EVEN-DAR, S. MANNOR, Y. MANSOUR, AND S. MAHADEVAN, *Action elimination and stopping conditions for the multi-armed bandit and reinforcement learning problems.*, Journal of Machine Learning Research, 7 (2006).

- [8] M. K. S. FARADONBEH, M. S. S. FARADONBEH, AND M. BAYATI, *Thompson sampling efficiently learns to control diffusion processes*, arXiv preprint arXiv:2206.09977, (2022).
- [9] D. FIROOZI AND S. JAIMUNGAL, *Exploratory LQG mean field games with entropy regularization*, *Automatica*, 139 (2022), p. 110177.
- [10] X. GAO AND X. Y. ZHOU, *Logarithmic regret bounds for continuous-time average-reward markov decision processes*, arXiv preprint arXiv:2205.11168, (2022).
- [11] M. GEIST, B. SCHERRER, AND O. PIETQUIN, *A theory of regularized markov decision processes*, in *International Conference on Machine Learning*, PMLR, 2019, pp. 2160–2169.
- [12] X. GUO, A. HU, AND Y. ZHANG, *Reinforcement learning for linear-convex models with jumps via stability analysis of feedback controls*, arXiv preprint arXiv:2104.09311, (2021).
- [13] X. GUO, R. XU, AND T. ZARIPHOUPOULOU, *Entropy regularization for mean field games with learning*, *Mathematics of Operations Research*, (2022).
- [14] S. JAIMUNGAL, *Reinforcement learning and stochastic optimisation*, *Finance and Stochastics*, 26 (2022), pp. 103–129.
- [15] Y. JIA AND X. Y. ZHOU, *Policy gradient and actor-critic learning in continuous time and space: Theory and algorithms*, arXiv preprint arXiv:2111.11232, (2021).
- [16] G. KALLIANPUR, *Stochastic filtering theory*, vol. 13, Springer Science & Business Media, 2013.
- [17] B. KERIMKULOV, D. ŠIŠKA, AND L. SZPRUCH, *A modified msa for stochastic control problems*, *Applied Mathematics & Optimization*, (2021), pp. 1–20.
- [18] N. B. KESKIN AND A. ZEEVI, *On incomplete learning and certainty-equivalence control*, *Operations Research*, 66 (2018), pp. 1136–1167.
- [19] R. S. LIPTSER AND A. N. SHIRIAEV, *Statistics of random processes I: General theory*, vol. 394, Springer, 1977.
- [20] H. MANIA, S. TU, AND B. RECHT, *Certainty equivalent control of LQR is efficient*, arXiv preprint arXiv:1902.07826, (2019).
- [21] K. NICOLE EL, N. DU’HŪŪ, AND J.-P. MONIQUE, *Compactification methods in the control of degenerate diffusions: existence of an optimal control*, *Stochastics: an international journal of probability and stochastic processes*, 20 (1987), pp. 169–219.
- [22] C. REISINGER AND Y. ZHANG, *Regularity and stability of feedback relaxed controls*, *SIAM Journal on Control and Optimization*, 59 (2021), pp. 3118–3151.
- [23] J. SCHULMAN, F. WOLSKI, P. DHARIWAL, A. RADFORD, AND O. KLIMOV, *Proximal policy optimization algorithms*, arXiv preprint arXiv:1707.06347, (2017).
- [24] M. SIMCHOWITZ AND D. FOSTER, *Naive exploration is optimal for online LQR*, in *International Conference on Machine Learning*, PMLR, 2020, pp. 8937–8948.
- [25] R. S. SUTTON, A. G. BARTO, ET AL., *Introduction to reinforcement learning*, vol. 2, MIT press Cambridge, 1998.
- [26] L. SZPRUCH, T. TREETANTHIPLOET, AND Y. ZHANG, *Exploration-exploitation trade-off for continuous-time episodic reinforcement learning with linear-convex models*, arXiv preprint arXiv:2112.10264, (2021).
- [27] R. VERSHYNIN, *High-dimensional probability: An introduction with applications in data science*, vol. 47, Cambridge university press, 2018.

- [28] H. WANG, T. ZARIPHOPOULOU, AND X. Y. ZHOU, *Reinforcement learning in continuous time and space: A stochastic control approach*, Journal of Machine Learning Research, 21 (2020), pp. 1–34.
- [29] H. WANG AND X. Y. ZHOU, *Continuous-time mean–variance portfolio selection: A reinforcement learning framework*, Mathematical Finance, 30 (2020), pp. 1273–1308.
- [30] J. YONG AND X. Y. ZHOU, *Stochastic controls: Hamiltonian systems and HJB equations*, vol. 43, Springer Science & Business Media, 1999.
- [31] L. C. YOUNG, *Lectures on the calculus of variations and optimal control theory*, vol. 304, American Mathematical Soc., 2000.
- [32] J. ZHANG, *Backward Stochastic Differential Equations*, Springer, 2017.