

An Improved Trickle Down Theorem for Partite Complexes

Dorna Abdolazimi ^{*} and Shayan Oveis Gharan [†]

University of Washington

September 5, 2022

Abstract

Given a $d + 1$ -partite d -dimensional simplicial complex, we prove a generalization of the trickle-down theorem. We show that if “on average” faces of co-dimension 2 are $\frac{1-\delta}{d}$ -(one-sided) spectral expanders, then any face of co-dimension k is an $O(\frac{1-\delta}{k\delta})$ -(one-sided) spectral expander, for all $3 \leq k \leq d + 1$. For an application, using our theorem as a black-box, we show that links of faces of co-dimension k in recent constructions of bounded degree high dimensional expanders have local spectral expansion at most $O(1/k)$ fraction of the local expansion of worst faces of co-dimension 2.

1 Introduction

A simplicial complex X on a finite ground set $[n] = \{0, \dots, n\}$ is a downwards closed collection of subsets of $[n]$, i.e. if $\tau \in X$ and $\sigma \subset \tau$, then $\sigma \in X$. The elements of X are called *faces*, and the maximal faces are called *facets*. We say a face τ is of dimension k if $|\tau| = k + 1$ and write $\dim(\tau) = k$. A simplicial complex X is a *pure* d -dimensional complex if every facet has dimension d . In this paper, all simplicial complexes are assumed to be pure. Given a d -dimensional complex X , for any $0 \leq i \leq d$, define $X(i) := \{\tau \in X : \dim(\tau) = i\}$. Moreover, the co-dimension of a face $\tau \in X$ is defined as $\text{codim}(\tau) := d - \dim(\tau)$. For a face $\tau \in X$, define the *link* of τ as the simplicial complex $X_\tau = \{\sigma \setminus \tau : \sigma \in X, \sigma \supset \tau\}$. Note that X_τ is a $(\text{codim}(\tau) - 1)$ -dimensional complex.

A $(d + 1)$ -partite complex is a d -dimensional complex such that $X(0)$ can be (uniquely) partitioned into sets $T_0 \cup \dots \cup T_d$ such that for every facet $\tau \in X(d)$, we have $|\tau \cap T_i| = 1$ for all $i \in [d]$. The type of any face $\tau \in X$ is defined as $\text{type}(\tau) = \{i \in [d] : |\tau \cap T_i| = 1\}$.

We equip X with a probability distribution π supported on all *facets* of X and we denote this pair by (X, π) . For a face $\tau \in X$, π induces a conditional distribution π_τ on facets of X_τ where for each facet $\sigma \in X_\tau$,

$$\pi_\tau(\sigma) = \frac{\pi(\sigma \cup \tau)}{\sum_{\text{facet } \sigma' \in X_\tau} \pi(\sigma' \cup \tau)}.$$

For each face τ of co-dimension at least 2 the 1-skeleton of (X_τ, π_τ) is a weighted graph with vertices $X_\tau(0)$, edges $X_\tau(1)$, and edge weights given by $\mathbb{P}_{\sigma \sim \pi_\tau}[\{x, y\} \subseteq \sigma]$ for each edge $\{x, y\}$.

^{*}dormaa@cs.washington.edu. Research supported by NSF grant CCF-1907845 and Air Force Office of Scientific Research grant FA9550-20-1-0212.

[†]shayan@cs.washington.edu. Research supported by Air Force Office of Scientific Research grant FA9550-20-1-0212, Simons Investigator grant.

Note that when τ is of co-dimension 2, the complex (X_τ, π_τ) is itself a weighted graph. We say a complex X is totally connected if the 1-skeleton of the link of any face τ of co-dimension at least 2 is connected.

Definition 1 (Local Spectral High Dimensional Expander). *A link (X_τ, π_τ) of co-dimension at least 2 of a d -dimensional (weighted) complex (X, π) is a λ -(one sided) spectral expander if the second (largest) eigenvalue of the simple random walk on the 1-skeleton of (X_τ, π_τ) is at most λ . We say (X, π) is a $(\gamma_2, \gamma_3, \dots, \gamma_{d+1})$ -local spectral expander if for any face τ of co-dimension at least 2, (X_τ, π_τ) is a $\gamma_{\text{codim}(\tau)}$ spectral expander. When the complex is clear in the context, for an integer $2 \leq k \leq d+1$, we write γ_k to denote the largest 2nd eigenvalue of all links of co-dimension k in x .*

Over the last few years study of local spectral HDX have revolutionized several areas of Math and theoretical CS, namely in analysis of Markov chains [Ana+19; ALO20; CLV21; ALO21], coding theory [Din+22], etc [AJT19; Din+21b; Din+21a]. One can generally divide the family of HDXes studied in recent works into two groups: (i) Dense Complexes. Here, we have a HDX with exponentially large number of facets, i.e., $|X(0)|^d$. One typically encounters these objects in studying Markov Chain Monte Carlo technique where we use a Markov Chain to sample from an exponentially large probability distribution. Perhaps the simplest such family is the complex of all independent sets of a matroid. (ii) Sparse/Ramanujan Complexes. Here we have a HDX where every vertex (of $X(0)$) only appear in constant number of facets, independent of $|X(0)|$. See, [LSV05; KO18; OP22] for explicit constructions. These objects have been useful in constructing double samplers [Din+21b], agreement testers [DD19], or locally testable codes [Din+22].

One of the main aspect of local spectral expanders is their local to global phenomenon often referred to as the Garland’s method or the trickle down theorem [Opp18].

Theorem 1.1 (Trickledown Theorem). *Given a totally connected complex (X, π) , if $\gamma_2 \leq \frac{1-\delta}{d}$ for some $0 < \delta \leq 1$, then $\gamma_k \leq \frac{1-\delta}{d\delta}$ for all $2 \leq k \leq d$.*

The trickle down theorem has found numerous applications in proving bounds on local spectral expansion of simplicial complexes. To invoke the theorem one needs to inspect all faces of co-dimension 2 to find the worst 2nd eigenvalue. If we get lucky and this number is below $1/d$, then, the trickledown theorem kicks in and inductively bounds local spectral expansion of all faces of the complex.

There are, however, two pitfalls for the theorem: i) The required bound on γ_2 is too small and often not satisfiable. In particular, for most dense complexes in counting and sampling applications (see e.g., [ALO20; CLV21]) we have $\gamma_k = O(1/d)$ for $k \geq \Omega(d)$, even though the faces of co-dimension 2 are only $\Theta(1)$ -spectral expanders. ii) Even if $\gamma_2 \ll 1/d$, the trickledown theorem only implies $\gamma_k \simeq \gamma_2$, i.e., γ_k does not increase too much. This is in contrast with the fact that one often observes a steep decrease in spectral expansion as the co-dimension increases in dense complexes, i.e., $\gamma_k \lesssim \gamma_2/k$.

These pitfalls, (especially (ii)) convinced some experts to conjecture that, perhaps, dense and sparse complexes exhibit a different pattern of local spectral expansion; in particular, unlike dense HDX, local spectral expansion does not decay for sparse complexes.

In this paper, we prove a generalization of the trickle-down theorem for *partite complexes* that shows that even if $\gamma_2 = \Theta(1)$, but “on average” the faces of co-dimension 2 are $< 1/d$ -spectral expanders, then we have $\gamma_k \leq O(1/k)$ for all $3 \leq k \leq d+1$. Surprising to us, our average condition is satisfied by some recent construction of (sparse) bounded degree high dimensional expanders [KO18]. In particular, as we explain below, one can use our theorem to prove a significantly better local spectral expansion for the Kaufman-Openheim construction in a black-box manner.

1.1 Main Contribution

For every integers $1 \leq n$, let H_n be the n -th harmonic number, i.e $H_n = \sum_{i=1}^n \frac{1}{i}$. Moreover, for any $1 \leq i \leq n$ define $H_n(i) := \sum_{j=i}^n \frac{1}{j}$ and let $H_n(0) = H_n(1)$.

Theorem 1.2 (Main). *Let (X, π) be a $(d+1)$ -partite (weighted) totally connected complex. For any distinct $i, j \in [d]$, let $\epsilon_{\{i,j\}} = \max_{\tau: \text{type}(\tau)=[d] \setminus \{i,j\}} \lambda_2(P_\tau)$ be the largest 2nd eigenvalue of the simple random walk matrices on (X_τ, π_τ) for all τ of type $[d] \setminus \{i, j\}$. Let $\Delta(i) := |\{j \in [d] \setminus i : \epsilon_{\{i,j\}} > 0\}|$ for all $i \in [d]$ and $\Delta := \max_{i \in [d]} \Delta(i)$. For some $0 < \delta < 1$, assume that for every $i \in [d]$,*

$$\epsilon_{\{i,j\}} \cdot H_{\Delta-1} \leq \frac{\delta^2}{10}, \forall j \neq i \quad \text{and} \quad (1)$$

$$\sum_{\ell=1}^{\Delta(i)} \epsilon_{\{i,j_\ell\}} \cdot H_{\Delta(i)-1}(\ell-1) \leq 1 - \delta, \quad (2)$$

where j_1, \dots, j_d is an ordering of $[d] \setminus i$ such that $\epsilon_{\{i,j_1\}} \leq \dots \leq \epsilon_{\{i,j_d\}}$. Then, X is $(\frac{c(1-\delta)}{\delta}, \dots, \frac{c(1-\delta)}{d\delta})$ -local spectral expander for $c = \frac{2(1+\frac{\delta^2}{10})}{(1+\delta)}$.

Remark 1. We remark that for every any $i \in [d]$, $\Delta(i) \leq \sum_{\ell=1}^{\Delta(i)} H_{\Delta(i)-1}(\ell-1) \leq \Delta(i) + \ln(\Delta(i))$. So, if $\epsilon_{\{i,j\}} = \epsilon_i$ for all $j \neq i$, i.e., they are invariant over j , the assumption of the theorem simplifies to $\epsilon_i(\Delta(i) + \ln(\Delta(i))) \leq 1 - \delta$. Roughly speaking, the latter condition implies that $\mathbb{E}_j[\epsilon_{\{i,j\}}] \leq \frac{1-\delta}{d}$ for every $i \in [d]$, where the expectation is over $j \neq i$ chosen uniformly at random. This is an improvement over the stronger condition in [Theorem 1.1](#).

Applications to Graph Coloring. Given a graph $G = (V, E)$ with degree function $\Delta : V \rightarrow \mathbb{Z}_{\geq 0}$ and maximum degree Δ , paired with a collection of color lists $\{L(v)\}_{v \in V}$ satisfying $L(v) \geq \Delta(v) + (1 + \eta)\Delta$ for all $v \in V$ and for some $0 < \eta \leq 0.9$ such that $\frac{1+\ln \Delta}{\Delta} \leq \frac{\eta^2}{40}$, if we apply the above theorem to the coloring complex $X(G, L)$ with $\delta = \frac{\eta}{2}$, we get that $X(G, L)$ is a $(\frac{4}{\eta}, \frac{4}{2\eta}, \dots, \frac{4}{(|V|-1)\eta})$ -local spectral expander, and thus the Glauber dynamics for sampling a random proper coloring mixes in polynomial time. This retrieves (up to constants) a theorem proved in [\[ALO21\]](#).

Applications to Sparse High Dimensional Expanders Kaufman and Oppenheim [\[KO18\]](#) obtained a simple construction of sparse $d + 1$ -partite complexes with $|X(0)| \geq p^s$ for any integer $s > d$ and prime power p such that every $x \in X(0)$ is in at most $p^{O(d^3)}$ many facets (hence the degree is independent of s). They argued that for any non-consecutive pair of parts $i, j \in [d]$, i.e., $j \neq i + 1$ and $i \neq j + 1 \pmod{d + 1}$, we have $\epsilon_{\{i,j\}} = 0$ but $\epsilon_{\{i,i+1\}} \leq \frac{1}{\sqrt{p}}$ for any $i \in [d]$ ($i + 1$ is taken modulo $d + 1$). Consequently, $\Delta(i) = 2$ for any $i \in [d]$. Then, using [Theorem 1.1](#), they show that the complex is a $(\frac{1}{\sqrt{p-(d-2)}}, \dots, \frac{1}{\sqrt{p-d-2}})$ -local spectral expander for $p > (d-2)^2$. Simply plugging in these values into the above theorem, for $\delta = 1 - \frac{2}{\sqrt{p}}$ and $p \geq 193$ (independent of d) the assumptions of the theorem are satisfied. The resulting complex is $(\frac{2c}{\sqrt{p\delta}}, \dots, \frac{2c}{d\sqrt{p\delta}})$ -local spectral expander for $c \approx 1.15$. In other words, not only does the Kaufman-Oppenheim construction give a HDX for constant values of p independent of d , but also its local spectral expansion improves inverse linearly with the co-dimension.

O'Donnell and Pratt [\[OP22\]](#) constructed $d + 1$ -partite (sparse) high-dimensional expanders, with unbounded dimension d , via root systems of simple Lie Algebras, namely families A_d for

$d \geq 1$, B_d for $d \geq 2$, C_d for $d \geq 3$ and D_d for $d \geq 4$. For explicit descriptions of these root systems, see e.g. [Car89, Sec. 3.6]. O’Donnell and Pratt [OP22] showed that, similar to the Kaufman-Oppenheim construction, the resulting d -dimensional complex X satisfies $|X(0)| \geq p^{\Theta(m)}$ whereas every vertex is only in $p^{\Theta(d^2)}$ many facets and for any $i, j \in [d]$, $\epsilon_{i,j} \leq \sqrt{2/p}$. Then, using [Theorem 1.1](#) they concluded that the complex is a $(\frac{1}{\sqrt{p/2-d+1}}, \dots, \frac{1}{\sqrt{p/2-d+1}})$ -local spectral expander. Upon further inspection of the explicit set of roots, one can verify that $\Delta \leq 2$ for complexes based on A_d, B_d, C_d root systems and $\Delta \leq 3$ for the D_d root system. Plugging in these values in the above theorem and setting $\delta = 1 - 2\sqrt{2/p}$ for A_d, B_d, C_d complexes and $\delta = 1 - 3.5\sqrt{2/p}$ for the D_d complex, if $p \geq 376$ for A_d, B_d, C_d complexes and $p \geq 729$ for the D_d complex, we get that these complexes are $(\frac{c'}{\sqrt{p\delta}}, \dots, \frac{c'}{d\sqrt{p\delta}})$ -local spectral expander for some constant $c' > 1$.

1.2 Proof Overview

At high-level our proof builds on the Matrix trickledown framework introduced in the work of the authors with Liu [ALO21]. Our technical contribution are two fold: First, we observe that the same framework can be extended to “sparse” complexes with a carefully chosen family of (diagonal) matrices that are recursively defined based on the spectral expansion of links of co-dimension 2. See proof of [Theorem 1.2](#) for the construction of these matrices. Second, we observe that for any two disjoint sets of parts $S, T \subseteq [d]$, if all links of co-dimension two that their types intersect both S, T are 0-spectral expanders then for any $\sigma \in X$ of type S and τ of type T ,

$$\mathbb{P}_{\eta \sim \pi}[\sigma \subset \eta | \tau \subset \eta] = \mathbb{P}_{\eta \sim \pi}[\sigma \subset \eta] \quad \text{and} \quad \mathbb{P}_{\eta \sim \pi}[\tau \subset \eta | \sigma \subset \eta] = \mathbb{P}_{\eta \sim \pi}[\tau \subset \eta].$$

namely the conditional distributions on these types are independent (see [Lemma 3.2](#) for details). This observation significantly simplifies invoking the Matrix trickledown framework.


1.3 Obstructions to Trickledown

We end this section by exhibiting some barriers to any class of trickledown theorems that only look at 2nd eigenvalue of links of co-dimension 2.

Example 1. We exhibit a family of partite complexes such that $\gamma_2 \leq \lambda$, but $\gamma_{2d+1} \geq \Omega(\lambda)$. Consider a $2d$ -partite ($2d - 1$ dimensional) complex (X, π) where $T_i = \{i, \bar{i}\}$. A set $\tau \in T_1 \times \dots \times T_{2d}$ is a facet of X iff whenever $i, j \in \tau$ we either have $i, j \leq d$ or $i, j \geq d + 1$. For simplicity of calculations, for such a τ , we define $w(\tau) = \lambda^{|\tau|_1}$ where $|\tau|_1$ is the number of $i \in \tau$, for some $0 < \lambda < 1$ that we choose later, i.e., we do not normalize the weights in a probability distribution.

As a side note, facets of this complex corresponds to the set of independent sets of $K_{d,d}$ the complete bipartite graph on the sets $\{1, \dots, d\}, \{d + 1, \dots, 2d\}$.

It is not hard to see that we have $\gamma_2 = \frac{\lambda}{1+\lambda}$. To see this, note that a worst link to take is $\tau = \{\bar{2}, \dots, \overline{2d-1}\}$ with the the following 1-skeleton. Notice that this graph is a $\frac{\lambda}{1+\lambda}$ -spectral expander.



We claim that the 2nd eigenvalue of the 1-skeleton of the link of the empty set is at least $\Omega(\lambda^2)$. First notice that this graph has $4d$ vertices. We partition its vertices into 4 sets, $A = \{1, \dots, d\}$, $\bar{A} = \{\bar{1}, \dots, \bar{d}\}$ and $B = \{d + 1, \dots, 2d\}$, $\bar{B} = \{\overline{d+1}, \dots, \overline{2d}\}$.

For simplicity of calculations, we can write the weight of every edge of the 1-skeleton as the sum of the weights of all facets that contain that edge. For every $i \in A, j \in \bar{A}$,

$$w_{i,j} = \sum_{k=0}^{d-2} \binom{d-2}{k} \lambda^{k+1} = \lambda(1+\lambda)^{d-2}$$

Running a similar calculation for all possible i, j pairs, we obtain the following 1-skeleton of the empty set (after dividing all edge weights by $(1+\lambda)^{d-2}$).




Figure 1: 1-skeleton of the Link of \emptyset for complex **Example 1**. Edges of the same color have the same weight.

We assume $1/d \ll \lambda \ll 1$ and $d \rightarrow \infty$ so we ignore $(1+\lambda)^{-(d-2)}$ low order term. It follows that $d_w(i) \approx 2d\lambda(1+\lambda)$ for $i \in X \cup Y$ and $d_w(\bar{i}) \approx 2d(2+3\lambda+\lambda^2)$. It follows that,


$$\left| w(E(A \cup B)) - \frac{\text{vol}(A \cup B)^2}{\text{vol}(V)} \right| \geq \left| d^2 \lambda^2 - \frac{(2d(2d\lambda(1+\lambda)))^2}{2d(2d(2+4\lambda+2\lambda^2))} \right| = |d^2 \lambda^2 - 2d^2 \lambda^2|$$

So, by **Fact 2.2** for $S = A \cup B$ we have

$$\lambda_2 \geq \frac{d^2 \lambda^2}{\text{vol}(A \cup B)} = \frac{d^2 \lambda^2}{4d^2 \lambda(1+\lambda)} \geq \frac{\lambda}{4(1+\lambda)}.$$


Note that if we apply **Theorem 1.2** to the setting of the above example, we obtain $\Delta(i) = d$ for all $1 \leq i \leq 2d$. So, if $\lambda \cdot d \leq 1 - \delta$, (and $\lambda d \log d \leq \delta^2/10$), our theorem implies that the 2nd eigenvalue of the link of the emptyset is at most $\frac{c(1-\delta)}{2d-\delta} \leq \frac{c\lambda}{\delta}$ which is consistent with the above calculations.

Example 2. In this example we construct a totally connected (non-partite) $d-1$ -dimensional weighted simplicial complex (X, π) such that links of co-dimension 2 are 0.5-spectral expanders, but the 1-skeleton of the link of the emptyset is only a $1 - \exp(-d)$ -spectral expander. This in particular shows that local spectral expansion can increase significantly for (non-partite) complexes. Let $B(V, E)$ be the following barbell graph: Consider two disjoint cliques K_1, K_2 each with $2d$ vertices and connect them with a path of length $d+2$ namely $x_0 \in K_1, x_1, \dots, x_d, x_{d+1} \in K_2$. Note that x_1, \dots, x_d do not belong to the cliques.



Now we define a $d - 1$ -dimensional weighted complex on vertices V . The facets of X are precisely sets $S \subseteq V$ with $|S| = d$ such that the induced graph $B[S]$ is connected. For simplicity of calculations we define the weight of every facet to be 1, i.e., we don't normalize the weights to be a probability distribution.

We claim that any link X_τ of co-dimension 2 is $\Omega(1)$ -spectral expander. Indeed the worst link is X_τ for $\tau = \{x_2, \dots, x_{d-1}\}$. The 1-skeleton of X_τ is the following graph with second eigenvalue 0.5.



Now, we claim that the 1-skeleton of X_\emptyset , G_0 , has 2nd eigenvalue $1 - \frac{1}{\exp(d)}$. First notice G_0 has the same vertex set as B . Now, consider the set $S = K_1 \cup \{x_0, x_1, \dots, x_{d/2}\}$. For simplicity of calculations, we let the weight of every edge $\{x, y\}$ of G_0 be sum of the weights of all facets that contains x, y . First, observe that the weight of every edge in K_1 is (at least) $\binom{2d-2}{d-2}$. On the other hand, the weight of any edge in the cut (S, \bar{S}) is at most $2\binom{2d}{d/2}$. Putting these together it follows that

$$\phi(S) = \frac{w(E(S, \bar{S}))}{\text{vol}(S)} \geq \frac{2\frac{d}{2} \cdot \frac{5d}{2} \cdot \binom{2d}{d/4}}{\binom{2d}{2} \cdot \binom{2d-2}{d-2}} \approx 2^{-d/2},$$

for a large enough d . Therefore, by Cheeger's inequality, [Lemma 2.1](#), the second eigenvalue of G_0 is at least $1 - 2^{-\Omega(d)}$ for d large enough.

1.4 Acknowledgments

The discussion that initiated this work took place at DIMACS Workshop on Entropy and Maximization. We would like to thank the DIMACS center and the workshop organizers for making this happen. In particular, we would like to thank Tali Kaufman for raising the question of an improved trickledown theorem for sparse simplicial complexes in that workshop. We also would like to thank Ryan O'Donnell and Kevin Pratt for helpful discussions on high dimensional expanders based on Chevalley groups.

2 Preliminaries

For any integer $n \geq 0$, let $[n] := \{0, \dots, n\}$. When it is clear from context, we write x to denote a singleton $\{x\}$. Given a set S , we write $v \in \mathbb{R}^S$ and $M \in \mathbb{R}^{S \times S}$ to respectively denote a vector and a matrix indexed by S . Given a probability distributions μ over a set S , we may view μ as a vector in $\mathbb{R}_{\geq 0}^S$. For a $n \times n$ matrix M with eigenvalues $\lambda_1, \dots, \lambda_n$, define $\rho(A) := \max_{1 \leq i \leq n} |\lambda_i|$.

Graphs Given a graph $G = (V, E)$, for any $v \in V$, let $\Delta_G(v)$ be the degree of v in G , and let Δ_G be the maximum degree of G . Moreover, given a subset $S \subseteq V$, $G[S]$ denotes the induced subgraph of G on the set of vertices S . For any $S \subseteq V$, define $G_S := G[V \setminus S]$. For simplicity of notation, when G is clear from context, we denote $\Delta_G(v)$ by $\Delta(v)$ for any $v \in V$, and for any $S \subseteq V$, we denote $\Delta_{G_S}(v)$ by $\Delta_S(v)$ for any $v \in V \setminus S$. Similarly, we denote the maximum degree of G and G_S by Δ and Δ_S respectively. Moreover, when G is clear from context, we write $u \sim v$ if u, v are adjacent vertices in G and $u \sim_S v$ if $u, v \in V \setminus S$ and $u \sim v$.

We say a graph $G = (V, E)$ paired with a weight function $w : E \rightarrow \mathbb{R}_{\geq 0}$ is ϵ -expander if $\lambda_2(P) \leq \epsilon$, where $P \in \mathbb{R}^{V \times V}$ is the transition probability matrix of the simple random walk on (G, w) defined as $P(x, y) = \frac{w(\{x, y\})}{\sum_z w(\{x, z\})}$ for any $x, y \in V$. For such a graph we write $d_w(x) =$

$\sum_{y \sim x} w(\{x, y\})$ to denote the weighted degree of a vertex x and $\text{vol}(S) = \sum_{x \in S} d_w(x)$ to denote the volume of a set $S \subseteq V$.

2.1 Linear Algebra

Lemma 2.1 (Cheeger's Inequality). *For any graph $G = (V, E)$ with weights $w : E \rightarrow \mathbb{R}_{\geq 0}$ and any $S \subseteq V$,*

$$\frac{w(E(S, \bar{S}))}{\min\{\text{vol}(S), \text{vol}(\bar{S})\}} \leq \sqrt{2(1 - \lambda_2)}$$

where λ_2 is the second largest eigenvalue of the simple random walk on G

Fact 2.2 (Expander Mixing Lemma). *Given a (weighted) graph $G = (V, E, w)$, for any set $S \subseteq V$,*

$$\left| w(E(S)) - \frac{\text{vol}(S)^2}{\text{vol}(V)} \right| \leq \lambda_2 \text{vol}(S),$$

where λ_2 is the second largest eigenvalue of the simple random walk on G .

2.2 Simplicial Complexes

We say a simplicial complex X is *gallery connected* if for any face τ of co-dimension at least 2 and any pair of facets σ, σ' of X_τ there is a sequence of facets of X_τ , $\sigma = \sigma_0, \sigma_1, \dots, \sigma_\ell = \sigma'$, such that for all $0 \leq i < \ell$, $|\sigma_i \Delta \sigma_{i+1}| = 1$. It is shown in [Opp18, Prop 3.6] that if X is totally connected then it is gallery connected.

Lemma 2.3. *Consider a totally connected $(d + 1)$ -partite complex X with parts indexed by $[d]$. For any $S \subseteq [d]$, The induced subgraph of the 1-skeleton of X on vertices of type S is connected.*

Proof. Take x, y of type $i, j \in S$ and facets η, η' such that $x \in \eta, y \in \eta'$. Total connectivity implies that there is a sequence $\eta = \eta_1, \dots, \eta_t = \eta'$ such that $\eta_i \cap \eta_{i+1} \neq \emptyset$ for all $1 \leq i \leq t - 1$. Let $\sigma_1 \subseteq \eta_1, \dots, \sigma_t \subseteq \eta_t$ be faces of type $\{i, j\}$. Then $\sigma_1, \dots, \sigma_t$ gives a path between x, y . \square

Given a (weighted) complex (X, π) , for integer $-1 \leq i \leq \dim(X) - 1$, π induces a distribution π_i on $X(i)$,

$$\pi_i(\sigma) = \frac{1}{\binom{\dim(X)+1}{i+1}} \Pr_{\tau \sim \pi}[\sigma \subset \tau] \quad \forall \sigma \in X(i).$$

Let $P_{(X, \pi), \tau} \in \mathbb{R}^{X(0) \times X(0)}$ denote the transition probability matrix of the simple random walk on the 1-skeleton of (X_τ, π_τ) padded with zeros outside the $X_\tau(0) \times X_\tau(0)$ block, i.e. $P_{(X, \pi), \tau}(x, y) = \frac{\mathbb{P}_{\sigma \sim \pi_\tau}[\{x, y\} \subset \sigma]}{\sum_{z \in X_\tau(0)} \mathbb{P}_{\sigma \sim \pi_\tau}[\{x, z\} \subset \sigma]}$ for $x, y \in X_\tau(0)$, and $P_\tau(x, y) = 0$ otherwise. When the weighted complex (X, π) is clear from context, we write P_τ to denote $P_{(X, \pi), \tau}$. For any τ of co-dimension at least 2, we define the diagonal matrix $\Pi_{(X, \pi), \tau} \in \mathbb{R}^{X(0) \times X(0)}$ as follows: $\Pi_{(X, \pi), \tau}(x, x) = \pi_{\tau, 0}(x)$ for $x \in X_\tau(0)$, and $\Pi_{(X, \pi), \tau}(x, x) = 0$ otherwise. When (X, π) is clear from context, we write Π_τ to denote $\Pi_{(X, \pi), \tau}$. Note that $\Pi_\tau P_\tau$ is a symmetric matrix.

Given a $(d + 1)$ -partite complex, we say an $x \in X(0)$ is of type i and write $\text{type}(x) = i$ if $x \in T_i$. Similarly, the type of a face $\tau \in X$ is defined as $\text{type}(\tau) = \{i \in [d] : |\tau \cap T_i| = 1\}$.

Fact 2.4. *Consider a weighted $(d + 1)$ partite complex (X, π) and a face τ of co-dimension $k \geq 1$. We have $k\pi_{\tau, 0}(x) = \Pr_{\sigma \sim \pi_\tau}[x \in \sigma]$ for all $x \in X_\tau(0)$.*

Fact 2.5. Consider a weighted $(d + 1)$ partite complex (X, π) with parts indexed by $[d]$ and a face τ of co-dimension $k \geq 1$. For any $i \in [d]$, $\sum_{x:\text{type}(x)=i} \Pr_{\sigma \sim \pi_\tau}[x \in \sigma] = 1$.

Fix $(d + 1)$ -partite complex X with parts indexed by $[d]$. Define a graph $G_{(X, \pi)}$ on the set of vertices $[d]$, where any distinct $i, j \in [d]$ are adjacent in $G_{(X, \pi)}$ if there exists τ of type $[d] \setminus \{i, j\}$ such that the second eigenvalue of (X_τ, π_τ) is positive. Note that if $\text{codim}(\tau) = k$, the link X_τ is a k -partite complex with parts indexed by $[d] \setminus S$. One can verify that given a face τ of type S , the set of edges of $G_{(X_\tau, \pi_\tau)}$ is a subset of the edges of $(G_{(X, \pi)})_S$, i.e., the induced subgraph of $G_{(X, \pi)}$ on $[d] \setminus S$. When (X, π) is clear from context, we write G for $G_{(X, \pi)}$ and G_S for $(G_{(X, \pi)})_S$.

Product of Weighted Complexes Given weighted complexes $(Y_1, \mu_1), \dots, (Y_\ell, \mu_\ell)$ defined on disjoint ground sets and of dimensions d_1, \dots, d_ℓ respectively, and a weighted complex (X, π) of dimension d , we write $(X, \pi) = (Y_1, \mu_1) \times \dots \times (Y_\ell, \mu_\ell)$ if $X(d) = \{\cup_{i \in [\ell]} \tau_i : \tau_1 \in Y_1(d_1), \dots, \tau_\ell \in Y_\ell(d_\ell)\}$ and $\pi(\cup_{i \in [\ell]} \tau_i) = \prod_{i \in [\ell]} \mu_i(\tau_i)$ for all $\tau_1 \in Y_1(d_1), \dots, \tau_\ell \in Y_\ell(d_\ell)$. We denote the generating polynomial of (X, π) by $g_{(X, \pi)}$, i.e. $g_{(X, \pi)} := \sum_{\tau \in X(d)} \pi(\tau) \prod_{x \in \tau} z_x$. One can verify that $(X, \pi) = (X_1, \mu_1) \times \dots \times (X_\ell, \mu_\ell)$ if and only if $g_{(X, \pi)} = g_{(X_1, \mu_1)} \times \dots \times g_{(X_\ell, \mu_\ell)}$. Note that this is true because we assume that for any weighted simplicial complex, the given distribution on facets is non-zero on all facets.

Matrix Trickle-Down Theorem We use the following theorem which is the main technical theorem in [ALO21].

Theorem 2.6 ([ALO21, Thm III.5]). Let (X, π_d) be a totally connected weighted complex. Suppose $\{M_\tau \in \mathbb{R}^{X(0) \times X(0)}\}_{\tau \in X(\leq d-2)}$ is a family of symmetric matrices satisfying the following:

1. **Base Case:** For every τ of co-dimension 2, we have the spectral inequality

$$\Pi_\tau P_\tau - 2\pi_\tau \pi_\tau^\top \preceq M_\tau \preceq \frac{1}{5} \Pi_\tau.$$

2. **Recursive Condition:** For every τ of co-dimension at least $k \geq 3$, at least one of the following holds: M_τ satisfies

$$M_\tau \preceq \frac{k-1}{3k-1} \Pi_\tau \quad \text{and} \quad \mathbb{E}_{x \sim \pi_\tau} M_{\tau \cup \{x\}} \preceq M_\tau - \frac{k-1}{k-2} M_\tau \Pi_\tau^{-1} M_\tau. \quad (3)$$

Or, $(X_\tau, \pi_{\tau, k-1})$ is a product of weighted simplicial complexes $(Y_1, \mu_1), \dots, (Y_t, \mu_t)$ and for every $\eta \in X_\tau(k-1)$,

$$M_\tau = \bigoplus_{1 \leq i \leq t: d_{Y_i} \geq 1} \frac{d_{Y_i}(d_{Y_i} + 1)}{k(k-1)} M_{\tau \cup \eta_{-i}},$$

where $\eta_{-i} = \eta \setminus Y_i(0)$.

Then for every $\tau \in X(\leq d-2)$, we have the bound $\lambda_2(\Pi_\tau P_\tau) \leq \rho(\Pi_\tau^{-1} M_\tau)$.

3 Proof of Main Theorem

Lemma 3.1. Consider a 2-partite complex (X, π) with parts S, T . If (X, π) is 0-expander, then $(X, \pi) = (X_z, \pi_z) \times (X_y, \pi_y)$ for any $y \in S$ and $z \in T$.

Proof. Note that (X, π) is a weighted bipartite graph with parts S, T . Let $A \in \mathbb{R}^{X^{(0)} \times X^{(0)}}$ be the adjacency matrix of (X, π) . Let $A_{S,T}(y, z) = A(y, z)$ for $y \in S, z \in T$ and 0 on other entries. Moreover, let $A_{T,S} = A - A_{S,T}$. Then, for any vector $v \in \mathbb{R}^{X^{(0)}}$, we get $A v = A_{S,T} v_T + A_{T,S} v_S$, where v_S, v_T are respectively supported on S, T and $v = v_S + v_T$. Thus, if $A v = \lambda v$, then $A v' = -\lambda v'$, for $v' = (-v_S + v_T)$. So if μ is an eigenvalue of A , then $-\mu$ is also an eigenvalue of A . Thus, if (X, π) is 0-expander, the rank of A is 2. This implies that there are vectors $w_S \in \mathbb{R}^S$ and $w_T \in \mathbb{R}^T$ such that $\pi(\{y, z\}) = A(y, z) = A(z, y) = w_S(y)w_T(z)$ for $y \in S, z \in T$. Without loss of generality, assume $\|w_S\|_1 = \|w_T\|_1 = 1$. Then, for any $y \in S$ and $z \in T$, we have $\pi_{\{z\}} = \sum_{y \in S} \pi(\{y, z\}) = w_S$, and similarly $\pi_{\{y\}} = w_T$. Thus $\pi(\{y, z\}) = \pi_y(z)\pi_z(y)$. This finishes the proof. \square

Lemma 3.2. Consider a totally connected $(d + 1)$ -partite complex (X, π) with parts indexed by $[d]$ and its associated graph $G := G_{(X, \pi)}$. Let $I_1 \cup \dots \cup I_\ell$ be a partition of $[d]$ such that for any $1 \leq i \leq \ell$ the induced graph $G[I_i]$ is the union of some connected components of G . Then $(X, \pi) = (X_{\sigma_{-1}}, \pi_{\sigma_{-1}}) \times \dots \times (X_{\sigma_{-\ell}}, \pi_{\sigma_{-\ell}})$, where σ_{-i} is an arbitrary face of type $[d] \setminus I_i$ for any $1 \leq i \leq \ell$.

Proof. We prove the statement by induction on d . For $d = 1$, the statement simply follows from [Lemma 3.1](#). Now, assume that $d > 1$. If $|I_i| = 1$ for all $1 \leq i \leq \ell$, then $\ell \geq 3$. In this case, let $S := I_1 \cup I_2$. Otherwise, WLOG assume that $|I_1| \geq 2$ and let $S := I_1$. First, we show that $g_{(X, \pi)}$ can be written as $g_{(X, \pi)} = h \cdot h'$, where h is a polynomial in $\{z_y : \text{type}(y) \in I \setminus S\}$ and h' is a polynomial in terms of variables in $\{z_y : \text{type}(y) \in S\}$. By induction hypothesis, for any $i \in S, x \in T_i$, and any face $\sigma \in X$ of type S such that $x \in \sigma$

$$\partial_{z_x} g_{(X, \pi)} = f^x \cdot g^x \tag{4}$$

where f^x is a polynomial in terms of variables in $\{z_y : \text{type}(y) \in S \setminus i\}$ and g^x is a polynomial in terms of variables in $\{z_y : \text{type}(y) \in I \setminus S\}$. Now, take arbitrary $i, j \in S$ such that $i \neq j$. Then, (4) implies that for any face $\{x, y\}$ of type $\{i, j\}$

$$\partial_{z_x} \partial_{z_y} g_{(X, \pi)} = (\partial_{z_y} f^x) g^x = (\partial_{z_x} f^y) g^y$$

It thus follows that g^x is a multiple of g^y . One can see this simply by substituting 1 for all variables in $\{z_y : \text{type}(y) \in S \setminus \{i, j\}\}$. Moreover, since g^x and g^y are generating polynomials of distributions, i.e. the coefficients sum up to 1, we get $g^x = g^y$. Therefore, we get that for any distinct x, y such that $\text{type}(x), \text{type}(y) \in S$ and $\{x, y\}$ is a face, $g^x = g^y$. Applying [Lemma 2.3](#), we get $g^x = g^y$ for all $x, y \in \cup_{i \in S} T_i$. Thus, there exist a polynomial h in variables $\{z_y : \text{type}(y) \in I \setminus S\}$ such that we can rewrite (4) for any x with $\text{type}(x) \in S$ as

$$\partial_{z_x} g_{(X, \pi)} = f^x \cdot h,$$

where f^x is a polynomial in terms of variables in $\{z_y : \text{type}(y) \in S \setminus i\}$. Finally, since X is a partite complex,

$$|S| g_{(X, \pi)} = \sum_{i \in S} \sum_{x \in T_i} z_x \partial_{z_x} g_{(X, \pi)} = h \cdot \sum_{i \in S} \sum_{x \in T_i} z_x f^x = h \cdot h', \tag{5}$$

where $h' = \sum_{i \in S} \sum_{x \in T_i} z_x f^x$ is a polynomial in $\{z_y : \text{type}(y) \in S\}$. It remains to show that for any face σ of type S , we have $h = g_{(X_\sigma, \pi_\sigma)}$, and for any τ of type $[d] \setminus S$, we have $h' = g_{(X_\tau, \pi_\tau)}$. Fix arbitrary faces σ of type S and τ of type $[d] \setminus S$. Noting that $g_{(X, \pi)}$ is a multiple of $h \cdot h'$, and that h' is in variables associated to elements whose types are in S and h is in variables associated to elements whose types are in $[d] \setminus S$, we conclude that h' has a monomial that is a multiple of $\prod_{x \in \sigma} z_x$ and h has a monomial that is a multiple of $\prod_{x \in \tau} z_x$. First, take $(\prod_{x \in \sigma} \partial_{z_x})$ from both

sides of (5). We get that $g_{(X_\sigma, \pi_\sigma)}$ is a positive multiple of h . Similarly, taking $(\prod_{x \in \tau} \partial_{z_x})$ from both sides of (5), we get that $g_{(X_\tau, \pi_\tau)}$ is a positive multiple of h' . Thus, noting that the coefficients of generating polynomials sum up to 1, we get $h = g_{(X_\sigma, \pi_\sigma)}$ and $h' = g_{(X_\tau, \pi_\tau)}$ as desired. Repeating the same argument inductively on the complex (X_σ, π_σ) proves the claim. \square

Definition 3.3. Given graph $G = (V, E)$, we say a family of vectors $\{f_S \in \mathbb{R}^V\}_{S \subset V: |S| \leq |V|-2}$ is G -consistent if for any $S \subset V$ of size at most $|V| - 2$, f_S is supported on $[d] \setminus S$ and, if G_S is disconnected, we have $f_S = \sum_{1 \leq i \leq \ell: |I_i| \geq 2} f_{V \setminus I_i}$, where $I_1 \cup \dots \cup I_\ell$ are the vertices of the connected components of G_S . Note that if $|I_i| = 1$ for all $1 \leq i \leq \ell$, then $f_S = 0$.

Proposition 3.4. Consider a totally connected $(d+1)$ -partite complex (X, π) with parts indexed by $[d]$ and graph $G := G_{(X, \pi)}$. Suppose $\{f_S \in \mathbb{R}^{[d]}\}_{S \subset [d], |S| < d}$ is a family of vectors that is G -consistent. Moreover, suppose that for all $k \geq 2$ and $S \subset [d]$ of size $(d+1) - k$, $\max_{i \in [d]} f_S(i) \leq \frac{(k-1)^2}{3k-1}$, and, if G_S is connected, the following holds:

1. Base Case: If $k = 2$, then for every face τ of type S , $\lambda_2(P_\tau) \leq f_S(i)$ for all $i \in [d] \setminus S$.
2. Recursive Condition: If $k \geq 3$, then

$$\sum_{j \in [d] \setminus (S \cup i)} f_{S \cup j}(i) \leq (k-2)f_S(i) - f_S^2(i),$$

for all $i \in [d] \setminus S$.

Then, for all $k \geq 2$ and τ of co-dimension k and type S , $\lambda_2(P_\tau) \leq \frac{\max_{i \in [d] \setminus S} f_S(i)}{k-1}$.

Proof. We apply **Theorem 2.6**. For every $S \subset [d]$ such that $|S| < d$ define a diagonal matrix $D_S \in \mathbb{R}^{X(0) \times X(0)}$ as $D_S(x, x) = f_S(\text{type}(x))$ for all $x \in X(0)$. We prove that the conditions of **Theorem 2.6** hold for $M_\tau := \frac{\Pi_\tau D_S}{k-1}$ for an arbitrary face $\tau \in X$ of co-dimension at least $k \geq 2$ and type S . Note that since $\max_{i \in [d]} f_S(i) \leq \frac{(k-1)^2}{3k-1}$, we get $D_\tau \preceq \frac{(k-1)^2}{3k-1} I$, and thus, $M_\tau \preceq \frac{k-1}{3k-1} \Pi_\tau$. To prove the rest of the conditions hold, first assume that $k = 2$. When G_S is disconnected the G -consistency condition implies $f_S = 0$, and therefore, $D_S = 0$. Thus, we get $\Pi_\tau P_\tau - \pi_{\tau,0} \pi_{\tau,0}^\top \preceq 0 = \Pi_\tau D_S = M_\tau$, as desired. When G_S is connected, $\lambda_2(P_\tau) \leq f_S(i)$ for all $i \in [d] \setminus S$ implies that $\lambda_2(P_\tau) \leq D_S(x, x)$ for all $x \in X_\tau(0)$. Therefore, $\Pi_\tau P_\tau - \pi_{\tau,0} \pi_{\tau,0}^\top \preceq \Pi_\tau D_S = M_\tau$. Now, assume $k \geq 3$. First assume that G_S is disconnected and $G[I_1], \dots, G[I_\ell]$ are its connected components for some partition $I_1 \cup \dots \cup I_\ell$ of $[d] \setminus S$. Fix any $\sigma \in X_\tau(k-1)$. By **Lemma 3.2**, $(X_\tau, \pi_\tau) = (X_{\tau \cup \sigma_{-1}}, \pi_{\tau \cup \sigma_{-1}}) \times \dots \times (X_{\tau \cup \sigma_{-\ell}}, \pi_{\tau \cup \sigma_{-\ell}})$ where for every $1 \leq j \leq \ell$, σ_{-j} is a subset of σ that has type $[d] \setminus (S \cup I_j)$. Therefore, we get $\Pr_{\eta \sim \pi_{\tau \cup \sigma_{-j}}}[x \in \eta] = \Pr_{\eta \sim \pi_\tau}[x \in \eta]$ for all $1 \leq j \leq \ell$ and $x \in X_{\tau \cup \sigma_{-j}}(0)$. Combining this with **Fact 2.4**, we get $k_j \cdot \pi_{\tau \cup \sigma_{-j},0}(x) = k \cdot \pi_{\tau,0}(x)$, where $k_j = |I_j|$ for all $1 \leq j \leq \ell$. Thus we can write

$$\begin{aligned} \sum_{1 \leq j \leq \ell: |I_j| \geq 2} \frac{(k_j - 1)k_j}{(k-1)k} M_{\tau \cup \sigma_{-j}} &\stackrel{\text{def of } M_{\tau \cup \sigma_{-j}}}{=} \sum_{1 \leq j \leq \ell: |I_j| \geq 2} \frac{(k_j - 1)k_j}{(k-1)k} \frac{\Pi_{\tau \cup \sigma_{-j}} D_{[d] \setminus I_j}}{k_j - 1} \\ &= \sum_{1 \leq j \leq \ell: |I_j| \geq 2} \frac{k_j}{k(k-1)} \frac{k}{k_j} \Pi_\tau D_{[d] \setminus I_j} \\ &= \frac{\Pi_\tau}{k-1} \sum_{1 \leq j \leq \ell: |I_j| \geq 2} D_{[d] \setminus I_j} = \frac{\Pi_\tau D_S}{k-1} = M_\tau, \end{aligned}$$

where in the second to last equality, we used the fact that $\sum_{1 \leq j \leq \ell: |I_j| \geq 2} f_{[d] \setminus I_j} = f_S$ due to the G -consistency condition, and thus $\sum_{1 \leq j \leq \ell: |I_j| \geq 2} D_{[d] \setminus I_j} = D_S$ by definition of D_S . Now, assume that

G_S is connected. It is enough to show that $\mathbb{E}_{x \sim \pi_{\tau,0}} M_{\tau \cup x} \preceq M_\tau - M_\tau \Pi_\tau^{-1} M_\tau$. This is equivalent to showing that for any $x \in X_\tau(0)$ and of type $\text{type}(x) = i$

$$\mathbb{E}_{x' \sim \pi_{\tau,0}} \left[\frac{(\Pi_\tau^{-1} \Pi_{\tau \cup x'} D_{S \cup \text{type}(x')})(x, x)}{k-2} \right] \leq \frac{D_S(x, x)}{k-1} - \frac{D_S^2(x, x)}{(k-2)(k-1)} \quad (6)$$

One can check that for any $x \in X_\tau(0)$

$$\begin{aligned} \mathbb{E}_{x' \sim \pi_{\tau,0}} \left[\frac{(\Pi_\tau^{-1} \Pi_{\tau \cup x'} D_{S \cup \text{type}(x')})(x, x)}{k-2} \right] &= \frac{\sum_{x' \in X_{\tau \cup x}(0)} \Pr_{\sigma \sim \pi_{\tau \cup x}} [x' \in \sigma] D_{S \cup \text{type}(x')}(x, x)}{(k-1)(k-2)} \\ &= \sum_{j \in [d] \setminus S} \frac{f_{\tau \cup j}(\text{type}(x))}{(k-1)(k-2)} \sum_{x' \in X_{\tau \cup x}(0): \text{type}(x')=j} \Pr_{\sigma \sim \pi_{\tau \cup x}} [x' \in \sigma] \\ &= \frac{\sum_{j \in [d] \setminus S} f_{\tau \cup j}(\text{type}(x))}{(k-1)(k-2)}, \end{aligned}$$

where in the last equality, we used **Fact 2.5**. Thus, substituting $D_S(x, x) = f_S(\text{type}(x))$ in the RHS of (6), it is enough to show that for any $i \in [d] \setminus S$

$$\frac{\sum_{j \in [d] \setminus S} f_{\tau \cup j}(i)}{(k-1)(k-2)} \leq \frac{f_S(i)}{k-1} - \frac{f_S^2(i)}{(k-1)(k-2)},$$

which holds by assumption. \square

We are ready to prove **Theorem 1.2**.

Proof of Theorem 1.2. We apply **Proposition 3.4**. Let $G := G_{(X, \pi)}$. Consider the following family of vectors $\{f_S \in \mathbb{R}^{[d]}\}_{S \subset [d]: |S| < d}$: for any $S \subset [d]$ such that $|S| < d$, let f_S be of the following form: for any $i \in S$, let $f_S(i, i) = 0$, and for any $i \in [d] \setminus S$ let

$$f_S(i) := \begin{cases} 0 & \text{if } \Delta_S(i) = 0, \\ \epsilon_{\{i,j\}} \cdot g_{i,j}(\Delta_S(j)) & \text{if } \Delta_S(i) = 1 \text{ and } i \sim_S j, \\ \sum_{j \sim_S i} \epsilon_{\{i,j\}} \cdot h_i(\Delta_S(i)) & \text{if } \Delta_S(i) \geq 2, \end{cases}$$

where functions $g_{i,j}, h_i : \{1, \dots, \Delta\} \rightarrow \mathbb{R}_{\geq 0}$ for every $i \in [d]$ and $j \sim i$, are defined later in a way that guarantees that $\{f_S\}_{S \subset [d]: |S| < d}$ satisfies the assumptions of **Proposition 3.4** (see (8), (10)). First, note that $\{f_S\}_{S \subset [d]: |S| < d}$ is G -consistent. This is because, for any $S, S' \subset [d]$ such that $|S|, |S'| < d$, if $i \notin S, S'$ and $\{j \in [d] : j \sim_S i\} = \{j \in [d] : j \sim_{S'} i\}$, then $f_S(i) = f_{S'}(i)$.

To verify the rest of the condition, take arbitrary $k \geq 2$ and $S \subset [d]$ of size $(d+1) - k$. Before verifying the condition $\max_{i \in [d]} f_S(i) \leq \frac{(k-1)^2}{3k-1}$, we verify the last set of conditions. For this purpose, assume G_S is connected. First, assume that $k = 2$. Let $[d] \setminus S = \{i, j\}$. By definition of $\epsilon_{\{i,j\}}$, for any τ of type S , $\lambda_2(P_\tau) \leq \epsilon_{\{i,j\}}$. Thus, if we define $g_{\ell,t}(1) = 1$ for all distinct $\ell, t \in [d]$, then we get $\lambda_2(P_\tau) \leq \epsilon_{\{i,j\}} = \epsilon_{\{i,j\}} g_{i,j}(1) = f_S(i) = f_S(j)$, as desired. Now, assume that $k \geq 3$. Fix an arbitrary $i \in [d] \setminus S$. Our goal is to define $f_{i,j}, h_i : \{1, \dots, \Delta\} \rightarrow \mathbb{R}_{\geq 0}$ for all $j \sim i$ such that $f_{i,j}(1) = 1$ for all $j \sim i$ and the following inequality is satisfied:

$$\sum_{j \in [d] \setminus (S \cup i)} f_{S \cup j}(i) \leq (k-2) f_S(i) - f_S^2(i)$$

To keep the notation concise, relabel the elements such that i is relabeled to 0 and $\epsilon_{\{0,1\}} \geq \dots \geq \epsilon_{\{0,d\}}$. Moreover, let $\epsilon_j := \epsilon_{\{0,j\}}$ for any $j \in [d] \setminus 0$.

Case 1: $\Delta_S(0) = 1$, and $j \sim_S 0$. Since G_S is connected and $(d+1) - |S| \geq 3$, we have $\Delta_S(j) \geq 2$. Let $t := \Delta_S(j)$. We have

$$\begin{aligned} \sum_{\ell \in [d] \setminus (S \cup 0)} f_{S \cup \ell}(0) &= f_{S \cup j}(0) + \sum_{\ell \in [d] \setminus (S \cup 0): \ell \sim_S j} f_{S \cup \ell}(0) + \sum_{\ell \in [d] \setminus (S \cup 0): \ell \not\sim_S j, \ell \neq j} f_{S \cup \ell}(0) \\ &= 0 + (t-1) \cdot \epsilon_j \cdot g_{0,j}(t-1) + (k-t-1) \cdot \epsilon_j \cdot g_{0,j}(t) \end{aligned}$$

On the other hand, $(k-2)f_S(0) - f_S(0)^2 = (k-2) \cdot \epsilon_j \cdot g_{0,j}(t) - \epsilon_j^2 \cdot g_{0,j}^2(t)$. So it is enough to satisfy

$$(t-1) \cdot \epsilon_j \cdot (g_{0,j}(t) - g_{0,j}(t-1)) \geq \epsilon_j^2 \cdot g_{0,j}^2(t). \quad (7)$$

Now, define $g_{0,j} : \{1, \dots, \Delta\} \rightarrow \mathbb{R}_{\geq 0}$ as follows: recall that we defined $g_{0,j}(1) = 1$. For any $2 \leq \ell \leq \Delta$, let

$$g_{0,j}(\ell) = 1 + 1.3 \cdot \epsilon_j \cdot H_{\ell-1}. \quad (8)$$

Using assumption (1), $\epsilon_j H_{\Delta-1} \leq \frac{\delta^2}{10} \leq \frac{1}{10}$. Thus

$$\epsilon_j^2 \cdot g_{0,j}^2(t) \leq \epsilon_j^2 (1 + 1.3\epsilon_j (1 + H_{\Delta-1}))^2 < 1.3\epsilon_j^2.$$

Substituting $g_{0,j}(t)$ according to (8) and using the above bound, one can verify that (7) holds.

Case 2: $\Delta_S(0) \geq 2$. For simplicity of notation, let $t := \Delta_S(0)$ and $\alpha := \sum_{j: j \sim_S 0} \epsilon_j$. Define $h_0(1) = \max_{j: j \sim 0} g_{0,j}(\Delta)$.

$$\begin{aligned} \sum_{j \in [d] \setminus (S \cup 0)} f_{S \cup j}(0) &= \sum_{j \in [d] \setminus (S \cup 0): j \sim_S 0} f_{S \cup j}(0) + \sum_{j \in [d] \setminus (S \cup 0): j \not\sim_S 0} f_{S \cup j}(0) \\ &\leq \left(\sum_{j \in [d] \setminus (S \cup 0): j \sim_S 0} (\alpha - \epsilon_{\{0,j\}}) \right) \cdot h_0(t-1) + (k-t-1) \cdot \alpha \cdot h_0(t) \\ &= (t-1) \cdot \alpha \cdot h_0(t-1) + (k-t-1) \cdot \alpha \cdot h_0(t). \end{aligned}$$

Note that if $t \geq 3$, the first inequality is an equality by definition. If $t = 2$, the first inequality follows from the definition of $h_0(1)$. Thus, it is enough to satisfy

$$\begin{aligned} \sum_{j \in [d] \setminus (S \cup 0)} f_{S \cup j}(0) &= (t-1) \cdot \alpha \cdot h_0(t-1) + (k-t-1) \cdot \alpha \cdot h_0(t) \\ &\leq (k-2) \cdot \alpha \cdot h_0(t) - \alpha^2 \cdot h_0^2(t) = (k-2)f_S(0) - f_S^2(0). \end{aligned}$$

Equivalently, it suffices to satisfy

$$(t-1)(h_0(t) - h_0(t-1)) \geq \alpha \cdot h_0^2(t). \quad (9)$$

Now, define $h_0 : \{1, \dots, \Delta\} \rightarrow \mathbb{R}_{\geq 0}$ as follows: recall that we defined $h_0(1) = \max_{j: j \sim 0} g_{0,j}(\Delta)$. For any $2 \leq \ell \leq \Delta$, let

$$h_0(\ell) := \frac{h_0(1)}{1 - c \left(\sum_{j=1}^{\ell} \epsilon_j H_{\ell-1}(j-1) \right)} \quad (10)$$

We need to prove (9) for a carefully chosen c . Let β be such that $h_0(t) = \frac{h_0(1)}{\beta}$. We get $h_0(t-1) = \frac{h_0(1)}{\beta + c(\sum_{j=1}^t \frac{\epsilon_j}{t-1})}$, and thus,

$$(t-1)(h_0(t) - h_0(t-1)) = \frac{h_0(1) \cdot c \sum_{j=1}^t \epsilon_j}{\beta \cdot (\beta + \frac{c \sum_{j=1}^t \epsilon_j}{t-1})}.$$

Note that $\alpha \cdot h_0^2(t) = \frac{\alpha \cdot h_0^2(1)}{\beta^2}$. Thus, to satisfy (9), it is enough to show that

$$\beta \cdot c \cdot \left(\sum_{j=1}^t \epsilon_j \right) \geq \alpha \cdot h_0(1) \cdot \left(\beta + \frac{c \sum_{j=1}^t \epsilon_j}{t-1} \right).$$

Note that

$$h_0(1) \leq \max_{j \sim i} g_{0,j}(\Delta) = 1 + 1.3\epsilon_1 H_{\Delta-1} \stackrel{\text{by (1)}}{\leq} 1 + 1.3 \frac{\delta^2}{10} \quad (11)$$

Moreover, $\sum_{j=1}^t \epsilon_j \geq \sum_{j:j \sim s_0} \epsilon_j = \alpha$. Thus, letting $c = 1 + c'\delta$ for some $c' > 0$ that we choose later, it is enough to show that

$$\beta \cdot (c' - 0.13\delta)\delta \geq (1 + 0.13\delta) \cdot \frac{(1 + c'\delta) \sum_{j=1}^t \epsilon_j}{t-1}$$

Using $\frac{\sum_{j=1}^t \epsilon_j}{t-1} \leq 2\epsilon_1 \stackrel{(1)}{\leq} \frac{\delta^2}{5}$, it is enough to show that

$$\beta \cdot (c' - 0.13\delta) \geq (1 + 0.13\delta)(1 + c'\delta) \frac{\delta}{5}. \quad (12)$$

On the other hand,

$$\beta \geq 1 - (1 + c'\delta) \left(\sum_{j=1}^{\Delta(0)} \epsilon_j H_{\Delta(0)-1}(j-1) \right) \stackrel{(2)}{\geq} 1 - (1 + c'\delta)(1 - \delta) = \delta(1 - c' + c'\delta), \quad (13)$$

Thus, to satisfy (12), it is enough to show that $(1 - c' + c'\delta)(c' - 0.13\delta) \geq (1 + 1.13\delta)(1 + c'\delta) \frac{1}{5}$. Letting $c' = \frac{1}{2}$, this inequality holds for every $0 < \delta < 1$.

Now, we verify the condition that $\max_{i \in [d]} f_S(i) \leq \frac{(k-1)^2}{3k-1}$ for an arbitrary $S \subset [d]$ such that $|S| = (d+1) - k$.

Note that $\sum_{j:j \sim i} \epsilon_{\{i,j\}} \leq \Delta_S \cdot \epsilon_1 \stackrel{(1)}{\leq} \Delta_S \cdot \frac{\delta^2}{10}$ for all $i \in [d] \setminus S$. Thus, we get $\max_{i \in [d]} f_S(i) \leq \Delta_S \cdot \frac{\delta^2}{10} \max_{i \in [d] \setminus S} h_i(\Delta_S(i))$. Moreover, using (11) and (13) with $c' = \frac{1}{2}$ (we can write this inequality for every i), we get

$$h_i(\Delta_S(i)) \leq h_i(\Delta(i)) \leq \frac{1 + \frac{\delta^2}{10}}{\delta(\frac{1}{2} + \frac{\delta}{2})}, \quad (14)$$

Thus, we can write

$$\max_{i \in [d]} f_S(i) \leq \Delta_S \cdot \frac{\delta^2}{10} \frac{1 + \frac{\delta^2}{10}}{\delta(\frac{1}{2} + \frac{\delta}{2})} \leq \frac{\Delta_S}{5} \leq \frac{k-1}{5} \leq \frac{(k-1)^2}{3k-1},$$

as desired. So we proved that $\{f_S\}_{S \subset [d]: |S| < d}$ satisfies the conditions of [Proposition 3.4](#). Now, we are ready to bound $\lambda_2(P_\tau)$ for any face τ of co-dimension $k \geq 2$ and type S . First, we show that for every $i \in [d] \setminus S$, $\sum_{j: j \sim_{Si}} \epsilon_{\{i,j\}} \leq 1 - \delta$. Note that

$$\sum_{\ell=1}^{\Delta(i)} H_{\Delta(i)-1}(\ell-1) = \sum_{\ell=2}^{\Delta(i)} \frac{\ell}{\ell-1} = 2 + \sum_{\ell=3}^{\Delta(i)} \frac{\ell}{\ell-1} \geq \Delta(i).$$

Thus, we can write

$$\sum_{j: j \sim_{Si}} \epsilon_{\{i,j\}} \leq \left(\sum_{\ell=1}^{\Delta(i)} \frac{H_{\Delta(i)-1}(\ell-1)}{\Delta(i)} \right) \left(\sum_{j \sim i} \epsilon_{\{i,j\}} \right) \leq \sum_{\ell=1}^{\Delta(i)} H_{\Delta(i)-1}(\ell-1) \cdot \epsilon_{\{i,j_\ell\}} \leq 1 - \delta. \quad (15)$$

where we assumed that i_1, \dots, j_d is an ordering of $[d] \setminus S$ such that $\epsilon_{j_1} \leq \dots \leq \epsilon_{j_d}$. Using this inequality and [\(14\)](#), we get

$$\begin{aligned} \lambda_2(P_\tau) &\leq \frac{\max_{i \in [d] \setminus S} f_S(i)}{k-1} \leq \frac{\max_{i \in [d]} (\sum_{j \sim_{Si}} \epsilon_{\{i,j\}}) \cdot h_i(\Delta_S(i))}{k-1} \\ &\leq \frac{(1-\delta) \cdot \max_{i \in [d]} h_i(\Delta(i))}{k-1} \leq \frac{(1-\delta) \cdot \frac{2(1+\frac{\delta^2}{10}\delta)}{\delta(\delta+1)}}{k-1}, \end{aligned}$$

as desired. □

References

- [AJT19] Vedat Levi Alev, Fernando Granha Jeronimo, and Madhur Tulsiani. “Approximating Constraint Satisfaction Problems on High-Dimensional Expanders”. In: *FOCS*. Ed. by David Zuckerman. IEEE Computer Society, 2019, pp. 180–201 (cit. on p. 2).
- [ALO20] Nima Anari, Kuikui Liu, and Shayan Oveis Gharan. “Spectral Independence in High-Dimensional Expanders and Applications to the Hardcore Model”. In: *SIAM Journal on Computing* 0.0 (2020), FOCS20-1-FOCS20–37. DOI: [10.1137/20M1367696](https://doi.org/10.1137/20M1367696). eprint: <https://doi.org/10.1137/20M1367696> (cit. on p. 2).
- [ALO21] Dorna Abdolazimi, Kuikui Liu, and Shayan Oveis Gharan. “A Matrix Trickle-Down Theorem on Simplicial Complexes and Applications to Sampling Colorings”. In: *FOCS*. IEEE, 2021, pp. 161–172 (cit. on pp. 2–4, 8).
- [Ana+19] Nima Anari, Kuikui Liu, Shayan Oveis Gharan, and Cynthia Vinzant. “Log-concave polynomials II: high-dimensional walks and an FPRAS for counting bases of a matroid”. In: *STOC*. Ed. by Moses Charikar and Edith Cohen. ACM, 2019, pp. 1–12 (cit. on p. 2).
- [Car89] Roger Carter. *Simple groups of Lie type*. John Wiley & Sons, 1989 (cit. on p. 4).

- [CLV21] Zongchen Chen, Kuikui Liu, and Eric Vigoda. “Optimal mixing of Glauber dynamics: entropy factorization via high-dimensional expansion”. In: *STOC*. Ed. by Samir Khuller and Virginia Vassilevska Williams. ACM, 2021, pp. 1537–1550 (cit. on p. 2).
- [DD19] Yotam Dikstein and Irit Dinur. “Agreement Testing Theorems on Layered Set Systems”. In: *FOCS*. Ed. by David Zuckerman. IEEE Computer Society, 2019, pp. 1495–1524 (cit. on p. 2).
- [Din+21a] Irit Dinur, Yuval Filmus, Prahladh Harsha, and Madhur Tulsiani. “Explicit SoS Lower Bounds from High-Dimensional Expanders”. In: *12th Innovations in Theoretical Computer Science Conference, ITCS 2021, January 6-8, 2021, Virtual Conference*. Ed. by James R. Lee. Vol. 185. LIPIcs. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021, 38:1–38:16 (cit. on p. 2).
- [Din+21b] Irit Dinur, Prahladh Harsha, Tali Kaufman, Inbal Livni Navon, and Amnon Ta-Shma. “List-Decoding with Double Samplers”. In: *SIAM J. Comput.* 50.2 (2021), pp. 301–349 (cit. on p. 2).
- [Din+22] Irit Dinur, Shai Evra, Ron Livne, Alexander Lubotzky, and Shahar Mozes. “Locally testable codes with constant rate, distance, and locality”. In: *STOC*. Ed. by Stefano Leonardi and Anupam Gupta. ACM, 2022, pp. 357–374 (cit. on p. 2).
- [KO18] Tali Kaufman and Izhar Oppenheim. “Construction of new local spectral high dimensional expanders”. In: *STOC*. Ed. by Ilias Diakonikolas, David Kempe, and Monika Henzinger. ACM, 2018, pp. 773–786. DOI: [10.1145/3188745.3188782](https://doi.org/10.1145/3188745.3188782) (cit. on pp. 2, 3).
- [LSV05] Alexander Lubotzky, Beth Samuels, and Uzi Vishne. “Explicit constructions of Ramanujan complexes of type \tilde{A}_d ”. In: *European Journal of Combinatorics* 26.6 (2005). Combinatorics and Representation Theory, pp. 965–993. ISSN: 0195-6698. DOI: <https://doi.org/10.1016/j.ejc.2005.07.001> (cit. on p. 2).
- [OP22] Ryan O’Donnell and Kevin Pratt. “High-Dimensional Expanders from Chevalley Groups”. In: *37th Computational Complexity Conference (CCC 2022)*. Ed. by Shachar Lovett. Vol. 234. Leibniz International Proceedings in Informatics (LIPIcs). Dagstuhl, Germany: Schloss Dagstuhl – Leibniz-Zentrum für Informatik, 2022, 18:1–18:26. ISBN: 978-3-95977-241-9 (cit. on pp. 2–4).
- [Opp18] Izhar Oppenheim. “Local spectral expansion approach to high dimensional expanders part I: Descent of spectral gaps”. In: *Discrete and Computational Geometry* 59.2 (2018), pp. 293–330 (cit. on pp. 2, 7).