

Dynamics of fractional N -soliton solutions with anomalous dispersions of integrable fractional higher-order nonlinear Schrödinger equations

Weifang Weng^{1,2}, Minghe Zhang^{1,2}, and Zhenya Yan^{1,2,*}

¹Key Lab of Mathematics Mechanization, Academy of Mathematics and Systems Science,
Chinese Academy of Sciences, Beijing 100190, China

²School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

In this paper, we explore the anomalous dispersive relations, inverse scattering transform and fractional N -soliton solutions of the integrable fractional higher-order nonlinear Schrödinger (fHONLS) equations, containing the fractional Hirota (fHirota), fractional complex mKdV (fcmKdV), and fractional Lakshmanan-Porsezian-Daniel (fLPD) equations, etc. The inverse scattering problem can be solved exactly by means of the matrix Riemann-Hilbert problem with simple poles. As a consequence, an explicit formula is found for the fractional N -soliton solutions of the fHONLS equations in the reflectionless case. In particular, we analyze the fractional one-, two- and three-soliton solutions with anomalous dispersions of fHirota and fcmKdV equations. The wave, group, and phase velocities of these envelope fractional 1-soliton solutions are related to the power laws of their amplitudes. These obtained fractional N -soliton solutions may be useful to explain the related super-dispersion transports of nonlinear waves in fractional nonlinear media.

Fractional nonlinear equations and integrable (integer-order) nonlinear equations are two kinds of important physical models in the fields of nonlinear dynamics and applications. The formers are used to describe physical phenomena with anomalous diffusion, and in general non-integrable such that they can be solved approximately by numerical methods, however, the latters are a class of important physical models and can be solved exactly by the inverse scattering transform (IST) to generate exact solitons, which can be used to compared with numerical and experimental results. More recently, based on two significant aspects, i.e., Riesz fractional derivative and IST integrability, Ablowitz *et al* presented the new types of integrable fractional nonlinear soliton equations such as the fractional KdV, fractional NLS, fractional mKdV, fractional sine-Gordon, and fractional sinh-Gordon equations. Moreover, their fractional one-soliton solutions were found. These solitons show the anomalous dispersions. In this paper, motivated by the idea, we will investigate the integrable fractional extensions of higher-order NLS (fHONLS) equations, containing the fractional Hirota (fHirota), fractional complex mKdV (fcmKdV), fractional LPD (fLPD) equations, and etc. We give the anomalous dispersive relations, and explicit forms of these fHONLS equations via the completeness of eigenfunctions. Based on the IST with matrix RH problems, we find a formula of fractional N -soliton solutions. In particular, we analyze the fractional one-, two- and three-soliton solutions with anomalous dispersions of fHirota and fcmKdV equations. The wave, group, and phase velocities of these fractional solitons are related to the power laws of their amplitudes. These obtained fractional N -soliton solutions may be useful to explain the super-dispersion transports of nonlinear waves in fractional nonlinear media.

1 Introduction

Since the inverse scattering transform (IST), as a nonlinear extension of Fourier transform, was presented by GGKM [1] in 1967, and then the Lax pairs were coined by Lax in 1968 [2], many types of nonlinear wave equations have been shown to be IST integrable (i.e., they can be solved exactly by the IST), containing the integrable nonlinear partial differential, differential-difference, difference, and differential-integrable equations (see, e.g., Refs. [3–6] and references therein), where the partial derivatives are usually integer-order derivatives. In fact,

*Email address: zyyan@mmrc.iss.ac.cn

since fractional calculus (FC), as an extension of integer-order one, was coined in the L'Hopital's letter written to Leibniz in 1659, more and more attention has been paid to FC and its application in many physical systems with anomalous diffusion such as quantum mechanics, nanofluids, geotechnical engineering, viscoelastic material, and polymer Science (see, e.g., Refs. [7–12] and reference therein). Up to now, there are many types of fractional linear equations such as the fractional Schrödinger equation [13–15], and fractional nonlinear wave (fNLW) equations such as the fractional nonlinear Schrödinger (NLS) equation [16–18]

$$i\psi_t + (-\nabla^2)^\epsilon \psi + V(\mathbf{r}, t)\psi + f(|\psi|^2)\psi = 0, \quad \mathbf{r} \in \mathbb{R}^d, \quad \epsilon \in (0, 1), \quad (1)$$

where ∇^2 denotes the d -dimensional Laplace operator, and the Riesz fractional derivative is defined by [19–21]

$$(-\nabla^2)^\epsilon \psi(\mathbf{r}, t) = \frac{1}{2\pi} \int_{\mathbb{R}^d} d\mathbf{k} |\mathbf{k}|^{2\epsilon} \int_{\mathbb{R}^d} e^{i\mathbf{k}(\mathbf{r}-\mathbf{r}')}\psi(\mathbf{r}')d\mathbf{r}',$$

and the fractional complex Ginzburg–Landau equation, etc [18]. However, these fNLW equations were not integrable in the sense of IST such that their exact solutions can usually not found, and their approximate solutions were given with the aid of numerical methods.

More recently, Ablowitz *et al* [22, 23] extended the well-established Riesz fractional derivative [19–21] (e.g., $|\partial^2|^\epsilon$, $0 < \epsilon < 1$) to present several new types of IST integrable fractional nonlinear evolution equations (fNLEEs) such as the fractional NLS (fNLS), fractional KdV (fKdV), fractional mKdV (fmKdV), fractional sine-Gordon (fsG), and fractional sinh-Gordon (fshG) equations, and found that they were integrable by the IST with GLM-type integral equations to admit the fractional one-soliton solutions.

When the ultra-short (e.g., 100 fs [24, 25]) optical pulse propagation is considered, the higher-order dispersive effects (e.g., third-order dispersion) and nonlinear effects (e.g., self-frequency shift and self-steepening) due to the stimulated Raman scattering can not be neglected [26–28]. As a result, a generation of the NLS equation called the Hirota equation [29]

$$iq_t + \alpha(q_{xx} + 2v|q|^2q) + i\beta(q_{xxx} + 6v|q|^2q_x) = 0, \quad \alpha, \beta \in \mathbb{R}, \quad v = \pm 1, \quad (x, t) \in \mathbb{R}^2 \quad (2)$$

was presented, which is also an important physical model, and IST integrable, where α , β stand for the second- and third-order dispersive coefficients, respectively, and $v = +1(-1)$ corresponds to the self-focusing (defocusing) interaction. The rogue wave solutions of the focusing Hirota equation were found using the Darboux transform [30–33]. At $\beta = 0$ and $\alpha = 1$, Eq. (2) becomes the NLS equation, while as $\alpha = 0$, $\beta = 1$, Eq. (2) is the complex modified KdV (cmKdV) equation

$$q_t + q_{xxx} + 6v|q|^2q_x = 0, \quad v = \pm 1, \quad (x, t) \in \mathbb{R}^2, \quad (3)$$

In fact, there exist higher-order NLS equation such as the fourth-order Lakshmanan-Porsezian-Daniel (LPD) [34], and fifth-order NLS equation [35, 36], and higher-order NLS equations [36].

In this paper, we would like to apply the idea (combination of Riesa fractional derivative and IST) [22] to study the integrable fractional Hirota (fHirota) and fractional cmKdV (fcmKdV) equations, and solve them by the IST with the matrix Riemann-Hilbert problems (not the GLM integral equations) such that we find their fractional N -soliton solutions. The rest of this paper is arranged as follows. In Sec. II, we give the integrable fHirota and fcmKdV equations, and their explicit forms in terms of the completeness relation of squared eigenfunctions. Similarly, the general higher-order fractional NLS equations are also presented, such as the fractional LPD equation and fractional fifth-order NLS equations. The anomalous dispersive relations are presented. Moreover, the IST with the matrix Riemann-Hilbert problem for the simple pole case is used to study their fractional solutions. The trace formulae are also studied. In Sec. 3, we give the formula of fractional N -soliton solutions for the reflectionless case. Some representative fractional one-, two- and three-soliton solutions are explored for the fHirota and fcmKdV equations. The wave, group, and phase velocities of these envelope fractional solitons are shown to be related to the power laws of their amplitudes. These obtained

fractional N -soliton solutions may be useful to explain the related super-dispersion transport of nonlinear wave phenomena in fractional nonlinear media. Moreover, we also present the corresponding formula for the fractional N -soliton solutions of the integrable fractional HONLS equations. Finally, some conclusions and discussions are given in Sec. 4.

2 Fractional Hirota equation and extensions: IST with RH problem

The fractional Hirota equation.—The 2×2 -matrix ZS-AKNS scattering problem [37,38] is given as

$$\Phi_x = X\Phi, \quad X(x, t; k) = -ik\sigma_3 + Q, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad Q = \begin{bmatrix} 0 & q(x, t) \\ r(x, t) & 0 \end{bmatrix}, \quad (4)$$

where $\Phi = \Phi(x, t; k)$ is a 2×2 matrix-valued eigenfunction, $k \in \mathbb{C}$ is a spectral parameter, and $r(x, t)$, $q(x, t)$ stand for the potentials. Starting from the spectral problem (4), one can find the integer-order integrable AKNS hierarchy [37,38]. Similarly, we here present the coupled fractional Hirota equations

$$\mathbf{q}_t + \sigma_3 \mathcal{N}(\widehat{\mathbf{L}})\mathbf{q} = 0, \quad \mathbf{q} = (r, q)^T, \quad (5)$$

where $r = r(x, t)$, $q = q(x, t)$, $\mathcal{N}(\widehat{\mathbf{L}}) = -4i(\alpha + 2\beta\widehat{\mathbf{L}})\widehat{\mathbf{L}}^2|4\widehat{\mathbf{L}}^2|^\epsilon$ with $\epsilon \in (0, 1)$, $\alpha, \beta \in \mathbb{R}$ and

$$\sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad \widehat{\mathbf{L}} = \frac{1}{2i} \begin{bmatrix} \partial - 2r\partial_{-1}^{-1}q & 2r\partial_{-1}^{-1}r \\ -2q\partial_{-1}^{-1}q & -\partial + 2q\partial_{-1}^{-1}r \end{bmatrix} \quad (6)$$

with $\partial = \partial/\partial x$, $\partial_{-1}^{-1} = \int_{-\infty}^x dy$. In particular, as $r = -\nu q^*$, $\nu = \pm 1$, where the star denotes the complex conjugate, we have the fHirota equation

$$i \begin{bmatrix} -\nu q^* \\ q \end{bmatrix}_t - \sigma_3 |4\widehat{\mathbf{L}}^2|^\epsilon \begin{bmatrix} -\alpha(\nu q_{xx}^* + 2|q|^2 q^*) + i\beta(\nu q_{xxx}^* + 6|q|^2 q_x^*) \\ \alpha(q_{xx} + 2\nu|q|^2 q) + i\beta(q_{xxx} + 6\nu|q|^2 q_x) \end{bmatrix} = 0. \quad (7)$$

In particular, when $\beta = 0$, one has the fractional NLS (fNLS) equation [22]

$$i \begin{bmatrix} \mp q^* \\ q \end{bmatrix}_t - \sigma_3 |4\widehat{\mathbf{L}}^2|^\epsilon \begin{bmatrix} -\alpha(\nu q_{xx}^* + 2|q|^2 q^*) \\ \alpha(q_{xx} + 2\nu|q|^2 q) \end{bmatrix} = 0. \quad (8)$$

As $\alpha = 0$, $\beta = 1$, one has the fractional complex mKdV (fcmKdV) equation

$$\begin{bmatrix} -\nu q^* \\ q \end{bmatrix}_t - \sigma_3 |4\widehat{\mathbf{L}}^2|^\epsilon \begin{bmatrix} -\nu q_{xxx}^* - 6|q|^2 q_x^* \\ q_{xxx} + 6\nu|q|^2 q_x \end{bmatrix} = 0. \quad (9)$$

Anomalous dispersive relation.—The formal solution $q(x, t) \sim e^{i[kx - w(k)t]}$ is employed into the linearization of Eq. (5) yields the dispersive relation of the linear fHirota equation

$$w(k) = i\mathcal{N}(-k/2). \quad (10)$$

We further consider the linearization of the fHirota Eq. (7)

$$iq_t + |-\partial^2|^\epsilon (\alpha q_{xx} + i\beta q_{xxx}) = 0,$$

where $|-\partial^2|^\epsilon$ denotes the Riesz fractional derivative, such that the anomalous dispersive relation is

$$w(k) = (\alpha - \beta k)k^2|k^2|^\epsilon, \quad (11)$$

in which we have

$$\mathcal{N}(k) = -iw(-2k) = -4i(\alpha + 2\beta k)k^2|4k^2|^\epsilon. \quad (12)$$

Fractional higher-order NLS equations and anomalous dispersive relations.—In fact, one can also extend the fHirota equation to other fractional higher-order NLS (fHONLS) equations in the form

$$\mathbf{q}_t + \sigma_3 \mathcal{N}_h(\widehat{\mathbf{L}}) \mathbf{q} = 0, \quad \mathcal{N}_h(\widehat{\mathbf{L}}) = \left(\sum_{j=2}^N \alpha_j \delta_j \widehat{\mathbf{L}}^j \right) |4\widehat{\mathbf{L}}^2|^\epsilon, \quad \mathbf{q} = (r, q)^T, \quad (13)$$

where $\alpha_j \in \mathbb{R}$ and

$$\delta_j = \begin{cases} i(2i)^j, & j = 2n, \quad n \in \mathbb{N}, \\ (2i)^j, & j = 2n + 1, \quad n \in \mathbb{N}. \end{cases}$$

Let $r = -\nu q^*$, $\nu = \pm 1$, then it follows from Eq. (13) that we have the fHONLS equation

$$i \begin{bmatrix} -\nu q^* \\ q \end{bmatrix}_t + i\sigma_3 |4\widehat{\mathbf{L}}^2|^\epsilon \left(\sum_{j=2}^N \alpha_j \delta_j \widehat{\mathbf{L}}^j \right) \begin{bmatrix} -\nu q^* \\ q \end{bmatrix} = 0. \quad (14)$$

In particular, as $N = 5$, we have the fractional fifth-order NLS equation

$$i \begin{bmatrix} \mp q^* \\ q \end{bmatrix}_t - \sigma_3 |4\widehat{\mathbf{L}}^2|^\epsilon \begin{bmatrix} \alpha_2(\mp q_{xx}^* - 2|q|^2 q^*) - i\alpha_3(\mp q_{xxx}^* - 6|q|^2 q_x^*) \mp \alpha_4 S_4^*[q] \pm i\alpha_5 S_5^*[q] \\ \alpha_2(q_{xx} \pm 2|q|^2 q) + i\alpha_3(q_{xxx} \pm 6|q|^2 q_x) + \alpha_4 S_4[q] + i\alpha_5 S_5[q] \end{bmatrix} = 0 \quad (15)$$

with

$$\begin{aligned} S_4[q] &= q_{xxxx} \pm 8|q|^2 q_{xx} + 6|q|^4 q \pm 4|q_x|^2 q \pm 6q^* q_x^2 \pm 2q^2 q_{xx}^*, \\ S_5[q] &= q_{5x} \pm 10|q|^2 q_{xxx} \pm 10(q|q_x|^2)_x \pm 10q^*(q_x^2)_x + 30|q|^4 q_x. \end{aligned}$$

which reduce to the fractional Lakshmanan-Porsezian-Daniel (fLPD) equation as $S_5 = 0$.

We further use the formal solution $q(x, t) \sim e^{i[kx - w_h(k)t]}$ to study the linearization of the fHONLS equation (14)

$$iq_t + |-\partial^2|^\epsilon \left(\sum_{j=2}^N \alpha_j i^{\zeta_j} q_{jx} \right) = 0, \quad \zeta = 0, \quad j = 2n; \quad \zeta = 1, \quad j = 2n + 1, \quad n \in \mathbb{N},$$

where $q_{jx} = \partial^j q / \partial x^j$, such that the dispersive relation is

$$w_h(k) = - \sum_{j=2}^N \alpha_j i^{\zeta_j + j} k^j |k^2|^\epsilon, \quad (16)$$

in which we have

$$\mathcal{N}_h(k) = -iw_h(-2k) = \sum_{j=2}^N \alpha_j i^{\zeta_j + j + 1} (-2k)^j |4k^2|^\epsilon. \quad (17)$$

In the following we mainly consider the fHirota and fcmKdV equations (7). In fact, one can also consider the fractional higher-order NLS equations (14).

The direct scattering with sufficient decay and smoothness of \mathbf{q} .—For the given ZS-AKNS spectral problem (4), following the idea [22], we now consider the time evolution of the matrix eigenfunction $\Phi(x, t; k)$ in the form

$$\Phi_t = T\Phi, \quad T(x, t; k) = \begin{bmatrix} T_1(x, t; k) & T_2(x, t; k) \\ T_3(x, t; k) & -T_1(x, t; k) \end{bmatrix}, \quad (18)$$

where $T_{1,2,3}(x, t; k)$ can not be explicitly presented in general for the fractional Hirota equation. But we here require the conditions $T_{2,3}(x, t; k) \rightarrow 0$ and $T_1(x, t; k) \rightarrow \frac{1}{2}\mathcal{N}(k)$ with $\mathcal{N}(k)$ given by Eq. (12) as $x \rightarrow \pm\infty$ (i.e.,

$q(x, t) \rightarrow 0$). Therefore, for the zero-boundary condition $q(x, t) \in L^1(\mathbb{R}^\pm)$ and $r = -q^*$, we consider the asymptotic problem ($x \rightarrow \pm\infty$) of the spectral problem (4) and time part (18)

$$\begin{aligned}\Phi_x &= X_\pm(k)\Phi, & X_\pm(k) &= \lim_{x \rightarrow \pm\infty} X(x, t; k) = -ik\sigma_3, \\ \Phi_t &= T_\pm(k)\Phi, & T_\pm(k) &= \lim_{x \rightarrow \pm\infty} T(x, t; k) = \frac{1}{2}\mathcal{N}(k)\sigma_3,\end{aligned}$$

which can deduce the fundamental solutions $\Phi_{bg}(x, t; k) = e^{[-ikx + \frac{1}{2}\mathcal{N}(k)t]\sigma_3}$. One may introduce the Jost solutions $\Phi_\pm(x, t; k)$ satisfying the following boundary conditions

$$\Phi_\pm(x, t; k) \sim e^{[-ikx + \frac{1}{2}\mathcal{N}(k)t]\sigma_3}, \quad x \rightarrow \pm\infty. \quad (19)$$

such that the modified Jost solutions $\phi_\pm(x, t; k)$ are defined as

$$\phi_\pm(x, t; k) = \Phi_\pm(x, t; k)e^{(ikx - \frac{1}{2}\mathcal{N}(k)t)\sigma_3} \rightarrow \mathbb{I}, \quad x \rightarrow \pm\infty \quad (20)$$

with

$$\phi_\pm(x, t; k) = \mathbb{I} + \int_{\pm\infty}^x dx' e^{-ik(x-x')\hat{\sigma}_3} Q(x', t) \phi_\pm(x', t; k), \quad e^{\hat{\sigma}_3} A = e^{\sigma_3} A e^{-\sigma_3}. \quad (21)$$

Let $D_+ = \{k | \text{Im } k > 0\}$, $D_- = \{k | \text{Im } k < 0\}$, $\Phi_\pm(x, t; k) = (\Phi_{\pm 1}, \Phi_{\pm 2})$ and $\phi_\pm(x, t; k) = (\phi_{\pm 1}, \phi_{\pm 2})$. Then one has the following conclusion (similarly to the NLS equation [37, 39] and Hirota equation [40]): For the given $q(x, t) \in L^1(\mathbb{R}^\pm)$, the vector-valued functions $\Phi_{\pm 2}$ and $\phi_{\pm 2}$ given by Eqs. (20) and (21) both have unique solutions in \mathbb{R} . Moreover, $\phi_{-1, +2}$, $\Phi_{-1, +2}(\phi_{-2, +1}, \Phi_{-2, +1})$ can be extended analytically to D_+ (D_-), and continuously to $D_+ \cup \mathbb{R}$ ($D_- \cup \mathbb{R}$). Because $\Phi_\pm(x, t; k)$ are both fundamental solutions of the spectral problem, thus based on the above properties, there exists a constant scattering matrix $S(k) = (s_{ij}(k))_{2 \times 2}$ between them obeying the relation

$$\Phi_+(x, t; k) = \Phi_-(x, t; k)S(k), \quad k \in \mathbb{R}, \quad (22)$$

which can generate $s_{11}s_{22} - s_{12}s_{21} = 1$ and

$$\begin{aligned}s_{11}(k) &= |\Phi_{+1}(x, t; k), \Phi_{-2}(x, t; k)|, & s_{22}(k) &= |\Phi_{-1}(x, t; k), \Phi_{+2}(x, t; k)|, \\ s_{12}(k) &= |\Phi_{+2}(x, t; k), \Phi_{-2}(x, t; k)|, & s_{21}(k) &= |\Phi_{-1}(x, t; k), \Phi_{+1}(x, t; k)|.\end{aligned} \quad (23)$$

According to the properties of $\Phi_{\pm j}(x, t; k)$, $j = 1, 2$, it can be seen that the scattering coefficient $s_{11}(k)$ ($s_{22}(k)$) in $k \in \mathbb{R}$ can be extended analytically to D_- (D_+), and continuously to $D_- \cup \mathbb{R}$ ($D_+ \cup \mathbb{R}$), whereas another two scattering coefficients $s_{12}(k)$ and $s_{21}(k)$ can not be analytically continued away from \mathbb{R} . To study the matrix RH problem of the inverse scattering, the potential is required not to admit spectral singularities [41], i.e. $s_{11}(k)s_{22}(k) \neq 0$ as $k \in \mathbb{R}$ such that the reflection coefficients are introduced as $\rho(k) = s_{21}(k)s_{11}^{-1}(k)$, $\hat{\rho}(z) = s_{12}(k)s_{22}^{-1}(k)$ as $k \in \mathbb{R}$.

Completeness of squared eigenfunctions and explicit fHirota equation and higher-order extensions.—Since $\hat{\mathbf{L}}$ is a adjoint of the matrix operator \mathbf{L}

$$\mathbf{L} = \frac{1}{2i} \begin{bmatrix} -\partial - 2q\partial_+^{-1}r & -2q\partial_+^{-1}q \\ 2r\partial_+^{-1}r & \partial + 2r\partial_+^{-1}q \end{bmatrix}, \quad \partial_+^{-1} = \int_x^\infty dy. \quad (24)$$

It follows from the ZS-AKNS spectral problem that the eigenfunctions Ψ_{+j} , $\hat{\Psi}_{-j}$ of \mathbf{L} and $\hat{\mathbf{L}}$ and eigenvalue k satisfy [38, 42]

$$\begin{aligned}\mathbf{L}\Psi_{+j} &= k\Psi_{+j}, & \hat{\mathbf{L}}\hat{\Psi}_{-j} &= k\hat{\Psi}_{-j}, \quad j = 1, 2, \\ \Psi_{+1} &= (\Phi_{+11}^2(x, t; k), \Phi_{+12}^2(x, t; k))^T, & \Psi_{+2} &= (\Phi_{+21}^2(x, t; k), \Phi_{+22}^2(x, t; k))^T, \\ \hat{\Psi}_{-1} &= (\Phi_{-12}^2(x, t; k), -\Phi_{-11}^2(x, t; k))^T, & \hat{\Psi}_{-2} &= (\Phi_{-22}^2(x, t; k), -\Phi_{-21}^2(x, t; k))^T,\end{aligned} \quad (25)$$

and they are complete.

Similarly,

$$\mathcal{N}(\widehat{\mathbf{L}})\widehat{\Psi}_{-j} = \mathcal{N}(k)\widehat{\Psi}_{-j}, \quad j = 1, 2. \quad (26)$$

Since the eigenfunctions $\widehat{\Psi}_{-j}$ are complete, therefore, one can use $\mathcal{N}(\widehat{\mathbf{L}})$ to act on a sufficiently smooth and decaying vector function $\mathbf{g}(x) = (g_1(x), g_2(x))^T$ to get

$$\mathcal{N}(\widehat{\mathbf{L}})\mathbf{g}(x) = \frac{1}{\pi} \sum_{\ell=1}^2 \int_{\Gamma_{\infty, \ell}} dk \mathcal{N}(k) f_{\ell}(k) \int_{\mathbb{R}} G_{\ell}(x, y, k) \mathbf{g}(y) dy, \quad (27)$$

where

$$\begin{aligned} G_1(x, y, k) &= \widehat{\Psi}_{-1}(x, k) \Psi_{+2}^T(y, k), \quad f_1(k) = -s_{22}^{-2}(k), \quad k \in D_+, \\ G_2(x, y, k) &= \widehat{\Psi}_{-2}(x, k) \Psi_{+1}^T(y, k), \quad f_2(k) = s_{11}^{-2}(k), \quad k \in D_-, \end{aligned}$$

and $\Gamma_{\infty, \ell} = \lim_{\mathbb{R} \rightarrow \infty} \Gamma_{\mathbb{R}, \ell}$, $\ell = 1, 2$ with $\Gamma_{\mathbb{R}, 1}$ ($\Gamma_{\mathbb{R}, 2}$) the semicircular contour in the upper (lower) half plane evaluated $-\mathbb{R}$ to \mathbb{R} .

Then, the operation of $\mathcal{N}(\widehat{\mathbf{L}}) = -4i(\alpha + 2\beta\widehat{\mathbf{L}})\widehat{\mathbf{L}}^2|4\widehat{\mathbf{L}}^2|^{\epsilon}$ on the $\mathbf{q} = (-q^*, q)^T$ yields

$$-4i(\alpha + 2\beta\widehat{\mathbf{L}})\widehat{\mathbf{L}}^2|4\widehat{\mathbf{L}}^2|^{\epsilon}\mathbf{q} = \frac{1}{\pi} \sum_{\ell=1}^2 \int_{\Gamma_{\infty, \ell}} dk |4k^2|^{\epsilon} f_{\ell}(k) \int_{\mathbb{R}} dy G_{\ell}(x, y, k) \begin{bmatrix} i\alpha(q_{yy}^* + 2|q|^2q^*) - \beta(q_{yyy}^* + 6|q|^2q_y^*) \\ i\alpha(q_{yy} + 2|q|^2q) - \beta(q_{yyy} + 6|q|^2q_y) \end{bmatrix}. \quad (28)$$

Therefore it follows from Eq. (28) and Eq. (7) with $\nu = 1$ that we have the explicit representation of the fHirota equation

$$i \begin{bmatrix} -q^* \\ q \end{bmatrix}_t - \frac{\sigma_3}{\pi} \sum_{\ell=1}^2 \int_{\Gamma_{\infty}^{\ell}} |4k^2|^{\epsilon} f_{\ell}(k) dk \int_{\mathbb{R}} G_{\ell}(x, y, k) \begin{bmatrix} -\alpha(q_{yy}^* + 2|q|^2q^*) + i\beta(q_{yyy}^* + 6|q|^2q_y^*) \\ \alpha(q_{yy} + 2|q|^2q) + i\beta(q_{yyy} + 6|q|^2q_y) \end{bmatrix} dy = 0,$$

that is the fHirota equation is given by

$$iq_t + \frac{1}{\pi} \sum_{\ell=1}^2 \int_{\Gamma_{\infty}^{\ell}} dk |4k^2|^{\epsilon} f_{\ell}(k) \int_{\mathbb{R}} \mathcal{G}_{\ell}(x, y, k) dy = 0, \quad (29)$$

where

$$\begin{aligned} \mathcal{G}_j(x, y, k) &= -\Phi_{-j1}^2(x, k) \left\{ \Phi_{+(3-j)1}^2(y, k) [-\alpha(q_{yy}^* + 2|q|^2q^*) + i\beta(q_{yyy}^* + 6|q|^2q_y^*)] \right. \\ &\quad \left. + \Phi_{+(3-j)2}^2(y, k) [\alpha(q_{yy} + 2|q|^2q) + i\beta(q_{yyy} + 6|q|^2q_y)] \right\}, \quad j = 1, 2. \end{aligned}$$

As $\alpha = 0$, $\beta = 1$, Eq. (29) reduces to the explicit fcmKdV equation. As $\alpha = 1$, $\beta = 0$, Eq. (29) reduces to the explicit fNLS equation. Moreover, at $\epsilon = 0$, one can recover the focusing Hirota equation (2) with $\nu = 1$ from Eq. (29).

Similarly, these results can also be extended to the higher-order case (14). Let

$$\begin{bmatrix} \widehat{q}_1(x, t; \alpha_j, \epsilon) \\ \widehat{q}_2(x, t; \alpha_j, \epsilon) \end{bmatrix} = \left(\sum_{j=2}^N \alpha_j \delta_j \widehat{\mathbf{L}}^j \right) \begin{bmatrix} -q^* \\ q \end{bmatrix}.$$

Then, according to the above-mentioned idea, we have the fractional HONLS equation (13) in the explicit form

$$iq_t + \frac{1}{\pi} \sum_{\ell=1}^2 \int_{\Gamma_{\infty}^{\ell}} dk |4k^2|^{\epsilon} f_{\ell}(k) \int_{\mathbb{R}} \widehat{\mathcal{G}}_{\ell}(x, y, k) dy = 0, \quad (30)$$

where

$$\widehat{G}_j(x, y, k) = -\Phi_{-j}^2(x, k) \left\{ \Phi_{+(3-j)1}^2(y, k) \widehat{q}_1(y, t; \alpha_j, \epsilon) + \Phi_{+(3-j)2}^2(y, k) \widehat{q}_2(y, t; \alpha_j, \epsilon) \right\}, \quad j = 1, 2.$$

Inverse scattering: symmetry, discrete spectrum, and residue conditions.—We consider the symmetry properties of scattering matrix as follows:

$$S(k) = \sigma S^*(k^*) \sigma^{-1}, \quad \sigma = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}. \quad (31)$$

The continuous spectrum set of X_{\pm} is $k \in \mathbb{R}$, which is the jump contour for the following considered Riemann-Hilbert (RH) problem. The discrete spectrum of the scattering problem is the set of $k \in \mathbb{C} \setminus \mathbb{R}$ such that they admit eigenfunctions in $L^2(\mathbb{R})$. For the fHirota equation (7) or fHONLS equation (14), they should satisfy $s_{jj}(k_j) = 0$, $j = 1, 2$ for $k_1 \in D_-, k_2 \in D_+$. We assume that $s_{22}(k)$ has N simple zeros in D_+ denoted by k_n , $n = 1, 2, \dots, N$, that is, $s_{22}(k_n) = 0$ and $s'_{22}(k_n) \neq 0$ ($n = 1, 2, \dots, N$). It follows from Eq. (31) that if $s_{22}(k_n) = 0$, then $s_{11}(k_n^*) = 0$. The discrete spectrum set is $K = \{ \{k_n, k_n^*\} | s_{22}(k_n) = 0, k_n \in D_+ \}$. Since $s_{22}(k_0) = 0$, $s'_{22}(k_0) \neq 0$, $s_{11}(k_0^*) = 0$ and $s'_{11}(k_0^*) \neq 0$ are required for $k_0 \in D_+$, $k_0^* \in D_-$, then it follows from Eq. (23) that there exist two norming constants $b_{\pm}(k_0)$ satisfying

$$\Phi_{+2}(x, t; k_0) = b_+(k_0) \Phi_{-1}(x, t; k_0), \quad \Phi_{+1}(x, t; k_0^*) = b_-(k_0^*) \Phi_{-2}(x, t; k_0^*),$$

which can generate the residue conditions

$$\begin{aligned} \operatorname{Res}_{k=k_0} \left[\frac{\Phi_{+2}(x, t; k)}{s_{22}(z)} \right] &= \frac{\Phi_{+2}(x, t; k_0)}{s'_{22}(k_0)} = \frac{b_+(k_0)}{s'_{22}(k_0)} \Phi_{-1}(x, t; k_0) = A_+[k_0] \Phi_{-1}(x, t; k_0), \\ \operatorname{Res}_{k=k_0^*} \left[\frac{\Phi_{+1}(x, t; k)}{s_{11}(k)} \right] &= \frac{\Phi_{+1}(x, t; k_0^*)}{s'_{11}(k_0^*)} = \frac{b_-(k_0^*)}{s'_{11}(k_0^*)} \Phi_{-2}(x, t; k_0^*) = A_-[k_0^*] \Phi_{-2}(x, t; k_0^*). \end{aligned} \quad (32)$$

Moreover, one has the symmetry $A_+[k_0] = -A_-^*[k_0^*]$, $k_0 \in D_+$, $k_0^* \in D_-$.

Riemann-Hilbert problem.—Let the sectionally meromorphic matrix $M(x, t; k)$ be

$$M(x, t; k) = \begin{cases} \left(\phi_{-1}(x, t; k), \frac{\phi_{+2}(x, t; k)}{s_{22}(k)} \right), & k \in D_+, \\ \left(\frac{\phi_{+1}(x, t; k)}{s_{11}(k)}, \phi_{-2}(x, t; k) \right), & k \in D_-. \end{cases}$$

where $M^{\pm}(x, t; k) = \lim_{\tilde{k} \rightarrow k, \tilde{k} \in D_{\pm}} M(x, t; \tilde{k})$, $k \in \mathbb{R}$. Then according to the properties of $\phi_{\pm j}(x, t; k)$, $j = 1, 2$ and

$$\left(\Phi_{-1}(x, t; k), \frac{\Phi_{+2}(x, t; k)}{s_{22}(k)} \right) = \left(\frac{\Phi_{+1}(x, t; k)}{s_{11}(k)}, \Phi_{-2}(x, t; k) \right) (\mathbb{I} - V_0(k)), \quad V_0(k) = \begin{pmatrix} 0 & -\hat{\rho}(k) \\ \rho(k) & \rho(k)\hat{\rho}(k) \end{pmatrix},$$

one has the following Riemann-Hilbert (RH) problem satisfied by the matrix function $M(x, t; k)$:

- *Analyticity:* $M(x, t; k)$ is analytic in $D_{\pm} \setminus K$, and has the simple poles in K ;
- *Jump relation:* $M^-(x, t; k) = M^+(x, t; k) (\mathbb{I} - V(x, t; k))$ with $V(x, t; k) = e^{(ikx - \frac{1}{2}N(k)t)\hat{\sigma}_3} V_0^{-1}(k)$, $k \in \mathbb{R}$;
- *Asymptoticity:* $M^{\pm}(x, t; k) \rightarrow \mathbb{I}$ for $k \rightarrow \infty$.

Considering the asymptotic behavior and the simple-pole contributions (cf. Eq. (32)) of $M(x, t; k)$

$$\begin{aligned} M_{sp}(x, t; k) &= \mathbb{I} + \sum_{n=1}^N \left[\frac{\operatorname{Res}_{k=k_n} M^+(x, t; k)}{k - k_n} + \frac{\operatorname{Res}_{k=k_n^*} M^-(x, t; k)}{k - k_n^*} \right] \\ &= \mathbb{I} + \left[\frac{A_-[k_n^*] e^{-2i\theta_{\epsilon}(x, t; k_n^*)}}{k - k_n^*} \phi_{-2}(x, t; k_n^*), \frac{A_+[k_n] e^{2i\theta_{\epsilon}(x, t; k_n)}}{k - k_n} \phi_{-1}(x, t; k_n) \right], \end{aligned} \quad (33)$$

where $\theta_\epsilon(x, t; k) = -kx + \frac{1}{2t}\mathcal{N}(k)t = -kx - 2(\alpha + 2\beta k)k^2|4k^2|^\epsilon t$ and then subtracting out $M_{sp}(x, t; k)$ from both sides of the above-given jump relation can generate

$$M^-(x, t; k) - M_{sp}(x, t; k) = M^+(x, t; k) - M_{sp}(x, t; k) - M^+(x, t; k) V(x, t; k), \quad (34)$$

where $M^\pm(x, t; k) - M_{sp}(x, t; k)$ are analytic in D_\pm . The Cauchy projectors $P^\pm[f](k) = \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{f(\zeta)}{\zeta - (k \pm i0)} d\zeta$ (where the signs $k \pm i0$ denotes the limit chosen from the left/right of k) and Plemelj's formulae are used to solve Eq. (34) to find the solution of the RH problem

$$M(x, t; k) = M_{sp}(x, t; k) + \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{M^+(x, t; \zeta) V(x, t; \zeta)}{\zeta - k} d\zeta, \quad k \in \mathbb{C} \setminus \mathbb{R}. \quad (35)$$

As $k = k_s (k_s^*)$, $s = 1, 2, \dots, N$, it follows from the first (second) column of $M(x, t; k)$ given by Eq. (35) with Eq. (33) that

$$\begin{aligned} \phi_{-1}(x, t; k_s) &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \sum_{n=1}^N \frac{A_-[k_n^*] e^{-2i\theta_\epsilon(x, t; k_n^*)}}{k_s - k_n^*} \phi_{-2}(x, t; k_n^*) + \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{(M^+V)_1(x, t; \zeta)}{\zeta - k_s} d\zeta, \quad s = 1, 2, \dots, N, \\ \phi_{-2}(x, t; k_s^*) &= \begin{bmatrix} 0 \\ 1 \end{bmatrix} + \sum_{n=1}^N \frac{A_+[k_n] e^{2i\theta_\epsilon(x, t; k_n)}}{k_s^* - k_n} \phi_{-1}(x, t; k_n) + \frac{1}{2\pi i} \int_{\mathbb{R}} \frac{(M^+V)_2(x, t; \zeta)}{\zeta - k_s^*} d\zeta, \quad s = 1, 2, \dots, N. \end{aligned} \quad (36)$$

According to the condition $q(x, t) = \lim_{k \rightarrow \infty} (kM)_{12}$ and the solutions $\phi_{-11}(x, t; k_n)$, $n = 1, 2, \dots, N$ given by system (36), we have the fractional simple-pole solutions of the fractional Hirota equation as

$$q(x, t) = 2i \sum_{n=1}^N A_+[k_n] e^{2i\theta_\epsilon(x, t; k_n)} \phi_{-11}(x, t; k_n) + \frac{1}{2\pi i} \int_{\mathbb{R}} (M^+V)_{12}(x, t; \zeta) d\zeta. \quad (37)$$

Since the scattering coefficients $s_{22}(k)$ and $s_{11}(k)$ are analytic in D_+ and D_- , respectively, and the discrete spectral points k_n 's and k_n^* 's are the simple zeros of $s_{22}(k)$ and $s_{11}(k)$, respectively, then one can also find the trace formulae for the fHirota equation

$$s_{22}(k) = e^{s(k)} s_0(k) \text{ for } k \in D_+, \quad s_{11}(k) = e^{-s(k)} / s_0(k) \text{ for } k \in D_-,$$

where $s(k) = \frac{i}{2\pi} \int_{\mathbb{R}} \frac{\log[1 + \rho(\zeta)\rho^*(\zeta^*)]}{\zeta - k} d\zeta$ and $s_0(k) = \prod_{n=1}^N \frac{(k - k_n)}{(k - k_n^*)}$.

3 Dynamics of fractional N -soliton solutions

Fractional N -soliton solutions.—We consider the case of reflectionless potential (i.e., $V = 0$) in system (36) such that we have the system of equations with respect to $\phi_{-11}(x, t; k_n)$, $n = 1, 2, \dots, N$

$$\left(1 - \sum_{n_1=1}^N \sum_{n_2=1}^N \frac{A_-[k_{n_1}^*] e^{-2i\theta_\epsilon(x, t; k_{n_1}^*)}}{k_n - k_{n_1}^*} \frac{A_+[k_{n_2}] e^{2i\theta_\epsilon(x, t; k_{n_2})}}{k_{n_1}^* - k_{n_2}} \right) \phi_{-11}(x, t; k_{n_2}) = 1, \quad n = 1, 2, \dots, N. \quad (38)$$

According to the condition $q(x, t) = \lim_{k \rightarrow \infty} (kM)_{12}$ and the solutions $\phi_{-11}(x, t; k_n)$, $n = 1, 2, \dots, N$ given by system (38), we have the fractional N -soliton solutions (reflectionless case) with simple poles of the fHirota equation as

$$q(x, t) = 2i \sum_{n=1}^N A_+[k_n] e^{2i\theta_\epsilon(x, t; k_n)} \phi_{-11}(x, t; k_n) = -2i \frac{\begin{vmatrix} \mathbb{I}_N + H & \mathbf{a}^T \\ \mathbf{b} & 0 \end{vmatrix}}{|\mathbb{I}_N + H|}, \quad (39)$$




Figure 1: Fractional 1-soliton solutions (40) for $k_1 = 1 + i$, $A_+[k_1] = 1$ and $\epsilon = 0, 0.5, 0.9$. (a1)-(a4) fHirota at $\alpha = 1$, $\beta = 0.05$; (b1)-(b4) fcmKdV at $\alpha = 0$, $\beta = 1$.

where \mathbb{I}_N is an $N \times N$ unit matrix, $\theta_\epsilon(x, t; k) = -kx - 2(\alpha + 2\beta k)k^2|4k^2|^\epsilon t$,

$$\mathbf{a} = (a_j)_{1 \times N} = (1, 1, \dots, 1), \quad \mathbf{b} = (b_j)_{1 \times N} = (b_1, b_2, \dots, b_N), \quad b_n = A_+[k_n]e^{2i\theta_\epsilon(x, t; k_n)}, \quad n = 1, 2, \dots, N,$$

$$H = (H_{(n, n_2)})_{N \times N}, \quad H_{(n, n_2)} = \sum_{n_1=1}^N \frac{A_-[k_{n_1}^*] e^{-2i\theta_\epsilon(x, t; k_{n_1}^*)}}{k_n - k_{n_1}^*} \frac{A_+[k_{n_2}] e^{2i\theta_\epsilon(x, t; k_{n_2})}}{k_{n_2} - k_{n_1}^*}.$$

In what follows we will discuss some fractional one- two-, and three-soliton solutions of the fHirota, fNLS and fcmKdV equations in terms of solution (39) with $N = 1, 2, 3$.

Fractional one-soliton solutions and velocities.—For $N = 1$, if we choose the spectral parameter $k_1 = \zeta + i\eta$ and $A_+[k_1] = a$ with $\zeta, \eta \in \mathbb{R}$ with $a\eta \neq 0$, then the expression of fractional 1-soliton solution of the fHirota equation can be written as:

$$q(x, t) = 2\eta \operatorname{sech} \left[2\eta x + 8\eta (\alpha\zeta + \beta(3\zeta^2 - \eta^2)) |2k_1|^{2\epsilon} t - \ln |2\eta/a| \right] e^{-2i\zeta x + 4i[\alpha(\eta^2 - \zeta^2) + 2\beta\zeta(3\eta^2 - \zeta^2)] |2k_1|^{2\epsilon} t + i(\pi/2 + \vartheta)}, \quad (40)$$

where $\vartheta = \operatorname{Arg}(a/2\eta)$.

Particularly, as $\beta = 0$, we have the fractional 1-soliton of the fNLS equation. When $\alpha = 0$, we have the fractional 1-soliton of the fcmKdV equation. It follows from the solution (40) that the wave velocity $v_{wv}(\zeta, \eta, \epsilon; \alpha, \beta)$, group velocity $v_{gv}(\zeta, \eta, \epsilon; \alpha, \beta)$ and phase velocity $v_{pv}(\zeta, \eta, \epsilon; \alpha, \beta)$ of the fractional 1-soliton solutions, respectively, are

$$\begin{aligned} v_{wv}(\zeta, \eta, \epsilon; \alpha, \beta) &= -4^{1+\epsilon} [\alpha\zeta + \beta(3\zeta^2 - \eta^2)] (\zeta^2 + \eta^2)^\epsilon, \\ v_{gv}(\zeta, \eta, \epsilon; \alpha, \beta) &= 4^{1+\epsilon} \left\{ \epsilon\zeta [\alpha(\eta^2 - \zeta^2) + 2\beta\zeta(3\eta^2 - \zeta^2)] - (\zeta^2 + \eta^2) [\alpha\zeta + 3\beta(\zeta^2 - \eta^2)] \right\} (\zeta^2 + \eta^2)^{\epsilon-1}, \\ v_{pv}(\zeta, \eta, \epsilon; \alpha, \beta) &= 2^{1+2\epsilon} [\alpha\zeta^{-1}(\eta^2 - \zeta^2) + 2\beta(3\eta^2 - \zeta^2)] (\zeta^2 + \eta^2)^\epsilon, \end{aligned} \quad (41)$$

which depend on the real (ζ) and imaginary (η) parts of the special parameter k_1 , fractional parameter ϵ and equation coefficients α, β .

For the given $\zeta = \eta = 1$, i.e., $k_1 = 1 + i$, Figs. 1(a1-a4) display the fractional 1-soliton solution of the fHirota equation as $\alpha = 1$, $\beta = 0.05$, which are left-going travelling-wave solitons ($v_{wv} = -1.1 \times 2^{2+3\epsilon} < 0$)




Figure 2: (a) wave velocities, (b) group velocities and (c) phase velocities given by Eq. (41) of fractional 1-soliton solutions of fHirota (dashed line, $\alpha = 1, \beta = 0.05$), fNLS (solid line, $\alpha = 1, \beta = 0$) and fcmKdV (dash-dotted line, $\alpha = 0, \beta = 1$) equations at $k_1 = 1 + i, A_+[k_1] = 1$.

at $\epsilon = 0, 0.5, 0.9$. Moreover, the absolute value of left-going travelling-wave velocity becomes larger as ϵ increases (see Fig. 2(a)). Figs. 1(b1-b4) display the fractional 1-soliton solution of the fcmKdV equation as $\alpha = 0, \beta = 1$, which are also left-going travelling-wave solitons ($v_{wv} = -2^{3+3\epsilon} < 0$) at $\epsilon = 0, 0.5, 0.9$. In fact, if we take $\alpha\zeta + \beta(3\zeta^2 - \eta^2) < 0$, then we have the right-going travelling-wave solitons of the fHirota equation with (e.g., $\zeta = \eta = 1, \alpha = -1, \beta = -0.05$), fNLS equation (e.g., $\zeta = \eta = 1, \alpha = -1, \beta = 0$), and fcmKdV equation (e.g., $\zeta = \eta = 1, \alpha = 0, \beta = -0.05$). Fig. 2(a) displays the wave velocities (see v_{wv} in Eq. (41)) of the fractional 1-solitons solutions of the fHirota, fNLS and fcmKdV equations, where the absolute values of the wave velocities increase as ϵ grows. Figs. 2(b, c) show the group velocities (see v_{gv} in Eq. (41)) and phase velocities (see v_{pv} in Eq. (41)) of the fractional 1-soliton solutions of the fHirota, fNLS and fcmKdV equations, respectively.

Elastic interactions of fractional two-soliton solutions.—For $N = 2$, if we take two spectral parameters k_1, k_2 , and $A_+[k_1] = a, A_+[k_2] = b$, then the fractional two-soliton solutions of the fHirota equation are

$$q(x, t) = -2i \frac{A(x, t)}{B(x, t)}, \quad (42)$$

where

$$A(x, t) = \begin{pmatrix} 1 - \frac{|a|^2 e^{2i\theta_\epsilon(k_1^*) - 2i\theta_\epsilon(k_1)}}{(k_1^* - k_1)^2} - \frac{a^* b e^{2i\theta_\epsilon(k_1^*) - 2i\theta_\epsilon(k_2)}}{(k_1^* - k_2)^2} & -\frac{ab^* e^{2i\theta_\epsilon(k_2^*) - 2i\theta_\epsilon(k_1)}}{(k_1^* - k_1)(k_2^* - k_1)} - \frac{|b|^2 e^{2i\theta_\epsilon(k_2^*) - 2i\theta_\epsilon(k_2)}}{(k_1^* - k_2)(k_2^* - k_2)} & 1 \\ -\frac{|a|^2 e^{2i\theta_\epsilon(k_1^*) - 2i\theta_\epsilon(k_1)}}{(k_1^* - k_1)(k_2^* - k_1)} - \frac{a^* b e^{2i\theta_\epsilon(k_1^*) - 2i\theta_\epsilon(k_2)}}{(k_1^* - k_2)(k_2^* - k_2)} & 1 - \frac{ab^* e^{2i\theta_\epsilon(k_2^*) - 2i\theta_\epsilon(k_1)}}{(k_2^* - k_1)^2} - \frac{|b|^2 e^{2i\theta_\epsilon(k_2^*) - 2i\theta_\epsilon(k_2)}}{(k_2^* - k_2)^2} & 1 \\ -a^* e^{2i\theta_\epsilon(k_1^*)} & -b^* e^{2i\theta_\epsilon(k_2^*)} & 0 \end{pmatrix},$$

$$B(x, t) = \begin{pmatrix} 1 - \frac{|a|^2 e^{2i\theta_\epsilon(k_1^*) - 2i\theta_\epsilon(k_1)}}{(k_1^* - k_1)^2} - \frac{a^* b e^{2i\theta_\epsilon(k_1^*) - 2i\theta_\epsilon(k_2)}}{(k_1^* - k_2)^2} & -\frac{ab^* e^{2i\theta_\epsilon(k_2^*) - 2i\theta_\epsilon(k_1)}}{(k_1^* - k_1)(k_2^* - k_1)} - \frac{|b|^2 e^{2i\theta_\epsilon(k_2^*) - 2i\theta_\epsilon(k_2)}}{(k_1^* - k_2)(k_2^* - k_2)} & 1 \\ -\frac{|a|^2 e^{2i\theta_\epsilon(k_1^*) - 2i\theta_\epsilon(k_1)}}{(k_1^* - k_1)(k_2^* - k_1)} - \frac{a^* b e^{2i\theta_\epsilon(k_1^*) - 2i\theta_\epsilon(k_2)}}{(k_1^* - k_2)(k_2^* - k_2)} & 1 - \frac{ab^* e^{2i\theta_\epsilon(k_2^*) - 2i\theta_\epsilon(k_1)}}{(k_2^* - k_1)^2} - \frac{|b|^2 e^{2i\theta_\epsilon(k_2^*) - 2i\theta_\epsilon(k_2)}}{(k_2^* - k_2)^2} & 1 \end{pmatrix}.$$

In particular, as $\beta = 0$, we have the fractional two-soliton solutions of the fNLS equation. When $\alpha = 0$, we get the fractional two-soliton solutions of the fcmKdV equation.

For the two chosen spectral parameters $k_1 = 1 + i, k_2 = 2i$ and $A_+[k_1] = A_+[k_2] = 1$, Figs. 3(a1-a4) display the fractional 2-soliton solution of the fHirota equation as $\alpha = 1, \beta = 0.05$, which imply the elastic interactions




Figure 3: Elastic interactions of fractional two-soliton solutions (42) for $k_1 = 1 + i$, $k_2 = 2i$, $A_+[k_1] = A_+[k_2] = 1$ and $\epsilon = 0, 0.5, 0.9$. (a1)-(a4) fHirota at $\alpha = 1$, $\beta = 0.05$; (b1)-(b4) fNLS at $\alpha = 1$, $\beta = 0$; (c1)-(c4) fcmKdV at $\alpha = 0$, $\beta = 1$.

of one right-going (left branch) and another left-going (right branch) travelling-wave solitons at $\epsilon = 0, 0.5, 0.9$. In particular, it follows from Figs. 3(a4) that before the elastic collision, the left-branch is a right-going travelling wave with larger amplitude, and the right branch is a left-going travelling wave with lower amplitude. Figs. 3(b1-b4) display the fractional two-soliton solutions of the fNLS equation as $\alpha = 0$, $\beta = 1$, which also imply the elastic interactions of one left-going and another center-going travelling-wave solitons at $\epsilon = 0, 0.5, 0.9$. Figs. 3(c1-c4) show the fractional two-soliton solutions of the fcmKdV equation as $\alpha = 0$, $\beta = 1$, which are also elastic interactions of one left-going and another right-going travelling-wave solitons at $\epsilon = 0, 0.5, 0.9$.

Elastic interactions of fractional three-soliton solutions.—For $N = 3$, if we take three spectral parameters k_1, k_2, k_3 and $A_+[k_1] = a$, $A_+[k_2] = b$, $A_+[k_3] = c$, then the fractional three-soliton solutions of the fHirota equation are given by Eq. (39). As $\beta = 0$, we have the fractional three-soliton solution of the fNLS equation. When $\alpha = 0$, we have the fractional three-soliton solution of the fcmKdV equation.

For the three taken spectral parameters $k_1 = 1 + i$, $k_2 = 2i$, $k_3 = -2 + i$ and $A_+[k_1] = A_+[k_2] = A_+[k_3] = 1$, Figs. 4(a1-a4) display the fractional 3-soliton solution of the fHirota equation as $\alpha = 1$, $\beta = 0.05$, which imply the elastic interactions of two right-going (left and center branches) and another left-going (right branch) travelling-wave solitons at $\epsilon = 0, 0.5, 0.9$. Figs. 4(b1-b4) display the fractional three-soliton solutions of the fNLS equation as $\alpha = 0$, $\beta = 1$, which also imply the elastic interactions of one left-going, one center-going and another right-going travelling-wave solitons at $\epsilon = 0, 0.5, 0.9$. Figs. 4(c1-c4) show the fractional three-soliton solutions of the fcmKdV equation as $\alpha = 0$, $\beta = 1$, which are also elastic interactions of two left-going and another right-going travelling-wave solitons at $\epsilon = 0, 0.5, 0.9$.




Figure 4: Elastic interactions of fractional three-soliton solutions (39) with $N = 3$ for $k_1 = 1 + i$, $k_2 = 2i$, $k_3 = -2 + i$, $A_+[k_1] = A_+[k_2] = A_+[k_3] = 1$ and $\epsilon = 0, 0.5, 0.9$. (a1)-(a4) fHirota at $\alpha = 1$, $\beta = 0.05$; (b1)-(b4) fNLS at $\alpha = 1$, $\beta = 0$; (c1)-(c4) fcmKdV at $\alpha = 0$, $\beta = 1$.

In fact, one can also study the other fractional N ($N > 3$)-soliton solutions in terms of the formula (39).

Fractional N -soliton solutions of fHONLS equations.—Similarly, we can also solve the integrable fractional HONLS equations (13) by the IST with the matrix RH problem such that the fractional N -soliton solutions of Eq. (13) is given by Eq. (39), where θ_ϵ is taken place by the general form

$$\theta_\epsilon(x, t; k) = -kx + \frac{1}{2} \left(\sum_{j=2}^N \alpha_j i^{\epsilon_j + j} (-2k)^j \right) |4k^2|^{\epsilon t}.$$

4 Conclusions and discussions

In conclusion, we have analyzed the explicit forms and anomalous dispersive relations of the fractional higher-order NLS equations containing the fHirota, fcmKdV, and fLPD equations. We investigate the IST with the RH problem to study fractional N -soliton solutions of the fHirota and fcmKdV equations. Moreover, we analyze the elastic interactions of the fractional two- and three-soliton solutions. The wave, group, and phase velocities of these envelope fractional one-soliton solutions are shown to be related to the power law of their amplitudes. We also deduce the general formula for the fractional N -soliton solutions of integrable generalized fractional higher-order NLS equations. The used idea can also be extended to other integrable fractional nonlinear evolution equations. These results will be useful to display the super-dispersive transports of nonlinear waves in fractional nonlinear media.

Acknowledgments

We thank G. Zhang for useful discussions. The work was supported by the National Natural Science Foundation of China (No. 11925108).

References

- [1] C. S. Gardner, J. M. Greene, M. D. Kruskal, and R. M. Miura, Method for solving the Korteweg-de Vries equation, *Phys. Rev. Lett.* 19, 1095-1097 (1967).
- [2] P. D. Lax, Integrals of nonlinear equations of evolution and solitary waves, *Commun. Pure Appl. Math.* 21 (1968) 467-490.
- [3] M. J. Ablowitz and H. Segur, *Solitons and the Inverse Scattering Transform* (SIAM, Philadelphia, 1981).
- [4] L. D. Faddeev and L. A. Takhtajan, *Hamiltonian Methods in the Theory of Solitons* (Springer, Berlin, 1987).
- [5] M. J. Ablowitz and P. A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering* (Cambridge University Press, Cambridge, 1991).
- [6] M. J. Ablowitz, B. Prinari, and A. D. Trubatch, *Discrete and continuous nonlinear Schrödinger systems* (Cambridge University Press, Cambridge, 2004).
- [7] K.B. Oldham, and J. Spanier, *The Fractional Calculus* (Academic Press, New York, 1974).
- [8] K.S. Miller, and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations* (John Wiley & Sons Inc., New York, 1993).
- [9] S. Das, *Introduction to Fractional Calculus* (Springer, Berlin, 2011).
- [10] R. Metzler and J. Klafter, The random walk's guide to anomalous diffusion: A fractional dynamics approach, *Phys. Rep.* 339, 1 (2000).
- [11] M. F. Shlesinger, B. J. West, and J. Klafter, Lévy dynamics of enhanced diffusion: Application to turbulence, *Phys. Rev. Lett.* 58, 1100 (1987).
- [12] B. J. West, P. Grigolini, R. Metzler, and T. F. Nonnenmacher, Fractional diffusion and Lévy stable processes, *Phys. Rev. E* 55 (1997) 99-106.
- [13] N. Laskin, Fractional Schrödinger equation, *Phys. Rev. E* 66, 056108 (2002).
- [14] S. Longhi, Fractional Schrödinger equation in optics, *Opt. Lett.* 40, 1117 (2015).
- [15] W. P. Zhong, M. R. Belić, B. A. Malomed, Y. Zhang, and T. Huang, Spatiotemporal accessible solitons in fractional dimensions, *Phys. Rev. E* 94, 012216 (2016).
- [16] U. Al Khawaja, M. Al-Refai, G. Shchedrin, and L. D. Carr, High-accuracy power series solutions with arbitrarily large radius of convergence for the fractional nonlinear Schrödinger-type equations, *J. Phys.* 51, 235201 (2018).
- [17] Y. Qiu, B. A. Malomed, D. Mihalache, X. Zhu, X. Peng, and Y. He, Stabilization of single-and multi-peak solitons in the fractional nonlinear Schrödinger equation with a trapping potential, *Chaos Solitons Fractals* 140, 110222 (2020).
- [18] B. A. Malomed, Optical solitons and vortices in fractional media: A mini-review of recent results, *Photonics* 8 (2021) 353.
- [19] M. Riesz, L'intégrale de Riemann-Liouville et le problème de Cauchy, *Acta Math.* 81 (1949) 1-222.
- [20] M. Cai and C.P. Li, On Riesz derivative, *Fractional Calculus Appl. Anal.* 22 (2019) 287-301.
- [21] A. Lischke, G. Pang, M. Gulian, F. Song, C. Glusa, X. Zheng, Z. Mao, W. Cai, M. M. Meerschaert, M. Ainsworth, and G. E. Karniadakis, What is the fractional Laplacian ? A comparative review with new results, *J. Comput. Phys.* 404, 109009 (2020).
- [22] M. J. Ablowitz, J. B. Been, and L. D. Carr, Fractional integrable nonlinear soliton equations, *Phys. Rev. Lett.* 128, 184101 (2022).

- [23] M. J. Ablowitz, J. B. Been, and L. D. Carr, Integrable fractional modified Korteweg-de Vries, sine-Gordon, and sinh-Gordon equations, arXiv: 2203.13755.
- [24] G. P. Agrawal, *Nonlinear Fiber Optics* (5th edn.) (New York, Academic Press, 2012).
- [25] Y. S. Kivshar and G. P. Agrawal, *Optical Solitons: from Fibers to Photonic Crystals* (New York, Academic Press, 2013).
- [26] Y. Kodama, Optical solitons in a monomode fiber, *J. Stat. Phys.* 39, 597 (1985).
- [27] Y. Kodama and A. Hasegawa, Nonlinear pulse propagation in a monomode dielectric guide, *IEEE J. Quantum Electron.* 23, 510 (1987).
- [28] D. Mihalache, Localized structures in optical and matter-wave media: a selection of recent studies, *Rom. Rep. Phys.* 73, 403 (2021)
- [29] R. Hirota, Exact envelope-soliton solutions of a nonlinear wave equation, *J. Math. Phys.* 14, 805 (1973).
- [30] A. Ankiewicz, J. M. Soto-Crespo, and N. Akhmediev, Rogue waves and rational solutions of the Hirota equation. *Phys. Rev. E* 81, 046602 (2010).
- [31] Y. Tao and J. He, Multisolitons, breathers, and rogue waves for the Hirota equation generated by the Darboux transformation, *Phys. Rev. E* 85, 026601 (2012).
- [32] Z. Yan and C. Dai, Optical rogue waves in the generalized inhomogeneous higher-order nonlinear Schrödinger equation with modulating coefficients, *J. Opt.* 15, 064012 (2013).
- [33] Y. Yang, Z. Yan, and B. A. Malomed, Rogue waves, rational solitons, and modulational instability in an integrable fifth-order nonlinear Schrödinger equation, *Chaos* 25, 103112 (2015).
- [34] M. Lakshmanan, K. Porsezian, M. Daniel Effect of discreteness on the continuum limit of the Heisenberg spin chain, *Phys. Lett. A*, 133 (1988) 483-488.
- [35] A. Chowdury, D. J. Kedziora, A. Ankiewicz, and N. Akhmediev, Soliton solutions of an integrable nonlinear Schrödinger equation with quintic terms, *Phys. Rev. E* 90, 032922 (2014).
- [36] T. Kano, Normal form of nonlinear Schrödinger equation, *J. Phys. Soc. Jpn.* 58, 4322 (1989).
- [37] V. E. Zakharov and A. B. Shabat, Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media, *Sov. Phys. JETP* 34, 62-69 (1972) [*Zh. Eksp. Teor. Fiz.* 61 (1971) 118-134].
- [38] M. J. Ablowitz, D. J. Kaup, A. C. Newell, and H. Segur, The inverse scattering transform-Fourier analysis for nonlinear problems, *Stud. Appl. Math.* 53 (1974) 249-315.
- [39] G. Biondini, and G. Kovacic, Inverse scattering transform for the focusing nonlinear Schrödinger equation with nonzero boundary conditions, *J. Math. Phys.* 55, 031506 (2014).
- [40] G. Zhang, S. Chen, and Z. Yan, Focusing and defocusing Hirota equations with non-zero boundary conditions: Inverse scattering transforms and soliton solutions, *Commun. Nonlinear Sci. Numer. Simulat.* 80 (2020) 104927.
- [41] X. Zhou, Direct and inverse scattering transforms with arbitrary spectral singularities, *Commun. Pure Appl. Math.* 42, 895-938 (1989).
- [42] D. J. Kaup. Closure of the squared Zakharov-Shabat eigenstates, *J. Math. Anal. Appl.* 54 (1976) 849-864.