

NILPOTENT PROBABILITY OF COMPACT GROUPS

ALIREZA ABDOLLAHI AND MEISAM SOLEIMANI MALEKAN

ABSTRACT. Let k be any positive integer and G a compact (Hausdorff) group. Let $\mathbf{np}_k(G)$ denote the probability that $k+1$ randomly chosen elements x_1, \dots, x_{k+1} satisfy $[x_1, x_2, \dots, x_{k+1}] = 1$. We study the following problem: If $\mathbf{np}_k(G) > 0$ then, does there exist an open nilpotent subgroup of class at most k ? The answer is positive for profinite groups and we give a new proof. We also prove that the connected component G^0 of G is abelian and there exists a closed normal nilpotent subgroup N of class at most k such that $G^0 N$ is open in G .

1. INTRODUCTION AND RESULTS

Let G be a compact (Hausdorff) group and denote by \mathbf{m}_G the (unique) Haar measure on G . The following question is proposed in [11, Question 1.20.]:

Question 1.1. Let G be a compact group, and suppose that

$$\mathcal{N}_k(G) := \{(x_1, \dots, x_{k+1}) \in G^{k+1} \mid [x_1, \dots, x_{k+1}] = 1\},$$

has positive Haar measure in G^{k+1} . Does G have an open k -step nilpotent subgroup?

Here $[x_1, x_2] = x_1^{-1}x_2^{-1}x_1x_2$ and $[x_1, \dots, x_{k+1}]$ is inductively defined by $[[x_1, \dots, x_k], x_{k+1}]$ for all $k > 1$. The Haar measure of $\mathcal{N}_k(G)$ is called the degree of k -nilpotence of the group G , and denoted in [11] by $\mathbf{dc}^k(G)$.

Denote by \mathbf{PA}_k the class of compact groups giving positive answer to Question 1.1. In [10] it is proved that \mathbf{PA}_1 contains profinite groups and in [8] it is proved that all compact groups includes \mathbf{PA}_1 . In [1] it is shown that \mathbf{PA}_2 contains all compact groups. Martino et al. have shown that \mathbf{PA}_k contains all profinite groups for any k (see [11, Theorem 1.19.]). Here in Theorem 2.12 below, we give another proof of the latter result without using submultiplicativity of the k -nilpotence degree of a finite group i.e. for a normal subgroup N of a finite group

2010 *Mathematics Subject Classification.* 20E18; 20P05.

Key words and phrases. Nilpotent probability; Compact group.

G , $\mathbf{dc}^k(G) \leq \mathbf{dc}^k(G/N)\mathbf{dc}^k(N)$ (see [11, Theorem 1.21.]).

Our main result is the following.

Theorem 1.2. *Let G be a compact group with $\mathbf{np}_k(G) > 0$. Then the connected component G^0 of G is abelian and there exists a closed normal nilpotent subgroup N of class at most k such that G^0N is open in G .*

2. RELATIVE NILPOTENT PROBABILITY AND SUBMULTIPLICATIVITY

We denote the set of continuous functions on a compact group G by $\mathcal{C}(G)$. If N is a closed normal subgroup of G , then G/N is a compact group, equipped with quotient topology. There exists a projection-like map $P : \mathcal{C}(G) \rightarrow \mathcal{C}(G/N)$ defined by

$$Pf(\bar{g}) := \int_N f(gy) \, dy,$$

where dy stands for the Haar measure on N . We will use the following two simple lemmas repeatedly:

Lemma 2.1. *Let G be a compact group and N be a closed normal subgroup of G . Then, for all $f \in \mathcal{C}(G)$*

$$\int_G f(g) \, dg = \int_{G/N} Pf(\bar{g}) \, d\bar{g}.$$

Proof. The functional

$$I : \mathcal{C}(G) \rightarrow \mathbb{C}, \quad f \mapsto \int_{G/N} Pf(\bar{g}) \, d\bar{g}$$

is nonzero, positive and left-invariant, also $I(1) = 1$, so by the uniqueness of Haar integral the result holds. \square

Remark 2.2. We have an analog of Lemma 2.1 whenever H is a closed not necessarily normal subgroup, see [4, (2.49) Theorem].

The Uniqueness of Harr measure has another achievement; consider a continuous epimorphism $\phi : G \rightarrow H$ between compact groups. Then

$$I : \mathcal{C}(H) \rightarrow \mathbb{C}, \quad f \mapsto \int_G f \circ \phi(g) \, dg$$

defines a nonzero, positive and left-invariant functional. Aslo, $I(1) = 1$, so we have

Lemma 2.3. *Let $\phi : G \rightarrow H$ be a continuous epimorphism between compact groups. Then*

$$\int_H f(h) dh = \int_G f \circ \phi(g) dg \quad (f \in \mathcal{C}(H)).$$

Lemma 2.4. *Let H be a closed subgroup of a group G . Then $yC_G(x) \cap H$ is either empty or a coset of $C_H(x)$, for all $x, y \in G$. In particular, we have $\mathbf{m}_H(yC_G(x) \cap H) \leq \mathbf{m}_H(C_H(x))$, for all $x, y \in G$.*

Proof. If $z \in C_G(x) \cap y^{-1}H$, then one can easily check that $C_G(x) \cap y^{-1}H = zC_H(x)$. \square

Proposition 2.5. *Let H be a closed subgroup of a compact group G . Then $\mathbf{np}(H; x, y) \leq \mathbf{cp}(H)$ for all x and y in G .*

Proof. By the above lemma, we have

$$\begin{aligned} \mathbf{np}(H; x, y) &= \int_H \int_H 1_{\mathcal{N}(H; x, y)}(\xi, \eta) d\eta d\xi \\ &= \int_H \mathbf{m}_H(x^{-1}C_G(y\eta) \cap N) d\eta \\ &\leq \int_H \mathbf{m}_H(C_H(y\eta)) d\eta \\ &= \int_H \mathbf{m}_H(y^{-1}C_G(\xi) \cap N) d\xi \\ &\leq \int_H \mathbf{m}_H(C_H(\xi)) d\xi = \mathbf{cp}(H). \end{aligned}$$

\square

Remark 2.6. In [11, Lemma 6.2.], using combinatorics methods, the authors have shown that for a normal subgroup N of a *finite* group G ,

$$\mathbf{np}(N; x_1, \dots, x_n) \leq \mathbf{np}(N; \overbrace{1, \dots, 1}^{n\text{-times}})$$

for all $x_k \in G$, $1 \leq k \leq n$. We could not implement these methods in the case of infinite compact groups to have the above inequality for a closed normal subgroup N , and of course we do not need this generalization.

Now, we are going to prove that $\mathbf{np}(N; x_1, \dots, x_n)$ is bounded away from 1, for all closed normal subgroup N of a compact group G which is not nilpotent of class at most k and all $x_k \in G$, $1 \leq k \leq n$.

Proposition 2.7. *Let H be a closed subgroup of a compact group G . Then*

$$\mathbf{np}(H; x_1, \dots, x_{n+1}) \leq \frac{1}{2} \left(1 + \mathbf{np} \left(\frac{H}{\mathbf{Z} \cap H}; \bar{x}_1, \dots, \bar{x}_n \right) \right),$$

for all $x_1, \dots, x_n \in G$, where \mathbf{Z} is the center of G , and \bar{x} denotes the image of x in $H/(\mathbf{Z} \cap H)$.

Proof. Put $K := \mathbf{Z} \cap H$,

$$X = \{\mathbf{y} = (y_1, \dots, y_n) \in H^n : [x_1 y_1, \dots, x_n y_n] \in K\}$$

and

$$\mathcal{N} = \mathcal{N} \left(\frac{H}{K}; \bar{x}_1, \dots, \bar{x}_n \right).$$

Then one can easily verify that

$$(2.1) \quad 1_X(\mathbf{y}\mathbf{z}) = 1_X(\mathbf{y}) \quad (\mathbf{y} \in H^n, \mathbf{z} \in K^n)$$

(1_X is the characteristic function of X) and

$$(2.2) \quad 1_X(\mathbf{y}) = 1 \quad \text{if and only if} \quad 1_{\mathcal{N}}(\bar{\mathbf{y}}) = 1.$$

Now, we have

$$\begin{aligned} \mathbf{np}(H; x_1, \dots, x_{n+1}) &= \int_{H^n} \mathbf{m}_H(x_{n+1}^{-1} C_G([x_1 y_1, \dots, x_n y_n]) \cap H) \, d\mathbf{y} \\ &\leq \int_{H^n} \mathbf{m}_H(C_H([x_1 y_1, \dots, x_n y_n])) \, d\mathbf{y} \\ &= \int_X \mathbf{m}_H(C_H([x_1 y_1, \dots, x_n y_n])) \, d\mathbf{y} \\ &\quad + \int_{H^n \setminus X} \mathbf{m}_H(C_H([x_1 y_1, \dots, x_n y_n])) \, d\mathbf{y} \\ &\leq \mathbf{m}_{H^n}(X) + \frac{1}{2}(1 - \mathbf{m}_{H^n}(X)) \\ &= \frac{1}{2}(1 + \mathbf{m}_{H^n}(X)). \end{aligned}$$

The latter inequality holds because on X^c , $C_G([x_1y_1, \dots, x_ny_n])$ has index at least 2. We compute $\mathbf{m}_{H^n}(X)$:

$$\begin{aligned} \mathbf{m}_{H^n}(X) &= \int_{H^n} 1_X(\mathbf{y}) \, d\mathbf{y} \\ &= \int_{H^n/K^n} \int_{K^n} 1_X(\mathbf{y}\mathbf{z}) \, d\mathbf{z} \, d\bar{\mathbf{y}}, \quad \text{by Lemma 2.1,} \\ &= \int_{H^n/K^n} 1_X(\mathbf{y}) \, d\bar{\mathbf{y}}, \quad \text{by (2.1),} \\ &= \int_{(H/K)^n} 1_{\mathcal{N}}(\bar{\mathbf{y}}) \, d\bar{\mathbf{y}}, \quad \text{by Lemma 2.3,} \\ &= \mathbf{np} \left(\frac{H}{\mathbf{Z} \cap H}; \bar{x}_1, \dots, \bar{x}_n \right). \end{aligned}$$

□

By Theorem 2.7 we obtain the following characterization of closed and nilpotent normal subgroups of a compact group:

Proposition 2.8. *Let H be a closed subgroup of a compact group G . Then H is nilpotent of class at most n if and only if*

$$\mathbf{np}(H; x_1, \dots, x_{n+1}) = 1$$

for some $x_k \in G$, $1 \leq k \leq n+1$.

Proof. Only one direction needs proof.

Using induction on n , we prove the result. For $n = 1$, assume that $\mathbf{np}(H; x, y) = 1$ for some $x, y \in G$, so by Proposition 2.5, $\mathbf{cp}(H) = 1$, whence implies that H is an abelian group. Assuming that the result is valid for $n - 1$, we prove its correctness for n . If

$$\mathbf{np}(H; x_1, \dots, x_{n+1}) = 1$$

for some $x_k \in G$, $1 \leq k \leq n+1$, then by Proposition 2.7, we get $\mathbf{np} \left(\frac{H}{\mathbf{Z} \cap H}; \bar{x}_1, \dots, \bar{x}_n \right) = 1$. Now, by induction hypothesis, $\frac{H}{\mathbf{Z} \cap H}$ is nilpotent of class at most $n - 1$, so, H is nilpotent of class at most n . □

Definition 2.9. Let H be a closed subgroup of a compact group G . The *relative k -nilpotence of H in G* , denoted by $\mathbf{np}_{k,G}(H)$, defined to be

$$\sup \{ \mathbf{np}(H, x_1, \dots, x_{k+1}) : x_1, \dots, x_k \in G \}.$$

When $H = G$, then we remove G from the index and simply write $\mathbf{np}_k(G)$.

Remark 2.10. Since the Haar measure is left-invariant, one can easily verify that $\mathbf{np}_k(G) = \mathbf{np}(G; x_1, \dots, x_{k+1})$ for all $x_1, \dots, x_{k+1} \in G$. So, $\mathbf{np}_k(G)$ is nothing but the degree of k -step nilpotence of G defined in [11]. Also, by Lemma 2.5, we have $\mathbf{np}_{1,G}(H) = \mathbf{cp}(H)$ for a closed subgroup H of a compact group G .

The following result follows inductively from Proposition 2.8, 2.5 and [7, Theorem].

Theorem 2.11. *Let H be closed subgroup of a compact group G which is not nilpotent of class at most n . Then $\mathbf{np}_{k,G}(H) \leq 1 - \frac{3}{2^{k+1}}$, for all positive integer k .*

By Theorem 2.11 in hand, we give another (simple) proof for the following statement.

Theorem 2.12. ([11, Theorem 1.19.]) *Let G is a profinite group with $\mathbf{np}_k(G) > 0$. Then G has an open k -step nilpotent subgroup.*

Proof. By [2, Lemma 2.4.], there is an open normal subgroup N along with elements $x_1, \dots, x_{k+1} \in G$ such that $\mathbf{np}(N; x_1, \dots, x_{k+1}) > 1 - \frac{3}{2^{k+1}}$, so by Theorem 2.11, N should be nilpotent of class at most k . \square

If B is a measurable subgroup of a compact group A , then $\mathbf{m}_A(B) = [A : B]^{-1}$, by convention on " $\frac{1}{\infty} = 0$ ". We have

Lemma 2.13. *Let G be a compact group, H be a closed subgroup of G and N be a normal closed subgroup of G . Then*

$$\mathbf{m}_G(H) = \mathbf{m}_N(N \cap H) \mathbf{m}_{G/N}(NH/N).$$

Proof. The following is well known

$$[G : H] = [G : NH][NH : H] = [G/N : NH/N][N : N \cap H].$$

This along with the explanation before the lemma implies the result. \square

Theorem 2.14. *Let G be a compact group with closed subgroups H and closed normal subgroup N contained in H . Then $\mathbf{np}_{k,G}(H) \leq \mathbf{np}_{k,G/N}(H/N) \mathbf{np}_{k,G}(N)$.*

Proof. For $x_1, \dots, x_{k+1} \in G$ we have

$$\mathbf{np}(H; x_1, \dots, x_{k+1}) \leq \int_{H^k} \mathbf{m}_H(C_H([x_1 y_1, \dots, x_k y_k])) \, \mathbf{d}y.$$

By Lemmas 2.1 and 2.13,

$$\begin{aligned}
\int_{H^k} \mathbf{m}_H(C_H([x_1y_1, \dots, x_ky_k])) \, d\mathbf{y} &= \int_{H^k/N^k} \int_{N^k} \mathbf{m}_H(C_H([x_1y_1z_1, \dots, x_ky_kz_k])) \, d\mathbf{z} \, d\bar{\mathbf{y}} \\
&= \int_{H^k/N^k} \int_{N^k} \mathbf{m}_N(C_N([x_1y_1z_1, \dots, x_ky_kz_k])) \mathbf{m}_{G/N}(NC_H([x_1y_1z_1, \dots, x_ky_kz_k])/N) \, d\mathbf{z} \, d\bar{\mathbf{y}} \\
&\stackrel{(*)}{\leq} \int_{H^k/N^k} \int_{N^k} \mathbf{m}_N(C_N([x_1y_1z_1, \dots, x_ky_kz_k])) \mathbf{m}_{H/N}(C_{H/N}([\overline{x_1y_1}, \dots, \overline{x_ky_k}])) \, d\mathbf{z} \, d\bar{\mathbf{y}} \\
&= \int_{H^k/N^k} \mathbf{m}_{H/N}(C_{H/N}([\overline{x_1y_1}, \dots, \overline{x_ky_k}])) \left(\int_{N^k} \mathbf{m}_N(C_N([x_1y_1z_1, \dots, x_ky_kz_k])) \, d\mathbf{z} \right) d\bar{\mathbf{y}} \\
&= \int_{H^k/N^k} \mathbf{m}_{G/N}(C_{G/N}([\overline{x_1y_1}, \dots, \overline{x_ky_k}])) \mathbf{np}(N; x_1y_1, \dots, x_ky_k, 1) \, d\bar{\mathbf{y}} \\
&\leq \mathbf{np}_{k,G}(N) \int_{(G/N)^k} \mathbf{m}_{G/N}(C_{G/N}([\overline{x_1y_1}, \dots, \overline{x_ky_k}])) \, d\bar{\mathbf{y}} \\
&\leq \mathbf{np}_{k,G}(N) \mathbf{np}_{k,G/N}(H/N).
\end{aligned}$$

The inequality (*) obtained by the inclusion

$$NC_H([x_1y_1, \dots, x_ky_k])/N \subset C_{H/N}([\overline{x_1y_1}, \dots, \overline{x_ky_k}]).$$

□

As a corollary of the above theorem, we state:

Corollary 2.15. *Let N be a closed normal subgroup of a compact group G . Then $\mathbf{np}_k(G) \leq \mathbf{np}_k(G/N) \mathbf{np}_{k,G}(N)$. In particular, $\mathbf{cp}(G) \leq \mathbf{cp}(G/N) \mathbf{cp}(N)$.*

3. PROOFS OF MAIN RESULTS

Using Corollaries 2.15 and 2.11, we prove the following.

Theorem 3.1. *Let G be a compact group and assume that $\mathbf{np}_k(G) > 0$. Then the length of all closed normal series of G which its factors are not nilpotent of class at most k should be bounded.*

Proof. Let

$$1 = G_{r+1} \leq G_r \leq \dots \leq G_1 \leq G_0 = G$$

be a closed normal series such that G_i/G_{i+1} , $0 \leq i \leq r$, is not nilpotent of class at most k . Using the Theorem 2.14 repeatedly, we have

$$\begin{aligned}
\mathbf{np}_k(G) &\leq \mathbf{np}_{k,G_0}(G_r) \mathbf{np}_k(G_0/G_r) \\
&\leq \mathbf{np}_{k,G_0}(G_r) \mathbf{np}_{k,G_0/G_r}(G_{r-1}/G_r) \mathbf{np}_k(G_0/G_{r-1}) \\
&\vdots \\
&\leq \prod_{i=2}^{r+1} \mathbf{np}_{k,G_0/G_i}(G_{i-1}/G_i).
\end{aligned}$$

The Corollary 2.11 implies that the right hand side of the above inequality is at most $(1 - \frac{3}{2^{k+1}})^r$. All in all, we get

$$r < \frac{\ln \mathbf{np}_k(G)}{\ln(1 - \frac{3}{2^{k+1}})}.$$

□

Theorem 3.2. *Let G be a compact group with $\mathbf{np}_k(G) > 0$. Assume that there exists a decreasing net (H_α) of closed normal subgroups of G such that $\bigcap_\alpha H_\alpha = 1$. Then H_α is nilpotent of class at most k for sufficiently large α .*

Proof. If none of H_α 's are nilpotent of class at most k , then we can construct an infinite closed normal series of G which its factors are not nilpotent of class at most k , violates Theorem 3.1; indeed, suppose we have constructed a closed normal series,

$$(3.1) \quad G = H_0 \geq H_{\alpha_1} \geq \cdots \geq H_{\alpha_n}$$

such that its factors are not nilpotent of class at most k . Since H_{α_n} is not k -step nilpotent, there exists (x_1, \dots, x_{k+1}) in $H_{\alpha_n}^{k+1}$ such that $[x_1, \dots, x_{k+1}] \neq 1$. By the assumption, we can choose $H_{\alpha_{n+1}}$ such that $[x_1, \dots, x_{k+1}] \notin H_{\alpha_{n+1}}$, so $H_{\alpha_{n+1}}$ can be added to (3.2). □

Since $\bigcap \{N : N \text{ is an open normal subgroup of } G\} = 1$, for a profinite group G , Theorem 3.2 gives another proof for the following well-known fact:

Corollary 3.3. ([11, Theorem 1.19.]) *Let G be a profinite group with $\mathbf{np}_k(G) > 0$. Then G has a k -step nilpotent open subgroup.*

Theorem 3.1 has also another implication:

Corollary 3.4. *Let $(G_\alpha)_\alpha$ be a collection of compact groups. If*

$$\mathbf{np}_k \left(\prod_\alpha G_\alpha \right) > 0,$$

then G_α 's are nilpotent of class at most k for all but finitely many α .

Now we're ready to show that the class \mathbf{PA}_k contains connected compact groups. A little preparation is needed before proving the later.

Lemma 3.5. *Let G be a compact group. Then G contains no nontrivial closed semi simple connected compact Lie subgroup H with $\mathbf{np}_{k,G}(H) > 0$.*

Proof. Suppose $\mathcal{N} = \mathcal{N}(H; x_1, \dots, x_{k+1})$ has positive measure for some $x_1, \dots, x_{k+1} \in G$. Since \mathcal{N} is Zariski-closed and has positive measure, it should be equal to G , so H should be nilpotent by Proposition 2.8 and hence trivial by semi simplicity. \square

Theorem 3.6. *Let G be a connected compact group with $\mathbf{np}_k(G) > 0$. Then G is an abelian group.*

Proof. Note first that simply connected compact groups are semi simple ([9, Theorem 9.29.]). Also, Corollary 2.15 implies that $\mathbf{np}_k(G/\mathbf{Z}) > 0$, where \mathbf{Z} stands for the center of G . Therefore by the proof of [9, Theorem 9.24] and Corollary 3.4, there is a finite family $\{L_1, \dots, L_n\}$ of simple simply connected compact Lie groups such that $\frac{G}{\mathbf{Z}} \cong \prod_i L_i$, whence implies $\prod_i \mathbf{np}_k(L_i) > 0$. Now the result follows from Lemma 3.5. \square

Also compact Lie groups belong to \mathbf{PA}_k . Next we use G^0 to denote the connected component of a compact group G .

Lemma 3.7. *Let G be a compact Lie group with $\mathbf{np}_k(G) > 0$. Then G^0 is torus. In particular G has an open k -step nilpotent subgroup.*

Proof. Denote by \mathbf{Z}_0 the center of G^0 . By Corollary 2.15 and Theorem 2.14, we obtain

$$0 < \mathbf{np}_k(G) \leq \mathbf{np}_{k,G}(G^0) \leq \mathbf{np}_{k,G/\mathbf{Z}_0}(G^0/\mathbf{Z}_0).$$

Now the result obtained with Corollary 3.5, because G^0/\mathbf{Z}_0 is a semi simple connected compact Lie subgroup of G/\mathbf{Z}_0 . The last claim of Lemma follows from the fact that G^0 is a closed subgroup of G of finite index ([9, Theorem 6.36.]), so it should be open. \square

Remark 3.8. The above lemma is attributed to Yves de Cornulier ([16]), which is given in response to a question posed by the second author ([15]). Of course, the above proof is somewhat different from the proof presented in [16].

The main result of the Lemma 3.7 is true for arbitrary compact groups. Denote by $\mathfrak{N}(G)$ the set of all closed normal subgroups N of G such that G/N is a Lie group.

Proof of Theorem 1.2. By Corollary 2.15, $\mathbf{np}_k(G/N) > 0$ for all $N \in \mathfrak{N}(G)$. Lemma 3.7, now implies that $(G/N)^0 = G^0N/N$ ([9, Lemma 9.18.]) is an open abelian group and so, G^0N becomes open and normal in G . If none of G^0N 's is k -step nilpotent, $N \in \mathfrak{N}(G)$, then we can construct a closed normal series of G whose factors are not nilpotent

of class at most k , violates Theorem 3.1; indeed, suppose we have constructed a closed normal series,

$$(3.2) \quad G \geq G^0 N_1 \geq \cdots \geq G^0 N_s$$

such that its factors are not nilpotent of class at most k and $N_1, \dots, N_s \in \mathfrak{N}(G)$. Since $G^0 N_s$ is not k -step nilpotent, there exists (x_1, \dots, x_{k+1}) in $(G^0 N_s)^{k+1}$ such that $[x_1, \dots, x_{k+1}] \neq 1$. But, $[x_1, \dots, x_{k+1}]$ is equal 1 modulus N_s , for $G^0 N_s / N_s$ is abelian. Therefore $[x_1, \dots, x_{k+1}] \in N_s$, and by the fact that $\bigcap \{N : N \in \mathfrak{N}(G)\} = 1$ we obtain N_{s+1} in $\mathfrak{N}(G)$, which does not contain $[x_1, \dots, x_{k+1}]$. \square

The following question is proposed in [1] in order to resolve Question 1.1.

Question 3.9 (Question 3.2 of [1]). Are there finitely many words $w_{i,j}$ ($1 \leq i \leq n$, $1 \leq j \leq k+1$) in the free group on $k+1$ generators such that if G is a compact group, (x_1, \dots, x_{k+1}) , $\mathbf{u} = (u_1, \dots, u_{k+1}) \in G^{k+1}$ and $[x_1 w_{i,1}(\mathbf{u}), \dots, x_{k+1} w_{i,k+1}(\mathbf{u})] = 1$ for all $i \in \{1, \dots, n\}$ then $[u_1, \dots, u_{k+1}] = 1$?

To settle Question 1.1, we will prove that using Theorem 1.2, we may study the following slightly simpler question on finite groups.

Question 3.10. Are there finitely many words $w_{i,j}$ ($1 \leq i \leq n$, $1 \leq j \leq k+1$) in the free group on $k+1$ generators such that if G is a finite group, (x_1, \dots, x_{k+1}) , $\mathbf{u} = (u_1, \dots, u_{k+1}) \in G^{k+1}$, $\langle u_1, \dots, u_{k+1} \rangle$ is abelian-by-nilpotent and $[x_1 w_{i,1}(\mathbf{u}), \dots, x_{k+1} w_{i,k+1}(\mathbf{u})] = 1$ for all $i \in \{1, \dots, n\}$ then $[u_1, \dots, u_{k+1}] = 1$?

Theorem 3.11. *If the answer of Question 3.10 is positive, the answer of Question 1.1 is also affirmative.*

Proof. First note that by Theorem 1.2 together with Hall Theorem, [14, 15.4.1], all finitely generated subgroups of G are residually finite.

Let n be the positive integer in Question 3.10. By [1, Theorem 3.1] there exists an open subset V of G containing 1 such that

$$(3.3) \quad \mathbf{m} \left(\bigcap_{i=1}^n \mathcal{N}_k(G) \mathbf{y}_i \right) > 0,$$

where $\mathbf{y}_i \in V^{k+1}$, $i = 1, \dots, n$. By continuity of word maps, we can find an open neighborhood U of 1, such that $w_{i,j}(\mathbf{u}) \in V$ for

$$\mathbf{u} = (u_1, \dots, u_{k+1}) \in U^{k+1},$$

and $1 \leq i \leq n$, $1 \leq j \leq k+1$. Now, for $\mathbf{u} = (u_1, \dots, u_{k+1}) \in U^{k+1}$, (3.3) ensures the existence of $x_1, \dots, x_{k+1} \in G$, such that

$$(3.4) \quad [x_1 w_{i,1}(\mathbf{u}), \dots, x_{k+1} w_{i,k+1}(\mathbf{u})] = 1, \quad (1 \leq i \leq n).$$

But $\langle x_i, u_i : i = 1, \dots, k+1 \rangle$ is residually finite, so by passing to its finite quotients and the correctness of the Question 3.10, we get from (3.4) that $[u_1, \dots, u_{k+1}] = 1$, whence the open subgroup $\langle U \rangle$ is a k -step open subgroup of G . \square

REFERENCES

- [1] A. Abdollahi, M. Soleimani Malekan, Compact groups with a set of positive Haar measure satisfying a nilpotent law, *Mathematical Proceedings of the Cambridge Philosophical Society* <https://doi.org/10.1017/S0305004121000542>
- [2] A. Abdollahi, M. Soleimani Malekan, Profinite groups with many elements of bounded order, to appear *Advances in Group Theory and Applications*. <https://arxiv.org/abs/2012.13886>.
- [3] A. Erfanian, R. Rezaei and P. Lescot, On the relative commutativity degree of a subgroup of a finite group, *Comm. Algebra*, 35, No. 12, (2007) 4183-4197.
- [4] G. B. Folland, *A course in abstract harmonic analysis*, *Studies in Advanced Mathematics*, CRC Press, Boca Raton, FL, 1995.
- [5] P. X. Gallagher, The number of conjugacy classes in a finite group, *Math. Z.* 118, (1970) 175-179.
- [6] R. M. Guralnick and G. R. Robinson, On the commuting probability in finite groups, *J. Algebra*, 300 (2006) 509-528.
- [7] W. H. Gustafson, What is the probability that two group elements commute? *Amer. Math. Monthly* 80 (1973), 1031-1034.
- [8] K. H. Hofmann and F. G. Russo, The probability that x and y commute in a compact group, *Math. Proc. Camb. Phil. Soc.* 153 (2012) 557-571.
- [9] Karl H. Hofmann and Sidney A. Morris. *The Structure of Compact Groups: A Primer for the Student – A Handbook for the Expert*, Berlin, Boston: De Gruyter, 2020. <https://doi.org/10.1515/9783110695991>
- [10] L. Lévai and L. Pyber, Profinite groups with many commuting pairs or involutions, *Arch. Math. (Basel)* 75 (2000) 1-7.
- [11] A. Martino, M. C. H. Tointon, M. Valiunas and E. Ventura, Probabilistic nilpotence in infinite groups, *Isr. J. Math.* (2021). <https://doi.org/10.1007/s11856-021-2168-3>
- [12] M. R. M. Moghaddam, A. R. Salemkar and K. Chiti, n -isoclinism classes and n -nilpotency degree of finite groups, *Algebra Colloquium* 12 (2005), 255-261.
- [13] H. Nagao, On a conjecture of Brauer for p -solvable groups, *Journal of Mathematics*, Osaka City University, Vol. 13, No. I. (1962) 35-38.
- [14] D. J. S. Robinson, *A course the theory of groups*, Second edition, *Graduate Texts in Mathematics*, 80. Springer-Verlag, New York, 1996.
- [15] Meisam Soleimani Malekan ([https://mathoverflow.net/users/84700/meisamsoleimani malekan](https://mathoverflow.net/users/84700/meisamsoleimani%20malekan)). Can the degree of k -nilpotence of a simple simply connected compact Lie group be in $(0, 1)$? *MathOverflow*. <https://mathoverflow.net/q/404459> (version: 2021-09-21).

- [16] YCor (<https://mathoverflow.net/users/14094/ycor>). Can the degree of k -nilpotence of a simple simply connected compact Lie group be in $(0, 1)$? <https://mathoverflow.net/q/404514> (version: 2021-09-22).

DEPARTMENT OF PURE MATHEMATICS, FACULTY OF MATHEMATICS AND STATISTICS, UNIVERSITY OF ISFAHAN, ISFAHAN 81746-73441, IRAN.

Email address: `a.abdollahi@math.ui.ac.ir`

SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES, TEHRAN, IRAN.

Email address: `msmalekan@gmail.com`