

POLYADIZATION OF ALGEBRAIC STRUCTURES

STEVEN DUPLIJ

*Center for Information Technology (WWU IT), Universität Münster, Röntgenstrasse 7-13
D-48149 Münster, Deutschland*

ABSTRACT. A generalization of the semisimplicity concept for polyadic algebraic structures is proposed. If semisimple structures can be presented in the block-diagonal matrix form (Wedderburn decomposition), a general form of polyadic structures is given by block-shift matrices. We combine these forms in a special way to get a general shape of semisimple nonderived polyadic structures.

We then introduce the polyadization concept (a “polyadic constructor”) according to which one can construct a nonderived polyadic algebraic structure of any arity from a given binary structure. The polyadization of supersymmetric structures is also discussed. The “deformation” by shifts of operations on the direct power of binary structures is defined and used to obtain a nonderived polyadic multiplication. Illustrative concrete examples for the new constructions are given.

CONTENTS

1. INTRODUCTION	2
2. PRELIMINARIES	2
3. POLYADIC SEMISIMPLICITY	3
3.1. Simple polyadic structures	3
3.2. Semisimple polyadic structures	4
4. POLYADIZATION CONCEPT	6
4.1. Polyadization of binary algebraic structures	7
4.2. Concrete examples of the polyadization procedure	9
4.2.1. Polyadization of $GL(2, \mathbb{C})$	9
4.2.2. Polyadization of $SO(2, \mathbb{R})$	11
4.3. “Deformation” of binary operations by shifts	12
4.4. Polyadization of binary supergroups	15
4.4.1. Polyadization of $GL(1 1, \Lambda)$	15
REFERENCES	17

E-mail address: douplii@uni-muenster.de; sduplij@gmail.com; <https://ivv5hpp.uni-muenster.de/u/douplii>.

Date: of start May 14, 2022. *Date: of completion* July 29, 2022.

Total: 32 references.

2010 Mathematics Subject Classification. 16T25, 17A42, 20B30, 20F36, 20M17, 20N15.

Key words and phrases. direct product, direct power, polyadic semigroup, arity, polyadic ring, polyadic field.

1. INTRODUCTION

I am no poet, but if you think for yourselves, as I proceed,
the facts will form a poem in your minds.

“*The Life and Letters of Faraday*” (1870) by Bence Jones
MICHAEL FARADAY

The concept of simple and semisimple rings, modules, and algebras (see, e.g., [ERDMANN AND HOLM \[2018\]](#), [HUNGERFORD \[1974\]](#), [LAMBEK \[1966\]](#), [ROTMAN \[2010\]](#)) plays a crucial role in the investigation of Lie algebras and representation theory [CURTIS AND REINER \[1962\]](#), [FULTON AND HARRIS \[2004\]](#), [KNAPP \[1986\]](#), as well as in category theory [HARADA \[1970\]](#), [KNOP \[2006\]](#), [SIMSON \[1977\]](#).

Here we first propose a generalization of this concept for polyadic algebraic structures [DUPLIJ \[2022a\]](#). If semisimple structures can be presented in the block-diagonal matrix form (resulting to the Wedderburn decomposition [WEDDERBURN \[1908\]](#), [HERSTEIN \[1996\]](#), [LAM \[1991\]](#)), a corresponding general form for polyadic rings can be decomposed to a kind of block-shift matrices [NIKITIN \[1984\]](#). We combine these forms and introduce a general shape of semisimple polyadic structures, which are nonderived in the sense that they cannot be obtained as a successive composition of binary operations, which can be treated as a polyadic (“double”) Wedderburn decomposition.

Second, going in the opposite direction, we define the polyadization concept (“polyadic constructor”) according to which one can construct a nonderived polyadic algebraic structure of any arity from a given binary structure. Then we briefly describe supersymmetric structure polyadization.

Third, we propose operations “deformed” by shifts to obtain a nonderived n -ary multiplication on the direct power of binary algebraic structures.

For these new constructions some illustrative concrete examples are given.

2. PRELIMINARIES

We use notation from [DUPLIJ \[2022a,b\]](#). In brief, a (one-set) *polyadic algebraic structure* \mathcal{A} is a set A closed with respect to polyadic operations (or n -ary multiplication) $\mu^{[n]} : A^n \rightarrow A$ (n -ary magma). We denote *polyads* [POST \[1940\]](#) by bold letters $\mathbf{a} = \mathbf{a}^{(n)} = (a_1, \dots, a_n)$, $a_i \in A$. A *polyadic zero* is defined by $\mu^{[n]}[\mathbf{a}^{(n-1)}, z] = z$, $z \in A$, $\mathbf{a}^{(n-1)} \in A^{n-1}$, where z can be on any place. A (positive) *polyadic power* $\ell_\mu \in \mathbb{N}$ is $a^{\langle \ell_\mu \rangle} = (\mu^{[n]})^{\circ \ell_\mu} [a^{\ell_\mu(n-1)+1}]$, $a \in A$. An element of a polyadic algebraic structure a is called ℓ_μ -*nilpotent* (or simply *nilpotent* for $\ell_\mu = 1$), if there exist ℓ_μ such that $a^{\langle \ell_\mu \rangle} = z$. A *polyadic (or n -ary) identity* (or neutral element) is defined by $\mu^{[n]}[a, e^{n-1}] = a$, $\forall a \in A$, where a can be on any place in the l.h.s. A one-set polyadic algebraic structure $\langle A \mid \mu^{[n]} \rangle$ is *totally associative*, if $(\mu^{[n]})^{\circ 2} [a, b, c] = \mu^{[n]} [a, \mu^{[n]} [b], c] = \text{invariant}$, with respect to placement of the internal multiplication on any of the n places, and a, b, c are polyads of the necessary sizes [DUPLIJ \[2018, 2019\]](#). A *polyadic semigroup* $\mathcal{S}^{(n)}$ is a one-set and one-operation structure in which $\mu^{[n]}$ is totally associative. A polyadic structure is *commutative*, if $\mu^{[n]} = \mu^{[n]} \circ \sigma$, or $\mu^{[n]} [a] = \mu^{[n]} [\sigma \circ a]$, $a \in A^n$, for all $\sigma \in S_n$.

A polyadic structure is *solvable*, if for all polyads \mathbf{b} , \mathbf{c} and an element x , one can (uniquely) resolve the equation (with respect to h) for $\mu^{[n]}[\mathbf{b}, x, \mathbf{c}] = a$, where x can be on any place, and \mathbf{b} , \mathbf{c} are polyads of the needed lengths. A solvable polyadic structure is called a *polyadic quasigroup* [BELOUSOV \[1972\]](#). An associative polyadic quasigroup is called a *n-ary (or polyadic) group* [GAL'MAK \[2003\]](#). In an *n-ary* group the only solution of

$$\mu^{[n]}[\mathbf{b}, \bar{a}] = a, \quad a, \bar{a} \in A, \quad \mathbf{b} \in A^{n-1} \quad (2.1)$$

is called a *querement* of a and denoted by \bar{a} [DÖRNTE \[1929\]](#), where \bar{a} can be on any place. Any idempotent a coincides with its querement $\bar{a} = a$. The relation (2.1) can be considered as a definition of the unary *querooperation* $\bar{\mu}^{(1)}[a] = \bar{a}$ [GLEICHGEWICHT AND GLĄZEK \[1967\]](#). For further details and definitions, see [DUPLIJ \[2022a\]](#).

3. POLYADIC SEMISIMPLICITY

In general, simple algebraic structures are building blocks (direct summands) for the semisimple ones satisfying special conditions (see, e.g., [ERDMANN AND HOLM \[2018\]](#), [LAMBEC \[1966\]](#)).

3.1. Simple polyadic structures. According to the Wedderburn-Artin theorem (see, e.g., [HERSTEIN \[1996\]](#), [LAM \[1991\]](#), [HAZEWINKEL AND GUBARENI \[2016\]](#)), a ring which is simple (having no two-sided ideals, except zero and the ring itself) and Artinian (having minimal right ideals) \mathcal{R}_{simple} is isomorphic to a full $d \times d$ matrix ring

$$\mathcal{R}_{simple} \cong Mat_{d \times d}^{full}(\mathcal{D}) \quad (3.1)$$

over a division ring \mathcal{D} . As a corollary,

$$\mathcal{R}_{simple} \cong \text{Hom}_{\mathcal{D}}(V(d | \mathcal{D}), V(d | \mathcal{D})) \equiv \text{End}_{\mathcal{D}}(V(d | \mathcal{D})), \quad (3.2)$$

where $V(d | \mathcal{D})$ is a d -finite-dimensional vector space (left module) over \mathcal{D} . In the same way, a finite-dimensional simple associative algebra \mathcal{A} over an algebraically closed field \mathcal{F} is

$$\mathcal{A} \cong Mat_{d \times d}^{full}(\mathcal{F}). \quad (3.3)$$

In the polyadic case, the structure of a simple Artinian $[2, n]$ -ring $\mathcal{R}_{simple}^{[2, n]}$ (with binary addition and n -ary multiplication $\mu^{[n]}$) was obtained in [NIKITIN \[1984\]](#), where the Wedderburn-Artin theorem for $[2, n]$ -rings was proved. So instead of one vector space $V(d | \mathcal{D})$, one should consider a direct sum of $(n - 1)$ vector spaces (over the same division ring \mathcal{D}), that is

$$V_1(d_1 | \mathcal{D}) \oplus V_2(d_2 | \mathcal{D}) \oplus \dots \oplus V_{n-1}(d_{n-1} | \mathcal{D}), \quad (3.4)$$

where $V_i(d_i | \mathcal{D})$ is a d_i -dimensional polyadic vector space [DUPLIJ \[2019\]](#), $i = 1, \dots, n - 1$. Then, instead of (3.2) we have the cyclic direct sum of homomorphisms

$$\begin{aligned} \mathcal{R}_{simple}^{[2, n]} \cong & \text{Hom}_{\mathcal{D}}(V_1(d_1 | \mathcal{D}), V_2(d_2 | \mathcal{D})) \oplus \text{Hom}_{\mathcal{D}}(V_2(d_2 | \mathcal{D}), V_3(d_3 | \mathcal{D})) \oplus \dots \\ & \dots \oplus \text{Hom}_{\mathcal{D}}(V_{n-1}(d_{n-1} | \mathcal{D}), V_1(d_1 | \mathcal{D})). \end{aligned} \quad (3.5)$$

This means that after choosing a suitable basis in terms of matrices (when the ring multiplication $\mu^{[n]}$ coincides with the product of n matrices) we have

Theorem 3.1. *The simple polyadic ring $\mathcal{R}_{simple}^{[2, n]}$ is isomorphic to the $d \times d$ matrix ring (cf. (3.1))*

$$\mathcal{R}_{simple}^{[2, n]} \cong Mat_{d \times d}^{Bshift(n)}(\mathcal{D}) = \{M^{Bshift(n)}(d \times d)\}, \quad (3.6)$$

where $M_{d \times d}^{Bshift}$ is the block-shift (traceless) matrix over \mathcal{D} of the form (which follows from (3.5))

$$M^{Bshift(n)}(d \times d) = \begin{pmatrix} 0 & B_1(d_1 \times d_2) & \dots & 0 & 0 \\ 0 & 0 & B_2(d_2 \times d_3) & \dots & 0 \\ 0 & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 & B_{n-2}(d_{n-2} \times d_{n-1}) \\ B_{n-1}(d_{n-1} \times d_1) & 0 & \dots & 0 & 0 \end{pmatrix}, \quad (3.7)$$

where $(n-1)$ blocks are full matrix rings $B_i(d' \times d'') \in Mat_{d' \times d''}^{full}(\mathcal{D})$, and $d = d_1 + d_2 + \dots + d_{n-1}$.

Note that the set of the fixed size blocks $\{B_i(d' \times d'')\}$ does not form a ring, because $d' \neq d''$, in general.

Assertion 3.2. *The block-shift matrices of the form (3.7) are closed with respect to n -ary multiplication and binary addition, and we call them n -ary matrices.*

Taking distributivity into account we arrive at the polyadic ring structure (3.6).

Corollary 3.3. In the limiting case $n = 2$, we have

$$M^{Bshift(2)}(d \times d) = B_1(d_1 \times d_1) \quad (3.8)$$

and $d = d_1$, giving a binary ring (3.1).

Assertion 3.4. *A finite-dimensional simple associative n -ary algebra $\mathcal{A}^{(n)}$ over an algebraically closed field \mathcal{F} CARLSSON [1980] is isomorphic to the block-shift n -ary matrix (3.7) over \mathcal{F}*

$$\mathcal{A}^{(n)} \cong Mat_{d \times d}^{Bshift(n)}(\mathcal{F}). \quad (3.9)$$

3.2. Semisimple polyadic structures. The Wedderburn-Artin theorem for semisimple Artinian rings \mathcal{R}_{semi} states that \mathcal{R}_{semi} is a finite direct sum of k simple rings, each of which has the form (3.1). Using (3.2) for every component, we decompose the d -finite-dimensional vector space (left module) into a direct sum of length k

$$V(d) = W^{(1)}(q^{(1)} | \mathcal{D}^{(1)}) \oplus W^{(2)}(q^{(2)} | \mathcal{D}^{(2)}) \oplus \dots \oplus W^{(k)}(q^{(k)} | \mathcal{D}^{(k)}), \quad (3.10)$$

where $d = q^{(1)} + q^{(2)} + \dots + q^{(k)}$. Then, instead of (3.2) we have the isomorphism¹

$$\mathcal{R}_{semi} \cong \text{End}_{\mathcal{D}^{(1)}} W^{(1)}(q^{(1)} | \mathcal{D}^{(1)}) \oplus \text{End}_{\mathcal{D}^{(2)}} W^{(2)}(q^{(2)} | \mathcal{D}^{(2)}) \oplus \dots \oplus \text{End}_{\mathcal{D}^{(k)}} W^{(k)}(q^{(k)} | \mathcal{D}^{(k)}). \quad (3.11)$$

In a suitable basis the Wedderburn-Artin theorem follows

Theorem 3.5. *A semisimple Artinian ring \mathcal{R}_{semi} is isomorphic to the $d \times d$ matrix ring*

$$\mathcal{R}_{semi} \cong Mat_{q^{(j)} \times q^{(j)}}^{Bdiag(k)} = \{M^{Bdiag(k)}(d \times d)\}, \quad (3.12)$$

where $M^{Bdiag(k)}(d \times d)$ are block-diagonal matrices of the form (which follows from (3.11))

$$M^{Bdiag(k)}(d \times d) = \begin{pmatrix} A^{(1)}(q^{(1)} \times q^{(1)}) & 0 & \dots & 0 \\ 0 & A^{(2)}(q^{(2)} \times q^{(2)}) & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \dots & A^{(k)}(q^{(k)} \times q^{(k)}) \end{pmatrix}, \quad (3.13)$$

¹We enumerate k simple components by an upper index in round brackets.

where k square blocks are full matrix rings over division rings $\mathcal{D}^{(j)}$

$$A^{(j)} \left(q^{(j)} \times q^{(j)} \right) \in \text{Mat}_{q^{(j)} \times q^{(j)}}^{\text{full}} \left(\mathcal{D}^{(j)} \right), \quad j = 1, \dots, k, \quad d = q^{(1)} + q^{(2)} + \dots + q^{(k)}. \quad (3.14)$$

The same matrix structure has a finite-dimensional semisimple associative algebra \mathcal{A} over an algebraically closed field \mathcal{F} (see (3.3)).

For further details, see, e.g., HERSTEIN [1996], LAM [1991], HAZEWINKEL AND GUBARENI [2016].

General properties of semisimple Artinian $[2, n]$ -rings were considered in NIKITIN [1984] (for ternary rings, see LISTER [1971], PROFERA [1982]). Here we propose a new manifest matrix structure for them.

Thus, our task is to decompose each of the $V_i(d_i)$, in (3.4) into components as in (3.10)

$$V_i(d_i) = W_i^{(1)} \left(q_i^{(1)} \mid \mathcal{D}^{(1)} \right) \oplus W_i^{(2)} \left(q_i^{(2)} \mid \mathcal{D}^{(2)} \right) \oplus \dots \oplus W_i^{(k)} \left(q_i^{(k)} \mid \mathcal{D}^{(k)} \right), \quad i = 1, \dots, n-1. \quad (3.15)$$

In matrix language this means that each block $B_{d' \times d''}$ from the polyadic ring (3.7) should have the semisimple decomposition (3.13), i.e. be a block-diagonal square matrix of the same size $p \times p$, where $p = d_1 = d_2 = \dots = d_{n-1}$ and the total matrix size becomes $d = (n-1)p$. Moreover, all the blocks B 's should have diagonal blocks A 's of the same size, and therefore $q^{(j)} \equiv q_1^{(j)} = q_2^{(j)} = \dots = q_{n-1}^{(j)}$ for all $j = 1, \dots, k$ and $p = q^{(1)} + q^{(2)} + \dots + q^{(k)}$, where k is the number of semisimple components. In this way the cyclic direct sum of homomorphisms for the semisimple polyadic rings becomes (we use different division rings for each semisimple component as in (3.14))

$$\begin{aligned} \mathcal{R}_{\text{semi}}^{[2,n]} &\cong \text{Hom}_{\mathcal{D}^{(1)}} \left(W_1^{(1)} \left(q^{(1)} \mid \mathcal{D}^{(1)} \right), W_2^{(1)} \left(q^{(1)} \mid \mathcal{D}^{(1)} \right) \right) \\ &\oplus \text{Hom}_{\mathcal{D}^{(2)}} \left(W_1^{(2)} \left(q^{(2)} \mid \mathcal{D}^{(2)} \right), W_2^{(2)} \left(q^{(2)} \mid \mathcal{D}^{(2)} \right) \right) \oplus \dots \\ &\dots \oplus \text{Hom}_{\mathcal{D}^{(k)}} \left(W_1^{(k)} \left(q^{(k)} \mid \mathcal{D}^{(k)} \right), W_2^{(k)} \left(q^{(k)} \mid \mathcal{D}^{(k)} \right) \right) \\ &\oplus \text{Hom}_{\mathcal{D}^{(1)}} \left(W_2^{(1)} \left(q^{(1)} \mid \mathcal{D}^{(1)} \right), W_3^{(1)} \left(q^{(1)} \mid \mathcal{D}^{(1)} \right) \right) \\ &\oplus \text{Hom}_{\mathcal{D}^{(2)}} \left(W_2^{(2)} \left(q^{(2)} \mid \mathcal{D}^{(2)} \right), W_3^{(2)} \left(q^{(2)} \mid \mathcal{D}^{(2)} \right) \right) \oplus \dots \\ &\dots \oplus \text{Hom}_{\mathcal{D}^{(k)}} \left(W_2^{(k)} \left(q^{(k)} \mid \mathcal{D}^{(k)} \right), W_3^{(k)} \left(q^{(k)} \mid \mathcal{D}^{(k)} \right) \right) \\ &\vdots \\ &\oplus \text{Hom}_{\mathcal{D}^{(1)}} \left(W_{n-2}^{(1)} \left(q^{(1)} \mid \mathcal{D}^{(1)} \right), W_{n-1}^{(1)} \left(q^{(1)} \mid \mathcal{D}^{(1)} \right) \right) \\ &\oplus \text{Hom}_{\mathcal{D}^{(2)}} \left(W_{n-2}^{(2)} \left(q^{(2)} \mid \mathcal{D}^{(2)} \right), W_{n-1}^{(2)} \left(q^{(2)} \mid \mathcal{D}^{(2)} \right) \right) \oplus \dots \\ &\dots \oplus \text{Hom}_{\mathcal{D}^{(k)}} \left(W_{n-2}^{(k)} \left(q^{(k)} \mid \mathcal{D}^{(k)} \right), W_{n-1}^{(k)} \left(q^{(k)} \mid \mathcal{D}^{(k)} \right) \right) \\ &\oplus \text{Hom}_{\mathcal{D}^{(1)}} \left(W_{n-1}^{(1)} \left(q^{(1)} \mid \mathcal{D}^{(1)} \right), W_1^{(1)} \left(q^{(1)} \mid \mathcal{D}^{(1)} \right) \right) \\ &\oplus \text{Hom}_{\mathcal{D}^{(2)}} \left(W_{n-1}^{(2)} \left(q^{(2)} \mid \mathcal{D}^{(2)} \right), W_1^{(2)} \left(q^{(2)} \mid \mathcal{D}^{(2)} \right) \right) \oplus \dots \\ &\dots \oplus \text{Hom}_{\mathcal{D}^{(k)}} \left(W_{n-1}^{(k)} \left(q^{(k)} \mid \mathcal{D}^{(k)} \right), W_1^{(k)} \left(q^{(k)} \mid \mathcal{D}^{(k)} \right) \right). \end{aligned} \quad (3.16)$$

After choosing a suitable basis we obtain a polyadic analog of the Wedderburn-Artin theorem for semisimple Artinian $[2, n]$ -rings $\mathcal{R}_{\text{semi}}^{[2,n]}$

Theorem 3.6. *The semisimple polyadic Artinian ring $\mathcal{R}_{semi}^{[2,n]}$ is isomorphic to the $d \times d$ matrix ring*

$$\mathcal{R}_{semi}^{[2,n]} \cong Mat_{d \times d}^{Bshiftdiag(n,k)} = \{N^{Bshiftdiag(n,k)}(d \times d)\}, \quad (3.17)$$

where $N_{d \times d}^{Bshiftdiag(n,k)}$ (n is the arity of the N 's and k is the number of simple components of the N 's) are the block-shift n -ary matrices with block-diagonal square blocks (which follows from (3.16))

$$N^{Bshiftdiag(n,k)}(d \times d) = \begin{pmatrix} 0 & B_1^{(k)}(p \times p) & \dots & 0 & 0 \\ 0 & 0 & B_2^{(k)}(p \times p) & \dots & 0 \\ 0 & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 & B_{n-2}^{(k)}(p \times p) \\ B_{n-1}(p \times p) & 0 & \dots & 0 & 0 \end{pmatrix}, \quad (3.18)$$

$$B_i^{(k)}(p \times p) = \begin{pmatrix} A_i^{(1)}(q^{(1)} \times q^{(1)}) & 0 & \dots & 0 \\ 0 & A_i^{(2)}(q^{(2)} \times q^{(2)}) & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 \\ 0 & 0 & \dots & A_i^{(k)}(q^{(k)} \times q^{(k)}) \end{pmatrix}, \quad (3.19)$$

where $d = (n-1)p$, $p = q^{(1)} + q^{(2)} + \dots + q^{(k)}$, and the k square blocks A 's are full matrix rings over the division rings $\mathcal{D}^{(j)}$

$$A_i^{(j)}(q^{(j)} \times q^{(j)}) \in Mat_{q^{(j)} \times q^{(j)}}^{full}(\mathcal{D}^{(j)}), \quad j = 1, \dots, k, \quad i = 1, \dots, n-1. \quad (3.20)$$

Let us generalize the above decompositions to superrings and superalgebras. For that we first assume that the constituent vector spaces (entering in (3.16)) are super vector spaces (\mathbb{Z}_2 -graded vector spaces) obeying the standard decomposition into even and odd parts

$$W_i^{(j)}(q^{(j)} | \mathcal{D}^{(j)}) = W_i^{(j)}(q_{even}^{(j)} | \mathcal{D}^{(j)})_{even} \oplus W_i^{(j)}(q_{odd}^{(j)} | \mathcal{D}^{(j)})_{odd}, \quad i = 1, \dots, n-1, \quad j = 1, \dots, k, \quad (3.21)$$

where $q_{even}^{(j)}$ and $q_{odd}^{(j)}$ are dimensions of the even and odd spaces, respectively, $q^{(j)} = q_{even}^{(j)} + q_{odd}^{(j)}$.

The parity of a homogeneous element of the vector space $v \in W_i^{(j)}(q^{(j)} | \mathcal{D}^{(j)})$ is defined by $|v| = \bar{0}$ (resp. $\bar{1}$), if $v \in W_i^{(j)}(q_{even}^{(j)} | \mathcal{D}^{(j)})_{even}$ (resp. $W_i^{(j)}(q_{odd}^{(j)} | \mathcal{D}^{(j)})_{odd}$), and $\bar{0}, \bar{1} \in \mathbb{Z}_2$. For details, see BEREZIN [1987], LEITES [1983]. In the graded case, the k square blocks A 's in (3.20) are full supermatrix rings of the size $(q_{even}^{(j)} | q_{odd}^{(j)}) \times (q_{even}^{(j)} | q_{odd}^{(j)})$, while the square B 's (3.19) are block-diagonal supermatrices, and the block-shift n -ary supermatrices have a nonstandard form (3.18).

We assume that in the super case the Wedderburn-Artin theorem for semisimple Artinian superrings is also valid, with the form of the decompositions (3.18)–(3.19) being the same, however now the blocks A 's and B 's are corresponding supermatrices.

4. POLYADIZATION CONCEPT

Here we propose a general procedure for how to construct new polyadic algebraic structures from binary (or lower arity) ones, using the “inverse” (informally) to the block-shift matrix decomposition (3.7). It can be considered as a polyadic analog of the inverse problem of the determination of an algebraic structure from the knowledge of its Wedderburn decomposition DIETZEL AND MITTAL [2021].

4.1. Polyadization of binary algebraic structures. Let a binary algebraic structure \mathcal{X} be represented by $p \times p$ matrices $B_{\mathbf{y}} \equiv B_{\mathbf{y}} (p \times p)$ over a ring \mathcal{R} (a linear representation), where \mathbf{y} is the set of $N_{\mathbf{y}}$ parameters corresponding to an element x of \mathcal{X} . Because the binary addition in \mathcal{R} transfers to the matrix addition without restrictions (as opposed to the polyadic case, see below), we will consider only the multiplicative part of the resulting polyadic matrix ring. In this way, we propose a special block-shift matrix method to obtain n -ary semigroups (n -ary groups) from the binary ones, but the former are not derived from the latter GAL'MAK [2003], DUPLIJ [2022a]. In general, this can lead to new algebraic structures that were not known before.

Definition 4.1. A (block-matrix) polyadization Φ_{pol} of a binary semigroup (or group) \mathcal{X} represented by square $p \times p$ matrices $B_{\mathbf{y}}$ is an n -ary semigroup (or an n -ary group) represented by the $d \times d$ block-shift matrices (over a ring \mathcal{R}) of the form (3.7) as follows

$$Q_{\mathbf{y}_1, \dots, \mathbf{y}_{n-1}} \equiv Q_{\mathbf{y}_1, \dots, \mathbf{y}_{n-1}}^{Bshift(n)} (d \times d) = \begin{pmatrix} 0 & B_{\mathbf{y}_1} & \dots & 0 & 0 \\ 0 & 0 & B_{\mathbf{y}_2} & \dots & 0 \\ 0 & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 & B_{\mathbf{y}_{n-2}} \\ B_{\mathbf{y}_{n-1}} & 0 & \dots & 0 & 0 \end{pmatrix}, \quad (4.1)$$

where $d = (n-1)p$, and the n -ary multiplication $\mu^{[[n]]}$ is given by the product of n matrices (4.1).

In terms of the block-matrices B's the multiplication

$$\mu^{[[n]]} \left[\overbrace{Q_{\mathbf{y}'_1, \dots, \mathbf{y}'_{n-1}}, Q_{\mathbf{y}''_1, \dots, \mathbf{y}''_{n-1}}, \dots, Q_{\mathbf{y}'''_1, \dots, \mathbf{y}'''_{n-1}}, Q_{\mathbf{y}''''_1, \dots, \mathbf{y}''''_{n-1}}}^n \right] = Q_{\mathbf{y}_1, \dots, \mathbf{y}_{n-1}} \quad (4.2)$$

has the cyclic product form (see DUPLIJ [2021])

$$\overbrace{B_{\mathbf{y}'_1} B_{\mathbf{y}''_2} \dots B_{\mathbf{y}'''_{n-1}} B_{\mathbf{y}''''_1}}^n = B_{\mathbf{y}_1}, \quad (4.3)$$

$$B_{\mathbf{y}'_2} B_{\mathbf{y}''_3} \dots B_{\mathbf{y}'''_1} B_{\mathbf{y}''''_2} = B_{\mathbf{y}_2}, \quad (4.4)$$

⋮

$$B_{\mathbf{y}'_{n-1}} B_{\mathbf{y}''_1} \dots B_{\mathbf{y}'''_{n-2}} B_{\mathbf{y}''''_{n-1}} = B_{\mathbf{y}_{n-1}}. \quad (4.5)$$

Remark 4.2. The number of parameters $N_{\mathbf{y}}$ describing an element $x \in \mathcal{X}$ increases to $(n-1)N_{\mathbf{y}}$, and the corresponding algebraic structure $\langle \{Q_{\mathbf{y}_1, \dots, \mathbf{y}_{n-1}}\} \mid \mu^{[[n]]} \rangle$ becomes n -ary, and so (4.1) can be treated as a new algebraic structure, which we denote by the same letter with the arities in double square brackets $\mathcal{X}^{[[n]]}$.

We now analyze some of the most general properties of the polyadization map Φ_{pol} which are independent of the concrete form of the block-matrices B's and over which algebraic structure (ring, field, etc...) they are defined. We then present some concrete examples.

Definition 4.3. A unique polyadization Φ_{Upol} is a polyadization where all sets of parameters coincide

$$\mathbf{y} = \mathbf{y}_1 = \mathbf{y}_2 \dots = \mathbf{y}_{n-1}. \quad (4.6)$$

Proposition 4.4. The unique polyadization is an n -ary-binary homomorphism.

Proof. In the case of (4.6) all $(n - 1)$ relations (4.3)–(4.5) coincide

$$\overbrace{B_{y'} B_{y''} \dots B_{y'''} B_{y''''}}^n = B_y, \quad (4.7)$$

which means that the ordinary (binary) product of n matrices B_y 's is mapped to the n -ary product of matrices Q_y 's (4.2)

$$\mu^{[[n]]} \left[\overbrace{Q_{y'}, Q_{y''}, \dots, Q_{y'''} Q_{y''''}}^n \right] = Q_y, \quad (4.8)$$

as it should be for an n -ary-binary homomorphism, but not for a homomorphism. \square

Assertion 4.5. *If matrices $B_y \equiv B_y$ ($p \times p$) contain the identity matrix E_p , then the n -ary identity $E_d^{(n)}$ in $\langle \{Q_y (d \times d)\} \mid \mu^{[[n]]} \rangle$, $d = (n - 1)p$ has the form*

$$E_d^{(n)} = \begin{pmatrix} 0 & E_p & \dots & 0 & 0 \\ 0 & 0 & E_p & \dots & 0 \\ 0 & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 & E_p \\ E_p & 0 & \dots & 0 & 0 \end{pmatrix}. \quad (4.9)$$

Proof. It follows from (4.1), (4.2) and (4.7). \square

In this case the unique polyadization maps the identity matrix to the n -ary identity $\Phi_{Upol} : E_p \rightarrow E_d^{(n)}$.

Assertion 4.6. *If the matrices B_y are invertible $B_y B_y^{-1} = B_y^{-1} B_y = E_p$, then each $Q_{y_1, \dots, y_{n-1}}$ has a querelement*

$$\overline{Q}_{y_1, \dots, y_{n-1}} = \begin{pmatrix} 0 & \overline{B}_{y_1} & \dots & 0 & 0 \\ 0 & 0 & \overline{B}_{y_2} & \dots & 0 \\ 0 & 0 & \ddots & \ddots & \vdots \\ \vdots & \vdots & \ddots & 0 & \overline{B}_{y_{n-2}} \\ \overline{B}_{y_{n-1}} & 0 & \dots & 0 & 0 \end{pmatrix}, \quad (4.10)$$

satisfying

$$\mu^{[[n]]} \left[\overbrace{Q_{y_1, \dots, y_{n-1}}, Q_{y_1, \dots, y_{n-1}}, \dots, Q_{y_1, \dots, y_{n-1}} \overline{Q}_{y_1, \dots, y_{n-1}}}^n \right] = Q_{y_1, \dots, y_{n-1}} \quad (4.11)$$

where $\overline{Q}_{y_1, \dots, y_{n-1}}$ can be on any places and

$$\overline{B}_{y_i} = B_{y_{i-1}}^{-1} B_{y_{i-2}}^{-1} \dots B_{y_2}^{-1} B_{y_1}^{-1} B_{y_{n-1}}^{-1} B_{y_{n-2}}^{-1} \dots B_{y_{i+2}}^{-1} B_{y_{i+1}}^{-1}. \quad (4.12)$$

Proof. This follows from (4.10)–(4.11) and (4.3)–(4.5), and then consequently applying $B_{y_i}^{-1}$ (with suitable indices) on both sides. \square

Let us suppose that on the set of matrices $\{B_y\}$ over a binary ring \mathcal{R} one can consider some analog of a multiplicative character $\chi : \{B_y\} \rightarrow \mathcal{R}$, being a (binary) homomorphism, such that

$$\chi(B_{y_1}) \chi(B_{y_2}) = \chi(B_{y_1} B_{y_2}). \quad (4.13)$$

For instance, in case $B \in GL(p, \mathbb{C})$, the determinant can be considered as a (binary) multiplicative character. Similarly, we can introduce

Definition 4.7. A polyadized multiplicative character $\chi : \{Q_{\mathbf{y}_1, \dots, \mathbf{y}_{n-1}}\} \rightarrow \mathcal{R}$ is proportional to a product of the binary multiplicative characters of the blocks $\chi(B_{\mathbf{y}_i})$

$$\chi(Q_{\mathbf{y}_1, \dots, \mathbf{y}_{n-1}}) = (-1)^n \chi(B_{\mathbf{y}_1}) \chi(B_{\mathbf{y}_2}) \cdots \chi(B_{\mathbf{y}_{n-1}}). \quad (4.14)$$

The normalization factor $(-1)^n$ in (4.14) is needed to be consistent with the case when \mathcal{R} is commutative, and the multiplicative characters are determinants. It can also be consistent in other cases.

Proposition 4.8. If the ring \mathcal{R} is commutative, then the polyadized multiplicative character χ is an n -ary-binary homomorphism.

Proof. It follows from (4.7)–(4.8), (4.14) and the commutativity of \mathcal{R} . \square

Proposition 4.9. If the ring \mathcal{R} is commutative and unital with the unit E_p , then the algebraic structure $\langle \{Q_{\mathbf{y}_1, \dots, \mathbf{y}_{n-1}}\} \mid \mu^{[[n]]} \rangle$ contains polyadic (n -ary) idempotents satisfying

$$B_{\mathbf{y}_1} B_{\mathbf{y}_2} \cdots B_{\mathbf{y}_{n-1}} = E_p. \quad (4.15)$$

Proof. It follows from (4.8) and (4.9). \square

4.2. Concrete examples of the polyadization procedure.

4.2.1. *Polyadization of $GL(2, \mathbb{C})$.* Consider the polyadization procedure for the general linear group $GL(2, \mathbb{C})$. We have for the 4-parameter block matrices $B_{\mathbf{y}_i} = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix} \in GL(2, \mathbb{C})$, $\mathbf{y}_i = (a_i, b_i, c_i, d_i) \in \mathbb{C} \times \mathbb{C} \times \mathbb{C} \times \mathbb{C}$, $i = 1, 2, 3$. Thus, the 12-parameter 4-ary group $GL^{[[4]]}(2, \mathbb{C}) = \langle \{Q_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3}\} \mid \mu^{[[4]]} \rangle$ is represented by the following 6×6 Q -matrices

$$Q_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3} = \begin{pmatrix} 0 & B_{\mathbf{y}_1} & 0 \\ 0 & 0 & B_{\mathbf{y}_2} \\ B_{\mathbf{y}_3} & 0 & 0 \end{pmatrix} \in GL^{[[4]]}(2, \mathbb{C}), \quad B_{\mathbf{y}_i} \in GL(2, \mathbb{C}), \quad i = 1, 2, 3, \quad (4.16)$$

obeying the 4-ary multiplication

$$\mu^{[[4]]} [Q_{\mathbf{y}'_1, \mathbf{y}'_2, \mathbf{y}'_3}, Q_{\mathbf{y}''_1, \mathbf{y}''_2, \mathbf{y}''_3}, Q_{\mathbf{y}'''_1, \mathbf{y}'''_2, \mathbf{y}'''_3}, Q_{\mathbf{y}''''_1, \mathbf{y}''''_2, \mathbf{y}''''_3}] = Q_{\mathbf{y}'_1, \mathbf{y}'_2, \mathbf{y}'_3} Q_{\mathbf{y}''_1, \mathbf{y}''_2, \mathbf{y}''_3} Q_{\mathbf{y}'''_1, \mathbf{y}'''_2, \mathbf{y}'''_3} Q_{\mathbf{y}''''_1, \mathbf{y}''''_2, \mathbf{y}''''_3} = Q_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3}. \quad (4.17)$$

In terms of the block matrices $B_{\mathbf{y}_i}$ the multiplication (4.17) becomes (see (4.2)–(4.5))

$$B_{\mathbf{y}'_1} B_{\mathbf{y}''_2} B_{\mathbf{y}'''_3} B_{\mathbf{y}''''_1} = B_{\mathbf{y}_1}, \quad (4.18)$$

$$B_{\mathbf{y}'_2} B_{\mathbf{y}''_3} B_{\mathbf{y}'''_1} B_{\mathbf{y}''''_2} = B_{\mathbf{y}_2}, \quad (4.19)$$

$$B_{\mathbf{y}'_3} B_{\mathbf{y}''_1} B_{\mathbf{y}'''_2} B_{\mathbf{y}''''_3} = B_{\mathbf{y}_3}, \quad (4.20)$$

which can be further expressed in the B -matrix entries (its manifest form is too cumbersome to give here).

For $\{Q_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3}\}$ to be a 4-ary group each Q -matrix should have the unique querelement determined by the equation (see (4.11))

$$Q_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3} Q_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3} Q_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3} \bar{Q}_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3} = Q_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3}, \quad (4.21)$$

which has the solution

$$\bar{Q}_{\mathbf{y}_1, \mathbf{y}_2, \mathbf{y}_3} = \begin{pmatrix} 0 & \bar{B}_{\mathbf{y}_1} & 0 \\ 0 & 0 & \bar{B}_{\mathbf{y}_2} \\ \bar{B}_{\mathbf{y}_3} & 0 & 0 \end{pmatrix}, \quad (4.22)$$

where (see (4.12))

$$\bar{B}_{\mathbf{y}_1} = B_{\mathbf{y}_3}^{-1} B_{\mathbf{y}_2}^{-1}, \quad \bar{B}_{\mathbf{y}_2} = B_{\mathbf{y}_1}^{-1} B_{\mathbf{y}_3}^{-1}, \quad \bar{B}_{\mathbf{y}_3} = B_{\mathbf{y}_2}^{-1} B_{\mathbf{y}_1}^{-1}. \quad (4.23)$$

In the manifest form the querelements of $GL^{[[4]]}(2, \mathbb{C})$ are (4.22), where

$$\overline{B}_{y_1} = \frac{1}{\Delta_3 \Delta_2} \begin{pmatrix} b_3 c_2 + d_3 d_2 & -b_3 a_2 - d_3 b_2 \\ -a_3 c_2 - c_3 d_2 & a_3 a_2 + c_3 b_2 \end{pmatrix} \quad (4.24)$$

$$\overline{B}_{y_2} = \frac{1}{\Delta_2 \Delta_3} \begin{pmatrix} b_1 c_3 + d_1 d_3 & -b_1 a_3 - d_1 b_3 \\ -a_1 c_3 - c_1 d_3 & a_1 a_3 + c_1 b_3 \end{pmatrix} \quad (4.25)$$

$$\overline{B}_{y_3} = \frac{1}{\Delta_2 \Delta_1} \begin{pmatrix} b_2 c_1 + d_2 d_1 & -b_2 a_1 - d_2 b_1 \\ -a_2 c_1 - c_2 d_1 & a_2 a_1 + c_2 b_1 \end{pmatrix}, \quad (4.26)$$

where $\Delta_i = a_i d_i - b_i c_i \neq 0$ are the (nonvanishing) determinants of B_{y_i} .

Definition 4.10. We call $GL^{[[4]]}(2, \mathbb{C})$ a polyadic (4-ary) general linear group.

If we take the binary multiplicative characters to be determinants $\chi(B_{y_i}) = \Delta_i \neq 0$, then the polyadized multiplicative character in $GL^{[[4]]}(2, \mathbb{C})$ becomes

$$\chi(Q_{y_1, y_2, y_3}) = \Delta_1 \Delta_2 \Delta_3, \quad (4.27)$$

which is a 4-ary-binary homomorphism, because (see (4.18)–(4.20))

$$\begin{aligned} & \chi(Q_{y'_1, y'_2, y'_3}) \chi(Q_{y''_1, y''_2, y''_3}) \chi(Q_{y'''_1, y'''_2, y'''_3}) \chi(Q_{y''''_1, y''''_2, y''''_3}) \\ &= (\Delta'_1 \Delta'_2 \Delta'_3) (\Delta''_1 \Delta''_2 \Delta''_3) (\Delta'''_1 \Delta'''_2 \Delta'''_3) (\Delta''''_1 \Delta''''_2 \Delta''''_3) \\ &= (\Delta'_1 \Delta'_2 \Delta'_3 \Delta''_1) (\Delta'_2 \Delta'_3 \Delta''_2 \Delta''_1) (\Delta'_3 \Delta'_1 \Delta''_3 \Delta''_2) \\ &= \chi(Q_{y'_1, y'_2, y'_3} Q_{y''_1, y''_2, y''_3} Q_{y'''_1, y'''_2, y'''_3} Q_{y''''_1, y''''_2, y''''_3}). \end{aligned} \quad (4.28)$$

The 4-ary identity $E_6^{(4)}$ of $GL^{[[4]]}(2, \mathbb{C})$ is unique and has the form (see (4.9))

$$E_6^{(4)} = \begin{pmatrix} 0 & E_2 & 0 \\ 0 & 0 & E_2 \\ E_2 & 0 & 0 \end{pmatrix}, \quad (4.29)$$

where E_2 is the identity of $GL(2, \mathbb{C})$. The 4-ary identity $E_6^{(4)}$ satisfies the 4-ary idempotence relation

$$E_6^{(4)} E_6^{(4)} E_6^{(4)} E_6^{(4)} = E_6^{(4)}. \quad (4.30)$$

In general, the 4-ary group $GL^{[[4]]}(2, \mathbb{C})$ contains an infinite number of 4-ary idempotents $Q_{y_1, y_2, y_3}^{idemp}$ defined by the system of equations

$$Q_{y_1, y_2, y_3}^{idemp} Q_{y_1, y_2, y_3}^{idemp} Q_{y_1, y_2, y_3}^{idemp} Q_{y_1, y_2, y_3}^{idemp} = Q_{y_1, y_2, y_3}^{idemp}, \quad (4.31)$$

which gives

$$B_{y_1}^{idemp} B_{y_2}^{idemp} B_{y_3}^{idemp} = E_2, \quad (4.32)$$

or manifestly

$$a_1 a_2 a_3 + a_1 b_2 c_3 + a_3 b_1 c_2 + b_1 c_3 d_2 = 1, \quad (4.33)$$

$$a_2 b_3 c_1 + b_2 c_1 d_3 + b_3 c_2 d_1 + d_1 d_2 d_3 = 1, \quad (4.34)$$

$$a_1 a_2 b_3 + a_1 b_2 d_3 + b_1 b_3 c_2 + b_1 d_2 d_3 = 0, \quad (4.35)$$

$$a_2 a_3 c_1 + a_3 c_2 d_1 + b_2 c_1 c_3 + c_3 d_1 d_2 = 0. \quad (4.36)$$

The infinite set of idempotents in $GL^{[[4]]}(2, \mathbb{C})$ is determined by $12 - 4 = 8$ complex parameters, because one block-matrix (with 4 complex parameters) can always be excluded using the equation (4.32).

Remark 4.11. The above example shows, how “far” polyadic groups can be formed from ordinary (binary) groups: the former can contain an infinite number of 4-ary idempotents determined by (4.33)–(4.36), in addition to the standard idempotent in any group, the 4-ary identity (4.29).

4.2.2. *Polyadization of $SO(2, \mathbb{R})$.* Here we provide a polyadization for the simplest subgroup of $GL(2, \mathbb{C})$, the special orthogonal group $SO(2, \mathbb{R})$. In the matrix form $SO(2, \mathbb{R})$ is represented by the one-parameter rotation matrix

$$B(\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \in SO(2, \mathbb{R}), \quad \alpha \in \mathbb{R} / 2\pi\mathbb{Z}, \quad (4.37)$$

satisfying the commutative multiplication

$$B(\alpha)B(\beta) = B(\alpha + \beta), \quad (4.38)$$

and the (binary) identity E_2 is $B(0)$. Therefore, the inverse element for $B(\alpha)$ is $B(-\alpha)$.

The 4-ary polyadization of $SO(2, \mathbb{R})$ is given by the 3-parameter 4-ary group of Q -matrices $SO^{[4]}(2, \mathbb{R}) = \langle \{Q(\alpha, \beta, \gamma)\} \mid \mu^{[4]}\rangle$, where (cf. (4.16))

$$Q(\alpha, \beta, \gamma) = \begin{pmatrix} 0 & B(\alpha) & 0 \\ 0 & 0 & B(\beta) \\ B(\gamma) & 0 & 0 \end{pmatrix} \quad (4.39)$$

$$= \begin{pmatrix} 0 & 0 & \cos \alpha & -\sin \alpha & 0 & 0 \\ 0 & 0 & \sin \alpha & \cos \alpha & 0 & 0 \\ 0 & 0 & 0 & 0 & \cos \beta & -\sin \beta \\ 0 & 0 & 0 & 0 & \sin \beta & \cos \beta \\ \cos \gamma & -\sin \gamma & 0 & 0 & 0 & 0 \\ \sin \gamma & \cos \gamma & 0 & 0 & 0 & 0 \end{pmatrix}, \quad \alpha, \beta, \gamma \in \mathbb{R} / 2\pi\mathbb{Z}, \quad (4.40)$$

and the 4-ary multiplication is

$$\begin{aligned} & \mu^{[4]} [Q(\alpha_1, \beta_1, \gamma_1), Q(\alpha_2, \beta_2, \gamma_2), Q(\alpha_3, \beta_3, \gamma_3), Q(\alpha_4, \beta_4, \gamma_4)] \\ &= Q(\alpha_1, \beta_1, \gamma_1) Q(\alpha_2, \beta_2, \gamma_2) Q(\alpha_3, \beta_3, \gamma_3) Q(\alpha_4, \beta_4, \gamma_4) \\ &= Q(\alpha_1 + \beta_2 + \gamma_3 + \alpha_4, \beta_1 + \gamma_2 + \alpha_3 + \beta_4, \gamma_1 + \alpha_2 + \beta_3 + \gamma_4) = Q(\alpha, \beta, \gamma), \end{aligned} \quad (4.41)$$

which is noncommutative, as opposed to the binary product of B -matrices (4.38).

The querelement $\overline{Q}(\alpha, \beta, \gamma)$ for a given $Q(\alpha, \beta, \gamma)$ is defined by the equation (see (4.21))

$$Q(\alpha, \beta, \gamma) Q(\alpha, \beta, \gamma) Q(\alpha, \beta, \gamma) \overline{Q}(\alpha, \beta, \gamma) = Q(\alpha, \beta, \gamma), \quad (4.42)$$

which has the solution

$$\overline{Q}(\alpha, \beta, \gamma) = Q(-\beta - \gamma, -\alpha - \gamma, -\alpha - \beta). \quad (4.43)$$

Definition 4.12. We call $SO^{[4]}(2, \mathbb{R})$ a polyadic (4-ary) special orthogonal group, and $Q(\alpha, \beta, \gamma)$ is called a polyadic (4-ary) rotation matrix.

Informally, the matrix $Q(\alpha, \beta, \gamma)$ represents the polyadic (4-ary) rotation. There are an infinite number of polyadic (4-ary) identities (neutral elements) $E(\alpha, \beta, \gamma)$ which are defined by

$$E(\alpha, \beta, \gamma) E(\alpha, \beta, \gamma) E(\alpha, \beta, \gamma) Q(\alpha, \beta, \gamma) = Q(\alpha, \beta, \gamma), \quad (4.44)$$

and the solution is

$$E(\alpha, \beta, \gamma) = Q(\alpha, \beta, \gamma), \quad \alpha + \beta + \gamma = 0. \quad (4.45)$$

It follows from (4.44) that $E(\alpha, \beta, \gamma)$ are 4-ary idempotents (see (4.30) and Remark 4.11).

The determinants of $B(\alpha)$ and $Q(\alpha, \beta, \gamma)$ are 1, and therefore the corresponding multiplicative characters and polyadized multiplicative characters (4.14) are also equal to 1.

Comparing with the successive products of four B -matrices (4.37)

$$B(\alpha_1) B(\alpha_2) B(\alpha_3) B(\alpha_4) = B(\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4), \quad (4.46)$$

we observe that 4-ary multiplication (4.41) gives a shifted sum of four angles.

More exactly, for the triple (α, β, γ) we introduce the circle (left) shift operator by

$$s\alpha = \beta, \quad s\beta = \gamma, \quad s\gamma = \alpha \quad (4.47)$$

with the property $s^3 = \text{id}$. Then the 4-ary multiplication (4.41) becomes

$$\begin{aligned} & \mu^{[4]} [Q(\alpha_1, \beta_1, \gamma_1), Q(\alpha_2, \beta_2, \gamma_2), Q(\alpha_3, \beta_3, \gamma_3), Q(\alpha_4, \beta_4, \gamma_4)] \\ &= Q(\alpha_1 + s\alpha_2 + s^2\alpha_3 + \alpha_4, \beta_1 + s\beta_2 + s^2\beta_3 + \beta_4, \gamma_1 + s\gamma_2 + s^2\gamma_3 + \gamma_4). \end{aligned} \quad (4.48)$$

The querelement has the form

$$\overline{Q}(\alpha, \beta, \gamma) = Q(-s\alpha - s^2\alpha, -s\beta - s^2\beta, -s\gamma - s^2\gamma). \quad (4.49)$$

The multiplication (4.48) can be (informally) expressed in terms of a new operation, the 4-ary "cyclic shift addition" defined on $\mathbb{R} \times \mathbb{R} \times \mathbb{R}$ by (see (4.41))

$$\begin{aligned} & \nu_s^{[4]} [(\alpha_1, \beta_1, \gamma_1), (\alpha_2, \beta_2, \gamma_2), (\alpha_3, \beta_3, \gamma_3), (\alpha_4, \beta_4, \gamma_4)] \\ &= (\alpha_1 + \beta_2 + \gamma_3 + \alpha_4, \beta_1 + \gamma_2 + \alpha_3 + \beta_4, \gamma_1 + \alpha_2 + \beta_3 + \gamma_4) \\ &= (\nu_s^{[4]} [\alpha_1, \alpha_2, \alpha_3, \alpha_4], \nu_s^{[4]} [\beta_1, \beta_2, \beta_3, \beta_4], \nu_s^{[4]} [\gamma_1, \gamma_2, \gamma_3, \gamma_4]), \end{aligned} \quad (4.50)$$

where $\nu_s^{[4]}$ is (informally)

$$\nu_s^{[4]} [\alpha_1, \alpha_2, \alpha_3, \alpha_4] = s^0\alpha_1 + s^1\alpha_2 + s^2\alpha_3 + s^3\alpha_4 = \alpha_1 + s\alpha_2 + s^2\alpha_3 + \alpha_4, \quad (4.51)$$

and $s^0 = \text{id}$. This can also be treated as some "deformation" of the repeated binary additions by shifts. It is seen that the 4-ary operation $\nu_s^{[4]}$ (4.50) is not derived and cannot be obtained by consequent binary operations on the triples (α, β, γ) as (4.46).

In terms of the 4-ary cyclic shift addition the 4-ary multiplication (4.48) becomes

$$\begin{aligned} & \mu^{[4]} [Q(\alpha_1, \beta_1, \gamma_1), Q(\alpha_2, \beta_2, \gamma_2), Q(\alpha_3, \beta_3, \gamma_3), Q(\alpha_4, \beta_4, \gamma_4)] \\ &= Q(\nu_s^{[4]} [(\alpha_1, \beta_1, \gamma_1), (\alpha_2, \beta_2, \gamma_2), (\alpha_3, \beta_3, \gamma_3), (\alpha_4, \beta_4, \gamma_4)]). \end{aligned} \quad (4.52)$$

The binary case corresponds to $s = \text{id}$, because in (4.37) we have only one angle α , as opposed to three angles in (4.47).

Thus, we conclude that just as the binary product of B -matrices corresponds to the ordinary angle addition (4.38), the 4-ary multiplication of polyadic rotation Q -matrices (4.39) corresponds to the 4-ary cyclic shift addition (4.51) through (4.52).

4.3. "Deformation" of binary operations by shifts. The concrete example from the previous subsection shows the strong connection (4.52) between the polyadization procedure and the shifted operations (4.51). Here we generalize it to an n -ary case for any semigroup.

Let $\mathcal{A} = \langle A \mid (+) \rangle$ be a binary semigroup, where A is its underlying set and $(+)$ is the binary operation (which can be noncommutative). The simplest way to construct an n -ary operation $\nu^{[n]} : A^n \rightarrow A$ is the consequent repetition of the binary operation (see (4.46))

$$\nu^{[n]} [\alpha_1, \alpha_2, \dots, \alpha_n] = \alpha_1 + \alpha_2 + \dots + \alpha_n, \quad (4.53)$$

where the n -ary multiplication $\nu^{[n]}$ (4.53) is called derived DÖRNTE [1929], ZUPNIK [1967].

To construct a nonderived operation, we now consider the (external) m th direct power \mathcal{A}^m of the semigroup \mathcal{A} by introducing m -tuples

$$\mathbf{a} \equiv \mathbf{a}^{(m)} = \left(\overbrace{\alpha, \beta, \dots, \gamma}^m \right), \quad \alpha, \beta, \dots, \gamma \in A, \quad \mathbf{a} \in A^m. \quad (4.54)$$

The m th direct power becomes a binary semigroup by endowing m -tuples with the componentwise binary operation $(\hat{+})$ as

$$\mathbf{a}_1 \hat{+} \mathbf{a}_2 = \left(\overbrace{\alpha_1, \beta_1, \dots, \gamma_1}^m \right) \hat{+} \left(\overbrace{\alpha_2, \beta_2, \dots, \gamma_2}^m \right) = \left(\overbrace{\alpha_1 + \alpha_2, \beta_1 + \beta_2, \dots, \gamma_1 + \gamma_2}^m \right). \quad (4.55)$$

The derived n -ary operation for m -tuples (on the m th direct power) is then defined componentwise by analogy with (4.53)

$$\nu^{[n]}[\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n] = \mathbf{a}_1 \hat{+} \mathbf{a}_2 \hat{+} \dots \hat{+} \mathbf{a}_n. \quad (4.56)$$

Now using shifts, instead of (4.56) we construct a nonderived n -ary operation on the direct power.

Definition 4.13. A cyclic m -shift operator s is defined for the m -tuple (4.54) by

$$\overbrace{s\alpha = \beta, s\beta = \gamma, \dots, s\gamma = \alpha}^m, \quad (4.57)$$

and $s^m = \text{id}$.

For instance, in this notation, if $m = 3$ and $\mathbf{a} = (\alpha, \beta, \gamma)$, then $s\mathbf{a} = (\gamma, \alpha, \beta)$, $s^2\mathbf{a} = (\beta, \gamma, \alpha)$, $s^3\mathbf{a} = \mathbf{a}$ (as in the previous subsection).

To obtain a nonderived n -ary operation, by analogy with (4.50), we deform by shifts the derived n -ary operation (4.56).

Definition 4.14. The shift deformation by (4.57) of the derived operation $\nu^{[n]}$ on the direct power \mathcal{A}^m is defined noncomponentwise by

$$\nu_s^{[n]}[\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n] = \sum_{i=1}^n s^{i-1} \mathbf{a}_i = \mathbf{a}_1 \hat{+} s\mathbf{a}_2 \hat{+} \dots \hat{+} s^{n-1} \mathbf{a}_n, \quad (4.58)$$

where $\mathbf{a} \in A^m$ (4.54) and $s^0 = \text{id}$.

Note that till now there exist no relations between n and m .

Proposition 4.15. The shift deformed operation $\nu_s^{[n]}$ is totally associative, if

$$s^{n-1} = \text{id}, \quad (4.59)$$

$$m = n - 1. \quad (4.60)$$

Proof. We compute

$$\begin{aligned} & \nu_s^{[n]} \left[\nu_s^{[n]} [\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n], \mathbf{a}_{n+1}, \mathbf{a}_{n+2}, \dots, \mathbf{a}_{2n-1} \right] \\ &= (\mathbf{a}_1 \hat{+} s\mathbf{a}_2 \hat{+} \dots \hat{+} s^{n-1} \mathbf{a}_n) \hat{+} s\mathbf{a}_{n+1} \hat{+} s^2 \mathbf{a}_{n+2} \hat{+} \dots \hat{+} s^{n-1} \mathbf{a}_{2n-1} \\ &= \mathbf{a}_1 \hat{+} s(\mathbf{a}_2 \hat{+} s\mathbf{a}_3 \hat{+} \dots \hat{+} s^{n-1} \mathbf{a}_{n+1}) \hat{+} s^2 \mathbf{a}_{n+2} \hat{+} s^3 \mathbf{a}_{n+3} \hat{+} \dots \hat{+} s^{n-1} \mathbf{a}_{2n-1} \\ &\vdots \\ & \mathbf{a}_1 \hat{+} s\mathbf{a}_2 \hat{+} \dots \hat{+} s^{n-2} \mathbf{a}_n \hat{+} s^{n-1} (\mathbf{a}_{n+1} \hat{+} s\mathbf{a}_{n+2} \hat{+} s^2 \mathbf{a}_{n+3} \hat{+} \dots \hat{+} s^{n-1} \mathbf{a}_{2n-1}) \\ & \nu_s^{[n]} [\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_{n-1}, \nu_s^{[n]} [\mathbf{a}_n, \mathbf{a}_{n+1}, \mathbf{a}_{n+2}, \dots, \mathbf{a}_{2n-1}]], \end{aligned} \quad (4.61)$$

which satisfy in all lines, if $s^{n-1} = \text{id}$ (4.59). \square

Corollary 4.16. The set of $(n-1)$ -tuples (4.54) with the shift deformed associative operation (4.58) is a nonderived n -ary semigroup $\mathcal{S}_{shift}^{[n]} = \langle \{\mathbf{a}\} \mid \nu_s^{[n]} \rangle$ constructed from the binary semigroup \mathcal{A} .

Proposition 4.17. If the binary semigroup \mathcal{A} is commutative, then $\mathcal{S}_{shift}^{[n]}$ becomes a nonderived n -ary group $\mathcal{G}_{shift}^{[n]} = \langle \{\mathbf{a}\} \mid \nu_s^{[n]}, \bar{\nu}_s^{[1]} \rangle$, such that each element $\mathbf{a} \in \mathcal{A}^{n-1}$ has a unique querelement $\bar{\mathbf{a}}$ (an analog of inverse) by

$$\bar{\mathbf{a}} = \bar{\nu}_s^{[1]}[\mathbf{a}] = -(\mathbf{sa} \hat{+} s^2 \mathbf{a} \hat{+} \dots \hat{+} s^{n-2} \mathbf{a}), \quad (4.62)$$

where $\bar{\nu}_s^{[1]} : \mathcal{A}^{n-1} \rightarrow \mathcal{A}^{n-1}$ is an unary quereoperation.

Proof. We have the definition of the querelement

$$\nu_s^{[n]}[\bar{\mathbf{a}}, \mathbf{a}, \dots, \mathbf{a}] = \mathbf{a}, \quad (4.63)$$

where $\bar{\mathbf{a}}$ can be on any place. So (4.58) gives the equation

$$\bar{\mathbf{a}} \hat{+} \mathbf{sa} \hat{+} s^2 \mathbf{a} \hat{+} \dots \hat{+} s^{n-2} \mathbf{a} \hat{+} \mathbf{a} = \mathbf{a}, \quad (4.64)$$

which can be resolved for the commutative and cancellative semigroup \mathcal{A} only, and the solution is (4.62). If $\bar{\mathbf{a}}$ is on the i th place in (4.63), then it has the coefficient s^{i-1} , and we multiply both sides by s^{n-i} to get $\bar{\mathbf{a}}$ without any shift operator coefficient using (4.59), which gives the same solution (4.62). \square

For $n = 4$ and $\mathbf{a} = (\alpha, \beta, \gamma)$, the equation (4.63) is

$$\bar{\mathbf{a}} \hat{+} \mathbf{sa} \hat{+} s^2 \mathbf{a} \hat{+} \mathbf{a} = \mathbf{a} \quad (4.65)$$

and (see(4.49))

$$\bar{\mathbf{a}} = -(\mathbf{sa} \hat{+} s^2 \mathbf{a}) \quad (4.66)$$

so (cf. (4.43))

$$\bar{\mathbf{a}} = \overline{(\alpha, \beta, \gamma)} = -(\gamma + \beta, \alpha + \gamma, \beta + \alpha). \quad (4.67)$$

It is known that the existence of an identity (as a neutral element) is not necessary for polyadic groups, and only a querelement is important DÖRNTE [1929], GLEICHGEWICHT AND GŁĄZEK [1967]. Nevertheless, we have

Proposition 4.18. If the commutative and cancellative semigroup \mathcal{A} has zero $0 \in A$, then the n -ary group $\mathcal{G}_{shift}^{[n]}$ has a set of polyadic identities (idempotents) satisfying

$$\mathbf{e} \hat{+} \mathbf{se} \hat{+} \dots \hat{+} s^{n-2} \mathbf{e} = \mathbf{0}, \quad (4.68)$$

where $\mathbf{0} = \left(\overbrace{0, 0, \dots, 0}^{n-1} \right)$ is the zero $(n-1)$ -tuple.

Proof. The definition of polyadic identity in terms of the deformed n -ary product in the direct power is

$$\nu_s^{[n]} \left[\overbrace{\mathbf{e}, \mathbf{e}, \dots, \mathbf{e}}^{n-1}, \mathbf{a} \right] = \mathbf{a}, \quad \forall \mathbf{a} \in \mathcal{A}^{n-1}. \quad (4.69)$$

Using (4.58) we get the equation

$$\mathbf{e} \hat{+} \mathbf{se} \hat{+} s^2 \mathbf{e} \hat{+} \dots \hat{+} s^{n-2} \mathbf{e} \hat{+} \mathbf{a} = \mathbf{a}. \quad (4.70)$$

After cancellation by \mathbf{a} we obtain (4.68). \square

For $n = 4$ and $e = (\alpha_0, \beta_0, \gamma_0)$ we obtain an infinite set of identities satisfying

$$e = (\alpha_0, \beta_0, \gamma_0), \quad \alpha_0 + \beta_0 + \gamma_0 = 0. \quad (4.71)$$

To see that they are 4-ary idempotents, insert $a = e$ into (4.69).

Thus, starting from a binary semigroup \mathcal{A} , using our polyadization procedure we have obtained a nonderived n -ary group on $(n - 1)$ th direct power \mathcal{A}^{n-1} with the shift deformed multiplication. This construction reminds the Post-like associative quiver from DUPLIJ [2018, 2022a], and allows us to construct a nonderived n -ary group from any semigroup in the unified way presented here.

4.4. Polyadization of binary supergroups. Here we consider a more exotic possibility, when the B -matrices are defined over the Grassmann algebra, and therefore can represent supergroups (see (3.21) and below). In this case B 's can be supermatrices of two kinds, even and odd, which have different properties BEREZIN [1987], LEITES [1983]. The general polyadization procedure remains the same, as for ordinary matrices considered before (see Definition 4.1), and therefore we confine ourselves to examples only.

Indeed, to obtain an n -ary matrix (semi)group represented now by the Q -supermatrices (4.1) of the nonstandard form, we should take $(n - 1)$ initial B -supermatrices which present a simple ($k = 1$ in (3.19)) binary (semi)supergroup, which now have different parameters $B_{y_i} \equiv B_{y_i}((p_{even} | p_{odd}) \times (p_{even} | p_{odd}))$, $i = 1, \dots, n - 1$, where p_{even} and p_{odd} are the even and odd dimensions of the B -supermatrix. The closure of the Q -supermatrix multiplication is governed by the closure of B -supermatrix multiplication (4.3)–(4.5) in the initial binary (semi)supergroup.

4.4.1. Polyadization of $GL(1 | 1, \Lambda)$. Let $\Lambda = \Lambda_{even} \oplus \Lambda_{odd}$ be a Grassmann algebra over \mathbb{C} , where Λ_{even} and Λ_{odd} are its even and odd parts (it can be also any commutative superalgebra). We provide (in brief) the polyadization procedure of the general linear supergroup $GL(1 | 1, \Lambda)$ for $n = 3$. The 4-parameter block (invertible) supermatrices become $B_{y_i} = \begin{pmatrix} a_i & \alpha_i \\ \beta_i & b_i \end{pmatrix} \in GL(1 | 1, \Lambda)$, where the parameters are $y_i = (a_i, b_i, \alpha_i, \beta_i) \in \Lambda_{even} \times \Lambda_{even} \times \Lambda_{odd} \times \Lambda_{odd}$, $i = 1, 2$. Thus, the 8-parameter ternary supergroup $GL^{[[3]]}(1 | 1, \Lambda) = \langle \{Q_{y_1, y_2}\} | \mu^{[[3]]} \rangle$ is represented by the following 4×4 Q -supermatrices

$$Q_{y_1, y_2} = \begin{pmatrix} 0 & B_{y_1} \\ B_{y_2} & 0 \end{pmatrix} = \begin{pmatrix} 0 & 0 & a_1 & \alpha_1 \\ 0 & 0 & \beta_1 & b_1 \\ a_2 & \alpha_2 & 0 & 0 \\ \beta_2 & b_2 & 0 & 0 \end{pmatrix} \in GL^{[[3]]}(1 | 1, \Lambda), \quad (4.72)$$

which satisfy the ternary (nonderived) multiplication

$$\mu^{[[3]]} [Q_{y'_1, y'_2}, Q_{y''_1, y''_2}, Q_{y'''_1, y'''_2}] = Q_{y'_1, y'_2} Q_{y''_1, y''_2} Q_{y'''_1, y'''_2} = Q_{y_1, y_2}. \quad (4.73)$$

In terms of the block matrices B_{y_i} the multiplication (4.17) becomes (see (4.2)–(4.5))

$$B_{y'_1} B_{y''_2} B_{y'''_1} = B_{y_1}, \quad (4.74)$$

$$B_{y'_2} B_{y'''_1} B_{y''_2} = B_{y_2}. \quad (4.75)$$

In terms of the B -supermatrix parameters the supergroup $GL^{[[3]]}(1 | 1, \Lambda)$ is defined by

$$\begin{aligned} \alpha'_1 \beta''_2 a'''_1 + a'_1 \alpha''_2 \beta'''_1 + \alpha'_1 b''_2 \beta'''_1 + a'_1 a''_2 a'''_1 &= a_1, & \beta'_1 a''_2 \alpha'''_1 + \beta'_1 \alpha''_2 b'''_1 + b'_1 \beta''_2 \alpha'''_1 + b'_1 b''_2 b'''_1 &= b_1, \\ \alpha'_1 \beta''_2 \alpha'''_1 + a'_1 a''_2 \alpha'''_1 + a'_1 \alpha''_2 b'''_1 + \alpha'_1 b''_2 b'''_1 &= \alpha_1, & \beta'_1 \alpha''_2 \beta'''_1 + \beta'_1 a''_2 a'''_1 + b'_1 \beta''_2 a'''_1 + b'_1 b''_2 \beta'''_1 &= \beta_1, \\ \alpha'_2 \beta''_1 a'''_2 + a'_2 \alpha''_1 \beta'''_2 + \alpha'_2 b''_1 \beta'''_2 + a'_2 a''_1 a'''_2 &= a_2, & \beta'_2 a''_1 \alpha'''_2 + \beta'_2 \alpha''_1 b'''_2 + b'_2 \beta''_1 \alpha'''_2 + b'_2 b''_1 b'''_2 &= b_2, \\ \alpha'_2 \beta''_1 \alpha'''_2 + a'_2 a''_1 \alpha'''_2 + a'_2 \alpha''_1 b'''_2 + \alpha'_2 b''_1 b'''_2 &= \alpha_2, & \beta'_2 \alpha''_1 \beta'''_2 + \beta'_2 a''_1 a'''_2 + b'_2 \beta''_1 a'''_2 + b'_2 b''_1 \beta'''_2 &= \beta_2. \end{aligned} \quad (4.76)$$

The unique querelement in $GL^{[[3]]}(1 | 1, \Lambda)$ can be found from the equation (see (4.11))

$$Q_{y_1, y_2} Q_{y_1, y_2} \bar{Q}_{y_1, y_2} = Q_{y_1, y_2}, \quad (4.77)$$

where the solution is

$$\bar{Q}_{y_1, y_2} = \begin{pmatrix} 0 & \bar{B}_{y_1} \\ \bar{B}_{y_2} & 0 \end{pmatrix}, \quad (4.78)$$

with (see (4.12))

$$\bar{B}_{y_1} = B_{y_2}^{-1}, \quad \bar{B}_{y_2} = B_{y_1}^{-1}, \quad (4.79)$$

and $B_{y_1}^{-1}, B_{y_2}^{-1} \in GL(1 | 1, \Lambda)$.

Definition 4.19. We call $GL^{[[3]]}(1 | 1, \Lambda)$ a polyadic (ternary) general linear supergroup obtained by the polyadization procedure from the binary linear supergroup $GL(1 | 1, \Lambda)$.

The ternary identity $E_4^{(3)}$ of $GL^{[[3]]}(1 | 1, \Lambda)$ has the form (see (4.9))

$$E_4^{(3)} = \begin{pmatrix} 0 & E_2 \\ E_2 & 0 \end{pmatrix}, \quad (4.80)$$

where E_2 is the identity of $GL(1 | 1, \Lambda)$, and is ternary idempotent

$$E_4^{(3)} E_4^{(3)} E_4^{(3)} = E_4^{(3)}. \quad (4.81)$$

The ternary supergroup $GL^{[[3]]}(1 | 1, \Lambda)$ contains the infinite number of ternary idempotents Q_{y_1, y_2}^{idemp} defined by the system of equations

$$Q_{y_1, y_2}^{idemp} Q_{y_1, y_2}^{idemp} Q_{y_1, y_2}^{idemp} = Q_{y_1, y_2}^{idemp}, \quad (4.82)$$

which gives

$$B_{y_1}^{idemp} B_{y_2}^{idemp} = E_2. \quad (4.83)$$

Therefore, the idempotents are determined by $8 - 4 = 4$ Grassmanian parameters. One of the ways to realize this is to exclude from (4.83) the 2×2 B-supermatrix. In this case, the idempotents in the supergroup $GL^{[[3]]}(1 | 1, \Lambda)$ become

$$Q_{y_1, y_2}^{idemp} = \begin{pmatrix} 0 & B_{y_1} \\ (B_{y_1})^{-1} & 0 \end{pmatrix}, \quad (4.84)$$

where $B_{y_i} \in GL(1 | 1, \Lambda)$ is an arbitrary 2×2 supermatrix of the standard form (see Remark 4.11).

In the same way one can polyadize any supergroup that can be presented by supermatrices.

Acknowledgements. The author is grateful to Vladimir Akulov, Mike Hewitt, Dimitrij Leites, Vladimir Tkach, Raimund Vogl and Alexander Voronov for useful discussions, and valuable help.

REFERENCES

- BELOUSOV, V. D. (1972). *n*-Ary Quasigroups. Kishinev: Shtintsa. [Cited on page 3]
- BEREZIN, F. A. (1987). *Introduction to Superanalysis*. Dordrecht: Reidel. [Cited on pages 6 and 15]
- CARLSSON, R. (1980). *N*-ary algebras. *Nagoya Math. J.* **78** (1), 45–56. [Cited on page 4]
- CURTIS, C. W. AND I. REINER (1962). *Representation theory of finite groups and associative algebras*. Providence: AMS. [Cited on page 2]
- DIETZEL, C. AND G. MITTAL (2021). Summands of finite group algebras. *Czech. Math. J.* **71** (4), 1011–1014. [Cited on page 6]
- DÖRNTE, W. (1929). Untersuchungen über einen verallgemeinerten Gruppenbegriff. *Math. Z.* **29**, 1–19. [Cited on pages 3, 12, and 14]
- DUPLIJ, S. (2018). Polyadic algebraic structures and their representations. In: S. DUPLIJ (Ed.), *Exotic Algebraic and Geometric Structures in Theoretical Physics*, New York: Nova Publishers, pp. 251–308. arXiv:math.RT/1308.4060. [Cited on pages 2 and 15]
- DUPLIJ, S. (2019). Arity shape of polyadic algebraic structures. *J. Math. Physics, Analysis, Geometry* **15** (1), 3–56. [Cited on pages 2 and 3]
- DUPLIJ, S. (2021). Higher braid groups and regular semigroups from polyadic-binary correspondence. *Mathematics* **9** (9), 972. [Cited on page 7]
- DUPLIJ, S. (2022a). *Polyadic Algebraic Structures*. Bristol: IOP Publishing. [Cited on pages 2, 3, 7, and 15]
- DUPLIJ, S. (2022b). Polyadic analogs of direct product. *Universe* **8** (4), 230. [Cited on page 2]
- ERDMANN, K. AND T. HOLM (2018). *Algebras and representation theory*. Cham: Springer. [Cited on pages 2 and 3]
- FULTON, W. AND J. HARRIS (2004). *Representation Theory: A First Course*. New York: Springer. [Cited on page 2]
- GAL'MAK, A. M. (2003). *n*-Ary Groups, Part 1. Gomel: Gomel University. [Cited on pages 3 and 7]
- GLEICHGEWICHT, B. AND K. GŁĄZEK (1967). Remarks on *n*-groups as abstract algebras. *Colloq. Math.* **17**, 209–219. [Cited on pages 3 and 14]
- HARADA, M. (1970). On semi-simple abelian categories. *Osaka J. Math.* **7**, 89–95. [Cited on page 2]
- HAZEWINKEL, M. AND N. M. GUBARENI (2016). *Algebras, Rings and Modules: Non-commutative Algebras and Rings*. Boca Raton: CRC Press. [Cited on pages 3 and 5]
- HERSTEIN, I. N. (1996). *Noncommutative rings*. Washington: Math. Assoc. America. [Cited on pages 2, 3, and 5]
- HUNGERFORD, T. W. (1974). *Algebra*. New York: Springer. [Cited on page 2]
- KNAPP, A. W. (1986). *Representation Theory of Semisimple Groups*. Princeton: Princeton Univ. Press. [Cited on page 2]
- KNOP, F. (2006). A construction of semisimple tensor categories. *Comptes Rendus Math.* **343** (1), 15–18. [Cited on page 2]
- LAM, T. Y. (1991). *A First Course in Noncommutative Rings*. New York: Springer. [Cited on pages 2, 3, and 5]
- LAMBEK, J. (1966). *Lectures on Rings and Modules*. Providence: Blaisdell. [Cited on pages 2 and 3]
- LEITES, D. (1983). *Supermanifold Theory*. Petrozavodsk: Math. Methods Sci. Invest. [Cited on pages 6 and 15]
- LISTER, W. G. (1971). Ternary rings. *Trans. Amer. Math. Soc.* **154**, 37–55. [Cited on page 5]
- NIKITIN, A. N. (1984). Semisimple Artinian (2,*n*)-rings. *Moscow Univ. Math. Bull.* **39** (6), 1–6. [Cited on pages 2, 3, and 5]
- POST, E. L. (1940). Polyadic groups. *Trans. Amer. Math. Soc.* **48**, 208–350. [Cited on page 2]
- PROFERA, L. (1982). Anelli ternari di Lister semisemplici. *Note Mat.* **11**, 1–56. [Cited on page 5]
- ROTMAN, J. J. (2010). *Advanced modern algebra* (2nd ed. ed.). Providence: AMS. [Cited on page 2]
- SIMSON, D. (1977). Pure semisimple categories and rings of finite representation type. *J. Algebra* **48** (2), 290–296. [Cited on page 2]
- WEDDERBURN, J. H. M. (1908). On hypercomplex numbers. *Proc. London Math. Soc. Second Series* **6** (1), 77–118. [Cited on page 2]
- ZUPNIK, D. (1967). Polyadic semigroups. *Publ. Math. (Debrecen)* **14**, 273–279. [Cited on page 12]