

Distances in $\frac{1}{|x-y|^{2d}}$ percolation models for all dimensions

Johannes Bäuml^{*}

August 10, 2022

Abstract. We study independent long-range percolation on \mathbb{Z}^d for all dimensions d , where the vertices u and v are connected with probability 1 for $\|u - v\|_\infty = 1$ and with probability $p(\beta, \{u, v\}) = 1 - e^{-\beta \int_{u+[0,1)^d} \int_{v+[0,1)^d} \frac{1}{\|x-y\|^{2d}} dx dy} \approx \frac{\beta}{\|u-v\|^{2d}}$ for $\|u - v\|_\infty \geq 2$. Let $u \in \mathbb{Z}^d$ be a point with $\|u\| = n$. We show that both the graph distance between the origin $\mathbf{0}$ and u and the diameter of the box $\{0, \dots, n\}^d$ grow like $n^{\theta(\beta)}$, where $0 < \theta(\beta) < 1$. We also show that the graph distance has the same asymptotic growth when two vertices u, v with $\|u - v\|_2 > 1$ are connected with a probability that is close enough to $p(\beta, \{u, v\})$. Furthermore, we determine the asymptotic behavior of $\theta(\beta)$ for large β .

Contents

1	Introduction and statement of main results	2
1.1	Main results	2
1.2	The continuous model	3
1.3	Notation	4
1.4	Related work	4
2	Asymptotic behavior of the distance exponent for large β	5
2.1	Bounds on connection probabilities	5
2.2	Submultiplicativity and the upper bound in Theorem 1.2	7
2.3	Spacing between points with long bonds	10
2.4	The lower bound in Theorem 1.2	13
3	Connected sets in graphs	15
4	Distances in V_0^n	21
4.1	Graph distances of far away points	21
4.2	The second moment bound	24
4.3	Graph distances between points and boxes	27
5	The proof of Theorem 1.1	35
5.1	Distances between certain points	36
5.2	Supermultiplicativity of $\Lambda(n, \beta)$	38
5.3	The diameter of boxes	43
6	Comparison with different inclusion probabilities	45

^{*}DEPARTMENT OF MATHEMATICS, TU MUNICH, GERMANY. E-Mail: johannes.baeumler@tum.de
 MSC Class: 05C12, 60K35, 82B27, 82B43

Key words: Long-range percolation, graph distance, diameter, renormalization

1 Introduction and statement of main results

Consider independent long-range bond percolation on \mathbb{Z}^d where all edges $\{u, v\}$ with $\|u - v\|_\infty = 1$ are open and an edge $\{u, v\}$ with $\|u - v\|_\infty \geq 2$ is open with probability

$$p(\beta, \{u, v\}) := 1 - e^{-\beta \int_{u+c} \int_{v+c} \frac{1}{\|x-y\|^{2d}} dx dy},$$

where $\mathcal{C} := [0, 1]^d$. We call the corresponding probability measure \mathbb{P}_β and denote its expectation by \mathbb{E}_β . The resulting graph is clearly connected and the graph distance $d(x, y)$ between two points $x, y \in \mathbb{Z}^d$ satisfies $d(x, y) \leq \|x - y\|_\infty$. We are interested in the scaling of the typical distance of two points $x, y \in \mathbb{Z}^d$ and the scaling of the diameter of boxes $\{0, \dots, N\}^d$. In [16] it is proven that the typical diameter of some box grows at most polynomially with some power strictly smaller than 1. More precisely, Coppersmith, Gamarnik, and Sviridenko proved that for all $\beta > 0$ there exists an exponent $\theta' = \theta'(\beta) < 1$ such that $\lim_{N \rightarrow \infty} \mathbb{P}_\beta \left(\text{Diam} \left(\{0, \dots, N\}^d \right) \leq N^{\theta'} \right) = 1$. However, the authors do not give any polynomial lower bound for dimensions $d \geq 2$. An analogous lower bound was already conjectured in [7, 12], and an exact lower bound was later proven to hold in one dimension. In [20] Ding and Sly showed that for the connection probability $p(\beta, \{u, v\})$ given by $p(\beta, \{u, v\}) = \frac{\beta}{|u-v|^2} \wedge 1$ for $|u - v| \geq 2$ and $p(\beta, \{u, v\}) = 1$ for $|u - v| = 1$ the typical distance between the two points $0, n \in \mathbb{Z}$ and the diameter of $\{0, \dots, n\}$ both grow like n^θ for some $\theta \in (0, 1]$, where $\theta = 1$ if and only if $\beta = 0$. More precisely, they proved that

$$n^\theta \approx_P d(0, n) \approx_P \text{Diam}(\{0, \dots, n\}) \approx_P \mathbb{E}[d(0, n)]$$

where the notation $A(n) \approx_P B(n)$ means that for all $\varepsilon > 0$ there exist $0 < c < C < \infty$ such that $\mathbb{P}(cB(n) \leq A(n) \leq CB(n)) > 1 - \varepsilon$ for all n . In this paper, we prove an analogous result for all dimensions.

1.1 Main results

Theorem 1.1. *For all dimensions d and all $\beta > 0$ there exists an exponent $\theta = \theta(\beta) < 1$ such that*

$$\|x\|^\theta \approx_P d(\mathbf{0}, x) \approx_P \mathbb{E}_\beta [d(\mathbf{0}, x)] \quad (1)$$

and

$$k^\theta \approx_P \text{Diam}(\{0, \dots, k\}^d) \approx_P \mathbb{E}_\beta \left[\text{Diam}(\{0, \dots, k\}^d) \right]. \quad (2)$$

We write $\mathbf{0}$ for the vector with all entries equal to 0 and the notation $A(x) \approx_P B(x)$ means that for all $\varepsilon > 0$ there exist $0 < c < C < \infty$ such that $\mathbb{P}_\beta(cB(x) \leq A(x) \leq CB(x)) > 1 - \varepsilon$ for all $x \in \mathbb{Z}^d$. We will show an analogous result for other inclusion probabilities,

like $\frac{\beta}{\|x-y\|^{2d}} \wedge 1$, in section 6. Furthermore, we give bounds on the asymptotic behavior of the function $\theta(\beta)$ for large β .

Theorem 1.2. *For all dimensions d there exist constants $0 < c < C < \infty$ such that*

$$\frac{c}{\log(\beta)} \leq \theta(\beta) \leq \frac{C}{\log(\beta)}$$

for all $\beta \geq 2$.

1.2 The continuous model

For $\beta > 0$, the described discrete percolation model has a self-similarity that comes from a coupling with the underlying continuous model, that we will now describe for any dimension. This will also explain our, at first sight complicated, choice of the connection probability. Consider a Poisson point process $\tilde{\mathcal{E}}$ on $\mathbb{R}^d \times \mathbb{R}^d$ with intensity $\frac{\beta}{2\|t-s\|^{2d}}$. Define the symmetrized version \mathcal{E} by $\mathcal{E} := \{(t, s) \in \mathbb{R}^d \times \mathbb{R}^d : (s, t) \in \tilde{\mathcal{E}}\} \cup \tilde{\mathcal{E}}$. For $u, v \in \mathbb{Z}^d$ with $\|u - v\|_\infty \geq 1$ we put an edge between u and v if and only if $((u + \mathcal{C}) \times (v + \mathcal{C})) \cap \mathcal{E} \neq \emptyset$, where we use the notation $\mathcal{C} = [0, 1)^d$. The cardinality of $((u + \mathcal{C}) \times (v + \mathcal{C})) \cap \tilde{\mathcal{E}}$ is a random variable with Poisson distribution and parameter $\int_{u+\mathcal{C}} \int_{v+\mathcal{C}} \frac{\beta}{2\|t-s\|^{2d}} ds dt$. Thus, by the properties of Poisson processes, the probability that $u \approx v$ equals

$$\begin{aligned} \mathbb{P}(((u + \mathcal{C}) \times (v + \mathcal{C})) \cap \mathcal{E} = \emptyset) &= \mathbb{P}(((u + \mathcal{C}) \times (v + \mathcal{C})) \cap \tilde{\mathcal{E}} = \emptyset)^2 \\ &= \left(e^{-\int_{u+\mathcal{C}} \int_{v+\mathcal{C}} \frac{\beta}{2\|t-s\|^{2d}} ds dt} \right)^2 = e^{-\int_{u+\mathcal{C}} \int_{v+\mathcal{C}} \frac{\beta}{\|t-s\|^{2d}} ds dt} = 1 - p(\beta, \{u, v\}) \end{aligned}$$

which is exactly the probability that $u \approx v$ under the measure \mathbb{P}_β . Note that for $\{u, v\}$ with $\|u - v\|_\infty = 1$ we have $\int_{u+\mathcal{C}} \int_{v+\mathcal{C}} \frac{\beta}{\|t-s\|^{2d}} ds dt = \infty$. So we really get that all edges of the form $\{u, v\}$ with $\|u - v\|_\infty = 1$ are present. The construction with the Poisson process also implies that the presence of different bonds is independent and thus the resulting measure of the random graph constructed above equals \mathbb{P}_β . The chosen inclusion probabilities have many advantages. First of all, the resulting model is invariant under translation and invariant under the reflection of coordinates, i.e., when we change the i -th component $p_i(x)$ of all $x \in \mathbb{Z}^d$ to $-p_i(x)$. Furthermore, the model has the following self-similarity: For some vector $u = (p_1(u), \dots, p_d(u)) \in \mathbb{Z}^d$ and $n \in \mathbb{N}_{>0}$ we define the translated boxes $V_u^n := \prod_{i=1}^d \{p_i(u)n, \dots, (p_i(u) + 1)n - 1\} = nu + \prod_{i=1}^d \{0, \dots, n - 1\}$. Then for all points $u, v \in \mathbb{Z}^d$, and all $n \in \mathbb{N}_{>0}$ one has

$$\begin{aligned} \mathbb{P}_\beta(V_u^n \approx V_v^n) &= \prod_{x \in V_u^n} \prod_{y \in V_v^n} \mathbb{P}_\beta(x \approx y) = \prod_{x \in V_u^n} \prod_{y \in V_v^n} e^{-\int_{x+\mathcal{C}} \int_{y+\mathcal{C}} \frac{\beta}{\|t-s\|^{2d}} ds dt} \\ &= e^{-\sum_{x \in V_u^n} \sum_{y \in V_v^n} \int_{x+\mathcal{C}} \int_{y+\mathcal{C}} \frac{\beta}{\|t-s\|^{2d}} ds dt} = e^{-\int_{nu+[0,n)^d} \int_{nv+[0,n)^d} \frac{\beta}{\|t-s\|^{2d}} ds dt} \\ &= e^{-\int_{u+\mathcal{C}} \int_{v+\mathcal{C}} \frac{\beta}{\|t-s\|^{2d}} ds dt} = \mathbb{P}_\beta(u \approx v) \end{aligned}$$

which shows the self-similarity of the model. Also observe that for any $\alpha \in \mathbb{R}_{>0}$ the process $\alpha\tilde{\mathcal{E}} := \{(x, y) \in \mathbb{R}^d \times \mathbb{R}^d : (\frac{1}{\alpha}x, \frac{1}{\alpha}y) \in \tilde{\mathcal{E}}\}$ is again a Poisson point process with intensity $\frac{\beta}{2\|x-y\|^{2d}}$.

1.3 Notation

We use the notation e_i for the i -th standard unit vector in \mathbb{R}^d . For a vector $y \in \mathbb{Z}^d$ we write $p_i(y)$ for the i -th coordinate of y , i.e., $p_i(y) = \langle e_i, y \rangle$. We also use the notation $\mathbf{0}$ for the vector with all entries equal to 0 and the notation $\mathbf{1}$ for the vector with all entries equal to 1. When we write $\|u\|$ we always mean the 2-norm of the vector u . We write \mathcal{S}_k for the set of points $\{x \in \mathbb{Z}^d : \|x\|_\infty = k\}$ and $\mathcal{S}_{\geq k}$ for the set $\{x \in \mathbb{Z}^d : \|x\|_\infty \geq k\}$. For the closed ball of radius r around $x \in \mathbb{Z}^d$ in the ∞ -norm we use the notation $B_r(x)$, i.e., $B_r(x) = \{y \in \mathbb{Z}^d : \|x - y\|_\infty \leq r\}$. When we want to emphasize that we work on certain subgraphs $A \subset \mathbb{Z}^d$ we will write $d_A(x, y)$ for the graph distance inside the set A , i.e., when we consider edges with both endpoints inside A only. Whenever we write $\text{Diam}(A)$ for some set $A \subset \mathbb{Z}^d$ we always mean the inside diameter of this set, i.e., $\text{Diam}(A) = \max_{x, y \in A} d_A(x, y)$. For a graph (V, E) we think of the percolation configuration as a random element $\omega \in \{0, 1\}^E$, where we say that the edge e exists or is open or present if $\omega(e) = 1$. For $\omega \in \{0, 1\}^E$ and $e \in E$ we define the configuration $\omega^{e^-} \in \{0, 1\}^E$ by

$$\omega^{e^-}(\tilde{e}) = \begin{cases} 0 & \tilde{e} = e \\ \omega(e) & \tilde{e} \neq e \end{cases},$$

so this is the configuration where we deleted the edge e .

1.4 Related work

The scaling of the graph distance, also called chemical distance or hop-count distance, is a central characteristic of a random graph and has also been examined for many different models of percolation, see for example [1, 3, 7, 11–14, 16, 18–21, 24, 25, 27, 32]. For all dimensions d one can also consider the long-range percolation model with connection probability $p(\beta, \{u, v\}) = \frac{\beta}{\|u-v\|^s}$. There is a first transition of the behavior at $s = d$, which is natural as the resulting random graph is locally finite if and only if $s > d$. For $s < d$ the graph distance between two points is at most $\lceil \frac{d}{d-s} \rceil$ [8], whereas for $s = d$, the diameter of the box $\{0, \dots, n\}^d$ is $\Theta\left(\frac{\log(n)}{\log(\log(n))}\right)$ [16]. In [7, 12–14] the authors proved that for $d < s < 2d$ the typical distance between two points of Euclidean distance n grows like $\log(n)^\Delta$, where $\Delta^{-1} = \log_2\left(\frac{2d}{s}\right)$. The behavior of the typical distance for long-range percolation on \mathbb{Z}^d also changes at $s = 2d$. For $s > 2d$ the graph distance grows at least linearly in the Euclidean distance of two points, as proven in [11]. In [20] it is shown that the typical distance for $d = 1, s = 2$ grows like n^θ for some $\theta \in (0, 1)$. For $d \geq 2$ and $s = 2d$ the authors in [16] proved a polynomial upper bound on the graph distance, but no lower bound. In this paper, we show a matching polynomial lower bound for all dimensions d , similar to the results of [20] in one dimension. For fixed dimension d , different characteristics of the function $\beta \mapsto \theta(\beta)$ are considered in the companion paper [4], where it is shown that the exponent is continuous and strictly decreasing in β . Together with the fact $\theta(0) = 1$ and Theorem 1.2 this shows that the long-range percolation model for $s = 2d$ interpolates between linear growth and subpolynomial growth as β goes from 0 to $+\infty$.

Another line of research is to investigate what happens when one drops the assumption that $p(\beta, \{u, v\}) = 1$ for all nearest neighbor edges $\{u, v\}$, but assigns i.i.d. random

variables to the nearest neighbor edges instead. For $d = 1$, there is a change of behavior at $s = 2$. As proven in [2, 35] or [22], an infinite cluster can not emerge for $s > 2$ and for $s = 2, \beta \leq 1$, no matter how small $\mathbb{P}(k \approx k + 1)$ is. On the other hand, an infinite cluster can emerge for $s < 2$ and $s = 2, \beta > 1$ (see [33]). In [2] the authors proved that there is a discontinuity in the percolation density for $s = 2$, contrary to the situation for $s < 2$, as proven in [10, 29]. For models, for which an infinite cluster exists the behavior of the percolation model at and near criticality is also a well-studied question (cf. [5, 6, 10, 15, 17, 29–31]). It is not known up to now how the typical distance in long-range percolation grows for $s = 2d$ and $p(\beta, \{u, v\}) < 1$ for nearest-neighbor edges $\{u, v\}$, but we conjecture also a polynomial growth in the Euclidean distance, whenever an infinite cluster exists.

Acknowledgements. I thank Noam Berger for many useful discussions. I also thank Yuki Tokushige for helpful comments on an earlier version of this paper. This work is supported by TopMath, the graduate program of the Elite Network of Bavaria and the graduate center of TUM Graduate School.

2 Asymptotic behavior of the distance exponent for large β

In this chapter, we prove Theorem 1.2. On the way, in section 2.1, we prove several elementary bounds on connection probabilities between certain points and boxes in the long-range percolation graph that will also be used in the following sections. In section 2.2 we prove a submultiplicative structure of the expected distance between two points, leading to the existence of a distance exponent, and also to the inverse logarithmic upper bound in Theorem 1.2. In section 2.3 we show that, typically, vertices inside a box are not connected to more than one box that is far away. This is necessary in order to prove strict positivity of the distance exponent $\theta(\beta)$ in section 2.4, and the lower bound on $\theta(\beta)$ in Theorem 1.2.

2.1 Bounds on connection probabilities

Lemma 2.1. *For all $\beta \geq 0$, all $n \in \mathbb{N}$, and all $u, v \in \mathbb{Z}^d$ with $\|u - v\|_\infty \geq 2$ one has the upper bound*

$$\mathbb{P}_\beta(u \sim v) = \mathbb{P}_\beta(V_u^n \sim V_v^n) \leq \frac{2^{2d}\beta}{\|u - v\|_\infty^{2d}} \quad (3)$$

and one has the lower bound

$$\mathbb{P}_\beta(u \sim v) = \mathbb{P}_\beta(V_u^n \sim V_v^n) \geq \frac{(4d)^{-2d}\beta}{\|u - v\|_\infty^{2d}} \wedge \frac{1}{2}. \quad (4)$$

For all $k \geq 2$ one has

$$\mathbb{P}_\beta(\mathbf{0} \sim \mathcal{S}_{\geq k}) \leq \beta 50^d k^{-d} \quad (5)$$

and for $m \in \mathbb{N}$, a vertex $x \in V_{\mathbf{0}}^m$, and a box V_w^m with $\|w\|_\infty \geq 2$ one has

$$\mathbb{P}_\beta(x \sim V_w^m) \leq \frac{\beta 4^{2d}}{\|w\|_\infty^{2d} m^d}. \quad (6)$$

Proof. The equality $\mathbb{P}_\beta(u \sim v) = \mathbb{P}_\beta(V_u^n \sim V_v^n)$ is clear from the discussion about the underlying continuous model in section 1.2. We start with the proof of (3). Applying the inequalities $1 - e^{-x} \leq x$ and $\|\cdot\|_2 \geq \|\cdot\|_\infty$ we get that for two vertices u, v with $\|u - v\|_\infty \geq 2$

$$\begin{aligned} \mathbb{P}_\beta(u \sim v) &= 1 - e^{-\beta \int_{u+c} \int_{v+c} \frac{1}{\|x-y\|^{2d}} dx dy} \leq \beta \int_{u+c} \int_{v+c} \frac{1}{\|x-y\|^{2d}} dx dy \\ &\leq \beta \int_{u+c} \int_{v+c} \frac{1}{\|x-y\|_\infty^{2d}} dx dy \leq \frac{\beta}{(\|u-v\|_\infty - 1)^{2d}} \leq \frac{2^{2d} \beta}{\|u-v\|_\infty^{2d}}. \end{aligned} \quad (7)$$

In order to bound the connection probability between u and v from below, first observe that $\|x\|_2 \leq \|x\|_1 \leq d\|x\|_\infty$ for all $x \in \mathbb{R}^d$. Thus we have

$$\begin{aligned} \int_{u+c} \int_{v+c} \frac{1}{\|t-s\|^{2d}} ds dt &\geq d^{-2d} \int_{u+c} \int_{v+c} \frac{1}{\|t-s\|_\infty^{2d}} ds dt \\ &\geq d^{-2d} \int_{u+c} \int_{v+c} \frac{1}{(\|u-v\|_\infty + 1)^{2d}} ds dt \geq (2d)^{-2d} \frac{1}{\|u-v\|_\infty^{2d}} \end{aligned}$$

and this already gives

$$\mathbb{P}_\beta(u \sim v) \geq 1 - e^{-(2d)^{-2d} \frac{\beta}{\|u-v\|_\infty^{2d}}} \geq \frac{(4d)^{-2d} \beta}{\|u-v\|_\infty^{2d}} \wedge \frac{1}{2}$$

as $1 - e^{-x} \geq \frac{x}{2} \wedge \frac{1}{2}$ for all $x \in \mathbb{R}_{\geq 0}$. So we showed (4).

For each point $x \in \mathcal{S}_k = \{z \in \mathbb{Z}^d : \|z\|_\infty = k\}$, at least one of its coordinates $p_i(x)$ equals $-k$ or $+k$. All other coordinates can be any integer between $-k$ and $+k$. Thus we can bound the size of the set $|\mathcal{S}_k|$ by $|\mathcal{S}_k| \leq 2d(2k+1)^{d-1}$. In the proof of (3) we showed that $\mathbb{P}_\beta(\mathbf{0} \sim x) \leq \frac{\beta}{(\|x\|_\infty - 1)^{2d}}$. This already implies that for $k \geq 2$

$$\mathbb{P}_\beta(\mathbf{0} \sim \mathcal{S}_k) \leq \sum_{x \in \mathcal{S}_k} \mathbb{P}_\beta(\mathbf{0} \sim x) \stackrel{(7)}{\leq} \sum_{x \in \mathcal{S}_k} \frac{\beta}{(\|x\|_\infty - 1)^{2d}} \leq 2d(2k+1)^{d-1} \frac{\beta}{(k-1)^{2d}}$$

and thus also

$$\begin{aligned} \mathbb{P}_\beta(\mathbf{0} \sim \mathcal{S}_{\geq k}) &\leq \sum_{k'=k}^{\infty} \mathbb{P}_\beta(\mathbf{0} \sim \mathcal{S}_{k'}) \leq \sum_{k'=k}^{\infty} 2d(2k'+1)^{d-1} \frac{\beta}{(k'-1)^{2d}} \\ &\leq \sum_{k'=k}^{\infty} 2^d 3^d (k')^{d-1} \frac{\beta 2^{2d}}{(k')^{2d}} = \beta 2^d \sum_{k'=k}^{\infty} (k')^{-d-1} \leq \beta 50^d k^{-d}, \end{aligned}$$

which already proves (5). For $m \in \mathbb{N}$, a vertex $x \in V_{\mathbf{0}}^m$, and a box V_w^m with $\|w\|_\infty \geq 2$ we have for all $z \in V_w^m$ that $\|x - z\|_\infty \geq (\|w\|_\infty - 1)m$ and this implies

$$\mathbb{P}_\beta(x \sim V_w^m) \stackrel{(3)}{\leq} \sum_{z \in V_w^m} \frac{2^{2d} \beta}{\|x-z\|_\infty^{2d}} \leq \sum_{z \in V_w^m} \frac{2^{2d} \beta}{((\|w\|_\infty - 1)m)^{2d}} \leq \frac{\beta 4^{2d}}{\|w\|_\infty^{2d} m^d}$$

which shows (6). \square

We will often condition on the event that two blocks V_u^n, V_v^n are connected. So if we write X for the number of edges between them, we condition on the event $X \geq 1$. This conditioning clearly increases the expected number of edges between V_u^n and V_v^n , but by at most $+1$, as shown in the next Lemma.

Lemma 2.2. *Let $u, v \in \mathbb{Z}^d$ with $u \neq v$ and let X be the number of edges between the blocks V_u^n and V_v^n . Then for all $\beta > 0$*

$$\mathbb{E}_\beta [X | X \geq 1] \leq 1 + \mathbb{E}_\beta [X]$$

Proof. The random variable X is a sum of independent Bernoulli random variables and we prove the statement for all random variables of this type. We use the notation $X = \sum_{i=1}^m X_i$, where $m \in \mathbb{N}$, and $(X_i)_{i \in \{1, \dots, m\}}$ are independent Bernoulli random variables. For $i \in \{1, \dots, m\}$, let A_i be the event that $X_i = 1$ and $X_j = 0$ for all $j \in \{1, \dots, i-1\}$. As $\{X \geq 1\}$ implies that there is a first index i such that $X_i = 1$, we get that

$$\{X \geq 1\} = \bigsqcup_{i=1}^m A_i,$$

where the symbol \bigsqcup means a disjoint union. On the event A_i , we know that all the random variables X_j with $j < i$ equal 0, but we have no information about random variables X_j with $j > i$. Thus we get that

$$\frac{\mathbb{E}_\beta [X \mathbb{1}_{A_i}]}{\mathbb{P}_\beta (A_i)} = \mathbb{E}_\beta [X | A_i] = \mathbb{E}_\beta \left[1 + \sum_{j=i+1}^m X_j \middle| A_i \right] = 1 + \mathbb{E}_\beta \left[\sum_{j=i+1}^m X_j \right] \leq 1 + \mathbb{E}_\beta [X]$$

and using this inequality we get that

$$\begin{aligned} \mathbb{E}_\beta [X | X \geq 1] &= \frac{\mathbb{E}_\beta [X \mathbb{1}_{\{X \geq 1\}}]}{\mathbb{P}_\beta (X \geq 1)} = \frac{\sum_{i=1}^m \mathbb{E}_\beta [X \mathbb{1}_{A_i}]}{\mathbb{P}_\beta (X \geq 1)} \\ &\leq \frac{\sum_{i=1}^m \mathbb{P}_\beta (A_i) (1 + \mathbb{E}_\beta [X])}{\mathbb{P}_\beta (X \geq 1)} = 1 + \mathbb{E}_\beta [X]. \end{aligned}$$

\square

2.2 Submultiplicativity and the upper bound in Theorem 1.2

In this section, we prove the submultiplicative structure in the model in Lemma 2.3. This allows us to define the distance growth exponent $\theta(\beta)$ and also helps to prove the upper bound on $\theta(\beta)$ in Theorem 1.2.

Lemma 2.3. *For all dimensions d and all $\beta \geq 0$ the sequence*

$$\Lambda(n) = \Lambda(n, \beta) := \max_{u, v \in \{0, \dots, n-1\}^d} \mathbb{E}_\beta [d_{V_0^n}(u, v)] + 1 \quad (8)$$

is submultiplicative and furthermore one has

$$\theta(\beta) = \inf_{n \geq 2} \frac{\log(\Lambda(n, \beta))}{\log(n)} .$$

Proof. We show the first claim in Lemma 2.3 using a renormalization argument. As before we define $V_u^n = \prod_{i=1}^d \{p_i(u)n, \dots, (p_i(u) + 1)n - 1\}$. The graph G' obtained by identifying all the vertices in V_u^n to one vertex $r(u)$ has the same connection probabilities as the original model. For $x, y \in \{0, \dots, mn - 1\}^d$, say with $x \in V_u^n$ and $y \in V_w^n$, we create a path from x to y as follows. First we consider the shortest path $\mathcal{P} = (r(u_0) = r(u), r(u_1), \dots, r(u_{l-1}), r(u_l) = r(w))$ from $r(u)$ to $r(w)$, where $l = d_{G'}(r(u), r(w))$ is the distance between $r(u)$ and $r(w)$ in the renormalized model. Inside $V_{u_i}^n$ we first fix two vertices z and v such that $z \sim V_{u_{i-1}}^n$ and $v \sim V_{u_{i+1}}^n$; for $i = 0$ set $z = x$ and for $i = l$ set $v = y$. In case there are several such vertices z and v , we choose the one with smallest coordinates, where we weigh the coordinates in decreasing order (any deterministic rule that does not depend on the environment would work here). For each i , there clearly is a path between z and v that is completely inside $V_{u_i}^n$. As no information has been revealed up to now about the edges with both endpoints inside $V_{u_i}^n$ the expected distance between v and z inside $V_{u_i}^n$ is at most

$$\max_{a, b \in V_{u_i}^n} \mathbb{E}_\beta \left[d_{V_{u_i}^n}(a, b) \right] = \Lambda(n, \beta) - 1 .$$

Now we glue all these paths together to get a path from x to y . To bound the total distance between x and y note that we have $l + 1$ sets $V_{u_i}^n$ in which we need to find a path between two vertices. Additionally, we need to add l for the steps that we make from $V_{u_i}^n$ to $V_{u_{i+1}}^n$. As the connections between the different vertices in the renormalized model G' and the connections between vertices inside $V_{u_i}^n$ are independent we obtain that

$$\begin{aligned} & \mathbb{E}_\beta \left[d_{V_0^{mn}}(x, y) \right] \\ & \leq (\mathbb{E}_\beta [d_{G'}(v_u, v_w)] + 1) \max_{a, b \in \{0, \dots, n-1\}^d} \mathbb{E}_\beta \left[d_{V_0^n}(a, b) \right] + \mathbb{E}_\beta [d_{G'}(r(u), r(w))] \\ & = (\mathbb{E}_\beta [d_{G'}(r(u), r(w))] + 1) \left(\max_{a, b \in \{0, \dots, n-1\}^d} \mathbb{E}_\beta \left[d_{V_0^n}(a, b) \right] + 1 \right) - 1 \\ & \leq \Lambda(m)\Lambda(n) - 1 , \end{aligned}$$

and as $x, y \in \{0, \dots, nm - 1\}^d$ were arbitrary we obtain

$$\Lambda(mn) \leq \Lambda(m)\Lambda(n) . \tag{9}$$

As the sequence is submultiplicative we can define

$$\theta = \theta(\beta) = \lim_{k \rightarrow \infty} \frac{\log(\Lambda(2^k, \beta))}{\log(2^k)} .$$

Actually, this limit exists not just along dyadic points of the form 2^k , for $k \in \mathbb{N}$, but even

when taking a limit along the integers, i.e.,

$$\theta = \theta(\beta) = \lim_{n \rightarrow \infty} \frac{\log(\Lambda(n, \beta))}{\log(n)},$$

which can be seen with Lemma 4.1 below. As a next step, we want to show that $\Lambda(n) \geq n^\theta$ for all n . We do this using a proof by contradiction. So assume the contrary, i.e., there exists a natural number N and a $c < 1$ with $\Lambda(N) = cN^\theta$. Using (9) we get that for every integer k

$$\Lambda(N^k) \leq \Lambda(N)^k = c^k N^{\theta k}$$

and thus

$$\theta = \lim_{k \rightarrow \infty} \frac{\log(\Lambda(N^k, \beta))}{\log(N^k)} \leq \limsup_{k \rightarrow \infty} \frac{\log(c^k N^{\theta k})}{\log(N^k)} = \frac{\log(c) + \theta \log(N)}{\log(N)} < \theta$$

which is a contradiction. Knowing this already gives us that for all positive numbers K we have

$$\theta = \lim_{n \rightarrow \infty} \frac{\log(\Lambda(n))}{\log(n)} = \inf_{n \geq 2} \frac{\log(\Lambda(n))}{\log(n)} = \inf_{n \geq K} \frac{\log(\Lambda(n))}{\log(n)}. \quad (10)$$

□

This lemma and its proof already have several interesting applications. First, we emphasize that $\Lambda(mn, \beta) \geq \Lambda(n, \beta)$ for all $m, n \in \mathbb{N}_{>0}$. This holds, as for arbitrary $x, y \in V_{\mathbf{0}}^n$, the distance $d_{V_{\mathbf{0}}^{mn}}(u, v)$ between $u \in V_x^m$ and $v \in V_y^m$ is at least the distance between $r(x)$ and $r(y)$ in G' . Using the coupling with the continuous model and taking expectations we thus get that

$$\mathbb{E}_\beta [d_{V_{\mathbf{0}}^{mn}}(u, v)] \geq \mathbb{E}_\beta [d_{G'}(r(x), r(y))] = \mathbb{E}_\beta [d_{V_{\mathbf{0}}^n}(x, y)]$$

which shows our claim. For $n = 3$ we have for all $u, v \in \{0, 1, 2\}^d$ with $u \neq v$ that

$$\mathbb{E}_\beta [d_{[0,2]^d}(u, v)] = 1 \cdot \mathbb{P}_\beta(u \sim v) + 2 \cdot \mathbb{P}_\beta(u \not\sim v) < 2$$

and this already implies that $\Lambda(3) =: 3^{\theta'} < 3$ for some $\theta' = \theta'(3) < 1$. Inductively, with a renormalization at scale 3, we get that

$$\Lambda(3^k N) \leq \Lambda(3)^k \Lambda(N) = 3^{k\theta'} \Lambda(N) \quad (11)$$

for all $k, N \in \mathbb{N}_{>0}$. This inequality already gives the upper bound in expectation for $s = 2d$, that was already observed in [16] with a very similar technique. Next, we do a renormalization at scale $\sqrt[2d]{\beta}$ instead of scale 3 in order to get the inverse logarithmic upper bound stated in Theorem 1.2.

Proof of the upper bound in Theorem 1.2. We want to show that there exists a constant $C < \infty$ such that for all $\beta \geq 2$

$$\theta(\beta) \leq \frac{C}{\log(\beta)}.$$

As the connection probability between two vertices $\mathbb{P}_\beta(u \sim v)$ is increasing in β the

distance exponent $\theta : \mathbb{R}_{\geq 0} \rightarrow [0, 1]$ is clearly decreasing by the *Harris coupling*, see for example [28]. Thus it suffices to show the upper bound for β large enough with $\sqrt[2d]{\beta} \in \mathbb{N}$. For such a β and all $u, v \in \{0, \dots, \sqrt[2d]{\beta} - 1\}^d$ we have for all $y \in u + \mathcal{C}$ and $x \in v + \mathcal{C}$ that

$$\|x - y\|^{2d} \leq d^{2d} \|x - y\|_{\infty}^{2d} \leq d^{2d} \sqrt[2d]{\beta}^{2d} = d^{2d} \beta \quad (12)$$

and this already implies

$$\int_{u+\mathcal{C}} \int_{v+\mathcal{C}} \frac{1}{\|x - y\|^{2d}} dx dy \geq \frac{1}{d^{2d} \beta}.$$

Inserting this into the definition $p(\beta, \{u, v\})$ and using that $1 - e^{-x} \geq \frac{x}{2}$ for all $x \leq 1$ we get that for large enough β that satisfy $\frac{1}{d^{2d} \beta} \leq 1$ we already have

$$\mathbb{P}_{\beta}(u \sim v) = 1 - e^{-\int_{u+\mathcal{C}} \int_{v+\mathcal{C}} \frac{1}{\|x-y\|^{2d}} dx dy} \stackrel{(12)}{\geq} 1 - e^{-d^{-2d}} \geq \frac{1}{2} d^{-2d} \geq (2d)^{-2d} \quad (13)$$

for all $u, v \in \{0, \dots, \sqrt[2d]{\beta} - 1\}^d$. Now let us bound the expected graph distance between u and v . We do this by comparing the distance to a geometric random variable. Let $(u = u_0, u_1, \dots, u_k = v)$ be a deterministic path between u and v inside $V_{\mathbf{0}}^{\sqrt[2d]{\beta}}$ with $\|u_i - u_{i-1}\|_{\infty} = 1$ for all $i \in \{1, \dots, k\}$ and $k \leq \sqrt[2d]{\beta}$. Starting from this, we build a shorter path between u and v as follows. We start at $u_0 = u$. Then for all i , if $u_i \sim v$, directly go to v . If $u_i \not\sim v$, then go to u_{i+1} . This gives a path P between u and v and for $l \in \{1, \dots, k\}$ this path has length of at least l if and only if $u_i \not\sim v$ for all $i \in \{0, \dots, l-2\}$. As the connections between v and different u_i -s are independent we get that

$$\begin{aligned} \mathbb{E}_{\beta} \left[d_{V_{\mathbf{0}}^{\sqrt[2d]{\beta}}}(u, v) \right] &= \sum_{l=1}^k \mathbb{P}_{\beta} \left(d_{V_{\mathbf{0}}^{\sqrt[2d]{\beta}}}(u, v) \geq l \right) \leq \sum_{l=1}^k \mathbb{P}_{\beta} (u_i \not\sim v \text{ for all } i \leq l-2) \\ &\stackrel{(13)}{\leq} \sum_{l=1}^k (1 - (2d)^{-2d})^{l-1} \leq \frac{1}{1 - (1 - (2d)^{-2d})} = (2d)^{2d}. \end{aligned}$$

This already implies that $\Lambda(\sqrt[2d]{\beta}, \beta) \leq (2d)^{2d} + 1 \leq (3d)^{2d}$. Applying the submultiplicativity of Λ iteratively we get that

$$\begin{aligned} \theta(\beta) &= \lim_{k \rightarrow \infty} \frac{\log \left(\Lambda \left(\sqrt[2d]{\beta^k}, \beta \right) \right)}{\log \left(\sqrt[2d]{\beta^k} \right)} \leq \limsup_{k \rightarrow \infty} \frac{\log \left(\Lambda \left(\sqrt[2d]{\beta}, \beta \right)^k \right)}{\log \left(\sqrt[2d]{\beta^k} \right)} \\ &= \frac{\log \left(\Lambda \left(\sqrt[2d]{\beta}, \beta \right) \right)}{\log \left(\sqrt[2d]{\beta} \right)} \leq \frac{\log \left((3d)^{2d} \right)}{\frac{1}{2d} \log(\beta)} = \frac{4d^2 \log(3d)}{\log(\beta)} \end{aligned}$$

which finishes the proof. \square

2.3 Spacing between points with long bonds

In this section, we investigate certain geometric properties of the cluster inside certain boxes. Mostly, we want to get upper bounds on the probability that a vertex is connected to two different long edges. As we will need it at a later point, namely in section 5.1, we will prove the statements for $\|x - y\|_{\infty} \leq 1$ instead of $x = y$. This does not cause major difficulties, as for each point $x \in \mathbb{Z}^d$, there are only 3^d many points $y \in \mathbb{Z}^d$ with

$\|x - y\|_\infty \leq 1$. We start showing that the probability that two near vertices x, y with $\|x - y\|_\infty \leq 1$ are both connected to far away boxes is very low.

Lemma 2.4. *For blocks V_u^m, V_v^m, V_w^m with $\|u - v\|_\infty, \|v - w\|_\infty \geq 2$ there exists a constant $C_d < \infty$ such that for all $\beta \geq 0$*

$$\mathbb{P}_\beta(\exists x, y \in V_v^m : \|x - y\|_\infty \leq 1, x \sim V_u^m, y \sim V_w^m) \leq \frac{C_d \beta^2}{\|u - v\|_\infty^{2d} \|w - v\|_\infty^{2d} m^d}.$$

Proof. By translational invariance we can assume that $v = \mathbf{0}$, and thus $\|u\|_\infty, \|w\|_\infty \geq 2$. For each $x \in V_{\mathbf{0}}^m$ there are at most 3^d vertices $y \in V_{\mathbf{0}}^m$ with $\|x - y\|_\infty \leq 1$. For each vertex $y \in V_{\mathbf{0}}^m$ the probability that $y \sim V_w^m$ or $x \sim V_u^m$ is bounded by $\frac{\beta 4^{2d}}{\|w\|_\infty^{2d} m^d}$, respectively by $\frac{\beta 4^{2d}}{\|u\|_\infty^{2d} m^d}$, by (6). Thus

$$\begin{aligned} \mathbb{P}_\beta(\exists x, y \in V_{\mathbf{0}}^m : \|x - y\|_\infty \leq 1, x \sim V_u^m, y \sim V_w^m) &\leq \sum_{x \in V_{\mathbf{0}}^m} \mathbb{P}_\beta(x \sim V_u^m) 3^d \frac{\beta 4^{2d}}{\|w\|_\infty^{2d} m^d} \\ &= \frac{48^d \beta}{\|w\|_\infty^{2d} m^d} \sum_{x \in V_{\mathbf{0}}^m} \mathbb{P}_\beta(x \sim V_u^m) \stackrel{(6)}{\leq} \frac{48^d \beta}{\|w\|_\infty^{2d} m^d} \frac{\beta 4^{2d}}{\|u\|_\infty^{2d} m^d} \leq \frac{\beta^2 1000^d}{\|w\|_\infty^{2d} \|u\|_\infty^{2d} m^d} \end{aligned}$$

which finishes the proof. \square

Lemma 2.5. *For blocks V_u^m, V_v^m, V_w^m with $\|v - w\|_\infty \geq 2$ and $\|u - v\|_\infty = 1$ there exists a constant $C_d < \infty$ such that for all $\beta \geq 0$*

$$\mathbb{P}_\beta(\exists x, y \in V_v^m : \|x - y\|_\infty \leq 1, x \sim V_u^m, y \sim V_w^m) \leq \begin{cases} \frac{C_d \beta [\beta] \log(m)}{\|v - w\|_\infty^{2d} m} & \text{for } d = 1 \\ \frac{C_d \beta [\beta]}{\|v - w\|_\infty^{2d} m} & \text{for } d \geq 2 \end{cases}.$$

Proof. By translational invariance we can assume that $v = \mathbf{0}$, and thus $\|u\|_\infty = 1, \|w\|_\infty \geq 2$. For each $x \in V_{\mathbf{0}}^m$ there are at most 3^d vertices $y \in V_{\mathbf{0}}^m$ with $\|x - y\|_\infty \leq 1$. For each vertex $y \in V_{\mathbf{0}}^m$ the probability that $y \sim V_w^m$ is bounded by $\frac{\beta 4^{2d}}{\|w\|_\infty^{2d} m^d}$ by (6). Thus

$$\begin{aligned} \mathbb{P}_\beta(\exists x, y \in V_{\mathbf{0}}^m : \|x - y\|_\infty \leq 1, x \sim V_u^m, y \sim V_w^m) &\leq \sum_{x \in V_{\mathbf{0}}^m} \mathbb{P}_\beta(x \sim V_u^m) 3^d \frac{\beta 4^{2d}}{\|w\|_\infty^{2d} m^d} \\ &= \frac{48^d \beta}{\|w\|_\infty^{2d} m^d} \sum_{x \in V_{\mathbf{0}}^m} \mathbb{P}_\beta(x \sim V_u^m). \end{aligned} \quad (14)$$

As $\|u\|_\infty = 1$ we have $d_\infty(x, V_u^m) \leq m$ for all $x \in V_{\mathbf{0}}^m$, where d_∞ is the distance with respect to the ∞ -norm. We furthermore have the inequality

$$|\{x \in V_{\mathbf{0}}^m : d_\infty(x, V_u^m) = k\}| \leq 6^d m^{d-1}$$

for all $k \in \mathbb{N}$. This is clear for $k > m$, as the relevant set is empty in this case. For $k \leq m$ the set $\{x \in \mathbb{Z}^d : d_\infty(x, V_u^m) = k\}$ is just the boundary of the box

$$\prod_{i=1}^d \{p_i(u)m - k, \dots, (p_i(u) + 1)m - 1 + k\},$$

which is a box of side length $m + 2k \leq 3m$. Thus the boundary has a cardinality of at

most $2d(3m)^{d-1} \leq 6^d m^{d-1}$. Using this observation we get that

$$\begin{aligned} \sum_{x \in V_0^m} \mathbb{P}_\beta(x \sim V_u^m) &= \sum_{k=1}^m \sum_{\substack{x \in V_0^m: \\ d_\infty(x, V_u^m)=k}} \mathbb{P}_\beta(x \sim V_u^m) \leq \sum_{k=1}^m 6^d m^{d-1} \mathbb{P}_\beta(x \sim x + \mathcal{S}_{\geq k}) \\ &\stackrel{(5)}{\leq} 6^d m^{d-1} + 6^d m^{d-1} \sum_{k=2}^m \beta 50^d k^{-d} \leq \begin{cases} m^{d-1} \log(m) \lceil \beta \rceil 400^d & \text{for } d = 1 \\ m^{d-1} \lceil \beta \rceil 400^d & \text{for } d \geq 2 \end{cases}. \end{aligned} \quad (15)$$

Inserting this into (14) we get that

$$\begin{aligned} \mathbb{P}_\beta(\exists x, y \in V_0^m : \|x - y\|_\infty \leq 1, x \sim V_u^m, y \sim V_w^m) &\leq \sum_{x \in V_0^m} \mathbb{P}_\beta(x \sim V_u^m) 3^d \frac{\beta 4^{2d}}{\|w\|_\infty^{2d} m^d} \\ &\leq \begin{cases} \frac{20000^d \beta \lceil \beta \rceil \log(m)}{\|w\|_\infty^{2d} m} & \text{for } d = 1 \\ \frac{20000^d \beta \lceil \beta \rceil}{\|w\|_\infty^{2d} m} & \text{for } d \geq 1 \end{cases} \end{aligned}$$

which finishes the proof. \square

Lemma 2.6. *Let $m \in \mathbb{N}, l \in \{1, \dots, 3^d - 1\}$, and let $v_0, v_1, \dots, v_{l+1} \in \mathbb{Z}^d$ be distinct with $\|v_{i+1} - v_i\|_\infty = 1$ for all $i \in \{0, \dots, l\}$, $\|v_i - v_0\|_\infty = 1$ for all $i \in \{0, \dots, l\}$ and $\|v_{l+1} - v_0\|_\infty = 2$. Then there exists a constant $C_d < \infty$ such that the two probabilities*

$$\begin{aligned} &\mathbb{P}_\beta\left(\exists i \in \{1, \dots, l\} \exists x, y \in V_{v_i}^m \text{ with } \|x - y\|_\infty \leq 1, x \sim V_{v_{i-1}}^m, y \sim V_{v_{i+1}}^m \cap \left(y + \mathcal{S}_{\geq \frac{m}{6^d}}\right)\right), \\ &\mathbb{P}_\beta\left(\exists i \in \{1, \dots, l\} \exists x, y \in V_{v_i}^m \text{ with } \|x - y\|_\infty \leq 1, x \sim V_{v_{i+1}}^m, y \sim V_{v_{i-1}}^m \cap \left(y + \mathcal{S}_{\geq \frac{m}{6^d}}\right)\right) \end{aligned}$$

are both bounded by $\frac{C_d \beta \lceil \beta \rceil \log(m)}{m}$ for $d = 1$ and by $\frac{C_d \beta \lceil \beta \rceil}{m}$ for $d \geq 2$.

Proof. By a union bound we have that

$$\begin{aligned} &\mathbb{P}_\beta\left(\exists i \in \{1, \dots, l\} \exists x, y \in V_{v_i}^m \text{ with } \|x - y\|_\infty \leq 1, x \sim V_{v_{i-1}}^m, y \sim V_{v_{i+1}}^m \cap \left(y + \mathcal{S}_{\geq \frac{m}{6^d}}\right)\right) \\ &\leq \sum_{i \in \{1, \dots, l\}} \sum_{\substack{x, y \in V_{v_i}^m: \\ \|x - y\|_\infty \leq 1}} \mathbb{P}_\beta\left(x \sim V_{v_{i-1}}^m, y \sim V_{v_{i+1}}^m \cap \left(y + \mathcal{S}_{\geq \frac{m}{6^d}}\right)\right) \\ &\leq \sum_{i \in \{1, \dots, l\}} \sum_{\substack{x, y \in V_{v_i}^m: \\ \|x - y\|_\infty \leq 1}} \mathbb{P}_\beta\left(x \sim V_{v_{i-1}}^m\right) \mathbb{P}_\beta\left(y \sim \left(y + \mathcal{S}_{\geq \frac{m}{6^d}}\right)\right) \\ &\stackrel{(5)}{\leq} \beta 50^d \left(\frac{m}{6^d}\right)^{-d} \sum_{i \in \{1, \dots, l\}} \sum_{\substack{x, y \in V_{v_i}^m: \\ \|x - y\|_\infty \leq 1}} \mathbb{P}_\beta\left(x \sim V_{v_{i-1}}^m\right) \\ &\leq \beta 150^d \left(\frac{m}{6^d}\right)^{-d} \sum_{i \in \{1, \dots, l\}} \sum_{x \in V_{v_i}^m} \mathbb{P}_\beta\left(x \sim V_{v_{i-1}}^m\right). \end{aligned}$$

The sum $\sum_{x \in V_{v_i}^m} \mathbb{P}_\beta\left(x \sim V_{v_{i-1}}^m\right)$ was already upper bounded in (15). Using this upper bound, $l \leq 3^d$, and inserting this into the line above we get that

$$\beta 150^d \left(\frac{m}{6^d}\right)^{-d} \sum_{i \in \{1, \dots, l\}} \sum_{x \in V_{v_i}^m} \mathbb{P}_\beta\left(x \sim V_{v_{i-1}}^m\right) \leq \begin{cases} \frac{\beta \lceil \beta \rceil (6^d 10^6)^d \log(m)}{m} & \text{for } d = 1 \\ \frac{\beta \lceil \beta \rceil (6^d 10^6)^d}{m} & \text{for } d \geq 2 \end{cases}$$

which finishes the proof for the first item in the statement of the lemma. The estimate for the second term works analogously. \square

2.4 The lower bound in Theorem 1.2

Finally, we developed all the necessary techniques in order to show the lower bound in Theorem 1.2, i.e., that there for all dimensions d , there exists a constant $c > 0$ such that $\theta(\beta) \geq \frac{c}{\log(\beta)}$ for all $\beta \geq 2$.

Proof of the lower bound in Theorem 1.2. Inequality (4) and Lemma 2.4 show that for all dimensions d there exists a constant $C_d < \infty$ such that for all $\beta \geq 2$ and all u, v, w with $\|u - v\|_\infty, \|v - w\|_\infty \geq 2$

$$\mathbb{P}_\beta (\exists x \in V_v^m : x \sim V_u^m, x \sim V_w^m \mid V_u^m \sim V_v^m \sim V_w^m) \leq \frac{C_d \beta^2}{m^{1/2}}. \quad (16)$$

Analogously, Lemma 2.5 shows that there exists a constant $C_d < \infty$ such that for all $\beta \geq 2$ and all u, v, w with $\|u - v\|_\infty \geq 2$ and $\|v - w\|_\infty = 1$

$$\mathbb{P}_\beta (\exists x \in V_v^m : x \sim V_u^m, x \sim V_w^m \mid V_u^m \sim V_v^m \sim V_w^m) \leq \frac{C_d \beta^2}{m^{1/2}} \quad (17)$$

where we also used that $\frac{\log(m)}{m} = \mathcal{O}(m^{1/2})$. Lemma 2.6 shows that for $l \in \{1, \dots, 3^d - 1\}$ and $v_0, v_1, \dots, v_{l+1} \in \mathbb{Z}^d$ distinct with $\|v_{i+1} - v_i\|_\infty = 1$ for all $i \in \{0, \dots, l\}$, $\|v_i - v_0\|_\infty = 1$ for all $i \in \{0, \dots, l\}$ and $\|v_{l+1} - v_0\|_\infty = 2$ one has the bound

$$\mathbb{P}_\beta (\exists x_1, \dots, x_l : x_i \in V_{v_i}^m, V_{v_0}^m \sim x_1 \sim x_2 \sim \dots \sim x_l \sim V_{v_{l+1}}^m) \leq \frac{C_d \beta^2}{m^{1/2}} \quad (18)$$

as a path from V_{v_0} to $V_{v_{l+1}}$ in $l + 1 \leq 3^d$ steps contains at least one edge $\{x_i, x_{i+1}\}$ with $\|x_i - x_{i+1}\|_\infty \geq \frac{m}{3^d}$ and thus $x_i \sim x_{i+1} + \mathcal{S}_{\geq \frac{m}{3^d}}$ in this case. We will now show that

$$\mathbb{E}_\beta [d_{V_{\mathbf{0}}^{mM}}(\mathbf{0}, (mM - 1)e_1)] \geq \left(1 + \frac{1}{3^{d+4}}\right) \mathbb{E}_\beta [d_{V_{\mathbf{0}}^M}(\mathbf{0}, (M - 1)e_1)] \quad (19)$$

for $m \geq (2000 \cdot \lceil \beta \rceil^{3 \cdot 5^d} C_d)^{(3^{4d})}$ and all large enough M . We will see later where this condition on m comes from. To see (19), we use a renormalization. For $u \in V_{\mathbf{0}}^M$ we identify the blocks V_u^m to vertices $r(u)$ and call the resulting graph G' . Then we will prove that

$$\mathbb{E}_\beta [d_{V_{\mathbf{0}}^{mM}}(\mathbf{0}, (mM - 1)e_1)] \geq \left(1 + \frac{1}{3^{d+4}}\right) \mathbb{E}_\beta [d_{G'}(r(\mathbf{0}), r((M - 1)e_1))]$$

for large enough M . This implies (19), as the random graphs G' and $V_{\mathbf{0}}^M$ have the same distribution. Now we condition on the graph G' , i.e., we already have the knowledge which blocks of the form V_u^m are connected in the original graph. Let $P' = (r(v_0), \dots, r(v_k))$ be a self-avoiding path in G' starting at the origin vertex, i.e., $v_0 = \mathbf{0}$. In this proof we will always assume that $k \geq 3^{d+3}$. Let $l = \lfloor \frac{k}{3^{d+1}} \rfloor$. For $j \in \{0, \dots, l\}$ we call the subsequence $R_j := (r(v_{2j \cdot 3^d}), r(v_{(2j+1) \cdot 3^d}), \dots, r(v_{(2j+2) \cdot 3^d}))$ *separated* if there does *not* exist a sequence

$(x_i)_{i=2j3^d+1}^{(2j+2) \cdot 3^d - 1}$ such that $x_i \in V_{v_i}^m$ for all $i \in \{2j3^d + 1, (2j+2) \cdot 3^d - 1\}$ and

$$V_{v_{j3^d}}^m \sim x_{j3^d+1} \sim x_{j3^d+2} \sim \dots \sim x_{(j+2)3^d-1} \sim V_{v_{(j+2)3^d}}^m .$$

For a given self-avoiding path $P' \subset G'$ and different values of $j \in \{0, \dots, l\}$ it is independent whether the subsequences $(r(v_{2j3^d}), r(v_{2j3^d+1}), \dots, r(v_{(2j+2)3^d}))$ are separated, and the probability that a specific subsequence $(r(v_{2j3^d}), r(v_{2j3^d+1}), \dots, r(v_{(2j+2)3^d}))$ is not separated is bounded by $\frac{C_d \beta^2}{m^{1/2}}$, as for every sequence $(v_{2j3^d}, v_{2j3^d+1}, \dots, v_{(2j+2)3^d})$ at least one of the situations of (16), (17) or (18) holds, as we will shortly show now. If none of the situations in (16), (17) holds, then the path makes only nearest neighbor-jumps within the subsequence R_j . However, as that there are only $3^d - 1$ many points $v \in \mathbb{Z}^d$ with $\|v - v_{2j3^d}\|_\infty = 1$, the situation of (18) must occur within the subsequence R_j .

In a separated subsequence, the walk on the original graph V_0^{mM} needs to take at least one additional step. For a fixed path $P' \subset G'$ of length k and $l = \lfloor \frac{k}{3^{d+1}} \rfloor$ we have that

$$\begin{aligned} & \mathbb{P}_\beta \left(|\{j \in \{0, \dots, l\} : R_j \text{ not separated}\}| > \frac{l}{2} \mid G' \right) \\ &= \mathbb{P}_\beta \left(\bigcup_{\substack{U \subset \{0, \dots, l\} \\ |U| > l/2}} \{R_j \text{ not separated for all } j \in U\} \mid G' \right) \leq 2^l \left(\frac{C_d \beta^2}{m^{1/2}} \right)^{l/2} . \end{aligned}$$

Next, we want to bound the expected degree of vertices in the long-range percolation graph from above. With the bound on the connection probability $\mathbb{P}_\beta(0 \sim u)$ (3) we get that

$$\begin{aligned} \mathbb{E}_\beta [\deg(\mathbf{0})] &= \sum_{u \in \mathbb{Z}^d \setminus \{0\}} \mathbb{P}_\beta(\mathbf{0} \sim u) \leq 3^d + \sum_{k=2}^{\infty} \sum_{u \in S_k} \frac{2^{2d} \beta}{k^{2d}} \leq 3^d + \sum_{k=2}^{\infty} 2d(2k+1)^{d-1} \frac{2^{2d} \beta}{k^{2d}} \\ &\leq 3^d + \beta 2^{3d} 3^d \sum_{k=2}^{\infty} k^{-d-1} \leq 3^d + \beta 3^{4d} \leq \lceil \beta \rceil 3^{5d} . \end{aligned} \quad (20)$$

Let \mathcal{P}'_k be the set of self-avoiding paths in G' starting at $r(\mathbf{0})$. With a comparison to the case of a Galton-Watson tree inequality (20) already gives that $\mathbb{E}_\beta [|\mathcal{P}'_k|] \leq (\lceil \beta \rceil 3^{5d})^k$. As $\lfloor \frac{k}{3^{d+2}} \rfloor \leq \frac{\lfloor \frac{k}{3^{d+1}} \rfloor}{2}$ we see that

$$\begin{aligned} & \mathbb{P}_\beta \left(\exists P' \in \mathcal{P}'_k \text{ with less than } \lfloor \frac{k}{3^{d+2}} \rfloor \text{ separated subpaths } R_j \right) \\ &= \mathbb{E}_\beta \left[\mathbb{P}_\beta \left(\exists P' \in \mathcal{P}'_k \text{ with less than } \lfloor \frac{k}{3^{d+2}} \rfloor \text{ separated subpaths } R_j \mid G' \right) \right] \\ &\leq \mathbb{E}_\beta \left[|\mathcal{P}'_k| 2^k \left(\frac{C_d \beta^2}{m^{1/2}} \right)^{\lfloor \frac{k}{3^{d+2}} \rfloor} \right] \leq (\lceil \beta \rceil 3^{5d})^k 2^k \left(\frac{C_d \beta^2}{m^{1/2}} \right)^{\lfloor \frac{k}{3^{d+2}} \rfloor} \\ &\leq (\lceil \beta \rceil 3^{5d})^k 2^k C_d^k \frac{1}{m^{\frac{k}{3^{4d}}}} \leq 0.01^k \end{aligned}$$

by the choice of $m \geq (2000 \cdot 3^{5d} C_d \lceil \beta \rceil^3)^{(3^{4d})}$. Next, we want to translate this bound on the probability of certain events to bounds on the expectation of the distances. For

this, let \mathcal{G}_k be the event that all self-avoiding paths $P' \subset G'$ starting at the origin and of length $\tilde{k} \geq k$ contain at least $\lfloor \frac{\tilde{k}}{3^{d+2}} \rfloor$ separated subpaths R_j . With the preceding inequality we directly get $\mathbb{P}_\beta(\mathcal{G}_k) \geq 1 - 0.1^k$. On the event \mathcal{G}_k , each path $P \subset V_{\mathbf{0}}^{mM}$ for which the loop-erased projection on G' goes through $\tilde{k} + 1$ different blocks of the form V_u^m , needs to have a length of at least $\tilde{k} + \lfloor \frac{\tilde{k}}{3^{d+2}} \rfloor \geq (1 + \frac{1}{3^{d+3}}) \tilde{k}$. Furthermore, if we have $d_{G'}(r(\mathbf{0}), r((M-1)e_1)) = \tilde{k}$, then the loop-erased projection of every path connecting $\mathbf{0}$ to $(mM-1)e_1$ into the original model goes through at least $\tilde{k} + 1$ different blocks of the form V_u^m . So we get that

$$\begin{aligned}
\mathbb{E}_\beta \left[d_{V_{\mathbf{0}}^{mM}}(\mathbf{0}, (mM-1)e_1) \right] &\geq \sum_{k=3^{d+3}}^{\infty} \mathbb{E}_\beta \left[d_{V_{\mathbf{0}}^{mM}}(\mathbf{0}, (mM-1)e_1) \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((M-1)e_1))=k\}} \right] \\
&\geq \sum_{k=3^{d+3}}^{\infty} \mathbb{E}_\beta \left[d_{V_{\mathbf{0}}^{mM}}(\mathbf{0}, (mM-1)e_1) \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((M-1)e_1))=k\}} \mathbb{1}_{\mathcal{G}_k} \right] \\
&\geq \sum_{k=3^{d+3}}^{\infty} \mathbb{E}_\beta \left[k \left(1 + \frac{1}{3^{d+3}} \right) \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((M-1)e_1))=k\}} \mathbb{1}_{\mathcal{G}_k} \right] \\
&\geq \sum_{k=3^{d+3}}^{\infty} \left(\mathbb{E}_\beta \left[k \left(1 + \frac{1}{3^{d+3}} \right) \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((M-1)e_1))=k\}} \right] - \mathbb{E}_\beta \left[2k \mathbb{1}_{\mathcal{G}_k^c} \right] \right) \\
&\geq \left(1 + \frac{1}{3^{d+3}} \right) \sum_{k=3^{d+3}}^{\infty} k \mathbb{E}_\beta \left[\mathbb{1}_{\{d_{V_{\mathbf{0}}^M}(\mathbf{0}, (M-1)e_1)=k\}} \right] - 2 \sum_{k=1}^{\infty} 0.1^k k \\
&\geq \left(1 + \frac{1}{3^{d+3}} \right) \mathbb{E}_\beta \left[d_{V_{\mathbf{0}}^M}(\mathbf{0}, (M-1)e_1) \right] - 3^{d+4} \mathbb{P}_\beta \left(d_{V_{\mathbf{0}}^M}(\mathbf{0}, (M-1)e_1) < 3^{d+3} \right) - 1 \\
&\geq \left(1 + \frac{1}{3^{d+4}} \right) \mathbb{E}_\beta \left[d_{V_{\mathbf{0}}^M}(\mathbf{0}, (M-1)e_1) \right]
\end{aligned}$$

where the last inequality holds for all large enough M , as the probability of the event $\{d_{V_{\mathbf{0}}^M}(\mathbf{0}, (M-1)e_1) < 3^{d+3}\}$ tends to 0 as $M \rightarrow \infty$. Say that it holds for all $M \geq m^N$ where $m = (2000 \cdot 3^{5d} C_d \lceil \beta \rceil^3)^{(3^{d+4})}$. The important property about the choice of m is, that its size is polynomial in β . This already implies that

$$\begin{aligned}
\theta(\beta) &\geq \lim_{n \rightarrow \infty} \frac{\log \left(\mathbb{E}_\beta \left[d_{V_{\mathbf{0}}^{m^n}}(\mathbf{0}, (m^n-1)e_1) \right] \right)}{\log(m^n)} \geq \lim_{n \rightarrow \infty} \frac{\log \left(\left(1 + \frac{1}{3^{d+4}} \right)^{n-N} \right)}{\log(m^n)} \\
&= \frac{\log \left(1 + \frac{1}{3^{d+4}} \right)}{\log(m)} \geq \frac{c}{\log(\beta)}
\end{aligned}$$

for some small $c > 0$ and all $\beta \geq 2$. \square

3 Connected sets in graphs

The expected number of open paths in the long range percolation model of length k , starting at $\mathbf{0}$, grows at most like $\mathbb{E}[\deg(\mathbf{0})]^k$, which can be easily proven by a comparison with a Galton-Watson tree. However, it is not clear how the number of connected subsets of \mathbb{Z}^d containing the origin grows. In particular, because the maximal degree of vertices is unbounded. In this chapter, we prove several results about the structure of connected sets in the long-range percolation graph. Mostly, we want to prove that with exponentially high probability in k , all connected sets of size k in the graph have not too many edges.

First, we need to define what we mean by a connected set. Formally, we define the a connected set as follows. For a graph $G = (V, E)$ we say that a subset $Z \subset V$ is *connected* if the graph (Z, E') with edge set $E' = \{\{x, y\} \in E : x, y \in Z\}$ is connected. As a first step, we bound the expected number of connected sets of certain size in Galton-Watson trees. This counting of connected sets plays an important role in section 5 and in the companion paper [4].

Lemma 3.1. *Let \mathcal{X} be a countable set with a total ordering and a minimal element, let X be a countable sum of independent Bernoulli-distributed random variables over this set, i.e., $X = \sum_{i \in \mathcal{X}} X_i$, and let μ be the expectation value of X . Say that $q(k) = \mathbb{P}(X_k = 1)$. Let T be a Galton-Watson tree with offspring distribution $\mathcal{L}(X)$. We denote the set of all open subtrees of size k containing the origin by \mathcal{T}_k . Then*

$$\mathbb{E}[|\mathcal{T}_k|] \leq 4^k \mu^k.$$

Proof. The choice of the set \mathcal{X} and the total ordering on it do not influence the outcome, so we will always work with $\mathcal{X} = \mathbb{N}$ from here on. We can think of the Galton-Watson tree as a model of independent bond percolation on the graph with vertex set $L = \bigcup_{n=0}^{\infty} L_n$ where $L_n = \mathbb{N}^n$ and with edge set $S = \{\{v, (v m)\} : v \in L, m \in \mathbb{N}\}$ where some edge of the form $\{v, (v m)\}$ is open with probability $q(m)$. Note that the graph $G = (L, S)$ is a tree, so in particular there exists a unique path from the origin \emptyset to every vertex. For a vertex v in the graph the number of open edges of the form $\{v, (v m)\}$ has the same law as X and thus we can identify the open cluster connected to the root \emptyset with a Galton-Watson tree with offspring distribution $\mathcal{L}(X)$. For a vertex $v \in L$, we call the vertices of the form $(v m)$ that are connected to v by an open bond the *children* of v . Vice versa, we say that v is the *parent* of the vertex $(v m)$, if $(v m)$ is connected to v . So in particular, the expected number of subtrees of the Galton-Watson tree of size k is the same as the expected number of connected sets of size k in (L, S) . For a connected set $L' \subset L$ of size k , we now describe an exploration process $(Y_i)_{i \in \{1, \dots, 2k-1\}}$ of it:

0. Start with $Y_1 = \emptyset$.
1. For $i = 1, \dots, 2k - 1$
 - (a) If there exists $m \in \mathbb{N}$ for which $(Y_i m) \in L'$ and $Y_j \neq (Y_i m)$ for all $j < i$, let m' be the minimal among those $m \in \mathbb{N}$ and set $Y_{i+1} = (Y_i m')$.
 - (b) If such an m does not exist, let Y_{i+1} be the parent of Y_i .

An example of this procedure is given in Figure 1. This exploration process traverses every edge exactly twice in opposite directions and starts and ends at the origin of the tree. We also say that the exploration process Y_i goes (one level) down if (a) occurs in the algorithm above and otherwise we say that the process goes (one level) up. We also define a different process $(Y'_i)_{i \in \{1, \dots, k\}}$, where Y'_i is the unique point Y_l such that $|\{Y_1, \dots, Y_{l-1}\}| < i$ and $|\{Y_1, \dots, Y_l\}| = i$. So the process Y' is like a depth-first search from left to right in the tree. We can encode all information contained the tree by the two sequences $(a_1, \dots, a_{2k-2}) \in \{u, d\}^{2k-2}$ and $(m_1, \dots, m_{k-1}) \in \mathbb{N}^{k-1}$. The first sequence (a_1, \dots, a_{2k-2}) encodes whether the process Y_i goes one level up or down at a certain point. Here $a_i = u$ if the process goes one level up after Y_i , i.e., if Y_{i+1} is the parent of




Figure 1: In the above tree, the process $(Y'_i)_{i \in \{1, \dots, 9\}}$ is written inside the vertices and the process $(Y_i)_{i \in \{1, \dots, 15\}}$ is written above the vertices. For this tree we have $(a_1, \dots, a_{14}) = (d, d, u, d, u, u, d, d, d, u, u, d, u, u)$.

Y_i . Otherwise we set $a_i = d$, i.e., if Y_{i+1} is the child of Y_i . The sequence (m_1, \dots, m_{k-1}) encodes the direction of the process, where the i -th coordinate gives the direction when the walk goes down for the i -th time. This happens when it touches the vertex Y'_{i+1} for the first time. So if v is the parent of Y'_{i+1} , then $Y'_{i+1} = (v \ m_i)$.

For fixed $\vec{a} = (a_1, \dots, a_{2k-2}) \in \{u, d\}^{2k-2}$, we want to upper bound the expected number of subtrees containing the origin with exactly this up-and-down structure. Assume that the exploration process Y_i visits exactly l children of some vertex Y'_j . Then the expected number of ways to choose these l children among the children of Y'_j in an increasing way is given by

$$\sum_{m_1 \in \mathbb{N}} q(m_1) \sum_{\substack{m_2 \in \mathbb{N} \\ m_2 > m_1}} q(m_2) \cdots \sum_{\substack{m_l \in \mathbb{N} \\ m_l > m_{l-1}}} q(m_l) \leq \mu^l.$$

We have this choice for all vertices Y'_j in the tree. The sum over the number of children of all the vertices is $k-1$, as every vertex, except the origin \emptyset , is the child of exactly one vertex. Thus the expected number of trees with a specified up-and-down structure can be bounded by

$$\sum_{m_1 \in \mathbb{N}} \cdots \sum_{m_{k-1} \in \mathbb{N}} \prod_{i=1}^{k-1} q(m_i) = \mu^{k-1}.$$

Up to now we considered a fixed up-and-down-structure. However, there are at most $|\{u, d\}^{2k-2}| = 2^{2k-2}$ possible up-and-down structures (a_1, \dots, a_{2k-2}) (In fact there are significantly less combinations, as one has additional constraints like $a_1 = d$). So in total, we get

$$\mathbb{E}[|\mathcal{T}_k|] \leq \sum_{\vec{a} \in \{u, d\}^{k-1}} \mu^{k-1} \leq (2^{2k-2}) \mu^{k-1} \leq 4^k \mu^k.$$

□

We now want to use the above lemma about the Galton-Watson tree in order to get results about the average degree of connected subsets of the long-range percolation graph. For this, we define the average degree of some set finite $Z \subset \mathbb{Z}^d$ by

$$\bar{d}(Z) = \frac{1}{|Z|} \sum_{v \in Z} \deg(v).$$

One elementary inequality we will use in the following controls the exponential moments of certain random variables. Assume that $(U_i)_{i \in \mathbb{N}}$ are independent Bernoulli random variables and $U = \sum_{i=1}^{\infty} U_i$. Then

$$\mathbb{E}[e^U] = \mathbb{E}\left[e^{\sum_{i \in \mathbb{N}} U_i}\right] = \prod_{i \in \mathbb{N}} \mathbb{E}[e^{U_i}] \leq \prod_{i \in \mathbb{N}} (1 + e\mathbb{E}[U_i]) \leq \prod_{i \in \mathbb{N}} e^{e\mathbb{E}[U_i]} = e^{e\mathbb{E}[U]} \quad (21)$$

and this already implies, by Markov's inequality, that for any $C > 1$

$$\mathbb{P}(U > C\mathbb{E}[U]) = \mathbb{P}\left(e^U > e^{C\mathbb{E}[U]}\right) \leq \mathbb{E}[e^U] e^{-C\mathbb{E}[U]} \stackrel{(21)}{\leq} e^{(e-C)\mathbb{E}[U]}. \quad (22)$$

Lemma 3.2. *Let $\mathcal{CS}_k = \mathcal{CS}_k(\mathbb{Z}^d)$ be all connected subsets of the long-range percolation graph with vertex set \mathbb{Z}^d of size k that contain the origin $\mathbf{0}$. We write μ_β for $\mathbb{E}_\beta[\deg(\mathbf{0})]$. Then for all $\beta > 0$*

$$\mathbb{P}_\beta(\exists Z \in \mathcal{CS}_k : \bar{d}(Z) \geq 20\mu_\beta) \leq e^{-4k\mu_\beta}.$$

Proof. Consider percolation on the tree $L = \bigcup_{n=0}^{\infty} L_n$, where $L_n = (\mathbb{Z}^d \setminus \{\mathbf{0}\})^n$, the edge set is given by $S = \{\{v, (v m)\} : v \in L, m \in \mathbb{Z}^d \setminus \{\mathbf{0}\}\}$ and an edge of the form $\{v, (v m)\}$ is open with probability $p(\beta, \{\mathbf{0}, m\})$. A total ordering on $\mathbb{Z}^d \setminus \{\mathbf{0}\}$ is given by considering an arbitrary deterministic bijection with \mathbb{N} . From Lemma 3.1 we know that the number of connected sets is bounded by $4^k \mu_\beta^k$. We want to project a finite tree $T \subset L$ of size k down to \mathbb{Z}^d . Remember the notation $(Y'_i)_{i \in \{1, \dots, k\}}$ for the depth-first search from left to right in the tree. The information contained in the structure of the tree, can be represented by the vectors $\vec{a} = (a_1, \dots, a_{2k-2}) \in \{u, d\}^{2k-2}$ and the direction of the vectors $\vec{m} = (m_1, \dots, m_{k-1}) \in (\mathbb{Z}^d \setminus \{\mathbf{0}\})^{k-1}$. We now define a subgraph $(Z(T), E(T))$ of the integer lattice and an exploration process $(X'_i)_{i \in \{1, \dots, k\}}$ as follows:

0. Start with $X'_1 = \mathbf{0}, E_1(T) = \emptyset$.
1. For $i = 2, \dots, k$:
 Let $j < i$ be such that $Y'_i = (Y'_j m)$ for some $m \in \mathbb{Z}^d \setminus \{\mathbf{0}\}$. Set $X'_i = X'_j + m$ and $E_i(T) = E_{i-1}(T) \cup \{\{X'_i, X'_j\}\}$
2. $Z(T) = \bigcup_{i=1}^k \{X'_i\}$ and $E(T) = E_k(T)$

See Figure 2 for an example of this projection. The graph $(Z(T), E(T))$ is clearly connected, but it is not necessarily a tree, as there can be $i \neq j$ with $X'_i = X'_j$, in which case there exists a loop containing X'_i . We call both the graph $(Z(T), E(T))$ and the tree T *admissible* if $(Z(T), E(T))$ is a tree. We also write \mathcal{TA}_k for the set of admissible trees $T \subset (L, S)$. For a tree $T \subset (L, S)$ of size k the condition $T \in \mathcal{TA}_k$ is equivalent to $|Z(T)| = k$, as every connected graph with k vertices and $k - 1$ edges is a tree. Assume




Figure 2: A tree T with 5 vertices, $(a_1, \dots, a_8) = (d, u, d, d, u, d, u, u)$, and $(m_1, \dots, m_4) = (-2, 1, -2, 2)$ and its projection on \mathbb{Z} . The vertices with thick boundary $\{-2, -1, 0, 1, 3\} \subset \mathbb{Z}$ are the set $Z(T)$ and the thick edges between them are the set $E(T)$. Note that $(Z(T), E(T))$ really is a tree for this example.

that the graph $(Z(T), E(T))$ is admissible, then the probability that all edges exist in the random graph equals $\prod_{i=1}^{k-1} p(\beta, \{\mathbf{0}, m_i\})$, which is exactly the probability that all edges of the tree T exist inside (L, S) . Every connected set $Z \subset \mathbb{Z}^d$ has a spanning tree. Thus there exists a tree $T \subset L$ with $Z = Z(T)$ such that all edges in $E(T)$ exist. This and the result of Lemma 3.1 imply that

$$\begin{aligned} \mathbb{E}_\beta [|\mathcal{CS}_k(\mathbb{Z}^d)|] &\leq \sum_{T \in \mathcal{TA}_k} \mathbb{P}_\beta (\text{all edges in } E(T) \text{ exist}) = \sum_{T \in \mathcal{TA}_k} \mathbb{P}_\beta (T \in \mathcal{T}_k) \\ &\leq \mathbb{E}_\beta [|\mathcal{T}_k|] \leq 4^k \mu_\beta^k. \end{aligned} \quad (23)$$

For an admissible tree T , the degree of each vertex $v \in Z(T)$ is the sum of an *inside degree* and an *outside degree*, which we will now define. The *inside degree* $\deg_{Z(T)}(v)$ of a vertex $v \in Z(T)$ is defined by

$$\deg_{Z(T)}(v) = \sum_{u \in Z(T)} \mathbb{1}_{\{\{v,u\} \in E(T)\}}$$

which is just the number of edges in $E(T)$ containing v . Note that for a given admissible tree T the inside degree is purely deterministic and does not depend on the environment. Also note that, by the handshaking lemma,

$$\sum_{v \in Z(T)} \deg_{Z(T)}(v) = 2|E(T)| = 2(|Z(T)| - 1), \quad (24)$$

where the last equality holds as $(Z(T), E(T))$ is a tree. Now let us turn to the *outside degree* $\deg_{Z(T)^c}(v)$ of a vertex $v \in Z(T)$, which we define by

$$\deg_{Z(T)^c}(v) = \sum_{\substack{u \in \mathbb{Z}^d \setminus \{v\} \\ \{u,v\} \notin E(T)}} \omega(\{v,u\})$$

which, contrary to $\deg_{Z(T)}(v)$, now depends on the random environment ω and is a non-constant random variable. Now we want to get bounds on the random variable $\sum_{v \in Z(T)} \deg_{Z(T)^c}(v)$. Note that $\{u, v\} \notin E(T)$ does not imply that $u \notin Z(T)$, but only that u and v are not neighbors in the graph induced by T . The random variable $\sum_{v \in Z(T)} \deg_{Z(T)^c}(v)$ is not the sum of independent Bernoulli random variables, as we might count some edges twice. But as one can count every edge at most twice in this sum, one has the bound

$$\frac{1}{2} \sum_{v \in Z(T)} \deg_{Z(T)^c}(v) \leq \sum_{\substack{\{u, v\} \notin E(T): \\ \{u, v\} \cap Z(T) \neq \emptyset}} \omega(\{u, v\}) \quad (25)$$

where the expression on the right-hand side is a sum of independent Bernoulli random variables with expectation at most $|Z(T)|\mu_\beta$. So for each admissible tree T we always have

$$\sum_{v \in Z(T)} \deg(v) = \sum_{v \in Z(T)} \deg_{Z(T)}(v) + \sum_{v \in Z(T)} \deg_{Z(T)^c}(v) \leq 2|Z(T)| + 2 \sum_{\substack{\{u, v\} \notin E(T): \\ \{u, v\} \cap Z(T) \neq \emptyset}} \omega(\{u, v\}).$$

Let us use the notation

$$U = U(T) := \sum_{\substack{\{u, v\} \notin E(T): \\ \{u, v\} \cap Z(T) \neq \emptyset}} \omega(\{u, v\}).$$

For a given finite admissible tree we have that

$$\begin{aligned} \mathbb{P}_\beta(\bar{d}(Z(T)) \geq 20\mu_\beta) &= \mathbb{P}_\beta\left(\sum_{v \in Z(T)} \deg(v) \geq 20|Z(T)|\mu_\beta\right) \leq \mathbb{P}_\beta(2U \geq 18|T|\mu_\beta) \\ &= \mathbb{P}_\beta(U \geq 9|T|\mu_\beta) \stackrel{(22)}{\leq} \mathbb{E}[e^U] e^{-9|T|\mu_\beta} \leq e^{e|T|\mu_\beta} e^{-9|T|\mu_\beta} \leq e^{-6|T|\mu_\beta}. \end{aligned} \quad (26)$$

So far we only got this bound for a fixed admissible tree $T \subset (L, S)$. Remember that every connected set $Z \in \mathcal{CS}_k$ has a spanning tree and there exists a tree $T \subset (L, S)$ so that $(Z(T), E(T))$ is exactly this spanning tree. Again, we use the notation \mathcal{TA}_k for the set of admissible trees $T \subset (L, S)$ of size k . With the observation from before we get that

$$\begin{aligned} \mathbb{P}_\beta(\exists Z \in \mathcal{CS}_k : \bar{d}(Z) \geq 20\mu_\beta) &\leq \sum_{T \in \mathcal{TA}_k} \mathbb{P}_\beta(\bar{d}(Z(T)) \geq 20\mu_\beta, \text{ all edges in } E(T) \text{ exist}) \\ &\leq \sum_{T \in \mathcal{TA}_k} \mathbb{P}_\beta(U(T) \geq 9k\mu_\beta, \text{ all edges in } E(T) \text{ exist}) \\ &= \sum_{T \in \mathcal{TA}_k} \mathbb{P}_\beta(U(T) \geq 9k\mu_\beta) \mathbb{P}_\beta(\text{all edges in } E(T) \text{ exist}) \\ &\stackrel{(26)}{\leq} e^{-6k\mu_\beta} \sum_{T \in \mathcal{TA}_k} \mathbb{P}_\beta(\text{all edges in } E(T) \text{ exist}) \\ &\stackrel{(23)}{\leq} e^{-6k\mu_\beta} 4^k \mu_\beta^k \leq e^{-6k\mu_\beta} e^{2k} e^{\mu_\beta k} \leq e^{-4k\mu_\beta} \end{aligned}$$

where we used that $\mu_\beta \geq 2$ in the last inequality. This holds for long-range percolation

with our parameters, as each vertex is always connected to its nearest neighbors. The final inequality is exactly the result that we wanted to show and thus finishes the proof. \square

4 Distances in $V_{\mathbf{0}}^n$

In this section, we give several bounds on the distribution of the graph distances between points, respectively sets, inside of certain boxes. In section 4.1, we determine several different properties of the function $(x, y) \mapsto \mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(x, y)]$. It is intuitively clear that the expression is large when x, y also have a big Euclidean distance, for example when $x = \mathbf{0}$ and $y = (n - 1)\mathbf{1}$. This intuition is made rigorous in Lemma 4.2. In section 4.2 we give bounds on the second moment of random variable of the form $d_{V_{\mathbf{0}}^n}(x, y)$. Then, in section 4.3 we use these results in order to bound the distance between certain points and sets in the long-range percolation graph.

4.1 Graph distances of far away points

From the definition of $\Lambda(n, \beta)$ in (8) it is not clear which pair u, v maximizes the expected distance and how the expected graph distances can be compared for different graphs $V_{\mathbf{0}}^n$ and $V_{\mathbf{0}}^{n'}$. In Lemma 4.1, we construct a coupling between the long-range percolation graph on $V_{\mathbf{0}}^n$ for different n . In Lemma 4.2, we show that, up to a constant factor, the maximum of $\Lambda(n, \beta)$ gets attained by the pair $\{\mathbf{0}, (n - 1)e_1\}$ or $\{\mathbf{0}, (n - 1)\mathbf{1}\}$.

Lemma 4.1. *Let $\beta \geq 0$ and $n', n \in \mathbb{N}_{>0}$ with $n' \leq n$. For $u, v \in V_{\mathbf{0}}^n$ define $u' := \lfloor \frac{n'}{n}u \rfloor, v' := \lfloor \frac{n'}{n}v \rfloor$, where the rounding operation is componentwise. There exists a coupling of the random graphs with vertex sets $V_{\mathbf{0}}^n$ and $V_{\mathbf{0}}^{n'}$ such that both are distributed according to \mathbb{P}_{β} and*

$$d_{V_{\mathbf{0}}^{n'}}(u', v') \leq 3d_{V_{\mathbf{0}}^n}(u, v) \quad (27)$$

for all $u, v \in V_{\mathbf{0}}^n$. The same holds true when one considers the graph \mathbb{Z}^d instead of $V_{\mathbf{0}}^n$ and this also implies that

$$\text{Diam}(V_{\mathbf{0}}^{n'}) \leq 3\text{Diam}(V_{\mathbf{0}}^n). \quad (28)$$

Proof. We prove the statement via a coupling with the underlying continuous model. As the claim clearly holds for $\beta = 0$ or for $u = v$, we can assume $\beta > 0, u \neq v$ from here on. Let $\tilde{\mathcal{E}}$ be a Poisson point process on $\mathbb{R}^d \times \mathbb{R}^d$ with intensity $\frac{\beta}{2\|t-s\|^2}$ and define $\mathcal{E} = \{(t, s) \in \mathbb{R}^d \times \mathbb{R}^d : (s, t) \in \tilde{\mathcal{E}}\} \cup \tilde{\mathcal{E}}$. Remember that this Poisson process has a scaling invariance, namely that for a constant $\alpha > 0$ the set $\alpha\mathcal{E}$ has exactly the same distribution as \mathcal{E} . We now define a random graph $G = (V, E)$: For $u, v \in V_{\mathbf{0}}^n =: V$ we place an edge between u and v if and only if $(u + \mathcal{C}) \times (v + \mathcal{C}) \cap n\mathcal{E} \neq \emptyset$. We have already seen in section 1.2 about the continuous model that this creates a sample of independent long-range percolation where the connection probability between the vertices u and v is given by $p(\beta, |v - u|) = 1 - e^{-\int_{u+\mathcal{C}} \int_{v+\mathcal{C}} \frac{\beta}{\|t-s\|^2} dt ds}$. We can do the same procedure for $V' := V_{\mathbf{0}}^{n'}$ and $n'\mathcal{E}$ to get a random graph $G' = (V', E')$. Formally, we place an edge between two vertices $u', v' \in V'$ if and only if $(u' + \mathcal{C}) \times (v' + \mathcal{C}) \cap n'\mathcal{E} \neq \emptyset$. We now claim that for any two vertices $u, v \in V$ with $u \neq v$ and u', v' defined as above one has $d_{G'}(u', v') \leq 2d_G(u, v) + 1$, which already implies (27). Assume that $(x_0 = u, x_1, \dots, x_l = v)$ is the shortest path

between u and v in G , where $l = d_G(u, v)$. Then for all $i = 1, \dots, l$ there are points

$$(y(i, 0), y(i, 1)) \in (x_{i-1} + \mathcal{C}) \times (x_i + \mathcal{C}) \cap n\mathcal{E} .$$

In particular one has

$$\|y(i-1, 1) - y(i, 0)\|_\infty < 1$$

for all $i = 2, \dots, l$, $\|y(1, 0) - u\|_\infty < 1$, and $\|y(l, 1) - v\|_\infty < 1$. For all $i = 1, \dots, l$ and $j \in \{0, 1\}$ define $y'(i, j) = \frac{y'}{n}y(i, j)$, which implies $(y'(i, 0), y'(i, 1)) \in n'\mathcal{E}$. With this definition one clearly has

$$\|y'(i-1, 1) - y'(i, 0)\|_\infty < 1$$

for all $i = 2, \dots, l$, $\|y'(1, 0) - \frac{u'}{n}u\|_\infty < 1$, and $\|y'(l, 1) - \frac{v'}{n}v\|_\infty < 1$. So in G' we can use the path from u' to v' that uses all the edges $\{[y'(i, 0)], [y'(i, 1)]\}$ and in the case where $[y'(i-1, 1)] \neq [y'(i, 0)]$ holds, respectively the analogous statement for u' and v' hold, we can use the nearest neighbor edge between those vertices, which exists as $\|y'(i-1, 1) - y'(i, 0)\|_\infty < 1$. So for each vertex that is touched by the shortest path between u and v in G one needs to make at most one additional step for the path between u' and v' in G' , which implies that $d_{G'}(u', v') \leq 2d_G(u, v) + 1$. If one does not restrict to the sets $V = V_0^n$ and $V' = V_0^{n'}$, but works on the graph with vertex set \mathbb{Z}^d instead, the same proof works. \square

Lemma 4.2. *For all $\beta \geq 0$, $n \in \mathbb{N}_{>0}$, and $u, v \in V_0^n$ we have*

$$\mathbb{E}_\beta [d_{V_0^n}(u, v)] \leq 6d\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)e_1)] \quad (29)$$

and

$$\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)e_1)] \leq 6\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] . \quad (30)$$

This lemma already has two interesting implications, that we want to discuss before going to the proof.

Remark 4.3. *Combining (29) and (30) already implies that for $\Lambda(n, \beta) = \max_{u, v \in V_0^n} \mathbb{E}_\beta [d_{V_0^n}(u, v)] + 1$ one has*

$$\begin{aligned} \mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)e_1)] + 1 &\leq \Lambda(n, \beta) \leq 6d\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)e_1)] + 1 \quad \text{and} \\ \mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] + 1 &\leq \Lambda(n, \beta) \leq 36d\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] + 1 . \end{aligned}$$

Remark 4.4. *For all bounded sets $K \subset \mathbb{R}_{\geq 0}$ there exists a constant $\theta^* > 0$ such that for all $\beta \in K$ and all M, N large enough one has*

$$\Lambda(MN, \beta) \geq M^{\theta^*} \Lambda(N, \beta)$$

Proof. Lemma 4.2 together with (19) already show the existence of such an θ^* along a subsequence of numbers of the form $M = m^k$. Lemma 4.1 shows the result for all large enough M . \square

Proof of Lemma 4.2. Using the triangle inequality and linearity of expectation we get for




Figure 3: Let $v = (6, 3) \in V_{\mathbf{0}}^8$. The points in the gray area are the set $V_{\mathbf{0}}^8$. The points in the hatched area are \mathcal{B}_1 .

all $u, v \in V_{\mathbf{0}}^n$ that

$$\mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(u, v)] \leq \mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(u, \mathbf{0})] + \mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(\mathbf{0}, v)]$$

and thus, in order to prove (29), it suffices to show that

$$\mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(\mathbf{0}, v)] \leq 3d\mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(\mathbf{0}, (n-1)e_1)]$$

for all $v \in V_{\mathbf{0}}^n$. By symmetry, we can assume that $p_1(v) \geq p_2(v) \geq \dots \geq p_d(v)$. For $k \in \{0, \dots, d\}$, we define the vectors $v(k) \in V_{\mathbf{0}}^n$ by

$$v(k) = \sum_{i=1}^k p_i(v) e_i,$$

i.e., the first k coordinates of $v(k)$ equal the corresponding coordinates of v and all other coordinates are 0. By the triangle inequality and linearity of expectation we clearly have

$$\mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(\mathbf{0}, v)] \leq \mathbb{E}_{\beta} \left[\sum_{i=0}^{d-1} d_{V_{\mathbf{0}}^n}(v(i), v(i+1)) \right] = \sum_{i=0}^{d-1} \mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(v(i), v(i+1))].$$

So in order to show (29), it suffices to show that

$$\mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(v(i), v(i+1))] \leq 3\mathbb{E}_{\beta} [d_{V_{\mathbf{0}}^n}(\mathbf{0}, (n-1)e_1)] \quad (31)$$

for all $i \in \{0, \dots, d-1\}$. For each such index i , the cube

$$\mathcal{B}_i = \prod_{j=1}^i \{p_j(v) - p_{i+1}(v), \dots, p_j(v)\} \times \{0, \dots, p_{i+1}(v)\}^{d-i}$$

is contained in the cube $V_{\mathbf{0}}^n$ and contains both the points $v(i)$ and $v(i+1)$, which lie on adjacent corners of the cube. See figure 3 for an example. Allowing the geodesic

to use less edges clearly increases the distance between two points this directly implies $d_{V_0^n}(v(i), v(i+1)) \leq d_{\mathcal{B}_i}(v(i), v(i+1))$ as $\mathcal{B}_i \subset V_0^n$. As the model is invariant under changing the coordinates and under the action $e_i \mapsto -e_i$ we already get for all $i \in \{0, \dots, d-1\}$

$$\mathbb{E}_\beta [d_{\mathcal{B}_i}(v(i), v(i+1))] = \mathbb{E}_\beta [d_{V_0^{p_{i+1}(v)+1}}(\mathbf{0}, p_{i+1}(v)e_1)] \leq 3\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)e_1)],$$

where we used Lemma 4.1 for the last inequality. This shows (31) and thus finishes the proof of (29). Now let us go to the proof of (30). Define $y \in \mathbb{Z}^d$ by $p_1(y) = 1, p_i(y) = -1$ for $i \geq 2$ and the cube \mathcal{B} by $\mathcal{B} = \{n-1, \dots, 2n-2\} \times \{0, \dots, n-1\}^{d-1}$. By the triangle inequality we have

$$d_{V_0^{2n-1}}(\mathbf{0}, (2n-2)e_1) \leq d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1}) + d_{\mathcal{B}}((n-1)\mathbf{1}, (2n-2)e_1).$$

Observe that $(2n-2)e_1 = (n-1)\mathbf{1} + (n-1)y$. The pairs of vertices $\mathbf{0}$ and $(n-1)\mathbf{1}$ lie on opposite corners of the cube V_0^n . The vertices $(n-1)\mathbf{1}$ and $(2n-2)e_1$ also lie on opposite corners of the cube \mathcal{B} . The two cubes V_0^n and \mathcal{B} are only different by a translation, so in particular they have the same side length. As the long-range percolation model is invariant under translation and reflection of any coordinate the two terms in the above sum have the same distribution which implies that

$$\mathbb{E}_\beta [d_{V_0^{2n-1}}(\mathbf{0}, (2n-2)e_1)] \leq 2\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})].$$

Using Lemma 4.1, we finally get

$$\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)e_1)] \leq 3\mathbb{E}_\beta [d_{V_0^{2n-1}}(\mathbf{0}, (2n-2)e_1)] \leq 6\mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})]$$

which shows (30). \square

4.2 The second moment bound

The next lemma relates the second moment of the distances to their first moment. We use a technique that has already been used in [20] before in a slightly different form for dimension $d = 1$ only. As we need the result in a uniform dependence on β in our companion paper [4], we directly prove the uniform statement here. The uniformity does not cause any complications for $d \geq 2$, but it causes minor technical difficulties for $d = 1$. So we give the proof for $d = 1$ in the appendix. The situation for $d \geq 2$ is easier, as there are no cut points, in the sense that for every $u, v \in V_0^n$ there exist two disjoint paths between u and v . For $d = 1$, and in particular for $\beta < 1$, such a statement will typically not be true.

Lemma 4.5. *For all $\beta \geq 0$ there exists a constant $C_\beta < \infty$ such that for all $n \in \mathbb{N}$, all $\varepsilon \in [0, 1]$ and all $u, v \in V_0^n$*

$$\mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(u, v)^2] \leq C_\beta \Lambda(n, \beta + \varepsilon)^2. \quad (32)$$

Proof of Lemma 4.5 for $d \geq 2$. Fix $\beta \geq 0$. We will prove that for all $\varepsilon \in [0, 1]$, all $m, n \in$

\mathbb{N} , and all $u, v \in V_{\mathbf{0}}^{mn}$

$$\mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^{mn}}(u, v)^2] \leq 170m^4 \Lambda(n, \beta + \varepsilon)^2 + 170 \max_{w, z \in V_{\mathbf{0}}^n} \mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^n}(w, z)^2]. \quad (33)$$

Iterating over this inequality one gets for some large enough N that

$$\begin{aligned} & \max_{u, v \in V_{\mathbf{0}}^{m^k N}} \mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^{m^k N}}(u, v)^2] \\ & \leq 170m^4 \sum_{i=0}^k 170^i \Lambda(m^{k-i} N, \beta + \varepsilon)^2 + 170^k \max_{u, v \in V_{\mathbf{0}}^N} \mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^N}(u, v)^2] \\ & \leq 170m^4 \sum_{i=0}^k 170^i \Lambda(m^{k-i} N, \beta + \varepsilon)^2 + 170^k N^2. \end{aligned} \quad (34)$$

for all $k \in \mathbb{N}$. By Remark 4.4 there exists $\theta^* = \theta^*(\beta) > 0$ such that for all $\varepsilon \in [0, 1]$ all $m, n \in \mathbb{N}$ large enough one has

$$\begin{aligned} \Lambda(mn, \beta + \varepsilon) &= \max_{u, v \in V_{\mathbf{0}}^{mn}} \mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^{mn}}(u, v)] + 1 \\ &\geq m^{\theta^*} \left(\max_{u, v \in V_{\mathbf{0}}^n} \mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^n}(u, v)] + 1 \right) = m^{\theta^*} \Lambda(n, \beta + \varepsilon). \end{aligned}$$

Take m large enough so that also $170m^{-\theta^*} < \frac{1}{2}$ is satisfied. Inserting this into (34) gives

$$\begin{aligned} \max_{u, v \in V_{\mathbf{0}}^{m^k N}} \mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^{m^k N}}(u, v)^2] &\leq 170m^4 \sum_{i=0}^k 170^i \Lambda(m^{k-i} N, \beta + \varepsilon)^2 + 170^k N^2 \\ &\leq 170m^4 \sum_{i=0}^k 170^i m^{-\theta^* i} \Lambda(m^k N, \beta + \varepsilon)^2 + N^2 \Lambda(m^k N, \beta + \varepsilon) \\ &\leq (340m^4 + N^2) \Lambda(m^k N, \beta + \varepsilon)^2 \end{aligned}$$

for large enough N . This shows (32) along the subsequence N, mN, m^2N, \dots . For general $n \in \mathbb{N}$, the desired result follows from Lemma 4.1. So we are left with showing (33). For this, we use an elementary observation, that was already used in [20]. Assume that $X_1, \dots, X_{\tilde{m}}$ are independent non-negative random variables and let $\tau = \arg \max_{i \in \{1, \dots, \tilde{m}\}} (X_i)$. Then

$$\mathbb{E} \left[\left(\max_{i \neq \tau} X_i \right)^2 \right] \leq \mathbb{E} \left[\sum_{i=1}^{\tilde{m}} X_i \left(\sum_{j \neq i} X_j \right) \right] = \sum_{i=1}^{\tilde{m}} \sum_{j \neq i} \mathbb{E} [X_i] \mathbb{E} [X_j] \leq \tilde{m}^2 \max_i \mathbb{E} [X_i]^2. \quad (35)$$

We still need to show inequality (33), i.e., that

$$\mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^{mn}}(u, v)^2] \leq 170m^4 \Lambda(n, \beta + \varepsilon)^2 + 170 \max_{w, z \in V_{\mathbf{0}}^n} \mathbb{E}_{\beta+\varepsilon} [d_{V_{\mathbf{0}}^n}(w, z)^2].$$

Let $u, v \in V_{\mathbf{0}}^{mn}$, say with $u \in V_x^n, v \in V_y^n$ where $x, y \in V_{\mathbf{0}}^m$. Inequality (33) clearly holds in the case where $x = y$. For the case $x \neq y$, let $x_0 = x, x_1, \dots, x_l = y$ and $x'_0 = x, x'_1, \dots, x'_l = y$ be two completely disjoint and deterministic paths between x and

y inside V_0^m that are of length at most $m + 1$ and use only nearest-neighbor edges, i.e., $\|x_i - x_{i-1}\|_\infty = 1$ and $\|x'_i - x'_{i-1}\|_\infty = 1$ for all suitable indices i . By completely disjoint we mean that $\{x_1, \dots, x_{l-1}\} \cap \{x'_1, \dots, x'_{l'-1}\} = \emptyset$; the starting point $x = x_0 = x'_0$ and the end point $y = x_l = x'_{l'}$ already need to agree by the construction. Now we iteratively define sequences $(L_i, R_i)_{i=0}^l$ and $(L'_i, R'_i)_{i=0}^{l'}$ as follows:

0. Set $L_0 = u, R_l = v$.

1. For $i = 1, \dots, l$ choose $R_{i-1} \in V_{x_{i-1}}^n$ and $L_i \in V_{x_i}^n$ such that $\|R_{i-1} - L_i\|_\infty = 1$.

Analogously we define $(L'_i, R'_i)_{i=0}^{l'}$ by

0. Set $L'_0 = u, R'_{l'} = v$.

1. For $i = 1, \dots, l'$ choose $R'_{i-1} \in V_{x'_{i-1}}^n$ and $L'_i \in V_{x'_i}^n$ such that $\|R'_{i-1} - L'_i\|_\infty = 1$.

The choice of these algorithms in step (1.) is typically not unique. If there are several possibilities, we always choose a the vertices with some deterministic rule that does not depend on the environment. So in particular we have $L_i, R_i \in V_{x_i}^n$ and $L'_i, R'_i \in V_{x'_i}^n$ for all $i \in \{0, \dots, l\}$, respectively $i \in \{0, \dots, l'\}$. Define

$$\begin{aligned} X_i &= d_{V_{x_i}^n}(L_i, R_i) \text{ for } i \in \{1, \dots, l-1\} \text{ and} \\ X'_i &= d_{V_{x'_i}^n}(L'_i, R'_i) \text{ for } i \in \{1, \dots, l'-1\}. \end{aligned}$$

This are at most $l - 1 + l' - 1 \leq 2m$ random variables and they are independent, as the boxes $V_{x'_i}^n$ and $V_{x_i}^n$ are disjoint. We order the random variables $\{X_i : i \in \{1, \dots, l-1\}\} \cup \{X'_i : i \in \{1, \dots, l'-1\}\}$ in a descending way and call them $Y_1, Y_2, \dots, Y_{l+l'-2}$. The idea in finding a short path between u and v is now to avoid the box where the maximum of the Y_i -s is attained. Assume that the maximum of them is one of the X_i -s, i.e., $X_i = Y_1$ for some $i \in \{1, \dots, l-1\}$. Then we consider the path that goes from $L'_0 = v$ to R'_0 and from there to L'_1 , and from there we go successively to $R'_{l'} = v$. Otherwise, we have $X'_i = Y_1$ for some $i \in \{1, \dots, l'-1\}$. In this situation, we consider the path that goes from $L_0 = v$ to R_0 and from there to L_1 and successively we go to $R_l = v$. In both cases we have

$$d_{V_0^{mn}}(u, v) \leq d_{V_x^n}(L_0, R_0) + d_{V_x^n}(L'_0, R'_0) + d_{V_y^n}(L_l, R_l) + d_{V_x^n}(L'_{l'}, R'_{l'}) + mY_2 + (m + 1), \quad (36)$$

where the summand $(m + 1)$ arises as one still needs to go from R_i to L_{i+1} for all $i \in \{0, \dots, l-1\}$, or from R'_i to L'_{i+1} for all $i \in \{0, \dots, l'-1\}$. But by assumption one has $l, l' \leq m + 1$, so one needs at most $m + 1$ additional steps. From (35) we know that

$$\mathbb{E}_{\beta+\varepsilon} [Y_2^2] \leq 4m^2 \max_{w, z \in V_0^n} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(w, z)]^2. \quad (37)$$

For the distance between L_0 and R_0 one clearly has

$$\mathbb{E}_{\beta+\varepsilon} [d_{V_x^n}(L_0, R_0)^2] \leq \max_{w, z \in V_0^n} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(w, z)^2]$$

and the same statements hold for $d_{V_x^n}(L'_0, R'_0)$, $d_{V_y^n}(L_l, R_l)$, and $d_{V_y^n}(L'_l, R'_l)$. Using the elementary inequality $\left(\sum_{i=1}^6 a_i\right)^2 \leq 36 \sum_{i=1}^6 a_i^2$ that holds for any six numbers $a_1, \dots, a_6 \in \mathbb{R}$ for the term in (36) we get that

$$\begin{aligned}
& \mathbb{E}_{\beta+\varepsilon} [d_{V_0^{mn}}(u, v)^2] \\
& \leq 36 \mathbb{E}_{\beta+\varepsilon} \left[d_{V_x^n}(L_0, R_0)^2 + d_{V_x^n}(L'_0, R'_0)^2 + d_{V_y^n}(L_l, R_l)^2 + d_{V_y^n}(L'_l, R'_l)^2 + m^2 Y_2^2 + (m+1)^2 \right] \\
& \leq 144 \max_{w, z \in V_0^n} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(w, z)^2] + 36m^2 \mathbb{E}_{\beta+\varepsilon} [Y_2^2] + 6(m+1)^2 \\
& \stackrel{(37)}{\leq} 170 \max_{w, z \in V_0^n} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(w, z)^2] + 170m^4 \Lambda(n, \beta + \varepsilon)^2
\end{aligned}$$

which shows (33) and thus finishes the proof. \square

Corollary 4.6. *Iterating this technique one can show that for all $k \in \mathbb{N}$ of the form $k = 2^l$ and for all $\beta > 0$ there exists a constant $C_\beta < \infty$ such that for all $n \in \mathbb{N}$, all $\varepsilon \in [0, 1]$, and all $u, v \in V_0^n$*

$$\mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(u, v)^k] \leq C_\beta \Lambda(n, \beta + \varepsilon)^k. \quad (38)$$

Then, one can extend this bound to all $k \in \mathbb{R}_{\geq 0}$ with Hölder's inequality.

4.3 Graph distances between points and boxes

So far we only considered distances between two different points in a box. In this section, we investigate the distance between certain points and boxes. For $n \in \mathbb{N}$ and $0 < \iota < \frac{1}{2}$ we define the boxes $L_\iota^n := [0, \iota n]^d$ and $R_\iota^n := [n-1-\iota n, n-1]^d$. This are boxes that lie in opposite corners of the box V_0^n , where L_ι^n lies in the corner containing $\mathbf{0}$ and R_ι^n lies in the corner containing $\mathbf{1}$. The next lemma deals with the graph distance of these two boxes. A similar statement of Lemma 4.7 for the continuous model and $d = 1$ was already proven in [20]. We follow the same strategy for the proof of this lemma. Again, we prove it uniformly for β in some compact intervals, as we will need this uniform statement in [4]. The uniformity does not make any complications in this proof here.

Lemma 4.7. *For all $\beta \geq 0$ there exists an $\iota > 0$ such that uniformly over all $\varepsilon \in [0, 1]$ and $n \in \mathbb{N}$*

$$\mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(L_\iota^n, R_\iota^n)] \geq \frac{1}{2} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})], \quad (39)$$

and there exists $c^* > 0$ such that uniformly over all $\varepsilon \in [0, 1]$ and $n \in \mathbb{N}$

$$\mathbb{P}_{\beta+\varepsilon} \left(d_{V_0^n}(L_\iota^n, R_\iota^n) \geq \frac{1}{4} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] \right) \geq c^*. \quad (40)$$

Proof. The statement clearly holds for small n , so we focus on $n \in \mathbb{N}$ large enough from here on. Let $x \in L_\iota^n$ and $y \in R_\iota^n$ be the minimizers of $d_{V_0^n}(L_\iota^n, R_\iota^n)$, i.e., $d_{V_0^n}(L_\iota^n, R_\iota^n) = d_{V_0^n}(x, y)$. If the minimizers are not unique, pick arbitrary ones in some fixed way not depending on the environment. The choice of x, y and the distance $d_{V_0^n}(L_\iota^n, R_\iota^n)$ depend only on edges with at least one endpoint in $V_0^n \setminus (L_\iota^n \cup R_\iota^n)$. The distances $d_{L_\iota^n}(\mathbf{0}, x)$,

respectively $d_{R_\iota^n}(y, (n-1)\mathbf{1})$, depend only on edges with both endpoints in L_ι^n , respectively R_ι^n . Thus we get that

$$\begin{aligned} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] &\leq \mathbb{E}_{\beta+\varepsilon} [d_{R_\iota^n}(\mathbf{0}, x)] + \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(L_\iota^n, R_\iota^n)] + \mathbb{E}_{\beta+\varepsilon} [d_{R_\iota^n}(y, (n-1)\mathbf{1})] \\ &\leq 2\Lambda(\lfloor \iota n \rfloor, \beta + \varepsilon) + \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(L_\iota^n, R_\iota^n)] . \end{aligned}$$

For ι small enough we get uniformly over $\varepsilon \in [0, 1]$ that

$$\Lambda(n, \beta + \varepsilon) \geq \left(\frac{1}{\iota}\right)^{\theta'} \Lambda(\lfloor \iota n \rfloor, \beta + \varepsilon)$$

for some $\theta' > 0$ by Remark 4.4. So by Lemma 4.2, respectively Remark 4.3, we can choose ι small enough so that uniformly over $n \in \mathbb{N}$ and $\varepsilon \in [0, 1]$

$$2\Lambda(\lfloor \iota n \rfloor, \beta + \varepsilon) \leq \frac{1}{2} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})]$$

and this implies that

$$\begin{aligned} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(L_\iota^n, R_\iota^n)] &\geq \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] - 2\Lambda(\lfloor \iota n \rfloor, \beta + \varepsilon) \\ &\geq \frac{1}{2} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] \end{aligned}$$

which proves (39). For such an ι , let \mathcal{A} be the event that $d_{V_0^n}(L_\iota^n, R_\iota^n) \geq \frac{1}{4} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})]$. By the Cauchy-Schwarz inequality we have

$$\begin{aligned} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] &\leq 2\mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(L_\iota^n, R_\iota^n)] \\ &= 2\mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(L_\iota^n, R_\iota^n) \mathbb{1}_{\mathcal{A}^c}] + 2\mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(L_\iota^n, R_\iota^n) \mathbb{1}_{\mathcal{A}}] \\ &\leq \frac{1}{2} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] + 2\mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})^2]^{1/2} \sqrt{\mathbb{P}_{\beta+\varepsilon}(\mathcal{A})} \\ &\leq \frac{1}{2} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] + C' \mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] \sqrt{\mathbb{P}_{\beta+\varepsilon}(\mathcal{A})}, \end{aligned}$$

where the last inequality holds for some $C' < \infty$, by Lemma 4.2 and Lemma 4.5. Solving the previous line of inequalities for $\mathbb{P}_{\beta+\varepsilon}(\mathcal{A})$ shows (40). \square

Lemma 4.8. *For all $\beta \geq 0$ and all dimensions d there exists a constant $c_1 > 0$ such that uniformly over all $n \in \mathbb{N}$ and all $x \in \mathcal{S}_n$*

$$\mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, x)] \geq c_1 \mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] \quad (41)$$

and the constant c_1 can be chosen in such a way so that it only depends on the dimension d and the value $\iota > 0$ in (39).

Proof. Let $v \in \mathcal{S}_n$ be one of the minimizers of $y \mapsto \mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, y)]$ among all vertices $y \in \mathcal{S}_n$. By reflection symmetry, we can assume that all coordinates of v are non-negative. With the notation $e_0 = e_d$ we define the vectors v_0, \dots, v_{d-1} by

$$\langle e_j, v_i \rangle = \langle e_{i+j \bmod d}, v \rangle$$

which are just versions of the vector v , where we cyclically permuted the coordinates. By

invariance under changes of coordinates, we have

$$\mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, v)] = \mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, v_i)]$$

for all $i \in \{0, \dots, d-1\}$. Define the vertices u_0, \dots, u_d by $u_j = \sum_{i=1}^j v_i$. By our construction we have $u_0 = \mathbf{0}$ and $u_d = \sum_{i=1}^d v_i = N\mathbf{1}$ for some integer $N \geq n$. The balls $B_n(u_i)$ are all contained in the cube $\Upsilon = \{-n, \dots, N+n\}^d$ for all $i \in \{0, \dots, d\}$. Thus we have

$$\mathbb{E}_\beta [d_\Upsilon(\mathbf{0}, N\mathbf{1})] \leq \sum_{i=1}^{d-1} \mathbb{E}_\beta [d_\Upsilon(u_{i-1}, u_{i-1} + v_i\mathbf{1})] \leq d\mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, v)]$$

and by translation invariance we also have for the cube $\Upsilon_1 = \{0, \dots, 2n+N\}^d$

$$\mathbb{E}_\beta [d_{\Upsilon_1}(n\mathbf{1}, (n+N)\mathbf{1})] \leq d \mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, v)].$$

Using the triangle inequality, we see that for all $k \in \mathbb{N}$ the expected distance between $n\mathbf{1}$ and $(n+kN)\mathbf{1}$ inside the cube $\Upsilon_k = \{0, \dots, kN+2n\}^d$ is upper bounded by

$$\mathbb{E}_\beta [d_{\Upsilon_k}(n\mathbf{1}, (n+kN)\mathbf{1})] \leq k \cdot d \mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, v)].$$

But Lemma 4.1 also gives that for $s = \frac{n}{kN+2n}$ and $w_1 = \lfloor sn\mathbf{1} \rfloor, w_2 = \lfloor s(n+kN)\mathbf{1} \rfloor$

$$\mathbb{E}_\beta [d_{V_0^n}(w_1, w_2)] \leq 3k \cdot d \mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, v)].$$

As $N \geq n$, for each fixed $\iota > 0$ we can choose k large enough so that $w_1 \in L_\iota^n$ and $w_2 \in R_\iota^n$ and thus $\mathbb{E}_\beta [d_{V_0^n}(w_1, w_2)] \geq \mathbb{E}_\beta [d_{V_0^n}(L_\iota^n, R_\iota^n)]$. Then we get by the lower bound on the expected distance between the boxes L_ι^n and R_ι^n (39) that for such a k

$$\begin{aligned} \mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, v)] &\geq \frac{1}{3kd} \mathbb{E}_\beta [d_{V_0^n}(w_1, w_2)] \geq \frac{1}{3kd} \mathbb{E}_\beta [d_{V_0^n}(L_\iota^n, R_\iota^n)] \\ &\stackrel{(39)}{\geq} \frac{1}{6kd} \mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)\mathbf{1})] \end{aligned}$$

which finishes the proof, as $v \in \mathcal{S}_n$ was assumed to minimize the expected distance $\mathbb{E}_\beta [d_{B_n(\mathbf{0})}(\mathbf{0}, y)]$ among all vertices $y \in \mathcal{S}_n$. \square

Lemma 4.9. *For all dimensions d and all $\beta > 0$ there exists an $\eta \in (0, \frac{1}{2})$ such that uniformly over all $n \in \mathbb{N}$ and all $x \in \mathcal{S}_n$*

$$\mathbb{E}_\beta [d_{B_n(\mathbf{0})}(B_{\eta n}(\mathbf{0}), B_{\eta n}(x))] \geq \frac{c_1}{2} \Lambda(n, \beta) \quad (42)$$

where c_1 is the constant from (41) and there exists a constant c_2 such that

$$\mathbb{P}_\beta \left(d_{B_n(\mathbf{0})}(B_{\eta n}(\mathbf{0}), B_{\eta n}(x)) \geq \frac{c_1}{4} \Lambda(n, \beta) \right) \geq c_2. \quad (43)$$

Furthermore, for each $\beta \geq 0$ there exist constants $c_3 > 0$ such that

$$\mathbb{P}_\beta \left(d(B_n(\mathbf{0}), B_{2n}(\mathbf{0})^C) \geq \frac{c_1}{4} \Lambda(n, \beta) \right) \geq c_3 \quad (44)$$

uniformly over all $n \in \mathbb{N}$.

Note that in the above lemma, for $x \in \mathcal{S}_n$ the boxes $B_{\eta n}(x)$ are not completely contained inside $B_n(\mathbf{0})$, but from the definition of $d_{B_n(\mathbf{0})}(\cdot, \cdot)$, we only consider the part that intersects $B_n(\mathbf{0})$.

Proof. The proof of (42) and (43) is the same as the proof of Lemma 4.7 and we omit it. Regarding the statement of (44) we will first prove that for $\eta > 0$ small enough

$$\mathbb{P}_\beta \left(d(B_{\eta n}(\mathbf{0}), B_n(\mathbf{0})^C) \geq \frac{c_1}{4} \Lambda(n, \beta) \right) \geq c_4 \quad (45)$$

for some constant $c_4 > 0$ and uniformly over all $n \in \mathbb{N}$. For this, we use the *FKG inequality*, see [28, Section 1.3] or [23, 26] for the original papers. We can cover the set $\bigcup_{x \in \mathcal{S}_n} B_{\eta n}(x)$ with uniformly (in n) finitely many sets of the form $B_{\eta n}(x)$. For example, we have

$$\bigcup_{\substack{x \in \mathcal{S}_n: \\ \langle x, e_1 \rangle = n}} B_{\eta n}(x) \subset \bigcup_{x \in F} B_{\eta n}(x)$$

where $F = \left\{ n e_1 + \sum_{i=2}^d k_i e_i : k_i \in \left\{ -\lceil \frac{n}{|\eta n|} \rceil, \dots, \lceil \frac{n}{|\eta n|} \rceil \right\} \text{ for all } i \in \{2, \dots, d\} \right\}$, and all other faces of the set $\bigcup_{x \in \mathcal{S}_n} B_{\eta n}(x)$ can be covered in a similar way. Suppose that $A'_n \subset \mathcal{S}_n$ is a sequence of finite sets with $\sup_n |A'_n| =: A' < \infty$ such that

$$\bigcup_{x \in \mathcal{S}_n} B_{\eta n}(x) = \bigcup_{x \in A'_n} B_{\eta n}(x)$$

for all $n \in \mathbb{N}$. So in particular we have

$$\begin{aligned} & \left\{ d_{B_n(\mathbf{0})}(B_{\eta n}(\mathbf{0}), B_{\eta n}(x)) \geq \frac{c_1}{4} \Lambda(n, \beta) \text{ for all } x \in \mathcal{S}_n \right\} \\ &= \left\{ d_{B_n(\mathbf{0})}(B_{\eta n}(\mathbf{0}), B_{\eta n}(x)) \geq \frac{c_1}{4} \Lambda(n, \beta) \text{ for all } x \in A'_n \right\}. \end{aligned}$$

The events $\left\{ d_{B_n(\mathbf{0})}(B_{\eta n}(\mathbf{0}), B_{\eta n}(x)) \geq \frac{c_1}{4} \Lambda(n, \beta) \right\}$ are decreasing for all $x \in \mathcal{S}_n$ in the sense that they are stable under the deletion of edges. Thus the FKG inequality and (43) imply that that

$$\mathbb{P}_\beta \left(d_{B_n(\mathbf{0})}(B_{\eta n}(\mathbf{0}), B_{\eta n}(x)) \geq \frac{c_1}{4} \Lambda(n, \beta) \text{ for all } x \in \mathcal{S}_n \right) \geq c_2^{|A'_n|} \geq c_2^{A'}.$$

Assume that there is no direct edge from $[-(n - \eta n), (n - \eta n)]^d$ to $\mathbb{Z}^d \setminus [-n, n]^d$. This has a uniform positive probability and is also a decreasing event. Then any path from $B_{\eta n}(\mathbf{0})$ to $B_n(\mathbf{0})^C$ goes through at least one box $B_{\eta n}(x) \cap B_n(\mathbf{0})$ for some $x \in \mathcal{S}_n$. So with another application of the FKG inequality we get that

$$\mathbb{P}_\beta \left(d(B_{\eta n}(\mathbf{0}), B_n(\mathbf{0})^C) \geq \frac{c_1}{4} \Lambda(n, \beta) \right) \geq c_5$$

for some $c_5 > 0$ and uniformly over all $n \in \mathbb{N}$. Next, let $A_n \subset B_n(\mathbf{0})$ be a sequence of sets such that $\bigcup_{x \in A_n} B_{\eta n}(x) = B_n(\mathbf{0})$ and $\sup_n |A_n| =: A < \infty$. Then $d(B_n(\mathbf{0}), B_{2n}(\mathbf{0})^C) < \frac{c_1}{4} \Lambda(n, \beta)$ already implies that there exists a point $x \in A_n$ such that $d(B_{\eta n}(x), B_n(x)^C) <$

$\frac{c_1}{4}\Lambda(n, \beta)$. By another application of the FKG inequality we have

$$\begin{aligned} & \mathbb{P}_\beta \left(d(B_n(\mathbf{0}), B_{2n}(\mathbf{0})^C) \geq \frac{c_1}{4}\Lambda(n, \beta) \right) \\ & \geq \mathbb{P}_\beta \left(d(B_{\eta_n}(x), B_n(x)^C) \geq \frac{c_1}{4}\Lambda(n, \beta) \text{ for all } x \in A_n \right) \geq c_5^{|A_n|} \geq c_5^A \end{aligned}$$

which proves (44). \square

Lemma 4.10. *For all $\beta \geq 0$ and all $\varepsilon > 0$ there exist $0 < c < C < \infty$ such that*

$$\mathbb{P}_\beta (c\Lambda(n, \beta) \leq d(\mathbf{0}, B_n(\mathbf{0})^C) \leq C\Lambda(n, \beta)) > 1 - \varepsilon \quad (46)$$

for all $n \in \mathbb{N}$.

Similar statements for one dimension and the continuous model were already proven in [20]. We follow a similar strategy here.

Proof. By the inequality $d(\mathbf{0}, B_n(\mathbf{0})^C) \leq d_{V_0^{n+2}}(\mathbf{0}, (n+1)\mathbf{1})$ we get that

$$\mathbb{E}_\beta [d(\mathbf{0}, B_n(\mathbf{0})^C)] \leq \Lambda(n+2, \beta) \leq \Lambda(n, \beta) + 2$$

and thus the probability $\mathbb{P}_\beta (d(\mathbf{0}, B_n(\mathbf{0})^C) \leq C\Lambda(n, \beta))$ can be made arbitrarily close to 1 by taking C large because of Markov's inequality. We will also refer to this case as the upper bound. The lower bound clearly holds for small n , so we will always focus on n large enough. Fix $K, N \in \mathbb{N}$ such that the function $i \mapsto \Lambda(K^{2^i}N, \beta)$ is increasing in i . This is possible by Remark 4.4. We now consider boxes of the form $B_{K^{2^i}N}(\mathbf{0})$. The probability of a direct edge from $B_{K^{2^{i-1}}N}(\mathbf{0})$ to $B_{K^{2^i}N}(\mathbf{0})^C$ equals the probability of a direct edge between $\mathbf{0}$ and $B_{K^2}(\mathbf{0})^C$ and is by (5) bounded by $\beta 50^d K^{-2}$. So the probability that there is some $i \in \{1, \dots, K\}$ for which there is a direct edge from $B_{K^{2^{i-1}}N}(\mathbf{0})$ to $B_{K^{2^i}N}(\mathbf{0})^C$ is bounded by $\beta 50^d K^{-1}$. We denote the complement of this event by \mathcal{A} . Conditioned on the event \mathcal{A} , where there exists no edge between $B_{K^{2^{i-1}}N}(\mathbf{0})$ and $B_{K^{2^i}N}(\mathbf{0})^C$ for all $i \in \{1, \dots, K\}$, each path from $B_N(\mathbf{0})$ to $B_{K^{2^K}N}(\mathbf{0})$ needs to cross all the distances from $B_{K^{2^{i-1}}N}(\mathbf{0})$ to $B_{2K^{2^{i-1}}N}(\mathbf{0})^C$. For odd i , these distances are independent. Remember that $i \mapsto \Lambda(K^{2^i}N, \beta)$ is increasing in i . So conditioned on the event \mathcal{A} we have the bound

$$\begin{aligned} & \mathbb{P}_\beta \left(d(\mathbf{0}, B_{K^{2^K}N}(\mathbf{0})^C) < \frac{c_1}{4}\Lambda(N, \beta) \middle| \mathcal{A} \right) \\ & \leq \mathbb{P}_\beta \left(d(B_{K^{2^{i-1}}N}(\mathbf{0}), B_{2K^{2^{i-1}}N}(\mathbf{0})^C) < \frac{c_1}{4}\Lambda(K^{2^{i-1}}N, \beta) \forall i \in \{1, \dots, K\} \text{ odd} \middle| \mathcal{A} \right) \\ & \leq (1 - c_3)^{\lfloor \frac{K}{2} \rfloor} \end{aligned}$$

and thus we have

$$\begin{aligned} & \mathbb{P}_\beta \left(d(\mathbf{0}, B_{K^{2^K}N}(\mathbf{0})^C) < \frac{c_1}{4}\Lambda(N, \beta) \right) \\ & \leq \mathbb{P}_\beta \left(d(\mathbf{0}, B_{K^{2^K}N}(\mathbf{0})^C) < \frac{c_1}{4}\Lambda(N, \beta) \middle| \mathcal{A} \right) + \mathbb{P}_\beta (\mathcal{A}^C) \leq (1 - c_3)^{\lfloor \frac{K}{2} \rfloor} + \beta 50^d K^{-1} \end{aligned}$$

and this quantity can be made arbitrary small by suitable choice of K . To finish the

proof, remember that $\Lambda(N, \beta)$ and $\Lambda(K^{2K}N, \beta)$ are off by a factor of at most K^{2K} , as

$$\Lambda(N, \beta) \leq \Lambda(K^{2K}N, \beta) \stackrel{(8)}{\leq} \Lambda(K^{2K}, \beta) \Lambda(N, \beta) \leq K^{2K} \Lambda(N, \beta).$$

Thus we have

$$\begin{aligned} \mathbb{P}_\beta \left(d(\mathbf{0}, B_{K^{2K}N}(\mathbf{0})^C) < \frac{c_1}{4K^{2K}} \Lambda(K^{2K}N, \beta) \right) &\leq \mathbb{P}_\beta \left(d(\mathbf{0}, B_{K^{2K}N}(\mathbf{0})^C) < \frac{c_1}{4} \Lambda(N, \beta) \right) \\ &\leq (1 - c_3)^{\lfloor \frac{K}{2} \rfloor} + \beta 50^d K^{-1}. \end{aligned}$$

Now, for fixed $\varepsilon > 0$ take K large enough so that $(1 - c_3)^{\lfloor \frac{K}{2} \rfloor} + \beta 50^d K^{-1} < \varepsilon$. For $n \in \mathbb{N}$, let N be the largest integer for which $K^{2K}N \leq n$. We know that $K^{2K}N \leq n \leq K^{2K}2N$ and this also yields, by Lemma 4.1, that

$$\Lambda(n, \beta) \leq 3\Lambda(K^{2K}2N, \beta) \leq 6\Lambda(K^{2K}N, \beta)$$

which already implies

$$\begin{aligned} \mathbb{P}_\beta \left(d(\mathbf{0}, B_n(\mathbf{0})^C) < \frac{c_1}{24K^{2K}} \Lambda(n, \beta) \right) &\leq \mathbb{P}_\beta \left(d(\mathbf{0}, B_n(\mathbf{0})^C) < \frac{c_1}{4K^{2K}} \Lambda(K^{2K}N, \beta) \right) \\ &\leq \mathbb{P}_\beta \left(d(\mathbf{0}, B_{K^{2K}N}(\mathbf{0})^C) < \frac{c_1}{4K^{2K}} \Lambda(K^{2K}N, \beta) \right) \leq (1 - c_3)^{\lfloor \frac{K}{2} \rfloor} + \beta 50^d K^{-1} \leq \varepsilon. \end{aligned}$$

□

The previous Lemma tells us that for fixed $\beta > 0$ all quantiles of $d(\mathbf{0}, B_n(\mathbf{0})^C)$ are of order $\Lambda(n, \beta)$. We want to prove a similar statement for the quantiles of $d(B_n(\mathbf{0}), B_{2n}(\mathbf{0})^C)$. However, an analogous statement can not be true, as there is a uniform positive probability of a direct edge between $B_n(\mathbf{0})$ and $B_{2n}(\mathbf{0})^C$. But if we condition on the event that there is no such direct edge, the statement still holds.

Lemma 4.11. *Let \mathcal{L} be the event that there is no direct edge between $B_n(\mathbf{0})$ and $B_{2n}(\mathbf{0})^C$. For all $\beta \geq 0$ and all $\varepsilon > 0$ there exist $0 < c < C < \infty$ such that*

$$\mathbb{P}_\beta \left(c\Lambda(n, \beta) \leq d(B_n(\mathbf{0}), B_{2n}(\mathbf{0})^C) \leq C\Lambda(n, \beta) \mid \mathcal{L} \right) > 1 - \varepsilon \quad (47)$$

for all $n \in \mathbb{N}$.

Proof. The upper bound follows with Markov's inequality, as

$$\begin{aligned} \mathbb{E}_\beta [d(B_n(\mathbf{0}), B_{2n}(\mathbf{0})^C)] &\leq \mathbb{E}_\beta [d(n\mathbf{1}, (2n+1)\mathbf{1})] \\ &\leq \mathbb{E}_\beta [d_{V_1^n}(n\mathbf{1}, (2n-1)\mathbf{1})] + 2 \leq \Lambda(n, \beta) + 1. \end{aligned}$$

For the lower bound, we first consider integers of the form $N_k = M^k N_0$, where we fix $M \in \mathbb{N}$ first. Let M be the smallest natural number such that $M \geq 100$ and $\Lambda(M, \beta) \leq \frac{M}{10}$. The inequality $\Lambda(M, \beta) \leq \frac{M}{10}$ holds for large enough M , as $\Lambda(M, \beta)$ asymptotically grows like a power of M that is strictly less than one, see section 2.2. As β is fixed for the rest of the proof, we simply write $\Lambda(n)$ for $\Lambda(n, \beta)$. We write C_n for the annulus $B_{2n}(\mathbf{0}) \setminus B_n(\mathbf{0})$. Let \mathcal{A}_δ denote the event that for all vertices $x \in C_{MN}$ for which there exists an edge $e = \{x, y\}$ with $\|x - y\|_\infty \geq N$ one has $d(x, C_{MN}^C; \omega^{e^-}) \geq \delta \Lambda(MN)$. We will now show

that the probability of the event \mathcal{A}_δ converges to 1 as $\delta \rightarrow 0$. Remember that $\Lambda(MN)$ and $\Lambda(N)$ differ by a factor of at most M . Let us first consider the event that for some $\delta_1 > 0$ there exists a vertex incident to a long edge in one of the boundary regions of thickness $\delta_1 N$ of C_{MN} , Formally, we define the boundary region $\partial^{\delta_1} C_{MN}$ of C_{MN} by

$$\partial^{\delta_1} C_{MN} = \{B_{MN+\delta_1 N}(\mathbf{0}) \setminus B_{MN}(\mathbf{0})\} \cup \{B_{2MN}(\mathbf{0}) \setminus B_{2MN-\delta_1 N}(\mathbf{0})\} .$$

Each of the two parts of $\partial^{\delta_1} C_{MN}$ has a size of at most $4d\delta_1 N (5MN)^{d-1}$, as one needs to fix one of the coordinates within the interval $(MN, MN + \delta_1 N]$, respectively in the interval $(2MN - \delta_1 N, 2MN]$, or one of the reflections of these intervals, and then has at most $4MN + 1$ possibilities for each of the remaining $d - 1$ coordinates. The probability that a vertex is incident to some edge of length $\geq N$ is proportional to $\frac{\beta}{N^d}$ as shown in (5). So together with (5) we get that

$$\mathbb{P}_\beta (\exists x \in \partial^{\delta_1} C_{MN}, y \in B_{N-1}(x)^C : x \sim y) \leq \delta_1 \cdot 4d(5MN)^d \beta 50^d N^{-d} \leq \delta_1 \cdot \beta (10^3 M)^d .$$

Furthermore, the expected number of points $x \in C_{MN}$ which are incident to a long edge is bounded by

$$\mathbb{E}_\beta [|\{x \in C_{MN} : x \sim B_{N-1}(x)^C\}|] \stackrel{(5)}{\leq} (5MN)^d \beta 50^d N^{-d} \leq \beta (250M)^d .$$

As the existence of a long edge e incident to some x and $d(x, B_{\delta_1 N}(x)^C; \omega^{e-})$ are independent we get with a union bound that

$$\begin{aligned} & \mathbb{P}_\beta (\exists x \in C_{MN}, y \in B_{N-1}(x)^C : x \sim y, d(x, B_{\delta_1 N}(x)^C; \omega^{e-}) < \delta \Lambda(MN)) \\ & \leq \sum_{x \in C_{MN}} \mathbb{P}_\beta (x \sim B_{N-1}(x)^C) \mathbb{P}_\beta (d(x, B_{\delta_1 N}(x)^C) < \delta \Lambda(MN)) \\ & \leq \beta (250M)^d \mathbb{P}_\beta (d(\mathbf{0}, B_{\delta_1 N}(\mathbf{0})^C) < \delta \Lambda(MN)) \end{aligned}$$

and thus we also get that

$$\begin{aligned} \mathbb{P}_\beta (\mathcal{A}_\delta^C) &= \mathbb{P}_\beta (\exists x \in C_{MN}, y \in B_{N-1}(x)^C : x \sim y, d(x, C_{MN}^C; \omega^{\{x,y\}^-}) < \delta \Lambda(MN)) \\ &\leq \mathbb{P}_\beta (\exists x \in \partial^{\delta_1} C_{MN}, y \in B_{N-1}(x)^C : x \sim y) \\ &\quad + \mathbb{P}_\beta (\exists x \in C_{MN}, y \in B_{N-1}(x)^C : x \sim y, d(x, B_{\delta_1 N}(x)^C; \omega^{\{x,y\}^-}) < \delta \Lambda(MN)) \\ &\leq \delta_1 \beta (10^3 M)^d + \beta (250M)^d \mathbb{P}_\beta (d(\mathbf{0}, B_{\delta_1 N}(\mathbf{0})^C) < \delta \Lambda(MN)) \end{aligned} \tag{48}$$

and this converges to 0 as $\delta \rightarrow 0$, for an appropriate choice of $\delta_1(\delta)$, by Lemma 4.10 uniformly over $N \in \mathbb{N}$. We write $f(\delta)$ for the supremum of $\mathbb{P}_\beta (\mathcal{A}_\delta^C)$ over all $N \in \mathbb{N}$ and for $A, B \subset V$ we write $d^*(A, B)$ for the indirect distance between A and B , i.e., the length of the shortest path between A and B that does not use a direct edge between A and B . Now assume that $d^*(B_{MN}(\mathbf{0}), B_{2MN}(\mathbf{0})^C) < \delta \Lambda(MN)$. We now consider the path between $B_{MN}(\mathbf{0})$ and $B_{2MN}(\mathbf{0})^C$ that achieves this distance. Either this path uses some long edge (of length greater than $N - 1$) or it only jumps from one block of the form V_v^N to directly neighboring blocks. The probability that there exists a point $x \in C_{MN}$ and a long edge e incident to it such that $d(x, C_{MN}^C; \omega^{e-}) < \delta \Lambda(MN)$ is relatively small

by (48). Any path that does not use long edges can only do jumps between neighboring blocks of the form V_v^N . Say the path uses the blocks $\left(V_{v_i}^N\right)_{i=0}^{L'}$. Consider the loop-erased trace of this walk on the blocks, i.e., say that the path uses the blocks $\left(V_{v_i}^N\right)_{i=0}^L \subset C_{MN}$ with $\|u_i - v_{i-1}\|_\infty = 1$ and never returns to $V_{v_i}^N$ after going to $V_{v_{i+1}}^N$. There need to be at least $\frac{M}{3}$ transitions between blocks $V_{u_i}^N$ and $V_{w_i}^N$ with $\|v_i - w_i\|_\infty = 2$ as the path needs to walk a distance in the infinity-norm of at least MN . So in particular we have

$$\begin{aligned} \sum_{i=1}^{M/3} d_{B_{2MN}(\mathbf{0})}^*(V_{u_i}^N, V_{w_i}^N) &\leq d_{B_{2MN}(\mathbf{0})}^*(B_{MN}(\mathbf{0}), B_{2MN}(\mathbf{0})^C) \\ &< \delta\Lambda(MN, \beta) \leq \delta\Lambda(M, \beta)\Lambda(N, \beta) \leq \frac{M}{10}\delta\Lambda(N, \beta) \end{aligned}$$

where we used our assumption on M for the last step. So in particular there need to be at least two transitions between $V_{u_i}^N$ and $V_{w_i}^N$ that satisfy $d_{B_{2MN}(\mathbf{0})}^*(V_{u_i}^N, V_{w_i}^N) < \delta\Lambda(N, \beta)$. In fact, there need to be some linear number in M many such transitions, but two are sufficient for our purposes here. All these transitions need to be disjoint, as shortest paths never use the same edge twice. Thus we get by the *BK inequality* (see [28, Section 1.3] or [9, 34]) that

$$\mathbb{P}_\beta \left(\sum_{i=1}^{M/3} d_{B_{2MN}(\mathbf{0})}^*(V_{u_i}^N, V_{w_i}^N) \leq \frac{M}{10}\delta\Lambda(N) \right) \leq M^2 \left(\min_i \mathbb{P}_\beta \left(d_{B_{2MN}(\mathbf{0})}^*(V_{u_i}^N, V_{w_i}^N) < \delta\Lambda(N) \right) \right)^2$$

For each combination of vectors u_i, w_i with $\|u_i - w_i\|_\infty = 2$ we can translate and rotate the boxes $V_{u_i}^N$ and $V_{w_i}^N$ to boxes $T(V_{u_i}^N)$ and $T(V_{w_i}^N)$ in such a way that $T(V_{u_i}^N) \subset B_N(\mathbf{0})$ and $T(V_{w_i}^N) \subset B_{2N}(\mathbf{0})^C$. By translational and rotational invariance of our long-range percolation model this already implies that

$$\min_i \mathbb{P}_\beta \left(d_{B_{2MN}(\mathbf{0})}^*(V_{u_i}^N, V_{w_i}^N) < \delta\Lambda(N) \right) \leq \mathbb{P}_\beta \left(d_{B_{2N}(\mathbf{0})}^*(B_N(\mathbf{0}), B_{2N}(\mathbf{0})^C) < \delta\Lambda(N) \right).$$

There are at most $((5M)^d)!$ choices for possible choice of vertices v_0, v_1, \dots, v_L , as there are at most $(5M)^d$ possibilities for v_0 and $(5M)^d - 1$ possibilities for v_1 and so on. Overall we see that the probability that there exists an indirect path between $B_{MN}(\mathbf{0})$ and $B_{2MN}(\mathbf{0})^C$ of length $\delta\Lambda(MN)$ that jumps between neighboring blocks of the form V_v^N only is bounded by

$$(5M^d)!M^2\mathbb{P}_\beta \left(d_{B_{2N}(\mathbf{0})}^*(B_N(\mathbf{0}), B_{2N}(\mathbf{0})^C) < \delta\Lambda(N) \right)^2.$$

We write D for the constant $(5M^d)!M^2$. Thus we get that

$$\begin{aligned} &\mathbb{P}_\beta \left(d_{B_{2MN}(\mathbf{0})}^*(B_{MN}(\mathbf{0}), B_{2MN}(\mathbf{0})^C) < \delta\Lambda(MN) \right) \\ &\leq D\mathbb{P}_\beta \left(d_{B_{2N}(\mathbf{0})}^*(B_N(\mathbf{0}), B_{2N}(\mathbf{0})^C) < \delta\Lambda(N) \right)^2 + \mathbb{P}_\beta(\mathcal{A}_\delta^C). \end{aligned}$$

We define the sequence $(a_n)_{n \in \mathbb{N}}$ by

$$a_0 = \mathbb{P}_\beta \left(d_{B_{2N}(\mathbf{0})}^*(B_N(\mathbf{0}), B_{2N}(\mathbf{0})^C) < \delta\Lambda(N) \right)$$

and $a_{n+1} = Da_n^2 + f(\delta)$. Inductively it follows that

$$\mathbb{P}_\beta \left(d_{B_{2M^k N}(\mathbf{0})}^* (B_{M^k N}(\mathbf{0}), B_{2M^k N}(\mathbf{0})^C) < \delta \Lambda(M^k N) \right) \leq a_k$$

for all $k \in \mathbb{N}$. For $f(\delta) < \frac{1}{4D}$, the equation $a = Da^2 + f(\delta)$ has the two solutions

$$a_- = \frac{1 - \sqrt{1 - 4Df(\delta)}}{2D} \text{ and } a_+ = \frac{1 + \sqrt{1 - 4Df(\delta)}}{2D} > \frac{1}{2D}.$$

For $a_0 \in [0, a_+)$, and thus in particular for $a_0 \in [0, \frac{1}{2D}]$, the sequence a_n converges to $a_- = \frac{1 - \sqrt{1 - 4Df(\delta)}}{2D} \approx f(\delta)$ and thus we get

$$\limsup_{k \rightarrow \infty} \mathbb{P}_\beta \left(d_{B_{2M^k N}(\mathbf{0})}^* (B_{M^k N}(\mathbf{0}), B_{2M^k N}(\mathbf{0})^C) < \delta \Lambda(M^k N) \right) \leq a_-.$$

For fixed $N \in \mathbb{N}$, the requirement

$$a_0 = \mathbb{P}_\beta \left(d_{B_{2N}(\mathbf{0})}^* (B_N(\mathbf{0}), B_{2N}(\mathbf{0})^C) < \delta \Lambda(N) \right) \leq \frac{1}{2D}$$

is satisfied for small enough $\delta > 0$ and this shows (47) along the subsequence $N_k = M^k N$. To get the statement for all integer numbers, one can use Lemma 4.1 and the fact that $\Lambda(n) \leq \Lambda(mn) \leq m\Lambda(n)$ for all integers m, n . \square

With the same technique as above one can also prove that the indirect distance between V_0^n and the set $B_n(V_0^n)^C = \{x \in \mathbb{Z}^d : d_\infty(x, V_0^n) > n\}$ scales like n^θ .

Corollary 4.12. *For all $\beta \geq 0$ and $\varepsilon > 0$ there exist $0 < c_\varepsilon < C_\varepsilon < \infty$ such that*

$$\mathbb{P}_\beta \left(c_\varepsilon \Lambda(n, \beta) \leq d^* \left(V_0^n, B_n(V_0^n)^C \leq C_\varepsilon \Lambda(n, \beta) \right) \right) \geq 1 - \varepsilon.$$

5 The proof of Theorem 1.1

We first give an outline of the proof of Theorem 1.1. In Lemma 4.10 above we showed that $d(\mathbf{0}, B_n(\mathbf{0})^C) \approx_P \Lambda(n, \beta)$ and Lemma 4.2 shows that $\Lambda(n, \beta) \approx \mathbb{E}_\beta [d_{V_0^n}(\mathbf{0}, (n-1)e_1)]$, meaning that the ratio of these two expressions is uniformly bounded from below and above by constants $0 < c < C < \infty$. In Lemma 5.4 below we prove supermultiplicativity of $\Lambda(n, \beta)$. Together with the submultiplicativity proven in Lemma 2.3 this shows that for each $\beta \geq 0$ there exists $c_\beta > 0$ such that $c_\beta \Lambda(m, \beta) \Lambda(n, \beta) \leq \Lambda(mn, \beta) \leq \Lambda(m, \beta) \Lambda(n, \beta)$. We define $a_k = \log(\Lambda(2^k, \beta))$. The sequence is subadditive and thus

$$\theta(\beta) = \lim_{k \rightarrow \infty} \frac{\log(\Lambda(2^k, \beta))}{\log(2^k)} = \lim_{k \rightarrow \infty} \frac{a_k}{\log(2)k} = \inf_{k \in \mathbb{N}} \frac{a_k}{\log(2)k}$$

exists, where the last inequality holds because of Fekete's Lemma. On the other hand, the sequence $b_k = \log(c_\beta \Lambda(2^k, \beta))$ satisfies

$$b_{k+l} = \log(c_\beta \Lambda(2^{k+l}, \beta)) \geq \log(c_\beta \Lambda(2^k, \beta) c_\beta \Lambda(2^l, \beta)) = b_k + b_l$$

and thus

$$\theta(\beta) = \lim_{k \rightarrow \infty} \frac{\log(c_\beta \Lambda(2^k, \beta))}{\log(2^k)} = \lim_{k \rightarrow \infty} \frac{b_k}{\log(2)k} = \sup_{k \in \mathbb{N}} \frac{b_k}{\log(2)k}.$$

This already implies that

$$2^{k\theta(\beta)} \leq \Lambda(2^k, \beta) \leq c_\beta^{-1} 2^{k\theta(\beta)}$$

for all $k \in \mathbb{N}$. These two inequalities can be extended from points of the form 2^k to all integers with Lemma 4.1. So there exists a constant $0 < C'_\beta < \infty$ such that for all $n \in \mathbb{N}$

$$\frac{1}{C'_\beta} n^{\theta(\beta)} \leq \Lambda(n, \beta) \leq C'_\beta n^{\theta(\beta)}.$$

which shows (1). So we still need to prove supermultiplicativity of $\Lambda(\cdot, \beta)$ in order to prove the first item in Theorem 1.1. The second item of Theorem 1.1, i.e., the bounds on the diameter (2) of certain boxes, we show in section 5.3.

5.1 Distances between certain points

In this chapter, we examine the typical behavior of distances between points that are connected to long edges. In Lemma 5.1 we consider the infinity distance between such points. Using a coupling argument with the continuous model, we compare the situation to the situations occurring in Lemma 2.4 and Lemma 2.5. Then, in Lemma 5.2 we translate these bounds on the infinity distance into bounds on the typical graph distance between points that are incident to long edges.

Fix the three blocks V_u^n, V_w^n and V_0^n with $\|u\|_\infty \geq 2$. The next lemma deals with the infinity distance between points $x, y \in V_0^n$ with $x \sim V_u^n, y \sim V_w^n$.

Lemma 5.1. *For all $\frac{1}{n} < \varepsilon \leq \frac{1}{4}$ and $u, w \in \mathbb{Z}^d \setminus \{0\}$ with $\|u\|_\infty \geq 2$ one has*

$$\mathbb{P}_\beta(\exists x, y \in V_0^n : \|x - y\|_\infty \leq \varepsilon n, x \sim V_u^n, y \sim V_w^n \mid V_0^n \sim V_u^n, V_0^n \sim V_w^n) \leq C'_d \varepsilon^{1/2} [\beta]^2$$

where C'_d is a constant that depends only on the dimension d .

Proof. Let \mathcal{E} be the symmetrized Poisson process constructed in subsection 1.2 about the continuous model, i.e., $\tilde{\mathcal{E}}$ is a Poisson process on $\mathbb{R}^d \times \mathbb{R}^d$ with intensity $\frac{\beta}{2\|t-s\|^{2d}}$ and \mathcal{E} is defined by $\mathcal{E} := \left\{ (s, t) \in \mathbb{R}^d \times \mathbb{R}^d : (t, s) \in \tilde{\mathcal{E}} \right\} \cup \tilde{\mathcal{E}}$. Now we place an edge between $x, y \in \mathbb{Z}^d$ if and only if

$$(x + \mathcal{C}) \times (y + \mathcal{C}) \cap n\mathcal{E} \neq \emptyset$$

and call this graph $G = (V, E)$. The distribution of the resulting graph is identical to \mathbb{P}_β by the dilation invariance of \mathcal{E} . We can do the same procedure for $\lfloor \frac{1}{2\varepsilon} - 1 \rfloor \mathcal{E}$, i.e., place an edge between $x', y' \in \mathbb{Z}^d$ if and only if

$$(x' + \mathcal{C}) \times (y' + \mathcal{C}) \cap \left\lfloor \frac{1}{2\varepsilon} - 1 \right\rfloor \mathcal{E} \neq \emptyset$$

and call the resulting graph $G' = (V', E')$. Now assume that in the graph G there exist

$x, y \in V_{\mathbf{0}}^n$ with $\|x - y\|_{\infty} \leq \varepsilon n$ such that $x \sim V_u^n$ and $y \sim V_w^n$ in G . Then there exist

$$x_c \in x + \mathcal{C}, u_c \in nu + [0, n]^d, y_c \in y + \mathcal{C}, w_c \in nw + [0, n]^d$$

with $(x_c, u_c), (y_c, w_c) \in n\mathcal{E}$. We also have $\|x_c - y_c\|_{\infty} \leq \varepsilon n + 1 \leq 2\varepsilon n$. Now we rescale the process from size n to size $\lfloor \frac{1}{2\varepsilon} - 1 \rfloor$. For $(\tilde{x}_c, \tilde{u}_c) = \frac{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor}{n}(x_c, u_c)$ and $(\tilde{y}_c, \tilde{w}_c) = \frac{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor}{n}(y_c, w_c)$ we have

$$\begin{aligned} (\tilde{x}_c, \tilde{u}_c) &\in \left(\frac{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor}{n}x + \left[0, \frac{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor}{n}\right]^d \right) \times \left(\left\lfloor \frac{1}{2\varepsilon} - 1 \right\rfloor u + \left[0, \left\lfloor \frac{1}{2\varepsilon} - 1 \right\rfloor\right]^d \right) \cap \left\lfloor \frac{1}{2\varepsilon} - 1 \right\rfloor \mathcal{E}, \\ (\tilde{y}_c, \tilde{w}_c) &\in \left(\frac{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor}{n}y + \left[0, \frac{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor}{n}\right]^d \right) \times \left(\left\lfloor \frac{1}{2\varepsilon} - 1 \right\rfloor w + \left[0, \left\lfloor \frac{1}{2\varepsilon} - 1 \right\rfloor\right]^d \right) \cap \left\lfloor \frac{1}{2\varepsilon} - 1 \right\rfloor \mathcal{E}. \end{aligned}$$

From the rescaling we also have $\|\tilde{x}_c - \tilde{y}_c\|_{\infty} \leq 2\varepsilon \lfloor \frac{1}{2\varepsilon} - 1 \rfloor < 1$. So in particular there are vertices $x', y' \in \{0, \dots, \lfloor \frac{1}{2\varepsilon} - 1 \rfloor - 1\}^d$ with $x' \sim V_u^{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor}, y' \sim V_w^{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor}$ in G' , and $\|x' - y'\|_{\infty} \leq 1$. Write $N = \lfloor \frac{1}{2\varepsilon} - 1 \rfloor$. From (16) and (17) we get

$$\begin{aligned} \mathbb{P}_{\beta} \left(\exists x, y \in V_{\mathbf{0}}^N : \|x - y\|_{\infty} \leq 1, x \sim V_k^N, y \sim V_j^N \mid V_{\mathbf{0}}^N \sim V_k^N, V_{\mathbf{0}}^N \sim V_j^N \right) \\ \leq \frac{C_d \lceil \beta \rceil^2}{N^{1/2}} = \frac{C_d \lceil \beta \rceil^2}{\lfloor \frac{1}{2\varepsilon} - 1 \rfloor^{1/2}} \leq C'_d \varepsilon^{1/2} \lceil \beta \rceil^2 \end{aligned}$$

for some $C'_d < \infty$. With the coupling argument from before we thus also get

$$\begin{aligned} \mathbb{P}_{\beta} \left(\exists x, y \in V_{\mathbf{0}}^n : \|x - y\|_{\infty} \leq \varepsilon n, x \sim V_u^n, y \sim V_w^n \mid V_{\mathbf{0}}^n \sim V_u^n, V_{\mathbf{0}}^n \sim V_w^n \right) \\ = \mathbb{P}_{\beta} \left(\exists x, y \in V_{\mathbf{0}}^N : \|x - y\|_{\infty} \leq 1, x \sim V_u^N, y \sim V_w^N \mid V_{\mathbf{0}}^N \sim V_u^N, V_{\mathbf{0}}^N \sim V_w^N \right) \leq C'_d \varepsilon^{1/2} \lceil \beta \rceil^2. \end{aligned}$$

□

Lemma 5.2. *For all dimensions d and all β there exists a function $g_1(\varepsilon)$ with $g_1(\varepsilon) \xrightarrow{\varepsilon \rightarrow 0} 1$ such that for all $u, w \in \mathbb{Z}^d \setminus \{\mathbf{0}\}$ with $\|u\|_{\infty} \geq 2$ and all large enough $n \geq n(\varepsilon)$*

$$\mathbb{P}_{\beta}(d_{V_{\mathbf{0}}^n}(x, y) > \varepsilon \Lambda(n, \beta) \text{ for all } x, y \in V_{\mathbf{0}}^n \text{ with } x \sim V_u^n, y \sim V_w^n \mid V_u^n \sim V_{\mathbf{0}}^n \sim V_w^n) \geq g_1(\varepsilon).$$

Proof. We write $\mathbb{P}_{\beta}^{u, w}(\cdot)$ for the conditional probability measure $\mathbb{P}_{\beta}(\cdot \mid V_u^n \sim V_{\mathbf{0}}^n \sim V_w^n)$. As β is fixed throughout the rest of the proof, we write $\Lambda(n)$ for $\Lambda(n, \beta)$. We define the event

$$\begin{aligned} \mathcal{A}(K, \varepsilon_1, \varepsilon) &= \{\|x - y\|_{\infty} > \varepsilon_1 n \text{ for all } x, y \in V_{\mathbf{0}}^n \text{ with } x \sim V_u^n, y \sim V_w^n\} \\ &\cap \{d_{V_{\mathbf{0}}^n}(x, B_{\varepsilon_1 n}(x)^C) > \varepsilon \Lambda(n) \text{ for all } x \in V_{\mathbf{0}}^n \text{ with } x \sim V_u^n\} \cap \{|\{x \in V_{\mathbf{0}}^n : x \sim V_u^n\}| \leq K\} \end{aligned}$$

and observe that

$$\{d_{V_{\mathbf{0}}^n}(x, y) > \varepsilon \Lambda(n) \text{ for all } x, y \in V_{\mathbf{0}}^n \text{ with } x \sim V_u^n, y \sim V_w^n\} \supset \{\mathcal{A}(K, \varepsilon_1, \varepsilon)\}.$$

Thus it suffices to show that $\mathbb{P}_{\beta}^{u, w}(\mathcal{A}(K, \varepsilon_1, \varepsilon))$ converges to 1 as $\varepsilon \rightarrow 0$ for an appropriate choice of $K = K(\varepsilon), \varepsilon_1 = \varepsilon_1(\varepsilon)$. Respectively, that $\mathbb{P}_{\beta}^{u, w}(\mathcal{A}(K, \varepsilon_1, \varepsilon)^C)$ converges to 0.

We have that

$$\begin{aligned} \mathcal{A}(K, \varepsilon_1, \varepsilon)^C &= \{|\{x \in V_{\mathbf{0}}^n : x \sim V_u^n\}| > K\} \\ &\cup \{\|x - y\|_\infty < \varepsilon_1 n \text{ for some } x, y \in V_{\mathbf{0}}^n \text{ with } x \sim V_u^n, y \sim V_w^n\} \\ &\cup \left(\{d_{V_{\mathbf{0}}^n}(x, B_{\varepsilon_1 n}(x)^C) \leq \varepsilon \Lambda(n) \text{ for some } x \in V_{\mathbf{0}}^n \text{ with } x \sim V_u^n\} \cap \{|\{x \in V_{\mathbf{0}}^n : x \sim V_u^n\}| \leq K\} \right) \end{aligned}$$

and thus we get with Lemma 5.1

$$\begin{aligned} \mathbb{P}_\beta^{u,w}(\mathcal{A}(K, \varepsilon_1, \varepsilon)^C) &\leq \mathbb{P}_\beta^{u,w}(|\{x \in V_{\mathbf{0}}^n : x \sim V_u^n\}| > K) + C'_d \varepsilon_1^{1/2} [\beta]^2 \\ &+ \mathbb{P}_\beta^{u,w}(\{d_{V_{\mathbf{0}}^n}(x, B_{\varepsilon_1 n}(x)^C) > \varepsilon \Lambda(n) \text{ for all } x \in V_{\mathbf{0}}^n \text{ with } x \sim V_u^n\} \cap \{|\{x \in V_{\mathbf{0}}^n : x \sim V_u^n\}| \leq K\}) \\ &\leq \mathbb{P}_\beta^{u,w}(|\{x \in V_{\mathbf{0}}^n : x \sim V_u^n\}| > K) + C'_d \varepsilon_1^{1/2} [\beta]^2 + K \mathbb{P}_\beta^{u,w}(d_{V_{\mathbf{0}}^n}(\mathbf{0}, B_{\varepsilon_1 n}(\mathbf{0})^C) \leq \varepsilon \Lambda(n)). \end{aligned}$$

The expression $\mathbb{P}_\beta^{u,w}(|\{x \in V_{\mathbf{0}}^n : x \sim V_u^n\}| > K)$ converges to 0 for $K \rightarrow \infty$, by Markov's inequality, as one has the bound

$$\begin{aligned} \mathbb{E}_\beta[|\{x \in V_{\mathbf{0}}^n, z \in V_u^n : x \sim z\}|] &= \sum_{x \in V_{\mathbf{0}}^n} \sum_{z \in V_u^n} \mathbb{P}_\beta(x \sim z) \stackrel{(3)}{\leq} \sum_{x \in V_{\mathbf{0}}^n} \sum_{z \in V_u^n} \frac{\beta}{(\|x - z\|_\infty - 1)^{2d}} \\ &\leq \sum_{x \in V_{\mathbf{0}}^n} \sum_{z \in V_u^n} \frac{\beta}{((\|u\|_\infty - 1)n)^{2d}} \leq \frac{\beta 2^{2d}}{\|u\|_\infty^{2d}} \leq \beta 2^{2d} \end{aligned}$$

and thus Lemma 2.2 already gives that

$$\mathbb{E}_\beta^{u,w}[|\{x \in V_{\mathbf{0}}^n : x \sim V_u^n\}|] \leq \mathbb{E}_\beta^{u,w}[|\{x \in V_{\mathbf{0}}^n, z \in V_u^n : x \sim z\}|] \leq \beta 2^{2d} + 1$$

and this upper bound does not depend on n or u . Using Lemma 4.10, we see that for fixed $\varepsilon_1 > 0$ the term $\mathbb{P}_\beta(d_{V_{\mathbf{0}}^n}(\mathbf{0}, B_{\varepsilon_1 n}(\mathbf{0})^C) \leq \varepsilon \Lambda(n))$ converges to 0 as $\varepsilon \rightarrow 0$ and thus we can take $K = K(\varepsilon)$ and $\varepsilon_1 = \varepsilon_1(\varepsilon)$ that converge to $+\infty$, respectively 0, slow enough such that $K \mathbb{P}_\beta(d_{V_{\mathbf{0}}^n}(\mathbf{0}, B_{\varepsilon_1 n}(\mathbf{0})^C) \leq \varepsilon \Lambda(n))$ also converges to 0 for $\varepsilon \rightarrow 0$. \square

We want a similar function for the indirect distance between boxes. Such a function exists by Corollary 4.12.

Definition 5.3. Let $g_2(\varepsilon)$ be a function with $g_2(\varepsilon) \xrightarrow{\varepsilon \rightarrow 0} 1$ such that the indirect distance d^* between the sets $V_{\mathbf{0}}^n$ and $B_n(V_{\mathbf{0}}^n)^C$ satisfies

$$\mathbb{P}_\beta(d^*(V_{\mathbf{0}}^n, B_n(V_{\mathbf{0}}^n)^C) > \varepsilon \Lambda(n, \beta)) \geq g_2(\varepsilon)$$

for all $n \geq n(\varepsilon)$ large enough.

5.2 Supermultiplicativity of $\Lambda(n, \beta)$

In this section, we prove the supermultiplicativity of $\Lambda(n, \beta)$. Our main tools for this are the results of the previous section and Lemma 3.2. We also use the the same notation as in Lemma 3.2, i.e., $\mu_\beta = \mathbb{E}_\beta[\deg(\mathbf{0})]$ and $\bar{d}(Z) = \frac{1}{|Z|} \sum_{v \in Z} \deg(v)$. We apply this lemma mostly to paths instead of connected sets here, but this is fine, as every path is also a connected set.

Lemma 5.4. For all $\beta > 0$ there exists a constant $c > 0$ such that for all $m, n \in \mathbb{N}$

$$\Lambda(mn, \beta) \geq c\Lambda(n, \beta)\Lambda(m, \beta) \quad (49)$$

Proof. Inequality (49) holds for all small m or $n \in \mathbb{N}$ for some $c > 0$, so it suffices to consider m and n large enough. We split the graph $V_{\mathbf{0}}^{mn}$ into blocks of the form V_v^n for $v \in V_{\mathbf{0}}^m$. We call the graph that results from contracting all the blocks of the form V_v^n $G' = (V', E')$. Then G' has the same distribution as long range percolation on $V_{\mathbf{0}}^m$ under the measure \mathbb{P}_β . By $r(u)$ we denote the vertex in G' that results from contracting the box V_u^n . We also define an analogy of the infinity-distance on G' by $\|r(u) - r(v)\|_\infty = \|u - v\|_\infty$. Our goal is to bound the expected distance between the vertices $\mathbf{0}$ and $(mn - 1)e_1$ from below, conditioned on the graph G' . For this, we consider all loop-erased walks $P' = (r(u_0), r(u_1), \dots, r(u_l))$ between $r(\mathbf{0})$ and $r((m - 1)e_1)$ in G' . In the following we always work on a certain event \mathcal{H}_t , which is defined by

$$\mathcal{H}_t = \bigcap_{k \geq t} \{|\mathcal{CS}_k(G')| \leq 10^k \mu_\beta^k\} \cap \bigcap_{k \geq t} \{\bar{d}(Z) \leq 20\mu_\beta \forall Z \in \mathcal{CS}_k(G')\} .$$

Note that, by Lemma 3.2, (23), and Markov's inequality one has

$$\begin{aligned} \mathbb{P}_\beta(\mathcal{H}_t^C) &\leq \sum_{k=t}^{\infty} \mathbb{P}_\beta(|\mathcal{CS}_k(G')| > 10^k \mu_\beta^k) + \sum_{k=t}^{\infty} \mathbb{P}_\beta(\exists Z \in \mathcal{CS}_k(G') : \bar{d}(Z) > 20\mu_\beta) \\ &\leq \sum_{k=t}^{\infty} 0.4^k + \sum_{k=t}^{\infty} e^{-4\mu_\beta k} \leq \sum_{k=t}^{\infty} 0.5^k = 2 \cdot 2^{-t}. \end{aligned} \quad (50)$$

There are two possibilities for a path of length $l \geq t$ in G' . Either the path contains at least $\frac{l}{2}$ steps of the form $(r(u_i), r(u_{i+1}))$ with $\|u_{i+1} - u_i\|_\infty \geq 2$ or it does not, but then it contains at least $\frac{l}{2}$ steps of the form $(r(u_i), r(u_{i+1}))$ with $\|u_{i+1} - u_i\|_\infty = 1$. We write $\mathcal{P}_{1,l}$ for the set of paths of length l in G' that are of the first form, i.e., have at least $\frac{l}{2}$ steps of length at least 2 (in the ∞ -norm). And we write $\mathcal{P}_{2,l}$ for paths of the second form of length l . Let us focus on paths of the first form for the moment. For such a path $P = (r(u_0), \dots, r(u_l))$, let $r(u_i)$ satisfy $\|u_{i+1} - u_i\|_\infty \geq 2$. We call the block $V_{u_i}^n$ an ε -influential block if

$$d_{V_{u_i}^n}(x, y) > \varepsilon \Lambda(n) \text{ for all } x, y \in V_{u_i}^n \text{ with } x \sim V_{u_{i+1}}^n, y \sim V_w^n$$

for all $w \notin \{u_i, u_{i+1}\}$ with $V_w^n \sim V_{u_i}^n$. For a block V_v^n we define its degree by the number of blocks V_u^n such that $V_v^n \sim V_u^n$. So in particular, conditioned on the graph G' and the path taken inside G' , the probability that $V_{u_i}^n$ is not an ε -influential block is bounded by $\deg(V_{u_i}^n)(1 - g_1(\varepsilon))$ for all large enough $n \geq n(\varepsilon)$. Now let $\varepsilon > 0$ be small enough so that

$$20\mu_\beta (100\mu_\beta(1 - g_1(\varepsilon)))^{\frac{1}{1000\mu_\beta}} \leq \frac{1}{2} \quad (51)$$

and let $n \geq n(\varepsilon)$ be large enough. We call a block $V_{u_i}^n$ *influential*, if it is ε -influential, and we call the vertex $r(u_i)$ an *influential vertex* if the underlying block $V_{u_i}^n$ is an influential block. For a path $P = (r(u_0), r(u_1), \dots, r(u_l)) \in \mathcal{P}_{1,l}$ we define the canonical separated low degree independent set $\mathcal{SLI}(P) = \mathcal{SLI}_{l-1}(P) \subset P$ as follows:

0. Start with $\mathcal{SLI}_0(P) = \emptyset$.

1. For $i = 1, \dots, l-1$ check whether $r(u_i) \approx \mathcal{SLI}_{i-1}(P)$, $\deg(r(u_i)) \leq 100\mu_\beta$, and $\|u_i - u_{i+1}\|_\infty \geq 2$.

If all three events occur, then $\mathcal{SLI}_i(P) = \mathcal{SLI}_{i-1}(P) \cup \{r(u_i)\}$.

So in particular the set $\mathcal{SLI}(P)$ is an independent set. On the event \mathcal{H}_t , each path $P \in \mathcal{P}_{1,l}$ satisfies $|\mathcal{SLI}(P)| \geq \frac{l}{500\mu_\beta}$, which we will show now. The average degree of the vertices $r(u_i) \in P$ with $\|u_i - u_{i+1}\|_\infty \geq 2$ is at most $40\mu_\beta$ (as there are more than $l/2$ of them) and thus more than $\frac{l}{4}$ such vertices have a degree of at most $100\mu_\beta$. When included to the set $\mathcal{SLI}(P)$, such a vertex can block at most $100\mu_\beta$ many other vertices and thus we obtain that $|\mathcal{SLI}(P)| \geq \frac{1}{100\mu_\beta+1} \frac{l}{4} - 1 \geq \frac{l}{500\mu_\beta}$, where the last inequality holds for all large enough l , for example $l \geq 4000\mu_\beta$. The -1 is necessary as the last vertex $r(u_l)$ can, by construction, never be in $\mathcal{SLI}(P)$.

Our next claim is that for a fixed self-avoiding path $P \in \mathcal{P}_{1,l}$ in the graph G' there is only a very low probability that the number of influential vertices in $\mathcal{SLI}(P)$ is less than $\frac{l}{1000\mu_\beta}$. Note that for the different vertices $\{x : x \in \mathcal{SLI}(P)\}$, it is independent whether the vertices are influential or not as there are no edges between those vertices. So in particular, conditioned on the degree of V_v^n for all v , the probability that there are less than $\frac{|\mathcal{SLI}(P)|}{2}$ influential vertices is bounded by

$$2^{|\mathcal{SLI}(P)|} (100\mu_\beta(1 - g_1(\varepsilon)))^{\frac{|\mathcal{SLI}(P)|}{2}} \leq 2^l (100\mu_\beta(1 - g_1(\varepsilon)))^{\frac{l}{1000\mu_\beta}} .$$

where the last inequality holds on the event \mathcal{H}_t for $l \geq t$ and l large enough. From now on we write $\mathcal{INF}(P) \subseteq \mathcal{SLI}(P)$ for the set of influential vertices inside $\mathcal{SLI}(P)$. Remember that the event \mathcal{H}_t is very likely, in particular $\mathbb{P}_\beta(\mathcal{H}_t) \geq 0.5$ for all $t \geq 3$, as shown in (50). So with the previous estimates we see that for such $t \geq 3$

$$\mathbb{P}_\beta \left(\exists P \in \mathcal{P}_{1,l} : |\mathcal{INF}(P)| \leq \frac{l}{1000\mu_\beta} \mid \mathcal{H}_t \right) \leq 10^l \mu_\beta^l 2^l (100\mu_\beta(1 - g_1(\varepsilon)))^{\frac{l}{1000\mu_\beta}} \leq 2^{-l}$$

where the last inequality holds because of our assumption on ε in (51). So in particular with exponentially high probability in l , every path $P \in \mathcal{P}_{1,l}$ has more than $\frac{l}{1000\mu_\beta}$ influential vertices inside $\mathcal{SLI}(P)$. Whenever a path in the original model crosses such an influential block $V_{u_i}^n$, it needs to walk a distance of at least $\varepsilon\Lambda(n, \beta)$ inside of this block, as it needs to walk inside $V_{u_i}^n$ from a vertex that is adjacent to a different block to a vertex that is adjacent to $V_{u_{i+1}}^n$. So we get that for t large enough, conditioned on the event \mathcal{H}_t and for any $l \geq t$, all paths in the original model, for which the loop-erased projection on G' is in $\mathcal{P}_{1,l}$, have a length of $l \frac{\varepsilon}{1000\mu_\beta} \Lambda(n, \beta)$ with probability at least $1 - 2^{-l}$.

Now let us consider paths for which the loop-erased projection on G' is in $\mathcal{P}_{2,l}$. For a path $P = (r(u_0), \dots, r(u_l)) \in \mathcal{P}_{2,l}$ and a vertex $r(u_i) \in P$ with $\|u_{i+1} - u_i\|_\infty = 1$, we define the neighborhood $\mathcal{N}(r(u_i))$ by

$$\mathcal{N}(r(u_i)) = \{r(v) \in G' : \|v - u_i\|_\infty \leq 1\} .$$

The set

$$P \cup \bigcup_{\substack{i \in \{0, \dots, l-1\}: \\ \|u_i - u_{i+1}\|_\infty = 1}} \mathcal{N}(r(u_i))$$

has a cardinality of at most $3^d l$, so its average degree is also bounded from above by $20\mu_\beta$.

We define the neighborhood-degree of $r(u_i)$ by

$$\deg^{\mathcal{N}}(r(u_i)) = \sum_{v \in \mathcal{N}(r(u_i))} \deg(r(v)) .$$

We know that there are at least $\frac{l}{2}$ many vertices $r(u_i) \in P$ with $\|r(u_{i+1}) - r(u_i)\|_\infty = 1$ and thus we can deterministically choose at least $\frac{1}{5^d} \frac{l}{2}$ many such vertices that have $r(u_i)$ that have a distance of at least 3, i.e., $\|r(u_i) - r(u_j)\|_\infty \geq 3$ for those vertices. This is possible, as the condition $\|r(u_i) - r(u_j)\|_\infty \geq 3$ is equivalent to the condition $r(u_j) \notin \{r(v) \in G' : \|r(v) - r(u_i)\|_\infty \leq 2\}$ and the set $\{r(v) \in G' : \|r(v) - r(u_i)\|_\infty \leq 2\}$ has a cardinality of at most 5^d , so with every vertex that we choose, we can block at most $5^d - 1$ other such vertices. Call the set of those vertices $\mathcal{SE}(P)$. For these vertices $r(u_i)$ the sets $\mathcal{N}(r(u_i))$ are disjoint. Thus we get that $\sum_{u \in \mathcal{SE}(P)} \deg^{\mathcal{N}}(r(u)) \leq 3^d l 20\mu_\beta$ and this already implies that

$$\left| \left\{ r(u) \in \mathcal{SE}(P) : \deg^{\mathcal{N}}(r(u)) > 4 \cdot 5^d 3^d 20\mu_\beta \right\} \right| \leq \frac{l}{4 \cdot 5^d}$$

and thus we already have

$$\left| \left\{ r(u) \in \mathcal{SE}(P) : \deg^{\mathcal{N}}(r(u)) \leq 4 \cdot 5^d 3^d 20\mu_\beta \right\} \right| \geq \frac{l}{4 \cdot 5^d} .$$

So with an algorithm similar to the one above we can deterministically extract a set $\mathcal{SEI}(P) \subset \mathcal{SE}(P)$ such that

- $\mathcal{N}(r(u)) \cap \mathcal{N}(r(v)) = \emptyset$ for disjoint $r(u), r(v) \in \mathcal{SEI}(P)$,
- $\deg^{\mathcal{N}}(r(u)) \leq 4 \cdot 5^d 3^d 20\mu_\beta$ for all $r(u) \in \mathcal{SEI}(P)$, and
- $|\mathcal{SEI}(P)| \geq \frac{l}{25000^d \mu_\beta}$.

For each vertex $r(u_i) \in \mathcal{SEI}(P)$, the path does not exit the block $V_{u_i}^n$ through a long edge at the last visit, as it goes to $V_{u_{i+1}}^n$ afterwards and $\|u_{i+1} - u_i\|_\infty = 1$. So when the path leaves $V_{u_i}^n$ for the last time, it goes to $\mathcal{N}(r(u_i))$. In order to exit the set $\mathcal{N}(r(u_i)) \setminus \{r(u_i)\}$, there are two possibilities: either the path exits to some box V_v^n with $\|v - u_i\|_\infty = 2$ or it exits to some box V_v^n with $\|v - u_i\|_\infty > 2$. Given the graph G' , the probability that there exists a path of length at most $\varepsilon \Lambda(n, \beta)$ starting from $V_{u_i}^n$ that exits the set $\bigcup_{r(w) \in \mathcal{N}(r(u_i)) \setminus \{r(u_i)\}} V_w^n$ through a long edge to some box V_v^n with $\|v - u_i\|_\infty > 2$ is bounded by

$$\deg^{\mathcal{N}}(r(u_i)) (1 - g_1(\varepsilon)) \leq 80 \cdot 15^d (1 - g_1(\varepsilon)).$$

The probability that there exists a path of length at most $\varepsilon \Lambda(n, \beta)$ that exits the set $\mathcal{N}(r(u_i)) \setminus \{r(u_i)\}$ and into a block V_v^n with $\|v - u_i\|_\infty = 2$ is, given the graph G' , bounded by $1 - g_2(\varepsilon)$. This is true, as we can not use the long edges between $V_{u_i}^n$ and

$\bigcup_{r(v):\|v-u_i\|_\infty=2} V_v^n$ for this path, and the information about their presence does not reveal any information about the inside structure of the set $\bigcup_{r(v):\|v-u_i\|_\infty \leq 2} V_v^n$. So in total we see that

$$\mathbb{P} \left(d^* \left(V_{u_i}^n, \bigcup_{r(v):\|v-u_i\|_\infty \geq 2} V_v^n \right) < \varepsilon \Lambda(n, \beta) \mid G' \right) \leq 80 \cdot 15^d \mu_\beta (1 - g_1(\varepsilon)) + (1 - g_2(\varepsilon))$$

for all large enough n , and this holds for all $u_i \in \mathcal{SEI}(P)$. In the following, we will use the abbreviation

$$g'(\varepsilon) := \min \{ 80 \cdot 15^d \mu_\beta (1 - g_1(\varepsilon)) + (1 - g_2(\varepsilon)), 1 \}.$$

Note that $\lim_{\varepsilon \rightarrow 0} g'(\varepsilon) = 0$. Choose ε small enough such that $2^l g'(\varepsilon)^{\frac{l}{50000^d \mu_\beta}} \leq 20^{-l} \mu_\beta^{-l}$. We say that $r(u_i) \in \mathcal{SEI}(P)$, respectively the block V_v^n , is *influential* if

$$d^* \left(V_{u_i}^n, \bigcup_{r(v):\|v-u_i\|_\infty \geq 2} V_v^n \right) \geq \varepsilon \Lambda(n, \beta)$$

and we use the notation $\mathcal{INF}(P)$ for the set of influential vertices of $\mathcal{SEI}(P)$. The distances $d^* \left(V_{u_i}^n, \bigcup_{r(v):\|v-u_i\|_\infty \geq 2} V_v^n \right)$ are independent for different vertices $r(u) \in \mathcal{SEI}(P)$, as the sets $\mathcal{N}(r(u))$ are not connected. So in particular, given the set $\mathcal{SEI}(P)$, the probability that there are less than $\frac{|\mathcal{SEI}(P)|}{2}$ influential vertices in it is bounded by

$$2^{|\mathcal{SEI}(P)|} g'(\varepsilon)^{\frac{|\mathcal{SEI}(P)|}{2}} \leq 2^l g'(\varepsilon)^{\frac{l}{50000^d \mu_\beta}}.$$

Each path in the original model, for which the loop-erased projected walk is in $\mathcal{P}_{2,l}$, needs to cross at least $\frac{l}{2 \cdot 25000^d \mu_\beta}$ influential vertices with very high probability, and thus has a length of at least $\frac{l}{2 \cdot 25000^d \mu_\beta} \varepsilon \Lambda(n, \beta)$ with very high probability. On the event \mathcal{H}_t , this bound holds for all paths $P \in \mathcal{P}_{2,l}$ simultaneously with high probability, as the probability that there exists a path without this property is bounded by

$$|\mathcal{CS}_l(G')| 20^{-l} \mu_\beta^{-l} \leq 10^l \mu_\beta^l 20^{-l} \mu_\beta^{-l} = 2^{-l}.$$

Overall, we see that on the event \mathcal{H}_t , there exist constants c_1, C_2 such that $|\mathcal{INF}(P)| > c_1 \cdot l$ for all $P \in \mathcal{P}_l$ with probability at least $1 - C_2 \cdot 2^{-l}$. Note that the set $\mathcal{INF}(P)$ is defined differently, depending on whether $P \in \mathcal{P}_{1,l}$ or $P \in \mathcal{P}_{2,l}$. But the idea is always that for each vertex $r(u) \in \mathcal{INF}(P)$, one needs to walk a distance of order $\Lambda(n)$ in or near the block V_u^n in the original model. We define the event

$$\mathcal{G}_t = \bigcap_{l=t}^{\infty} \{ |\mathcal{INF}(P)| > c_1 l \text{ for all } P \in \mathcal{P}_l \} \cap \mathcal{H}_t$$

and note that

$$\mathbb{P}_\beta(\mathcal{G}_t^C) \leq C_3 2^{-t} \tag{52}$$

for some constant $C_3 < \infty$. For large enough t , say for all $t \geq K$, on the event $\mathcal{G}_t \cap$

$\{d_{G'}(r(\mathbf{0}), r((m-1)e_1)) \geq t\}$ every path between $r(\mathbf{0})$ and $r((m-1)e_1)$ needs to cross at least $c_1 \cdot t$ many influential vertices, with some constant $c_1 > 0$, and thus every path in the original model between $V_{\mathbf{0}}^n$ and $V_{(m-1)e_1}^n$ needs to walk a distance of at least $c_1 \varepsilon \Lambda(n, \beta)t$. So in total we get that for some large enough K

$$\begin{aligned} \mathbb{E}_\beta [d_{V_{\mathbf{0}}^{mn}}(\mathbf{0}, (mn-1)e_1)] &\geq \sum_{t=K}^{\infty} \mathbb{E}_\beta [d_{V_{\mathbf{0}}^{mn}}(\mathbf{0}, (mn-1)e_1) \mathbb{1}_{\mathcal{G}_t} \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((m-1)e_1))=t\}}] \\ &\geq c\varepsilon \Lambda(n, \beta) \sum_{t=K}^{\infty} t \mathbb{E}_\beta [\mathbb{1}_{\mathcal{G}_t} \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((m-1)e_1))=t\}}] \end{aligned} \quad (53)$$

and we can further bound the last sum by

$$\begin{aligned} &\sum_{t=K}^{\infty} t \mathbb{E}_\beta [\mathbb{1}_{\mathcal{G}_t} \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((m-1)e_1))=t\}}] \\ &= \sum_{t=K}^{\infty} t \mathbb{E}_\beta [\mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((m-1)e_1))=t\}}] - \sum_{t=K}^{\infty} t \mathbb{E}_\beta [\mathbb{1}_{\{\mathcal{G}_t^c\}} \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((m-1)e_1))=t\}}] \\ &= \sum_{t=K}^{\infty} t \mathbb{E}_\beta [\mathbb{1}_{\{d_{V_{\mathbf{0}}^m}(\mathbf{0}, (m-1)e_1)=t\}}] - \sum_{t=K}^{\infty} t \mathbb{E}_\beta [\mathbb{1}_{\{\mathcal{G}_t^c\}} \mathbb{1}_{\{d_{G'}(r(\mathbf{0}), r((m-1)e_1))=t\}}] \\ &\geq \sum_{t=1}^{\infty} t \mathbb{E}_\beta [\mathbb{1}_{\{d_{V_{\mathbf{0}}^m}(\mathbf{0}, (m-1)e_1)=t\}}] - \sum_{t=1}^{K-1} t \mathbb{E}_\beta [\mathbb{1}_{\{d_{V_{\mathbf{0}}^m}(\mathbf{0}, (m-1)e_1)=t\}}] - C_3 \sum_{t=K}^{\infty} t 2^{-t} \\ &\geq \mathbb{E}_\beta [d_{V_{\mathbf{0}}^m}(\mathbf{0}, (m-1)e_1)] - K - 2C_3 \geq c' \Lambda(m, \beta) \end{aligned}$$

for small enough $c' > 0$ and m large enough by Lemma 4.2. Inserting this into (53) finishes the proof. \square

5.3 The diameter of boxes

In this section, we prove the second item of Theorem 1.1, i.e., that the diameter of the box $\{0, \dots, n-1\}^d$ and its expectation both grow like n^θ .

Lemma 5.5. *For all $\beta \geq 0$ one has*

$$n^{\theta(\beta)} \approx_P \text{Diam}(\{0, \dots, n-1\}^d) \approx_P \mathbb{E}_\beta [\text{Diam}(\{0, \dots, n-1\}^d)].$$

Proof. By Lemma 4.1 it suffices to consider the case when $n = 2^k$. We have

$$\text{Diam}(\{0, \dots, n-1\}^d) \geq d_{V_{\mathbf{0}}^n}(\mathbf{0}, (n-1)e_1)$$

and this already implies that for each $\varepsilon > 0$ there exist $c, c_\varepsilon > 0$ such that

$$\mathbb{P}_\beta (c_\varepsilon n^{\theta(\beta)} < \text{Diam}(\{0, \dots, n-1\}^d)) > 1 - \varepsilon$$

and

$$c n^{\theta(\beta)} \leq \mathbb{E}_\beta [\text{Diam}(\{0, \dots, n-1\}^d)]$$

uniformly over n . For the upper bound, we make a dyadic decomposition of the box $V_{\mathbf{0}}^n$.

Similar ideas are also used in [20] for one dimension or in [4]. For a constant $S \geq 1$ we say that a box $V_y^{2^l} \subset V_0^{2^k}$ is S -good if

$$d_{V_y^{2^l}}(2^l y, 2^l y + (2^l - 1)e) \leq S \left(\frac{3}{2}\right)^{(l-k)\theta} 2^{k\theta}$$

for all $e \in \{0, 1\}^d$, where we simply write θ for $\theta(\beta)$ from here on. We use the notation

$$\Omega_l^S = \bigcap_{y \in V_0^{2^{k-l}}} \left\{ V_y^{2^l} \text{ is } S\text{-good} \right\} \quad \text{and} \quad \Omega^S = \bigcap_{l=1}^k \Omega_l^S.$$

On the event Ω^S , we can bound the graph distance between $\mathbf{0}$ and any $y \in V_0^{2^k}$ by considering a path that goes along the boxes in a dyadic decomposition. Let $y_0, \dots, y_k \in \mathbb{Z}^d$ be such that $y \in V_{y_i}^{2^i}$ for all i . So in particular $y_0 = y$ and $y_k = \mathbf{0}$. We also have that $V_{y_0}^{2^0} \subset V_{y_1}^{2^1} \subset \dots \subset V_{y_k}^{2^k}$ and thus also $2^{i-1}y_{i-1} \in V_{y_i}^{2^i}$ for all $i \geq 1$. This implies that $2^{i-1}y_{i-1} = 2^i y_i + 2^{i-1}e$ for some $e \in \{0, 1\}^d$. As all the boxes inside $V_0^{2^k}$ were assumed to be S -good we have

$$\begin{aligned} d_{V_0^{2^k}}(2^i y_i, 2^{i-1} y_{i-1}) &\leq d_{V_0^{2^k}}(2^i y_i, 2^i y_i + (2^{i-1} - 1)e) + 1 \\ &= d_{V_{2^{i-1}y_{i-1}}^{2^{i-1}}}(2^{i-1} 2y_i, 2^{i-1} 2y_i + (2^{i-1} - 1)e) + 1 \leq S \left(\frac{3}{2}\right)^{(i-1-k)\theta} 2^{k\theta} + 1. \end{aligned}$$

Now we have by the triangle inequality

$$\begin{aligned} d_{V_0^{2^k}}(\mathbf{0}, v) &\leq \sum_{l=1}^k \left(S \left(\frac{3}{2}\right)^{(l-1-k)\theta} 2^{k\theta} + 1 \right) \leq S 2^{k\theta} \sum_{l=1}^k \left(\left(\frac{3}{2}\right)^{(l-1-k)\theta} + \frac{1}{2^{k\theta}} \right) \\ &\leq C_\theta S 2^{k\theta} \end{aligned}$$

where the constant C_θ depends only on θ . As $d(u, v) \leq d(\mathbf{0}, u) + d(\mathbf{0}, v)$ for all $u, v \in V_0^{2^k}$, the previous bound already implies that on the event Ω^S one has

$$\text{Diam}(V_0^{2^k}) \leq 2C_\theta S 2^{k\theta} \tag{54}$$

and thus it suffices to bound the probability of $(\Omega^S)^C$. We know from Corollary 4.6 that the r -th moment of $d_{V_y^{2^l}}(2^l y, 2^l y + (2^l - 1)e)$ is of order $2^{r\theta}$, for all $r \geq 0$. So by a union bound and Markov's inequality we get that for every fixed box $V_y^{2^l}$

$$\begin{aligned} \mathbb{P}_\beta(V_y^{2^l} \text{ is not } S\text{-good}) &\leq \sum_{e \in \{0, 1\}^d} \mathbb{P}_\beta \left(d_{V_y^{2^l}}(2^l y, 2^l y + (2^l - 1)e) > S \left(\frac{3}{2}\right)^{(l-k)\theta} 2^{k\theta} \right) \\ &= \sum_{e \in \{0, 1\}^d} \mathbb{P}_\beta \left(d_{V_0^{2^l}}(\mathbf{0}, \mathbf{0} + (2^l - 1)e)^{\frac{4d}{\theta}} > S^{\frac{4d}{\theta}} \left(\frac{3}{2}\right)^{(l-k)4d} 2^{4dk} \right) \\ &\leq \sum_{e \in \{0, 1\}^d} \frac{\mathbb{E}_\beta \left[d_{V_0^{2^l}}(\mathbf{0}, \mathbf{0} + (2^l - 1)e)^{\frac{4d}{\theta}} \right]}{S^{\frac{4d}{\theta}} \left(\frac{3}{2}\right)^{(l-k)4d} 2^{4dk}} \leq \frac{C \cdot 2^{l\theta \frac{4d}{\theta}}}{S^{\frac{4d}{\theta}} \left(\frac{3}{2}\right)^{(l-k)4d} 2^{4dk}} = \frac{C \cdot 2^{4dl}}{S^{\frac{4d}{\theta}} \left(\frac{3}{2}\right)^{(l-k)4d} 2^{4dk}} \end{aligned}$$

$$\leq \frac{C}{S^{\frac{2d}{\theta}}} \left(\frac{2}{3}\right)^{(l-k)4d} 2^{(l-k)4d} = \frac{C}{S^{\frac{2d}{\theta}}} \left(\frac{4}{3}\right)^{(l-k)4d}$$

for some constant $C < \infty$. With another union bound that we get that

$$\begin{aligned} \mathbb{P}_\beta \left((\Omega_l^S)^C \right) &\leq \sum_{y \in V_0^{2^{k-l}}} \mathbb{P}_\beta \left(V_y^{2^l} \text{ is not } S\text{-good} \right) \leq \sum_{y \in V_0^{2^{k-l}}} \frac{C}{S^{\frac{2d}{\theta}}} \left(\frac{4}{3}\right)^{(l-k)4d} \\ &= \frac{C}{S^{\frac{2d}{\theta}}} 2^{(k-l)d} \left(\frac{4}{3}\right)^{(l-k)4d} = \frac{C}{S^{\frac{2d}{\theta}}} \left(\frac{81}{128}\right)^{(k-l)d} \end{aligned}$$

which implies that

$$\mathbb{P}_\beta \left((\Omega^S)^C \right) \leq \sum_{l=1}^k \frac{C}{S^{\frac{2d}{\theta}}} \left(\frac{81}{128}\right)^{(k-l)d} \leq \frac{C'}{S^{\frac{2d}{\theta}}} \quad (55)$$

for some constant $C' < \infty$. Together with (54) this proves that $\text{Diam} \left(V_0^{2^k} \right) \approx_P 2^{k\theta}$. Inequality (54) also implies that

$$\left\{ \frac{\text{Diam} \left(V_0^{2^k} \right)}{2^{k\theta}} > S \right\} \subset \left(\Omega^{\frac{S}{2C_\theta}} \right)^C$$

whenever $\frac{S}{2C_\theta} > 1$ and this implies that for some finite $K \in \mathbb{N}$ and all $k \in \mathbb{N}$

$$\begin{aligned} \mathbb{E}_\beta \left[\frac{\text{Diam} \left(V_0^{2^k} \right)}{2^{k\theta}} \right] &\leq K + \sum_{S=K}^{\infty} \mathbb{P}_\beta \left(\frac{\text{Diam} \left(V_0^{2^k} \right)}{2^{k\theta}} > S \right) \leq K + \sum_{S=K}^{\infty} \mathbb{P}_\beta \left(\left(\Omega^{\frac{S}{2C_\theta}} \right)^C \right) \\ &\leq K + \sum_{S=1}^{\infty} \frac{C' (2C_\theta)^{\frac{2d}{\theta}}}{S^{\frac{2d}{\theta}}} < \infty \end{aligned}$$

and the last term is finite, as $\frac{2d}{\theta} > 1$. This also shows that

$$\mathbb{E}_\beta \left[\text{Diam} \left(V_0^{2^k} \right) \right] = \mathcal{O} \left(2^{k\theta} \right)$$

and thus finishes the proof of Lemma 5.5. \square

6 Comparison with different inclusion probabilities

In this section, we compare the graph distances that result from percolation with measure \mathbb{P}_β to the graph distances that result from independent bond percolation on \mathbb{Z}^d where two vertices $u, v \in \mathbb{Z}^d$ are connected with probability $p^*(\beta, \{u, v\})$, which is assumed to be close enough to $p(\beta, \{u, v\})$. The precise condition required for the function $p^*(\beta, \{u, v\})$ is that for fixed β it satisfies that

$$p^*(\beta, \{u, v\}) = 1 \text{ for } \|u - v\| = 1 \text{ and } p^*(\beta, \{u, v\}) = p(\beta, \{u, v\}) + \mathcal{O} \left(\frac{1}{\|u - v\|^{2d+1}} \right) \quad (56)$$

as $\|u - v\| \rightarrow \infty$. An example of such a set of inclusion probabilities $p^*(\beta, \{u, v\})$ is given by

$$p^*(\beta, \{u, v\}) = \begin{cases} 1 & \text{for } \|u - v\| = 1 \\ \frac{\beta}{\|u - v\|^{2d}} \wedge 1 & \text{for } \|u - v\| > 1 \end{cases}$$

where we prove in Example 6.2 that (56) is satisfied. These inclusion probabilities were for example also used in [20] for $d = 1$.

We write \mathbb{P}_β^* for the probability measure resulting from independent bond percolation with inclusion probabilities $(p^*(\beta, \{u, v\}))_{u, v \in \mathbb{Z}^d}$ satisfying (56). We now give a proof that the graph distance $d(\mathbf{0}, x)$ also scales like $\|x\|^{\theta(\beta)}$ under the measure \mathbb{P}_β^* . We follow a technique that was already used in [20] in a similar form for a comparison between the discrete and the continuous model of percolation.

Theorem 6.1. *For fixed $\beta \geq 0$, suppose that $p^*(\beta, \{u, v\})$ satisfies (56). Then the graph distances between the origin $\mathbf{0}$ and $x \in \mathbb{Z}^d$ satisfies*

$$d(\mathbf{0}, x) \approx_P \|x\|^{\theta(\beta)}$$

under the measure \mathbb{P}_β^* .

Proof. We fix the dimension d and β from here on and consider them as constants. We write $E_{u,v}^*$ for the event when there exists an edge between u and v in the graph sampled with the measure \mathbb{P}_β^* , and we write $E_{u,v}$ if there exists an edge between u and v in the graph sampled with the measure \mathbb{P}_β . With the standard coupling for percolation we can couple the measures \mathbb{P}_β and \mathbb{P}_β^* so that uniformly over all $u \in \mathbb{Z}^d, v \in \mathbb{Z}^d \setminus \{u\}$

$$\mathbb{P}(E_{u,v}^* \setminus E_{u,v}) + \mathbb{P}(E_{u,v} \setminus E_{u,v}^*) \leq C_1 \frac{1}{\|u - v\|^{2d+1}}$$

where $C_1 < \infty$ is a constant. Thus we also get

$$\mathbb{P}\left(\left(E_{u,v}^*\right)^C \mid E_{u,v}\right) + \mathbb{P}\left(\left(E_{u,v}\right)^C \mid E_{u,v}^*\right) \leq C_2 \frac{1}{\|u - v\|}$$

for some constant $C_2 < \infty$. We write ω^* for the percolation configuration sampled by \mathbb{P}_β^* and ω for the percolation configuration sampled by \mathbb{P}_β . For two points $x, y \in \mathbb{Z}^d$, let P be a geodesic between x and y for the environment ω . We construct a path between x and y in the environment ω^* as follows:

- For $\{u, v\} \in P$, if $E_{u,v}^*$ occurs we use the direct edge between u and v .
- For $\{u, v\} \in P$, if $E_{u,v}^*$ does not occur go from u to v using $\|u - v\|_1$ many nearest-neighbor edges.

This gives a path P^* between x and y in the environment ω^* . The length of this path equals

$$\sum_{\substack{\{u,v\} \in P: \\ E_{u,v}^* \text{ occurs}}} 1 + \sum_{\substack{\{u,v\} \in P: \\ (E_{u,v}^*)^C \text{ occurs}}} \|u - v\|_1 = \sum_{\{u,v\} \in P} \left(\mathbb{1}_{E_{u,v}^*} + \|u - v\|_1 \mathbb{1}_{(E_{u,v}^*)^C} \right)$$

and thus we get that

$$\begin{aligned} \mathbb{E} [d(x, y; \omega^*) \mid \omega] &\leq \sum_{\{u, v\} \in P} \mathbb{E} \left[1 + \|u - v\|_1 \mathbb{1}_{(E_{u, v}^*)^c} \mid \omega \right] \\ &\leq \sum_{\{u, v\} \in P} \left(1 + \|u - v\|_1 C_2 \frac{1}{\|u - v\|} \right) \leq C_3 d(x, y; \omega) \end{aligned}$$

and this yields together with Markov's inequality for the conditional measure $\mathbb{P}(\cdot \mid \omega)$ that for each $\varepsilon > 0$ there exists a constant C_ε such that

$$\mathbb{P}(d(x, y; \omega^*) \leq C_\varepsilon d(x, y; \omega)) \geq 1 - \varepsilon .$$

Interchanging the roles of ω and ω^* one gets that for each $\varepsilon > 0$ there exists a constant C_ε^* such that

$$\mathbb{P}(d(x, y; \omega) \leq C_\varepsilon^* d(x, y; \omega^*)) \geq 1 - \varepsilon .$$

which finishes the proof. \square

Example 6.2. *The inclusion probabilities given by*

$$p^*(\beta, \{u, v\}) = \begin{cases} 1 & \text{for } \|u - v\| = 1 \\ \frac{\beta}{\|u - v\|^{2d}} \wedge 1 & \text{for } \|u - v\| > 1 \end{cases}$$

satisfy (56).

Proof. For each $x \in v + \mathcal{C}, y \in u + \mathcal{C}$ we have by the triangle inequality

$$\|u - v\| - \sqrt{d} \leq \|x - y\| \leq \|u - v\| + \sqrt{d}$$

and this already implies that for $\|u - v\| > \sqrt{d}$

$$\frac{1}{(\|u - v\| + \sqrt{d})^{2d}} \leq \int_{v+\mathcal{C}} \int_{u+\mathcal{C}} \frac{1}{\|x - y\|^{2d}} dy dx \leq \frac{1}{(\|u - v\| - \sqrt{d})^{2d}} .$$

With a Taylor expansion we see that

$$\begin{aligned} \frac{1}{\|u - v\| \pm \sqrt{d}} &= \frac{1}{\|u - v\|} \frac{1}{1 \pm \frac{\sqrt{d}}{\|u - v\|}} = \frac{1}{\|u - v\|} \left(1 + \mathcal{O}\left(\frac{1}{\|u - v\|}\right) \right) \\ &= \frac{1}{\|u - v\|} + \mathcal{O}\left(\frac{1}{\|u - v\|^2}\right) \end{aligned}$$

and raising this expression to the $2d$ -th power already gives that

$$\int_{v+\mathcal{C}} \int_{u+\mathcal{C}} \frac{1}{\|x - y\|^{2d}} dy dx = \frac{1}{\|u - v\|^{2d}} + \mathcal{O}\left(\frac{1}{\|u - v\|^{2d+1}}\right) \quad (57)$$

for $\|u - v\| \rightarrow \infty$. With the Taylor expansion of the exponential function we have $1 - e^{-s} = s + \mathcal{O}(s^2)$ for small s and thus by inserting (57) into the definition of $p(\beta, \{u, v\})$

we get

$$p(\beta, \{u, v\}) = \frac{\beta}{\|u-v\|^{2d}} + \mathcal{O}\left(\frac{1}{\|u-v\|^{2d+1}}\right) \quad (58)$$

which is equivalent to

$$p^*(\beta, \{u, v\}) = \frac{\beta}{\|u-v\|^{2d}} = p(\beta, \{u, v\}) + \mathcal{O}\left(\frac{1}{\|u-v\|^{2d+1}}\right).$$

□

Example 6.3. *The inclusion probabilities given by*

$$\tilde{p}(\beta, \{u, v\}) = \begin{cases} 1 & \text{for } \|u-v\| = 1 \\ 1 - e^{-\frac{\beta}{\|u-v\|^{2d}}} & \text{for } \|u-v\| > 1 \end{cases}.$$

satisfy (56).

Proof. By a Taylor expansion of the exponential function we get

$$1 - e^{-\frac{\beta}{\|u-v\|^{2d}}} = \frac{\beta}{\|u-v\|^{2d}} + \mathcal{O}\left(\frac{1}{\|u-v\|^{4d}}\right) = p^*(\beta, \{u, v\}) + \mathcal{O}\left(\frac{1}{\|u-v\|^{2d+1}}\right)$$

and we already know from Example 6.2 that $p^*(\beta, \{u, v\})$ satisfies (56). Thus we directly get that also $\tilde{p}(\beta, \{u, v\})$ satisfies (56). □

7 Appendix: Proofs for $d = 1$

In this appendix, we show a few lemmas for $d = 1$, where slightly different techniques compared to $d \geq 2$ are needed. It is well-known that for fixed $\beta < 1$ one has $\mathbb{E}_\beta [d(0, n)] = \Omega(n^{1-\beta})$. The next lemma gives a more uniform bound on the growth of $\mathbb{E}_\beta [d(0, n)]$ that holds for all $\beta \in [0, 1]$ simultaneously.

Lemma 7.1. *There exists a $c > 0$ such that for all $M, n \in \mathbb{N}$ and $\beta \in [0, 1]$*

$$\mathbb{E}_\beta [d_{[0, Mn-1]}(0, Mn-1)] \geq cM^{1-\beta} \mathbb{E}_\beta [d_{[0, n-1]}(0, n-1)]. \quad (59)$$

Proof. First note that the proof of (39) does not depend on a uniform bound on the second moment, so we can safely apply it in our argumentation here. By (39) we can choose $\iota > 0$ small enough so that

$$\mathbb{E}_\beta [d_{[0, n-1]}([0, \iota n], [n - \iota n - 1, n - 1])] \geq \frac{1}{2} \mathbb{E}_\beta [d_{[0, n-1]}(0, n-1)]$$

uniformly over $\beta \in [0, 1]$. This implies the existence a $c' > 0$ such that

$$\mathbb{E}_\beta [d_{[-n, 2n-1]}(V_{-1}^n, V_1^n)] \geq c' \mathbb{E}_\beta [d_{[0, n-1]}(0, n-1)] \quad (60)$$

uniformly over $\beta \in [0, 1]$ and $n \in \mathbb{N}$ large enough, as we will argue now. For fixed $\iota > 0$ there is a uniform positive probability (in $\beta \in [0, 1]$ and $n \in \mathbb{N}$) that the rightmost vertex




Figure 4: The vertex w is a separation point if all edges e with $|e| \geq 2$ are either strictly to the left or right of w , as above.

incident to V_{-1}^n lies inside $[0, \iota n]$ and that the leftmost vertex incident to V_1^n lies inside $[n - \iota n - 1, n - 1]$. Call this event A . Whenever the event A holds, one already has

$$d_{[-n, 2n-1]}(V_{-1}^n, V_1^n) \geq d_{[0, n-1]}([0, \iota n], [n - \iota n - 1, n - 1])$$

and as both the event A and the distance $d_{[0, n-1]}([0, \iota n], [n - \iota n - 1, n - 1])$ are decreasing one has by the FKG inequality

$$\begin{aligned} \mathbb{E}_\beta [d_{[-n, 2n-1]}(V_{-1}^n, V_1^n)] &\geq \mathbb{E}_\beta [d_{[-n, 2n-1]}(V_{-1}^n, V_1^n) \mathbb{1}_A] \\ &\geq \mathbb{E}_\beta [d_{[0, n-1]}([0, \iota n], [n - \iota n - 1, n - 1]) \mathbb{1}_A] \\ &\geq \mathbb{E}_\beta [d_{[0, n-1]}([0, \iota n], [n - \iota n - 1, n - 1])] \mathbb{P}_\beta(A) \geq \frac{\mathbb{P}_\beta(A)}{2} \mathbb{E}_\beta [d_{[0, n-1]}(0, n - 1)] \end{aligned}$$

which shows (60). For long-range percolation on the line segment $\{0, \dots, M - 1\}$, we call an odd point $w \in \{1, \dots, M - 2\}$ a *separation point* if $w \asymp \{0, \dots, w - 2\}$, $w \asymp \{w + 2, \dots, M - 1\}$ and $\{0, \dots, w - 1\} \asymp \{w + 1, \dots, M - 1\}$; See Figure 4 for an illustration. Even points can simply never be separation points with our definition. These three events are independent and we can bound the probability of the first event by

$$\mathbb{P}_\beta (w \asymp \{0, \dots, w - 2\}) \geq e^{-\beta \int_{-\infty}^0 \int_1^2 \frac{1}{|t-s|^2} dt ds} \geq e^{-1}.$$

The same proof also works for the second event and shows that $\mathbb{P}_\beta (w \asymp \{w + 2, \dots, M - 1\}) \geq e^{-1}$ for all $\beta \in [0, 1]$. The probability of the event $\{0, \dots, w - 1\} \asymp \{w + 1, \dots, M - 1\}$ can be bounded by

$$\begin{aligned} \prod_{0 \leq u < w} \prod_{w < v \leq M-1} e^{-\beta \int_u^{u+1} \int_v^{v+1} \frac{1}{|x-y|^2} dx dy} &= e^{-\beta \int_0^w \int_{w+1}^M \frac{1}{|x-y|^2} dx dy} \\ &\geq e^{-\beta \int_0^w \int_{w+1}^\infty \frac{1}{|x-y|^2} dx dy} = e^{-\beta \int_0^w \frac{1}{w+1-y} dy} = e^{-\beta \log(w+1)} \geq M^{-\beta}. \end{aligned}$$

uniformly over $\beta \in [0, 1]$. Now, using the independence of the three relevant events we get that

$$\begin{aligned} \mathbb{P}_\beta (w \text{ is a separation point}) &= \mathbb{P}_\beta (w \asymp \{0, \dots, w - 2\}) \cdot \mathbb{P}_\beta (w \asymp \{w + 2, \dots, M - 1\}) \\ &\quad \cdot \mathbb{P}_\beta (\{0, \dots, w - 1\} \asymp \{w + 1, \dots, M - 1\}) \geq e^{-2} M^{-\beta} \geq 0.1 M^{-\beta}. \end{aligned}$$

For w odd, we call the set V_w^n a *separation interval* if $V_w^n \asymp [0, (w - 1)n - 1]$, $V_w^n \asymp [(w + 2)n, Mn - 1]$ and $\{0, \dots, wn - 1\} \asymp \{(w + 1)n, \dots, Mn - 1\}$. Again, an even w can

never be a separation interval. By the scaling invariance of the underlying continuous model, the probability that V_w^n is a separation interval is exactly the probability that w is a separation point for the line segment $\{0, \dots, M-1\}$ and this probability is bounded from below by $0.1M^{-\beta}$. Let $w_1, \dots, w_l \in \{1, \dots, M-2\}$ be integers such that $V_{w_i}^n$ is a separation interval for all i . Then each path between 0 and $Mn-1$ in the graph $\{0, \dots, Mn-1\}$ needs to cross all separation intervals of this form and in particular

$$d_{[0, Mn-1]}(0, Mn-1) \geq \sum_{i=1}^l d_{[(w_i-1)n, (w_i+2)n-1]}(V_{w_i-1}^n, V_{w_i+1}^n).$$

A block V_w^n being a separation interval and the inside structure of $[(w-1)n, (w+2)n-1]$ are independent. The fact that V_w^n is a separation interval reveals no information about the edges with both endpoints in $\{(w-1)n, \dots, (w+2)n-1\}$, except that there is no direct edge from $\{(w-1)n, \dots, wn-1\}$ to $\{(w+1)n, \dots, (w+2)n-1\}$. Thus, by taking expectations in the above inequality and using that both the event $\{V_w^n \text{ is a sep. int.}\}$ and the random distance $d_{[(w-1)n, (w+2)n-1]}(V_{w-1}^n, V_{w+1}^n)$ are decreasing, we get by the FKG-inequality

$$\begin{aligned} \mathbb{E}_\beta [d_{[0, Mn-1]}(0, Mn-1)] &\geq \mathbb{E}_\beta \left[\sum_{w=1}^{M-2} \mathbb{1}_{\{V_w^n \text{ is a sep. int.}\}} d_{[(w-1)n, (w+2)n-1]}(V_{w-1}^n, V_{w+1}^n) \right] \\ &\geq \sum_{w=1}^{M-2} \mathbb{E}_\beta [\mathbb{1}_{\{V_w^n \text{ is a sep. int.}\}}] \mathbb{E}_\beta [d_{[-n, 2n-1]}(V_{-1}^n, V_1^n)] \\ &\stackrel{(60)}{\geq} \sum_{\substack{w \in \{1, \dots, M-2\}: \\ w \text{ odd}}} 0.1M^{-\beta} c' \mathbb{E}_\beta [d_{[0, n-1]}(0, n-1)] \geq cM^{1-\beta} \mathbb{E}_\beta [d_{[0, n-1]}(0, n-1)] \end{aligned}$$

for some small constant $c > 0$ and M large enough. For small M , the statement holds for a small enough c by Lemma 4.1. \square

With this we are now ready to go to the proof of Lemma 4.5 for $d = 1$.

Proof of Lemma 4.5 for $d = 1$. We say that the vertex $w \in \{1, \dots, m-2\}$ is a cut point (for the interval $\{0, \dots, m-1\}$) if there exists no edge of the form $\{u, v\}$ with $0 \leq u < w < v \leq m-1$. For $w < \frac{m}{2}$ and $\beta \leq 2$ we have

$$\begin{aligned} \mathbb{P}_\beta(w \text{ is a cut point}) &= \prod_{0 \leq u < w} \prod_{w < v \leq m-1} e^{-\beta \int_u^{u+1} \int_v^{v+1} \frac{1}{|x-y|^2} dx dy} = e^{-\beta \int_0^w \int_{w+1}^m \frac{1}{|x-y|^2} dx dy} \\ &\leq e^{-\beta \int_0^w \int_{w+1}^{2w+1} \frac{1}{|x-y|^2} dx dy} = e^{-\beta \int_0^w \frac{1}{w+1-y} - \frac{1}{2w+1-y} dy} \\ &= e^{-\beta(-\log(1)+2\log(w+1)-\log(2w+1))} = e^{-\beta \log\left(\frac{(w+1)^2}{2w+1}\right)} \leq e^{-\beta \log\left(\frac{w+1}{2}\right)} \\ &\leq e^{-\beta \log(w+1)} e^{\beta \log(2)} \leq 4w^{-\beta} \end{aligned}$$

and with this we get, by linearity of expectation and symmetry of the process, that

$$\mathbb{E}_\beta [|\{w \in \{1, \dots, m-2\} : w \text{ is a cut point}\}|] \leq 1 + 2 \sum_{w=1}^{\lfloor m/2 \rfloor} \mathbb{P}_\beta(w \text{ is a cut point})$$




Figure 5: The long edges inside the box $\{0, \dots, 7^{n+1} - 1\}$. The set \mathcal{B} are the bold black edges.

$$\leq 1 + 8 \sum_{w=1}^{\lfloor m/2 \rfloor} w^{-\beta} \leq 10 + 8 \int_1^m w^{-\beta} dw = \begin{cases} 10 + 8 \left[\frac{w^{-\beta+1}}{-\beta+1} \right]_1^m & \beta \in [0, 2] \setminus \{1\} \\ 10 + 8 \log(m) & \beta = 1 \end{cases}.$$

As the expected number of cut points is monotone decreasing in β we get that for the function $f(\beta, m) := \mathbb{E}_\beta [|\{w \in \{1, \dots, m-2\} : w \text{ is a cut point}\}|]$ we have the upper bound

$$f(\beta, m) \leq \begin{cases} \frac{20}{1-\beta} m^{1-\beta} & \beta < 1 \\ 10 + 8 \log(m) & 1 \leq \beta \leq 2 \\ 20 & \beta > 2 \end{cases}. \quad (61)$$

We now use a method (that was already used in [20] in a similar form for the continuous model) in order to bound the second moment of $d_{V_0^{m^{n+1}}}(0, m^{n+1} - 1)$. We say that an interval $V_k^{m^n}$ is *unbridged* if there exists no edge $\{u, v\}$ with both endpoints in $\{0, \dots, m^{n+1} - 1\}$ and $u < km^n, v \geq (k+1)m^n$; Contrary, if there exists such an edge we say that the interval is *bridged*. In this case, we also say that the interval is bridged by the edge $\{u, v\}$. So clearly the intervals $V_0^{m^n}, V_{m-1}^{m^n}$ are unbridged and the probability that $V_w^{m^n}$ is unbridged for $w \in \{1, \dots, m-2\}$ is exactly the probability that w is a cut point for the interval $\{0, \dots, m-1\}$. We now define a set of edges \mathcal{B} as follows: Let $i < j \in \{0, \dots, m-1\}$ with $|i-j| > 1$ satisfy $V_i^{m^n} \sim V_j^{m^n}$ and $V_{i-l_1}^{m^n} \approx V_{j+l_2}^{m^n}$ for all $(l_1, l_2) \in \{0, \dots, i\} \times \{0, \dots, m-1-j\} \setminus \{(0,0)\}$. In this situation, we add one edge between $V_i^{m^n}$ and $V_j^{m^n}$ to \mathcal{B} . If there are several edges between $V_i^{m^n}$ and $V_j^{m^n}$ we choose the left-most shortest such edge (this rule is arbitrary, any deterministic rule would work here). An example of this construction is given in Figure 5. So the set \mathcal{B} is the set of possible bridges where we already delete edges that are furthermore bridged by even longer edges. With this construction, we get $|\mathcal{B}| \leq m$, as each interval $V_j^{m^n}$ can be adjacent to at most two edges in \mathcal{B} and each edge in \mathcal{B} touches two intervals. Furthermore, if an interval $V_j^{m^n}$ is bridged, then there exists an edge $e \in \mathcal{B}$ so that $V_j^{m^n}$ is bridged by e . Let \mathcal{U}' be the set of endpoints of edges in \mathcal{B} and let

$$\mathcal{U} := \mathcal{U}' \cup \{0, m^n, \dots, (m-1)m^n\} \cup \{m^n - 1, 2m^n - 1, \dots, m^{n+1} - 1\}.$$

Let $\mathcal{U} = \{x_0, x_1, \dots, x_u\}$, where $x_0 < \dots < x_u$. By the construction we have $|\mathcal{U}| \leq 4m$ and $|x_{i+1} - x_i| \leq m^n - 1$. For x_{i-1}, x_i with $(x_{i-1}, x_i) \neq (km^n - 1, km^n)$ for all k we say that $[x_i, x_{i+1}]$ is bridged, if there exists an edge $\{u, v\} \in \mathcal{B}$ with $u \leq x_i < x_{i+1} \leq v$. Assume we have (x_{i-1}, x_i) which is not of the form $(km^n - 1, km^n)$, say with $[x_i, x_{i+1}] \subset V_j^{m^n}$ for some $j \in \{0, \dots, m-1\}$, and $[x_i, x_{i+1}]$ is not bridged, then also $V_j^{m^n}$ is not bridged. On the other hand, if $[x_i, x_{i+1}]$ is bridged, then also $V_j^{m^n}$ is bridged by some edge in \mathcal{B} .

In each interval $V_j^{m^n}$ there are at most two points in $V_j^{m^n} \cap \mathcal{U}$ that come from endpoints of edges in \mathcal{B} ; Furthermore, the two endpoints of the interval are also in $V_j^{m^n} \cap \mathcal{U}$. So in total there are at most 4 points in $V_j^{m^n} \cap \mathcal{U}$ for all $j \in \{0, \dots, m-1\}$, and thus there are at most 3 intervals $[x_i, x_{i+1}]$ inside each $V_j^{m^n}$. This already implies that

$$|\{i \in \{1, \dots, u\} : [x_{i-1}, x_i] \text{ is not bridged}\}| \leq 3 \left| \{j \in \{0, \dots, m-1\} : V_j^{m^n} \text{ is not bridged}\} \right|. \quad (62)$$

Now we want to construct a path between 0 and $m^{n+1} - 1$. Let

$$\tau = \arg \max_{i \in \{1, \dots, u\}} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i).$$

If there are multiple maximizers, we pick one with $x_i \neq km^n$ for all k and with minimal x_i among those maximizers. So in particular $[x_{i-1}, x_i]$ always lies inside some interval $V_j^{m^n}$. If $[x_{\tau-1}, x_\tau]$ is bridged by some edge $e = \{x_{\tau_1}, x_{\tau_2}\} \in \mathcal{B}$, say with $x_{\tau_1} < x_{\tau_2}$, then we consider the path that goes from $0 = x_0$ to x_{τ_1} , then directly jumps to x_{τ_2} and from there goes to $x_u = m^{n+1} - 1$. This implies that

$$\begin{aligned} d_{[0, m^{n+1}-1]}(0, m^{n+1} - 1) &\leq \sum_{i=1}^{\tau_1} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) + 1 + \sum_{i=\tau_2+1}^u d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \\ &\leq u \max_{i \neq \tau} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \leq 4m \max_{i \neq \tau} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \end{aligned}$$

in this case. For the case where $[x_{\tau-1}, x_\tau]$ is not bridged, we consider the path that goes iteratively from x_0 to x_u . Here we have

$$\begin{aligned} &d_{[0, m^{n+1}-1]}(0, m^{n+1} - 1) \\ &\leq \sum_{i=1}^{\tau-1} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) + d_{[x_{\tau-1}, x_\tau]}(x_{\tau-1}, x_\tau) + \sum_{i=\tau+1}^u d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \\ &\leq 4m \max_{i \neq \tau} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) + \max_{[x_{i-1}, x_i] \text{ not bridged}} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \end{aligned}$$

and thus we have in both cases that

$$\begin{aligned} &(d_{[0, m^{n+1}-1]}(0, m^{n+1} - 1))^2 \\ &\leq 2 \left(4m \max_{i \neq \tau} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \right)^2 + 2 \left(\max_{[x_{i-1}, x_i] \text{ not bridged}} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \right)^2 \\ &\leq 32m^2 \left(\max_{i \neq \tau} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \right)^2 + 2 \sum_{[x_{i-1}, x_i] \text{ not bridged}} (d_{[x_{i-1}, x_i]}(x_{i-1}, x_i))^2. \quad (63) \end{aligned}$$

Now we want to bound both terms in the above sum in expectation. To bound the first term, we use the following observation: If $X_1, \dots, X_{\bar{m}}$ are independent non-negative

random variables and $\tau = \arg \max_{i \in \{1, \dots, \tilde{m}\}}$, then

$$\mathbb{E} \left[\left(\max_{i \neq \tau} X_i \right)^2 \right] \leq \mathbb{E} \left[\sum_{i=1}^{\tilde{m}} X_i \left(\sum_{j \neq i} X_j \right) \right] = \sum_{i=1}^{\tilde{m}} \sum_{j \neq i} \mathbb{E} [X_i] \mathbb{E} [X_j] \leq \tilde{m}^2 \max_i \mathbb{E} [X_i]^2. \quad (64)$$

Conditioned on \mathcal{U} , the random variables $d_{[x_{i-1}, x_i]}(x_{i-1}, x_i)$ are independent and by Lemma 4.1 the second moment of them is bounded by the second moment of $(d_{V_0^{m^n}}(0, m^n - 1))$, up to a factor of $9 = 3^2$. As $u \leq 4m$ we get with (64) and Lemma 4.1 that

$$\begin{aligned} \mathbb{E}_\beta \left[\max_{i \neq \tau} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i)^2 \right] &= \mathbb{E}_\beta \left[\mathbb{E} \left[\max_{i \neq \tau} d_{[x_{i-1}, x_i]}(x_{i-1}, x_i)^2 \mid \mathcal{U} \right] \right] \\ &\leq \mathbb{E}_\beta \left[u^2 \mathbb{E} \left[\max_i d_{[x_{i-1}, x_i]}(x_{i-1}, x_i) \mid \mathcal{U} \right]^2 \right] \leq 144m^2 \mathbb{E}_\beta \left[d_{V_0^{m^n}}(0, m^n - 1) \right]^2. \end{aligned} \quad (65)$$

In order to bound the second summand in (63) in expectation we use the bound on the number of unbridged segments (62). With this we get that

$$\begin{aligned} &\mathbb{E}_\beta \left[\sum_{[x_{i-1}, x_i] \text{ not bridged}} (d_{[x_{i-1}, x_i]}(x_{i-1}, x_i))^2 \right] \\ &= \mathbb{E}_\beta \left[\mathbb{E}_\beta \left[\sum_{[x_{i-1}, x_i] \text{ not bridged}} (d_{[x_{i-1}, x_i]}(x_{i-1}, x_i))^2 \mid \mathcal{U} \right] \right] \\ &\leq 9 \mathbb{E}_\beta \left[d_{V_0^{m^n}}(0, m^n - 1)^2 \right] \mathbb{E}_\beta \left[\sum_{[x_{i-1}, x_i] \text{ not bridged}} 1 \right] \\ &\leq 27 \mathbb{E}_\beta \left[d_{V_0^{m^n}}(0, m^n - 1)^2 \right] \mathbb{E}_\beta \left[\left| \{j \in \{0, \dots, m-1\} : V_j^{m^n} \text{ unbridged}\} \right| \right] \\ &\leq 27 \mathbb{E}_\beta \left[d_{V_0^{m^n}}(0, m^n - 1)^2 \right] (2 + f(\beta, m)) =: \mathbb{E}_\beta \left[d(0, m^n - 1)^2 \right] \tilde{f}(\beta, m), \end{aligned} \quad (66)$$

where $\tilde{f}(\beta, m) = 27(2 + f(\beta, m))$. Combining (65) and (66) and inserting it into (63) we obtain that

$$\begin{aligned} \mathbb{E}_\beta \left[d_{V_0^{m^{n+1}}}(0, m^{n+1} - 1)^2 \right] &\leq 5000m^4 \mathbb{E}_\beta \left[d_{V_0^{m^n}}(0, m^n - 1) \right]^2 \\ &\quad + 2\tilde{f}(\beta, m) \mathbb{E}_\beta \left[d_{V_0^{m^n}}(0, m^n - 1)^2 \right] \end{aligned}$$

and iterating this inequality over all $k = 1, \dots, n$ we get

$$\mathbb{E}_\beta \left[d_{V_0^{m^{n+1}}}(0, m^{n+1} - 1)^2 \right] \leq 5000m^4 \sum_{k=1}^n \left(2\tilde{f}(\beta, m) \right)^{n+1-k} \mathbb{E}_\beta \left[d_{V_0^{m^k}}(0, m^k - 1) \right]^2. \quad (67)$$

Using the bounds on $f(\beta, m)$ from (61) we see that function $\tilde{f}(\beta, m)$ clearly satisfies

$$\tilde{f}(\beta, m) = 27(2 + f(\beta, m)) \leq \begin{cases} \frac{600}{1-\beta} m^{1-\beta} & \beta < 1 \\ 600(1 + \log(m)) & 1 \leq \beta \leq 2 \\ 600 & \beta > 2 \end{cases} .$$

By compactness of each interval $[\beta, \beta + 1]$ it suffices to show that the uniform bound on the second moment (32) holds for all $\beta > 0$ and $\varepsilon \in (-c_\beta, c_\beta)$ for some $c_\beta > 0$ small enough, respectively for $\beta = 0$ and $\varepsilon \in [0, c_\beta)$. To extend the inequality from open sets to compact intervals, one can cover each compact interval with finitely many open sets and then take the largest among these finitely many constants that arose from this procedure. So we are left to show that for all $\beta \geq 0$ there exists a constant $C_\beta < \infty$ such that for all $n \in \mathbb{N}$, all $\varepsilon \in (-c_\beta, c_\beta)$, respectively all $\varepsilon \in [0, c_\beta)$ for $\beta = 0$, and all $u, v \in V_0^n$

$$\mathbb{E}_{\beta+\varepsilon} [d_{V_0^n}(u, v)^2] \leq C_\beta \Lambda(n, \beta + \varepsilon)^2. \quad (68)$$

We start with the case $\beta \geq 1$. By Remark 4.4, there exists a $\theta' = \theta'(\beta) > 0$ such that

$$\mathbb{E}_{\beta+\varepsilon} [d_{V_0^{m^{n+1}}}(0, m \cdot m^n - 1)] \geq m^{\theta'} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^{m^n}}(0, m^n - 1)]$$

for all $n \in \mathbb{N}$, m large enough, and $|\varepsilon| \leq \frac{1}{2}$. Inserting this into (67) we get

$$\begin{aligned} & \mathbb{E}_\beta [d_{V_0^{m^{n+1}}}(0, m^{n+1} - 1)^2] \\ & \leq 5000m^4 \sum_{k=1}^n \left(2\tilde{f}(\beta, m)\right)^{n+1-k} \left(m^{-2\theta'}\right)^{n-k} \mathbb{E}_\beta [d_{V_0^{m^n}}(0, m^n - 1)]^2. \end{aligned}$$

Now choose $m \in \mathbb{N}$ large enough and $0 < c_\beta < 0.1$ small enough so that $2\tilde{f}(\beta + \varepsilon, m)m^{-2\theta'(\beta)} \leq 0.5$ for all $\varepsilon \in (-c_\beta, c_\beta)$. This is clearly possible for $\beta > 1$. For $\beta = 1$, we can choose c_β small enough so that $c_\beta < \theta'(1)$ and by monotonicity in the first argument of the function $\tilde{f}(\cdot, \cdot)$ one then has $\tilde{f}(1 + \varepsilon, m) \leq \frac{600}{c_\beta} m^{c_\beta}$ for all $\varepsilon \in (-c_\beta, c_\beta)$, which shows that one can find m, c_β so that $2\tilde{f}(1 + \varepsilon, m)m^{-2\theta'(1)} \leq 0.5$ for all $\varepsilon \in (-c_\beta, c_\beta)$. This then gives that

$$\begin{aligned} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^{m^{n+1}}}(0, m^{n+1} - 1)^2] & \leq 10000\tilde{f}(\beta - c_\beta, m)m^4 \sum_{k=1}^n 0.5^{n-k} \mathbb{E}_{\beta+\varepsilon} [d_{V_0^{m^n}}(0, m^n - 1)]^2 \\ & \leq 20000\tilde{f}(\beta - c_\beta, m)m^4 \mathbb{E}_{\beta+\varepsilon} [d_{V_0^{m^n}}(0, m^n - 1)]^2 \leq 20000\tilde{f}(\beta - c_\beta, m)m^4 \Lambda(m^n, \beta + \varepsilon)^2 \end{aligned}$$

for all $\varepsilon \in (-c_\beta, c_\beta)$. This shows (68) along the subsequence m, m^2, m^3, \dots . To extend inequality (68) from this subsequence to all integers, use Lemma 4.1.

Now let us consider the case where $\beta < 1$. From the definition of the function $f(\cdot, \cdot)$ it is clear that $f(\beta, m) \leq m - 2$ for all $\beta \in \mathbb{R}_{\geq 0}$ and thus we also have $\tilde{f}(\beta, m) = 27(2 + f(\beta, m)) \leq 27m$. Using Lemma 7.1, we know that there is a constant $c > 0$ such that

$$\mathbb{E}_\beta [d_{V_0^{m^n}}(0, m^n - 1)] \geq cm^{(n-k)(1-\beta)} \mathbb{E}_\beta [d_{V_0^{m^{n-k}}}(0, m^{n-k} - 1)]$$

$$\geq (cm^{1-\beta})^{n-k} \mathbb{E}_\beta \left[d_{V_0^{m^{n-k}}} (0, m^{n-k} - 1) \right]$$

for all $n \geq k$ and $m \in \mathbb{N}$. Now take m large enough and c_β small enough so that $\frac{1200 m^{\beta+\varepsilon-1}}{c(1-\beta-\varepsilon)} < 0.5$ for all $\varepsilon \in (-c_\beta, c_\beta)$, and that $0 < \beta - c_\beta < \beta + c_\beta < 1$. Using (67) we get that for such m and $\varepsilon \in (-c_\beta, c_\beta)$

$$\begin{aligned} \mathbb{E}_{\beta+\varepsilon} \left[d_{V_0^{m^{n+1}}} (0, m^{n+1} - 1)^2 \right] &\leq 5000m^4 \sum_{k=1}^n \left(2\tilde{f}(\beta, m) \right)^{n+1-k} \mathbb{E}_\beta \left[d_{V_0^{m^k}} (0, m^k - 1) \right]^2 \\ &\leq 5000m^4 \sum_{k=1}^n \left(2\tilde{f}(\beta + \varepsilon, m) \right)^{n+1-k} (cm^{1-\beta-\varepsilon})^{2(k-n)} \mathbb{E}_{\beta+\varepsilon} \left[d_{V_0^{m^n}} (0, m^n - 1) \right]^2 \\ &\leq 10000\tilde{f}(\beta - c_\beta, m)m^4 \sum_{k=1}^n \left(\frac{1200 m^{\beta+\varepsilon-1}}{c(1-\beta-\varepsilon)} \right)^{n-k} \mathbb{E}_{\beta+\varepsilon} \left[d_{V_0^{m^n}} (0, m^n - 1) \right]^2 \\ &\leq 10000\tilde{f}(\beta - c_\beta, m)m^4 \sum_{k=1}^n 0.5^{n-k} \mathbb{E}_{\beta+\varepsilon} \left[d_{V_0^{m^n}} (0, m^n - 1) \right]^2 \\ &\leq 20000\tilde{f}(\beta - c_\beta, m)m^4 \mathbb{E}_{\beta+\varepsilon} \left[d_{V_0^{m^n}} (0, m^n - 1) \right]^2 \leq 10^6 m^5 \mathbb{E}_{\beta+\varepsilon} \left[d_{V_0^{m^n}} (0, m^n - 1) \right]^2. \end{aligned}$$

which shows (68) for numbers of the form m, m^2, m^3, \dots . To extend inequality (68) from this subsequence to all integers, use Lemma 4.1. \square

8 References

- [1] Louigi Addario-Berry, Nicolas Broutin, and Christina Goldschmidt. The continuum limit of critical random graphs. *Probability Theory and Related Fields*, 152(3-4):367–406, 2012.
- [2] Michael Aizenman and Charles M. Newman. Discontinuity of the percolation density in one dimensional $1/|x - y|^2$ percolation models. *Communications in Mathematical Physics*, 107(4):611–647, 1986.
- [3] Peter Antal and Agoston Pisztora. On the chemical distance for supercritical bernoulli percolation. *The Annals of Probability*, pages 1036–1048, 1996.
- [4] Johannes Bäumler. Behavior of the distance exponent for $1/|x - y|^{2d}$ long-range percolation. *Preprint*, 2022.
- [5] Johannes Bäumler and Noam Berger. Isoperimetric lower bounds for critical exponents for long-range percolation. *arXiv preprint arXiv:2204.12410*, 2022.
- [6] David J. Barsky and Michael Aizenman. Percolation critical exponents under the triangle condition. *The Annals of Probability*, pages 1520–1536, 1991.
- [7] Itai Benjamini and Noam Berger. The diameter of long-range percolation clusters on finite cycles. *Random Structures & Algorithms*, 19(2):102–111, 2001.
- [8] Itai Benjamini, Harry Kesten, Yuval Peres, and Oded Schramm. Geometry of the uniform spanning forest: transitions in dimensions 4, 8, 12, \dots . In *Selected Works of Oded Schramm*, pages 751–777. Springer, 2011.
- [9] Jacob Van Den Berg and Harry Kesten. Inequalities with applications to percolation and reliability. *Journal of applied probability*, 22(3):556–569, 1985.

- [10] Noam Berger. Transience, recurrence and critical behavior for long-range percolation. *Communications in mathematical physics*, 226(3):531–558, 2002.
- [11] Noam Berger. A lower bound for the chemical distance in sparse long-range percolation models. *arXiv preprint math/0409021*, 2004.
- [12] Marek Biskup. On the scaling of the chemical distance in long-range percolation models. *The Annals of Probability*, 32(4):2938–2977, 2004.
- [13] Marek Biskup. Graph diameter in long-range percolation. *Random Structures & Algorithms*, 39(2):210–227, 2011.
- [14] Marek Biskup and Jeffrey Lin. Sharp asymptotic for the chemical distance in long-range percolation. *Random Structures & Algorithms*, 55(3):560–583, 2019.
- [15] Christian Borgs, Jennifer T. Chayes, Gordon Slade, Joel Spencer, and Remco van der Hofstad. Random subgraphs of finite graphs. ii. the lace expansion and the triangle condition. *The Annals of Probability*, 33(5):1886–1944, 2005.
- [16] Don Coppersmith, David Gamarnik, and Maxim Sviridenko. The diameter of a long-range percolation graph. In *Mathematics and computer science II*, pages 147–159. Springer, 2002.
- [17] Michael Damron, Jack Hanson, and Philippe Sosoe. On the chemical distance in critical percolation. *Electronic Journal of Probability*, 22:1–43, 2017.
- [18] Barbara Dembin. The variance of the graph distance in the infinite cluster of percolation is sublinear. *arXiv preprint arXiv:2203.01083*, 2022.
- [19] Jian Ding and Li Li. Chemical distances for percolation of planar gaussian free fields and critical random walk loop soups. *Communications in Mathematical Physics*, 360(2):523–553, 2018.
- [20] Jian Ding and Allan Sly. Distances in critical long range percolation. *arXiv preprint arXiv:1303.3995*, 2013.
- [21] Alexander Drewitz, Balázs Ráth, and Artëm Sapozhnikov. On chemical distances and shape theorems in percolation models with long-range correlations. *Journal of Mathematical Physics*, 55(8):083307, 2014.
- [22] Hugo Duminil-Copin, Christophe Garban, and Vincent Tassion. Long-range models in 1d revisited. *arXiv preprint arXiv:2011.04642*, 2020.
- [23] Cees M. Fortuin, Pieter W. Kasteleyn, and Jean Ginibre. Correlation inequalities on some partially ordered sets. *Communications in Mathematical Physics*, 22(2):89–103, 1971.
- [24] Olivier Garet and Régine Marchand. Large deviations for the chemical distance in supercritical bernoulli percolation. *The Annals of Probability*, 35(3):833–866, 2007.
- [25] Nannan Hao and Markus Heydenreich. Graph distances in scale-free percolation: the logarithmic case. *arXiv preprint arXiv:2105.05709*, 2021.
- [26] Theodore E. Harris. A lower bound for the critical probability in a certain percolation process. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 56, pages 13–20. Cambridge University Press, 1960.
- [27] Sarai Hernandez-Torres, Eviatar B Procaccia, and Ron Rosenthal. The chemical distance in random interacements in the low-intensity regime. *arXiv preprint arXiv:2112.13390*, 2021.
- [28] Markus Heydenreich and Remco Van der Hofstad. *Progress in high-dimensional percolation and random graphs*. Springer, 2017.

- [29] Tom Hutchcroft. Power-law bounds for critical long-range percolation below the upper-critical dimension. *Probability Theory and Related Fields*, pages 1–38, 2021.
- [30] Tom Hutchcroft. The critical two-point function for long-range percolation on the hierarchical lattice. *arXiv preprint arXiv:2103.17013*, 2021.
- [31] Tom Hutchcroft. Sharp hierarchical upper bounds on the critical two-point function for long-range percolation on \mathbb{Z}^d . *arXiv preprint arXiv:2202.07634*, 2022.
- [32] Asaf Nachmias and Yuval Peres. Critical random graphs: diameter and mixing time. *The Annals of Probability*, 36(4):1267–1286, 2008.
- [33] Charles M. Newman and Lawrence S. Schulman. One dimensional $1/|j-i|^s$ percolation models: The existence of a transition for $s \leq 2$. *Communications in Mathematical Physics*, 104(4):547–571, 1986.
- [34] David Reimer. Proof of the van den berg–kesten conjecture. *Combinatorics, Probability and Computing*, 9(1):27–32, 2000.
- [35] Lawrence S. Schulman. Long range percolation in one dimension. *Journal of Physics A: Mathematical and General*, 16(17):L639, 1983.