

# The genus distribution of cubic graphs and asymptotic number of rooted cubic maps with high genus

Zhicheng Gao\*

School of Mathematics and Statistics  
Carleton University  
Ottawa, Ontario K1S5B6, Canada  
zgao@math.carleton.ca

August 16, 2022

## Abstract

Let  $C_{n,g}$  be the number of rooted cubic maps with  $2n$  vertices on the orientable surface of genus  $g$ . We show that the sequence  $(C_{n,g} : g \geq 0)$  is asymptotically normal with mean and variance asymptotic to  $(1/2)(n - \ln n)$  and  $(1/4) \ln n$ , respectively. We derive an asymptotic expression for  $C_{n,g}$  when  $(n - 2g)/\ln n$  lies in any closed subinterval of  $(0, 1)$ . Using rotation systems and Bender's theorem about generating functions with fast-growing coefficients, we derive simple asymptotic expressions for the numbers of rooted regular maps, disregarding the genus. In particular, we show that the number of rooted cubic maps with  $2n$  vertices, disregarding the genus, is asymptotic to  $\frac{3}{\pi} n!6^n$ .

**Keywords:** graph, map, genus, normal distribution, asymptotics.  
Mathematics Subject Classification: 05A16, 05C89

---

\*Research supported by Carleton University Research Development Fund

# 1 Introduction

This paper is motivated by two lines of research. The first is about genus distributions of graphs, which is an active research area in topological graph theory. The second is about asymptotic properties of rooted maps of high genus, which has attracted much attention recently. There is a close connection between genus distribution of graphs and enumeration of rooted maps, which we shall briefly describe below. Let  $\Sigma_g$  denote the orientable surface of genus  $g$ . A *map* on  $\Sigma_g$  is a connected graph  $G$  that is *embedded* on  $\Sigma_g$  in such a way that each component of  $\Sigma_g - G$ , called a *face*, is a simply connected region (i.e., a topological disk). Such an embedding is known as *2-cell* (or *cellular*) embedding. Throughout the paper, all maps and graph embeddings are cellular embeddings on orientable surfaces. A map on  $\Sigma_g$  will be called a map with genus  $g$ . A map is called *cubic* if all its vertices have degree 3. The dual of a cubic map is known as a *triangular map* (or *triangulation*), whose faces all have degree 3. Throughout the paper, a map is always *rooted*, meaning that a vertex and an edge incident to it are distinguished. The notion of rooted map was first introduced by Tutte in his seminal paper [26] on enumeration of planar maps. We emphasize that there is no labeling (neither vertex nor edge) in rooted maps. Rooting a map trivializes the automorphism group which makes enumeration easier.

On the other hand, in topological graph theory [6, 7], one trivializes the automorphism group of an embedded graph by labeling and orienting the edges. A 2-cell embedding can be described combinatorially in terms of a permutation  $\phi$  of the ends of the edges, known as *rotation system*. The cycles of  $\phi$  describes the cyclic ordering (say, clockwise) of the (ends of) edges at each vertex. For each rooted map with  $n$  edges, there are  $(n-1)!2^{n-1}$  ways to label and to orient the  $n-1$  non-root edges of the underlying graph. This gives the following connection between rooted maps and rotation systems [19, Lemma 2.3].

**Proposition 1** *Each rooted map with  $n$  edges corresponds to exactly  $2^{n-1}(n-1)!$  rotation systems.*

For a given family  $\mathcal{F}$  of edge-labeled directed graphs, let  $F_{n,g}$  denote the number of embeddings of the graphs in  $\mathcal{F}$  of  $n$  edges and genus  $g$ . The sequence  $(F_{n,g}, g \geq 0)$  is called the *genus distribution* of  $\mathcal{F}$ . By Proposition 1,  $F_{n,g}2^{1-n}/(n-1)!$  is the number of rooted maps whose underlying graphs are in  $\mathcal{F}$ . Genus distributions of graphs were first introduced by Gross and

Furst [6]. Jackson [19] derived a simple recursion for the number of rooted *bouquets* (maps with only one vertex, whose duals are known as *unicellular maps*) using rotation system and group characters. Using Jackson's result (also known as Harer-Zagier's formula [17]), Gross et al. [15] showed that the genus distribution of bouquets is strongly unimodal, and Stahl [25] derived an asymptotic relation between the number of rooted bouquets and the Stirling cycle numbers. It follows from Stahl's result (and the well-known fact that the distribution of Stirling cycle numbers is asymptotically normal) that the genus distribution of bouquets is asymptotically normal. An exact expression for the genus distribution of bouquets was given in [20].

Algorithms for computing the genus distributions of some families of cubic graphs were studied in [13, 14, 16]. It was shown in [27] that the genus distribution of a general graph can be computed using the genus distributions of some cubic graphs.

Much work has been done on the enumeration of rooted maps of constant genus since Tutte's pioneering work on planar maps [26]. Many families of non-planar maps have also been enumerated; see, e.g., [3, 4, 9]. In contrast, enumerative results on maps with high genus (both the genus and the number of edges go to infinity) are rare. It is clear that studying the genus distribution involves enumerating maps of high genus. Properties of large unicellular maps of high genus were studied in [1, 23, 24].

Let  $C_{n,g}$  be the number of rooted cubic maps with  $2n$  vertices and genus  $g$ , and  $J_{n,f}$  be the number of rooted cubic maps with  $2n$  vertices and  $f$  faces. It follows from Euler's formula that

$$f = n + 2 - 2g, \tag{1}$$

$$C_{n,g} = J_{n,n+2-2g}. \tag{2}$$

By Proposition 1,  $(C_{n,g}2^{3n-1}(3n-1)! : g \geq 0)$  is the the genus distribution of cubic graphs. The sequence  $(J_{n,f}2^{3n-1}(3n-1)! : f \geq 1)$  is known as the *region distribution* of cubic graphs.

More recently properties of large triangulations (duals of cubic maps) of high genus were studied in [5], where an asymptotic formula, which is accurate up to a sub-exponential factor, was obtained. An asymptotic formula for such triangulations was reported in [10]; however, there is a gap in the proof. In this paper we will derive an asymptotic formula for the number of rooted cubic maps with high genus, which is accurate up to a constant factor. As in [10], we use the Goulden-Jackson recursion for the number of rooted

cubic maps; however, we shall focus on the genus polynomial  $\sum_g C_{n,g} x^g$  in the current paper instead of the generating function  $\sum_n C_{n,g} x^n$  in [10].

The rest of the paper is organized as follows. In Section 2, we use rotation systems and Bender's theorem about generating functions with fast growing coefficients to derive asymptotic expressions for the numbers of all maps and regular maps, disregarding the genus. In particular, a simple asymptotic formula is obtained for the number of rooted cubic maps, disregarding the genus. Our main results about the genus distribution of cubic graphs and asymptotic number of rooted cubic maps of high genus are stated at the end of this section. In section 3, we analyze the asymptotic behavior of the genus polynomials of cubic graphs using the Goulden-Jackson recursion. This is the most technical part of the paper due to the complexity of the nonlinear recursion. The proofs of our main results are then completed by using the limit theorems from [11]. Section 4 concludes our paper.

## 2 The results

Since there are  $(2n - 1)!$  permutations of  $2n$  elements which have exactly one cycle, by Proposition 1 the total number of rooted bouquets with  $n$  edges is equal to

$$\frac{(2n - 1)!}{(n - 1)!2^{n-1}}.$$

Proposition 1 can also be used to find the asymptotic number of rooted maps, disregarding the genus, in some other families. To the best of our knowledge, such asymptotic results have not appeared in the literature. We will use the following result about the generating functions with fast growing coefficients, which is an immediate consequence of [2, Corollary 4]. We say that a sequence  $(a_n)$  of positive numbers *grows super-exponentially* if  $a_n/a_{n-1} \rightarrow \infty$  as  $n \rightarrow \infty$ .

**Proposition 2 (Bender)** *Suppose  $(a_n)$  grows super-exponentially. Then*

$$[z^n] \ln \left( 1 + \sum_{n \geq 1} a_n z^n \right) \sim a_n.$$

**Proposition 3** (a) *The total number of rooted maps with  $n$  edges is asymptotic to*

$$\frac{(2n)!}{(n - 1)!} 2^{1-n}.$$

(b) For each fixed integer  $d \geq 2$  and as  $m \rightarrow \infty$ , the number of rooted  $2d$ -regular maps with  $m$  vertices is asymptotic to

$$\frac{(2md)!}{m!(md-1)!} (2d)^{-m} 2^{1-md}. \quad (3)$$

(c) For each fixed odd integer  $r \geq 3$  and as  $k \rightarrow \infty$ , the number of rooted  $r$ -regular maps with  $2k$  vertices is asymptotic to

$$\frac{(2kr)!}{(2k)!(kr-1)!} r^{-2k} 2^{1-kr}. \quad (4)$$

**Proof** For part (a), we note [21] that the exponential generating function of (transitive) rotation systems is given by

$$\ln \left( \sum_{k \geq 0} \frac{(2k)!}{k!} z^k \right).$$

It follows from Proposition 1 that the total number of rooted  $n$ -edged maps is equal to

$$\frac{1}{(n-1)!} 2^{1-n} n! [z^n] \ln \left( \sum_{k \geq 0} \frac{(2k)!}{k!} z^k \right).$$

Since  $\frac{(2k)!}{2^k k!}$  grows super-exponentially, by Proposition 2, we have

$$[z^n] \ln \left( \sum_{k \geq 0} \frac{(2k)!}{k!} z^k \right) \sim \frac{(2n)!}{n!}.$$

Hence the total number of rooted  $n$ -edged maps is asymptotic to

$$\frac{1}{(n-1)!} 2^{1-n} n! \frac{(2n)!}{n!},$$

which completes the proof of part (a).

For part (b), we first note that each  $2d$ -regular graph with  $m$  vertices has  $md$  edges, and there are

$$\frac{(2md)!}{(2d)^m m!}$$

permutations of  $2md$  elements with exactly  $m$  cycles of length  $2d$ . Hence the corresponding exponential generating function of (transitive) rotation systems is given by

$$\ln \left( \sum_{m \geq 0} \frac{(2md)!}{(2d)^m m! (md)!} z^{md} \right).$$

For each fixed  $d \geq 2$ ,  $\frac{(2md)!}{(2d)^m m! (md)!}$  grows super-exponentially. Applying Proposition 2 again (in terms of the variable  $w := z^d$ ), we obtain, for each fixed  $d$  and as  $m \rightarrow \infty$ ,

$$[z^{md}] \ln \left( \sum_{m \geq 0} \frac{(2md)!}{(2d)^m m! (md)!} z^{md} \right) \sim \frac{(2md)!}{(2d)^m m! (md)!}.$$

It follows from Proposition 1 that the number of rooted  $2d$ -regular maps with  $m$  vertices is asymptotic to

$$\frac{(2md)!}{m!(md-1)!} (2d)^{-m} 2^{1-md},$$

which gives (3).

For part (c), since  $r \geq 3$  is odd, each  $r$ -regular graph must have even number of vertices, and each  $r$ -regular graph with  $2k$  vertices has exactly  $kr$  edges. Since there are

$$\frac{(kr)!}{r^{2k} (2k)!}$$

permutations of  $2kr$  elements with exactly  $2k$  cycles of length  $r$ , the corresponding exponential generating function of (transitive) rotation systems is given by

$$\ln \left( \sum_{k \geq 0} \frac{(2kr)!}{r^{2k} (2k)! (kr)!} z^{kr} \right).$$

Since  $\frac{(2kr)!}{r^{2k} (2k)! (kr)!}$  grows super-exponentially, it follows from Proposition 2 (in terms of the variable  $w := z^r$ ) that the number of rooted  $r$ -regular maps with  $2k$  vertices is asymptotic to

$$\frac{(2kr)!}{r^{2k} (2k)! (kr-1)!} 2^{1-kr},$$

which gives (4). ■

Let  $C_n = \sum_{g \geq 0} C_{n,g}$  be the total number of rooted cubic maps with  $2n$  vertices. The next result follows immediately from Proposition 3(c) and Stirling's formula.

**Corollary 1** *The total number of rooted cubic maps with  $2n$  vertices is*

$$C_n \sim \frac{3}{\pi} n! 6^n \sim \frac{6}{\sqrt{2\pi}} \left(\frac{6}{e}\right)^n n^{n+\frac{1}{2}}.$$

Our main results are summarized in the following two theorems.

**Theorem 1** (a) *The genus distribution of cubic graphs is asymptotically normal with mean and variance, respectively, asymptotic to  $\frac{n-\ln n}{2}$  and  $\frac{\ln n}{4}$ . That is,*

$$\sum_{g \leq \frac{n-\ln n}{2} + \frac{t\sqrt{\ln n}}{2}} \frac{C_{n,g}}{C_n} \sim \frac{1}{\sqrt{2\pi}} \int_{-\infty}^t \exp\left(-\frac{x^2}{2}\right) dx.$$

(b) *The region distribution of cubic graphs is asymptotically normal with mean and variance both asymptotic to  $\ln n$ .*

**Theorem 2** *Let  $\varepsilon$  be any small positive constant. There is a function  $K(y)$  which is analytic in  $[\varepsilon, 1 - \varepsilon]$  such that*

$$C_{n,g} \sim \frac{\sqrt{2}}{3} K\left(\frac{n-2g}{\ln n}\right) \left(\frac{\ln n}{n-2g}\right)^2 6^n \frac{(n-1)!}{(n-2g)!} (\ln n)^{n-2g}, \quad (5)$$

*uniformly for all  $g, n \rightarrow \infty$  satisfying*

$$\frac{n-2g}{\ln n} \in [\varepsilon, 1 - \varepsilon].$$

### 3 Proofs of Theorems 1 and 2

Define

$$H_{n,g} = (3n+2)C_{n,g} \quad \text{for } n \geq 1, \quad (6)$$

$$H_{-1,g} = \llbracket g = 0 \rrbracket / 2, \quad H_{0,g} = 2\llbracket g = 0 \rrbracket, \quad H_{n,-1} = 0. \quad (7)$$

Goulden and Jackson [12] derived the following recursion for  $(n, g) \neq (-1, 0)$ :

$$H_{n,g} = \frac{4n(3n+2)(3n-2)}{n+1}H_{n-2,g-1} + \frac{4(3n+2)}{n+1} \sum_{k=-1}^{n-1} \sum_{h=0}^g H_{k,h}H_{n-2-k,g-h}. \quad (8)$$

Define

$$H_n(x) = \frac{3n+2}{n!}6^{-n} \sum_{g \geq 0} C_{n,g}x^g = \frac{1}{n!}6^{-n} \sum_{g \geq 0} H_{n,g}x^g, \\ J_n(x) = \frac{3n+2}{n!}6^{-n} \sum_{f \geq 1} J_{n,f}x^f.$$

Using (2), we obtain

$$J_n(x) = \frac{3n+2}{n!}6^{-n} \sum_{g \geq 0} C_{n,g}x^{n+2-2g} \\ = H_n(1/x^2)x^{n+2}. \quad (9)$$

It follows from (7) and (8) that

$$H_0(x) = 2, \quad (10)$$

$$H_1(x) = \frac{20+5x}{6}, \quad (11)$$

$$H_2(x) = \frac{32+28x}{9}, \quad (12)$$

$$H_n(x) = \frac{2(3n+2)}{3n(n+1)}H_{n-1}(x) + \frac{9n^2-4}{9(n^2-1)}xH_{n-2}(x) \\ + \frac{3n+2}{9n(n^2-1)} \sum_{i=0}^{n-2} \frac{1}{\binom{n-2}{i}} H_i(x)H_{n-2-i}(x). \quad (n \geq 2) \quad (13)$$

Recursion (13) suggests that  $H_n(x)$  is of the order  $n^{1/\sqrt{x}}x^{n/2}$  (see Theorem 3 below). Hence it is convenient to set

$$y := \frac{1}{\sqrt{x}}$$



and consider

$$h_n(y) := H_n(1/y^2)n^{-y}y^{n+2}. \quad (n \geq 1) \quad (14)$$

It follows from (9) that

$$h_n(y) = J_n(y)n^{-y}. \quad (15)$$

Using (10)–(14) (with the help of the Computer Algebra System *Maple*), we obtain

$$h_1(y) = \frac{y(20y^2 + 5)}{6}, \quad (16)$$

$$h_2(y) = \frac{4y^2(8y^2 + 7)}{9}2^{-y}, \quad (17)$$

$$h_3(y) = \frac{11y(336y^4 + 664y^2 + 105)}{1296}3^{-y}, \quad (18)$$

$$h_4(y) = y^2 \left( \frac{448}{243}y^4 + \frac{1631}{243}y^2 + \frac{1183}{324} \right) 4^{-y}, \quad (19)$$

$$h_5(y) = \frac{17y(27456y^6 + 163248y^4 + 198396y^2 + 25025)}{466560}5^{-y}, \quad (20)$$

$$h_6(y) = \frac{y^2(3072y^6 + 27532y^4 + 61185y^2 + 26261)}{6561}6^{-y}. \quad (21)$$

In terms of  $h_n(y)$  and for  $n \geq 7$ , we can rewrite (13) as (separating the six terms corresponding to  $k \in \{0, 1, 2, n-2, n-3, n-4\}$  in the last summation)

$$\begin{aligned} h_n(y) &= \left( \frac{2y}{n} - \frac{2y}{3n(n+1)} \right) \left( \frac{n-1}{n} \right)^y h_{n-1}(y) \\ &+ \left( 1 + \frac{5}{9(n^2-1)} + \frac{4(3n+2)y^2}{9n(n^2-1)} \right) \left( \frac{n-2}{n} \right)^y h_{n-2}(y) \\ &+ \frac{2(3n+2)}{9n(n^2-1)(n-2)} \left( \frac{n-3}{n} \right)^y h_1(y)h_{n-3}(y) \\ &+ \frac{2(3n+2)}{9n(n^2-1)(n-2)} \left( \frac{n-4}{n} \right)^y h_2(y)h_{n-4}(y) \\ &+ \frac{(3n+2)}{9n(n^2-1)} \sum_{k=3}^{n-5} \frac{1}{\binom{n-2}{k}} \left( \frac{k(n-2-k)}{n} \right)^y h_k(y)h_{n-2-k}(y). \end{aligned} \quad (22)$$

We note that the RHS of (22) is of the form

$$\frac{2y}{n}h_{n-1}(y) + \left(\frac{n-2}{n}\right)^y h_{n-2}(y) + O\left(\frac{1}{n^2}\right),$$

provided that  $h_n(y)$  is bounded.

We first show that  $h_n(y)$  is bounded when  $0 \leq y \leq 1$ .

**Lemma 1** *For  $n \geq 2$ , we have*

$$h_n(1) \leq \frac{10}{3}, \quad (23)$$

$$h_n(y) \leq \exp(5 - 5/n). \quad (0 \leq y < 1) \quad (24)$$

**Proof** Using (16)–(21), we obtain

$$h_1(1) = \frac{25}{6}, h_2(1) = \frac{10}{3} > h_3(1) > h_4(1) > h_5(1) > h_6(1) < 3. \quad (25)$$

Now we assume  $n \geq 7$  and move on to the inductive step. Dividing both sides of (22) by  $10/3$ , using  $3n + 2 < 3(n + 1)$  and induction hypothesis, we obtain

$$\begin{aligned} \frac{h_n(1)}{10/3} &\leq \frac{2(n-1)}{n^2} + \left( \frac{n-2}{n} + \frac{5}{9(n^2-1)n} + \frac{4(n-2)}{3n^2(n-1)} \right) \\ &\quad + \frac{25}{9n^2(n-1)} + \frac{80}{9n^2(n-1)(n-2)} \\ &\quad + \frac{10}{9n^2(n-1)} \sum_{k=3}^{n-5} \frac{k(n-2-k)}{\binom{n-2}{k}} \end{aligned} \quad (26)$$

$$= 1 - \frac{6n^2 - 34n - 35}{9n^2(n^2-1)} + \frac{10}{9n^2(n-1)} \sum_{k=3}^{n-5} \frac{k(n-2-k)}{\binom{n-2}{k}}. \quad (27)$$

Noting that  $\frac{k(n-2-k)}{\binom{n-2}{k}}$  is a decreasing function of  $k$  when  $k \leq \lfloor n/2 \rfloor - 1$ , and using Maple, it is easy to verify

$$\sum_{k=3}^{n-5} \frac{k(n-2-k)}{\binom{n-2}{k}} \leq \frac{2}{25}. \quad (28)$$

It follows from (27) that, for  $n \geq 7$ ,

$$\begin{aligned} \frac{h_n(1)}{10/3} &\leq 1 - \frac{6n^2 - 34n - 35}{9n^2(n^2 - 1)} + \frac{4}{45n^2(n - 1)} \\ &\leq 1 - \frac{30n^2 - 174n - 179}{45n^2(n^2 - 1)} \\ &< 1. \end{aligned}$$

This completes the proof of (23).

Next we consider the case  $0 \leq y < 1$ . Noting that  $J_n(y)$  is a polynomial of  $y$  with non-negative coefficients, we obtain from (15) and (23) that

$$h_n(y) \leq J_n(1)n^{-s} = h_n(1)n^{1-s} \leq \frac{10}{3}n^{1-s}. \quad (n \geq 2, 0 \leq y \leq 1) \quad (29)$$

It is easy to check  $10n/3 < \exp(5 - 5/n)$  for all  $2 \leq n \leq 35$ . It follows from (29) that (24) holds for  $2 \leq n \leq 35$ . Hence we may assume  $n \geq 36$  and move on to the inductive step. Dividing both sides of (22) by  $\exp(5 - 5/n)$  and applying the upper bound from (29) as well as the inequality  $3n + 2 < 3n + 3$  in the last two lines in (22), we obtain

$$\begin{aligned} \frac{h_n(y)}{\exp(5 - 5/n)} &\leq \frac{2y}{n} + \left(\frac{n-2}{n}\right)^y \exp\left(\frac{-10}{n^2}\right) \\ &\quad + \frac{5}{9(n^2 - 1)} + \frac{4y^2}{3n(n-1)} \\ &\quad + \frac{2}{3n(n-1)} \left( \frac{h_1(y)}{n-2} + \frac{2h_2(y)}{(n-2)(n-3)} \right) \\ &\quad + \frac{10}{9n(n-1)} \sum_{k=3}^{n-5} \frac{k}{\binom{n-2}{k}}. \end{aligned} \quad (30)$$

Using

$$(1-t)^y \leq 1 - yt, \quad (0 \leq y, t \leq 1) \quad (31)$$

$$e^{-t} \leq 1 - \frac{4t}{5}, \quad (0 \leq t \leq 0.45) \quad (32)$$

$$h_1(y) + h_2(y) \leq \frac{10}{3}(1+2) = 10,$$

$$\sum_{k=3}^{n-5} \frac{k}{\binom{n-2}{k}} \leq \frac{1}{100}, \quad (n \geq 36)$$

we obtain from (30) that (noting  $n \geq 36$ )

$$\begin{aligned} \frac{h_n(y)}{\exp(5 - 5/n)} &\leq \frac{2y}{n} + \left(1 - \frac{2y}{n}\right) \left(1 - \frac{8}{n^2}\right) \\ &\quad + \frac{5}{9(n^2 - 1)} + \frac{4}{3n(n - 1)} \\ &\quad + \frac{1}{n(n - 1)} + \frac{1}{90n(n - 1)} \\ &\leq 1 - \frac{1}{n^2} < 1. \end{aligned}$$

■

The range of  $y$  in Lemma 1 can be extended, and we illustrate this in the next lemma. We remark that the upper end  $y = 2$  below is chosen for simplicity, and it can be extended further. In fact we believe that  $h_n(y)$  is bounded by a function of  $y$  for all  $y \geq 0$ , but we are unable to prove this at this stage.

**Lemma 2** *For  $n \geq 2$  and  $1 < y \leq 2$ , we have*

$$h_n(y) \leq 9n, \tag{33}$$

$$h_n(y) \leq \exp(10 - 10/n). \tag{34}$$

**Proof** Using (16)–(21) and (22), it is easy to check that both (33) and (34) hold for  $2 \leq n \leq 19$ . So we assume  $n \geq 20$  and move on to the inductive step. Using (22), (32), and

$$(1 - t)^y \leq e^{-yt} \leq 1 - yt + \frac{y^2 t^2}{2}, \quad (0 < t < 1, y \geq 0) \tag{35}$$

we obtain

$$\begin{aligned} \frac{h_n(y)}{9n} &\leq \frac{2y}{n} + \left(1 - \frac{2(y+1)}{n} + \frac{2(y+1)^2}{n^2}\right) \\ &\quad + \frac{5}{9(n^2 - 1)} + \frac{4y^2}{3n(n - 1)} \\ &\quad + \frac{2}{3n(n - 1)} \left(\frac{h_1(y)}{n - 2} + \frac{2h_2(y)}{(n - 2)(n - 3)}\right) \\ &\quad + \frac{3}{n(n - 1)} \sum_{k=3}^{n-5} \frac{1}{\binom{n-2}{k}} \left(\frac{k(n - 2 - k)}{n}\right)^{y+1}. \end{aligned} \tag{36}$$

With the help of Maple, it can be checked that, for  $n \geq 20$  and  $1 < y \leq 2$ ,

$$\sum_{k=3}^{n-5} \frac{1}{\binom{n-2}{k}} \left( \frac{k(n-2-k)}{n} \right)^{y+1} < \frac{1}{4}. \quad (37)$$

Substituting this into (36) and using  $n \geq 20, 1 < y \leq 2$ , we obtain

$$\frac{h_n(y)}{9n} \leq 1 - \frac{2}{n} \left( 1 - \frac{13}{n} \right) < 1.$$

This completes the proof of (33).

The proof of (34) is similar. Using (35), (33), (32) and (22), we obtain

$$\begin{aligned} \frac{h_n(y)}{\exp(10 - 10/n)} &\leq \frac{2y}{n} + \left( 1 - \frac{2y}{n} + \frac{2y^2}{n^2} \right) \left( 1 - \frac{16}{n^2} \right) \\ &\quad + \frac{5}{9(n^2 - 1)} + \frac{4y^2}{3n(n-1)} \\ &\quad + \frac{2}{3n(n-1)} \left( \frac{h_1(y)}{n-2} + \frac{2h_2(y)}{(n-2)(n-3)} \right) \\ &\quad + \frac{3}{n(n-1)} \sum_{k=3}^{n-5} \frac{k}{\binom{n-2}{k}} \left( \frac{k(n-2-k)}{n} \right)^y. \end{aligned}$$

Using  $1 < y \leq 2, n \geq 20, h_1(y) \leq 9, h_2(y) \leq 18$  and (37), we obtain

$$\frac{h_n(y)}{\exp(10 - 10/n)} \leq 1 - \frac{1}{n^2} < 1.$$

This completes the proof of (34). ■

The next result gives asymptotics for  $h_n(y)$  which will be used to prove our main results (Theorems 2 and 3). In what follows,  $\Re(z)$  and  $\Im(z)$  stand for the real and imaginary parts of a complex number  $z$ , respectively,  $\mathcal{S}$  shall denote the set  $\{y \in \mathbb{C} : |y| \leq 2, \Re(y) \geq 0, |y| - \Re(y) < 1\}$ , and all the big- $O$  terms are independent of  $y$ . We shall also use the Iverson bracket  $\llbracket P \rrbracket$ , which is equal to 1 if the predicate  $P$  is true and 0 otherwise.

**Theorem 3** *There is a function  $K(y)$  which is analytic in  $\mathcal{S}$  such that*

$$h_n(y) = K(y) + o(1) + \llbracket \Re(y) = 0 \rrbracket O(1). \quad (y \in \mathcal{S}) \quad (38)$$

**Proof** We have

$$|h_n(y)| = |J_n(y)| n^{-\Re(y)} \leq J_n(|y|^2) n^{-\Re(y)} = h_n(|y|) n^{|y|-\Re(y)}. \quad (39)$$

Using Lemmas 1 and 2, we have

$$|h_n(y)| \leq e^{10} n^{|y|-\Re(y)}. \quad (|y| \leq 2) \quad (40)$$

It follows from (22) that

$$h_n(y) - h_{n-2}(y) = \frac{2y}{n} (h_{n-1}(y) - h_{n-2}(y)) + O\left(n^{-(2+\Re(y)-|y|)}\right), \quad (41)$$

or

$$h_n(y) - h_{n-1}(y) = \left(\frac{2y}{n} - 1\right) (h_{n-1}(y) - h_{n-2}(y)) + O\left(n^{-(2+\Re(y)-|y|)}\right). \quad (42)$$

Define

$$d_n(y) := |h_n(y) - h_{n-1}(y)|.$$

Using (42) and

$$\begin{aligned} \left|\frac{2y}{n} - 1\right| &= \left(\left(\frac{2\Re(y)}{n} - 1\right)^2 + \left(\frac{2\Im(y)}{n}\right)^2\right)^{1/2} \\ &= 1 - \frac{2\Re(y)}{n} + O\left(\frac{1}{n^2}\right), \end{aligned}$$

we obtain

$$d_n(y) \leq \left(1 - \frac{2\Re(y)}{n}\right) d_{n-1}(y) + O\left(n^{-(2+\Re(y)-|y|)}\right).$$

It is known (use “rsolve” in Maple or [22, Section 2.2]) that the solutions to the recursion

$$d_n = \left(1 - \frac{a}{n}\right) d_{n-1} + n^{-b} \quad (a \geq 0, n \geq 1)$$

satisfy

$$d_n = O\left(n^{-a} (1 + \llbracket b - a = 1 \rrbracket \ln n) + n^{-(b-1)}\right).$$

Hence

$$d_n(y) = O\left(n^{-2\Re(y)} (1 + \llbracket \Re(y) + |y| = 1 \rrbracket \ln n) + n^{-(1+\Re(y)-|y|)}\right).$$

That is,

$$h_n(y) - h_{n-1}(y) = o(1) + \llbracket \Re(y) = 0 \rrbracket O(1) \quad (y \in \mathcal{S}). \quad (43)$$

On the other hand, using (41) and

$$\frac{h_{n-2}(y)}{n-1} - \frac{h_{n-2}(y)}{n} = \frac{h_{n-2}(y)}{n^2} = O\left(n^{-(2+\Re(y)-|y|)}\right),$$

we obtain

$$h_n(y) - h_{n-2}(y) - 2y \left( \frac{h_{n-1}(y)}{n} - \frac{h_{n-2}(y)}{n-1} \right) = O\left(n^{-(2+\Re(y)-|y|)}\right). \quad (44)$$

Set

$$R_n(y) := h_n(y) - h_{n-2}(y) - 2y \left( \frac{h_{n-1}(y)}{n} - \frac{h_{n-2}(y)}{n-1} \right).$$

By Lemma 1,  $R_n(y)$  is analytic everywhere for each  $n \geq 3$ . It follows from (44) and Weierstrass M-test that the series  $\sum_{k \geq 3} R_k(y) = K_1(y)$  is analytic when  $\Re(y) - |y| > -1$ , and

$$\sum_{k=3}^n R_k(y) = K_1(y) + O\left(n^{-(1+\Re(y)-|y|)}\right). \quad (45)$$

Summing both sides of (44) from 3 to  $n$  (noting the cancellations from the telescoping sum), we obtain

$$h_n(y) + h_{n-1}(y) - h_1(y) - h_2(y) + y h_1(y) = K_1(y) + O\left(n^{-(1+\Re(y)-|y|)}\right),$$

or

$$h_n(y) + h_{n-1}(y) = K_1(y) + (1-y)h_1(y) + h_2(y) + O\left(n^{-(1+\Re(y)-|y|)}\right). \quad (46)$$

Combining this with (43), and setting

$$K(y) := \frac{K_1(y) + (1-y)h_1(y) + h_2(y)}{2}, \quad (47)$$

we obtain (38). ■

**Remark** Using Theorem 3 and (14), we have

$$H_n(1) \sim K(1)n.$$

Noting

$$C_n = \sum_{g \geq 0} C_{n,g} = \frac{1}{3n+2} \sum_{g \geq 0} H_{n,g} = \frac{n!}{3n+2} 6^n H_n(1),$$

we obtain

$$C_n \sim \frac{K(1)}{3} n! 6^n.$$

Comparing this with the asymptotic expression in Corollary 1, we obtain

$$K(1) = \frac{9}{\pi}.$$

**Proof of Theorem 1** Using (15) and Theorem 3, we have, uniformly for  $x$  in a small neighborhood of 1, that

$$H_n(x) \sim K(1/\sqrt{x}) n^{1/\sqrt{x}} x^{(n+2)/2}. \quad (48)$$

We would like to point out that Hwang's quasi-power theorem [18] does not apply directly here because of the factor  $x^{(n+2)/2}$  appearing in (48). One can apply it to  $J_n(y)$  to derive asymptotic normal distribution for the sequence  $(J_{n,f} : f \geq 1)$ , and then use (2) to obtain (a). This will be illustrated below in the proof of Theorem 2.

Theorem 2 (case (2)) from [11] can be applied directly here by noting (in the notation of [11])

$$\begin{aligned} s &= \ln x, \\ m_r(s) &= \frac{d}{ds} \frac{s}{2} = \frac{1}{2}, \\ m_\alpha(s) &= \frac{d}{ds} e^{-s/2} = -\frac{1}{2} e^{-s/2}, \\ B_r(s) &= 0, \\ B_\alpha(s) &= \frac{d}{ds} m_\alpha(s) = \frac{1}{4} e^{-s/2}. \end{aligned}$$



Now part (a) follows by noting

$$\begin{aligned} m_r(1)n + m_\alpha(1) \ln n &= \frac{1}{2}(n - \ln n), \\ B_\alpha(1) \ln n &= \frac{1}{4} \ln n. \end{aligned}$$

Part (b) follows immediately from part (a) and (2). ■

**Proof of Theorem 2** We may use Theorem 3 and apply the standard saddle-point method directly. In what follows, we shall apply [11, Theorem 4] to  $J_n(x)$ . More precisely, to avoid the parity issue, we shall consider

$$G_n(x) = \left(\sqrt{x}\right)^{\llbracket 2\uparrow n \rrbracket} J_n(\sqrt{x}). \quad (49)$$

By Lemma 1,  $G_n(x)$  is a polynomial in  $x$ . For  $x \neq 0$ , write

$$\begin{aligned} x &= \rho e^{i\theta}, & (-\pi < \theta \leq \pi) \\ \sqrt{x} &= \sqrt{\rho} \exp(i\theta/2). \end{aligned}$$

Then

$$\Re(\sqrt{x}) = \sqrt{\rho} \cos(\theta/2) \geq 0.$$

Using Theorem 3, we obtain, for some small positive constants  $\varepsilon$  and  $\delta$ ,

$$\begin{aligned} G_n(x) &\sim K(\sqrt{x}) \left(\sqrt{x}\right)^{\llbracket 2\uparrow n \rrbracket} n^{\sqrt{x}}, & (\rho < 1, |\theta| \leq \pi - \varepsilon) \\ \frac{G_n(x)}{G_n(\rho)} &= O\left(n^{-\delta}\right). & (\rho < 1, |\theta| \geq \pi - \varepsilon) \end{aligned}$$

Applying [11, Theorem 4(2)] with  $\alpha(s) = e^{s/2}$ , we obtain

$$[x^k]G_n(x) \sim \frac{K(\sqrt{\rho})}{\sqrt{2\pi(\ln n)\sqrt{\rho}/4}} (\sqrt{\rho})^{\llbracket 2\uparrow n \rrbracket} n^{\sqrt{\rho}} \rho^{-k}, \quad (50)$$

where  $\rho$  satisfies

$$\frac{\sqrt{\rho}}{2} = \frac{k}{\ln n} \quad (0 < \rho < 1). \quad (51)$$

Substituting (51) into (50), we obtain

$$[x^k]G_n(x) \sim \frac{K(2k/\ln n)}{\sqrt{k\pi}} e^{2k} \left(\frac{2k}{\ln n}\right)^{\llbracket 2\uparrow n \rrbracket - 2k}. \quad (52)$$

Using (2) and (49), we obtain

$$\begin{aligned}
[x^k]G_n(x) &= \llbracket 2 \mid n \rrbracket [x^{2k}]J_n(\sqrt{x}) + \llbracket 2 \nmid n \rrbracket [x^{2k-1}]J_n(\sqrt{x}) \\
&= \frac{3n+2}{n!} 6^{-n} \left( \llbracket 2 \mid n \rrbracket C_{n, \frac{n}{2}+1-k} + \llbracket 2 \nmid n \rrbracket C_{n, \frac{n+1}{2}+1-k} \right) \\
&= \frac{3n+2}{n!} 6^{-n} C_{n, \frac{n+\llbracket 2 \nmid n \rrbracket}{2}+1-k}
\end{aligned}$$

Combining this with (50) and setting  $k = \frac{n+\llbracket 2 \nmid n \rrbracket}{2} + 1 - g$ , we obtain

$$\begin{aligned}
C_{n,g} &\sim \frac{(n-1)!}{3} 6^n \frac{K(2k/\ln n)}{\sqrt{k\pi}} e^{2k} \left( \frac{2k}{\ln n} \right)^{\llbracket 2 \nmid n \rrbracket - 2k} \\
&\sim \frac{(n-1)!}{3} 6^n \frac{\sqrt{2} K((n-2g)/\ln n)}{\sqrt{(n-2g)\pi}} e^{n-2g+2+\llbracket 2 \nmid n \rrbracket} \\
&\quad \times \left( \frac{n-2g+2+\llbracket 2 \nmid n \rrbracket}{\ln n} \right)^{2g-n-2} \\
&\sim \frac{(n-1)!}{3} 6^n \frac{\sqrt{2} K((n-2g)/\ln n)}{\sqrt{(n-2g)\pi}} \left( \frac{n-2g}{\ln n} \right)^{2g-n-2} \\
&\quad \times e^{n-2g+2+\llbracket 2 \nmid n \rrbracket} \left( 1 + \frac{2+\llbracket 2 \nmid n \rrbracket}{n-2g} \right)^{-(n-2g)} \\
&\sim \frac{(n-1)!}{3} 6^n \frac{\sqrt{2} K((n-2g)/\ln n)}{\sqrt{(n-2g)\pi}} \left( \frac{n-2g}{\ln n} \right)^{2g-n-2} e^{n-2g}.
\end{aligned}$$

Now Theorem 2 follows by using Stirling's formula:

$$(n-2g)! \sim \sqrt{2\pi(n-2g)} \left( \frac{n-2g}{e} \right)^{n-2g} \quad (n-2g \rightarrow \infty).$$

■

## 4 Conclusion

Using the Goulden-Jackson recursion for the number of rooted cubic maps, we derived an asymptotic formula for the number of rooted cubic maps with  $2n$  vertices and genus  $g$  when  $n, g \rightarrow \infty$  and  $(n-2g)/\ln n$  lies in any closed

subinterval of  $(0, 1)$ . The asymptotic formula is accurate up to a constant factor. We also showed that the genus distribution of cubic graphs is asymptotically normal with mean and variance, respectively, asymptotic to  $\frac{1}{2}(n - \ln n)$  and  $\frac{1}{4} \ln n$ . Asymptotic formulas were also obtained for the number of rooted regular maps, disregarding the genus, for constant degree and as the number of vertices going to infinity.

## Acknowledgements

The author would like to thank Mihyun Kang for helpful comments on an earlier draft which improves the presentation.

## References

- [1] O. Angel, G. Chapuy, N. Curien, and G. Ray. The local limit of unicellular maps in high genus. *Electron. Commun. Probab.*, 18:paper no. 86, 8 pp., 2013.
- [2] E. A. Bender, An Asymptotic Expansion for the Coefficients of Some Formal Power Series, *J. London Math. Soc.* **9 (2)** (1975), 451–458.
- [3] E. A. Bender and E. R. Canfield. The asymptotic number of rooted maps on a surface. *J. Combin. Theory Ser. A*, **43(2)** (1986), 244–257.
- [4] E. A. Bender, E. R. Canfield, and L. B. Richmond. The asymptotic number of rooted maps on a surface. II. Enumeration by vertices and faces. *J. Combin. Theory Ser. A*, **63(2)** (1993), 318–329.
- [5] T. Budzinski and B. Louf. Local limits of uniform triangulations in high genus. [arXiv:1902.00492v1](https://arxiv.org/abs/1902.00492v1).
- [6] J. L. Gross and M. L. Furst, Hierarchy of imbedding distribution invariants of a graph, *J. Graph Theory*, **11** (1987), 205–220.
- [7] J. L. Gross and T.W. Tucker, *Topological Graph Theory*, Dover, 2001; (original edn. Wiley, 1987).
- [8] M. L. Furst, J. L. Gross and R. Statman, Genus distributions for two classes of graphs, *J. Combin. Theory (B)* **46** (1989), 22–36.

- [9] Z. Gao. A pattern for the asymptotic number of rooted maps on surfaces. *J. Combin. Theory Ser. A*, 64(2):246–264, 1993.
- [10] Z. Gao and M. Kang, Counting triangulations with high genus, *Proceedings of the 31st International Conference on Probabilistic, Combinatorial and Asymptotic Methods for the Analysis of Algorithms*, Austria, July, 2020.
- [11] Z. Gao and L. B. Richmond, Central and local limit theorems applied to asymptotic enumeration IV: multivariate generating functions, *Journal of Computational and Applied Mathematics* **41** (1992) 177–186.
- [12] I. Goulden and D.M. Jackson, The KP hierarchy, branched covers and triangulations, *Advance in Mathematics* **219**(3) (2008), 932–951.
- [13] J. L. Gross, Genus Distributions of Cubic Outerplanar Graphs, *Journal of Graph Algorithms and Applications*, <http://jgaa.info/> vol. 15, no. 2, pp. 295-316 (2011).
- [14] J. L. Gross, Embeddings of cubic Halin graphs: Genus distributions, *Ars Math. Contemp.* **6** (2013) 37–56.
- [15] J. L. Gross, D.P. Robbins and T.W. Tucker, Genus distributions for bouquets of circles, *J. Combin. Theory B* **47** (1989), 292–306.
- [16] J. L. Gross, M.Kotrbcík and T. Sun, Genus distributions of cubic series-parallel graphs, *Discrete mathematics & theoretical computer science DMTCS* **16**(3), (2014), 129–146.
- [17] J. Harer and D. Zagier, The Euler characteristic of the moduli space of curves, *Invent. Math.*, **85** (1986) 457–485.
- [18] H. K. Hwang, On convergence rates in the central limit theorems for combinatorial structures. *Eur. J. Combin.* **19** (1998), 329–343.
- [19] D. M. Jackson, Counting cycles in permutations by group characters, with an application to a topological problem, *Trans. Amer. Math. Soc.* **299** (1987), 785–801.
- [20] D. M. Jackson, The genus series for maps, *Journal of Pure and Applied Algebra* **105** (1995) 293–297.

- [21] D. M. Jackson and T. I. Visentin, A character theoretic approach to embeddings of rooted maps in an orientable surface of given genus, *Trans. Amer. Math. Soc.* **322** (1990), 343–363.
- [22] R. E. Mickens, Difference equations, New York : Van Nostrand Reinhold, 2nd ed., 1990.
- [23] G. Ray, Large unicellular maps in high genus, *Ann. Inst. Henri Poincaré Probab. Stat.*, **51**(4) (2015), 1432–1456.
- [24] G. Ray, Large unicellular maps in high genus, arXiv:1307.1224v2.
- [25] S. Stahl, Region distributions of graph embeddings and stirling numbers, *Discrete Math.* **82** (1990), 57–78.
- [26] W. T. Tutte, A census of planar maps, *Canadian J. Math.*, **15** (1963), 249–271.
- [27] L. Wan, H. J. Lai and Y. Liu, Genus Distribution for a Graph, *International J.Math. Combin.* Vol.4 (2009), 99–109.