

**CERTAIN PROPERTIES INVOLVING THE
UNBOUNDED OPERATORS $p(T)$, TT^* , AND T^*T ; AND
SOME APPLICATIONS TO POWERS AND n th ROOTS
OF UNBOUNDED OPERATORS**

MOHAMMED HICHEM MORTAD

ABSTRACT. In this paper, we are concerned with conditions under which $[p(T)]^* = \bar{p}(T^*)$, where $p(z)$ is a one-variable complex polynomial, and T is an unbounded, densely defined, and linear operator. Then, we deal with the validity of the identities $\sigma(AB) = \sigma(BA)$, where A and B are two unbounded operators. The equations $(TT^*)^* = TT^*$ and $(T^*T)^* = T^*T$, where T is a densely defined closable operator, are also studied. A particular interest will be paid to the equation $T^*T = p(T)$ and its variants. Then, we have certain results concerning n th roots of classes of normal and nonnormal (unbounded) operators. Some further consequences and counterexamples accompany our results.

1. PRELIMINARIES

First, we assume readers have some familiarity with linear bounded and unbounded operators on Hilbert spaces. Some useful references are [47], [49], and [63]. Classical definitions and notations follow those in [63]. We do recall the definition of the spectrum that will be used here (as this could be different elsewhere): Let A be an operator on a complex Hilbert space H . The resolvent set of A , denoted by $\rho(A)$, is defined by

$$\rho(A) = \{\lambda \in \mathbb{C} : \lambda I - A \text{ is bijective and } (\lambda I - A)^{-1} \in B(H)\}.$$

The complement of $\rho(A)$, denoted by $\sigma(A)$, i.e.,

$$\sigma(A) = \mathbb{C} \setminus \rho(A)$$

is called the spectrum of A .

2020 *Mathematics Subject Classification*. Primary 47B25. Secondary 47B15, 47A05, 47A10, 47B20, 47A08.

Key words and phrases. Closed operator; Symmetric operator; Self-adjoint operator; Normal operator; Quasinormal operator; Hyponormal operator; Operator polynomials; Spectrum; Square roots of operators; n th roots of operators.

Therefore, if A is a linear operator that is not closed, then necessarily $\sigma(A) = \mathbb{C}$.

We also recall the definitions of unbounded nonnormal operators.

Say that a linear operator T is quasinormal, provided it is closed, densely defined, and $TT^*T = T^*TT$ (as in [36]). This condition was weakened to $TT^*T \subset T^*TT$ in [33], but not $TT^*T \supset T^*TT$. By [5], the quasinormality of T is, in fact, equivalent to $T|T| = |T|T$. In [73], T is quasinormal iff $U|T| \subset |T|U$, where $T = U|T|$ is the usual polar decomposition of T in terms of partial isometries. By, e.g., Lemma 2.2 in [52], it is seen that $U|T| \subset |T|U$ is equivalent to $U|T| = |T|U$.

A densely defined linear operator A with domain $D(A) \subset H$, is said to be subnormal when there are a Hilbert space K with $H \subset K$, and a normal operator N with $D(N) \subset K$ such that

$$D(A) \subset D(N) \text{ and } Ax = Nx \text{ for all } x \in D(A).$$

A densely defined A is said to be formally normal if

$$\|Ax\| = \|A^*x\|, \forall x \in D(A) \subset D(A^*).$$

A densely defined operator A with domain $D(A)$ is called hyponormal provided

$$D(A) \subset D(A^*) \text{ and } \|A^*x\| \leq \|Ax\|, \forall x \in D(A).$$

Recall that a linear operator $A : D(A) \subset H \rightarrow H$ is said to be paranormal when

$$\|Ax\|^2 \leq \|A^2x\| \|x\|$$

for all $x \in D(A^2)$. It is straightforward to check that a hyponormal operator is paranormal. However, and unlike the class of hyponormal operators, a densely defined paranormal operator need not be closable. Also, the closure of a densely defined closable paranormal operator fails, in general, to be paranormal. See [9] or [50]. In the latter reference, an example of a densely defined paranormal operator T that satisfies $D(T^*) = \{0\}$ is provided. Also, readers could find an example of a closed densely defined operator T such that both T and T^* are one-to-one and paranormal, yet T is not normal (cf. [1] and [84]).

It is widely known that a normal operator is quasinormal, a quasinormal operator is subnormal, a subnormal operator is hyponormal, and a hyponormal operator is paranormal. It is also known that one cannot go backward in the last implications, even for bounded operators (see [49] for counterexamples). Readers may consult, e.g., [77] to see how all those (and more classical) notions interplay.

2. INTRODUCTION

Taking the adjoint of the product or the sum of densely defined unbounded operators can be an arduous task in some cases, and it can be a major impediment to advances in many proofs, which makes us frustrated before such situations. Recall that when A , B and AB are all densely defined operators, then $A^*B^* \subset (BA)^*$, and the equality

$$(BA)^* = A^*B^*$$

holds true if, e.g., $B \in B(H)$. It also holds when $A^{-1} \in B(H)$.

The following lemma, which will be needed below, was shown in Corollary 1.7 in [30].

Lemma 2.1. *Let A and B be two densely defined closed operators such that $\sigma(AB) \neq \mathbb{C}$ and $\sigma(BA) \neq \mathbb{C}$. Then AB and BA are two densely defined closed operators that obey*

$$(AB)^* = B^*A^* \text{ and } (BA)^* = A^*B^*.$$

Some very related papers are [3], [7], [27], [70], and certain references therein. As for sums, readers may consult, e.g., [31], [45], and [70].

Notice that conditions like $\sigma(AB) \neq \mathbb{C}$ and $\sigma(BA) \neq \mathbb{C}$ are strong, but they entail interesting consequences as regards the adjoint of the product of two unbounded operators and their spectrum. Remember that when $A, B \in B(H)$, then $\sigma(AB) \neq \mathbb{C}$ and $\sigma(BA) \neq \mathbb{C}$. Some related papers are [2], [12], and [54].

Let T be a densely defined closed operator. One of the most fundamental properties in unbounded operator theory is the fact that T^*T (and also TT^*) is a densely defined, self-adjoint and positive operator. This is a very well-known von-Neumann's theorem. In fact, von-Neumann's result may be obtained via the so-called Nelson's trick (cf. [79]). In particular,

$$(TT^*)^* = TT^* \text{ and } (T^*T)^* = T^*T.$$

This result then enables us to define the very important notion of the modulus of an operator which, and as it is known, intervenes in the definition of the polar decomposition of an operator. We may also find it in an abstract version of the damped wave equation of the form $\ddot{u} + R\dot{u} + T^*Tu = 0$ under certain conditions and in some separable Hilbert space, where T is a densely defined closed operator and R is some perturbation of T^*T . This equation has been extensively investigated in [22]. Operator theorists are also well aware of other uses of the above result. Some related results may be found in [6], [49], [59], and [66].

Recently, Z. Sebestyén and Zs. Tarcsay have discovered that if TT^* and T^*T are both self-adjoint, then T must be closed (see [68], cf.

[14] and [23]). In this paper, we deal with the validity of the identities $(TT^*)^* = TT^*$ and $(T^*T)^* = T^*T$, where T is a densely defined closable operator. In the end, we supply a few counterexamples that show interesting pathological properties of TT^* and T^*T .

But, before that, we will be first investigating the validity of the equality $[p(T)]^* = \bar{p}(T^*)$, where $p(z)$ is a one-variable complex polynomial, and T is densely defined. Then, a few consequences about powers of operators are obtained. Readers might be interested in papers dealing with polynomials of closed (or other classes of) operators such as: [10], [53], [60], [72], and [75].

In addition, we have some useful results concerning n th roots of certain classes of normal and nonnormal (unbounded) operators.

We close the paper with several interesting counterexamples related to certain of the results to be shown here.

3. ON THE ADJOINT OF $p(T)$

Before giving the general case, we present a simple consequence of Lemma 2.1. First, recall that it is well known that $(T^n)^* = T^{*n}$ for any $T \in B(H)$ and any n . The unbounded version need not hold even when $n = 2$. See, e.g., Question 20.2.14 in [49] for a counterexample. Remember that if T^n (hence T) is densely defined, then only $(T^*)^n \subset (T^n)^*$ holds. If T is normal, then one does have $(T^n)^* = T^{*n}$ for any n (a proof may be consulted in say [55]). Jabłoński et al. constructed in [33], a quasinormal operator T such that $T^{*n} \subsetneq (T^n)^*$ for all $n \geq 2$ (cf. Lemma 3.8 in [14]).

By Lemma 2.1, it is seen that when T is closed and $\sigma(T^2) \neq \mathbb{C}$, then $(T^2)^* = T^{*2}$. However, a naive direct generalization to the case T^n , $n \in \mathbb{N}$, requires an induction argument, some standard facts, and rather strong assumptions involving the closedness and spectra of T^p , $p \leq n$.

In fact, we can prove a much stronger result, which will also allow us to present a new proof of a slightly improved Lemma 2.1.

Let $p(z) = a_n z^n + \cdots + a_1 z + a_0$, where a_n, \dots, a_1, a_0 are complex numbers, and z is a complex variable. Set $\bar{p}(z) = \bar{a}_n z^n + \cdots + \bar{a}_1 z + \bar{a}_0$.

It is easy to see that $D[p(T)] = D(T^n)$. So, when T^n is densely defined, then T is a densely defined operator. But, in general, only $[p(T)]^* \supset \bar{p}(T^*)$ holds. The coming result seems therefore interesting.

Theorem 3.1. *Let $n \in \mathbb{N}$ and let T be a linear closable operator on a Hilbert space such that T^n is densely defined. If $\sigma[p(T)] \neq \mathbb{C}$, then $[p(T)]^* = \bar{p}(T^*)$, where $p(z)$ is a given complex polynomial of degree n .*

Remark. Recall that the closedness of T alone does not, in general, yield the closedness of T^n , neither it implies its closability. For instance, in Question 22.2.6 in [49], there is densely defined closed operator T such that T^2 is also densely defined but $(T^2)^*$ is not densely defined, i.e., T^2 is not closable.

The proof seems to need the coming lemma, whose proof is essentially contained in the proof of Theorem 2.1 in [14].

Lemma 3.2. *Let p be a complex polynomial of one variable of degree n (whose leading coefficient equals 1, WLOG). Assume that A is a closable operator in a Banach space such that $\sigma[p(A)] \neq \mathbb{C}$. If $\lambda \in \rho[p(A)]$ is such that*

$$p(A) - \lambda I = (A - \mu_1 I)(A - \mu_2 I) \cdots (A - \mu_n I)$$

for certain complex numbers $\mu_1, \mu_2, \dots, \mu_n$, then all of $A - \mu_1 I$, $A - \mu_2 I$, ..., and $A - \mu_n I$ are boundedly invertible. In particular, $\sigma(A) \neq \mathbb{C}$, and so A is closed.

Proof. By the fundamental theorem of algebra, we know that there always exist complex numbers $\mu_1, \mu_2, \dots, \mu_n$ such that

$$p(z) - \lambda = (z - \mu_1)(z - \mu_2) \cdots (z - \mu_n),$$

where $z \in \mathbb{C}$. Hence

$$p(A) - \lambda I = (A - \mu_1 I)(A - \mu_2 I) \cdots (A - \mu_n I)$$

with $D[p(A) - \lambda I] = D(A^n)$. In [14], it was shown that $A - \mu_1 I$ is boundedly invertible (hence $\sigma(A) \neq \mathbb{C}$, and so A is closed). But since $(A - \mu_1 I)(A - \mu_2 I) \cdots (A - \mu_n I) = (A - \mu_2 I)(A - \mu_3 I) \cdots (A - \mu_n I)(A - \mu_1 I)$, $A - \mu_2 I$ too is boundedly invertible. A similar argument gives the bounded invertibility of the remaining terms. The proof is complete. \square

Now, we show Theorem 3.1:

Proof. First, observe that since T is closed and $\rho(T) \neq \emptyset$, $p(T)$ is densely defined and closed (cf. [41]).

Let λ be in $\mathbb{C} \setminus \sigma[p(T)]$. Write

$$p(z) - \lambda = (z - \mu_1)(z - \mu_2) \cdots (z - \mu_n),$$

where $\mu_1, \mu_2, \dots, \mu_n$ are complex numbers. Then

$$\overline{p(z) - \lambda} = \overline{p(\bar{z}) - \bar{\lambda}} = (\bar{z} - \bar{\mu}_1)(\bar{z} - \bar{\mu}_2) \cdots (\bar{z} - \bar{\mu}_n).$$

Symbolically,

$$p(T) - \lambda I = (T - \mu_1 I)(T - \mu_2 I) \cdots (T - \mu_n I)$$

and

$$\overline{p}(T^*) - \overline{\lambda}I = (T^* - \overline{\mu_1}I)(T^* - \overline{\mu_2}I) \cdots (T^* - \overline{\mu_n}I).$$

Since each of $T - \mu_1I$, $T - \mu_2I$, ..., and $T - \mu_nI$ is boundedly invertible, we obtain

$$\begin{aligned} [p(T)]^* - \overline{\lambda}I &= (p(T) - \lambda I)^* \\ &= [(T - \mu_1I)(T - \mu_2I) \cdots (T - \mu_nI)]^* \\ &= (T - \mu_nI)^* [(T - \mu_1I)(T - \mu_2I) \cdots (T - \mu_{n-1}I)]^* \\ &= \cdots \\ &= (T - \mu_nI)^* \cdots (T - \mu_2I)^* (T - \mu_1I)^* \\ &= (T^* - \overline{\mu_n}I) \cdots (T^* - \overline{\mu_2}I)(T^* - \overline{\mu_1}I) \\ &= (T^* - \overline{\mu_1}I)(T^* - \overline{\mu_2}I) \cdots (T^* - \overline{\mu_n}I) \\ &= \overline{p}(T^*) - \overline{\lambda}I. \end{aligned}$$

Accordingly, $[p(T)]^* = \overline{p}(T^*)$, as wished. \square

Corollary 3.3. *Let T be a densely defined linear closable operator on a Hilbert space. If $\sigma[p(T)] \neq \mathbb{C}$, then*

$$\overline{p(\overline{T})} = p(\overline{T}),$$

where $p(z)$ is a given complex polynomial.

Proof. By Theorem 3.1, $[p(T)]^* = \overline{p}(T^*)$. Since T^* is closed, and $\sigma[(p(T))^*] \neq \mathbb{C}$, so to speak, $\sigma[\overline{p}(T^*)] \neq \mathbb{C}$, it follows that

$$\overline{p(\overline{T})} = [p(T)]^{**} = [\overline{p}(T^*)]^* = \overline{\overline{p}(T^*)} = p(\overline{T}),$$

as desired. \square

We have yet another consequence, which first appeared in Theorem 2.3 in [14] (cf. [53], [75], and [69]).

Corollary 3.4. *Let p be a real polynomial of one variable of degree $n \geq 1$. Assume that T is a symmetric (not necessarily densely defined) operator in a Hilbert space H such that $p(T)$ is self-adjoint. Then $f(T)$ is self-adjoint for any real-valued Borel function on \mathbb{R} , where $D[f(T)] = \{x \in H : \int_{\mathbb{R}} |f(t)|^2 d\langle E_t x, x \rangle < \infty\}$ and E_t is the associated spectral measure with T .*

Proof. Since p is a non-constant polynomial, we know that there exists an $\alpha \in \mathbb{C} \setminus \mathbb{R}$ such that $p(\alpha) := \lambda \in \mathbb{C} \setminus \mathbb{R}$ (see [75]). By the self-adjointness of $p(T)$, T is closed. Now, let $\lambda \notin \sigma[p(T)]$. Then by the spectral mapping theorem (see, e.g., [24] or Section 5.7 in [78]), $\alpha \notin \sigma(T)$.

Then, we may write

$$p(z) - \lambda = (z - \alpha)q(z) = q(z)(z - \alpha),$$

where $q(z)$ is a polynomial of degree $n - 1$. Hence

$$p(T) - \lambda I = (T - \alpha I)q(T) = q(T)(T - \alpha I)$$

and

$$p(T^*) - \lambda I = (T^* - \alpha I)q(T^*) = q(T^*)(T^* - \alpha I)$$

Since $p(T) = [p(T)]^* = p(T^*)$ and $T - \alpha I$ is boundedly invertible, it ensues that

$$p(T) - \bar{\lambda}I = [p(T) - \lambda I]^* = (T^* - \bar{\alpha}I)[q(T)]^*.$$

The self-adjointness of $p(T)$ with a glance at the basic criterion for self-adjointness imply

$$\ker(p(T) - \lambda I) = \ker(p(T) - \bar{\lambda}I) = \{0\},$$

from which we derive

$$\ker(T^* - \alpha I) = \ker(T^* - \bar{\alpha}I) = \{0\}.$$

Thus T is self-adjoint by the basic criterion for self-adjointness given that T is already closed and symmetric. The last statement about the self-adjoint of $f(T)$ is a mere consequence of the spectral theorem for unbounded self-adjoint operators. \square

The next consequence is Corollary 19 in [75]:

Corollary 3.5. *Let p be a real polynomial of one variable of degree $n \geq 1$. Assume that T is a symmetric operator in a Hilbert space H such that $p(T)$ is essentially self-adjoint. Then $f(\bar{T})$ is self-adjoint for any real-valued Borel function on \mathbb{R} , where $D[f(\bar{T})] = \{x \in H : \int_{\mathbb{R}} |f(t)|^2 d\langle E_t x, x \rangle < \infty\}$ and E_t is the associated spectral measure with \bar{T} .*

Proof. Since a symmetric T is formally normal, $p(\bar{T}) = \overline{p(T)}$ by [75]. Hence $p(\bar{T})$ is self-adjoint. Thus, \bar{T} is self-adjoint by Corollary 3.4. \square

4. ON THE EQUALITIES $(AB)^* = B^*A^*$ AND $\sigma(AB) = \sigma(BA)$ FOR UNBOUNDED OPERATORS

We begin by reproving Lemma 2.1 under slightly weaker assumptions, namely the closability of A and B instead of their closedness.

Proposition 4.1. *Let A and B be two densely defined closable operators in a Hilbert space H , such that $\sigma(AB) \neq \mathbb{C}$ and $\sigma(BA) \neq \mathbb{C}$. Then A and B are closed. Besides, AB and BA are two densely defined closed operators that obey*

$$(AB)^* = B^*A^* \text{ and } (BA)^* = A^*B^*.$$

Hence

$$(A^*B^*)^* = BA \text{ and } (B^*A^*)^* = AB.$$

Proof. Set

$$T = \begin{pmatrix} 0 & A \\ B & 0 \end{pmatrix}$$

with $D(T) = D(A) \oplus D(B)$. Because A and B are closable, T is closable, too. But

$$T^2 = \begin{pmatrix} AB & 0 \\ 0 & BA \end{pmatrix}.$$

By assumption, $\sigma(AB) \neq \mathbb{C}$ and $\sigma(BA) \neq \mathbb{C}$. Lemma 2.4 in [29] then yields $\sigma(AB) \cup \sigma(BA) \neq \mathbb{C}$, i.e., it is seen that $\sigma(T^2) \neq \mathbb{C}$. Hence T closed, from which we may derive the closedness of A as well as that of B . So, T^2 is densely defined, which forces both AB and BA to be densely defined.

Finally, since $T^* = \begin{pmatrix} 0 & B^* \\ A^* & 0 \end{pmatrix}$, and as $(T^2)^* = T^{*2}$, inspection of the entries of the two corresponding operator matrices gives the desired identities

$$(AB)^* = B^*A^* \text{ and } (BA)^* = A^*B^*,$$

as needed.

To obtain the last claim, just take adjoints, by remembering that AB and BA are closed. \square

The next similar result is stated without proof.

Corollary 4.2. *Let T and S be linear closable operators on a Hilbert space. Let p and q be two one-variable complex polynomials of degrees n and m , respectively. If $\sigma[p(T)] \neq \mathbb{C}$ and $\sigma[q(S)] \neq \mathbb{C}$, $\sigma[p(T)q(S)] \neq \mathbb{C}$, and $\sigma[q(S)p(T)] \neq \mathbb{C}$, then $p(T)q(S)$ and $q(S)p(T)$ are densely defined, and*

$$[p(T)q(S)]^* = \bar{q}(S^*)\bar{p}(T^*) \text{ and } [q(S)p(T)]^* = \bar{p}(T^*)\bar{q}(S^*).$$

It is well-known that an expression like $AB = BA$, on some common dense domain, does not always yield the strong commutativity of A and B (which are assumed to be unbounded and self-adjoint). This is the famous Nelson's counterexample (see [51] or [58]). Nelson's example

is developed in detail in [62], pp 257-258. The simplest Nelson-like example in the literature is due to K. Schmüdgen in [61]. So, the next consequence of Proposition 4.1 is worth stating.

Corollary 4.3. (Cf. [18], [46]) *Let A and B be two unbounded self-adjoint operators. If $AB = BA$ and $\sigma(AB) \neq \mathbb{C}$, then A strongly commutes with B .*

Proof. By assumption, $\sigma(AB) \neq \mathbb{C}$ and $\sigma(BA) \neq \mathbb{C}$. Hence $(AB)^* = BA = AB$, i.e., AB is self-adjoint. By, e.g. [18], A and B strongly commute. \square

It is widely known that

$$\sigma(AB) - \{0\} = \sigma(BA) - \{0\}$$

for any $A, B \in B(H)$, with $\dim H = \infty$ (cf. [17] and [29]). Imposing certain conditions on A and/or B gives the full equality $\sigma(AB) = \sigma(BA)$. For instance, when A is normal. See [4]. However, such a result was at least known at the time of [8], and this was missed by some authors, including myself. See [32] and [47] for other instances of when this is true, and [49] for related counterexamples. It appears, however, that the most general condition guaranteeing the equality of spectra is $\ker A = \ker A^*$ (or $\ker B = \ker B^*$). This appeared in [12], in the case B is in $B(H)$ and A is a densely defined, non-necessarily bounded, and closed operator. Below we generalize this result to two unbounded operators.

Theorem 4.4. *Let A and B be two densely defined closable operators in a Hilbert space H , such that $\sigma(AB) \neq \mathbb{C}$ and $\sigma(BA) \neq \mathbb{C}$. If $\ker(A^*) = \ker(A)$ and $\ker(B) = \ker(B^*)$, then*

$$\sigma(BA) = \sigma(AB).$$

Remark. The preceding theorem improves Equation 3.2 in Proposition 3.1 in [2].

Proof. By Proposition 4.1, A and B are closed. Then, Theorem 1.1 in [29] gives $\sigma(AB) - \{0\} = \sigma(BA) - \{0\}$. What remains to show is:

$$BA \text{ is boundedly invertible} \iff AB \text{ is boundedly invertible.}$$

Assume that BA is boundedly invertible, and so BA is right invertible. That is, $BAT = I$ for some $T \in B(H)$. Since AT is closed and $D(AT) = H$, $AT \in B(H)$. So, B is right invertible, but because $\ker B \subset \ker B^*$, it follows that B is boundedly invertible (Theorem 2.3 in [12]). Since $(BA)^*$ is boundedly invertible, so is $A^*B^* = (BA)^*$. In particular, A^*B^* is right invertible, and a similar argument as above

yields the right invertibility of A^* . Since $\ker A^* \subset \ker A$, Theorem 2.3 in [12] implies the bounded invertibility of A^* , or that of A . In other words, AB is boundedly invertible.

Conversely, suppose AB is boundedly invertible. Hence $(AB)^* = B^*A^*$ too is boundedly invertible. Now, apply the first part of the proof together with the conditions $\ker B^* \subset \ker B$ and $\ker A \subset \ker A^*$ to obtain that A^*B^* is boundedly invertible. Accordingly, $BA = (A^*B^*)^*$ is boundedly invertible, establishing the result. \square

Corollary 4.5. *Let A and B be two unbounded normal operators such that $\sigma(AB) \neq \mathbb{C}$ and $\sigma(BA) \neq \mathbb{C}$. Then*

$$\sigma(BA) = \sigma(AB).$$

The following simple and related observation could interest some readers:

Proposition 4.6. *Let A and B be two self-adjoint operators such that $B \in B(H)$ is positive and BA is closed. Then*

$$\sigma(AB) \neq \mathbb{C} \iff \sigma(AB) \subset \mathbb{R}.$$

Proof. We are only concerned with the implication " \Rightarrow ", so suppose $\sigma(AB) \neq \mathbb{C}$. Let \sqrt{B} be the unique positive square root of B . Since BA is closed on $D(A)$, by Proposition 3.7 in [12], $\sqrt{B}A$ is also closed. Since $\sqrt{B}A$ is domain-dense, $(\sqrt{B}A)^* = A\sqrt{B}$. By the closedness of $\sqrt{B}A$, $A\sqrt{B}$ is densely defined. Therefore,

$$\sqrt{B}A = (\sqrt{B}A)^{**} = (A\sqrt{B})^*.$$

Since $\sqrt{B} \in B(H)$, it is seen that

$$(\sqrt{B}A\sqrt{B})^* = (A\sqrt{B})^*\sqrt{B} = \sqrt{B}A\sqrt{B},$$

which means that $\sqrt{B}A\sqrt{B}$ is self-adjoint. So $\sigma(\sqrt{B}A\sqrt{B}) \subset \mathbb{R}$ or $\sigma(\sqrt{B}A\sqrt{B}) \neq \mathbb{C}$. Since $\sigma(AB) \neq \mathbb{C}$ means that $\sigma(A\sqrt{B}\sqrt{B}) \neq \mathbb{C}$, we infer that

$$\sigma(AB) - \{0\} = \sigma(A\sqrt{B}\sqrt{B}) - \{0\} = \sigma(\sqrt{B}A\sqrt{B}) - \{0\}$$

(utilizing Theorem 1.1 in [29]). Thus, $\sigma(AB) \subset \mathbb{R}$, as needed. \square

Remark. It is worth recalling that if the closedness of BA is replaced by the normality of AB in the result above (resp. the normality of BA), then AB is self-adjoint (resp. BA and AB are self-adjoint). In particular, $\sigma(AB) \subset \mathbb{R}$. See [43] for a proof. Related results about unbounded operators may be consulted in [15], [28], [34], and [44].

Similarly, we have:

Proposition 4.7. *Let A and B be two self-adjoint operators such that $B \in B(H)$ is positive. If \sqrt{B} has a closed range, and has finite dimensional kernel, then*

$$\sigma(AB) \neq \mathbb{C} \iff \sigma(AB) \subset \mathbb{R}.$$

Proof. By [26], $\sqrt{B}A\sqrt{B}$ is self-adjoint. The rest of the proof is as above. \square

5. ON n th ROOTS OF SOME CLASSES OF UNBOUNDED OPERATORS

It is plain that if $T \in B(H)$, then T^2 is self-adjoint if and only if $T^2 = T^{*2}$. This is not the case anymore when T is closed and densely defined. Indeed, there is a closed densely defined operator T such that

$$D(T^2) = D(T^{*2}) = \{0\}$$

(hence T^2 cannot be self-adjoint). Such an example may be found in [13]. The situation is not better when $D(T^2)$ is dense (see [14]), neither it is for higher powers (witness [48]). Nonetheless, we have:

Corollary 5.1. *Let $n \in \mathbb{N}$ be fixed and let T be a linear closable operator on a Hilbert space. Then*

$$T^n \text{ is self-adjoint} \iff T^n = T^{*n} \text{ and } \sigma(T^n) \neq \mathbb{C}.$$

Proof. If T^n is self-adjoint, then $\sigma(T^n) \neq \mathbb{C}$ and

$$T^n = (T^n)^* = T^{*n}.$$

Conversely, $T^n = T^{*n}$ and $\sigma(T^n) \neq \mathbb{C}$ imply $(T^n)^* = T^n$. Consequently, T^n is self-adjoint. \square

Before giving a result that generalizes Proposition 3.7 in [14], recall the ensuing result, which might be known under different proofs.

Lemma 5.2. *Let T be a quasinormal (unbounded) operator. Then T^n is closed and densely defined, for each $n \in \mathbb{N}$.*

Proof. Let $n \in \mathbb{N}$. Write $T = U|T| = |T|U$, where $U \in B(H)$ is a partial isometry. Then

$$T^n = (U|T|)^n = U^n|T|^n = |T|^nU^n.$$

Since T is closed, $|T|$ is self-adjoint, as are its powers $|T|^n$. Since $D(T^n) = D(|T|^n)$, T^n is densely defined.

Since $|T|^nU^n$ is closed for each n , T^n too is closed. \square

Here is the promised generalization.

Proposition 5.3. *Let T be a quasinormal (unbounded) operator such that $T^n = T^{*n}$ for some natural number n . Then T^n is self-adjoint, and T is normal.*

In particular, and when $n = 2$, then T is either self-adjoint or skew-adjoint.

Proof. We claim that

$$|T|^{2n} = |T^*|^{2n} = T^{2n}.$$

By Corollary 3.1 in [81], we know that

$$T^{*n}T^n = (T^*T)^n \geq (TT^*)^n \geq T^nT^{*n},$$

whenever T is quasinormal, and for all $n \geq 2$. Hence

$$T^{2n} = (T^*T)^n = |T|^{2n} \geq |T^*|^{2n} = (TT^*)^n \geq T^{2n},$$

still for any $n \geq 2$. Therefore, we obtain $|T|^{2n} = |T^*|^{2n} = T^{2n}$.

So, T^{2n} is self-adjoint and positive. Since T is quasinormal, T^n is quasinormal and densely defined. Since a quasinormal operator is hyponormal, Corollary 3.6 in [14] gives the self-adjointness of T^n .

Now, since $|T|^{2n} = |T^*|^{2n}$, upon passing to the unique self-adjoint positive n -th root, we get $|T|^2 = |T^*|^2$, that is, T is normal.

Finally, let $n = 2$. Then T^2 is self-adjoint and T is normal. Let $\lambda \in \sigma(T)$, and so $\lambda^2 \in \sigma(T^2) \subset \mathbb{R}$. Then, either $\text{Im } \lambda = 0$ or $\text{Re } \lambda = 0$. In the latter case, T is skew-adjoint, and in the former case, T is self-adjoint. This marks the end of the proof. \square

Remark. Another way of seeing that T^n is self-adjoint is: The condition $T^n = T^{*n}$ implies that T^n is symmetric. Since T^n is quasinormal, the corollary to Theorem 3.3 in [81] gives the self-adjointness of T^n .

From the previous proof and Theorem 3.1, we have:

Corollary 5.4. *Let $n \in \mathbb{N}$ and let T be a quasinormal (unbounded) operator. Then*

$$T^n \text{ is self-adjoint} \iff T^n = T^{*n}.$$

The next result to be given below these lines is already known in the case of normal operators, and it is most probably known to specialists in the case of paranormal operators, which is the version to be shown. We include a simple proof for ease of reference.

Lemma 5.5. *Let $n \in \mathbb{N}$ and let T be a non-necessarily bounded paranormal operator. Then*

$$\ker T = \ker T^2 = \dots = \ker T^n.$$

Hence $\overline{\text{ran } T^n} = \dots = \overline{\text{ran } T^2} = \overline{\text{ran } T}$.

Proof. First, observe that we always have $\ker T \subset \ker T^n$ for any n (and for *any* linear operator). Indeed, let $x \in \ker T$, i.e., $x \in D(T)$ and $Tx = 0$. Hence $Tx \in D(T^{n-1})$ and $T^n x = 0$, i.e., $x \in \ker T^n$.

To show the converse, recall that from [76], we know that for all $x \in D(T^{n+1})$, the following inequalities hold:

$$\|T^n x\| \leq \|T^{n+1} x\|^{n/(n+1)} \|x\|^{1/(n+1)}$$

and

$$\|Tx\| \leq \|T^{n+1} x\|^{1/(n+1)} \|x\|^{n/(n+1)}.$$

Let $x \in \ker T^n$, that is, $T^n x = 0$ with $x \in D(T^n)$ (hence $x \in D(T)$). Therefore, $x \in \ker T$, as wished.

The last statement holds by standard arguments. \square

Corollary 5.6. *Let T be a quasinormal (unbounded) operator such that $T^n = T^{*n}$ for some natural n . Then T is normal.*

Proof. Since T is quasinormal and $T^n = T^{*n}$, it is seen that

$$\ker T = \ker T^n = \ker T^{*n} \supset \ker T^*.$$

But, a quasinormal operator is hyponormal, and so $\ker T \subset \ker T^*$, i.e. $\ker T = \ker T^*$. By Corollary 3.2 in [81], T is normal. \square

The next consequence appeared in Theorem 3.9 in [14]. The proof here is, however, much shorter.

Corollary 5.7. *Let T be a quasinormal (unbounded) operator such that T^n is normal for a certain $n \geq 2$. Then T is normal.*

Proof. Since T is quasinormal, $\ker T = \ker T^n$. Since T^n is normal, $\ker(T^n) = \ker[(T^n)^*]$. As $T^{*n} \subset (T^n)^*$, $\ker T^* \subset \ker[(T^n)^*]$. Therefore, $\ker T^* \subset \ker T$, and because we already have $\ker T \subset \ker T^*$, we obtain $\ker T^* = \ker T$. Thus, and by invoking Corollary 3.2 in [81], T is normal, as needed. \square

Remark. There is yet another way of showing the last result. According to Corollary 3.3 in [81], we obtain the normality of T from the quasinormality of T , $\ker T^* = \ker T^{*2}$, and $\ker T^* \subset D(T^*T)$. Let us then show the last two conditions by supposing T is quasinormal and T^n is normal for some $n \geq 2$:

That $\ker T^* \subset \ker T^{*2}$ is clear. Then

$$\ker T^{*2} \subset \ker T^{*n} \subset \ker(T^n)^* = \ker T^n = \ker T \subset \ker T^*,$$

whereby $\ker T^* = \ker T^{*2}$.

Now, let $x \in \ker T^*$. As in the above proof, $x \in \ker T$, i.e., $x \in D(T)$ and $Tx = 0$. Hence $Tx \in D(T^*)$, that is, $x \in D(T^*T)$, as needed.

Corollary 5.7 has a certain generalization to polynomials.

Proposition 5.8. *Let T be a quasinormal (unbounded) operator in a complex Hilbert space H . Let $p(z)$ be a non-constant complex polynomial such that $p(0) = 0$, and which is expressed as*

$$p(z) = z^{n_1}(z - \lambda_2)^{n_2} \cdots (z - \lambda_k)^{n_k},$$

where $0, \lambda_2, \dots, \lambda_k$ are distinct roots. Assume that $\lambda_2, \dots, \lambda_k \notin \sigma_p(T)$. If $p(T)$ is normal, then T is normal.

Proof. We have

$$p(T) = T^{n_1}(T - \lambda_2 I)^{n_2} \cdots (T - \lambda_k I)^{n_k}.$$

By [25] (or Lemma 3.1.15 in [39]), it follows that

$$\ker p(T) = \ker(T^{n_1}) \oplus \ker(T - \lambda_2 I)^{n_2} \oplus \cdots \oplus \ker(T - \lambda_k I)^{n_k}.$$

Since a quasinormal operator is hyponormal, it ensues that all of $T, T - \lambda_2 I, \dots, T - \lambda_k I$ are hyponormal. So, Lemma 5.5 implies

$$\ker p(T) = \ker T \oplus \ker(T - \lambda_2 I) \oplus \cdots \oplus \ker(T - \lambda_k I).$$

But $\lambda_2, \dots, \lambda_k \notin \sigma_p(T)$, and so

$$\ker(T - \lambda_2 I) = \cdots = \ker(T - \lambda_k I) = \{0_H\}.$$

Therefore, $\ker p(T) = \ker T$. By the normality of $p(T)$, we know that $\ker p(T) = \ker[(p(T))^*]$. Since $p(0) = 0$, it is seen that

$$\ker T^* \subset \ker \bar{p}(T^*) \subset \ker[(p(T))^*] = \ker p(T) = \ker T.$$

Consequently, T is normal by proceeding as in the end of the proof of Corollary 5.7. \square

The verbatim extension to symmetric operators is also valid. We need, however, to add an assumption but then the conclusion is stronger.

First, we have an auxiliary lemma, which characterizes self-adjointness. It is interesting enough to be singled out.

Lemma 5.9. *Let T be an unbounded, densely defined, and symmetric operator in a Hilbert space H , i.e., $T \subset T^*$. If $\text{ran } T$ is closed and $\ker T = \ker T^*$, then T is self-adjoint.*

Proof. Observe that

$$\ker T = \ker T^* = (\text{ran } T)^\perp.$$

Hence $(\ker T)^\perp = \text{ran } T$ thanks to the closedness of $\text{ran } T$. Therefore, $\ker T \oplus \text{ran } T = H$. By Corollary 6.5 in [71], T is self-adjoint. \square

Remark. There is another way to establish the preceding lemma. First, remember that $\text{ran } T$ is closed if and only if $\text{ran } T^*$ is closed (see Theorem 5.13, IV, [35]). Then apply Proposition 2.1 in [67].

Proposition 5.10. *Let T be an unbounded, densely defined, and symmetric operator in a complex Hilbert space H such that its range $\text{ran } T$ is closed. Let $p(z)$ be a non-constant complex polynomial such that $p(0) = 0$, and which is expressed as*

$$p(z) = z^{n_1}(z - \lambda_2)^{n_2} \cdots (z - \lambda_k)^{n_k},$$

where $0, \lambda_2, \dots, \lambda_k$ are distinct roots. Assume that $\lambda_2, \dots, \lambda_k \notin \sigma_p(T)$. If $p(T)$ is normal, then T is self-adjoint.

Proof. Since a symmetric operator is hyponormal, obtain, as in the proof of Proposition 5.8, that $\ker T = \ker T^*$. Then, apply Lemma 5.9 to get the self-adjointness of T , as wished. \square

The following result gives a simple characterization of orthogonal projections (cf. [67]).

Proposition 5.11. *Let T be a symmetric operator with domain $D(T) \subset H$, a priori non-densely defined. Assume $T^2 \subset T$. Then the following assertions are equivalent:*

- (1) $T \in B(H)$ is an orthogonal projection,
- (2) T^n is self-adjoint for some $n \geq 2$.

Proof. We only show the implication "(2) \Rightarrow (1)". Since $T^2 \subset T$, it follows that $T^n \subset T^2 \subset T$ for any $n \geq 2$. Since T^n is self-adjoint, T is densely defined. So, T is a densely defined symmetric operator. By maximality, $T^n = T$, and so T is self-adjoint. Besides, $T = T^2$, i.e., T is idempotent. Since $D(T) = D(T^2)$ and T is self-adjoint, it ensues that $T \in B(H)$ (by Lemma 2.1 in [65], say), as desired. \square

It is known that the condition $D(A) = D(B)$ does not imply $D(A^2) = D(B^2)$, even when A and B are self-adjoint positive operators. See Question 21.2.51 in [49] for a counterexample. However, it is shown in ([83], Theorem 9.4) that if A and B are two self-adjoint positive operators with domains $D(A)$ and $D(B)$ respectively, then

$$D(A) = D(B) \implies D(\sqrt{A}) = D(\sqrt{B}).$$

It is therefore natural to wonder whether this property remains valid for arbitrary square roots. That is, if A and B are square roots of some S , i.e. $A^2 = B^2 = S$, is it true that $D(A) = D(B)$? This question,

which is meaningless over $B(H)$, has a negative answer for unbounded operators. See Example 8.4.

Nonetheless, if the square roots are self-adjoint then they necessarily have equal domains. In fact, a much better result holds true, and for arbitrary n th roots. It reads:

Proposition 5.12. *Let A and B be two quasinormal operators such that $A^n = B^n$ for some $n \in \mathbb{N}$, $n \geq 2$. Then $D(A) = D(B)$.*

Proof. Since A and B are quasinormal, we have

$$A^n = B^n \implies |A|^n = |A^n| = |B^n| = |B|^n$$

(see, e.g., Corollary 3.8 in [33]). Upon passing to the unique positive self-adjoint n th root implies $|A| = |B|$. Therefore, $D(A) = D(B)$, as desired. \square

The generalization to unbounded closed hyponormal operators seems difficult to obtain (if it is ever true), however, we have the following result, whose proof is omitted.

Proposition 5.13. *Let A and B be two (closed) hyponormal operators such that $A^2 = B^2$. Assume further that A^2 is self-adjoint and positive. Then $D(A) = D(B)$.*

We finish this section with a result in the same spirit but the approach is different. More precisely, we show that a square root of a symmetric operator is normal, if some conditions are imposed on its real and imaginary parts.

Theorem 5.14. *(Cf. [21]) Let $T = A + iB$ where A and B are self-adjoint (one of them is also positive). If T^2 is symmetric, then*

$$T \text{ is normal} \iff D(A) = D(B) \text{ and } D(AB) = D(BA).$$

If this is the case, then T^2 is self-adjoint. If we further assume the positiveness of T^2 , then T is self-adjoint and positive.

Before proving this result, we give some auxiliary result whose proof is very simple, thus omitted. It is worth noticing in passing that there are unbounded self-adjoint operators A and B such that $A + iB \subset 0$ (where 0 designates the zero operator on all of H), yet $A \not\subset 0$ and $B \not\subset 0$. For example, let A and B be unbounded self-adjoint operators such that $D(A) \cap D(B) = \{0_H\}$ (as in, e.g., [49]). Assuming $D(A) = D(B)$ makes the whole difference. Indeed:

Proposition 5.15. *Let A and B be two densely defined symmetric operators with domains $D(A), D(B) \subset H$ respectively. Assume that*

$D(A) = D(B)$. If $A + iB \subset 0$, then $A \subset 0$ and $B \subset 0$. If A (or B) is further taken to be closed, then $A = B = 0$ everywhere on H .

Now, we prove Theorem 5.14.

Proof. The implication " \Rightarrow " follows from one of the versions of the spectral theorem for unbounded normal operators.

Now, assume $D(A) = D(B)$ and $D(AB) = D(BA)$. Also, suppose A is positive (the proof in the case of the positiveness of B is similar).

We have

$$A^2 - B^2 + i(AB + BA) \subset (A + iB)A + i(A + iB)B = T^2, \quad ^1$$

thereby

$$A^2 - B^2 - T^2 + i(AB + BA) \subset 0.$$

Since $D(A) = D(B)$, $D(A^2) = D(BA)$ and $D(B^2) = D(AB)$. Hence

$$D(A^2 - B^2) = D(AB + BA).$$

Since $D(AB) = D(BA)$, we have

$$D(A^2 - B^2 - T^2) = D(AB + BA) = D(A^2) = D(B^2).$$

Since A is self-adjoint, A^2 is densely defined. Then both $AB + BA$ and $A^2 - B^2 - T^2$ are densely defined. By the symmetricity of A , B , and T^2 , we have

$$AB + BA \subset (AB + BA)^* \text{ and } A^2 - B^2 - T^2 \subset (A^2 - B^2 - T^2)^*.$$

That is, both $AB + BA$ and $A^2 - B^2 - T^2$ are symmetric. Proposition 5.15 then yields $AB + BA \subset 0$. Hence $AB = -BA$ (for $D(AB) = D(BA)$) and so

$$A^2B = -ABA = BA^2.$$

As A is positive, we obtain $AB = BA$ by [5]. Hence $(A + I)B \subset B(A + I)$. But $D[B(A + I)] = \{x \in D(A) : Ax + x \in D(B)\}$. So, if $x \in D[B(A + I)]$, it follows that $x \in D(A) = D(B)$ and $Ax \in D(B)$, i.e., $x \in D(BA)$. Since $D(AB) = D(BA)$, we have $x \in D(AB) = D[(A + I)B]$. Thus

$$(A + I)B = B(A + I).$$

Since A is self-adjoint and positive, it results that $A + I$ is boundedly invertible. Then $(A + I)^{-1}B \subset B(A + I)^{-1}$. By Proposition 5.27 in [63], this means that A commutes strongly with B . Accordingly T is normal.

Hence, T^2 is normal, and so it is self-adjoint because it is already symmetric.

¹ In fact " \subset " is a full equality due to the conditions on domains, but this observation does not help much for the rest of the proof.

Finally, we show the last statement. Assume that T^2 is positive, and remember that T is normal and T^2 is self-adjoint. Let λ be a complex number in $\sigma(T)$. Then

$$\lambda^2 \in [\sigma(T)]^2 = \sigma(T^2).$$

That is, $\lambda^2 \geq 0$ and so the only possible outcome is $\lambda \in \mathbb{R}$. Therefore, T is self-adjoint. Since in this case

$$0 \leq A = \operatorname{Re} T = \frac{T + T^*}{2} = T,$$

it follows that T is also positive. This marks the end of the proof. \square

Corollary 5.16. (Cf. [56]) *Let $T = A + iB$ where A and B are self-adjoint (one of them is also positive) where $D(A) = D(B)$. If $T^2 = 0$ on $D(T)$, then $T \in B(H)$ is normal and so $T = 0$ everywhere on H .*

Proof. What prevents us a priori from using Theorem 5.14 is that the condition $D(AB) = D(BA)$ is missing. But, writing $A = (T + T^*)/2$ and $B = (T - T^*)/2i$ (and so $D(T) \subset D(T^*)$), we see that if $x \in D(T)$, then

$$Tx + T^*x \in D(T) \iff Tx - T^*x \in D(T)$$

for $Tx \in D(T)$ (because $D(T^2) = D(T)$). In other language, $D(AB) = D(BA)$, as needed. \square

6. ON THE OPERATOR EQUATION $T^*T = p(T)$

In [16], it was shown that if A is densely defined closed operator that satisfies the equation $A^2 = A^*A$, then A is self-adjoint. This result now is a consequence of Theorem 3.1. Recall that such a problem was first mooted in [38] (see also [82] and [40], cf. [57] and [80]).

Corollary 6.1. *Let T be a closed and densely defined operator verifying $T^*T = T^2$. Then T is self-adjoint.*

Proof. Clearly

$$T^*T = T^2 \implies TT^*T = T^3 \implies TT^*T = T^2T \implies TT^*T = T^*TT,$$

from which we derive the quasinormality of T . Since T^2 is self-adjoint, Theorem 3.1 yields $T^2 = T^{*2}$. By Proposition 5.3, T is normal. The way of showing that $\sigma(T) \subset \mathbb{R}$ is as in the proof of Theorem 3.2 in [16]. Thus, T is self-adjoint. \square

Corollary 6.2. *Let T be a densely defined closed operator satisfying $TT^* = T^2$. Then T is self-adjoint.*

Proof. Because T is closed, it is seen that TT^* , or T^2 , is self-adjoint. Theorem 3.1, used in the special case $p(z) = z^2$, then gives $T^2 = (T^2)^* = T^{*2}$. Hence $TT^* = T^{*2}$. Setting $S = T^*$, which is closed and densely defined, the previous equation becomes $S^*S = S^2$. Therefore, S must be self-adjoint by Corollary 6.1. Thus,

$$T = \overline{T} = T^{**} = T^*,$$

as wished. □

Corollary 6.3. *Let T be a densely defined closed operator satisfying $T^*T = T^{*2}$. Then T is self-adjoint.*

Corollary 6.4. *Let T be a densely defined closed operator satisfying $TT^* = T^{*2}$. Then T is self-adjoint.*

We have an additional result about square roots of symmetric operators.

Corollary 6.5. *Let T be a densely defined symmetric operator. Then*

$$T \text{ is essentially self-adjoint} \iff T^{*2} \text{ is symmetric.}$$

Proof. Assume T^{*2} is symmetric. Since T is symmetric, it ensues that $TT^* \subset \overline{TT^*} \subset T^{*2}$. So, T^{*2} is densely defined as $\overline{TT^*}$ is self-adjoint. By the same token, $\overline{TT^*} \subset T^{*2}$ becomes $\overline{TT^*} = T^{*2}$ for self-adjoint operators are maximally symmetric. The preceding equation may be rewritten as $\overline{TT^*} = (\overline{T})^{*2}$. Therefore, \overline{T} is self-adjoint by Corollary 6.4.

The converse, which is plain, is left to readers. □

Remark. It is worth noticing that the equation $T^*T = T^2$ yields, thanks to Theorem 3.1, $T^*T = T^{*2}$. Hence $(T^*T)^2 = T^{*2}T^2 (= T^2T^{*2})$. This equation has been investigated in detail in [33].

What about the more general equation $T^*T = p(T)$? We will deal directly with the case of densely defined closed operators.

Theorem 6.6. *Let T be a densely defined closed operator with domain $D(T) \subset H$, and such that*

$$T^*T = p(T),$$

where $p(z)$ is a polynomial of degree $n \geq 1$. Then the following assertions hold:

- (1) T is quasinormal.
- (2) If $n \neq 2$, then $T \in B(H)$.
- (3) Let $n = 2$. Assume that $p(z)$ has the property that whenever there is (a priori complex) z such that $p(z) \geq 0$, then $z \in \mathbb{R}$. Then T is self-adjoint.

Proof.

- (1) The observation

$$T^*T = p(T) \implies TT^*T = Tp(T) = p(T)T = T^*TT$$

shows the quasinormality of T .

- (2) If $n = 1$, then $D(T^*T) = D(T)$, which gives $T \in B(H)$ because T is closed (Exercise 11 on Page 427 in [20], see also Lemma 2.1 in [65]).

Assume now that $n \geq 3$. Since T is quasinormal and closed, $|T^n| = |T|^n$. Then

$$D(|T|^2) = D(T^*T) = D[p(T)] = D(T^n) = D(|T^n|) = D(|T|^n),$$

and because $|T|$ is self-adjoint and positive, Lemma A.1 in [65] yields $|T| \in B(H)$. Therefore, for a certain $M \geq 0$ and all $x \in H$

$$\|Tx\| = \||T|x\| \leq M\|x\|,$$

that is, $T \in B(H)$.

- (3) First, recall that a closed hyponormal operator with real spectrum is self-adjoint (see the proof of Theorem 8 in [11]). It is also known that quasinormality yields hyponormality. So, we need only show that $\sigma(T) \subset \mathbb{R}$. Let $\lambda \in \sigma(T)$. Then,

$$p(\lambda) \in p[\sigma(T)] = \sigma[p(T)] = \sigma(T^*T) \subset [0, \infty).$$

By the hypothesis on $p(z)$, we immediately obtain $\sigma(T) \subset \mathbb{R}$, as desired. □

Remark. There are certain interpretations of the condition on $p(z)$ above. For instance, it could be replaced by $p(z) = az^2 + bz + c$, with $a > 0$ and $4ac - b^2 \leq 0$, as kindly indicated by Prof. Spiros Konstantogiannis. In the counterexamples section, I give simple examples showing the indispensability of these conditions on the coefficients.

Example 6.7. Let T be a densely defined closed operator such that $T^*T = T^2 - 3T + 2I$. Then T is self-adjoint. We need only check that $\sigma(T) \subset \mathbb{R}$. Let $\lambda := \alpha + i\beta \in \sigma(T)$. Then $\lambda^2 - 3\lambda + 2 \geq 0$. For the inequality to make sense, we must have $\beta = 0$ (while $\alpha^2 - 3\alpha + 2 \geq 0$). In consequence, $\lambda \in \mathbb{R}$, as needed.

Remark. It is clear that $T^*T = p(T)$, where $\deg p(z) = 1$, directly gives the self-adjointness of T when p has real coefficients only. If $p(z)$ has at least one complex coefficient, then T is normal.

Corollary 6.8. *Let T be a densely defined, closed, and symmetric operator. If*

$$T^*T = p(T),$$

where $p(z)$ is a polynomial of degree $n \geq 1$, then T is self-adjoint.

Proof. By Theorem 6.6, T is quasinormal, and we have already recalled that a quasinormal symmetric operator is self-adjoint. \square

Corollary 6.9. *Let T be a densely defined, closed, and symmetric operator. If*

$$TT^* = p(T),$$

where $p(z)$ is a polynomial of degree $n \geq 1$, then T is self-adjoint.

Proof. In this case, we cannot get the quasinormality of T out of $TT^* = p(T)$. Nonetheless, T is self-adjoint by Corollary 3.4. \square

7. ON THE ADJOINT OF TT^* AND T^*T

Let T be densely defined and closable. We will deal with the validity of, e.g., $(TT^*)^* = TT^*$, where TT^* is densely defined. Recall that TT^* is symmetric. If TT^* is further assumed to be closed, then, as is known, either $\sigma(TT^*) = \mathbb{C}$; or $\sigma(TT^*) = \{\lambda \in \mathbb{C} : \text{Im } \lambda \geq 0\}$; or $\sigma(TT^*) = \{\lambda \in \mathbb{C} : \text{Im } \lambda \leq 0\}$; or $\sigma(TT^*) \subset \mathbb{R}$. By assuming below that $\sigma(TT^*) \neq \mathbb{C}$, and so TT^* is closed; we show that the only option left is $\sigma(TT^*) \subset \mathbb{R}$.

Theorem 7.1. *Let T be a densely defined closable operator. If TT^* is densely defined, then*

$$\sigma(TT^*) \neq \mathbb{C} \iff (TT^*)^* = \overline{TT^*} = TT^*.$$

*Similarly, when T^*T is densely defined, then*

$$\sigma(T^*T) \neq \mathbb{C} \iff (T^*T)^* = T^*\overline{T} = T^*T.$$

Proof. Clearly

$$\overline{TT^*} \subset (TT^*)^*.$$

Since $\sigma(TT^*) \neq \mathbb{C}$, consider a complex number λ such that $TT^* - \lambda I$ is (boundedly) invertible. Then

$$(TT^* - \lambda I)^* = (TT^*)^* - \overline{\lambda}I$$

remains invertible.

On the other hand, since $\overline{TT^*}$ is self-adjoint, $\overline{TT^*} - \overline{\lambda}I$ too is (boundedly) invertible. Therefore

$$\overline{TT^*} \subset (TT^*)^* \implies \overline{TT^*} - \overline{\lambda}I \subset (TT^*)^* - \overline{\lambda}I.$$

Since, e.g., $\overline{TT^*} - \overline{\lambda}I$ is surjective and $(TT^*)^* - \overline{\lambda}I$ is injective, Lemma 1.3 in [63] gives $\overline{TT^*} - \overline{\lambda}I = (TT^*)^* - \overline{\lambda}I$ or merely $\overline{TT^*} = (TT^*)^*$.

To show the other equality, we may reason as above by using $TT^* \subset \overline{TT^*}$. Alternatively, here is a different approach: Since $\overline{TT^*}$ is self-adjoint, we have

$$\overline{TT^*} = (\overline{TT^*})^* = (TT^*)^{**} = \overline{\overline{TT^*}} = TT^*$$

where the closedness of TT^* is obtained from $\sigma(TT^*) \neq \mathbb{C}$. This settles the implication " \Rightarrow ".

Conversely, if $(TT^*)^* = \overline{TT^*} = TT^*$, then because TT^* is self-adjoint and positive, $\sigma(TT^*) \subset [0, \infty)$, that is, $\sigma(TT^*) \neq \mathbb{C}$.

The second assertion of the theorem may be shown analogously, thus omitted. \square

Remark. Recall that a condition like $\sigma(A) \neq \mathbb{C}$ does not necessarily give the density of $D(A)$ (e.g., let $A := I_E$ be the identity operator restricted to an unclosed non-dense domain E).

Remark. Despite the fact the previous proof is short and elementary, there is yet another proof of the implication " \Rightarrow ", which was communicated to me by a reader, who prefers to remain anonymous. It reads: By assumptions, $S := TT^*$ is a densely defined, and symmetric operator with non-empty resolvent set, so S is closed. Since S is positive as well, it has equal deficiency indices which, together with $\sigma(S) \neq \mathbb{C}$, imply $\sigma(S) \subset \mathbb{R}$. Hence S is self-adjoint.

The following consequence is presented without proof.

Corollary 7.2. *Let T be a densely defined closable operator. If TT^* is densely defined, then*

$$\sigma(TT^*) \neq \mathbb{C} \iff \sigma(TT^*) \subset [0, \infty).$$

*Similarly, when T^*T is densely defined, then*

$$\sigma(T^*T) \neq \mathbb{C} \iff \sigma(T^*T) \subset [0, \infty).$$

By scrutinizing the previous proof, it turns out that we may show a stronger result, given that we have not required the closedness of the spectrum in this paper.

Proposition 7.3. *Let T be a densely defined closable operator. If TT^* is densely defined and $(TT^*)^*$ does not possess at least one complex eigenvalue, then*

$$(TT^*)^* = \overline{TT^*} = TT^*.$$

In particular, TT^ is self-adjoint.*

Remark. Mutatis mutandis, a similar result holds for T^*T .

Proof. It suffices to let λ be a complex number such that $(TT^*)^* - \bar{\lambda}I$ is injective, and the rest of the proof is as that of Theorem 7.1. \square

Remark. When $\sigma(TT^*) \neq \mathbb{C}$ and $\sigma(T^*T) \neq \mathbb{C}$ simultaneously, then both TT^* and T^*T are densely defined, and besides

$$(TT^*)^* = TT^* \text{ and } (T^*T)^* = T^*T.$$

This may be consulted in [29] and [30].

A. Devinatz, A. E. Nussbaum, and J. von Neumann obtained in [19] the following maximality result (cf. [42]):

Theorem 7.4. *Let T , T_1 and T_2 be self-adjoint operators. Then*

$$T \subset T_1T_2 \implies T = T_1T_2.$$

A somewhat similar idea as above may be reused to reach an akin conclusion, namely:

Proposition 7.5. *Let T , T_1 and T_2 be unbounded operators. If T is self-adjoint and $T \subset T_1T_2$, then*

$$\sigma(T_1T_2) \neq \mathbb{C} \iff T = T_1T_2.$$

Proof. Pick a complex λ such that $T_1T_2 - \lambda I$ is boundedly invertible. By the self-adjointness of T , $T - \lambda I$ is always boundedly invertible. So

$$T \subset T_1T_2 \implies T - \lambda I \subset T_1T_2 - \lambda I.$$

Hence $(T - \lambda I)^{-1} = (T_1T_2 - \lambda I)^{-1}$, as they are both everywhere defined. Thus, $T - \lambda I = T_1T_2 - \lambda I$, or merely $T = T_1T_2$. This shows the implication " \implies ". The backward implication is evident, and the proof is complete. \square

8. COUNTEREXAMPLES

- (1) (Cf. Theorem 6.6) The equation $T^*T = p(T)$, if $\deg p(z) = 2$ (and without any other condition on $p(z)$), does not always yield the self-adjointness of T , even when $\dim H < \infty$. Remember that in a such context, quasinormality coincide with normality, so the matrix must first be normal.

Example 8.1. On a bi-dimensional space, consider

$$T = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

If $p(z) = z^2 + 2$, then it is seen that

$$T^*T = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = T^2 + 2I.$$

Observe in the end that the complex number i , which is an eigenvalue for T , is such that $p(i) \geq 0$.

The next example shows that even if $p(z)$ has two real roots, we still require the positivity of the leading coefficient. Indeed:

Example 8.2. Let T be a non-zero skew-adjoint matrix, i.e. $T^* = -T$ and $T^* \neq T$. Then $T^*T = -T^2$, yet T is not self-adjoint.

- (2) Corollary 3.2 in [81] was extremely helpful to establish some of the results in this paper. Recall that it states that a quasinormal operator T is normal if (and only if) $\ker T = \ker T^*$.

It would therefore be practical to have a result like it for other classes of operators. The answer is, alas, negative. We first give an example borrowed from Answer 21.2.5 in [49], where readers can find more details about it.

Example 8.3. Consider $Tf(x) = xf(x)$ on

$$D(T) = \left\{ f \in L^2(\mathbb{R}) : xf \in L^2(\mathbb{R}), \int_{\mathbb{R}} f(x)dx = 0 \right\}.$$

Then T is densely defined, closed, and symmetric. However, T is not self-adjoint, i.e., $D(T) \neq D(T^*)$. It is also seen that

$$\ker T = \ker T^* = \{0\}.$$

In other words, the condition $\ker T = \ker T^*$ ($= \{0\}$) combined with the symmetricity and the closedness of T do not suffice to force T to be self-adjoint. In fact, T is not even normal.

Now, a densely defined symmetric operator is formally normal. Also, a symmetric operator always has a self-adjoint extension, possibly in a larger Hilbert space (see, e.g., Proposition 3.17 in [63]). Put differently, a symmetric operator is subnormal, and it is patently hyponormal. That being said, and due to the above example, a formally normal or subnormal operator T with the condition $\ker T = \ker T^*$ ($= \{0\}$) need not be normal.

- (3) As alluded to above, below we give an unbounded self-adjoint positive operator whose square roots have pairwise different domains.

Example 8.4. Let T be an unbounded, self-adjoint, and positive operator with domain $D(T) \subsetneq H$, then define $S = \begin{pmatrix} T & 0 \\ 0 & T \end{pmatrix}$, where $D(S) = D(T) \oplus D(T)$. It is plain that S too is unbounded, self-adjoint, and positive. Letting \sqrt{T} represent the unique self-adjoint positive square root of T , readers may readily check that each of

$$A = \begin{pmatrix} 0 & T \\ I & 0 \end{pmatrix}, B = \begin{pmatrix} 0 & I \\ T & 0 \end{pmatrix}, \text{ and } C = \begin{pmatrix} \sqrt{T} & 0 \\ 0 & \sqrt{T} \end{pmatrix}$$

is in effect a square root of S . Observe in the end that $D(A) = H \oplus D(T)$, $D(B) = D(T) \oplus H$, and $D(C) = D(\sqrt{T}) \oplus D(\sqrt{T})$. In other words, these three domains are pairwise different.

- (4) It is natural to ask whether the identity $(TT^*)^* = TT^*$, which holds for densely defined closed operators T , still holds for closable operators. The following example answers this question in the negative.

Example 8.5. Let A be a densely defined positive operator in a Hilbert space H such that it is not essentially self-adjoint. Assume further that $\text{ran} A$ is dense in H . Define an inner product space on $\text{ran} A$ by

$$\langle Ax, Ay \rangle_A = \langle Ax, y \rangle, \quad x, y \in D(A).$$

Denote the completion of this pre-Hilbert space by H_A . Define the canonical embedding operator $T : H_A \supseteq \text{ran} A \rightarrow H$ by

$$T(Ax) := Ax, \quad x \in D(A).$$

It may then be shown that $D(A) \subset D(T^*)$ and

$$T^*x = Ax \in H_A, \quad x \in D(A).$$

In particular, $T : H_A \supset \text{ran} A \rightarrow H$ is a densely defined and closable linear operator. Besides, $T^{**}T^*$ is a self-adjoint positive extension of A .

On the other hand, notice that

$$\ker T^* = (\text{ran} T)^\perp = (\text{ran} A)^\perp = \{0\}.$$

In other words, T^* is one-to-one. So, $T^*y \in \text{ran} A (=D(T))$, where $y \in D(T^*)$, implies $y \in D(A)$. Accordingly,

$$D(TT^*) = \{y \in D(T^*) : T^*y \in D(T)\} = D(A).$$

Thus, $TT^* = A$. This signifies that

$$(TT^*) = A^* \neq T^{**}T^* = \overline{T}T^*$$

for the latter operator is self-adjoint, whilst the former is not.

- (5) The preceding example may be beefed up to obtain a stronger counterexample, namely:

Example 8.6. Let T be a closable densely defined operator such that $(TT^*)^* \neq \overline{T}T^*$ (as just before), then set

$$S = \begin{pmatrix} 0 & T \\ T^* & 0 \end{pmatrix}$$

with $D(S) = D(T^*) \oplus D(T)$. So S is densely defined, and since

$$S^* = \begin{pmatrix} 0 & \overline{T} \\ T^* & 0 \end{pmatrix}, S \text{ is symmetric.}$$

Now,

$$SS^* = \begin{pmatrix} TT^* & 0 \\ 0 & T^*\overline{T} \end{pmatrix} \text{ and } \overline{S}S^* = \begin{pmatrix} \overline{T}T^* & 0 \\ 0 & T^*\overline{T} \end{pmatrix}.$$

But

$$(SS^*)^* = \begin{pmatrix} (TT^*)^* & 0 \\ 0 & (T^*\overline{T})^* \end{pmatrix} = \begin{pmatrix} (TT^*)^* & 0 \\ 0 & T^*\overline{T} \end{pmatrix}$$

because $T^*\overline{T}$ is self-adjoint. Since we already know that $(TT^*)^* \neq \overline{T}T^*$, it ensues that $(SS^*)^* \neq \overline{S}S^*$ and yet S is densely defined and symmetric.

- (6) We have been assuming that TT^* is densely defined as, in general, $D(TT^*)$ could be non-dense. For instance, in [48] (or [49]), we have found an example of a densely defined T that obeys

$$D(T^2) = D(T^*) = D(TT^*) = D(T^*T) = \{0\}.$$

Obviously such an operator T cannot be closable. So, does the closability of T suffice to make TT^* densely defined? The answer is negative even when T is symmetric. This is seen next:

Example 8.7. Let E be a dense linear proper subspace of H , let u be a non-zero element in H but not in E , and define T to be projection on the 1-dimensional subspace spanned by u with $D(T) = E$. Then T is a bounded, non-everywhere defined, unclosed, and symmetric operator. Also, T^* is the same projection, defined on the entire H . Then T^*x is in E only if it is 0. So,

$$D(TT^*) = \{u\}^\perp,$$

which is not dense.

Remark. In the previous example, one sees that $D(T^*T)$ is dense. This is, however, not peculiar to this example. Indeed, if T is a densely defined, symmetric and bounded (non-everywhere defined) operator with domain $D(T) \subsetneq H$, then $D(T^*T)$ is always dense. This is easy to see, as $T \subset T_H$ where T_H is the extension of T to all of H . Then $T_H^* \subset T^*$, and so $D(T^*) = H$. Therefore,

$$D(T^*T) = D(T),$$

which is dense.

- (7) Last but not least, we supply a densely defined (unclosed) symmetric operator T such that T^*T is not densely defined. The same idea could have been used to construct an akin counterexample in the case of TT^* .

Example 8.8. Let A and B be self-adjoint operators, where $B \in B[L^2(\mathbb{R})]$, yet to be chosen. Then, define

$$T = \begin{pmatrix} 0 & AB \\ BA & 0 \end{pmatrix},$$

with $D(T) = D(A) \oplus D(AB)$. A priori, BA need not be closed, and we will choose BA as such. We also take AB to be densely defined. Then T will be densely defined, and symmetric for

$$T^* = \begin{pmatrix} 0 & AB \\ (AB)^* & 0 \end{pmatrix} \supset \begin{pmatrix} 0 & AB \\ BA & 0 \end{pmatrix} = T.$$

Moreover,

$$T^*T = \begin{pmatrix} AB^2A & 0 \\ 0 & (AB)^*AB \end{pmatrix}.$$

We are done as soon as $D(AB^2A)$ is not dense. Now, we give a pair of A and B that meet the above conditions. Some details about the remaining portion of this example may have to be consulted in Answer 21.2.24 in [49] (and also [37]).

Consider the self-adjoint operator S

$$Sf(x) = e^{\frac{x^2}{2}}f(x),$$

defined on $D(S) = \{f \in L^2(\mathbb{R}) : e^{\frac{x^2}{2}}f \in L^2(\mathbb{R})\}$. Then, set $A := \mathcal{F}^*S\mathcal{F}$, where \mathcal{F} is the usual $L^2(\mathbb{R})$ -Fourier transform. So, A is also self-adjoint.

Clearly S is boundedly invertible (hence so is A). Set

$$B^2 f(x) := S^{-1} f(x) = e^{\frac{-x^2}{2}} f(x),$$

which is defined from $L^2(\mathbb{R})$ onto $D(S)$.

We know that $D(AB^2)$ is trivial if $D(A) \cap \text{ran}(B^2) = \{0\}$, and B^2 is injective (which is the case). But,

$$D(A) \cap \text{ran}(B^2) = D(A) \cap D(S) = \{0\}$$

as this is already available to us from Answer 21.2.17 in [49] (or [37]). So, $D(AB^2) = \{0\}$. Thus,

$$D(AB^2 A) = \{f \in D(A) : Af \in D(AB^2) = \{0\}\} = \ker A = \{0\},$$

i.e., $D(AB^2 A)$ is not dense in $L^2(\mathbb{R})$. It only remains to check that $D(AB)$ is indeed dense in $L^2(\mathbb{R})$. Observe that by uniqueness of the positive square root that

$$Bf(x) = e^{\frac{-x^2}{4}} f(x),$$

with $D(B) = L^2(\mathbb{R})$. Now

$$D(AB) = \{f \in L^2(\mathbb{R}) : \widehat{e^{\frac{-x^2}{4}} f} \in L^2(\mathbb{R}), e^{\frac{x^2}{2}} \widehat{e^{\frac{-x^2}{4}} f} \in L^2(\mathbb{R})\},$$

which is dense by, say, an idea borrowed from the proof of Corollary 9 in [37].

Remark. It seems that there is no magic result that guarantees the density of $D(TT^*)$ or $D(T^*T)$, when T is symmetric, except perhaps to assume that T^2 is densely defined.

ACKNOWLEDGEMENT

The author wishes to thank Professor Zsigmond Tarcsay for Example 8.5, which was based on a construction by Z. Sebestyén and J. Stochel (see [64]).

REFERENCES

- [1] T. Ando. Operators with a norm condition, *Acta Sci. Math. (Szeged)*, **33** (1972) 169-178.
- [2] T. Ya. Azizov, M. Denisov, F. Philipp. Spectral functions of products of selfadjoint operators, *Math. Nachr.*, **285/14-15** (2012) 1711-1728.
- [3] T. Ya. Azizov, A. Dijksma. Closedness and adjoints of products of operators, and compressions, *Integral Equations Operator Theory*, **74/2** (2012) 259-269.
- [4] M. Barraa, M. Boumazghour. Numerical range submultiplicity, *Linear Multilinear Algebra*, **63/11** (2015), 2311-2317.
- [5] S. J. Bernau. The square root of a positive self-adjoint operator, *J. Austral. Math. Soc.*, **8** (1968) 17-36.

- [6] I. Boucif, S. Dehimi and M. H. Mortad. On the absolute value of unbounded operators, *J. Operator Theory*, **82/2** (2019) 285-306.
- [7] J. A. W. van Casteren, S. Goldberg. The conjugate of the product of operators, *Studia Math.*, **38** (1970) 125-130.
- [8] M. Chō, R. E. Curto, T. Huruya. n -tuples of operators satisfying $\sigma_T(AB) = \sigma_T(BA)$. Special issue dedicated to Professor T. Ando., *Linear Algebra Appl.*, **341** (2002) 291-298.
- [9] A. Daniluk. On the closability of paranormal operators, *J. Math. Anal. Appl.*, **376/1** (2011) 342-348.
- [10] R. Dautray, J.L. Lions. Mathematical analysis and numerical methods for science and technology. Vol. **2**. Functional and variational methods. With the collaboration of Michel Artola, Marc Authier, Philippe Bénilan, Michel Cessenat, Jean Michel Combes, Hélène Lanchon, Bertrand Mercier, Claude Wild and Claude Zuily. Translated from the French by Ian N. Sneddon. *Springer-Verlag, Berlin*, 1988.
- [11] S. Dehimi, M. H. Mortad. Bounded and unbounded operators similar to their adjoints, *Bull. Korean Math. Soc.*, **54/1** (2017) 215-223.
- [12] S. Dehimi, M. H. Mortad. Right (or left) invertibility of bounded and unbounded operators and applications to the spectrum of products, *Complex Anal. Oper. Theory*, **12/3** (2018) 589-597.
- [13] S. Dehimi, M. H. Mortad. Chernoff-like counterexamples related to unbounded operators, *Kyushu J. Math.*, **74/1** (2020) 105-108.
- [14] S. Dehimi, M. H. Mortad. Unbounded operators having self-adjoint, subnormal or hyponormal powers, *Math. Nachr.*, (to appear). DOI: 10.1002/mana.202100390
- [15] S. Dehimi, M. H. Mortad, A. Bachir. On the commutativity of closed symmetric operators (submitted). arXiv:2203.07266
- [16] S. Dehimi, M. H. Mortad, Z. Tarcsay. On the operator equations $A^n = A^*A$, *Linear Multilinear Algebra*, **69/9** (2021) 1771-1778.
- [17] P. A. Deift. Applications of a commutation formula, *Duke Math. J.*, **45/2** (1978) 267-310.
- [18] A. Devinatz, A. E. Nussbaum. On the permutability of normal operators, *Ann. of Math. (2)*, **65** (1957) 144-152.
- [19] A. Devinatz, A. E. Nussbaum, J. von Neumann. On the permutability of self-adjoint operators, *Ann. of Math. (2)*, **62** (1955) 199-203.
- [20] J. Dieudonné. *Treatise on analysis. Vol. II*. Enlarged and corrected printing. Translated by I. G. Macdonald. With a loose erratum. Pure and Applied Mathematics, 10-II. Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1976.
- [21] N. Frid, M. H. Mortad, S. Dehimi. When nilpotence implies the zeroness of linear operators, *Khayyam J. Math.*, (to appear).
- [22] F. Gesztesy, J. A. Goldstein, H. Holden, G. Teschl. Abstract wave equations and associated Dirac-type operators, *Ann. Mat. Pura Appl. (4)*, **191/4** (2012) 631-676.
- [23] F. Gesztesy, K. Schmüdgen. On a theorem of Z. Sebestyén and Zs. Tarcsay, *Acta Sci. Math. (Szeged)*, **85/1-2** (2019) 291-293.
- [24] H. A. Gindler. Classroom Notes: A Spectral Mapping Theorem for Polynomials, *Amer. Math. Monthly*, **72/5** (1965) 528-530.

- [25] M. González. Null spaces and ranges of polynomials of operators, *Publ. Mat.*, **32/2** (1988) 167-170.
- [26] K. Gustafson. On projections of self-adjoint operators and operator product adjoints, *Bull. Amer. Math. Soc.*, **75** (1969) 739-741.
- [27] K. Gustafson, M. H. Mortad. Unbounded products of operators and connections to Dirac-type operators, *Bull. Sci. Math.*, **138/5** (2014), 626-642.
- [28] K. Gustafson, M. H. Mortad. Conditions implying commutativity of unbounded self-adjoint operators and related topics, *J. Operator Theory*, **76/1**, (2016) 159-169.
- [29] V. Hardt, A. Konstantinov, R. Mennicken. On the spectrum of the product of closed operators, *Math. Nachr.*, **215**, (2000) 91-102.
- [30] V. Hardt, R. Mennicken. On the spectrum of unbounded off-diagonal 2×2 operator matrices in Banach spaces. Recent advances in operator theory (Groningen, 1998), 243-266, *Oper. Theory Adv. Appl.*, **124**, Birkhäuser, Basel, 2001.
- [31] P. Hess, T. Kato. Perturbation of Closed Operators and Their Adjoints, *Comment. Math. Helv.*, **45** (1970) 524-529.
- [32] M. Hladnik, M. Omladič. Spectrum of the product of operators, *Proc. Amer. Math. Soc.*, **102/2**, (1988) 300-302.
- [33] Z. J. Jabłoński, Il B. Jung, J. Stochel. Unbounded quasinormal operators revisited, *Integral Equations Operator Theory*, **79/1** (2014) 135-149.
- [34] Il B. Jung, M. H. Mortad, J. Stochel. On normal products of selfadjoint operators, *Kyungpook Math. J.*, **57** (2017) 457-471.
- [35] T. Kato. Perturbation theory for linear operators, *Springer*, 1980 (2nd edition).
- [36] W. E. Kaufman. Closed operators and pure contractions in Hilbert space, *Proc. Amer. Math. Soc.*, **87/1** (1983) 83-87.
- [37] H. Kosaki. On intersections of domains of unbounded positive operators, *Kyushu J. Math.*, **60/1** (2006) 3-25.
- [38] K. R. Laberteux. Problem 10377, *Amer. Math. Monthly*, **101** (1994) 362.
- [39] K. B. Laursen, M. M. Neumann. An introduction to local spectral theory. London Mathematical Society Monographs. New Series, **20**. *The Clarendon Press, Oxford University Press, New York*, 2000.
- [40] S. A. McCullough, L. Rodman. Hereditary classes of operators and matrices, *Amer. Math. Monthly*, **104/5** (1997) 415-430.
- [41] A. McIntosh. Operators which have an H_∞ functional calculus. *Miniconference on operator theory and partial differential equations* (North Ryde, 1986), 210-231, Proc. Centre Math. Anal. Austral. Nat. Univ., 14, Austral. Nat. Univ., Canberra, 1986.
- [42] M. Mezzanese, M. H. Mortad. Maximality of linear operators, *Rend. Circ. Mat. Palermo, Ser II.*, **68/3** (2019) 441-451.
- [43] M. H. Mortad. An application of the Putnam-Fuglede theorem to normal products of self-adjoint operators, *Proc. Amer. Math. Soc.*, **131/10**, (2003) 3135-3141.
- [44] M. H. Mortad. On some product of two unbounded self-adjoint operators, *Integral Equations Operator Theory*, **64/3** (2009) 399-408.
- [45] M. H. Mortad. On the adjoint and the closure of the sum of two unbounded operators, *Canad. Math. Bull.*, **54/3** (2011) 498-505.
- [46] M. H. Mortad. Commutativity of unbounded normal and self-adjoint operators and applications, *Oper. Matrices*, **8/2** (2014) 563-571.

- [47] M. H. Mortad. *An operator theory problem book*, World Scientific Publishing Co., (2018).
- [48] M. H. Mortad. On the triviality of domains of powers and adjoints of closed operators, *Acta Sci. Math. (Szeged)*, **85** (2019) 651-658.
- [49] M. H. Mortad. *Counterexamples in operator theory*, Birkhäuser/Springer, Cham (2022).
- [50] M. H. Mortad. Counterexamples related to unbounded paranormal operators, *Examples and Counterexamples* (to appear). <https://doi.org/10.1016/j.exco.2021.100017>
- [51] E. Nelson. Analytic vectors, *Ann. of Math. (2)*, **70** (1959) 572-615.
- [52] S. Ôta. Some classes of q -deformed operators, *J. Operator Theory*, **48/1** (2002) 151-186.
- [53] S. Ôta, K. Schmüdgen. Some selfadjoint 2×2 operator matrices associated with closed operators, *Integral Equations Operator Theory*, **45/4** (2003) 475-484.
- [54] F. Philipp, A. C. M. Ran, M. Wojtylak. Local definitizability of $T^{[*]}T$ and $TT^{[*]}$, *Integral Equations Operator Theory*, **71/4** (2011) 491-508.
- [55] P. Pietrzycki, J. Stochel. On n th roots of bounded and unbounded quasinormal operators. arXiv:2103.09961v3
- [56] C. R. Putnam. On square roots of normal operators, *Proc. Amer. Math. Soc.*, **8** (1957) 768-769.
- [57] M. Roman, A. Sandovici. Multivalued linear operator equation $A^*A = \lambda A^n$, *Complex Anal. Oper. Theory*, **15/1** (2021) Paper No. 6, 14 pp.
- [58] M. Reed, B. Simon. *Methods of modern mathematical physics, Vol. 1: Functional analysis*, Academic Press. 1972.
- [59] M. Reed, B. Simon. *Methods of modern mathematical physics, Vol. 2: Fourier analysis, self-adjointness*, Academic Press. 1975.
- [60] K. Schmüdgen. On domains of powers of closed symmetric operators, *J. Operator Theory*, **9/1** (1983) 53-75.
- [61] K. Schmüdgen. On commuting unbounded selfadjoint operators. I, *Acta Sci. Math. (Szeged)*, **47/1-2** (1984) 131-146.
- [62] K. Schmüdgen. *Unbounded operator algebras and representation theory*. Operator Theory: Advances and Applications, **37**. Birkhäuser Verlag, Basel, 1990.
- [63] K. Schmüdgen. *Unbounded self-adjoint operators on Hilbert space*, Springer. GTM **265** (2012).
- [64] Z. Sebestyén, J. Stochel. Restrictions of positive selfadjoint operators, *Acta Sci. Math. (Szeged)*, **55/1-2** (1991) 149-154.
- [65] Z. Sebestyén, J. Stochel. On suboperators with codimension one domains, *J. Math. Anal. Appl.*, **360/2** (2009) 391-397.
- [66] Z. Sebestyén, Zs. Tarcsay. T^*T always has a positive selfadjoint extension, *Acta Math. Hungar.*, **135/1-2** (2012) 116-129.
- [67] Z. Sebestyén, Zs. Tarcsay. Characterizations of selfadjoint operators, *Studia Sci. Math. Hungar.*, **50/4** (2013) 423-435.
- [68] Z. Sebestyén, Zs. Tarcsay. A reversed von Neumann theorem, *Acta Sci. Math. (Szeged)*, **80/3-4** (2014) 659-664.
- [69] Z. Sebestyén, Zs. Tarcsay. Operators having selfadjoint squares, *Ann. Univ. Sci. Budapest. Eötvös Sect. Math.*, **58** (2015) 105-110.
- [70] Z. Sebestyén, Zs. Tarcsay. Adjoint of sums and products of operators in Hilbert spaces, *Acta Sci. Math. (Szeged)*, **82/1-2** (2016) 175-191.

- [71] Z. Sebestyén, Zs. Tarcsay. On the adjoint of Hilbert space operators, *Linear Multilinear Algebra*, **67/3** (2019) 625-645.
- [72] J. Stochel. Lifting strong commutants of unbounded subnormal operators, *Integral Equations Operator Theory*, **43/2** (2002) 189-214.
- [73] J. Stochel, F. H. Szafraniec. On normal extensions of unbounded operators. II, *Acta Sci. Math. (Szeged)*, **53/1-2** (1989) 153-177.
- [74] J. Stochel, F. H. Szafraniec. C^∞ -vectors and boundedness. Volume dedicated to the memory of Włodzimierz Mlak, *Ann. Polon. Math.*, **66** (1997) 223-238.
- [75] J. Stochel, F. H. Szafraniec. Domination of unbounded operators and commutativity, *J. Math. Soc. Japan*, **55/2** (2003) 405-437.
- [76] F. H. Szafraniec. Kato-Protter type inequalities, bounded vectors and the exponential function, *Ann. Polon. Math.*, **51** (1990) 303-312.
- [77] F. H. Szafraniec. Normals, subnormals and an open question, *Oper. Matrices*, **4/4** (2010) 485-510.
- [78] A. E. Taylor. *Introduction to functional analysis*, John Wiley & Sons, Inc., New York; Chapman & Hall, Ltd., London 1958.
- [79] B. Thaller. *The Dirac equation*. Texts and Monographs in Physics. Springer-Verlag, Berlin, 1992.
- [80] Y. Tian. A note on two-sided removal and cancellation properties associated with Hermitian matrix, *preprint* (2021).
- [81] M. Uchiyama. Operators which have commutative polar decompositions. Contributions to operator theory and its applications, 197-208, *Oper. Theory Adv. Appl.*, **62**, Birkhäuser, Basel, 1993.
- [82] B-Y. Wang, F. Zhang. Words and normality of matrices, *Linear and Multilinear Algebra*, **40/2** (1995) 111-118.
- [83] J. Weidmann. *Linear Operators in Hilbert Spaces*, Springer, 1980.
- [84] T. Yamazaki, M. Yanagida. Relations between two operator inequalities and their applications to paranormal operators, *Acta Sci. Math. (Szeged)*, **69/1-2** (2003) 377-389.

LABORATORY OF MATHEMATICAL ANALYSIS AND APPLICATIONS. DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ORAN 1, AHMED BEN BELLA, B.P. 1524, EL MENOVAR, ORAN 31000, ALGERIA.

Email address: mhmortad@gmail.com, mortad.hichem@univ-oran1.dz.