

Detecting coherence with respect to general quantum measurements

Yu-Cheng Chen,¹ Jiong Cheng,¹ Wen-Zhao Zhang,^{1,*} and Cheng-Jie Zhang^{1,2,†}

¹*School of Physical Science and Technology, Ningbo University, Ningbo, 315211, China*

²*State Key Laboratory of Precision Spectroscopy, School of Physics and Electronic Science, East China Normal University, Shanghai 200241, China*

(Dated: August 11, 2022)

Quantum coherence is a crucial resource in quantum resource theory. Previous study mainly focused on standard coherence under a complete orthogonal reference basis. The standard coherence has recently been extended to general positive-operator-valued measure (POVM)-based coherence, including block coherence as a special case. Therefore, it is necessary to construct block coherence and POVM-based coherence witnesses to detect them. In this work, we present witnesses for block coherence and POVM-based coherence, and obtain the necessary and sufficient conditions for arbitrary block coherence and POVM-based coherence witnesses. We also discuss possible realizations of some block coherence and POVM-based coherence witnesses in experiments, and present examples of measuring block coherence witnesses based on real experimental data. Furthermore, an application of block coherence witnesses has been presented in a quantum parameter estimation task with a degenerate Hamiltonian, and one can estimate the unknown parameter by measuring our block coherence witnesses if the input state is block coherent. Last but not least, we prove that the quantum Fisher information of any block incoherent state is equal to zero, which coincides with the result from measuring block coherence witnesses.

I. INTRODUCTION

Quantum coherence plays a significant role in quantum theory, which has a number of applications in quantum optics, quantum information processing, nanoscale thermodynamics and biological systems [1–15]. Recently, it has been recognized that coherence can be treated as one kind of quantum resources. Therefore, detecting coherence is crucial in quantum physics.

In Ref. [16], quantitative investigations of quantum coherence have been launched, and several coherence measures for standard coherence, with respect to von Neumann measurements, have been proposed. For standard coherence, consider a d -dimensional Hilbert space \mathcal{H} , a state δ is incoherent under a chosen reference basis $\{|i\rangle\}_{i=1}^d$ if and only if δ is diagonal under the reference basis [16], i.e.,

$$\delta = \sum_{i=1}^d p_i |i\rangle\langle i|, \quad (1)$$

with probabilities $\{p_i\}$. One can define the standard dephasing operation Δ as

$$\Delta(\rho) := \sum_{i=1}^d |i\rangle\langle i| \rho |i\rangle\langle i|. \quad (2)$$

Thus, a state δ is incoherent under a chosen reference basis $\{|i\rangle\}_{i=1}^d$ if and only if the following condition holds,

$$\delta = \Delta(\delta), \quad (3)$$

where Δ is defined in Eq. (2).

In Ref. [17], Åberg proposed a framework which actually defined block coherence with respect to projective measurements. The standard coherence can be viewed as a special case of block coherence. Moreover, Bischof *et al.* generalized block coherence to coherence with respect to the most general quantum measurements [18], i.e., the positive-operator-valued measure (POVM)-based coherence. Therefore, the resource theory of coherence was generalized by extending the standard coherence to block coherence and even POVM-based coherence. It is worth noticing that POVMs describe the most general type of quantum measurements, and they may be more advantageous compared to von Neumann measurements.

However, unlike the standard coherence, there are only a few results reported on quantifying block coherence and POVM-based coherence [18–20]. The methods to detect whether a state has nonzero block coherence and POVM-based coherence are missing. For the standard coherence, Ref. [21] first introduced the standard coherence witness W . Similar to entanglement witnesses, a Hermitian operator W is a standard coherence witness if $\text{Tr}(\delta W) \geq 0$ holds for all incoherent states δ . If one finds $\text{Tr}(\rho W) < 0$ for a state ρ , then the state ρ must be a standard coherent state. Compared with coherence measures, which usually need full information of the state by using quantum state tomography, coherence witnesses can be measured with much less measurements since quantum state tomography requires exponentially growing measurements with the number of qubits. Thus, it is necessary to construct block coherence witnesses and POVM-based coherence witnesses to detect them without quantum state tomography, especially for experimentally unknown states.

The purpose of this work is two-fold. On the one

*Electronic address: zhangwenzhao@nbu.edu.cn

†Electronic address: chengjie.zhang@gmail.com

hand, we present witnesses for block coherence and POVM-based coherence, and obtain the necessary and sufficient conditions for arbitrary block coherence witnesses and POVM-based coherence witnesses. Moreover, we discuss possible realizations of some block coherence and POVM-based coherence witnesses in real experiments, and present examples to detect block coherence coherence by measuring block coherence witnesses. On the other hand, we provide an application of block coherence witnesses in a quantum parameter estimation task with a degenerate Hamiltonian, and one can estimate the unknown parameter by measuring our block coherence witnesses if the input state is block coherent. We also prove that the quantum Fisher information of any block incoherent state is equal to zero, which coincides with the result from measuring block coherence witnesses.

II. DETECTING BLOCK COHERENCE BASED ON BLOCK COHERENCE WITNESSES

Before embarking on our main results, let us first review the definition of block incoherent states. In Refs. [17–20, 22, 23], block incoherent state has been defined as follows.

Given a d -dimensional Hilbert space \mathcal{H} , which has been divided into n ($n \leq d$) subspaces, and subspace projectors are $\mathbf{P} := \{P_s\}_{s=1}^n$ with $\sum_{s=1}^n P_s = \mathbb{1}$ (where $\mathbb{1}$ is the identity operator). A state $\tilde{\delta}$ is block incoherent under the reference subspace projectors \mathbf{P} , if and only if $\tilde{\delta}$ is block diagonal under the reference \mathbf{P} , i.e.,

$$\tilde{\delta} = \sum_{s=1}^n P_s \tilde{\delta} P_s := \tilde{\Delta}(\tilde{\delta}), \quad (4)$$

where we define the modified dephasing operation as

$$\tilde{\Delta}(\rho) := \sum_s P_s \rho P_s. \quad (5)$$

Similar to the witnesses for standard coherence [21, 24], we can construct block coherence witnesses as follows.

Theorem 1. (a) For any Hermitian operator A , we can construct a block coherence witness

$$\tilde{W}_A = \tilde{\Delta}(A) - A. \quad (6)$$

(b) An arbitrary Hermitian operator \tilde{W} is a block coherence witness if and only if $\tilde{\Delta}(\tilde{W}) \geq 0$.

Proof.— (a) We first prove that \tilde{W}_A is a block coherence witness. Since A is a Hermitian operator, \tilde{W}_A must be Hermitian as well. Thus, for an arbitrary block inco-

herent state $\tilde{\delta} = \sum_s P_s \tilde{\delta} P_s$, we can obtain

$$\begin{aligned} \text{Tr}(\tilde{\delta} \tilde{W}_A) &= \text{Tr}[\tilde{\delta} \tilde{\Delta}(A)] - \text{Tr}[\tilde{\delta} A] \\ &= \text{Tr}[\tilde{\delta} \sum_s P_s A P_s] - \text{Tr}[\tilde{\delta} A] \\ &= \text{Tr}[\sum_s P_s \tilde{\delta} P_s A] - \text{Tr}[\tilde{\delta} A] \\ &= 0, \end{aligned} \quad (7)$$

which means \tilde{W}_A is a block coherence witness.

(b) It is worth noticing that a Hermitian operator W is a coherence witness for standard coherence if and only if $\Delta(W) \geq 0$ [21]. Similarly, we can prove that a Hermitian operator \tilde{W} is a block coherence witness if and only if $\tilde{\Delta}(\tilde{W}) \geq 0$.

Firstly, if $\tilde{\Delta}(\tilde{W}) \geq 0$ holds, then for any block incoherent state $\tilde{\delta}$ we can obtain that

$$\begin{aligned} \text{Tr}[\tilde{\delta} \tilde{W}] &= \text{Tr}[\tilde{\Delta}(\tilde{\delta}) \tilde{W}] \\ &= \text{Tr}[\tilde{\Delta}(\tilde{W}) \tilde{\delta}] \\ &\geq 0, \end{aligned} \quad (8)$$

i.e., \tilde{W} is a block coherence witness.

Conversely, we prove that if for any block incoherent state $\tilde{\delta}$ $\text{Tr}[\tilde{\delta} \tilde{W}] \geq 0$ holds, then $\tilde{\Delta}(\tilde{W}) \geq 0$. For any quantum state ρ , one can obtain

$$\begin{aligned} \text{Tr}[\rho \tilde{\Delta}(\tilde{W})] &= \text{Tr}[\tilde{\Delta}(\rho) \tilde{W}] \\ &= \text{Tr}[\tilde{\delta}_\rho \tilde{W}] \\ &\geq 0, \end{aligned} \quad (9)$$

where $\tilde{\delta}_\rho := \tilde{\Delta}(\rho)$ is a block incoherent state. Thus, $\tilde{\Delta}(\tilde{W})$ is positive-semidefinite, i.e., $\tilde{\Delta}(\tilde{W}) \geq 0$.

Therefore, it can be given that \tilde{W} is block coherence witness if and only if $\tilde{\Delta}(\tilde{W}) \geq 0$. \square

Remark.— Based on Theorem 1(a), one can construct a block coherence witness \tilde{W}_σ by using any density matrix σ as the Hermitian operator A in Eq. (6),

$$\tilde{W}_\sigma = \tilde{\Delta}(\sigma) - \sigma. \quad (10)$$

Moreover, if σ is a pure state $|\phi\rangle$, we can obtain that

$$\tilde{W}_\phi = \tilde{\Delta}(|\phi\rangle\langle\phi|) - |\phi\rangle\langle\phi|, \quad (11)$$

and

$$\begin{aligned} \text{Tr}[\rho \tilde{W}_\phi] &= \text{Tr}[\rho(\tilde{\Delta}(|\phi\rangle\langle\phi|) - |\phi\rangle\langle\phi|)] \\ &= \langle\phi|\tilde{\Delta}(\rho)|\phi\rangle - \langle\phi|\rho|\phi\rangle \\ &= F(\tilde{\Delta}(\rho), |\phi\rangle) - F(\rho, |\phi\rangle), \end{aligned} \quad (12)$$

where $F(\rho, |\phi\rangle) := \langle\phi|\rho|\phi\rangle$ is the fidelity between the state ρ and the pure state $|\phi\rangle$. Therefore, the expect value of block coherence witness \tilde{W}_ϕ is related to the two fidelities.

III. COHERENCE WITNESS WITH RESPECT TO GENERAL MEASUREMENTS

Recently, Bishof *et al.* introduced a generalization to a resource theory of coherence with respect to the general quantum measurement. The POVM-based coherence is defined as follows [18–20].

Let \mathbf{E} be a collection of n positive operators $\mathbf{E} := \{E_i\}_{i=1}^n$ with $\sum_{i=1}^n E_i = \mathbb{1}$. The corresponding measurement operator of each E_i is denoted by A_i , such that $E_i = A_i^\dagger A_i$ holds. Thus, a state $\bar{\delta}$ is called incoherent state with respect to the general measurement \mathbf{E} if and only if

$$E_i \bar{\delta} E_{i'} = 0, \quad \forall i \neq i'. \quad (13)$$

Note that it is equivalent to [18]

$$A_i \bar{\delta} A_{i'}^\dagger = 0, \quad \forall i \neq i'. \quad (14)$$

Therefore, any POVM-based incoherent state $\bar{\delta}$ should satisfy

$$\bar{\delta} = \sum_i E_i \bar{\delta} E_i := \bar{\Delta}(\bar{\delta}), \quad (15)$$

where $\bar{\Delta}$ is defined as

$$\bar{\Delta}(\rho) := \sum_i E_i \rho E_i. \quad (16)$$

It is worth noticing that Eq. (15) can be easily proved from the definition of POVM-based incoherent state, since for any POVM-based incoherent state $\bar{\delta}$, one can obtain

$$\begin{aligned} \bar{\delta} &= \left(\sum_i E_i \right) \bar{\delta} \left(\sum_j E_j \right) \\ &= \sum_i E_i \bar{\delta} E_i + \sum_{i \neq j} E_i \bar{\delta} E_j \\ &= \sum_i E_i \bar{\delta} E_i, \end{aligned} \quad (17)$$

where we have used $E_i \bar{\delta} E_j = 0, \forall i \neq j$.

Theorem 2. (a) For any Hermitian operator A , we can construct a POVM-based coherence witness \bar{W}_A as follows,

$$\bar{W}_A = \bar{\Delta}(A) - A. \quad (18)$$

(b) An arbitrary Hermitian operator \bar{W} is a POVM-based coherence witness if and only if $\bar{\Delta}(\bar{W}) \geq 0$.

Proof.— (a) We provide that \bar{W}_A is a POVM-based coherence witness. As A is a Hermitian operator, \bar{W}_A must be Hermitian. For any incoherent state $\bar{\delta}$ with respect to $\{E_i\}$, $\bar{\delta} = \sum_i E_i \bar{\delta} E_i$ holds and thus

$$\begin{aligned} \text{Tr}(\bar{\delta} \bar{W}_A) &= \text{Tr}[\bar{\delta} \bar{\Delta}(A)] - \text{Tr}[\bar{\delta} A] \\ &= \text{Tr}[\bar{\delta} \sum_i E_i A E_i] - \text{Tr}[\bar{\delta} A] \\ &= \text{Tr}[\sum_i E_i \bar{\delta} E_i A] - \text{Tr}[\bar{\delta} A] \\ &= 0, \end{aligned} \quad (19)$$

which means \bar{W}_A is a POVM-based coherence witness.

(b) Firstly, if $\bar{\Delta}(\bar{W}) \geq 0$ holds, then for any POVM-based incoherent state $\bar{\delta}$ we can obtain that

$$\begin{aligned} \text{Tr}[\bar{\delta} \bar{W}] &= \text{Tr}[\bar{\Delta}(\bar{\delta}) \bar{W}] \\ &= \text{Tr}[\bar{\Delta}(\bar{W}) \bar{\delta}] \\ &\geq 0. \end{aligned} \quad (20)$$

Thus, \bar{W} is a POVM-based coherence witness.

Conversely, we prove that if $\text{Tr}[\bar{\delta} \bar{W}] \geq 0$, then $\bar{\Delta}(\bar{W}) \geq 0$. For any quantum state ρ , one can obtain that

$$\begin{aligned} \text{Tr}[\rho \bar{\Delta}(\bar{W})] &= \text{Tr}[\bar{\Delta}(\rho) \bar{W}] \\ &= \text{Tr}[\bar{\delta}_\rho \bar{W}] \\ &\geq 0, \end{aligned} \quad (21)$$

where $\bar{\delta}_\rho := \bar{\Delta}(\rho)$ is a POVM-based incoherent state. Thus, $\bar{\Delta}(\bar{W})$ is positive-semidefinite, i.e., $\bar{\Delta}(\bar{W}) \geq 0$.

Therefore, we proved that \bar{W} is a POVM-based coherence witness if and only if $\bar{\Delta}(\bar{W}) \geq 0$. \square

Remark.— Based on Theorem 2(a), we can also construct a POVM-based coherence witness \bar{W}_σ by choosing any density matrix σ as the Hermitian operator A ,

$$\bar{W}_\sigma = \bar{\Delta}(\sigma) - \sigma. \quad (22)$$

Moreover, if σ is a pure state $|\phi\rangle$, one can obtain that

$$\bar{W}_\phi = \bar{\Delta}(|\phi\rangle\langle\phi|) - |\phi\rangle\langle\phi|, \quad (23)$$

and

$$\begin{aligned} \text{Tr}[\rho \bar{W}_\phi] &= \text{Tr}[\rho (\bar{\Delta}(|\phi\rangle\langle\phi|) - |\phi\rangle\langle\phi|)] \\ &= \text{Tr}[\rho \bar{\Delta}(|\phi\rangle\langle\phi|)] - \text{Tr}[\rho |\phi\rangle\langle\phi|] \\ &= F(\bar{\Delta}(\rho), |\phi\rangle) - F(\rho, |\phi\rangle), \end{aligned} \quad (24)$$

where it demonstrates the relationship of the POVM-based coherence witness \bar{W}_ϕ and fidelities between the state ρ (or $\bar{\Delta}(\rho)$) and the pure state $|\phi\rangle$.

IV. POSSIBLE EXPERIMENTAL REALIZATION FOR WITNESSES AND EXAMPLES

Many experiments have measured fidelities $F = \langle\phi|\rho_{exp}|\phi\rangle$ between the experimental state ρ_{exp} and a target pure state $|\phi\rangle$ [25–28]. For bipartite and multipartite systems, one possible way to measure fidelities is to decompose the operator $|\phi\rangle\langle\phi|$ as sum of tensor products of local observables [26–28]. Therefore, one can measure our witnesses (11) and (23) in the same manner.

In the following, we will present examples of N -qubit W states $|W_N\rangle$ from real experimental data, where

$$|W_N\rangle = (|0 \cdots 001\rangle + |0 \cdots 010\rangle + \cdots + |10 \cdots 0\rangle) / \sqrt{N}. \quad (25)$$

We use the block coherence witness (11) to detect block coherence of W states. In Ref. [25], N -qubit W states

$ W_N\rangle$	4	5	6	7	8
F_N	0.846	0.759	0.788	0.763	0.722
$\langle \tilde{\Delta}(W_N\rangle\langle W_N) \rangle$	0.321	0.207	0.173	0.141	0.119
$-\text{Tr}(\rho'_{exp}\tilde{W})$	0.525	0.552	0.615	0.622	0.603

TABLE I: The fidelity F_N is the result of overlap $\langle W_N|\rho'_{exp}|W_N\rangle$ from Ref. [25]. The values $\langle \tilde{\Delta}(|W_N\rangle\langle W_N|) \rangle$ and $-\text{Tr}(\rho'_{exp}\tilde{W})$ are obtained from the density matrices of experiment states in Ref. [25]. One can see that $-\text{Tr}(\rho'_{exp}\tilde{W})$ is always greater than zero.

($4 \leq N \leq 8$) have been experimentally generated by trapped ions, with fidelities between experimental states and perfect W states being $F_4 = 0.846$, $F_5 = 0.759$, $F_6 = 0.788$, $F_7 = 0.763$, $F_8 = 0.722$ for the 4-, 5-, 6-, 7- and 8-ion W states, respectively. Moreover, numerical values of the density matrices of experiment states with $4 \leq N \leq 8$ have been presented in Ref. [25]. It is worth noticing that the experimental states have local phases, and one can find the local phases by maximizing the fidelities $F = \langle \tilde{W}_N|\rho_{exp}|\tilde{W}_N\rangle$ with $|\tilde{W}_N\rangle$ being W states containing local phases shown in [25]. After choosing local unitary transformations based on local phases, we can transform ρ_{exp} to ρ'_{exp} such that $F = \langle \tilde{W}_N|\rho_{exp}|\tilde{W}_N\rangle = \langle W_N|\rho'_{exp}|W_N\rangle$. We consider the following reference subspace projectors \mathbf{P} with

$$\begin{aligned}
P_0 &= |\phi^-\rangle\langle\phi^-|, \\
P_1 &= |\phi^+\rangle\langle\phi^+|, \\
P_2 &= |0\dots 010\rangle\langle 0\dots 010|, \\
&\dots, \\
P_{N-1} &= |010\dots 0\rangle\langle 010\dots 0|, \\
P_N &= \mathbb{1} - \sum_{i=0}^{N-1} P_i,
\end{aligned}$$

where $|\phi^\pm\rangle := (|00\dots 01\rangle \pm |10\dots 0\rangle)/\sqrt{2}$. Thus, our block coherence witness is as follows,

$$\begin{aligned}
\tilde{W} &= \tilde{\Delta}(|W_N\rangle\langle W_N|) - |W_N\rangle\langle W_N| \\
&= \sum_{i=0}^N P_i|W_N\rangle\langle W_N|P_i - |W_N\rangle\langle W_N|. \quad (26)
\end{aligned}$$

From Table I, we can see that $-\text{Tr}(\rho'_{exp}\tilde{W})$ is always greater than zero, which means all the experimental states in Ref. [25] contain the block coherence under the above reference subspace projectors \mathbf{P} .

V. QUANTUM PARAMETER ESTIMATION TASK WITH DEGENERATE HAMILTONIANS

For quantum metrology, one of the main tasks is parameter estimation in a quantum channel to improve accuracy that is different from the standard quantum limits [29, 30]. Quantum coherence plays a fundamental

role in quantum parameter estimation [31]. For the case of unitary evolution with a degenerate Hamiltonian, we will propose a simple application of quantum block coherence, and find that the quantum Fisher information is strongly related to the block coherence.

A. Quantum parameter estimation by using block coherent states

Now we consider a d -dimensional Hilbert space, suppose that H is a degenerate Hamiltonian,

$$H = \sum_{s=1}^n \sum_{g=1}^{k_s} E_s |s, g\rangle\langle s, g|, \quad (27)$$

where H has n different eigenvalues $\{E_s\}_{s=1}^n$, and each eigenvalue E_s has k_s degenerate eigenstates $\{|s, g\rangle\}_{g=1}^{k_s}$. Here the index s is for the s -th different eigenvalue, and the index g is denoted as the g -th E_s (the total number of the same eigenvalue E_s is k_s). $|s, g\rangle$ is the eigenstate corresponding to the eigenvalue, i.e., the g -th E_s . According to the definition of block coherence, one can naturally choose the degenerate subspaces of H in Eq. (27) as the reference subspaces. Therefore,

$$P_s = \sum_{g=1}^{k_s} |s, g\rangle\langle s, g|, \quad (28)$$

is the s -th subspace projector, where every pure state in this subspace is an eigenstate of H with the eigenvalue being E_s .

Proposition 3. With a degenerate Hamiltonian H in Eq. (27), we choose $\{P_s\}_{s=1}^n$ in (28) as the reference subspace projectors \mathbf{P} . The output state $\rho_{out} = U_\varphi \rho_{in} U_\varphi^\dagger := \rho_\varphi$ can be used to estimate the unknown parameter φ in the black box in Fig. 1, if and only if ρ_{in} and ρ_{out} have nonzero block coherence under the reference subspaces \mathbf{P} .

Proof.— For an arbitrary input state ρ_{in} , it can be expressed under the eigenstates of H (27), i.e.,

$$\rho_{in} = \sum_{s,s'} \sum_{g,g'} \rho_{(s,g),(s',g')} |s, g\rangle\langle s', g'|, \quad (29)$$

where we used $\rho_{(s,g),(s',g')} := \langle s, g|\rho_{in}|s', g'\rangle$ with $\sum_{s,g} \rho_{(s,g),(s,g)} = 1$. Thus, the corresponding output state $\rho_{out} = U_\varphi \rho_{in} U_\varphi^\dagger := \rho_\varphi$ can be expressed as follows,

$$\begin{aligned}
\rho_{out} &= U_\varphi \rho_{in} U_\varphi^\dagger \\
&= \sum_{\substack{s,s',g,g' \\ s \neq s'}} \rho_{(s,g),(s',g')} e^{-i(E_s - E_{s'})\varphi} |s, g\rangle\langle s', g'| \\
&\quad + \sum_s \sum_{g,g'} \rho_{(s,g),(s,g')} |s, g\rangle\langle s, g'|. \quad (30)
\end{aligned}$$




FIG. 1: The black box implements an unitary evolution $U_\varphi = e^{-iH\varphi}$ on the input state ρ_{in} , and the unknown parameter φ of the black box must be estimated by measuring $\rho_{out} = U_\varphi \rho_{in} U_\varphi^\dagger := \rho_\varphi$. Suppose that the Hamiltonian H is degenerate and already known, the quantum parameter estimation task is estimating the unknown parameter φ from the output state ρ_{out} .

We can see that ρ_{out} is dependent on φ if and only if there exists nonzero $\rho_{(s,g),(s',g')}$ with $s \neq s'$, i.e., ρ_{out} and ρ_{in} have nonzero block coherence under the reference subspaces. \square

Remark.— It is worth noticing that witnesses for standard coherence may be useless to estimate the unknown parameter when the Hamiltonian H is degenerate, although the input state has standard coherence. Consider a special case, if we use an input state which is block incoherent, but its density matrix contains off-diagonal nonzero elements in some degenerate subspaces under a chosen eigenvectors of H as reference basis, i.e., this input state contains standard coherence but no block coherence. The expect values of standard coherence witnesses has no information of the parameter, and one cannot estimate it by measuring witnesses of standard coherence, although the input state contains standard coherence.

B. Quantum Fisher information of block incoherent states

For an arbitrary output state ρ_φ , the quantum Fisher information F_q can be obtained [32–35]

$$\begin{aligned} F_q &= \text{Tr}[\rho_\varphi L_\varphi^2] \\ &= \sum_{m,n} 4c_m \left(\frac{c_n - c_m}{c_n + c_m} \right)^2 |\langle m|H|n\rangle|^2, \end{aligned} \quad (31)$$

where $U_\varphi = e^{-iH\varphi}$ is a unitary operator and H is the corresponding Hermitian Hamiltonian, and the output state $\rho_\varphi = U_\varphi \rho_{in} U_\varphi^\dagger = U_\varphi (\sum_n c_n |n\rangle \langle n|) U_\varphi^\dagger$ with $\rho_{in} = \sum_n c_n |n\rangle \langle n|$. $\{c_n\}$ and $\{|n\rangle\}$ are the eigenvalues and eigenvectors of ρ_{in} , respectively. The symmetric logarithmic derivative operator is $L_\varphi = U_\varphi (-2i \sum_{m,n} \frac{\langle m|[H,\rho_{in}]|n\rangle}{c_n + c_m} |m\rangle \langle n|) U_\varphi^\dagger$ [32–35]. Furthermore, we will discuss the quantum Fisher information with a block incoherent state.

Proposition 4. Consider a degenerate Hamiltonian H in Eq. (27) and the quantum parameter estimation task

in Fig. 1, one can choose the degenerate subspaces of H as the reference subspaces. If the input state ρ_{in} is a block incoherent state, then the quantum Fisher information for the output state is $F_q = 0$.

Proof.— Consider an arbitrary block incoherent state as the input state ρ_{in} ,

$$\begin{aligned} \rho_{in} &= \sum_i \sum_{g,g'} \rho_{(i,g),(i,g')} |i,g\rangle \langle i,g'| \\ &= \sum_i \sum_{\tilde{g}} A_{\tilde{g}}^{(i)} |i,\tilde{g}\rangle \langle i,\tilde{g}|, \end{aligned} \quad (32)$$

where the second equation holds since we diagonalize ρ_{in} in each subspace. Thus, the eigenvalues of ρ_{in} are $\{A_{\tilde{g}}^{(i)}\}$ with corresponding eigenvectors $\{|i,\tilde{g}\rangle\}$. The output state ρ_φ can be expressed as

$$\begin{aligned} \rho_\varphi &= U_\varphi \rho_{in} U_\varphi^\dagger \\ &= U_\varphi \sum_i \sum_{\tilde{g}} A_{\tilde{g}}^{(i)} |i,\tilde{g}\rangle \langle i,\tilde{g}| U_\varphi^\dagger, \end{aligned} \quad (33)$$

where $U_\varphi = e^{-iH\varphi}$ is a unitary operator. Furthermore, the symmetric logarithmic derivative operator L_φ can be expressed as [32–35],

$$\begin{aligned} L_\varphi &= U_\varphi \left(-2i \sum_{i,j} \sum_{\tilde{g},\tilde{h}} \frac{\langle i,\tilde{g}|[H,\rho_{in}]|j,\tilde{h}\rangle}{A_{\tilde{g}}^{(i)} + A_{\tilde{h}}^{(j)}} |i,\tilde{g}\rangle \langle j,\tilde{h}| \right) U_\varphi^\dagger \\ &= U_\varphi \left(-2i \sum_{i,j} \sum_{\tilde{g},\tilde{h}} \frac{A_{\tilde{h}}^{(j)} - A_{\tilde{g}}^{(i)}}{A_{\tilde{g}}^{(i)} + A_{\tilde{h}}^{(j)}} \langle i,\tilde{g}|H|j,\tilde{h}\rangle |i,\tilde{g}\rangle \langle j,\tilde{h}| \right) U_\varphi^\dagger. \end{aligned}$$

Finally, we obtain the quantum Fisher information F_q as

$$\begin{aligned} F_q &= \text{Tr}(\rho_\varphi L_\varphi^2) \\ &= 4 \sum_{i,j} \sum_{\tilde{g},\tilde{h}} A_{\tilde{g}}^{(i)} |\langle i,\tilde{g}|H|j,\tilde{h}\rangle|^2 \left(\frac{A_{\tilde{h}}^{(j)} - A_{\tilde{g}}^{(i)}}{A_{\tilde{g}}^{(i)} + A_{\tilde{h}}^{(j)}} \right)^2 \\ &= 4 \sum_{i,j} \sum_{\tilde{g},\tilde{h}} A_{\tilde{g}}^{(i)} |\langle i,\tilde{g}|H|j,\tilde{h}\rangle|^2 \left(\frac{A_{\tilde{h}}^{(j)} - A_{\tilde{g}}^{(i)}}{A_{\tilde{g}}^{(i)} + A_{\tilde{h}}^{(j)}} \right)^2 \\ &\quad + 4 \sum_i \sum_{\tilde{g},\tilde{h}} A_{\tilde{g}}^{(i)} |\langle i,\tilde{g}|H|i,\tilde{h}\rangle|^2 \left(\frac{A_{\tilde{h}}^{(i)} - A_{\tilde{g}}^{(i)}}{A_{\tilde{g}}^{(i)} + A_{\tilde{h}}^{(i)}} \right)^2 \\ &= 0, \end{aligned} \quad (34)$$

where we have used $\langle i,\tilde{g}|H|j,\tilde{h}\rangle = 0$ with $i \neq j$, and $\langle i,\tilde{g}|H|i,\tilde{h}\rangle = E_i \delta_{\tilde{g}\tilde{h}}$. \square

Remark.— When the input state is a block incoherent state, the quantum Fisher information of the output state is equal to zero, which means one cannot estimate the parameter from the output state, i.e., the output state contains no information of the parameter. That coincides with the result from Proposition 3.

VI. DISCUSSIONS AND CONCLUSIONS

Besides the Fisher information estimation mentioned in the above example, when performing quantum coherence operations on any degenerate quantum system, we also need to take into account the measure of coherence. In the practical systems, degeneracy is a very common situation and has been widely reported in the microscopic field, e.g., orbital degenerate states and degenerate spin states in λ -type and the chainlike structures atomic system [36], near-degenerate states in few-electron ions system [37], etc. Moreover, degeneracy also exists in macroscopic systems, e.g., macroscopically degenerate in intrinsic quasi-crystals [38], and been regarded to be appeared during the conversion or transition of black holes [39]. In addition, degeneracy plays an important role in both quantum computing and quantum networks construction [40, 41]. It is often used as a protected qubit in fault-tolerant quantum operation [40]. And it can also be used as a control device to control the coherence between quantum systems [42]. In the discussion of topological structure and phase transition, the study of degeneracy and near-degeneracy also plays an important role [43–45]. These studies related

to the issue of how to correctly measure the coherence of degenerate quantum systems. Therefore, our coherence detection scheme has potential value in the measurement and application of degenerate states.

In conclusion, we have discussed witnesses for block coherence and POVM-based coherence. The necessary and sufficient conditions for arbitrary block coherence and POVM-based coherence witnesses have been obtained. Furthermore, we have shown that block coherent states can be used in a quantum parameter estimation task with a degenerate Hamiltonian if the input state is block coherent. The quantum Fisher information of block incoherent states is equal to zero, which coincides with the result from the block coherence witness.

ACKNOWLEDGMENTS

This work is supported by the National Natural Science Foundation of China (Grants No. 11734015, No. 11704205, and No. 12074206), the open funding program from State Key Laboratory of Precision Spectroscopy (East China Normal University), and K.C. Wong Magna Fund in Ningbo University.

-
- [1] M. Piani, M. Cianciaruso, T. R. Bromley, C. Napoli, N. Johnston, and G. Adesso, Robustness of asymmetry and coherence of quantum states, *Phys. Rev. A* **93**, 042107 (2016).
 - [2] S. F. Huelga and M.B. Plenio, Vibrations quanta and biology, *Contemp. Phys.* **54**, 181 (2013).
 - [3] F. G. S. L. Brandão and M. B. Plenio, Entanglement theory and the second law of thermodynamics, *Nat. Phys.* **4**, 873 (2008).
 - [4] F. G. S. L. Brandão and M. B. Plenio, A Reversible Theory of Entanglement and its Relation to the Second Law, *Commun. Math. Phys.* **295**, 829 (2010).
 - [5] M. Horodecki and J. Oppenheim, (Quantumness in the context of) Resource Theories, *Int. J. Mod. Phys. B* **27**, 1345019 (2013).
 - [6] Y. Dai, Y. Dong, Z. Xu, W. You, C. Zhang, and O. Gühne, Experimentally Accessible Lower Bounds for Genuine Multipartite Entanglement and Coherence Measures, *Phys. Rev. Applied* **13**, 054022 (2020).
 - [7] M. B. Plenio and S. S. Virmani, An Introduction to Entanglement Theory, *Quantum. Inf. Coher.* 173–209 (2014).
 - [8] D. J. Zhang, C. L. Liu, X. D. Yu, and D. M. Tong, Estimating Coherence Measures from Limited Experimental Data Available, *Phys. Rev. Lett.* **120**, 170501 (2018).
 - [9] K. D. Wu, Z. Hou, Y. Y. Zhao, G. Y. Xiang, C. F. Li, G. C. Guo, J. Ma, Q. Y. He, J. Thompson, and M. Gu, Experimental Cyclic Interconversion between Coherence and Quantum Correlations, *Phys. Rev. Lett.* **121**, 050401 (2018).
 - [10] C. Zhang, T. R. Bromley, Y. F. Huang, H. Cao, W. M. Lv, B. H. Liu, C. F. Li, G. C. Guo, M. Cianciaruso, and G. Adesso, Demonstrating Quantum Coherence and Metrology that is Resilient to Transversal Noise, *Phys. Rev. Lett.* **123**, 180504 (2019).
 - [11] D. Buono, G. Nocerino, G. Petrillo, G. Torre, G. Zonzo, and F. Illuminati, Quantum coherence of Gaussian states, arXiv:1609.00913 (2016).
 - [12] T. R. Bromley, M. Cianciaruso, and G. Adesso, Frozen Quantum Coherence, *Phys. Rev. Lett.* **114**, 210401 (2015).
 - [13] C. M. Li, L. Neill, Y. N. Chen, G. Y. Chen, and N. Franco, Witnessing quantum coherence: from solid-state to biological systems, *Sci. Rep.* **2**, 885 (2012).
 - [14] K. von Prillwitz, L. Rudnicki, and F. Mintert, Contrast in multipath interference and quantum coherence, *Phys. Rev. A* **92**, 052114 (2015).
 - [15] M. Lostaglio, K. Korzekwa, D. Jennings, and T. Rudolph, Quantum Coherence, Time-Translation Symmetry and Thermodynamics, *Phys. Rev. X* **5**, 021001 (2015).
 - [16] T. Baumgratz, M. Cramer, and M. B. Plenio, Quantifying coherence, *Phys. Rev. Lett.* **113**, 140401 (2014).
 - [17] J. Åberg, Quantifying Superposition, arXiv:quant-ph/0612146 (2006).
 - [18] F. Bischof, H. Kampermann, and D. Bruß, Resource theory of coherence based on positive-operator-valued measures, *Phys. Rev. Lett.* **123**, 110402 (2019).
 - [19] J. W. Xu, L. H. Shao, and S. M. Fei, Coherence measures with respect to general quantum measurements, *Phys. Rev. A* **102**, 012411 (2020).
 - [20] F. Bischof, H. Kampermann, and D. Bruß, Quantifying coherence with respect to general quantum measurements, *Phys. Rev. A* **103**, 032429 (2021).
 - [21] C. Napoli, T. R. Bromley, M. Cianciaruso, M. Piani, N. Johnston, and G. Adesso, Robustness of coherence: an operational and observable measure of quantum coherence, *Phys. Rev. Lett.* **116**, 150502 (2016).
 - [22] S. H. Kim, C. H. Xiong, A. Kuma, and J. D. Wu, Convert-

- ing coherence based on positive-operator-valued measures into entanglement, *Phys. Rev. A* **103**, 052418 (2021).
- [23] S. Kim, C. Xiong, A. Kumar, G. Zhang, and J. Wu, Quantifying dynamical coherence with coherence measures, *Phys. Rev. A* **104**, 012404 (2021).
- [24] Z. Ma, Y. Zhou, Y. Dai, Y. D. Dong, and C. J. Zhang, Detecting and estimating coherence based on coherence witnesses, *Phys. Rev. A* **103**, 012409 (2021).
- [25] H. Häffner, *et al.*, Scalable multiparticle entanglement of trapped ions, *Nature* **438**, 643 (2005).
- [26] X.-L. Wang, L.-K. Chen, W. Li, H.-L. Huang, C. Liu, C. Chen, Y.-H. Luo, Z.-E. Su, D. Wu, Z.-D. Li, H. Lu, Y. Hu, X. Jiang, C.-Z. Peng, L. Li, N.-L. Liu, Y.-A. Chen, C.-Y. Lu, and J.-W. Pan, Experimental ten-photon entanglement, *Phys. Rev. Lett.* **117**, 210502 (2016).
- [27] X.-C. Yao, T.-X. Wang, P. Xu, H. Lu, G.-S Pan, X.-H. Bao, C.-Z. Peng, C.-Y. Lu, Y.-A. Chen, J.-W. Pan, Observation of eight-photon entanglement, *Nature Photon.* **6**, 225-228 (2012).
- [28] Y.-F. Huang, B.-H. Liu, L. Peng, Y.-H. Li, L. Li, C.-F. Li and G.-C. Guo, Experimental generation of an eight-photon Greenberger-Horne-Zeilinger state, *Nature Commun.* **2**, 546-552 (2011).
- [29] V. Giovannetti, S. Lloyd, and L. Maccone, Quantum metrology, *Phys. Rev. Lett.* **96**, 010401 (2006).
- [30] J. P. Dowling, Quantum Optical Metrology—The Lowdown on High-N00N States, *Contemp Phys.* **49**, 125 (2008).
- [31] G. Tóth and I. Apellaniz, Quantum metrology from a quantum information science perspective, *J. Phys. A Math. Theor.* **47**, 424006 (2014).
- [32] S. L. Braunstein and C. M. Caves, Statistical distance and the geometry of quantum states, *Phys. Rev. Lett.* **72**, 3439 (1994).
- [33] S. L. Braunstein, C. M. Caves, and G. J. Milburn, Generalized uncertainty relations: theory, examples, and Lorentz invariance, *Ann. Phys.* **247**, 135 (1996).
- [34] M. G. A. PARIS, Quantum estimation for quantum technology, *Int. J. Quantum. Inf.* **7**, 125 (2009).
- [35] V. Giovannetti, S. Lloyd, and L. Maccone, Advance in quantum metrology, *Nat. Photonics.* **5**, 222–229 (2011).
- [36] V. D. B. Jeroen and K. Daniel, Double Exchange via Degenerate Orbitals, *Phys. Rev. Lett.* **82**, 1016 (1999).
- [37] M. Togawa, *et al.*, Observation of strong two electron one photon transitions in few electron ions, *Phys. Rev. A* **102**, 052831 (2020).
- [38] H. Ha and B.-J. Yang, Macroscopically degenerate localized zero-energy states of quasicrystalline bilayer systems in the strong coupling limit, *Phys. Rev. B* **104**, 165112 (2021).
- [39] J. Yang and H. Huang, Trapping horizons of the evolving charged wormhole and black bounce, *Phys. Rev. D* **104**, 084005 (2021).
- [40] A. R. Klots and L. B. Ioffe, Set of holonomic and protected gates on topological qubits for a realistic quantum computer, *Phys. Rev. B* **104**, 144502 (2021).
- [41] M. Schneider, J. Ostmeier, K. Jansen, T. Luu, and C. Urbach, Simulating both parity sectors of the Hubbard model with tensor networks, *Phys. Rev. B* **104**, 155118 (2021).
- [42] D. Christian, *et al.*, Collective-Mode Enhanced Matter-Wave Optics, *Phys. Rev. Lett.* **127**, 100401 (2021).
- [43] X.-X. Zhang, D. Manske, and N. Nagaosa, Ultrafast excitation and topological soliton formation in incommensurate charge density wave states, *Phys. Rev. B* **104**, 125132 (2021).
- [44] Z. Ge, S. Slizovskiy, F. Joucken, E. A. Quezada, T. Taniguchi, K. Watanabe, V. I. Fal'ko, and J. V. Jr., Control of Giant Topological Magnetic Moment and Valley Splitting in Trilayer Graphene, *Phys. Rev. Lett.* **127**, 136402 (2021).
- [45] A. L. Corps and A. Relano, Constant of Motion Identifying Excited-State Quantum Phases, *Phys. Rev. Lett.* **127**, 130602 (2021).