

On interactions of massless spin 3 and scalar fields

P.M. Lavrov^{(a,b)1}

^(a) *Center of Theoretical Physics,
Tomsk State Pedagogical University,
Kievskaya St. 60, 634061 Tomsk, Russia*
^(b) *National Research Tomsk State University,
Lenin Av. 36, 634050 Tomsk, Russia*

Abstract

Using new approach for the deformation procedure in the case of reducible gauge theories (Lavrov in Eur. Phys. J. C 82:429, 2022), it is shown that in the model of massless spin 3 fields and a real scalar field local cubic vertices invariant under initial gauge transformations do not exist while local quartic gauge-invariant vertices can be constructed.

Keywords: BV-formalism, deformation procedure, anticanonical transformations

PACS numbers: 11.10.Ef, 11.15.Bt

¹E-mail: lavrov@tspu.edu.ru

1 Introduction

The Batalin-Vilkovisky (BV) formalism [1, 2] being the most powerful covariant quantization method plays the very important role in solving the deformation problem which is considered as a way in construction of suitable interactions among fields. The reason is related with embedding the problem into solutions to the classical master-equation of the BV-formalism. Standard approach [3, 4] operates infinite number of equations appeared in expansion of the solution with respect to the deformation parameter. Then the system of these equations is analyzed in each order with respect to the deformation parameter using cohomological methods. Quite recently, the new method to find solutions of the deformation procedure within the BV-formalism has been proposed [5, 6, 7].² The method is based on the invariance of the classical master-equation under anticanonical transformations acting in minimal antisymplectic space of the BV-formalism. The anticanonical transformations (for recent developments see [12, 13]) by itself present an effective tool in studying principal properties of general gauge theories. So, the gauge fixing procedure and the gauge dependence of Green functions can be described in terms of special anticanonical transformations observed firstly in [14]. Moreover, the anticanonical transformations admit to describe the renormalization procedure of general gauge theories [14] that, in its turn, allowed to introduce new understanding the renormalization problem beyond the usual index arguments [15]. Here, we meet with a new face of anticanonical transformations in the solution of the deformation problem, which make it possible to represent solutions in an explicit and closed form.

In the present paper, we apply the new method [5, 6, 7] to construct suitable interactions between massless spin 3 and scalar fields as the result of deformation of initial free action. The initial action is sum of the Fronsdal action for massless spin 3 fields [16] and the action for a free real scalar field. The choice of initial model allows to use the BV-formalism and the method [7] directly because the model belongs to the class of reducible gauge theories. In the case of massless integer spin $s > 3$ fields, the direct application of the BV-formalism is impossible due to the traceless conditions on fields. Fortunately, the situation is not so hopeless because there exist unconstrained formulations for higher spin fields [17, 18, 19] which make possible application of the BV-formalism and new method. In the case under consideration we study cubic and quartic vertices containing one spin 3 field and two or three scalar fields using special anticanonical transformations acting non-trivially in the sector of spin 3 fields only. It is shown that the gauge algebra does not deform under proposed transformations. It has the following consequence: a) the cubic vertices appearing as a local part of the deformed action disappears under requirement of the gauge invariance, b) the local quartic vertices invariant under the gauge transformations do exist.

The paper is organized as follows. In section 2, a short presentation of new approach to the deformation procedure in the BV-formalism for reducible gauge theories is given. In section 3, the Fronsdal action for spin 3 fields as a gauge action of the first-stage reducible theory is discussed. In section 4, deformations of free model of massless spin 3 and scalar fields responsible for local cubic and quartic vertices are studied. In section 5, we summarize the results.

We use the DeWitt's condensed notations [20] and employ the symbols $\varepsilon(A)$ for the Grassmann parity and $\text{gh}(A)$ for the ghost number respectively. The right and left functional derivatives are marked by special symbols " \leftarrow " and " \rightarrow " respectively. Arguments of any functional are enclosed in square brackets $[]$, and arguments of any function are enclosed in parentheses, $()$. The partial right derivative of a function $F(A)$ with respect to field A^i is denoted as $F_{,i}(A)$.

²Analog of this approach in canonical formalism for dynamical systems with constraints within the Batalin-Fradkin-Vilkovisky formalism [8, 9, 10] is known as well [11].

2 Deformations in BV-formalism

In this section, we consider in short main statements of the new approach [5, 6, 7] in solving the deformation problem within the BV-formalism. Starting point of the BV-formalism is a gauge theory of the fields $A = \{A^i\}$ with Grassmann parities $\varepsilon(A^i) = \varepsilon_i$ and ghost numbers $\text{gh}(A^i) = 0$. The theory is described by an initial action $S_0[A]$. It is assumed that the action is invariant under the gauge transformations³

$$S_{0,i}[A]R_\alpha^i(A) = 0, \quad \delta A^i = R_\alpha^i(A)\xi^\alpha, \quad \alpha = 1, 2, \dots, m \quad (1)$$

where $R_\alpha^i(A)$ ($\varepsilon(R_\alpha^i(A)) = \varepsilon_i + \varepsilon_\alpha$, $\text{gh}(R_\alpha^i(A)) = 0$) are gauge generators, and gauge parameters ξ^α ($\varepsilon(\xi^\alpha) = \varepsilon_\alpha$) are arbitrary functions of space-time coordinates. It is assumed that the fields $A = \{A^i\}$ are linear independent with respect to the index i however, in general, the generators $R_\alpha^i(A)$ may be linear dependent with respect to index α . Linear dependence of $R_\alpha^i(A)$ implies that the matrix $R_\alpha^i(A)$ has at the extremals $S_{0,j}[A] = 0$ zero-eigenvalue eigenvectors $Z_{\alpha_1}^\alpha = Z_{\alpha_1}^\alpha(A)$, such that

$$R_\alpha^i(A)Z_{\alpha_1}^\alpha(A) = S_{0,j}[A]K_{\alpha_1}^{ji}(A), \quad \alpha_1 = 1, \dots, m_1, \quad (2)$$

and the number $\varepsilon_{\alpha_1} = 0, 1$ can be found in such a way that $\varepsilon(Z_{\alpha_1}^\alpha) = \varepsilon_\alpha + \varepsilon_{\alpha_1}$. Matrices $K_{\alpha_1}^{ij} = K_{\alpha_1}^{ij}(A)$ can be chosen to possess the properties:

$$K_{\alpha_1}^{ij} = -(-1)^{\varepsilon_i\varepsilon_j}K_{\alpha_1}^{ji}, \quad \varepsilon(K_{\alpha_1}^{ji}) = \varepsilon_i + \varepsilon_j + \varepsilon_{\alpha_1}.$$

Here, we restrict ourself to the case of first-stage reducibility when the set of zero-eigenvalue eigenvectors $Z_{\alpha_1}^\alpha$ is linear independent with respect to the index α_1 .

In general, the generators $R_\alpha^i(A)$ satisfy the following relations

$$R_{\alpha,j}^i(A)R_\beta^j(A) - (-1)^{\varepsilon_\alpha\varepsilon_\beta}R_{\beta,j}^i(A)R_\alpha^j(A) = -R_\gamma^i(A)F_{\alpha\beta}^\gamma(A) - S_{0,j}[A]M_{\alpha\beta}^{ji}(A), \quad (3)$$

where $F_{\alpha\beta}^\gamma(A) = F_{\alpha\beta}^\gamma$ ($\varepsilon(F_{\alpha\beta}^\gamma) = \varepsilon_\alpha + \varepsilon_\beta + \varepsilon_\gamma$, $\text{gh}(F_{\alpha\beta}^\gamma) = 0$) are the structure coefficients which may depend on the fields A^i . They obey the following symmetry properties $F_{\alpha\beta}^\gamma = -(-1)^{\varepsilon_\alpha\varepsilon_\beta}F_{\beta\alpha}^\gamma$, and $M_{\alpha\beta}^{ij}(A) = M_{\alpha\beta}^{ij}$ satisfy the conditions $M_{\alpha\beta}^{ij} = -(-1)^{\varepsilon_i\varepsilon_j}M_{\alpha\beta}^{ji} = -(-1)^{\varepsilon_\alpha\varepsilon_\beta}M_{\beta\alpha}^{ij}$. If $M_{\alpha\beta}^{ij} = 0$ then the gauge algebra is called closed.

According to main statements of papers [5, 6, 7], the deformation procedure for general gauge theories can be described in terms of the set of generating functions $h^i(A)$ of anticanonical transformations depending on initial fields A^i only and having the same transformation properties and quantum numbers as for A^i . The deformed initial action, $\tilde{S}_0[A]$, is defined by the relation

$$\tilde{S}_0[A] = S_0[A + h(A)] \quad (4)$$

being invariant under the deformed gauge transformation

$$\tilde{S}_{0,i}[A]\tilde{R}_\alpha^i(A) = 0, \quad \tilde{\delta}A = \tilde{R}_\alpha^i(A)\xi^\alpha, \quad (5)$$

where $\tilde{R}_\alpha^i(A)$ are the deformed gauge generators,

$$\tilde{R}_\alpha^i(A) = (M^{-1}(A))^i_j R_\alpha^j(A + h(A)). \quad (6)$$

Here, the matrix $(M^{-1}(A))^i_j$ is inverse to

$$M_j^i(A) = \delta_j^i + h^i_{,j}(A), \quad (7)$$

³To simplify presentation of all relations containing the right functional derivative of functional $S_0[A]$ with respect to field A^i we will use the symbol $S_{0,i}[A]$.

The deformed gauge generators on extremals of the deformed action are linear dependent,

$$\tilde{R}_\alpha^i(A)\tilde{Z}_{\alpha_1}^\alpha(A) = \tilde{S}_{0,j}[A]\tilde{K}_{\alpha_1}^{ji}(A), \quad (8)$$

where the functions $\tilde{Z}_{\alpha_1}^\alpha(A)$ and $\tilde{K}_{\alpha_1}^{ji}(A)$ are

$$\tilde{Z}_{\alpha_1}^\alpha(A) = Z_{\alpha_1}^\alpha(A + h(A)), \quad \tilde{K}_{\alpha_1}^{ji}(A) = -(M^{-1}(A))_l^j(M^{-1}(A))_k^i K_{\alpha_1}^{kl}(A + h(A))(-1)^{\varepsilon_l \varepsilon_i}. \quad (9)$$

The deformed gauge generators satisfy the relations

$$\tilde{R}_{\alpha,j}^i(A)\tilde{R}_\beta^j(A) - (-1)^{\varepsilon_\alpha \varepsilon_\beta} \tilde{R}_{\beta,j}^i(A)\tilde{R}_\alpha^j(A) = -\tilde{R}_\gamma^i(A)\tilde{F}_{\alpha\beta}^\gamma(A) - \tilde{S}_{0,j}[A]\tilde{M}_{\alpha\beta}^{ji}(A), \quad (10)$$

where $\tilde{F}_{\alpha\beta}^\gamma(A) = F_{\alpha\beta}^\gamma(A + h(A))$ and $\tilde{M}_{\alpha\beta}^{ji}(A) = -(M^{-1}(A))_l^j(M^{-1}(A))_k^i M_{\alpha\beta}^{kl}(A + h(A))(-1)^{\varepsilon_l \varepsilon_i}$. Therefore, the deformed theory looks like as the gauge theory of the first-stage reducibility similar to the initial one. It is remarkable fact that the deformation of an initial gauge theory is described in the explicit and closed form with the help of generating functions $h^i(A)$ of special anticanonical transformations in the BV-formalism. Moreover, all arbitrariness in the deformation procedure is controlled by the generating functions $h^i(A)$ only. Any choice of these functions guarantees that the deformed action will be invariant under the corresponding deformed gauge transformations.

Deformation of initial action looks like as the shift of argument $A \rightarrow A + h(A)$. Such kind of transformations is trivial in the case of local function $h(A)$. To obtain non-trivial deformations, the function $h(A)$ must be non-local. It means the non-locality of the deformed action. It may happen that there exists a local sector of the deformed action. If this local part of the deformed action will be invariant under local part of the deformed gauge generators then the deformation procedure creates a consistent local gauge-invariant theory. Namely, in paper [5] it was exactly shown that the Yang-Mills theory has been reproduced by special non-local deformation of free vector field model.

3 Fronsdal action for massless spin 3 fields

The Fronsdal action for massless spin 3 fields has the form

$$\begin{aligned} S_0[\varphi] = & \varphi^{\mu\nu\rho}(x)\square\varphi_{\mu\nu\rho}(x) - 3\eta_{\mu\nu}\eta^{\rho\sigma}\varphi^{\mu\nu\delta}(x)\square\varphi_{\rho\sigma\delta}(x) - \frac{3}{2}\varphi_{\mu\nu\lambda}\eta^{\mu\nu}\partial^\lambda\partial_\alpha\varphi^{\alpha\beta\gamma}\eta_{\beta\gamma} - \\ & - 3\varphi^{\mu\rho\sigma}(x)\partial_\mu\partial^\nu\varphi_{\nu\rho\sigma}(x) + 6\eta_{\mu\nu}\varphi^{\mu\nu\delta}(x)\partial^\rho\partial^\sigma\varphi_{\rho\sigma\delta}(x), \end{aligned} \quad (11)$$

where $\varphi^{\mu\nu\lambda}$ is completely symmetric third rank tensor, \square is the D'Alembertian, $\square = \partial_\mu\partial^\mu$, and $\eta_{\mu\nu}$ is the metric tensor of flat Minkowski space of the dimension d . The action (11) is invariant under the gauge transformations⁴

$$\delta\varphi^{\mu\nu\lambda} = \partial^{(\mu}\xi^{\nu\lambda)} = R_{\rho\sigma}^{\mu\nu\lambda}\xi^{\rho\sigma} \quad (12)$$

when gauge parameters $\xi^{\mu\nu}$ are subjected to the traceless condition,

$$\eta_{\mu\nu}\xi^{\mu\nu} = 0. \quad (13)$$

The gauge generators, $R_{\rho\sigma}^{\mu\nu\lambda} = \partial^{(\mu}\delta_\rho^\nu\delta_\sigma^\lambda)$, do not dependent on fields. They are symmetric in lower indices, $R_{\rho\sigma}^{\mu\nu\lambda} = R_{\sigma\rho}^{\mu\nu\lambda}$, and, therefore, satisfy the relations

$$R_{\rho\sigma}^{\mu\nu\lambda}Z_\ell^{\rho\sigma} = 0, \quad Z_\ell^{\rho\sigma} = -Z_\ell^{\sigma\rho}, \quad \ell = 1, 2, \dots, \frac{d(d-1)}{2}, \quad (14)$$

⁴The symbol (\dots) means the cycle permutation of indexes involved.

with antisymmetric matrices $Z_\ell^{\rho\sigma}$ which are linear independent with respect to index ℓ . We assume that $Z_\ell^{\rho\sigma}$ are Lorentz second-rank tensor with respect to upper indices. Therefore, in the BV-formalism we have to consider the case $s = 3$ as a gauge theory with first-stage reducible algebra specified by additional restrictions, $F_{\alpha\beta}^\gamma(A) = 0$, $M_{\alpha\beta}^{ij}(A) = 0$, $K_{\alpha_1}^{ij}(A) = 0$. The zero-eigenvalue eigenvectors, $Z_\ell^{\rho\sigma}$, do not depend on fields. In particular, it means that they do not change in the deformation process.

4 Deformations of initial action

We are going to study a possibility in construction of interactions of massless spin 3 fields with a real scalar field $\phi = \phi(x)$. We assume that the initial action has the form

$$S_0[\varphi, \phi] = S_0[\varphi] + S_0[\phi], \quad (15)$$

where $S_0[\varphi]$ is defined in (11) and $S_0[\phi]$ is the action of a free real scalar field,

$$S_0[\phi] = \frac{1}{2}(\partial_\mu\phi \partial^\mu\phi - m^2\phi^2). \quad (16)$$

The action (15) is invariant under the following gauge transformations

$$\delta\varphi^{\mu\nu\lambda} = \partial^{(\mu}\xi^{\nu\lambda)}, \quad \delta\phi = 0, \quad \eta_{\mu\nu}\xi^{\mu\nu} = 0, \quad (17)$$

and, therefore, belongs to the class of first-stage reducible theories. Relations with notations used in the Sec.2 are

$$\begin{aligned} A^i &= (\varphi^{\mu\nu\lambda}, \phi), \quad R_\alpha^i(A) = (R_{\nu_1\nu_2}^{\mu_1\mu_2\mu_3}, 0), \quad Z_{\alpha_1}^\alpha(A) = (Z_\ell^{\nu_1\nu_2}, 0), \quad \xi^\alpha = (\xi^{\mu\nu}, 0), \\ F_{\alpha\beta}^\gamma(A) &= 0, \quad M_{\alpha\beta}^{ij}(A) = 0, \quad K_{\alpha_1}^{ij}(A) = 0. \end{aligned} \quad (18)$$

Now, we consider possible deformations of initial action using the procedure which is ruled out by the generating functions $h^i(A)$. Here, we restrict ourself by the case of anticanonical transformations acting effectively in the sector of fields $\varphi^{\mu\nu\lambda}$ of the initial theory. It means the following structure of generating functions $h^i(A) = (h^{\mu\nu\lambda}(\varphi, \phi), 0)$. In construction of suitable generating functions $h^{\mu\nu\lambda} = (h^{\mu\nu\lambda}(\varphi, \phi))$, we have to take into account the dimensions of quantities involved in the initial action $S_0[\varphi, \phi]$,

$$\dim(\varphi^{\mu\nu\lambda}) = \dim(\phi) = \frac{d-2}{2}, \quad \dim(\xi^{\mu\nu}) = \frac{d-4}{2}, \quad \dim(\partial_\mu) = 1, \quad \dim(\square) = 2. \quad (19)$$

The generating function $h^{\mu\nu\lambda}$ should be symmetric and non-local with the dimension equals to $(d-2)/2$. The non-locality will be achieved by using the operator $1/\square$. To construct cubic vertices $\sim \varphi\phi\phi$, the $h^{\mu\nu\lambda}$ should be at least quadratic in fields ϕ . The tensor structure of $h^{\mu\nu\lambda}$ is obeyed by using partial derivatives ∂_μ and the metric tensor $\eta_{\mu\nu}$. The minimal number of derivatives equals to 3. Therefore, the more general form of $h^{\mu\nu\lambda} = h^{\mu\nu\lambda}(\phi)$ satisfying requirements listed above reads

$$\begin{aligned} h^{\mu\nu\lambda} &= a_0 \frac{1}{\square} (c_1 \partial^\mu \partial^\nu \partial^\lambda \phi \phi + c_2 \partial^{(\mu} \partial^\nu \phi \partial^{\lambda)} \phi + c_3 \eta^{(\mu\nu} \square \partial^{\lambda)} \phi \phi + \\ &\quad + c_4 \square \phi \eta^{(\mu\nu} \partial^{\lambda)} \phi + c_5 \eta^{(\mu\nu} \partial^{\lambda)} \partial_\sigma \phi \partial^\sigma \phi), \end{aligned} \quad (20)$$

where a_0 is the coupling constant with $\dim(a_0) = -d/2$ and c_i , $i = 1, 2, \dots, 5$ are arbitrary dimensionless constants. Local part of the deformed action has the form

$$S_{loc}[\varphi, \phi] = S_0[\varphi] + S_0[\phi] + S_{1\ loc}[\varphi, \phi] \quad (21)$$

where $S_{1\text{ loc}} = S_{1\text{ loc}}[\varphi, \phi]$,

$$S_{1\text{ loc}} = 2a_0\varphi_{\mu\nu\lambda} \left[c_1\partial^\mu\partial^\nu\partial^\lambda\phi\phi + c_2\partial^{(\mu}\partial^\nu\phi\partial^{\lambda)}\phi - (c_1 + c_3(d+1))\eta^{(\mu\nu}\square\partial^{\lambda)}\phi\phi - \right. \\ \left. - (c_2 + c_4(d+1))\square\phi\eta^{(\mu\nu}\partial^{\lambda)}\phi - (2c_2 + c_5(d+1))\eta^{(\mu\nu}\partial^{\lambda)}\partial_\sigma\phi\partial^\sigma\phi \right], \quad (22)$$

corresponds to possible cubic vertices. Due to the special structure of generating functions $h^i(A) = (h^{\mu\nu\lambda}(\phi), 0)$, the matrix $(M^{-1}(A))^i_j$ can be found in the explicit form

$$(M^{-1}(A))^i_j = \begin{pmatrix} E^{\mu\nu\lambda}_{\rho\sigma\gamma} & -h^{\mu\nu\lambda}(\phi)\overleftarrow{\partial}_\phi \\ 0 & 1 \end{pmatrix}, \quad (23)$$

where $E^{\mu\nu\lambda}_{\rho\sigma\gamma}$ are elements of the unit matrix in the space of third rank symmetric tensors. Therefore, the deformed gauge generators coincide with initial ones, $\tilde{R}_\alpha^i(A) = R_\alpha^i(A)$. Consider the variation of $S_{1\text{ loc}}$ under the gauge transformations (17). We have

$$\delta S_{1\text{ loc}} = -6a_0\xi_{\nu\lambda} \left[- (c_1 + 2c_3(d+1))\square\partial^\nu\partial^\lambda\phi\phi + (c_1 - 3c_2 - 2c_5(d+1))\partial^\mu\partial^\nu\partial^\lambda\phi\partial_\mu\phi - \right. \\ \left. - 2c_4(d+1)\square\partial^\nu\phi\partial^\lambda\phi - -2(c_2 + c_5(d+1))\partial^\lambda\partial^\mu\phi\partial_\mu\partial^\nu\phi - \right. \\ \left. - (c_2 + 2c_4(d+1))\square\phi\partial^\nu\partial^\lambda\phi \right]. \quad (24)$$

The system of algebraic equations appears

$$c_1 + 2c_3(d+1) = 0, \quad c_1 - 3c_2 - 2c_5(d+1) = 0, \quad c_4(d+1) = 0, \quad (25)$$

$$c_2 + c_5(d+1) = 0, \quad c_2 + 2c_4(d+1) = 0 \quad (26)$$

as consequence of requirement $\delta S_{1\text{ loc}} = 0$. This system has the solution

$$c_i = 0, \quad i = 1, 2, \dots, 5, \quad (27)$$

which leads to the important conclusion

$$S_{1\text{ loc}}[\varphi, \phi] = 0. \quad (28)$$

Therefore, we have proved impossibility in the theory of massless spin 3 and scalar fields to construct cubic vertices $\varphi\phi\phi$ being invariant under initial gauge transformations (17).

In connection with the obtained result, a natural question arises: Do gauge invariant quartic vertices $\varphi\phi\phi\phi$ in the theory under consideration exist? Repeating the main arguments that led to the construction of the generating function (20) of the anticanonical transformation, the most general form of the generating function $h^{\mu\nu\lambda}$ with three derivatives responsible for the generation of quartic vertices reads

$$h^{\mu\nu\lambda} = a_1\frac{1}{\square} \left[c_1\partial^\mu\partial^\nu\partial^\lambda\phi\phi^2 + c_2\partial^{(\mu}\partial^\nu\phi\partial^{\lambda)}\phi\phi + c_3\partial^\mu\phi\partial^\nu\phi\partial^\lambda\phi + c_4\eta^{(\mu\nu}\partial^{\lambda)}\square\phi\phi^2 + \right. \\ \left. + c_5\eta^{(\mu\nu}\partial^{\lambda)}\partial_\sigma\phi\partial^\sigma\phi\phi + c_6\eta^{(\mu\nu}\partial^{\lambda)}\phi\partial_\sigma\phi\partial^\sigma\phi + c_7\square\phi\eta^{(\mu\nu}\partial^{\lambda)}\phi\phi \right], \quad (29)$$

where a_1 is a coupling constant with $\dim(a_1) = -(d-1)$ and c_i , $i = 1, 2, \dots, 7$ are arbitrary dimensionless constants. For the local addition, $S_{2\text{ loc}} = S_{2\text{ loc}}[\varphi, \phi]$, to the initial action (15), we obtain

$$S_{2\text{ loc}} = 2a_1\varphi_{\mu\nu\lambda} \left[c_1\partial^\mu\partial^\nu\partial^\lambda\phi\phi^2 + c_2\partial^{(\mu}\partial^\nu\phi\partial^{\lambda)}\phi\phi + c_3\partial^\mu\phi\partial^\nu\phi\partial^\lambda\phi - \right. \\ \left. - (c_1 + c_4(d+1))\eta^{(\mu\nu}\partial^{\lambda)}\square\phi\phi^2 - (2c_2 + c_5(d+1))\eta^{(\mu\nu}\partial^{\lambda)}\partial_\sigma\phi\partial^\sigma\phi\phi - \right. \\ \left. - (c_3 + c_6(d+1))\eta^{(\mu\nu}\partial^{\lambda)}\phi\partial_\sigma\phi\partial^\sigma\phi - (c_2 + c_7(d+1))\square\phi\eta^{(\mu\nu}\partial^{\lambda)}\phi\phi \right]. \quad (30)$$

Notice that as in previous case, the gauge generators do not transform under anticanonical transformations generated by functions (29).

Omitting the algebraic manipulations, we find the variation $S_{2\text{ loc}}$ under gauge transformations (17) in the form

$$\begin{aligned} \delta S_{2\text{ loc}} = & -6a_1\xi_{\nu\lambda} \left[-(c_1 + 2c_4(d+1))\partial^\nu\partial^\lambda\Box\phi\phi^2 + (2c_1 - 3c_2 - 2c_5(d+1))\partial^\sigma\partial^\nu\partial^\lambda\phi\partial_\sigma\phi\phi - \right. \\ & -2(2c_1 + 2c_4(d+1) + c_7(d+1))\Box\partial^\nu\phi\partial^\lambda\phi\phi - 2(c_2 + c_5(d+1))\partial^\sigma\partial^\nu\phi\partial^\lambda\partial_\sigma\phi\phi - \\ & -(c_2 + 2c_7(d+1))\partial^\nu\partial^\lambda\phi\Box\phi\phi - 2(c_2 + c_3 + (c_5 + 2c_6)(d+1))\partial^\sigma\partial^\nu\phi\partial^\lambda\phi\partial_\sigma\phi + \\ & \left. +(c_2 - 2c_3 - 2c_6(d+1))\partial^\nu\partial^\lambda\phi\partial_\sigma\phi\partial^{\sigma\phi} + (c_3 - 2c_2 - 2c_7(d+1))\Box\phi\partial^\nu\phi\partial^\lambda\phi \right]. \quad (31) \end{aligned}$$

Note that the system of algebraic equations

$$\begin{aligned} c_1 + 2c_4(d+1) = 0, \quad 2c_1 - 3c_2 - 2c_5(d+1) = 0, \quad 2c_1 + 2c_4(d+1) + c_7(d+1) = 0, \\ c_2 + c_5(d+1) = 0, \quad c_2 + 2c_7(d+1) = 0, \quad c_2 + c_3 + (c_5 + 2c_6)(d+1) = 0, \\ c_2 - 2c_3 - 2c_6(d+1) = 0, \quad c_3 - 2c_2 - 2c_7(d+1) = 0 \end{aligned} \quad (32)$$

has non-trivial solution

$$c_2 = c_3 = 2c_1, \quad c_4 = -\frac{1}{2(d+1)}c_1, \quad c_5 = -\frac{2}{d+1}c_1, \quad c_6 = c_7 = -\frac{1}{d+1}c_1. \quad (33)$$

Therefore, the functional

$$\begin{aligned} S_{2\text{ loc}} = & 2a_1\varphi_{\mu\nu\lambda} \left[\partial^\mu\partial^\nu\partial^\lambda\phi\phi^2 + 2\partial^{(\mu}\partial^\nu\phi\partial^{\lambda)}\phi\phi + 2\partial^\mu\phi\partial^\nu\phi\partial^\lambda\phi - \frac{1}{2}\eta^{(\mu\nu}\partial^{\lambda)}\Box\phi\phi^2 - \right. \\ & \left. -2\eta^{(\mu\nu}\partial^{\lambda)}\partial_\sigma\phi\partial^\sigma\phi\phi - \eta^{(\mu\nu}\partial^{\lambda)}\phi\partial_\sigma\phi\partial^\sigma\phi - \Box\phi\eta^{(\mu\nu}\partial^{\lambda)}\phi\phi \right] \end{aligned} \quad (34)$$

is gauge invariant,

$$\delta S_{2\text{ loc}} = 0, \quad (35)$$

and presents the quartic vertices.

The local action

$$S[\varphi, \phi] = S_0[\varphi] + S_0[\phi] + S_{2\text{ loc}}[\varphi, \phi] \quad (36)$$

describes the model of interacting $\varphi^{\mu\nu\lambda}$ and ϕ fields which is invariant under gauge transformations (17) and belongs to the class of first-stage reducible theories. In fact, the model (36) is the first new example of local gauge theory constructed with using the method of our papers [5, 6, 7].

5 Discussion

In the present paper, the new approach [5, 6, 7] to the deformation procedure [3, 4] has been applied to construct local cubic and quartic vertices for massless spin 3 and scalar fields. It was shown that non-local anticanonical transformations acting non-trivial only in the sector of spin 3 fields forbid existence of local cubic vertices being invariant under initial gauge transformations while local gauge-invariant quartic vertices do exist. In fact, the new local gauge-invariant model in the theory of interacting massless integer higher spin fields has been proposed. It needs to stress that the obtained result cannot be reproduced in the Noether procedure adopted for construction of interactions in the theory of higher spin fields.

On the first sight, it seems that the obtained result (28) contradicts with the statement of paper [21]⁵ that the gauge invariant cubic vertices $\varphi\phi\phi$ in the theory of massless spin 3 and a pair of scalars ϕ^i , $i = 1, 2$ can be constructed. The corresponding action $S_{1Zin} = S_{1Zin}[\varphi, \phi]$ has the form

$$S_{1Zin} = \frac{a_0}{6} \varepsilon^{ij} \varphi_{\mu\nu\lambda} \left[-2\partial^\mu \partial^\nu \phi_i \partial^\lambda \phi_j + 2\eta^{\nu\lambda} \square \phi_i \partial^\mu \phi_j + \eta^{\nu\lambda} \partial^\mu \partial_\sigma \phi_i \partial^\sigma \phi_j \right]. \quad (37)$$

Indeed, this action is not invariant under the gauge transformations (17),

$$\delta S_{1Zin} = -\frac{a_0}{6} \varepsilon^{ij} \xi_{\nu\lambda} \left[-2\partial^\sigma \partial^\nu \phi_i \partial^\lambda \partial_\sigma \phi_j - 2\partial^\nu \partial^\lambda \phi_i \square \phi_j + 4\square \phi_i \partial^\nu \partial^\lambda \phi_j \right] \neq 0. \quad (38)$$

But it needs to have in mind that the cubic vertices for the initial free model of massless scalar fields and massless spin 3 fields were studied using the Noether's procedure when scalar fields were subjected to gauge transformations too. Then the gauge invariance of the theory using action (37) is supported in the linear approximation only. In the approach applied in Section 4 it corresponds to anticanonical transformations when the generating function responsible for change of massless scalar field ϕ ($m = 0$) is not equal to zero, $h \neq 0$. In its turn, it will lead to non-trivial deformations of scalar fields and gauge algebra.

We can analyze the interactions of massless spin 3 field with the pair of scalars ϕ^i , $i = 1, 2$ within the new approach leading to the cubic vertex which will be anti-symmetric on them. The most general form of generating function containing three partial derivatives reads

$$h^{\mu\nu\lambda} = a_0 \frac{1}{\square} \varepsilon^{ij} (c_1 \partial^\mu \partial^\nu \partial^\lambda \phi_i \phi_j + c_2 \partial^{(\mu} \partial^\nu \phi_i \partial^{\lambda)} \phi_j + c_3 \eta^{(\mu\nu} \square \partial^{\lambda)} \phi_i \phi_j + c_4 \square \phi_i \eta^{(\mu\nu} \partial^{\lambda)} \phi_j + c_5 \eta^{(\mu\nu} \partial^{\lambda)} \partial_\sigma \phi_i \partial^\sigma \phi_j), \quad (39)$$

where a_0 is the coupling constant with $\dim(a_0) = -d/2$ and c_i , $i = 1, 2, \dots, 5$ are arbitrary dimensionless constants. Then, the local addition to the initial action (15) has the form

$$S_{1loc} = 2a_0 \varphi_{\mu\nu\lambda} \varepsilon^{ij} \left[c_1 \partial^\mu \partial^\nu \partial^\lambda \phi_i \phi_j + c_2 \partial^{(\mu} \partial^\nu \phi_i \partial^{\lambda)} \phi_j - (c_1 + c_3(d+1)) \eta^{(\mu\nu} \square \partial^{\lambda)} \phi_i \phi_j - (c_2 + c_4(d+1)) \square \phi_i \eta^{(\mu\nu} \partial^{\lambda)} \phi_j - (2c_2 + c_5(d+1)) \eta^{(\mu\nu} \partial^{\lambda)} \partial_\sigma \phi_i \partial^\sigma \phi_j \right], \quad (40)$$

Due to the fact that the gauge algebra does not deform under anticanonical transformations with the generating function (39), the functional (40) must be invariant under original gauge transformations, $\delta S_{1loc} = 0$, $\delta \varphi_{\mu\nu\lambda} = \partial_{(\mu} \xi_{\nu\lambda)}$, $\eta_{\mu\nu} \xi^{\mu\nu} = 0$, $\delta \phi_i = 0$, $i = 1, 2$. Analysis of this requirement leads to the result $S_{1loc} = 0$. We can also discuss the situation with quartic vertices, $\sim \varphi\phi\phi\phi$, in the case under consideration. To construct such interactions, in general, one has to use the generating functions $h^{\mu\nu\lambda}$ in the sector of fields $\varphi^{\mu\nu\lambda}$ and h_i in the sector of fields ϕ_i . In turn, the structure of generating functions should be as follows $h^{\mu\nu\lambda} \sim \varepsilon^{jk} \partial^\mu \partial^\nu \partial^\lambda \phi_i \phi_j \phi_k$ and $h_i \sim \varepsilon^{jk} \partial_\mu \partial_\nu \partial_\lambda \varphi^{\mu\nu\lambda} \phi_j \phi_k$ that is contrary to the tensor properties for these functions. Therefore, quartic vertices are forbidden as well.

Quite recently, it was shown that non-trivial quartic vertices describing interactions among massless spin 3 field, $\varphi^{\mu\nu\lambda}$, and massive vector, A_μ , and real scalar, ϕ , fields of the form $\sim \varphi A \phi \phi$ can be constructed [23]. These vertices are invariant under original gauge transformations, similar to the case considered in Section 4.

Acknowledgments

The author thanks I.L. Buchbinder for stimulating discussions of different aspects concerning the problem of interactions for higher spin fields. The work is supported by the Ministry of Education of the Russian Federation, project FEWF-2020-0003.

⁵Notice that in [22] $N = 2$ supersymmetric generalization of cubic vertices containing in bosonic sector interactions of spin 0 field with massless integer spin s fields has been considered.

References

- [1] I.A. Batalin, G.A. Vilkovisky, *Gauge algebra and quantization*, Phys. Lett. **B102** (1981) 27.
- [2] I.A. Batalin, G.A. Vilkovisky, *Quantization of gauge theories with linearly dependent generators*, Phys. Rev. **D28** (1983) 2567.
- [3] G. Barnich, M. Henneaux, *Consistent coupling between fields with gauge freedom and deformation of master equation*, Phys. Lett. **B** 311 (1993) 123-129, arXiv:hep-th/9304057.
- [4] M. Henneaux, *Consistent interactions between gauge fields: The cohomological approach*, Contemp. Math. **219** (1998) 93-110, arXiv:hep-th/9712226.
- [5] I.L. Buchbinder, P.M. Lavrov, *On a gauge-invariant deformation of a classical gauge-invariant theory*, JHEP 06 (2021) 854, arXiv:2104.11930 [hep-th].
- [6] I.L. Buchbinder, P.M. Lavrov, *On classical and quantum deformations of gauge theories*, Eur. Phys. J. **C** 81 (2021) 856, arXiv:2108.09968 [hep-th].
- [7] P.M. Lavrov, *On gauge-invariant deformation of reducible gauge theories*, Eur. Phys. J. **C** 82 (2022) 429, arXiv:2201.07505 [hep-th].
- [8] E.S. Fradkin, G.A. Vilkovisky, *Quantization of relativistic systems with constraints*, Phys. Lett. **B** 55 (1975) 224.
- [9] I.A. Batalin, G.A. Vilkovisky, *Relativistic S-matrix of dynamical systems with boson and fermion constraints*, Phys. Lett. **B** 69 (1977) 309.
- [10] I.A. Batalin, E.S. Fradkin, *Operator quantization of relativistic dynamical system subject to first class constraints*, Phys. Lett. **B** 128 (1983) 303.
- [11] I.L. Buchbinder, P.M. Lavrov, *On deformations of constrained Hamiltonian systems in BFV-formalism*, arXiv:2203.05313 [hep-th].
- [12] I.A. Batalin, P.M. Lavrov, I.V. Tyutin, *Finite anticanonical transformations in field-antifield formalism*, Eur. Phys. J. **C75** (2015) 270.
- [13] I.A. Batalin, P.M. Lavrov, I.V. Tyutin, *Anticanonical transformations and Grand Jacobian*, Russ. Phys. J. **64** (2021) 688.
- [14] B.L. Voronov, P.M. Lavrov, I.V. Tyutin, *Canonical transformations and gauge dependence in general gauge theories*, Sov. J. Nucl. Phys. **36** (1982) 292.
- [15] J. Gomis, S. Weinberg, *Are nonrenormalizable gauge theories renormalizable?*, Nucl. Phys. B **469** (1996) 473.
- [16] C. Fronsdal, *Massless field with integer spin*, Phys. Rev. **D18** (1978) 3624.
- [17] I.L. Buchbinder, A.V. Galajinsky, V.A. Krykhtin, *Quartet unconstrained formulation for massless higher spin fields*, Nucl. Phys. **B779** (2007) 155.
- [18] I.L. Buchbinder, A.V. Galajinsky, *Quartet unconstrained formulation for massive higher spin fields*, JHEP 11 (2008) 081, arXiv: 0810.2852 [hep-th].
- [19] I. Buchbinder, E. Ivanov, N. Zaigraev, *Unconstrained off-shell superfield formulation of 4D, N = 2 supersymmetric higher spins*, JHEP 12 (2021) 016, arXiv:2109.07639 [hep-th].
- [20] B.S. DeWitt, *Dynamical theory of groups and fields*, (Gordon and Breach, 1965).
- [21] Yu.M. Zinoviev, *Spin 3 cubic vertices in a frame-like formalism*, JHEP 08 (2010) 084, arXiv:1007.0158 [hep-th].
- [22] I. Buchbinder, E. Ivanov, N. Zaigraev, *Off-shell cubic hypermultiplet couplings to N=2 higher spin gauge superfields*, JHEP 05 (2022) 104, arXiv:2202.08196 [hep-th].
- [23] P.M. Lavrov, *Gauge-invariant models of interacting fields with spins 3,1 and 0*. arXiv:2209.03678 [hep-th].