

Propagation of spinors on a noncommutative spacetime: equivalence of the formal and the effective approach

Marija Dimitrijević-Čirić,^{1,*} Nikola Konjik,^{1,†} and Andjelo Samsarov^{2,‡}

¹ Faculty of Physics, University of Belgrade, Studentski trg 12, 11000 Beograd, Serbia

² Rudjer Bošković Institute, Bijenička c.54, HR-10002 Zagreb, Croatia

(Dated: August 15, 2022)

Some noncommutative (NC) theories possess a certain type of dualities that are implicitly built within their structure. In this paper we establish still another example of this kind. More precisely, we show that the noncommutative $U(1)_*$ gauge theory coupled to a NC scalar field and to a classical geometry of the Reissner–Nordström (RN) type is completely equivalent at the level of equations of motion to the commutative $U(1)$ gauge theory coupled to a commutative scalar field and to a classical geometry background, different from the starting RN background. The new (effective) metric is obtained from the RN metric by switching on an additional nonvanishing $r - \phi$ component. Using this duality between two theories and physical systems they describe, we formulate an effective approach to studying a dynamics of spin $\frac{1}{2}$ fields on the curved background of RN type with an abiding noncommutative structure. As opposed to that, we also study the dynamics of spin $\frac{1}{2}$ fields in a more formal way, by studying the semiclassical theory which describes the NC $U(1)_*$ gauge field coupled with NC spin $\frac{1}{2}$ field and also with gravity which is however treated classically. Upon utilising the Seiberg–Witten map in order to write the NC spinor and NC gauge fields in terms of their corresponding commutative degrees of freedom, we find that the equation of motion for the fermion field obtained within the formal approach exactly coincides with the equation of motion obtained within the effective approach that utilises noncommutative duality. We then use these results to analyze the problem of stability of solutions of the equations of motion and the associated issue of superradiance, as related to fermions in RN spacetime with an all-pervasive noncommutative structure.

I. INTRODUCTION

Many distinct approaches to a unification of quantum mechanics with gravity in the ultraviolet sector point toward the existence of absolute minimal length scale, whose very presence puts a lower bound to the minimal possible resolution of space as its intrinsic property. As a consequence, one of the cornerstones of quantum mechanics, the Heisenberg uncertainty relations, start to call for a revision, resulting in a generalized uncertainty principle [1],[2], which is also suggested by perturbative string theory [3],[4], quantum gravity [5],[6] and black hole physics [7]. Moreover, in loop quantum gravity a process of quantization gives rise to the area and volume operators which have discrete spectra, whose lowest possible eigenvalues are being proportional to the square and cube of the Planck length, respectively [8],[9]. This, together with the generalized uncertainty principle, implies the existence of a minimum uncertainty in position [7],[10],[11],[12].

One of the more known patterns to implement a minimal length scale in quantum mechanics, quantum field theory (QFT) and gravity is provided by the frame of noncommutative (NC) geometry, which is characterized by the fact that the spacetime coordinates get raised to the level of the operators and thus generally fail to commute [13–17]. The idea of noncommutativity of spacetime was first clearly articulated by Snyder [18] and expounded further in terms of geometric notions by Alain Connes [19],[20]. Correspondingly, a possibility to observe the consequences of spacetime noncommutativity and the existence of a minimal length scale led to the intensive study of noncommutative versions of quantum mechanics, QFT and gravity with the aim of revising the standard theories as diverse as the gauge theory, particle physics and the Quantum Hall effect (QHE), thermodynamics and the black-hole physics and cosmology, so that they may keep the track with and accommodate the eventual novel features into their framework [21–32].

It is known in general that various noncommutative models, including models of noncommutative field theory and noncommutative gravity allow for a representation in terms of classical commutative fields either in a compact and closed form or in a form of a perturbation expansion up to a certain order in a deformation parameter. The most known

*Electronic address: dmarija@ipb.ac.rs

†Electronic address: konjik@ipb.ac.rs

‡Electronic address: asamsarov@irb.hr

example of this kind is obtained by means of the Seiberg-Witten (SW) map [15], which is a field transformation that allows to rewrite a gauge theory on noncommutative space as a gauge theory on commutative space. In an attempt to map a noncommutative gauge field theory to its commutative counterpart from which it emerged as a result of deforming an associated Poisson structure, the SW map undoubtedly becomes highly important.

Related to this, it is noteworthy to recall that the SW map helped to resolve some of the issues that had appeared by the introduction of the star-product into the action, among them being the advent of the field operator ordering ambiguities, as well as the breaking of the ordinary gauge invariance and the problem with the charge quantisation. This map also ensures that by going from a set of degrees of freedom describing noncommutative gauge symmetry to a corresponding set describing local commutative gauge symmetry the number of degrees of freedom stays the same. This way a number of NC deformed QFT's could be properly defined for arbitrary gauge group representations, which has facilitated the building of the whole range of semi-realistic NC deformed particle physics [33–41] and gravity models [42–46].

Another interesting situation where the NC theory allows for an interpretation in terms of commutative degrees of freedom involves a class of noncommutative models which have been shown to exhibit a specific type of duality relations that are implicitly built-in at the level of equations of motion. This feature has been observed in some hybrid or semi-hybrid noncommutative models investigated within a so called realization framework [47–53], which utilizes the representation of NC coordinates and NC field operators in terms of formal power series of generators of the undeformed Heisenberg algebra. The Lagrangian density in these models, which is expressed partially or completely in terms of NC degrees of freedom (that's why these models are being labelled as hybrid or semi-hybrid), allowed not only for a reinterpretation of the Lagrangian density in terms of commutative degrees of freedom, but also allowed for a radical refashioning of the initial semi-hybrid NC model so that it could take on a form of an effective commutative model realized within a similar or even the same physical setting, but with modified system parameters. Such reinterpretation was then able to give noncommutativity a definite physical meaning. For example, in [54–58] it has been found that the semi-hybrid model of NC massless scalar field coupled to a classical nonrotational BTZ geometry is dual to the model of massive commutative scalar field probing the geometry of a rotating BTZ black hole. In this way, the noncommutativity took on the role of an agent medium that has put a black hole into a state of rotation. Besides, the noncommutativity was shown to be responsible for a mass generating mechanism, as applied to a scalar probe, and for inducing certain back-reaction effects.

A similar situation has been encountered in [59], where the analogy between NC version of the Schwarzschild black hole and the commutative Reissner–Nordström black hole with a stretched horizon was drawn. Likewise, in [60] the authors have shown that the minimal $U(1)$ NCQED based on a reversible Seiberg-Witten (SW) map is equivalent to the Moyal NCQED without SW map, as manifested at the level of tree-level scattering amplitudes [60]. In this case the equivalence between two models comes as a result of a mutual cancellation between terms induced by the reversible SW map, which might also be viewed as being due to a presence of the specific duality relations that are inherent to the noncommutative model being considered. Similarly, the features of this kind can be found in supersymmetric noncommutative field theories related by the theta-exact Seiberg-Witten map [61]. In most of these cases (certainly for the case studied in [54–58]) the notion of duality refers to an exact mathematical correspondence that may be drawn between two different physical systems having different system parameters, though governed by the same Lagrangian density and the associated equation of motion. A duality understood that way gives an example of the equivalence that can be established between noncommutative and commutative model, where each of these models separately describes its own respective physical system. As these two systems that commutative and noncommutative models refer to are actually being governed by the same equations of motion and the same Lagrangian density, they may too be characterized as being equivalent with each other. Therefore, referring in the current context to a duality itself that exists between two different physical systems, it shouldn't come as a surprise that it gets manifested through a set of exact mathematical transformations that connect the parameters of these two physical systems, thus making them dual with each other.

In this paper we set out to find another example of this kind within the semi-hybrid NC model studied in [62–64]. In particular, here we show that noncommutative $U(1)_*$ gauge theory coupled to NC scalar field and to a classical geometry of the Reissner–Nordström type is equivalent to a commutative $U(1)$ gauge theory coupled to a commutative scalar field and to a classical geometry background, which however does not coincide with the initial RN metric, but instead represents an effective metric which encodes the impacts of the spacetime deformation. In other words the former model can be recast into a latter by redefining the components of the initial RN metric. In this way we end up with an effective, but equivalent description in which the redefined metric has an important role, forming one of the crucial building blocks that characterize the dual system and the duality transformation itself. From now on we refer to this redefined metric as the effective metric characterizing the dual picture, or the effective dual metric in short. It can be viewed as having ensued from a specific deformation of the RN metric which brings in a nonvanishing $r - \phi$ component. This whole scheme thus establishes a duality relation between two models, one

semi-hybrid noncommutative and the other fully commutative, and two physical systems they describe.

As a further step, we study a dynamics of fermions on a curved background with a deformed spacetime structure, where as an exemplar for the curved geometry we use the RN black hole background. This study has been carried out within two different approaches: effective and formal. The effective approach uses standard notions of commutative differential geometry, with parallel implementation of the noncommutative-born effects through the utilisation of the type of duality just explained, where the effective dual metric mimicks the impacts of noncommutativity. The formal approach is a level beyond more rigorous in a sense that it attempts to stay in line and be compatible with the requirements that the NC $U(1)_*$ Dirac action on a curved background remains invariant under NC gauge transformations, as well as under local $SO(1,3)_*$. With that in mind we come out with a proposal for the NC $U(1)_*$ Dirac action that could meet these requirements. As the NC $U(1)_*$ Dirac action that we have proposed is being expressed in terms of the NC spin $\frac{1}{2}$ field and the NC $U(1)_*$ gauge field that are both coupled to gravitational degrees of freedom, it is important to stress that this action is going to stay invariant under the NC gauge transformations as long as we assume that the gravity is unaffected by them, i.e. $\delta_* g_{\mu\nu} = 0$ (or alternatively $\delta_* e^a{}_\mu = \delta_* \omega_\mu{}^{ab} = 0$). In other words, in our formal approach we shall assume that the NC spin $\frac{1}{2}$ field, and the NC $U(1)$ gauge field are the only degrees of freedom that get affected by the NC gauge transformations. The NC $U(1)_*$ Dirac action that we propose is in a line with the proposals made in [42],[43].

The main result of this paper is to show that these two different approaches surprisingly give rise to the same equation of motion describing a dynamics of fermions on a curved space in presence of a NC structure of spacetime, therefore explicitly demonstrating the equivalence between the effective and formal, more rigorous approach. Namely, what has been shown in the present paper is that the equation of motion obtained by using the SW map for spin $\frac{1}{2}$ and gauge fields, and by varying the NC $U(1)_*$ Dirac action that is invariant under NC gauge transformations and $S(1,3)_*$ group at the end of the day appears to be the same as the equation of motion obtained by simply writing the Dirac equation on a geometric background described by the effective dual metric.

Dirac equation in the context of noncommutative spaces is important from many reasons and was studied intensively in the literature. The range of topics where it finds application is vast, going from the high energy physics, all through the gravitational physics and cosmology and all the way down to the problems in condensed matter. In particular, the problems related to high energy physics involve a study of the hydrogen atom spectrum on Moyal [65] and kappa-Minkowski space [66], the impact of quantum deformation on the spin- $\frac{1}{2}$ Aharonov–Bohm effect [67], the problem of Yukawa couplings and seesaw neutrino masses in noncommutative gauge theory [39], photon-neutrino interaction in noncommutative field theory [68], renormalizability and dispersion of chiral fermions in NCQED [69] and the impact on neutrino oscillations due to noncommutativity of spacetime [70] to name just a few. Besides canonical and kappa-Minkowski type of noncommutativity (for a recent review see [71]), other types of spacetime noncommutativity that have been frequently studied in the past include that of Snyder type (for the review see [72, 73]), as well as that which is usually referred to as the “spin noncommutativity”, first introduced in [74, 75], and which could be theoretically understood as a non-relativistic analog of the original Snyder’s model [18]. In [76], the spin noncommutativity was obtained by means of a consistent deformation of the Berezin-Marinov pseudoclassical model for the spinning particle [77]. Like the Snyder model, spin noncommutativity exhibits preservation of the Lorentz symmetry. Within this framework a modification of the Dirac equation was proposed and a dynamics of a Dirac fermion in the presence of spin noncommutativity was studied in [76, 78].

In condensed matter, the topic of special interest is the integer and in particular the fractional quantum Hall effect and a related attempt to procure the explanation for the latter in terms of the Dirac oscillator [79] and especially in terms of the Dirac oscillator on NC space [80–88]. Due to the same reasons the study of relativistic Landau levels or Dirac-Landau levels, including the breaking of their degeneracy, becomes increasingly more important and even more so as these results were being applied to the graphene and the related nanostructures [89],[90].

Another intriguing field where the Dirac equation under conditions of discretized spacetime was being analyzed involves deformed relativistic wave equations, namely the Klein–Gordon and Dirac equations in a Doubly Special Relativity (DSR) scenario [91–93]. Besides the algebraic approach to this problem, which originally started by considering the standard real form of the quantum anti-de Sitter algebra, $SO_q(3,2)$ and then by consistently modifying the related coproduct [94], there recently appeared a geometric approach [95–98] to the same problem, which is based on the geometry of a curved momentum space [99–105]. While the Dirac equation obtained in [94] is invariant under the spin-half representation of the κ -Poincaré algebra, it unlike the commutative case doesn’t yield the Casimir after squaring. Instead, its square gives rise to the κ -deformed Pauli-Lubanski vector. Contrary to that, the Dirac equation obtained in [98] gives rise to the κ -Poincaré Casimir upon squaring, along with having the required symmetry properties. This geometric approach should be seen as complementary to the more spread algebraic one [106–112]. Finally, it is worthy to mention that the Dirac equation in a presence of noncommutativity has an important role in studying the impact of NC spacetime deformation on quasinormal modes (QNM) spectrum of the fermionic perturbations of black holes [57].

The paper is organized as follows. After a very brief review of NC deformation that we analyze in this paper, in Section III we study the NC $U(1)_*$ gauge theory coupled with NC spin $\frac{1}{2}$ field and NC gravitational degrees of freedom, whose action is invariant under NC gauge transformations and local $SO(1, 3)_*$ group. Upon utilising the Seiberg-Witten map in order to write the NC spinor and NC gauge fields in terms of their corresponding commutative degrees of freedom, the equation of motion for the fermion field is obtained by varying the action over $\bar{\Psi}$. In Section IV the equivalence between NC $U(1)_*$ gauge theory with NC scalar field on a classical RN background and the commutative $U(1)$ gauge theory with ordinary scalar field on the background with the effective dual metric is established. Moreover, the explicit form of the effective dual metric is found. This effective dual metric is then used in Section V to write a noncommutative version of the equation of motion for the fermions in a curved background of RN type, thus putting forth and unrolling the effective approach of the same problem that was treated in a formal, more rigorous way in Section III. Here we find that both of these two approaches, formal as well as effective, surprisingly yield the same final result. In Section VI we show that the resulting equation of motion is separable, yielding two pairs of equations, one for the angular part and the other for the radial part. Noncommutative deformation appears to affect only the radial part, with the angular part being solved by the same spin $\frac{1}{2}$ spherical harmonics as in the case of Dirac equation for the hydrogen atom in flat or Schwarzschild case. At the end, we utilise the general properties of the fermionic solutions deduced here to investigate the problem of their stability. In particular, the issue of superradiance is considered as related to the solutions of the Dirac equation in RN spacetime, and especially the impact of noncommutative deformation on the effect of superradiance is addressed.

II. PRELIMINARY SETTINGS

A solution to Einstein equations representing a charged non-rotating black hole with mass M and charge Q is given by the Reissner–Nordström (RN) metric

$$ds^2 = \left(1 - \frac{2MG}{r} + \frac{Q^2G}{r^2}\right)dt^2 - \frac{dr^2}{1 - \frac{2MG}{r} + \frac{Q^2G}{r^2}} - r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (1)$$

Being static and spherically symmetric, the spacetime of RN black hole has four Killing vectors, among which ∂_t and ∂_ϕ are included, and t and ϕ are the time and polar variables of the spherical coordinate system $x^\mu = (t, r, \theta, \phi)$.

In the previous paper [62] we have introduced a semiclassical model describing a charged NC scalar field $\hat{\Phi}$ and NC $U(1)$ gauge field \hat{A} on a classical gravitational background of RN type. By semiclassical we mean that while the gravitational field in this model was assumed to be a classical degree of freedom (i.e. not deformed by noncommutativity), the scalar and gauge field propagating in that classical gravitational background were assumed to be affected by noncommutative nature of spacetime. In a sense, we are therefore dealing with a situation where the scalar and gauge field are quantized and gravitational field is not. It is however important to stress that the gauge and scalar field are not quantized in a sense of quantum field theory.

The model was built by using deformation quantization techniques based on Drinfeld twist operator and the explicit twist operator that was used in the construction was the so called angular twist operator [62, 63]

$$\begin{aligned} \mathcal{F} &= e^{-\frac{i}{2}\theta^{\alpha\beta}\partial_\alpha\otimes\partial_\beta} \\ &= e^{-\frac{ia}{2}(\partial_t\otimes\partial_\phi - \partial_\phi\otimes\partial_t)}, \end{aligned} \quad (2)$$

with $\alpha, \beta = t, r, \theta, \phi$ and $\theta^{t\phi} = -\theta^{\phi t} = a$ as the only non-zero components of the deformation tensor $\theta^{\alpha\beta}$. The small constant parameter a is the deformation parameter that sets up the NC scale, commonly related to the Planck length. This twist operator is a Killing twist, since it is built from the vector fields that are actually Killing vectors for the metric (1). In this way it is ensured that the geometry (1) stays unaffected by the deformation as the twist (2) does not act on the RN metric.

The star product, the wedge star product between forms, the coproduct and other structural maps of the related symmetry algebra can all be obtained from the twist operator (2). In particular, the star product between functions is given by

$$\begin{aligned} f \star g &= \mu \circ \mathcal{F}^{-1} (f \otimes g) = \mu \{ e^{\frac{ia}{2}(\partial_t\otimes\partial_\phi - \partial_\phi\otimes\partial_t)} f \otimes g \} \\ &= fg + \frac{ia}{2}(\partial_t f(\partial_\phi g) - \partial_t g(\partial_\phi f)) + \mathcal{O}(a^2), \end{aligned} \quad (3)$$

where the map μ represents the usual pointwise multiplication. The remaining ingredients of the differential calculus are described in [62].

III. SPINOR FIELD ON THE NONCOMMUTATIVE RN BACKGROUND: FORMAL ANALYSIS

A massive and charged spinor field Ψ on a fixed gravitational background can be described by the following action

$$S = \int d^4x |e| \bar{\Psi} \left(i\gamma^\mu D_\mu \Psi - m\Psi \right). \quad (4)$$

The mass and the charge of the spinor field Ψ are respectively m and q . The determinant of the vierbein $e^a{}_\mu$ we label with $|e| = \sqrt{-g}$ and the covariant derivative D_μ includes both the spin connection ω_μ and the $U(1)$ gauge field A_μ

$$D_\mu \Psi = \partial_\mu \Psi - \frac{i}{2} \omega_\mu{}^{ab} \Sigma_{ab} \Psi - iqA_\mu \Psi. \quad (5)$$

Matrices Σ^{ab} are the (hermitean) generators of the local Lorentz transformations and they close the Lorentz algebra

$$[\Sigma^{ab}, \Sigma^{cd}] = i(\eta^{ad}\Sigma^{bc} + \eta^{bc}\Sigma^{ad} - \eta^{ac}\Sigma^{bd} - \eta^{bd}\Sigma^{ac}).$$

The spin connection is not an independent field, but a function of $e^a{}_\mu$, calculated from the torsion free condition

$$T_{\mu\nu}^a = \nabla_\mu e^a{}_\nu - \nabla_\nu e^a{}_\mu,$$

with $\nabla_\mu e^a{}_\nu = \partial_\mu e^a{}_\nu + \omega_\mu{}^a{}_b e^b{}_\nu$. For more details on the spin connection, vierbeins and the notation we use, we refer to the beginning of Section V.

The action (4) is invariant under the local $U(1)$ transformations

$$\delta_\alpha \Psi = i\alpha(x)\Psi, \quad \delta_\alpha \bar{\Psi} = -i\bar{\Psi}\alpha(x), \quad \delta_\alpha A_\mu = \frac{1}{q} \partial_\mu \alpha(x). \quad (6)$$

Note that these transformations do not act on the gravitational background, that is on $e^a{}_\mu$ and ω_μ . The action (4) is also invariant under the diffeomorphisms and the local $SO(1,3)$ symmetry. In this paper we use the semiclassical analysis and promote only the $U(1)$ gauge symmetry to the noncommutative $U(1)_\star$ gauge symmetry. In our future work we will lift this approximation and allow for the noncommutative local $SO(1,3)_\star$ symmetry. The equation of motion for the spinor field Ψ is obtained by varying the action (4) with respect to $\bar{\Psi}$ and it is given by

$$i\gamma^\mu \left(\partial_\mu \Psi - i\omega_\mu \Psi - iqA_\mu \Psi \right) - m\Psi = 0. \quad (7)$$

Following the steps from [62], we now introduce an action functional that describes the NC $U(1)_\star$ gauge theory of a charged spinor field on the RN background¹ (1)

$$S_\star = \int d^4x |e| \star \bar{\hat{\Psi}} \star \left(i\gamma^\mu \left(\partial_\mu \hat{\Psi} - i\omega_\mu \star \hat{\Psi} - iq\hat{A}_\mu \star \hat{\Psi} \right) - m\hat{\Psi} \right). \quad (8)$$

Noncommutative fields are now labeled with a $\hat{}$ and the \star -product is given by (3). One can show that this action is invariant under the following infinitesimal $U(1)_\star$ gauge transformations:

$$\begin{aligned} \delta_\star \hat{\Psi} &= i\hat{\Lambda} \star \hat{\Psi}, \\ \delta_\star \hat{A}_\mu &= \partial_\mu \hat{\Lambda} + i(\hat{\Lambda} \star \hat{A}_\mu - \hat{A}_\mu \star \hat{\Lambda}), \\ \delta_\star \omega_\mu &= \delta_\star e^a{}_\mu = 0, \end{aligned} \quad (9)$$

where $\hat{\Lambda}$ is the NC gauge parameter. In particular, note that

$$\delta_\star D_\mu \hat{\Psi} = i\hat{\Lambda} \star D_\mu \hat{\Psi}$$

since the twist (2) does not act on the gravitational field and therefore $\omega_\mu \star \Lambda = \omega_\mu \cdot \Lambda = \Lambda \star \omega_\mu$. To simplify the calculation, from now on we redefine $A_\mu = q\hat{A}_\mu$. Then we use the Seiberg-Witten (SW)-map [15],[26] in order to express NC fields $\hat{\Psi}$ and \hat{A}_μ as functions of the corresponding commutative fields and the deformation parameter a .

¹ Actually, it can be any background with Killing vectors ∂_t and ∂_ϕ .

The SW-map assumes an expansion in orders of the deformation parameter and this expansion is known to all orders for an arbitrary Abelian twist deformation [26],[43],[44] of which the twist (2) is only one example. For the twist operator (2), SW-map gives rise to the following expansions for the fields:

$$\hat{\Psi} = \Psi - \frac{1}{2}\theta^{\rho\sigma}A_\rho(\partial_\sigma\Psi), \quad (10)$$

$$\hat{A}_\mu = A_\mu - \frac{1}{2}\theta^{\rho\sigma}A_\rho(\partial_\sigma A_\mu + F_{\sigma\mu}). \quad (11)$$

The expanded action up to first order in a is given by

$$S_\star = \int d^4x |e|\bar{\Psi}\left(i\gamma^\mu D_\mu\Psi - m\Psi\right) + \frac{1}{2}\theta^{\alpha\beta}\left(-iF_{\mu\alpha}\bar{\Psi}\gamma^\mu D_\beta^{U(1)}\Psi - \frac{i}{2}\bar{\Psi}\gamma^\mu\omega_\mu F_{\alpha\beta}\Psi - \frac{1}{2}F_{\alpha\beta}\bar{\Psi}(i\gamma^\mu D_\mu^{U(1)}\Psi - m\Psi)\right).$$

Remembering that $F_{\alpha\beta} = \partial_\alpha A_\beta - \partial_\beta A_\alpha$ and that the only non-zero component of A_μ is $A_t = -\frac{qQ}{r}$, the only non-zero component of $F_{\alpha\beta}$ is $F_{rt} = \frac{qQ}{r^2}$. This leads to a simplified NC action

$$S_\star = \int d^4x |e|\bar{\Psi}\left(i\gamma^\mu D_\mu\Psi - m\Psi\right) - \frac{i}{2}\theta^{\alpha\beta}\bar{\Psi}F_{\mu\alpha}\gamma^\mu D_\beta^{U(1)}\Psi \quad (12)$$

and the corresponding equation of motion for the spinor Ψ

$$i\gamma^\mu\left(\partial_\mu\Psi - i\omega_\mu\Psi - iA_\mu\Psi\right) - m\Psi - \frac{ia}{2}F_{rt}\gamma^r\partial_\phi\Psi = 0. \quad (13)$$

Inserting the explicit expressions for F_{rt} and $\gamma^r = e_a{}^r\gamma^a$, this equation reduces to

$$i\gamma^\mu\left(\partial_\mu\Psi - i\omega_\mu\Psi - iA_\mu\Psi\right) - m\Psi - \frac{ia}{2}\frac{qQ}{r^2}\sqrt{f}\gamma^1\partial_\phi\Psi = 0. \quad (14)$$

The semiclassical approximation manifests itself in (12) in the following way: the covariant derivative $D_\mu\Psi = \partial_\mu\Psi - iA_\mu\Psi - i\omega_\mu\Psi$ includes both the electromagnetic ($U(1)$) and the gravitational part, while the covariant derivative $D_\beta^{U(1)}\Psi = \partial_\beta\Psi - iA_\beta\Psi$ has only the electromagnetic part. In the NC correction only the $U(1)$ part appears.

IV. EFFECTIVE METRIC FROM NONCOMMUTATIVE DUALITY

In our previous paper [62] we analyzed the propagation of the NC scalar field in the RN background. The equation of motion governing the evolution of the scalar field is given by

$$\left(\frac{1}{f}\partial_t^2 - \Delta + (1-f)\partial_r^2 + \frac{2MG}{r^2}\partial_r + 2iqQ\frac{1}{rf}\partial_t - \frac{q^2Q^2}{r^2f} - \mu^2\right)\Phi + \frac{aqQ}{r^3}\left(\frac{MG}{r} - \frac{GQ^2}{r^2}\right)\partial_\phi + rf\partial_r\partial_\phi\Phi = 0, \quad (15)$$

with $f = 1 - \frac{2MG}{r} + \frac{Q^2G}{r^2}$. In [?] an equivalence between a NC scalar field on a non-rotating BTZ background and the commutative scalar field on the rotating BTZ background was established. We will follow that idea here and try to understand if there is an effective description of the NC scalar field on the RN background.

Equation (15) can be symbolically written in terms of an extended Klein-Gordon operator extended to include a coupling to a gauge field

$$(\square_{g'} + \mathcal{O}(a))\Phi \equiv \left(g'^{\mu\nu}(\nabla'_\mu - iA_\mu)(\nabla'_\nu - iA_\nu) + \mathcal{O}(a)\right)\Phi = 0. \quad (16)$$

Corrections are included in $\mathcal{O}(a)$ that is a generic expression and it designates symbolically a whole set of correction terms in the equation (15) that are induced by the noncommutativity and are therefore linear in NC parameter a . Likewise, ∇'_μ is a covariant derivative with respect to the metric $g'^{\mu\nu}$ (1) and $\square_{g'}$ is the Klein-Gordon operator for

² $\nabla'_\mu A^\nu = \partial_\mu A^\nu + \Gamma'^{\lambda}_{\mu\nu}A^\lambda$ and $\nabla'_\mu A_\nu = \partial_\mu A_\nu - \Gamma'^{\lambda}_{\mu\nu}A_\lambda$.

the metric $g'_{\mu\nu}$. Note that by switching off a noncommutativity by letting $a \rightarrow 0$, all corrections that scale with a disappear, and the KG equation reduces to

$$\square_{g'}\Phi \equiv g'^{\mu\nu}(\nabla'_\mu - iA_\mu)(\nabla'_\nu - iA_\nu)\Phi = \frac{1}{\sqrt{-g'}}(\partial_\mu - iA_\mu)\left(\sqrt{-g'}g'^{\mu\nu}(\partial_\nu - iA_\nu)\right)\Phi = 0. \quad (17)$$

At this stage one is naturally led to ponder over a possibility that the terms in (15) which scale linearly with the NC parameter a can actually be soaked up by the already present KG operator $\square_{g'}$ to yield a KG operator \square_g with a redefined metric that has managed to absorb within itself noncommutative features of the original problem. More concisely, the question to be posed is if there exists a metric which is able to meet the requirement

$$\begin{aligned} (\square_{g'} + \mathcal{O}(a))\Phi &\equiv \left(g'^{\mu\nu}(\nabla'_\mu - iA_\mu)(\nabla'_\nu - iA_\nu) + \mathcal{O}(a)\right)\Phi \\ &= \square_g\Phi = g^{\mu\nu}(\nabla_\mu - iA_\mu)(\nabla_\nu - iA_\nu)\Phi = \frac{1}{\sqrt{-g}}(\partial_\mu - iA_\mu)\left(\sqrt{-g}g^{\mu\nu}(\partial_\nu - iA_\nu)\right)\Phi = 0. \end{aligned} \quad (18)$$

where ∇_μ is a covariant derivative with respect to the new, effective metric $g_{\mu\nu}$. We point out that the gauge potential did not change upon switching to a new setting and rewriting dynamics of the system in terms of the effective metric.

In order to find the metric tensor which satisfies the requirement (18), one may try with the following ansatz

$$g_{\mu\nu} = \begin{pmatrix} f & 0 & 0 & 0 \\ 0 & -\frac{1}{f} & 0 & g_{r\phi} \\ 0 & 0 & -r^2 & 0 \\ 0 & g_{r\phi} & 0 & -r^2 \sin\theta \end{pmatrix}, \quad (19)$$

The novel nonvanishing entry $g_{r\phi}$ is assumed to depend only on variables r and θ , since we expect that ∂_t and ∂_ϕ are Killing vectors for the effective metric as well. Moreover, it is assumed to be at least linear in NC parameter a , $g_{r\phi} \sim \mathcal{O}(a)$ since the effective metric $g_{\mu\nu}$ has to reduce to the original RN metric $g'_{\mu\nu}$ in the limiting case $a \rightarrow 0$. The inverse of the metric tensor (19) has nonvanishing entries at the same places

$$g^{\mu\nu} = \begin{pmatrix} \frac{1}{f} & 0 & 0 & 0 \\ 0 & -f + \frac{f^2 g_{r\phi}^2}{f g_{r\phi}^2 - r^2 \sin^2 \theta} & 0 & \frac{f g_{r\phi}}{f g_{r\phi}^2 - r^2 \sin^2 \theta} \\ 0 & 0 & -\frac{1}{r^2} & 0 \\ 0 & \frac{f g_{r\phi}}{f g_{r\phi}^2 - r^2 \sin^2 \theta} & 0 & \frac{1}{f g_{r\phi}^2 - r^2 \sin^2 \theta} \end{pmatrix}, \quad (20)$$

It can be seen that while off-diagonal elements have a leading correction term that is linear in a , the diagonal elements g^{rr} and $g^{\phi\phi}$ have a leading correction term that is quadratic in a . Likewise, the determinant and the square-root of the determinant of the effective metric (19) have a leading correction term that is quadratic in the NC parameter a , $\sqrt{-g} = r^2 \sin\theta + \mathcal{O}(a^2)$. These observations will have a crucial role in the subsequent analysis, whose aim is to deduce the metric $g_{\mu\nu}$, satisfying the requirement (18).

The form of the metric (19) dictates which terms are going to survive after the equation (18) is written out explicitly

$$\begin{aligned} \square_g\Phi &= \frac{1}{\sqrt{-g}}(\partial_\mu - iA_\mu)\left(\sqrt{-g}g^{\mu\nu}(\partial_\nu - iA_\nu)\right)\Phi \\ &= \frac{1}{\sqrt{-g}}\left[(\partial_t - iA_t)\left(\sqrt{-g}g^{tt}(\partial_t - iA_t)\right) + (\partial_r - iA_r)\left(\sqrt{-g}g^{rr}(\partial_r - iA_r)\right) \right. \\ &\quad + (\partial_r - iA_r)\left(\sqrt{-g}g^{r\phi}(\partial_\phi - iA_\phi)\right) + (\partial_\theta - iA_\theta)\left(\sqrt{-g}g^{\theta\theta}(\partial_\theta - iA_\theta)\right) \\ &\quad \left. + (\partial_\phi - iA_\phi)\left(\sqrt{-g}g^{\phi r}(\partial_r - iA_r)\right) + (\partial_\phi - iA_\phi)\left(\sqrt{-g}g^{\phi\phi}(\partial_\phi - iA_\phi)\right)\right]\Phi. \end{aligned}$$

Taking into account the fact that the gauge potential has only time component, one finds that the equation of motion (18) further boils down to

$$\begin{aligned} \frac{1}{f}\left[\partial_t^2\Phi - 2iA_t\partial_t\Phi\right] &+ \frac{1}{\sqrt{-g}}\left[\partial_r(\sqrt{-g}g^{rr})\right]\partial_r\Phi + g^{rr}\partial_r^2\Phi + \frac{1}{\sqrt{-g}}\left[\partial_r(\sqrt{-g}g^{r\phi})\right]\partial_\phi\Phi \\ &+ 2g^{r\phi}\partial_r\partial_\phi\Phi + \frac{1}{\sqrt{-g}}\left[\partial_\theta(\sqrt{-g}g^{\theta\theta})\right]\partial_\theta\Phi + g^{\theta\theta}\partial_\theta^2\Phi + g^{\phi\phi}\partial_\phi^2\Phi = 0. \end{aligned}$$

Focusing only on terms in the above equation that are at most linear in a , and stacking it up against the equation (15) leads to the following two relations:

$$\begin{aligned} \frac{aqQ}{r^3} \left(\frac{MG}{r} - \frac{GQ^2}{r^2} \right) \partial_\phi \Phi &= \frac{1}{\sqrt{-g}} \left[\partial_r (\sqrt{-g} g^{r\phi}) \right] \partial_\phi \Phi, \\ \frac{aqQ}{r^2} f \partial_r \partial_\phi \Phi &= 2g^{r\phi} \partial_r \partial_\phi \Phi. \end{aligned} \quad (21)$$

The solution to this set of relations, which is consistent with the requirement $g^{r\phi} = \frac{f g_{r\phi}}{f g_{r\phi}^2 - r^2 \sin^2 \theta}$, finally gives for the dual effective metric

$$g_{\mu\nu} = \begin{pmatrix} f & 0 & 0 & 0 \\ 0 & -\frac{1}{f} & 0 & -\frac{aqQ}{2} \sin^2 \theta \\ 0 & 0 & -r^2 & 0 \\ 0 & -\frac{aqQ}{2} \sin^2 \theta & 0 & -r^2 \sin^2 \theta \end{pmatrix} \quad (22)$$

and for its inverse metric

$$g^{\mu\nu} = \begin{pmatrix} \frac{1}{f} & 0 & 0 & 0 \\ 0 & -f & 0 & \frac{aqQ}{2r^2} f \\ 0 & 0 & -\frac{1}{r^2} & 0 \\ 0 & \frac{aqQ}{2r^2} f & 0 & -\frac{1}{r^2 \sin^2 \theta} \end{pmatrix}. \quad (23)$$

Note that we demand $g_{\mu\nu} g^{\nu\rho} = \delta_\mu^\rho + \mathcal{O}(a^2)$.

We have thus shown that the equation of motion for a charged NC scalar field in a classical RN background, coupled to NC $U(1)$ gauge field may be rewritten in terms of the equation of motion governing behaviour of a charged commutative scalar field (having the same charge q as its NC counterpart), propagating in a modified RN geometry

$$ds^2 = \left(1 - \frac{2MG}{r} + \frac{Q^2 G}{r^2} \right) dt^2 - \frac{dr^2}{1 - \frac{2MG}{r} + \frac{Q^2 G}{r^2}} - aqQ \sin^2 \theta dr d\phi - r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (24)$$

It appears that new, effective metric (24) acquires an additional off-diagonal term which is induced purely by non-commutative nature of spacetime. This feature comes into play only in the presence of charged matter. Unlike in the case of scalar field in the (NC) BTZ background, in this case the effective metric cannot be interpreted as a metric of a background rotating RN (or any other) geometry.

V. SPINOR FIELDS ON THE NONCOMMUTATIVE RN BACKGROUND: EFFECTIVE APPROACH

Having found the effective metric (24), let us now investigate the propagation of a charged massive spinor field ψ in this geometry. In particular, it is interesting to see if this effective approach agrees with the more rigorous approach from Section II. The Dirac equation in a curved background given by the effective noncommutative (NC) metric

$$(i\gamma^a \nabla_a - m)\Psi = 0, \quad (25)$$

where the Latin indices such as a , ($a = 0, 1, 2, 3$) refer to intrinsic coordinates and γ^a are the standard flat space Dirac gamma matrices, $\{\gamma_a, \gamma_b\} = 2\eta_{ab}$, where

$$\eta_{ab} = \eta^{ab} = \begin{pmatrix} +1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}. \quad (26)$$

If in addition, the spinor field is charged, this gives rise to a Dirac equation in which the gauge potential A_μ is minimally coupled to a Dirac operator on a curved background

$$\left(i\gamma^a (\nabla_a - iA_a) - m \right) \Psi = \left(i\gamma^a e_a{}^\mu (\nabla_\mu - iA_\mu) - m \right) \Psi = 0. \quad (27)$$

The gravitational covariant derivative ∇_μ is defined as $\nabla_\mu \Psi = \partial_\mu \Psi - \frac{i}{2} \omega_\mu^{ab} \Sigma_{ab} \Psi$. The Dirac operator $\gamma^a \nabla_a$ on a curved space is introduced in terms of tetrads (vierbeins) $e^a{}_\mu$ and their inverse $e_a{}^\mu$, satisfying $e^a{}_\mu e_a{}^\nu = \delta^\nu{}_\mu$ and $e^a{}_\mu e_b{}^\mu = \delta^a{}_b$. Tetrads written in components are $e^a{}_\mu = (e^a{}_t, e^a{}_r, e^a{}_\theta, e^a{}_\phi)$ and $e_a{}^\mu = (e_0{}^\mu, e_1{}^\mu, e_2{}^\mu, e_3{}^\mu)$. They also satisfy $g_{\mu\nu} = e^a{}_\mu e^b{}_\nu \eta_{ab}$ and $g^{\mu\nu} = e_a{}^\mu e_b{}^\nu \eta^{ab}$. In what follows we use the setting defined in [113] with the vierbein frame chosen to be

$$e^a{}_\mu = \begin{pmatrix} \sqrt{f} & 0 & 0 & 0 \\ 0 & \frac{1}{\sqrt{f}} & 0 & 0 \\ 0 & 0 & r & 0 \\ 0 & \frac{aqQ}{2r} \sin \theta & 0 & r \sin \theta \end{pmatrix} \quad (28)$$

with the corresponding inverse matrix

$$e_a{}^\mu = \begin{pmatrix} \frac{1}{\sqrt{f}} & 0 & 0 & 0 \\ 0 & \sqrt{f} & 0 & -\frac{aqQ}{2r^2} \sqrt{f} \\ 0 & 0 & \frac{1}{r} & 0 \\ 0 & 0 & 0 & \frac{1}{r \sin \theta} \end{pmatrix}. \quad (29)$$

The representation of gamma matrices is

$$\begin{aligned} \gamma^0 &= i\tilde{\gamma}^0 = i \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, & \gamma^1 &= i\tilde{\gamma}^3 = i \begin{pmatrix} 0 & \sigma_3 \\ -\sigma_3 & 0 \end{pmatrix}, \\ \gamma^2 &= i\tilde{\gamma}^1 = i \begin{pmatrix} 0 & \sigma_1 \\ -\sigma_1 & 0 \end{pmatrix}, & \gamma^3 &= i\tilde{\gamma}^2 = i \begin{pmatrix} 0 & \sigma_2 \\ -\sigma_2 & 0 \end{pmatrix}, \end{aligned} \quad (30)$$

where $\tilde{\gamma}^0, \tilde{\gamma}^1, \tilde{\gamma}^2$ and $\tilde{\gamma}^3$ are gamma matrices in chiral (Weyl) representation, while σ_i , ($i = 1, 2, 3$) are the usual Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (31)$$

By writing out a detailed structure of the covariant derivative ∇_a , the Dirac equation (27) takes the form³

$$\left[i\gamma^a e_a{}^\mu \left(\partial_\mu + \frac{i}{2} \omega_\mu{}^{cd} \Sigma_{cd} - iA_\mu \right) - m \right] \Psi = 0. \quad (32)$$

Here $\Sigma_{cd} = \frac{i}{4} [\gamma_c, \gamma_d]$ and the coefficients of the spin connection $\omega_\mu{}^{ab}$ are given by

$$\begin{aligned} \omega_\mu{}^{ab} &= e^a{}_\nu \eta^{bc} \partial_\mu e_c{}^\nu + e^a{}_\nu \eta^{bc} e_c{}^\lambda \Gamma^\nu{}_{\mu\lambda} \\ &= \frac{1}{2} e^{a\nu} \left(\partial_\mu e^b{}_\nu - \partial_\nu e^b{}_\mu \right) - \frac{1}{2} e^{b\nu} \left(\partial_\mu e^a{}_\nu - \partial_\nu e^a{}_\mu \right) - \frac{1}{2} e^{a\rho} e^{b\sigma} \left(\partial_\rho e_{c\sigma} - \partial_\sigma e_{c\rho} \right) e^c{}_\mu, \end{aligned}$$

where $\Gamma^\nu{}_{\mu\lambda} = \frac{1}{2} g^{\nu\delta} \left(\partial_\mu g_{\delta\lambda} + \partial_\lambda g_{\mu\delta} - \partial_\delta g_{\mu\lambda} \right)$ are the coefficients of the affine connection. Note that $\omega_\mu{}^{ab} = -\omega_\mu{}^{ba}$.

With the tetrads given in (28), one gets that the only non zero components of the spin connection are

$$\begin{aligned} \omega_t{}^{01} &= -\omega_t{}^{10} = -\frac{Mr-Q^2}{r^3}, & \omega_\theta{}^{12} &= -\omega_\theta{}^{21} = \sqrt{f}, \\ \omega_\phi{}^{13} &= -\omega_\phi{}^{31} = \sqrt{f} \sin \theta, & \omega_\phi{}^{23} &= -\omega_\phi{}^{32} = \cos \theta, \\ \omega_r{}^{23} &= -\omega_r{}^{32} = \frac{aqQ}{2r^2} \cos \theta, & \omega_r{}^{13} &= -\omega_r{}^{31} = \frac{aqQ\sqrt{f}}{2r^2} \sin \theta. \end{aligned} \quad (33)$$

³ Since implementation of (30) as our representation of γ -matrices involves a flip between their hermitian and antihermitian properties, the covariant derivative gets changed, $\nabla_\mu = \partial_\mu - \frac{i}{2} \omega_\mu{}^{cd} \frac{i}{4} [\gamma_c, \gamma_d] \rightarrow \nabla_\mu = \partial_\mu - \frac{1}{2} \omega_\mu{}^{cd} \frac{1}{4} [\gamma_c, \gamma_d]$, which amounts to changing the sign in the covariant derivative in front of the spin part, $\nabla_\mu = \partial_\mu + \frac{i}{2} \omega_\mu{}^{cd} \Sigma_{cd} = \partial_\mu + \frac{i}{2} \omega_\mu{}^{cd} \frac{i}{4} [\gamma_c, \gamma_d]$.

In subsequent analysis we will also use the sums $\omega_\mu{}^{cd}\Sigma_{cd}$:

$$\begin{aligned}
\omega_t{}^{cd}\Sigma_{cd} &= 2\omega_t{}^{01}\Sigma_{01} = -2\frac{Mr - Q^2}{r^3}i[\gamma_0, \gamma_1] = -i\frac{Mr - Q^2}{r^3}\begin{pmatrix} -\sigma_3 & 0 \\ 0 & \sigma_3 \end{pmatrix}, \\
\omega_r{}^{cd}\Sigma_{cd} &= 2\omega_r{}^{23}\Sigma_{23} + 2\omega_r{}^{13}\Sigma_{13} = -\frac{aqQ}{2r^2}\cos\theta\begin{pmatrix} \sigma_3 & 0 \\ 0 & \sigma_3 \end{pmatrix} + \frac{aqQ\sqrt{f}}{2r^2}\sin\theta\begin{pmatrix} \sigma_1 & 0 \\ 0 & \sigma_1 \end{pmatrix}, \\
\omega_\theta{}^{cd}\Sigma_{cd} &= 2\omega_\theta{}^{12}\Sigma_{12} = -\sqrt{f}\begin{pmatrix} \sigma_2 & 0 \\ 0 & \sigma_2 \end{pmatrix}, \\
\omega_\phi{}^{cd}\Sigma_{cd} &= 2\omega_\phi{}^{13}\Sigma_{13} + 2\omega_\phi{}^{23}\Sigma_{23} = \sqrt{f}\sin\theta\begin{pmatrix} \sigma_1 & 0 \\ 0 & \sigma_1 \end{pmatrix} - \cos\theta\begin{pmatrix} \sigma_3 & 0 \\ 0 & \sigma_3 \end{pmatrix},
\end{aligned} \tag{34}$$

Inserting these into (32) leads to the Dirac equation

$$\begin{aligned}
&\left[i\gamma^0 e_0{}^t \left(\partial_t - iA_t + \frac{i}{2}\omega_t{}^{cd}\Sigma_{cd} \right) + i\gamma^1 e_1{}^r \left(\partial_r - iA_r + \frac{i}{2}\omega_r{}^{cd}\Sigma_{cd} \right) + i\gamma^2 e_2{}^\theta \left(\partial_\theta - iA_\theta + \frac{i}{2}\omega_\theta{}^{cd}\Sigma_{cd} \right) \right. \\
&\left. + i\gamma^3 e_3{}^\phi \left(\partial_\phi - iA_\phi + \frac{i}{2}\omega_\phi{}^{cd}\Sigma_{cd} \right) + i\gamma^3 e_3{}^\phi \left(\partial_\phi - iA_\phi + \frac{i}{2}\omega_\phi{}^{cd}\Sigma_{cd} \right) - m \right] \Psi = 0.
\end{aligned} \tag{35}$$

With the spinor field Ψ written in terms of two two-component spinors Ψ_1 and Ψ_2 , namely $\Psi = \begin{pmatrix} \Psi_1 \\ \Psi_2 \end{pmatrix}$ and the gauge potential $A_\mu = (A_t, \vec{A}) = (-\frac{aQ}{r}, \vec{0})$, the equation (35) splits into two two-component equations

$$\begin{aligned}
&\left[-\frac{1}{\sqrt{f}}\mathbb{1}\partial_t - \sqrt{f}\sigma_3\partial_r - \frac{1}{2}\frac{Mr - Q^2}{r^3}\frac{1}{\sqrt{f}}\sigma_3 - \frac{\sqrt{f}}{r}\sigma_3 - \frac{1}{r}\sigma_1\partial_\theta \right. \\
&\quad \left. + \frac{aqQ}{2r^2}\sqrt{f}\sigma_3\partial_\phi - \frac{1}{r\sin\theta}\sigma_2\partial_\phi - \frac{1}{2r}\cot\theta\sigma_1 - \frac{iqQ}{r\sqrt{f}}\mathbb{1} \right] \Psi_2 - m\mathbb{1}\Psi_1 = 0, \\
&\left[-\frac{1}{\sqrt{f}}\mathbb{1}\partial_t + \frac{1}{2}\frac{Mr - Q^2}{r^3}\frac{1}{\sqrt{f}}\sigma_3 + \sqrt{f}\sigma_3\partial_r + \frac{\sqrt{f}}{r}\sigma_3 + \frac{1}{r}\sigma_1\partial_\theta \right. \\
&\quad \left. - \frac{aqQ}{2r^2}\sqrt{f}\sigma_3\partial_\phi + \frac{1}{r\sin\theta}\sigma_2\partial_\phi + \frac{1}{2r}\cot\theta\sigma_1 - \frac{iqQ}{r\sqrt{f}}\mathbb{1} \right] \Psi_1 - m\mathbb{1}\Psi_2 = 0.
\end{aligned} \tag{36}$$

We see that these equations have the same form as the equation (14). The only NC correction is of the form $\frac{aqQ}{2r^2}\sqrt{f}\gamma^1\partial_\phi\Psi$. Therefore we can conclude that the rigorous approach of the NC gauge theory and the SW expansion described in Section III and the effective approach described here lead to the same result. However, two comments are in order. Firstly, our results are valid up to first order in the deformation parameter a . Secondly, the result in Section III was deduced using the semiclassical approximation. In our future work we plan to investigate if this duality holds more generally.

VI. DISCUSSION AND OUTLOOK

Let us analyze the equation (36) in more detail. This equation can be used to study various effects, such as NC spinor bound states or quasinormal modes in the RN background.

In order to solve (36) for the wavefunction $\Psi \equiv \Psi(t, r, \theta, \phi)$, we follow [113] and take the ansatz

$$\Psi = \begin{pmatrix} \Psi_1 \\ \Psi_2 \end{pmatrix} = \begin{pmatrix} \Psi_1^{(1)} \\ \Psi_1^{(2)} \\ \Psi_2^{(1)} \\ \Psi_2^{(2)} \end{pmatrix} = \frac{1}{r}f^{-1/4}\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = (r^4 f)^{-1/4}\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \left(r^4 - 2Mr^3 + Q^2 r^2 \right)^{-1/4} \begin{pmatrix} \psi_1^{(1)} \\ \psi_1^{(2)} \\ \psi_2^{(1)} \\ \psi_2^{(2)} \end{pmatrix}. \tag{37}$$

After plugging this ansatz into (36) and performing some simplifications, the set of equations (36) reduces to

$$\begin{aligned}
&\left[-\frac{r}{\sqrt{f}}\mathbb{1}\partial_t - r\sqrt{f}\sigma_3\partial_r - \sigma_1\partial_\theta - \frac{1}{\sin\theta}\sigma_2\partial_\phi - \frac{1}{2}\cot\theta\sigma_1 + \frac{aqQ}{2r}\sqrt{f}\sigma_3\partial_\phi - \frac{iqQ}{\sqrt{f}}\mathbb{1} \right] \psi_2 - mr\mathbb{1}\psi_1 = 0, \\
&\left[-\frac{r}{\sqrt{f}}\mathbb{1}\partial_t + r\sqrt{f}\sigma_3\partial_r + \sigma_1\partial_\theta + \frac{1}{\sin\theta}\sigma_2\partial_\phi + \frac{1}{2}\cot\theta\sigma_1 - \frac{aqQ}{2r}\sqrt{f}\sigma_3\partial_\phi - \frac{iqQ}{\sqrt{f}}\mathbb{1} \right] \psi_1 - mr\mathbb{1}\psi_2 = 0.
\end{aligned} \tag{38}$$

We further make a factorization of the spinor wavefunctions ψ_1 and ψ_2 according to

$$\begin{aligned}\psi_1 &\equiv \psi_1(t, r, \theta, \phi) = e^{i(\nu\phi - \omega t)} \begin{pmatrix} \psi_1^{(1)}(r, \theta) \\ \psi_1^{(2)}(r, \theta) \end{pmatrix} = e^{i(\nu\phi - \omega t)} \begin{pmatrix} -R_2(r)S_1(\theta) \\ -R_1(r)S_2(\theta) \end{pmatrix}, \\ \psi_2 &\equiv \psi_2(t, r, \theta, \phi) = e^{i(\nu\phi - \omega t)} \begin{pmatrix} \psi_2^{(1)}(r, \theta) \\ \psi_2^{(2)}(r, \theta) \end{pmatrix} = e^{i(\nu\phi - \omega t)} \begin{pmatrix} R_1(r)S_1(\theta) \\ R_2(r)S_2(\theta) \end{pmatrix},\end{aligned}\quad (39)$$

where ω and ν are respectively energy and projection of the angular momentum of the spin 1/2 particle. The first step in utilizing the factorization (39), which includes a separation of the azimuthal and time variables, gives rise to the set of two 2-component equations

$$\begin{aligned}\left[\frac{i\omega r}{\sqrt{f}} \mathbb{1} - r\sqrt{f}\sigma_3\partial_r - \sigma_1\partial_\theta - \frac{i\nu}{\sin\theta}\sigma_2 - \frac{1}{2}\cot\theta\sigma_1 + i\nu\frac{aqQ}{2r}\sqrt{f}\sigma_3 - \frac{iqQ}{\sqrt{f}}\mathbb{1} \right] \psi_2(r, \theta) - mr\mathbb{1}\psi_1(r, \theta) &= 0, \\ \left[\frac{i\omega r}{\sqrt{f}} \mathbb{1} + r\sqrt{f}\sigma_3\partial_r + \sigma_1\partial_\theta + \frac{i\nu}{\sin\theta}\sigma_2 + \frac{1}{2}\cot\theta\sigma_1 - i\nu\frac{aqQ}{2r}\sqrt{f}\sigma_3 - \frac{iqQ}{\sqrt{f}}\mathbb{1} \right] \psi_1(r, \theta) - mr\mathbb{1}\psi_2(r, \theta) &= 0.\end{aligned}\quad (40)$$

The second step, which involves a separation of the radial and polar angle variables, leads to the set of four coupled partial differential equations

$$\begin{aligned}\frac{i\omega r}{\sqrt{f}}R_1S_1 - r\sqrt{f}(\partial_r R_1)S_1 - R_2\partial_\theta S_2 - \frac{i\nu}{\sin\theta}(-i)R_2S_2 - \frac{1}{2}\cot\theta R_2S_2 \\ + i\nu\frac{aqQ}{2r}\sqrt{f}R_1S_1 + mrR_2S_1 - \frac{iqQ}{\sqrt{f}}R_1S_1 &= 0, \\ \frac{i\omega r}{\sqrt{f}}R_2S_2 + r\sqrt{f}(\partial_r R_2)S_2 - R_1\partial_\theta S_1 - \frac{i\nu}{\sin\theta}iR_1S_1 - \frac{1}{2}\cot\theta R_1S_1 \\ - i\nu\frac{aqQ}{2r}\sqrt{f}R_2S_2 + mrR_1S_2 - \frac{iqQ}{\sqrt{f}}R_2S_2 &= 0, \\ -\frac{i\omega r}{\sqrt{f}}R_2S_1 - r\sqrt{f}(\partial_r R_2)S_1 - R_1\partial_\theta S_2 - \frac{i\nu}{\sin\theta}(-i)R_1S_2 - \frac{1}{2}\cot\theta R_1S_2 \\ + i\nu\frac{aqQ}{2r}\sqrt{f}R_2S_1 - mrR_1S_1 + \frac{iqQ}{\sqrt{f}}R_2S_1 &= 0, \\ -\frac{i\omega r}{\sqrt{f}}R_1S_2 + r\sqrt{f}(\partial_r R_1)S_2 - R_2\partial_\theta S_1 - \frac{i\nu}{\sin\theta}iR_2S_1 - \frac{1}{2}\cot\theta R_2S_1 \\ - i\nu\frac{aqQ}{2r}\sqrt{f}R_1S_2 - mrR_2S_2 + \frac{iqQ}{\sqrt{f}}R_1S_2 &= 0.\end{aligned}\quad (41)$$

After dividing above equations respectively with R_2S_1 , R_1S_2 , R_1S_1 , R_2S_2 one finds that this system of equations is completely separable,

$$\begin{aligned}\frac{i\omega r}{\sqrt{f}}\frac{R_1}{R_2} - r\sqrt{f}\frac{\partial_r R_1}{R_2} + i\nu\frac{aqQ}{2r}\sqrt{f}\frac{R_1}{R_2} - \frac{iqQ}{\sqrt{f}}\frac{R_1}{R_2} + mr &= \frac{\partial_\theta S_2}{S_1} + \frac{\nu}{\sin\theta}\frac{S_2}{S_1} + \frac{1}{2}\cot\theta\frac{S_2}{S_1} \equiv \lambda, \\ \frac{i\omega r}{\sqrt{f}}\frac{R_2}{R_1} + r\sqrt{f}\frac{\partial_r R_2}{R_1} - i\nu\frac{aqQ}{2r}\sqrt{f}\frac{R_2}{R_1} - \frac{iqQ}{\sqrt{f}}\frac{R_2}{R_1} + mr &= \frac{\partial_\theta S_1}{S_2} - \frac{\nu}{\sin\theta}\frac{S_1}{S_2} + \frac{1}{2}\cot\theta\frac{S_1}{S_2} \equiv \lambda_1, \\ -\frac{i\omega r}{\sqrt{f}}\frac{R_2}{R_1} - r\sqrt{f}\frac{\partial_r R_2}{R_1} + i\nu\frac{aqQ}{2r}\sqrt{f}\frac{R_2}{R_1} + \frac{iqQ}{\sqrt{f}}\frac{R_2}{R_1} - mr &= \frac{\partial_\theta S_2}{S_1} + \frac{\nu}{\sin\theta}\frac{S_2}{S_1} + \frac{1}{2}\cot\theta\frac{S_2}{S_1} = \lambda, \\ -\frac{i\omega r}{\sqrt{f}}\frac{R_1}{R_2} + r\sqrt{f}\frac{\partial_r R_1}{R_2} - i\nu\frac{aqQ}{2r}\sqrt{f}\frac{R_1}{R_2} + \frac{iqQ}{\sqrt{f}}\frac{R_1}{R_2} - mr &= \frac{\partial_\theta S_1}{S_2} - \frac{\nu}{\sin\theta}\frac{S_1}{S_2} + \frac{1}{2}\cot\theta\frac{S_1}{S_2} = \lambda_1.\end{aligned}\quad (42)$$

Moreover, it is easily seen that two separation constants λ and λ_1 , which have appeared in a process of separation are not mutually independent, but subject to the requirement $\lambda = -\lambda_1$. In effect, the system of equations (42) gives rise to two angular equations

$$\begin{aligned}\partial_\theta S_2 + \frac{\nu}{\sin\theta}S_2 + \frac{1}{2}\cot\theta S_2 &= \lambda S_1, \\ \partial_\theta S_1 - \frac{\nu}{\sin\theta}S_1 + \frac{1}{2}\cot\theta S_1 &= -\lambda S_2,\end{aligned}\quad (43)$$

and two radial equations

$$\begin{aligned}\frac{i\omega r}{\sqrt{f}}R_1 - r\sqrt{f}\partial_r R_1 + i\nu\frac{aqQ}{2r}\sqrt{f}R_1 - \frac{iqQ}{\sqrt{f}}R_1 &= (\lambda - mr)R_2, \\ \frac{i\omega r}{\sqrt{f}}R_2 + r\sqrt{f}\partial_r R_2 - i\nu\frac{aqQ}{2r}\sqrt{f}R_2 - \frac{iqQ}{\sqrt{f}}R_2 &= -(\lambda + mr)R_1,\end{aligned}\quad (44)$$

This system of radial equations can be used to study the behaviour of spinor quasinormal modes in the RN background.

This discussion we close with the analysis of the stability of chargeless, but massive fermionic modes. For that purpose we recall that for the bosonic fields on Kerr spacetime there exists a regime in which bosonic modes become unstable, due to superradiant growth [114–118]. Contrary to that, in the same regime where the bosonic modes manifest instability, the fermionic fields on Kerr spacetime under condition of extra slow rotation do not, resulting in them being stable and subject only to a decay [119],[120],[121],[122],[123]. In other words, unlike the equations of motion governing the bosonic fields, the single-particle Dirac equation is not subject to superradiance, and thus all modes decay in that particular regime, which includes a setting where $mM \lesssim (l + \frac{1}{2})$, as well as the limit of slow rotation, $\Omega/M \ll 1^4$. Interestingly, fermionic fields on Schwarzschild or Reissner–Nordström spacetime display somewhat different characteristics, making them more likely to exhibit the effect of superradiance and thus not remain stable. These observations may be drawn by inspecting the bosonic and fermionic modes in question, either by inspecting the imaginary part of their bound state frequencies or by investigating the properties of the flux passing into the horizon and the corresponding conservation law.

Here we set to examine a possibility that a noncommutative deformation of RN spacetime, realized in a form of the effective metric (22), introduces certain changes to the above statements. To start with, let us recall the form of the wave function that solves the Dirac equation

$$\Psi = e^{i(\nu\phi - \omega t)} (r^4 f)^{-1/4} \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = e^{i(\nu\phi - \omega t)} \left(r^4 - 2Mr^3 + Q^2 r^2 \right)^{-1/4} \begin{pmatrix} -R_2(r)S_1(\theta) \\ -R_1(r)S_2(\theta) \\ R_1(r)S_1(\theta) \\ R_2(r)S_2(\theta) \end{pmatrix}. \quad (45)$$

Its corresponding hermitian conjugate is defined as

$$\bar{\Psi} = -\Psi^\dagger \gamma^0 = -i\Psi^\dagger \begin{pmatrix} 0 & \mathbb{1} \\ \mathbb{1} & 0 \end{pmatrix}, \quad (46)$$

and the covariant derivatives that include the spin connection part are given by

$$\begin{aligned}\nabla_\mu \Psi &= \partial_\mu \Psi - \Gamma_\mu \Psi = \partial_\mu \Psi - \frac{1}{4}\omega_\mu{}^{bc}\gamma_b\gamma_c\Psi, \\ \nabla_\mu \bar{\Psi} &= \partial_\mu \bar{\Psi} + \bar{\Psi}\Gamma_\mu = \partial_\mu \bar{\Psi} + \frac{1}{4}\omega_\mu{}^{bc}\bar{\Psi}\gamma_b\gamma_c.\end{aligned}$$

With these quantities at hand, one may define the stress-energy tensor as

$$T_{\mu\nu} = \frac{i}{4} \left[\bar{\Psi}\gamma_\mu\nabla_\nu\Psi + \bar{\Psi}\gamma_\nu\nabla_\mu\Psi - (\nabla_\mu\bar{\Psi})\gamma_\nu\Psi - (\nabla_\nu\bar{\Psi})\gamma_\mu\Psi \right]. \quad (47)$$

For the fermionic field on the Kerr spacetime the form of the solution for the wave function essentially (up to a different prefactor) has the same general form as (45). The radial Dirac current and the corresponding conservation law in this case give rise to the condition

$$\frac{dN}{dt} = \left(|R_1|^2 - |R_2|^2 \right)_{r=r_h} \leq 0, \quad (48)$$

where N is the number density and r_h is the outer horizon radius, signalling the absence of superradiance [113]. For the case considered in this paper, i.e. deformed RN metric (22), the radial component of the Dirac current

⁴ Note that m is the mass of the perturbing field, M is the mass of the black hole, l is the orbital angular momentum number and ΩM is the angular momentum of a black hole.

$J^\mu = \bar{\Psi}\gamma^\mu\Psi = \bar{\Psi}e_a{}^\mu\gamma^a\Psi$ may be shown to have the form

$$\begin{aligned}
J^r &= \bar{\Psi}\gamma^r\Psi = \bar{\Psi}e_a{}^r\gamma^a\Psi \\
&= -i\left(r^4 - 2Mr^3 + Q^2r^2\right)^{-1/2}\sqrt{f}\begin{pmatrix} R_1^*S_1^* & R_2^*S_2^* & -R_2^*S_1^* & -R_1^*S_2^* \end{pmatrix} \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \\ -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix} \begin{pmatrix} -R_2S_1 \\ -R_1S_2 \\ R_1S_1 \\ R_2S_2 \end{pmatrix} \\
&= \frac{1}{r^2}\left(|R_1|^2 - |R_2|^2\right)\left(|S_1|^2 + |S_2|^2\right).
\end{aligned} \tag{49}$$

At first glance, with the result (49) at hand, one might be led to think that the fermionic field on deformed RN spacetime might also exhibit the absence of superradiance. This is due to the fact that the integral of the radial current evaluated at the lower bound, i.e. at the outer horizon $r = r_h$, has a natural interpretation as the flux passing into the horizon and the expression on the rhs of (49) not being positive-definite.

However, a deeper inspection seems not to confirm this conclusion. In fact, a superradiance may be seen as a direct consequence of the second law of black hole thermodynamics. On the other hand, a crucial assumption that underlies the second law of black hole thermodynamics is that the condition $T_{\mu\nu}t^\mu t^\nu \geq 0$ must hold for any time-like vector field, $t_\mu t^\mu > 0$, with $T_{\mu\nu}$ being the stress-energy tensor. This condition is known as the weak energy condition. Therefore, for the purpose of demonstrating that a superradiance is absent in a given system, or at least that it doesn't show up for a particular choice of the system parameters, we would first have to make evident that there exists a sector in the parameter space of that system where the weak energy condition is violated. In order to examine if the weak energy condition might possibly be violated within the context of fermionic field on NC deformed RN background, opening in this way a window for a possible violation of the second law of thermodynamics and a consequent loss of the effect of superradiance within the same context, we take the time-like vector $t^\mu \equiv e_0{}^\mu = (e_0{}^t, e_0{}^r, e_0{}^\theta, e_0{}^\phi) = (\frac{1}{\sqrt{f}}, 0, 0, 0)$, $t_\mu t^\mu > 0$, and first evaluate and then analyse the bilinear form $T_{\mu\nu}t^\mu t^\nu = T_{\mu\nu}e_0{}^\mu e_0{}^\nu = T_{tt}e_0{}^t e_0{}^t$. This gives

$$\begin{aligned}
T_{\mu\nu}t^\mu t^\nu &= \frac{i}{4}\left[\bar{\Psi}\gamma_t\nabla_t\Psi + \bar{\Psi}\gamma_t\nabla_t\Psi - (\nabla_t\bar{\Psi})\gamma_t\Psi - (\nabla_t\bar{\Psi})\gamma_t\Psi\right]e_0{}^t e_0{}^t \\
&= 2\left[\frac{i}{4}\bar{\Psi}\gamma_t\nabla_t\Psi e_0{}^t e_0{}^t - \frac{i}{4}(\nabla_t\bar{\Psi})\gamma_t\Psi e_0{}^t e_0{}^t\right].
\end{aligned} \tag{50}$$

Inserting (28),(30) and (45) into this equation leads to

$$\begin{aligned}
T_{\mu\nu}t^\mu t^\nu &= 2\frac{\left(r^4 - 2Mr^3 + Q^2r^2\right)^{-1/2}}{4\sqrt{f}}\left[\omega\left(|R_1|^2 + |R_2|^2\right)\left(|S_1|^2 + |S_2|^2\right) \right. \\
&\quad \left. + i\frac{Mr - Q^2}{2r^3}\left(|R_1|^2 - |R_2|^2\right)\left(|S_1|^2 + |S_2|^2\right)\right] \\
&\quad - 2\frac{\left(r^4 - 2Mr^3 + Q^2r^2\right)^{-1/2}}{4\sqrt{f}}\left[-\omega\left(|R_1|^2 + |R_2|^2\right)\left(|S_1|^2 + |S_2|^2\right) \right. \\
&\quad \left. + i\frac{Mr - Q^2}{2r^3}\left(|R_1|^2 - |R_2|^2\right)\left(|S_1|^2 + |S_2|^2\right)\right] \\
&= \frac{\left(r^4 - 2Mr^3 + Q^2r^2\right)^{-1/2}}{2\sqrt{f}}2\omega\left(|R_1|^2 + |R_2|^2\right)\left(|S_1|^2 + |S_2|^2\right).
\end{aligned} \tag{51}$$

It is clear that outside the outer horizon the expression under the square root is greater than zero. Moreover, the expression (51) as a whole is strictly positive-definite, that is $T_{\mu\nu}t^\mu t^\nu \geq 0$, implying that the noncommutative deformation of the Reissner–Nordström spacetime does not violate the weak energy condition for the fermionic field. This in turn implies that the weak energy condition is not violated for the Dirac particle in a Reissner Nordstrom spacetime subject to a noncommutative deformation. Since the key assumption for the second law of black hole

thermodynamics is not violated, the law continues to hold and the superradiance is expected to occur for the fermionic field in the spacetime described by the effective (deformed RN) metric, contrary to a first naive impression obtained by considering the radial component of the Dirac current. This result is different from the case of the fermionic field on Kerr spacetime in the near horizon region $f(r) \rightarrow 0$ and in the superradiant⁵ regime $\omega < \nu \frac{\Omega}{2Mr_h}$, where ω is the frequency of the mode and ν is its azimuthal number, $\frac{\Omega}{2Mr_h}$ is the angular frequency of the horizon and r_h is the horizon radius. In the latter case the weak-energy condition is violated for the Dirac field on Kerr spacetime and consequently the effect of superradiance is absent. In our future work we plan to use the results obtained in this paper in order to study the massless as well as the massive fermionic perturbations of RN black hole in the presence of spacetime noncommutativity.

Acknowledgment We would like to thank Kumar Gupta, Tajron Juricé and Ivica Smolié for fruitful discussion and useful comments. This research was supported by the Croatian Science Foundation Project No. IP-2020-02-9614 *Search for Quantum spacetime in Black Hole QNM spectrum and Gamma Ray Bursts*. The work of M.D.C. and N.K. is supported by project 451-03-9/2021-14/200162 of the Serbian Ministry of Education and Science. This work is partially supported by ICTP-SEENET-MTP Project NT-03 "Cosmology-Classical and Quantum Challenges" in frame of the Southeastern European Network in Theoretical and Mathematical Physics and the COST action CA18108 *Quantum gravity phenomenology in the multimessenger approach*.

-
- [1] A. Kempf, G. Mangano and R. B. Mann, Phys. Rev. D 52, 1108 (1995).
 - [2] E. Witten, Phys. Today 49 (4), 24 (1996).
 - [3] D. Amati, M. Ciafaloni and G. Veneziano, Phys. Lett. B 197, 81 (1987).
 - [4] D. J. Gross and P. F. Mende, Nucl. Phys. B 303, 407 (1988).
 - [5] L. J. Garay, Int. J. Mod. Phys. A 10, 145 (1995).
 - [6] Spin networks and quantum gravity, Carlo Rovelli and Lee Smolin, Phys. Rev. **D52** (1995) 5743.
 - [7] M. Maggiore, Phys. Lett. B 304, 65 (1993).
 - [8] C. Rovelli, "Quantum Gravity," Cambridge, UK: Univ. Pr. (2004) 455 p.
 - [9] T. Thiemann, Lect. Notes Phys. 631 (2003) 41, arXiv:gr-qc/0210094.
 - [10] D. Amati, M. Ciafaloni, and G. Veneziano, Phys. Lett. B 216, 41 (1989).
 - [11] S. Hossenfelder, Living Rev. Relativity 16, 2 (2013).
 - [12] M. Sprenger, P. Nicolini and M. Bleicher, Eur. J. Phys. 33, 853 (2012).
 - [13] D. V. Ahluwalia, Phys. Lett. B 339 (1994) 301.
 - [14] S. Doplicher, K. Fredenhagen and J. E. Roberts, Phys. Lett. B 331 (1994) 39; Commun. Math. Phys. 172 (1995) 187, hep-th/0303037.
 - [15] N. Seiberg and E. Witten, JHEP 09 (1999) 032, hep-th/9908142.
 - [16] M. R. Douglas and N. A. Nekrasov, Noncommutative field theory, Rev. Mod. Phys. 73, 977 (2001) [hep-th/0106048].
 - [17] R. J. Szabo, Quantum field theory on noncommutative spaces, Phys. Rept. 378, 207 (2003) [hep-th/0109162];
 - [18] H. S. Snyder, Quantized space-time," Phys. Rev. 71, 38 (1947).
 - [19] A. Connes, Noncommutative Geometry," ISBN- 9780121858605.
 - [20] A. Connes and J. Lott, Nucl. Phys. Proc. Suppl. 18B, 29 (1991).
 - [21] M. Chaichian, A. Demichev and P. Presnajder, Nucl. Phys. B **567** (2000), 360-390, [arXiv:hep-th/9812180 [hep-th]].
 - [22] M. Hayakawa, [arXiv:hep-th/9912167 [hep-th]].
 - [23] I. F. Riad, M. M. Sheikh-Jabbari, JHEP 0008 (2000) 045.
 - [24] I. Ya. Aref'eva, D. M. Belov, A. S. Koshelev, Phys. Lett. B 476 (2000) 431.
 - [25] A. Micu and M. M. Sheikh Jabbari, JHEP **01** (2001), 025, [arXiv:hep-th/0008057 [hep-th]].
 - [26] B. Jurčo, L. Möller, S. Schraml, P. Schupp and J. Wess, *Construction of non-Abelian gauge theories on noncommutative spaces*, Eur. Phys. J. C**21**, 383 (2001), [hep-th/0104153].
 - [27] M. Chaichian, P. Presnajder, M. M. Sheikh-Jabbari and A. Tureanu, Eur. Phys. J. C **29** (2003), 413-432, [arXiv:hep-th/0107055 [hep-th]].
 - [28] M. Chaichian, P. P. Kulish, K. Nishijima and A. Tureanu, Phys. Lett. B **604** (2004), 98-102, [arXiv:hep-th/0408069 [hep-th]].
 - [29] P. Aschieri, C. Blohmann, M. Dimitrijevic, F. Meyer, P. Schupp and J. Wess, Class. Quant. Grav. **22** (2005), 3511-3532, [arXiv:hep-th/0504183 [hep-th]].

⁵ The term superradiant is here used because in this regime bosonic fields rapidly grow in time, thus exhibiting a superradiance. Dirac fields though remain stable in this regime, as they are not superradiant there.

- [30] P. Aschieri, M. Dimitrijevic, F. Meyer and J. Wess, *Class. Quant. Grav.* **23** (2006), 1883-1912, [arXiv:hep-th/0510059 [hep-th]].
- [31] B. Harms and O. Micu, *J. Phys. A* **40**, 10337 (2007) [hep-th/0610081].
- [32] F. G. Scholtz, B. Chakraborty, S. Gangopadhyay and J. Govaerts, *J. Phys. A* **38**, 9849 (2005) [cond-mat/0509331 [cond-mat.mes-hall]].
- [33] Y. Okawa and H. Ooguri, An Exact solution to Seiberg-Witten equation of noncommutative gauge theory, *Phys. Rev. D* **64** (2001) 046009, [hep-th/0104036].
- [34] G. Barnich, F. Brandt and M. Grigoriev, Seiberg-Witten maps and noncommutative Yang-Mills theories for arbitrary gauge groups, *JHEP* **0208** (2002) 023, [hep-th/0206003].
- [35] M. Buric and V. Radovanovic, *JHEP* **02** (2004), 040, [arXiv:hep-th/0401103 [hep-th]].
- [36] M. Buric, V. Radovanovic and J. Trampetic, *JHEP* **03** (2007), 030, [arXiv:hep-th/0609073 [hep-th]].
- [37] D. Latas, V. Radovanovic and J. Trampetic, *Phys. Rev. D* **76** (2007), 085006, [arXiv:hep-th/0703018 [hep-th]].
- [38] C. P. Martin, Computing the θ -exact Seiberg-Witten map for arbitrary gauge groups, *Phys. Rev. D* **86** (2012) 065010, [hep-th]].
- [39] R. Horvat, A. Ilakovac, P. Schupp, J. Trampetić, and J. You, Yukawa couplings and seesaw neutrino masses in noncommutative gauge theory, *Phys. Lett. B* **715**, 340-347 (2012),
- [40] M. Dimitrijevic, L. Jonke and A. Pachol, *SIGMA* **10** (2014), 063, [arXiv:1403.1857 [hep-th]].
- [41] M. Dimitrijević Ćirić, D. Gočanin, N. Konjik and V. Radovanović, *Eur. Phys. J. C* **78** (2018) no.7, 548, [arXiv:1804.00608 [hep-th]].
- [42] P. Aschieri and L. Castellani, *JHEP* **06** (2009), 086, [arXiv:0902.3817 [hep-th]].
- [43] P. Aschieri and L. Castellani, “Noncommutative gravity coupled to fermions: second order expansion via Seiberg-Witten map,” *JHEP* **07** (2012), 184, [arXiv:1111.4822 [hep-th]].
- [44] P. Aschieri, L. Castellani and M. Dimitrijević, *Phys. Rev. D* **87** (2013) no.2, 024017, [arXiv:1207.4346 [hep-th]].
- [45] P. Aschieri and L. Castellani, *JHEP* **11** (2014), 103, [arXiv:1406.4896 [hep-th]].
- [46] M. Dimitrijević Ćirić, B. Nikolić and V. Radovanović, *Phys. Rev. D* **96** (2017) no.6, 064029, [arXiv:1612.00768 [hep-th]].
- [47] S. Meljanac and M. Stojić, “New realizations of Lie algebra kappa-deformed Euclidean space,” *Eur. Phys. J. C* **47**, 531 (2006) [hep-th/0605133].
- [48] S. Kresic-Jurić, S. Meljanac and M. Stojić, “Covariant realizations of kappa-deformed space,” *Eur. Phys. J. C* **51**, 229 (2007) [hep-th/0702215].
- [49] S. Meljanac, A. Samsarov, M. Stojić and K. S. Gupta, “Kappa-Minkowski space-time and the star product realizations,” *Eur. Phys. J. C* **53**, 295 (2008) [arXiv:0705.2471 [hep-th]].
- [50] S. Meljanac and S. Kresic-Juric, *J. Phys. A* **41** (2008), 235203 [arXiv:0804.3072 [hep-th]].
- [51] T. Jurić, S. Meljanac and R. Strajn, “Twists, realizations and Hopf algebroid structure of kappa-deformed phase space,” *Int. J. Mod. Phys. A* **29**, no. 5, 1450022 (2014) [arXiv:1305.3088 [hep-th]].
- [52] E. Harikumar, T. Jurić, and S. Meljanac, *Phys. Rev. D* **86** (2012) 045002.
- [53] T. Jurić, S. Meljanac, D. Pikutić and R. Strajn, *JHEP* **1507** (2015) 055.
- [54] K. S. Gupta, E. Harikumar, T. Jurić, S. Meljanac and A. Samsarov, *Adv. High Energy Phys.* **2014** (2014) 139172.
- [55] K. S. Gupta, E. Harikumar, T. Jurić, S. Meljanac and A. Samsarov, *JHEP* **1509** (2015) 025.
- [56] T. Jurić and A. Samsarov, *Phys. Rev. D* **93** (2016) no.10, 104033.
- [57] K. S. Gupta, T. Jurić and A. Samsarov, *JHEP* **1706** (2017) 107.
- [58] K. S. Gupta, T. Jurić, A. Samsarov and I. Smolić, *JHEP* **10** (2019), 170, [arXiv:1908.07402 [hep-th]].
- [59] R. Bufalo and A. Tureanu, *Phys. Rev. D* **92** (2015) no.6, 065017.
- [60] J. Trampetić and J. You, *Phys. Rev. D* **105** (2022) no.7, 075016, [arXiv:2111.04154 [hep-th]].
- [61] R. Horvat, C. P. Martin, J. Trampetic and J. You, [arXiv:1609.08065 [hep-th]].
- [62] M. D. Ćirić, N. Konjik and A. Samsarov, *Class. Quant. Grav.* **35** (2018) no.17, 175005, [arXiv:1708.04066 [hep-th]].
- [63] M. Dimitrijević Ćirić, N. Konjik and A. Samsarov, *Phys. Rev. D* **101** (2020) no.11, 116009, [arXiv:1904.04053 [hep-th]].
- [64] M. D. Ćirić, N. Konjik and A. Samsarov, [arXiv:1910.13342 [hep-th]].
- [65] T. C. Adorno, M. C. Baldiotti, M. Chaichian, D. M. Gitman and A. Tureanu, *Phys. Lett. B* **682** (2009), 235-239, [arXiv:0904.2836 [hep-th]].
- [66] E. Harikumar and M. Sivakumar, *Mod. Phys. Lett. A* **26** (2011), 1103-1115, [arXiv:0910.5778 [hep-th]].
- [67] F. M. Andrade, E. O. Silva, Effects of quantum deformation on the spin-1/2 Aharonov-Bohm problem, *Phys. Lett. B* **719** (2013) 467.
- [68] R. Horvat, D. Kekez, P. Schupp, J. Trampetic and J. You, *Phys. Rev. D* **84** (2011), 045004, [arXiv:1103.3383 [hep-ph]].
- [69] M. Buric, D. Latas, V. Radovanovic and J. Trampetic, *Phys. Rev. D* **83** (2011), 045023, [arXiv:1009.4603 [hep-th]]. [70]
- [70] C. L. Ching, C. X. Yeo and W. K. Ng, *Int. J. Mod. Phys. A* **32** (2017) no.02n03, 1750009, [arXiv:1601.04420 [gr-qc]].
- [71] M. Arzano and J. Kowalski-Glikman, An Introduction to κ -Deformed Symmetries, *Phase Spaces and Field Theory, Symmetry* **2021**, 13(6), 946.
- [72] S. Mignemi, Classical and quantum mechanics of the nonrelativistic Snyder model. *Phys. Rev. D* **84**, 025021 (2011).
- [73] S. Mignemi, Classical and quantum mechanics of the nonrelativistic Snyder model in curved space. *Class. Quantum Gravity* **29**, 215019 (2012).
- [74] H. Falomir, J. Gamboa, J. Lopez-Sarrion, F. Mendez, and P. A. G. Pisani, *Phys. Lett. B* **680**, 384 (2009).
- [75] A. Das, H. Falomir, M. Nieto, J. Gamboa, and F. Mendez, *Phys. Rev. D* **84**, 045002 (2011).

- [76] M. Gomes, V. G. Kupriyanov, and A. J. da Silva, Phys. Rev. D **81**, 085024 (2010).
- [77] F. Berezin and M. Marinov, Annals Phys. **104**, 336 (1977).
- [78] A. F. Ferrari, M. Gomes, V. G. Kupriyanov and C. A. Stechhahn, Phys. Lett. B **718** (2013), 1475-1480, [arXiv:1207.0412 [hep-th]].
- [79] M. Moshinsky, A. Szczepaniak, J. Phys. A: Math. Gen. **22** L817 (1989).
- [80] B. P. Mandal and S. Verma, Phys. Lett. A **374** (2010), 1021-1023, [arXiv:0907.4544 [hep-th]].
- [81] B. P. Mandal and S. K. Rai, Phys. Lett. A **376** (2012), 2467-2470, [arXiv:1203.2714 [hep-th]].
- [82] C. Quesne, V. M. Tkachuk, Dirac oscillator with nonzero minimal uncertainty in position, J. Phys. A **38** (2005) 1747.
- [83] C. Quesne, V. M. Tkachuk, Lorentz-covariant deformed algebra with minimal length and application to the (1 + 1)-dimensional Dirac oscillator, J. Phys. A **39** (2006) 10909.
- [84] M. M. Stetsko, J. Math. Phys. **56** (2015), 012101, [arXiv:1310.0706 [quant-ph]].
- [85] M. M. Stetsko, Mod. Phys. Lett. A **34** (2019) no.36, 1950300, [arXiv:1810.00056 [quant-ph]].
- [86] F. M. Andrade, E. O. Silva, M. M. Ferreira and E. C. Rodrigues, Phys. Lett. B **731** (2014), 327-330, [arXiv:1312.2973 [hep-th]].
- [87] F. M. Andrade and E. O. Silva, Phys. Lett. B **738** (2014), 44-47, [arXiv:1406.3547 [hep-th]].
- [88] M. Merad and M. Hadj Moussa, Exact Solution of Klein–Gordon and Dirac Equations with Snyder–de Sitter Algebra, Few-Body Syst **59**, 5 (2018).
- [89] B. Hamil and M. Merad, Few Body Syst. **60** (2019) no.2, 36.
- [90] R. R. S. Oliveira, G. Alencar and R. R. Landim, [arXiv:2204.06057 [hep-th]].
- [91] G. Amelino-Camelia, Nature **418** 34 (2002); Phys. Lett. B **510** (2001) 255; Int. J. Mod. Phys. A **11** (2002) 35.
- [92] J. Magueijo and L. Smolin, Phys. Rev. Lett. **88**(2002) 190403; Phys. Rev. D **67** (2003) 044017.
- [93] S. Mignemi, Phys. Lett. B **551** (2003), 169-172, [arXiv:hep-th/0208062 [hep-th]].
- [94] A. Nowicki, E. Sorace, and M. Tarlini, The Quantum deformed Dirac equation from the kappa Poincare algebra, Phys. Lett. B **302**, 419 (1993), arXiv:hep-th/9212065.
- [95] J. J. Relancio and S. Liberati, Phys. Rev. D **101** (2020) no.6, 064062, [arXiv:2002.10833 [gr-qc]].
- [96] J. J. Relancio and S. Liberati, Class. Quant. Grav. **38** (2021) no.13, 135028, [arXiv:2010.15734 [gr-qc]].
- [97] C. Pfeifer and J. J. Relancio, Eur. Phys. J. C **82** (2022) no.2, 150, [arXiv:2103.16626 [gr-qc]].
- [98] S. A. Franchino-Viñas and J. J. Relancio, [arXiv:2203.12286 [hep-th]].
- [99] G. Amelino-Camelia, L. Freidel, J. Kowalski-Glikman and L. Smolin, Phys. Rev. D **84** (2011), 084010, [arXiv:1101.0931 [hep-th]].
- [100] G. Amelino-Camelia, M. Arzano, J. Kowalski-Glikman, G. Rosati and G. Trevisan, Class. Quant. Grav. **29** (2012), 075007, [arXiv:1107.1724 [hep-th]].
- [101] G. Gubitosi and F. Mercati, Class. Quant. Grav. **30** (2013), 145002, [arXiv:1106.5710 [gr-qc]].
- [102] J. M. Carmona, J. L. Cortes and F. Mercati, Phys. Rev. D **86** (2012), 084032, [arXiv:1206.5961 [hep-th]].
- [103] J. M. Carmona, J. L. Cortes and J. J. Relancio, Phys. Rev. D **94** (2016) no.8, 084008, [arXiv:1609.01347 [hep-th]].
- [104] J. M. Carmona, J. L. Cortés and J. J. Relancio, Phys. Rev. D **100** (2019) no.10, 104031, [arXiv:1907.12298 [hep-th]].
- [105] B. Ivetic, S. Mignemi and A. Samsarov, Phys. Rev. D **94** (2016) no.6, 064064. [arXiv:1606.04692 [hep-th]].
- [106] A. Agostini, G. Amelino-Camelia and M. Arzano, Class. Quant. Grav. **21** (2004), 2179-2202, [arXiv:gr-qc/0207003 [gr-qc]].
- [107] R. Aloisio, J. M. Carmona, J. L. Cortes, A. Galante, A. F. Grillo and F. Mendez, JHEP **05** (2004), 028, [arXiv:hep-th/0404111 [hep-th]].
- [108] P. Gosselin, A. Berard, H. Mohrbach and S. Ghosh, Phys. Lett. B **660** (2008), 267-274, [arXiv:0709.0579 [hep-th]].
- [109] Z. Belhadi, F. Menas, A. Berard, P. Gosselin and H. Mohrbach, Int. J. Mod. Phys. A **27** (2012), 1250031, [arXiv:1106.1302 [hep-th]].
- [110] P. N. Bibikov, J. Phys. A **31** (1998) 6437.
- [111] K. Kosinski, J. Lukierski and P. Maslanka, Nucl. Phys. Proc. Suppl. **102** (2001) 161.
- [112] P. Kosinski, P. Maslamka, J. Lukierski and A. Sitarz, Czech. Journ. Phys. **48** (1998) 11.
- [113] S. R. Dolan and D. Dempsey, Class. Quant. Grav. **32** (2015) no.18, 184001, [arXiv:1504.03190 [gr-qc]].
- [114] S. L. Detweiler, Phys. Rev. D **22** (1980) 2323–2326.
- [115] J.G. Rosa and S. R. Dolan, Phys. Rev. D **85** (2012) 044043 (Preprint 1110.4494).
- [116] H. Witek, V. Cardoso, A. Ishibashi and U. Sperhake, Phys. Rev. D **87** (2013) 043513 (Preprint 1212.0551).
- [117] P. Pani, V. Cardoso, L. Gualtieri, E. Berti and A. Ishibashi, Phys. Rev. Lett. **109** (2012) 131102 (Preprint 1209.0465).
- [118] P. Pani, V. Cardoso, L. Gualtieri, E. Berti and A. Ishibashi, Phys. Rev. D **86** (2012) 104017 (Preprint 1209.0773).
- [119] M. Martellini and A. Treves, Phys. Rev. D **15** (1977) 3060–3061.
- [120] B. R. Iyer and A. Kumar, Pramana **8** (1977) 500–511.
- [121] B. R. Iyer and A. Kumar, Phys. Rev. D **18** (1978) 4799–4801.
- [122] I. Ternov, A. B. Gaina and G. Chizhov, Sov. Phys. J. **23** (1980) 695–700.
- [123] W. Unruh, Phys. Rev. Lett. **31**(20) (1973) 1265–1267.