

Measuring incompatibility and clustering quantum observables with a quantum switch

Ning Gao,¹ Dantong Li,¹ Anchit Mishra,¹ Junchen Yan,¹ Kyrylo Simonov,^{2,3} and Giulio Chiribella^{1,4,5}

¹*QICI Quantum Information and Computation Initiative, Department of Computer Science,
The University of Hong Kong, Pokfulam Road, Hong Kong*

²*s7 rail technology GmbH, Lastenstraße 36, 4020 Linz, Austria*

³*Fakultät für Mathematik, Universität Wien, Oskar-Morgenstern-Platz 1, 1090 Vienna, Austria*

⁴*Quantum Group, Department of Computer Science, University of Oxford,
Wolfson Building, Parks Road, Oxford, OX1 3QD, United Kingdom*

⁵*Perimeter Institute for Theoretical Physics, 31 Caroline Street North, Waterloo, Ontario, Canada*

(Dated: August 15, 2022)

The existence of incompatible observables is a cornerstone of quantum mechanics and a valuable resource in quantum technologies. Here we introduce a measure of incompatibility, called the mutual eigenspace disturbance (MED), which quantifies the amount of disturbance induced by the measurement of a sharp observable on the eigenspaces of another. The MED is a faithful measure of incompatibility for sharp observables and provides a metric on the space of von Neumann measurements. It can be efficiently estimated by letting the measurements act in an indefinite order, using a setup known as the quantum switch. Thanks to these features, the MED can be used in quantum machine learning tasks, such as clustering quantum measurement devices based on their mutual compatibility. We demonstrate this application by providing an unsupervised algorithm that clusters unknown von Neumann measurements. Our algorithm is robust to noise and can be used to identify groups of observers that share approximately the same measurement context.

Introduction.— One of the most striking features of quantum mechanics is the existence of incompatible observables, such as the position and velocity of a particle, or the projections of a spin along two different axes. Incompatible observables are at the heart of Bohr’s notion of complementarity [1] and of Heisenberg’s uncertainty principle [2]. In addition to their foundational relevance, they play center stage in quantum information technologies [3]. For example, the existence of incompatible observables enables secure key distribution in the BB84 protocol [4], in which any attempt at eavesdropping induces a disturbance that can be detected by the communicating parties. In general, quantum incompatibility is a valuable resource [5], in a similar way as quantum entanglement and coherence.

Several measures of incompatibility have been proposed in the past years, capturing different aspects, such as robustness to noise [6, 7], nonorthogonality [8], and the disturbance on the measurement statistics [9]. These measures are conceptually important, because they provide quantitative insights into the operational features associated to incompatibility. On the other hand, they are sometimes hard to compute analytically, because they are defined in terms of optimization problems that generally require numerical optimization. In addition, the existing incompatibility measures cannot be obtained experimentally in a direct way: in most cases, the best known way to infer the incompatibility of two unknown observables is to perform a full tomography, which is challenging for systems of many particles.

In this paper we introduce a measure of incompatibility for sharp observables [10], that is, quantum measurements associated to the spectral resolution of self-adjoint operators. This measure, called the Mutual Eigenspace Disturbance (MED), quantifies the amount of disturbance induced by the measurement of an observable on the eigenspaces of another. The MED has a simple closed-form analytical expression, and, at the same time, it retains most of the key

conceptual properties of other incompatibility measures: it is faithful (that is, strictly positive if and only if the observables are incompatible), decreasing under coarse-graining, and maximal for complementary observables. It constitutes a metric on the space of von Neumann measurements, and, in addition, it can be estimated experimentally by testing the measurement processes in a setup, known as the quantum switch [11, 12], that probes the measurement processes in a superposition of alternative causal orders. This realization is practically appealing, as the quantum switch can be reproduced on various photonic setups, as demonstrated in a series of experiments [13–19] (see also [20] for a review), and is potentially implementable also on other experimental platforms, such as superconducting qubits [21]. Using the quantum switch, the incompatibility can be estimated with a number of experiments that is independent of the dimension of the measured system, thereby avoiding the exponentially growing complexity of quantum process tomography. An additional feature is that the quantum switch setup does not require access to the measurement outcomes: the incompatibility between two unknown observables can be estimated by probing only the average evolution associated to the two measurement processes, without examining the outcome statistics.

The ability to experimentally measure the MED, together with the metric properties for nondegenerate observables, make the MED suitable for applications in quantum machine learning. To illustrate the idea, we provide a quantum algorithm that clusters nondegenerate observables based on their mutual compatibility. At the foundational level, this algorithm can be used to identify clusters of observers who share approximately the same measurement context [22–25], and thereby could share the same notion of an emergent classical reality [26–30]. At a more practical level, we show that the algorithm is robust to the presence of noise in the data, and to deviations of the measurements performed in the laboratory

from ideal projective measurements.

Mutual Eigenspace Disturbance.—Standard quantum observables, sometimes called sharp observables [10], are associated to the spectral resolution of self-adjoint operators in terms of orthogonal projectors. For general (possibly non-projective) quantum measurements, compatibility is defined as the existence of a joint measurement [3]. For projective measurements, this notion is equivalent to commutativity of the projectors associated to the two observables [31].

We now introduce the MED. The intuition behind the definition is that commuting observables have a common set of eigenstates, and therefore the measurement of one observable leaves the eigenspaces of the other observable invariant. The violation of this condition can then be regarded as an indicator of incompatibility. Let A and B be two observables, and let $\mathbf{P} = (P_i)_{i=1}^{k_A}$ and $\mathbf{Q} = (Q_j)_{j=1}^{k_B}$ be the $k_{A/B}$ projectors associated to the eigenspaces of A and B , respectively ($k_{A/B} \leq d$, where d is the total Hilbert space's dimension). To quantify the disturbance of a measurement of B on the eigenspaces of A , we imagine that the system is initially in an eigenstate of A , say $|\alpha_i\rangle$, picked uniformly at random from the i -th eigenspace, with i distributed according to the probability distribution $p_i := d_{A,i}/d$, where $d_{A,i}$ is the eigenspace's dimension. Then, the system undergoes the canonical (Lüders) measurement process associated to the observable B . On average over all outcomes, the canonical measurement of B induces a dephasing channel \mathcal{B} defined by the relation $\mathcal{B}(\rho) = \sum_j Q_j \rho Q_j$ for generic density matrices ρ . Finally, a measurement of the observable A is performed. The probability to find the outcome i , associated to the original subspace, is $\text{Tr}[P_i \mathcal{B}(|\alpha_i\rangle\langle\alpha_i|)]$. On average, the probability that a random eigenstate of A appears to be still inside its original eigenspace is

$$\text{Prob}(A, \mathcal{B}) := \sum_i p_i \int \pi_i(d\alpha_i) \text{Tr}[P_i \mathcal{B}(|\alpha_i\rangle\langle\alpha_i|)], \quad (1)$$

where $\pi_i(d\alpha_i)$ is the uniform probability distribution over the pure states in the i -th subspace. Explicit calculation yields the expression

$$\text{Prob}(A, \mathcal{B}) := \frac{1}{d} \sum_{ij} \text{Tr}[P_i Q_j P_i Q_j]. \quad (2)$$

Note that the role of the projectors \mathbf{P} and \mathbf{Q} is completely symmetric. Operationally, this symmetry implies to the relation

$$\text{Prob}(A, \mathcal{B}) = \text{Prob}(B, \mathcal{A}), \quad (3)$$

where $\text{Prob}(B, \mathcal{A})$ is the average probability that a randomly chosen state $|\beta_j\rangle$ from the j -th eigenspace of B , drawn with probability $q_j := d_{B,j}/d$ (where $d_{B,j}$ is the eigenspace's dimension), is still found in the same eigenspace after the action of the dephasing channel $\mathcal{A} : \rho \mapsto \mathcal{A}(\rho) = \sum_j P_j \rho P_j$.

Note also that the probabilities in Eq. (3) depend only on the dephasing channels \mathcal{A} and \mathcal{B} . Accordingly, we will denote by $D(\mathcal{A}, \mathcal{B}) := 1 - \text{Prob}(A, \mathcal{B}) \equiv 1 - \text{Prob}(B, \mathcal{A})$ the

average probability of eigenstate disturbance. We then define the MED of the two observables A and B as

$$\text{MED}(\mathcal{A}, \mathcal{B}) := \sqrt{D(\mathcal{A}, \mathcal{B})}. \quad (4)$$

The reason for the square root is that it guarantees that the MED is a distance measure on the set of von Neumann measurements, associated to rank-one projectors.

The MED exhibits several appealing properties for a measure of incompatibility:

1. it is *symmetric and nonnegative*, namely $\text{MED}(\mathcal{A}, \mathcal{B}) = \text{MED}(\mathcal{B}, \mathcal{A}) \geq 0$ for every pair of dephasing channels \mathcal{A} and \mathcal{B} ,
2. it is *faithful*, namely $\text{MED}(\mathcal{A}, \mathcal{B}) > 0$ if and only if A and B are incompatible,
3. it is *maximal for maximally complementary observables* [32], that is, observables such that their eigenstates form mutually unbiased bases [33, 34]. In general, one has the bound $\text{MED}(\mathcal{A}, \mathcal{B}) \leq \sqrt{1 - 1/\min\{k_A, k_B\}}$, and the maximum value $\text{MED}(\mathcal{A}, \mathcal{B}) = \sqrt{1 - 1/d}$ and is attained if and only if A and B are maximally complementary,
4. it is *decreasing under coarse-graining*,
5. it is a *metric on von Neumann measurements*, corresponding to the channels of the form $\mathcal{A}(\rho) = \sum_{i=1}^d \langle\alpha_i|\rho|\alpha_i\rangle |\alpha_i\rangle\langle\alpha_i|$ for some orthonormal basis $\{|\alpha_i\rangle\}_{i=1}^d$. In this case, one has $\text{MED}(\mathcal{A}, \mathcal{B}) = 0$ if and only if $\mathcal{A} = \mathcal{B}$, and the triangle inequality $\text{MED}(\mathcal{A}, \mathcal{C}) \leq \text{MED}(\mathcal{A}, \mathcal{B}) + \text{MED}(\mathcal{B}, \mathcal{C})$ holds for every \mathcal{A}, \mathcal{B} , and \mathcal{C} , and
6. it is *robust to noise*, that is, it remains faithful even if one of the channels \mathcal{A} and \mathcal{B} is replaced by the evolution resulting from a noisy measurement of the corresponding observable.

The proof of the above properties is provided in the Appendix A.

Experimental setups for estimating the MED.—The closed-form expression (4) can be used to design experimental setups that estimate the MED of two unknown observables. A naïve approach would be to estimate the channels \mathcal{A} and \mathcal{B} via process tomography [35–40], or the projective measurements \mathbf{P} and \mathbf{Q} via measurement tomography [41]. However, process and measurement tomography require a number of experimental settings that grows polynomially with the dimension, and therefore exponentially in the number of particles for multiparticle quantum systems.

Another way to estimate the MED is to perform sequential measurements. The scheme works as follows: first, the system is prepared in the maximally mixed state. Then, it undergoes a canonical measurement of the observable A , a canonical measurement of the observable B , and finally, another measurement of the observable A . The total probability that the two measurements of A give equal outcomes is equal

to $\text{Prob}(A, \mathcal{B}) = 1 - \text{MED}(\mathcal{A}, \mathcal{B})^2$. In this case, the complexity of estimating the MED does not grow exponentially with the system's size. However, this estimation strategy requires access to the outcomes of two measurements of the observable A . Access to the outcomes may not be available in certain situations, *e.g.* if the experimenter wants to discover the incompatibility of two observables measured by two parties in their local laboratories. In the following we show that, fortunately, there exists a way to estimate the incompatibility of two observables without any access to the measurement outcomes.

The measurement protocol uses the quantum switch [11, 12], an operation that combines two unknown channels in a coherent superposition of two alternative orders. The quantum switch was previously shown to be able to distinguish between pairs of quantum channels with commuting or anti-commuting Kraus operators [42], a task that can be practically achieved with photonic systems [13, 43]. We now show that the quantum switch can be used to estimate the amount of commutativity of two unknown quantum channels.

When the quantum switch is applied to two generic quantum channels \mathcal{C} and \mathcal{D} , it produces a new quantum channel $\mathcal{S}_{\mathcal{C}, \mathcal{D}}$ with Kraus operators

$$S_{ij} = C_i D_j \otimes |0\rangle\langle 0| + D_j C_i \otimes |1\rangle\langle 1| \quad (5)$$

where (C_i) and (D_j) are Kraus operators for \mathcal{C} and \mathcal{D} , respectively, and $\{|0\rangle, |1\rangle\}$ are basis states of a control qubit, which controls the order of the channels \mathcal{C} and \mathcal{D} . The action of the channel $\mathcal{S}_{\mathcal{C}, \mathcal{D}}$ on a generic product state $\rho \otimes \omega$ is

$$\begin{aligned} \mathcal{S}_{\mathcal{C}, \mathcal{D}}(\rho \otimes \omega) = & \frac{1}{4} \sum_{ij} \left(\{C_i, D_j\} \rho \{C_i, D_j\}^\dagger \otimes \omega \right. \\ & + \{C_i, D_j\} \rho [C_i, D_j]^\dagger \otimes \omega Z \\ & + [C_i, D_j] \rho \{C_i, D_j\}^\dagger \otimes Z \omega \\ & \left. + [C_i, D_j] \rho [C_i, D_j]^\dagger \otimes Z \omega Z \right), \quad (6) \end{aligned}$$

where $[C_i, D_j] := C_i D_j - D_j C_i$ denotes the commutator, $\{C_i, D_j\} := C_i D_j + D_j C_i$ denotes the anti-commutator, and $Z := |0\rangle\langle 0| - |1\rangle\langle 1|$.

To estimate the MED, we apply the quantum switch with the system initialized in the maximally mixed state $\rho = I/d$ and the control initialized in the maximally coherent state $\omega = |+\rangle\langle +|$, using the standard notation for the Fourier basis $|\pm\rangle = (|0\rangle \pm |1\rangle) / \sqrt{2}$. Then, we measure the control qubit in the Fourier basis, obtaining the outcome $-$ with probability

$$p_- = \frac{1}{4d} \sum_{ij} \left\| [C_i, D_j] \right\|_2^2, \quad (7)$$

where $\|O\|_2^2 := \text{Tr}[O^\dagger O]$ for a generic operator O .

We define the *non-commutativity* of two generic quantum channels \mathcal{C} and \mathcal{D} as

$$\text{NCOM}(\mathcal{C}, \mathcal{D}) := \sqrt{\frac{\sum_{ij} \left\| [C_i, D_j] \right\|_2^2}{2d}} \equiv \sqrt{2p_-} \quad (8)$$




FIG. 1. Clustering of 100 qubit observables, using a k -medoids algorithm based on the values of the MED with k -means++ style initial seeding. The algorithm correctly identifies two clusters, one centered around the Pauli X observable (Bloch vectors in yellow), and one centered around the Pauli Z observable (Bloch vectors in blue).

(in the special case $\mathcal{C} = \mathcal{D}$, a related definition was used in [44] to quantify the degree of non-commutativity of the Kraus operators of a given channel). It is evident from the definition that the noncommutativity is symmetric, positive, and faithful, meaning that $\text{NCOM}(\mathcal{C}, \mathcal{D}) = 0$ if and only if every Kraus operator \mathcal{C} commutes with every Kraus operator of \mathcal{D} . In the special case where \mathcal{C} and \mathcal{D} are two dephasing channels \mathcal{A} and \mathcal{B} , explicit calculation yields the relation

$$\text{NCOM}(\mathcal{A}, \mathcal{B}) = \text{MED}(\mathcal{A}, \mathcal{B}). \quad (9)$$

Eqs. (8) and (9) imply that the MED of two unknown observables can be estimated from the probability p_- in the quantum switch setup. The estimation can be done efficiently: in the Appendix B we show that the $\text{MED}(\mathcal{A}, \mathcal{B})$ can be guaranteed to have error at most ϵ with probability at least $1 - \delta$ by repeating the experiment for $n = \log \frac{2}{\delta} / (2\epsilon^2)$ times.

Clustering algorithm for quantum observables.—We now provide a machine learning algorithm for identifying clusters of observables that are approximately compatible with one another. The algorithm is unsupervised: the learner does not need to be trained with labelled examples of observables belonging to different clusters.

The input of the algorithm is the access to m black boxes, implementing m unknown dephasing channels $\mathcal{A}_1, \dots, \mathcal{A}_m$ associated to non-degenerate quantum observables A_1, \dots, A_m . The quantum part of the algorithm is the estimation of the MED for every pair of observables. Then,




FIG. 2. Clustering of 100 noisy qubit measurements, using a k -medoids algorithm based on the values of the noncommutativity with k -means++ style initial seeding. The algorithm correctly identifies two clusters, centered around the Pauli X and Z observables, respectively. The three plots refer to numerical experiments with maximum noise level 0.25, 0.5, and 0.75, respectively.

the estimated values of the MEDs are fed into a classical clustering algorithm. Here we choose k -medoids clustering with k -means++ style initial seeding [45, 46]). Compared to the popular k -means method, k -medoids works better with arbitrary dissimilarity measures other than Euclidean distances in terms of the convergence and efficiency of the clustering algorithm.

To illustrate the algorithm, we generate $m = 100$ random qubit observables, of the form $A_l = b_x^{(l)} X + b_y^{(l)} Y + b_z^{(l)} Z$, $l \in \{1, \dots, m\}$, where X, Y, Z are the three Pauli matrices and $\mathbf{b}_l = (b_x^{(l)}, b_y^{(l)}, b_z^{(l)}) \in \mathbb{R}^3$ is a unit vector (the Bloch vector of the l -th observable). The vectors are generated in the following way: for 50 observables, we start from the Bloch vector $(1, 0, 0)$ and apply a rotation by a random angle θ with $|\theta| \leq 22.5^\circ$ about a random rotation axis. For the remaining 50 observables, we start from the Bloch vector $(0, 0, 1)$ and apply the same procedure. In this way, the 100 observables are naturally divided into two clusters, as in Figure 1.

We performed a numerical experiment on the classical part of the algorithm, feeding the values of the MED into the k -medoids algorithm. For improved reliability, we repeated the experiment for 50 times, finding that in each repetition all the 100 observables are correctly clustered. Note that, while we fed the algorithm with the exact values of the MED, the robustness of the k -medoids [45, 46] algorithm implies that the results are robust to errors in the estimation of the MED from actual experimental data.

Clustering with noisy observables.—Here we extend the results of the previous section to noisy measurements. Following [3], the noise is modelled by randomizing the measurement of each ideal observable with a trivial measurement, which produces the same outcome statistics for every possible input state. Mathematically, this means that the projective measurement $\mathbf{P}^{(l)}$, associated to the l -th observable, is replaced by a non-projective measurement $\mathbf{N}^{(l)} = (1 - \lambda_l) \mathbf{P}^{(l)} + \lambda_l \mathbf{T}^{(l)}$, where λ_l is the noise probability, and $\mathbf{T}^{(l)}$ is a trivial measurement, with POVM operators $T_i^{(l)} = p_i^{(l)} I$,

for a fixed probability distribution $\mathbf{p}^{(l)} = (p_i^{(l)})$. For the measurement process associated to the noisy measurement $\mathbf{N}^{(l)}$, we take quantum instruments with Kraus operators of the form $N_{ij}^{(l)} = \sqrt{a_{ij}^{(l)} p_i^{(l)} + b_{ij}^{(l)} I}$, where $a_{ij}^{(l)}$ and $b_{ij}^{(l)}$ are arbitrary nonnegative coefficients subject to the constraints $\sum_j a_{ij}^{(l)} = 1 - \lambda_l$ and $\sum_j b_{ij}^{(l)} = \lambda_l p_i^{(l)}$ for every i and l . The k -medoid algorithm can then be applied, using the noncommutativity (8) of the noisy channels $\mathcal{N}_1, \dots, \mathcal{N}_m$ defined by $\mathcal{N}_l(\rho) := \sum_{ij} N_{ij}^{(l)} \rho N_{ij}^{(l)\dagger}$ for $l \in \{1, \dots, m\}$.

To test the algorithm, we performed a numerical experiment on 100 randomly generated noisy qubit observables. For simplicity, we chose isotropic noise [47] and set the noise randomly following a uniform distribution, picking each probability $p_i^{(l)}$ uniformly at random in the interval $(0, 1)$, subject to the constraint $\sum_i p_i^{(l)} = 1, \forall l$. For the original observables, we generated the Bloch vectors as in the noiseless case. For the noise, we first defined a maximum noise level η and then we picked a random noise probability $\lambda_l = \eta R_l$ where each R_l is chosen independently, uniformly at random in the interval $[0, 1]$. The coefficients $a_{ij}^{(l)}$ and $b_{ij}^{(l)}$ are then chosen uniformly at random, subject to the constraints. The experiment has been performed for $\eta = 0.25, 0.5$, and 0.75 , and for each setting the k -medoids algorithm has been run 50 times. The results of the experiment, plotted on Fig. 2, show that perfect clustering is still achieved in the presence of noise.

Conclusions and outlook.—In this paper we introduced the MED, an experimentally accessible measure of incompatibility for sharp observables. The MED quantifies the noncommutativity of the projectors associated to a pair of observables, and can be directly measured without access to the measurement outcomes, by letting the two measurement processes act in an indefinite order. Thanks to its properties, the MED can be used in quantum machine learning tasks, such as clustering unknown observables based on their degree of compatibility. To illustrate this application, we

demonstrated a clustering of noisy qubit observables using an unsupervised learning algorithm.

An interesting direction of future research is to extend the results of this paper to infinite dimensional systems, unsharp observables, and general quantum channels. For sharp observables with discrete spectrum, our approach can be easily extended by taking a limit of finite dimensional subspaces. For observables with continuous spectrum, however, the situation is more complex, due to the fact that no repeatable measurement exists [48]. Regarding the extension to unsharp observables and general channels, one approach is to consider measures of commutativity, which is a sufficient condition for joint measurability. On the other hand, commutativity is not a necessary condition, and an open question is to determine whether there exists a measure of incompatibility that can be measured experimentally like the MED.

Another direction is the extension of the MED from pairs to arbitrary numbers of observables. One option would be to generalize our definition, examining the amount of disturbance on the eigenspace of one observable induced by measurements of the other observables. An appealing feature is that the resulting quantity could be estimated by placing the measurements in a superposition of orders, in a similar way as it was done in our paper for the case of two observables.

ACKNOWLEDGEMENTS

We acknowledge support of the Hong Kong Research Grant Council through grants 17326616 and 17300918, and through the Senior Research Fellowship Scheme via SRFS2021-7S02, and support of Croucher foundation and Grant No. 61466 from the John Templeton Foundation as part of the QISS project.

Appendix A: Properties of the MED

Symmetry and nonnegativity. Nonnegativity is immediate from the definition of the MED in the main text, being square root of a probability. Symmetry follows from the symmetry condition $P(A, \mathcal{B}) = P(B, \mathcal{A})$.

Faithfulness. We now show that, for dephasing channels \mathcal{A} and \mathcal{B} , $\text{MED}(\mathcal{A}, \mathcal{B}) > 0$ if and only if the observables A and B are incompatible.

Proof. Recall the definition of MED:

$$\text{MED}(\mathcal{A}, \mathcal{B}) = \sqrt{1 - \text{Prob}(A, \mathcal{B})}, \quad (\text{A1})$$

where

$$\text{Prob}(A, \mathcal{B}) = \frac{1}{d} \sum_{ij} \text{Tr}[P_i Q_j P_i Q_j] \quad (\text{A2})$$

is the probability that a random eigenstate of A is not disturbed by \mathcal{B} .

We now show that $\text{Prob}(A, \mathcal{B}) = 1$ if and only if \mathcal{A} and \mathcal{B} are compatible. Using the Cauchy-Schwarz inequality with

respect to the Hilbert-Schmidt product, we obtain

$$\text{Tr}[P_i Q_j P_i Q_j] \leq \|Q_j P_i\|_2 \cdot \|P_i Q_j\|_2, \quad (\text{A3})$$

where $\|A\|_2 = \sqrt{\text{Tr}[A^\dagger A]}$ is the Hilbert-Schmidt norm. Explicitly, we have

$$\begin{aligned} \|Q_j P_i\|_2^2 &= \text{Tr}[P_i Q_j Q_j P_i] = \text{Tr}[P_i Q_j] \\ \|P_i Q_j\|_2^2 &= \text{Tr}[Q_j P_i P_i Q_j] = \text{Tr}[P_i Q_j], \end{aligned} \quad (\text{A4})$$

where we used the idempotence of P_i and Q_j and the cyclicity of the trace. Inserting the above expressions into Eq. (A3) and summing both sides of over i and j , we obtain

$$\text{Prob}(A, \mathcal{B}) \leq \frac{1}{d} \sum_{i,j} \text{Tr}[P_i Q_j] = 1, \quad (\text{A5})$$

the last equality following from the normalization condition $\sum_i P_i = \sum_j Q_j = I$.

In order for the inequality (A5) to hold with the equality sign, each Cauchy-Schwarz inequality (A3) must hold with the equality sign, which happens if and only if $P_i Q_j$ and $Q_j P_i$ are proportional, i.e., $Q_j P_i = \alpha_{ij} P_i Q_j$ for some $\alpha_{ij} \in \mathbb{C}$. Taking the trace on both sides, we obtain $\text{Tr}[Q_j P_i] = \alpha_{ij} \text{Tr}[Q_j P_i]$, which implies $\alpha_{ij} = 1$ whenever $\text{Tr}[Q_j P_i] \neq 0$. In this case, P_i and Q_j commute. To conclude the proof, recall that $\|Q_j P_i\|_2 = \text{Tr}[Q_j P_i]$ and therefore $\text{Tr}[Q_j P_i] = 0$ implies $Q_j P_i = 0$ and, taking the adjoint on both sides, $P_i Q_j = 0 = Q_j P_i$. Summarizing, the equality $\text{Prob}(A, \mathcal{B})$ holds if and only if the projectors P_i and Q_j commute for every i and j , that is, if and only if the observables A and B are compatible. ■

Maximality for maximally complementary observables. We now show the inequality $\text{MED}(\mathcal{A}, \mathcal{B}) \leq \sqrt{1 - 1/\min\{k_A, k_B\}}$, where $k_{A/B}$ is the number of projectors in the observables A and B , respectively. The maximum value is given by $\text{MED}(\mathcal{A}, \mathcal{B}) = \sqrt{1 - 1/d}$ and attained if and only if A and B are maximally complementary [32] or in other words, their POVM operators are rank-one projectors onto the the basis vectors of two mutually unbiased bases

The proof uses the following lemma from [49]:

Lemma 1 *Let A, B be $n \times n$ Hermitian matrices: If A and B are positive semi-definite, then AB is diagonalizable and has non-negative eigenvalues.*

Proof. Using the definition (A1), we focus on a term $\sum_j \text{Tr}[P_i Q_j P_i Q_j]$, fixing thereby the index i . If P_i is a projector of rank r_i ,

$$\text{rank}(P_i Q_j) < r_i. \quad (\text{A6})$$

Both P_i and Q_j are orthogonal projectors, they are Hermitian and positive semi-definite operators. Therefore, we can apply the following

Using the above lemma, we can diagonalize the operator $P_i Q_j$ as

$$P_i Q_j = X_{ij} \Delta_{ij} X_{ij}^{-1},$$

where Δ_{ij} is a diagonal matrix with non-negative entries. Let r_{ij} be the rank of Δ_{ij} and assume that the first r_{ij} diagonal entries of Δ_{ij} , denoted by $\lambda_{i,j,1}, \dots, \lambda_{i,j,r_{ij}}$, are non-zero. Then, we can write

$$\text{Tr}[P_i Q_j] = \sum_{s=1}^{r_{ij}} \lambda_{i,j,s}. \quad (\text{A7})$$

Taking into account the relation $\sum_{j=1}^{k_B} \text{Tr}[P_i Q_j] = \text{Tr}[P_i] =: r_i$, we find

$$\sum_{j=1}^{k_B} \sum_{s=1}^{r_{ij}} \lambda_{i,j,s} = r_i. \quad (\text{A8})$$

On the other hand,

$$\sum_{j=1}^{k_B} \text{Tr}[P_i Q_j P_i Q_j] = \sum_{j=1}^{k_B} \sum_{s=1}^{r_{ij}} (\lambda_{i,j,s})^2. \quad (\text{A9})$$

The minimum of (A9) under the constraint (A8) can be computed with the method of Lagrange multipliers. Defining the vector λ_i with entries $(\lambda_{i,j,s})_{j \in \{1, \dots, k_B\}, s \in \{1, \dots, r_{ij}\}}$, we find that the minimum is achieved for $\lambda_i = \eta(1, \dots, 1)/2$, where η is the Lagrange multiplier. Plugging the optimal λ_i into (A8), we find $\eta = \frac{2}{k_B}$, and, thus, $\lambda_i = \frac{1}{k_B}(1, \dots, 1)$. Therefore,

$$\sum_{j=1}^{k_B} \text{Tr}[P_i Q_j P_i Q_j] \geq \frac{r_i}{k_B}, \quad (\text{A10})$$

and summing over i ,

$$\sum_{i=1}^{k_A} \sum_{j=1}^{k_B} \text{Tr}[P_i Q_j P_i Q_j] \geq \frac{d}{k_B}. \quad (\text{A11})$$

The lower bound can be achieved only if all the vectors λ_i have constant entries. Since $\text{Tr}[P_i Q_j] = \sum_s \lambda_{i,j,s}$, a necessary condition for achieving the bound is that, for every i , one has

$$\text{Tr}[P_i Q_j] = \text{Tr}[P_i Q_{j'}] \quad \forall j, j' \in \{1, \dots, k_B\}. \quad (\text{A12})$$

Repeating the above argument, this time using the relation $\sum_{i=1}^{k_A} \text{Tr}[P_i Q_j] = \text{Tr}[Q_j] =: s_j$, we can derive the bound

$$\sum_{i=1}^{k_A} \sum_{j=1}^{k_B} \text{Tr}[P_i Q_j P_i Q_j] \geq \frac{d}{k_A}, \quad (\text{A13})$$

which can be achieved only if, for every j ,

$$\text{Tr}[P_i Q_j] = \text{Tr}[P_{i'} Q_j] \quad \forall i, i' \in \{1, \dots, k_A\}. \quad (\text{A14})$$

Combining (A11) and (A13) together, we obtain

$$\sum_{i=1}^{k_A} \sum_{j=1}^{k_B} \text{Tr}[P_i Q_j P_i Q_j] \geq \frac{d}{\min(k_A, k_B)}.$$

and therefore

$$\text{MED}(\mathcal{A}, \mathcal{B}) \leq \sqrt{1 - \frac{1}{\min(k_A, k_B)}}. \quad (\text{A15})$$

The right hand side is maximized for $k_A = k_B = d$, for which we have

$$\text{MED}(\mathcal{A}, \mathcal{B}) \leq \sqrt{1 - \frac{1}{d}}. \quad (\text{A16})$$

We now determine when the bound is achieved. First, note that the bound can only be achieved by rank-1 projectors, because the number of outcomes of the two measurements must be d . Now, Eqs. (A12) and (A14) imply

$$\begin{aligned} \text{Tr}[P_i Q_j] &= \text{Tr}[P_{i'} Q_{j'}] & \forall i, i' \in \{1, \dots, k_A\} \\ & & \forall j, j' \in \{1, \dots, k_B\}, \end{aligned} \quad (\text{A17})$$

that is, $\text{Tr}[P_i Q_j] = 1/d$ for every i, j . Since P_i and Q_j are rank-1 projectors, this condition implies that they are projectors on the basis vectors of two mutually unbiased bases.

Monotonicity under coarse-graining. Let $\mathbf{P}' := (P'_k)_{k=1}^{o'}$ be a coarse-graining of the measurement $\mathbf{P} = (P_i)_{i=1}^o$, meaning that there is a surjective function $f : \{1, \dots, o\} \rightarrow \{1, \dots, o'\}$ such that $P'_k = \sum_{i:f(i)=k} P_i$. Note that, by definition, one has the operator inequality $P_i \leq P'_{f(i)}$. Hence, for every set of operators (D_j) , one has the bound

$$\begin{aligned} \sum_{i,j} \text{Tr}[P_i Q_j P_i Q_j] &\leq \sum_{i,j} \text{Tr}[P'_{f(i)} Q_j P_i Q_j] \\ &= \sum_{k,j} \text{Tr}[P'_k Q_j P'_k Q_j], \end{aligned} \quad (\text{A18})$$

which implies the inequality $\text{MED}(\mathcal{A}, \mathcal{B}) \geq \text{MED}(\mathcal{A}', \mathcal{B})$, where \mathcal{A}' is the dephasing channel associated to the coarse-grained measurement \mathbf{P}' , namely $\mathcal{A}'(\rho) = \sum_k P'_k \rho P'_k$.

Metric on von Neumann measurements. We now show that MED is a metric on the set of rank-one projective measurements, which are in one-to-one correspondence with dephasing channels are of the form $\mathcal{A}(\rho) = \sum_{i=1}^d \langle \alpha_i | \rho | \alpha_i \rangle | \alpha_i \rangle \langle \alpha_i |$ for some orthonormal basis $\{|\alpha_i\rangle\}_{i=1}^d$. Specifically, we show the MED for dephasing channels \mathcal{A} and \mathcal{B} associated to rank-one measurements has the expression

$$\text{MED}(\mathcal{A}, \mathcal{B}) = \frac{1}{\sqrt{2d}} \|\tau_{\mathcal{A}} - \tau_{\mathcal{B}}\|_2, \quad (\text{A19})$$

where $\|X\|_2 = \sqrt{\text{Tr}[X^\dagger X]}$ is the Hilbert-Schmidt norm, and $\tau_{\mathcal{A}}$ and $\tau_{\mathcal{B}}$ are the Choi operators of \mathcal{A} and \mathcal{B} , given by

$$\begin{aligned} \tau_{\mathcal{A}} &:= \sum_{m,n} \mathcal{A}(|m\rangle\langle n|) \otimes |m\rangle\langle n| = \sum_i |\alpha_i\rangle\langle\alpha_i| \otimes |\bar{\alpha}_i\rangle\langle\bar{\alpha}_i| \\ \tau_{\mathcal{B}} &:= \sum_{m,n} \mathcal{B}(|m\rangle\langle n|) \otimes |m\rangle\langle n| = \sum_j |\beta_j\rangle\langle\beta_j| \otimes |\bar{\beta}_j\rangle\langle\bar{\beta}_j|, \end{aligned}$$

$|\bar{\psi}\rangle$ being the complex conjugate of a generic vector $|\psi\rangle$.

Proof. Denoting by $\langle X, Y \rangle = \text{Tr}[X^\dagger Y]$ the Hilbert-Schmidt product of two operators X and Y , we have

$$\begin{aligned} \|\tau_{\mathcal{A}} - \tau_{\mathcal{B}}\|_2^2 &= \langle \tau_{\mathcal{A}} - \tau_{\mathcal{B}}, \tau_{\mathcal{A}} - \tau_{\mathcal{B}} \rangle \\ &= \langle \tau_{\mathcal{A}}, \tau_{\mathcal{A}} \rangle + \langle \tau_{\mathcal{B}}, \tau_{\mathcal{B}} \rangle - 2\langle \tau_{\mathcal{A}}, \tau_{\mathcal{B}} \rangle \end{aligned} \quad (\text{A20})$$

and

$$\begin{aligned} \langle \tau_{\mathcal{A}}, \tau_{\mathcal{A}} \rangle &= \sum_{i,i'} \text{Tr} ([|\alpha_i\rangle\langle\alpha_i| \otimes |\bar{\alpha}_i\rangle\langle\bar{\alpha}_i|] (|\alpha'_i\rangle\langle\alpha'_i| \otimes |\bar{\alpha}'_i\rangle\langle\bar{\alpha}'_i|)) \\ &= \sum_i \text{Tr} [|\alpha_i\rangle\langle\alpha_i| \otimes |\bar{\alpha}_i\rangle\langle\bar{\alpha}_i|] \\ &= d \\ &= \langle \tau_{\mathcal{B}}, \tau_{\mathcal{B}} \rangle, \end{aligned} \quad (\text{A21})$$

and

$$\begin{aligned} \langle \tau_{\mathcal{A}}, \tau_{\mathcal{B}} \rangle &= \text{Tr} [\tau_{\mathcal{A}}^\dagger \tau_{\mathcal{B}}] \\ &= \sum_{i,j} \text{Tr} [|\alpha_i\rangle\langle\alpha_i| \beta_j \rangle \langle \beta_j| \otimes |\bar{\alpha}_i\rangle\langle\bar{\alpha}_i| \bar{\beta}_j \rangle \langle \bar{\beta}_j|] \\ &= \sum_{i,j} |\langle \alpha_i | \beta_j \rangle|^4 \\ &= \sum_{i,j} \text{Tr} [P_i Q_j P_i Q_j] \end{aligned} \quad (\text{A22})$$

Hence,

$$\begin{aligned} \|\tau_{\mathcal{A}} - \tau_{\mathcal{B}}\|_2^2 &= 2d - 2 \sum_{i,j} \text{Tr} [P_i Q_j P_i Q_j] \\ &= 2d (\text{MED}(\mathcal{A}, \mathcal{B}))^2 \end{aligned} \quad (\text{A23})$$

Eq. (A23) shows that $\text{MED}(\mathcal{A}, \mathcal{B})$ coincides with the Hilbert-Schmidt norm $\|\tau_{\mathcal{A}} - \tau_{\mathcal{B}}\|_2$ up to a constant factor. ■

Robustness to noise. The MED remains a faithful measure of incompatibility even if the canonical channels \mathcal{A} and \mathcal{B} are replaced by quantum channels \mathcal{A}' and \mathcal{B}' associated to noisy measurements of \mathcal{A} and \mathcal{B} , that is, channels of the form $\mathcal{A}' = \sum_{i,k} A_{i,k} \rho A_{i,k}^\dagger$ and $\mathcal{B}'(\rho) = \sum_{j,l} B_{j,l} \rho B_{j,l}^\dagger$, with $\sum_k A_{i,k}^\dagger A_{i,k} = (1 - \lambda) P_i + \lambda p_i I$ and $\sum_l B_{j,l}^\dagger B_{j,l} = (1 - \mu) Q_j + \mu q_j I$, with suitable probabilities $\lambda, \mu \in [0, 1]$ and suitable probability distributions \mathbf{p} and \mathbf{q} . To see this fact, consider the disturbance of a noisy measurement of B on the eigenspaces of A . Adopting the same settings of the main text, the average probability of no eigenspace disturbance is given by

$$\begin{aligned} \text{Prob}(A, \mathcal{B}') &= \frac{1}{d} \sum_{i,j,l} \text{Tr} [P_i B_{j,l} P_i B_{j,l}^\dagger] \\ &= 1 - \sum_{i,j,l} \frac{\text{Tr} [|P_i, B_{j,l}|]^2}{2d}. \end{aligned} \quad (\text{A24})$$

In order for this probability to be 1, each commutator must be zero. Hence, one must have $[P_i, Q_j] = \sum_l [P_i, B_{j,l}^\dagger B_{j,l}] = 0$ for every i and j , that is, the original noiseless observables must be compatible.

The same argument applies for the disturbance of a noisy measurement of A on the eigenspaces of B .

Appendix B: Performance of the protocol

In each run on the estimation protocol in the main text, an outcome of the measurement of the control qubit is obtained in a constant time, and it is possible to estimate $p(-)$ (and, hence, $\text{MED}(\mathcal{A}, \mathcal{B})$) with an arbitrary accuracy by repeating it. In particular, the number of accesses to the measurement devices is independent of the dimension d and inverse polynomial on the accuracy we need. In order to show it, we model the result of the quantum Fourier measurement as a Bernoulli variable. A series of such results could be modelled as random variables X_1, X_2, \dots, X_n , where $\forall 1 \leq i \leq n, X_i \stackrel{\text{i.i.d.}}{\sim} \text{Ber}(p)$,

$$S = \sum_i X_i \sim \text{Bin}(n, p),$$

and the Hoeffding's inequality implies the following bound,

$$\forall \epsilon > 0, P \left(\left| \frac{S}{n} - p \right| < \epsilon \right) \geq 1 - 2e^{-2\epsilon^2 n}.$$

Now, we want to guarantee that with a probability arbitrarily close to 1, namely, $1 - \delta$, our estimated value deviates within an error of ϵ . From the above inequality, we find that the minimum number of repetitions of the experiment needed to achieve error parameters ϵ and δ is

$$n(\epsilon, \delta) = -\frac{1}{2\epsilon^2} \log \frac{\delta}{2}. \quad (\text{B1})$$

After multiplying by a constant term, we conclude that the sample complexity of the estimation protocol using quantum switches is of $O \left(\frac{1}{\epsilon^2} \log \frac{\delta}{2} \right)$.

[1] N. Bohr, *The Quantum Postulate and the Recent Development of Atomic Theory*, Nature **121**, 580 (1928).

[2] W. Heisenberg, *Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik*, Z. Phys. **43**, 172 (1927).

- [3] T. Heinosaari, T. Miyadera, and M. Ziman, *An invitation to quantum incompatibility*, J. Phys. A: Math. Theor. **49**, 123001 (2016).
- [4] C. H. Bennett and G. Brassard, *Quantum cryptography: Public key distribution and coin tossing*, Theor. Comput. Sci. **560**, 7 (2014).
- [5] F. Buscemi, E. Chitambar, and W. Zhou, *Complete Resource Theory of Quantum Incompatibility as Quantum Programmability*, Phys. Rev. Lett. **124**, 120401 (2020).
- [6] P. Busch, T. Heinosaari, J. Schultz, and N. Stevens, *Comparing the degrees of incompatibility inherent in probabilistic physical theories*, Europhys. Lett. **103**, 10002 (2013).
- [7] T. Heinosaari, J. Schultz, A. Toigo, and M. Ziman, *Maximally incompatible quantum observables*, Phys. Lett. A **378**, 1695 (2014).
- [8] S. Bandyopadhyay and P. Mandayam, *Operational measure of incompatibility of noncommuting observables*, Phys. Rev. A **87**, 042120 (2013).
- [9] P. Mandayam and M. Srinivas, *Measures of disturbance and incompatibility for quantum measurements*, Phys. Rev. A **89**, 062112 (2014).
- [10] P. Busch, P. J. Lahti, and P. Mittelstaedt, *The quantum theory of measurement* (Springer, 1996).
- [11] G. Chiribella, G. M. D’Ariano, P. Perinotti, and B. Valiron, *Beyond quantum computers*, arXiv:0912.0195 [quant-ph].
- [12] G. Chiribella, G. M. D’Ariano, P. Perinotti, and B. Valiron, *Quantum computations without definite causal structure*, Phys. Rev. A **88**, 022318 (2013).
- [13] L. M. Procopio, A. Moqanaki, M. Araújo, F. Costa, I. Alonso Calafell, E. G. Dowd, D. R. Hamel, L. A. Rozema, Č. Brukner, and P. Walther, *Experimental superposition of orders of quantum gates*, Nat. Commun. **6**, 7913 (2015).
- [14] G. Rubino, L. A. Rozema, A. Feix, M. Araújo, J. M. Zeuner, L. M. Procopio, Č. Brukner, and P. Walther, *Experimental verification of an indefinite causal order*, Sci. Adv. **3**, e1602589 (2017).
- [15] K. Goswami, C. Giarmatzi, M. Kewming, F. Costa, C. Branciard, J. Romero, and A. G. White, *Indefinite Causal Order in a Quantum Switch*, Phys. Rev. Lett. **121**, 090503 (2018).
- [16] K. Wei, N. Tischler, S.-R. Zhao, Y.-H. Li, J. M. Arrazola, Y. Liu, W. Zhang, H. Li, L. You, Z. Wang, Y.-A. Chen, B. C. Sanders, Q. Zhang, G. J. Pryde, F. Xu, and J.-W. Pan, *Experimental Quantum Switching for Exponentially Superior Quantum Communication Complexity*, Phys. Rev. Lett. **122**, 120504 (2019).
- [17] Y. Guo, X.-M. Hu, Z.-B. Hou, H. Cao, J.-M. Cui, B.-H. Liu, Y.-F. Huang, C.-F. Li, G.-C. Guo, and G. Chiribella, *Experimental Transmission of Quantum Information Using a Superposition of Causal Orders*, Phys. Rev. Lett. **124**, 030502 (2020).
- [18] K. Goswami, Y. Cao, G. Paz-Silva, J. Romero, and A. White, *Increasing communication capacity via superposition of order*, Phys. Rev. Res. **2**, 033292 (2020).
- [19] G. Rubino, L. A. Rozema, D. Ebler, H. Kristjánsson, S. Salek, P. Allard Guérin, A. A. Abbott, C. Branciard, Č. Brukner, G. Chiribella, and P. Walther, *Experimental quantum communication enhancement by superposing trajectories*, Phys. Rev. Research **3**, 013093 (2021).
- [20] K. Goswami and J. Romero, *Experiments on quantum causality*, AVS Quantum Science **2**, 037101 (2020).
- [21] N. Friis, A. A. Melnikov, G. Kirchmair, and H. J. Briegel, *Coherent controlization using superconducting qubits*, Sci. Rep. **5**, 18036 (2015).
- [22] E. Specker, *Die Logik nicht gleichzeitig entscheidbarer Aussagen*, Dialectica **14**, 239 (1960).
- [23] S. Kochen and E. P. Specker, *The problem of hidden variables in quantum mechanics*, J. Math. Mech. **17**, 59 (1967).
- [24] C. Budroni, A. Cabello, O. Gühne, M. Kleinmann, and J.-Å. Larsson, *Quantum contextuality*, arXiv:2102.13036 (2021).
- [25] S. Abramsky and A. Brandenburger, *The sheaf-theoretic structure of non-locality and contextuality*, New J. Phys. **13**, 113036 (2011).
- [26] B. d’Espagnat, *Conceptual foundations of quantum mechanics* (CRC Press, 2018).
- [27] M. Schlosshauer, *Decoherence, the measurement problem, and interpretations of quantum mechanics*, Rev. Mod. Phys. **76**, 1267 (2005).
- [28] P. Grangier, *Contextual objectivity: a realistic interpretation of quantum mechanics*, Eur. J. Phys. **23**, 331 (2002).
- [29] W. H. Zurek, *Quantum Darwinism*, Nat. Phys. **5**, 181 (2009).
- [30] R. Horodecki, J. Korbicz, and P. Horodecki, *Quantum origins of objectivity*, Phys. Rev. A **91**, 032122 (2015).
- [31] P. Lahti, *Coexistence and joint measurability in quantum mechanics*, Int. J. Theor. Phys. **42**, 893 (2003).
- [32] K. Kraus, *Complementary observables and uncertainty relations*, Phys. Rev. D **35**, 3070 (1987).
- [33] I. Ivonovic, *Geometrical description of quantal state determination*, J. Phys. A: Math. Gen. **14**, 3241 (1981).
- [34] W. K. Wootters and B. D. Fields, *Optimal state-determination by mutually unbiased measurements*, Annals of Physics **191**, 363 (1989).
- [35] I. L. Chuang and M. A. Nielsen, *Prescription for experimental determination of the dynamics of a quantum black box*, J. Mod. Opt. **44**, 2455 (1997).
- [36] J. Poyatos, J. I. Cirac, and P. Zoller, *Complete characterization of a quantum process: the two-bit quantum gate*, Phys. Rev. Lett. **78**, 390 (1997).
- [37] D. W. Leung, *Choi’s proof as a recipe for quantum process tomography*, J. Math. Phys. **44**, 528 (2003).
- [38] G. D’Ariano and P. L. Presti, *Quantum tomography for measuring experimentally the matrix elements of an arbitrary quantum operation*, Phys. Rev. Lett. **86**, 4195 (2001).
- [39] W. Dür and J. Cirac, *Nonlocal operations: Purification, storage, compression, tomography, and probabilistic implementation*, Phys. Rev. A **64**, 012317 (2001).
- [40] J. B. Altepeter, D. Branning, E. Jeffrey, T. Wei, P. G. Kwiat, R. T. Thew, J. L. O’Brien, M. A. Nielsen, and A. G. White, *Ancilla-assisted quantum process tomography*, Phys. Rev. Lett. **90**, 193601 (2003).
- [41] G. M. D’Ariano, L. Maccone, and P. L. Presti, *Quantum calibration of measurement instrumentation*, Phys. Rev. Lett. **93**, 250407 (2004).
- [42] G. Chiribella, *Perfect discrimination of no-signalling channels via quantum superposition of causal structures*, Phys. Rev. A **86**, 040301(R) (2012).
- [43] E. Andersson, J. Bergou, and I. Jex, *Comparison of unitary transforms using Franson interferometry*, J. Mod. Opt. **52**, 1485 (2005).
- [44] N. Loizeau and A. Grinbaum, *Channel capacity enhancement with indefinite causal order*, Phys. Rev. A **101**, 012340 (2020).
- [45] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, and E. Duchesnay, *Scikit-learn: Machine Learning in Python*, J. Mach. Learn. Res. **12**, 2825 (2011).
- [46] D. Arthur and S. Vassilvitskii, *k-means++: the advantages of careful seeding*, in *SODA ’07: Proceedings of the eighteenth annual ACM-SIAM symposium on Discrete algorithms* (Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 2007) pp. 1027–1035.
- [47] M. Dall’Arno, G. M. D’Ariano, and M. F. Sacchi, *Purification of noisy quantum measurements*, Phys. Rev. A **82**, 042315 (2010).

- [48] M. Ozawa, *Quantum measuring processes of continuous observables*, J. Math. Phys. **25**, 79 (1984).
- [49] R. A. Horn and C. R. Johnson, *Matrix Analysis* (Cambridge University Press, 1990).