

On the non-measurability of ω -categorical Hrushovski constructions

Paolo Marimon

August 15, 2022

Abstract

We study ω -categorical MS -measurable structures. Our main result is that a class of ω -categorical Hrushovski constructions, supersimple of finite SU -rank is not MS -measurable. These results complement the work of Evans [5] on a conjecture of Macpherson and Elwes [2]. Differently from Evans' work, our structures may satisfy independent n -amalgamation for all n . We also prove some general results in the context of ω -categorical MS -measurable structures. Firstly, in these structures the dimension in the MS -dimension-measure can be chosen to be SU -rank. Secondly, non-forking independence implies a form of probabilistic independence in the measure. The latter follows from more general unpublished results of Hrushovski [10], but we provide a self-contained proof.

1 Introduction

This work focuses on ω -categorical MS -measurable structures and on the role of supersimple ω -categorical Hrushovski constructions in settling some conjectures about them. In particular, we prove that a certain class of ω -categorical supersimple finite SU -rank Hrushovski constructions is not MS -measurable. The question of whether in general such structures are not MS -measurable is still open.

Let \mathcal{L} be a first order language. A complete \mathcal{L} -theory is **ω -categorical** if it has a unique countable model up to isomorphism. We say that an \mathcal{L} -structure is ω -categorical if its theory is. We define MS -measurable structures in Definition 2.6. But the basic idea of an MS -measurable structure is that we may associate with it a dimension-measure function on its definable sets, where for the sets of a given dimension there is an invariant, additive measure satisfying Fubini's theorem with respect to the dimension and giving such sets positive measures. Standard examples of MS -measurable structures are pseudofinite fields, the random graph, and ω -categorical ω -stable structures [2]. In this article we focus on whether ω -categorical Hrushovski constructions are MS -measurable. These are a class of relational structures, which generalise Fraïssé limits, and whose construction depends on a choice of a parameter $\alpha \in \mathbb{R}^{>0}$ and of a non-decreasing function $f : \mathbb{R}^{>0} \rightarrow \mathbb{R}^{>0}$. We discuss their construction in section 4.

In their review article on MS -measurable structures [2], Elwes and Macpherson ask the following questions which require the study of ω -categorical Hrushovski constructions:

Question 1.1. Is there any ω -categorical supersimple theory of finite SU -rank which is not MS -measurable?

Question 1.2. Is every ω -categorical measurable structure one-based?

Being MS -measurable implies being supersimple of finite SU -rank [2]. Furthermore, in such structures D -rank, $S1$ -rank and SU -rank are the same. Hence the first question simply consists in asking whether supersimplicity and finite rank imply MS -measurability in an

ω -categorical context.

Supersimple ω -categorical Hrushovski constructions of finite SU -rank are essential structures for the study of these questions. These are the only examples of ω -categorical supersimple not one-based structures which we know of. Hence, finding out that any of these structures are not MS -measurable would answer the first question positively. Finding out that any of these structures is MS -measurable would answer negatively the second question. We already know that some ω -categorical Hrushovski constructions are not MS -measurable from the work of Evans [5]. There, Evans first proves that any MS -measurable structure must satisfy a weak form of independent n -amalgamation using a version by Towsner of the Hypergraph Removal Lemma [20][21][7][18]. Then, he proves that for a class of Hrushovski constructions (with a ternary relation and $\alpha = 1$), any dimension is a scaled version of the natural Hrushovski dimension. With these tools, Evans shows that some such structures do not satisfy the weak n -amalgamation property.

While Evans' dimension theorem can be generalised to other classes of Hrushovski constructions, it is possible for Hrushovski constructions to satisfy the weak amalgamation condition (and even independent n amalgamation for all n) as long as f is slow-growing enough [12]. Indeed, a natural question which arises from Evans' paper is whether MS -measurability for an ω -categorical finite rank structure is implied by satisfying some strong enough form of independent n -amalgamation.

In this paper we develop new tools to study ω -categorical MS -measurable structures. These will allow us to study how an MS -dimension-measure would behave in an ω -categorical Hrushovski construction if it was MS -measurable. In particular, in Section 5 we introduce the ω -categorical supersimple finite rank Hrushovski constructions \mathcal{M}_f for which we prove the main theorem of this paper:

Theorem 5.9. *The structures \mathcal{M}_f satisfying the conditions of Construction 5.1 are not MS -measurable. Hence, there are ω -categorical supersimple finite SU -rank structures which are not MS -measurable. Indeed, there are supersimple ω -categorical structures of finite SU -rank and with independent n -amalgamation for all n which are not MS -measurable.*

The proof of this result takes all of subsection 5.2.

The structure of this paper is as follows. In Section 2 we introduce some basic notions concerning MS -measurable structures. We also prove the folklore theorem that the natural notion of independence in MS -measurable structures corresponds to non-forking independence. In Corollary 2.20 we find a set of equations that the measure of an MS -measurable structure must satisfy. We also prove that in an ω -categorical context, we may take the dimension of an MS -measurable structure to be SU -rank (Corollary 2.22). This allows us to circumvent the need to prove a dimension theorem for the class of Hrushovski constructions we will consider.

Section 3 yields another set of equations for the measure in ω -categorical MS -measurable structures which show how non-forking independence yields probabilistic independence in the measure (Theorem 3.5). These results are a special case of the probabilistic independence theorem in the unpublished [10].

In Section 4 we introduce Hrushovski constructions and in subsection 4.2 we specify the form of the equations from section 2 in the context of such structures.

In Section 5 we introduce in Remark 5.1 various supersimple ω -categorical Hrushovski constructions of finite SU -rank, some of which satisfy independent n -amalgamation for all n . We will prove these are not MS -measurable in subsection 5.2. Various properties of these structures (especially supersimplicity) are proven in the appendix. The most relevant results for understanding the proof of our main theorem are Corollaries 2.20 and 2.22, Theorems 3.5 and 4.8, and subsection 5.2.

I assume some basic knowledge of model theory, especially regarding ω -categorical structures. All of the relevant material is covered in Chapters 1 to 4 of Tent and Ziegler's book [19]. Section 3 uses some local stability theory for which the first chapter of [17] is sufficient. We work in countable languages and only with complete theories. We shall make frequent use of the equivalent conditions to ω -categoricity from the Ryll-Nardzewski theorem. We also assume some basic knowledge of simple theories and SU -rank. Chapter 2 of [13] covers the relevant material.

Regarding notation, for a language \mathcal{L} , we work on an \mathcal{L} -structure \mathcal{M} and write M to denote the underlying set. The monster model for the \mathcal{L} -theory of \mathcal{M} is denoted by \mathbb{M} . We use overlined lowercase letters at the beginning of the alphabet \bar{a}, \bar{b}, \dots to denote finite tuples from a model, and use \bar{x}, \bar{y}, \dots to denote tuples of variables. We use the non-overlined versions when speaking of 1-tuples is sufficient. We use letters A, B, C, \dots to denote (usually finite) subsets of our model. We use the greek letters $\phi, \psi, \chi \dots$ to denote \mathcal{L} -formulas, which we often write in the form $\phi(\bar{x}, \bar{y})$ to specify their free variables.

2 Measurable ω -categorical structures

In this section we introduce MS -measurable structures, and some basic results about them. We also provide some original results in the last part of this section regarding ω -categorical MS -measurable structures.

We begin with Subsection 2.1, where we introduce the notion of MS -measurability following [15] and [2]. Then, in Subsection 2.2 we prove the folklore result that dimension independence in MS -measurable structures corresponds to non-forking independence. Finally, in Subsection 2.3, we find a set of equations which hold in ω -categorical MS -measurable structures. We also show that if an ω -categorical structure is MS -measurable, then it is MS -measurable with dimension given by SU -rank. This will allow us to avoid proving Evans' dimension theorem for the class of ω -categorical Hrushovski constructions we will consider.

2.1 Basic definitions

Let \mathcal{M} be a structure. Recall that by $\text{Def}(M)$ we mean the set of non-empty definable subsets of M (defined by formulas with parameters from M). Meanwhile, $\text{Def}_{\bar{x}}(M)$ denotes the definable subsets of M in the variable \bar{x} . For $\bar{a}, \bar{a}' \in M^{|\bar{y}|}$ and $A \subset M$, we write $\bar{a} \equiv_A \bar{a}'$ to say that \bar{a} and \bar{a}' have the same type over A . If $A = \emptyset$, we simply write $\bar{a} \equiv \bar{a}'$.

Definition 2.1. Let X be any set and consider a function $g : \text{Def}(M) \rightarrow X$. For $A \subseteq M$, we say that g is **A -invariant** if $g(\phi(M^{\bar{x}}, \bar{a})) = g(\phi(M^{\bar{x}}, \bar{a}'))$ whenever $\bar{a} \equiv_A \bar{a}'$. When $A = \emptyset$, we say g is **invariant**. The function g is **definable** if for any formula $\phi(\bar{x}, \bar{y})$ and $k \in X$, the set of $\bar{a} \in M^{|\bar{y}|}$ such that $g(\phi(M^{\bar{x}}, \bar{a})) = k$ is definable over the empty set. We say that g is **finite** if the set of values of $g(\phi(M^{\bar{x}}, \bar{a}))$ for $\bar{a} \in M^{|\bar{y}|}$ is finite.

To avoid cumbersome notation, when the model \mathcal{M} is clear, we sometimes write $g(\phi(\bar{x}, \bar{a}))$ instead of $g(\phi(M^{\bar{x}}, \bar{a}))$.

Note that definability always implies invariance. In an ω -categorical context, by Ryll-Nardzewski, invariance implies definability and finiteness.

Before introducing the notion of an MS -measurable structure we follow the notation of [22] to speak of a dimension function:

Definition 2.2. We call $\dim : \text{Def}(M) \rightarrow \mathbb{N}$ a **dimension** if it satisfies the following conditions:

- (Algebraicity) for X finite and non-empty, $\dim(X) = 0$;
- (Union) For $X, Y \in \text{Def}(M)$, $\dim(X \cup Y) = \max\{\dim(X), \dim(Y)\}$; and

- (Additivity) For finite tuples $\bar{a}, \bar{b}, \bar{c}$ from M , $\dim(\bar{a}\bar{b}/\bar{c}) = \dim(\bar{a}/\bar{b}\bar{c}) + \dim(\bar{b}/\bar{c})$.

Remark 2.3. In the additivity condition, by $\dim(\bar{a}/B)$ we mean $\dim(\text{tp}(\bar{a}/B))$. For a partial type $\pi(\bar{x})$ over $B \subseteq M$, we define

$$\dim(\pi(\bar{x})) = \min\{\dim(\phi(M^{|\bar{x}|}, \bar{b}) \mid \pi(\bar{x}) \vdash \phi(\bar{x}, \bar{b}))\}.$$

Remark 2.4. In the course of this paper we work with definable dimensions in an ω -categorical context. Hence, for \bar{a}, \bar{b} finite tuples from M we have that $\dim(\bar{a}/\bar{b}) = \dim(\phi(M^{|\bar{x}|}, \bar{b}))$, where $\phi(\bar{x}, \bar{b})$ is a formula isolating $\text{tp}(\bar{a}/\bar{b})$.

Definition 2.5. We say that the function $\mu : \text{Def}_{\bar{x}}(M) \rightarrow \mathbb{R}^{\geq 0} \cup \{\infty\}$ is a **measure** if it is a finitely additive function, i.e. for X and Y definable and disjoint in the same variable,

$$\mu(X \cup Y) = \mu(X) + \mu(Y).$$

We say μ is a **Keisler measure** if it takes values in $[0, 1]$.

We give here a slightly simplified definition of *MS*-measurability than the original following [15][Proposition 5.7]:

Definition 2.6. The \mathcal{L} -structure \mathcal{M} is **MS-measurable** if there is a dimension-measure function $h : \text{Def}(\mathcal{M}) \rightarrow \mathbb{N} \times \mathbb{R}^{>0}$, with notation $h(X) = (d(X), \mu(X))$, satisfying the following conditions:

1. The function h is finite and definable.
2. For $\bar{a} \in M^n$, $h(\{\bar{a}\}) = (0, 1)$.
3. (Additivity) For $X, Y \subseteq M^n$ definable and disjoint,

$$\mu(X \cup Y) = \begin{cases} \mu(X) + \mu(Y), & \text{for } d(X) = d(Y); \\ \mu(Y) & \text{for } d(X) < d(Y). \end{cases}$$

4. (Fubini) For $n \geq 2$, let $X \subseteq M^n$ be definable and for $1 \leq m < n$, let $\pi : M^n \rightarrow M^m$ be a projection of M^n to m -many coordinates. Suppose there is $(k, \nu) \in \mathbb{N} \times \mathbb{R}^{>0}$ such that for any $\bar{a} \in \pi(X)$, we have that $h(\pi^{-1}(\bar{a}) \cap X) = (k, \nu)$. Then, $d(X) = d(\pi(X)) + k$ and $\mu(X) = \mu(\pi(X))\nu$.

Remark 2.7. From the definition it follows that being *MS*-measurable is a property of a theory, i.e. if \mathcal{M} is *MS*-measurable, then so is any elementarily equivalent structure. We say that a many sorted structure \mathcal{M}^* is *MS*-measurable if every restriction of \mathcal{M}^* to finitely many sorts is *MS*-measurable. We have that if \mathcal{M} is *MS*-measurable, then so is \mathcal{M}^{eq} [15][Proposition 5.10].

Remark 2.8. An important, and somehow hidden, assumption of the definition of *MS*-measurability is that of **positivity**. That is, for $h(X) = (k, \nu)$, $\nu > 0$. This will be vital to our proof of non-measurability since we will show that a given definable set must be assigned measure zero.

Remark 2.9. As noted in [15][Proposition 5.3], for A -definable X , we have an induced A -invariant probability measure on the definable subsets of X .

Remark 2.10. The dimension part of an *MS*-function is a definable dimension function, as defined above. Additivity follows from [22][p.919].

In an *MS*-measurable context, for $\pi(\bar{x})$ a partial type over $B \subseteq M$, we define

$$\mu(\pi(\bar{x})) = \inf\{\mu(\phi(\bar{x}, \bar{b})) \mid \pi(\bar{x}) \vdash \phi(\bar{x}, \bar{b}), d(\phi(\bar{x}, \bar{b})) = d(\pi(\bar{x}))\},$$

and we write $\mu(\bar{a}/B)$ for $\mu(\text{tp}(\bar{a}/B))$. For an ω -categorical *MS*-measurable structure, $\mu(\bar{a}/\bar{b}) = \mu(\phi(M^{|\bar{x}|}, \bar{b}))$, for any formula $\phi(\bar{x}, \bar{b})$ isolating $\text{tp}(\bar{a}/\bar{b})$.

Remark 2.11. When speaking of the dimension and measure of types we shall also set the convention that $d(\emptyset) = 0$ and $\mu(\emptyset) = 1$. This is helpful to treat the case of types over \emptyset analogously to that of types over sets of parameters when dealing with Fubini. In fact, we shall make frequent use of the fact that in an ω -categorical MS-measurable structure Fubini implies that

$$h(\bar{a}\bar{b}) = (\dim(\bar{a}\bar{b}), \mu(\bar{a}\bar{b})) = (\dim(\bar{a}/\bar{b}) + \dim(\bar{b}), \mu(\bar{a}/\bar{b})\mu(\bar{b})).$$

Hence, we adopt the conventions $d(\emptyset) = 0$ and $\mu(\emptyset) = 1$ so that we may write

$$\dim(\bar{a}) = \dim(\bar{a}/\emptyset) + \dim(\emptyset) = \dim(\bar{a}/\emptyset), \text{ and}$$

$$\mu(\bar{a}) = \mu(\bar{a}/\emptyset)\mu(\emptyset) = \mu(\bar{a}/\emptyset).$$

2.2 Dimension-independence

It is easy to obtain a notion of independence from an invariant dimension. In particular, in an MS-measurable structure the dimension-independence corresponds to non-forking independence. In [2] this result is attributed to unpublished work of Ben-Yaacov. In this section we prove it briefly. In the next section this will help us showing that in ω -categorical MS-measurable structures we may take the dimension to be SU-rank without harm.

Definition 2.12. Let $d : \text{Dim}(\mathbb{M}) \rightarrow \mathbb{N}$ be an invariant dimension, where \mathbb{M} is a monster model. Let \bar{a} be a tuple and B, C be small subsets of \mathbb{M} . We say that \bar{a} is d -independent from B over C , writing $\bar{a} \downarrow_C^d B$ if

$$d(\bar{a}/BC) = d(\bar{a}/C).$$

The following is easy to prove from the basic properties of an invariant dimension:

Proposition 2.13. *Let $d : \text{Dim}(\mathbb{M}) \rightarrow \mathbb{N}$ be an invariant dimension. The relation of d -independence is a notion of independence in the sense of [14][Def. 4.1].*

Hence, in an MS-measurable context the dimension part of an MS-dimension-measure yields a notion of independence. We shall prove this coincides with non-forking independence. In order to prove this, first recall Lemma 3.5 from [2]:

Lemma 2.14. *Let \mathcal{M} be MS-measurable. Let $X \subseteq M^{|\bar{x}|}$ be definable and $\phi(\bar{x}, \bar{y})$ be such that there is an indiscernible sequence $(\bar{b}_i)_{i < \omega}$ such that $\{\phi(\bar{x}, \bar{b}_i) \mid i < \omega\}$ is inconsistent and $\phi(M^{|\bar{x}|}, \bar{b}_i) \subseteq X$ for each $i < \omega$. Then, $d(X) > d(\phi(\bar{x}, \bar{b}_i))$.*

From this result it is easy to prove that dimension independence implies non-forking independence.

Proposition 2.15. *Suppose that $\bar{a} \downarrow_C^d B$. Then $\text{tp}(\bar{a}/BC)$ does not divide over C . Hence, by simplicity, $\bar{a} \downarrow_C^d B$ implies $\bar{a} \downarrow_C^f B$, where \downarrow_C^f is non-forking independence.*

Proof. Suppose by contrapositive that $\text{tp}(\bar{a}/BC)$ divides over C . So there is $\phi(\bar{x}, \bar{d}) \in \text{tp}(\bar{a}/BC)$ such that $\{\phi(\bar{x}, \bar{d}_i) \mid i < \omega\}$ is k -inconsistent and $(\bar{d}_i)_{i < \omega}$ is C -indiscernible with $\bar{d}_0 = \bar{d}$. Let $\chi(\bar{x})$ be a formula defined over C witnessing $d(\bar{a}/C) = k$. Consider $\phi(\bar{x}, \bar{d}_i) \wedge \chi(\bar{x})$. Since $\phi(\bar{x}, \bar{d}) \in \text{tp}(\bar{a}/BC)$, we have that $\models \exists \bar{x} \phi(\bar{x}, \bar{d}) \wedge \chi(\bar{x})$, and since $\bar{d}_i \equiv_C \bar{d}$, $\models \exists \bar{x} \phi(\bar{x}, \bar{d}_i) \wedge \chi(\bar{x})$. Hence, $\phi(\mathbb{M}^{|\bar{x}|}, \bar{d}_i) \wedge \chi(\mathbb{M}^{|\bar{x}|}) \subseteq \chi(\mathbb{M}^{|\bar{x}|})$, and $\{\phi(\bar{x}, \bar{d}_i) \wedge \chi(\bar{x}) \mid i < \omega\}$ is inconsistent. The conditions of Lemma 2.14 are met and

$$d(\chi(\mathbb{M}^{|\bar{x}|})) > d(\phi(\mathbb{M}^{|\bar{x}|}, \bar{d}_i) \wedge \chi(\mathbb{M}^{|\bar{x}|})) = d(\phi(\mathbb{M}^{|\bar{x}|}, \bar{d}) \wedge \chi(\mathbb{M}^{|\bar{x}|})) = d(\phi(\mathbb{M}^{|\bar{x}|}, \bar{d})).$$

Where the second equality holds by invariance of the dimension and the last equality holds since $\phi(\mathbb{M}^{|\bar{x}|}, \bar{d}) \subseteq \chi(\mathbb{M}^{|\bar{x}|})$. Hence, by our choice of $\chi(\bar{x})$, we have that $d(\bar{a}/BC) < d(\bar{a}/C)$, implying that \bar{a} is not d -independent from B over C . Our claim then holds by contrapositive. \square

The other implication holds by the following result of Kim and Pillay [14][Theorem 4.2, Claim I]:

Lemma 2.16. *Let T be an arbitrary theory and \perp^* be a notion of independence. Suppose that $a \perp_C^f B$. Then, $a \perp_C^* B$.*

And this yields the desired result:

Theorem 2.17. *Let \mathcal{M} be an MS-measurable structure. Then, dimension independence is the same as non-forking independence.*

Remark 2.18. An alternative way to prove that dimension independence is the same of non-forking independence is by noting how the independence theorem is substantially proven in Theorem 2.18 of [8].

2.3 MS-measurability in an ω -categorical context

In this subsection we introduce some basic equations which must be satisfied in ω -categorical MS-measurable structures through Corollary 2.20. Observations regarding these equations allow us to prove that in an ω -categorical context we may always take the dimension part of an MS-dimension-measure to be SU-rank, as shown in corollary 2.22.

The following result is a standard fact about ω -categorical MS-measurable structures. As far as I can tell, a version of this is first proven in Elwes' PhD thesis [1][Lemma 5.2.1]:

Lemma 2.19. *Let \mathcal{M} be ω -categorical and MS-measurable. Let $B \supseteq A$ be finite subsets of M . Suppose p is a partial type over A . By ω -categoricity, p has finitely many complete extensions to B . Let p_1, \dots, p_n be those which do not fork over A . Then,*

$$\mu(p) = \sum_{i=1}^n \mu(p_i). \quad (1)$$

Proof. Suppose the complete extensions of p to B are p_1, \dots, p_m with $m \geq n$ where p_1, \dots, p_n have maximal dimension. By Theorem 2.17, these are the non-forking extensions of p to B . Consider the sets

$$p_i(M^{|\bar{x}|}) := \left\{ \bar{c} \in M^{|\bar{x}|} \mid \bar{c} \models p(\bar{x}) \right\}.$$

By ω -categoricity, these are disjoint definable sets such that $\bigcup_{i=1}^m p_i(M^{|\bar{x}|}) = p(M^{|\bar{x}|})$. Hence, by additivity:

$$\mu(p) = \mu \left(\bigcup_{i=1}^n p_i(M^{|\bar{x}|}) \cup \bigcup_{i>n}^m p_i(M^{|\bar{x}|}) \right) = \mu \left(\bigcup_{i=1}^n p_i(M^{|\bar{x}|}) \right) = \sum_{i=1}^n \mu(p_i).$$

Where the second last equality holds by additivity in the case $d(X) < d(Y)$, and the last equality due to additivity in the case of $d(X) = d(Y)$. \square

By an easy application of Fubini (see Remark 2.11) we get equations in terms of types over the empty set and obtain:

Corollary 2.20. *Let \mathcal{M} be ω -categorical and MS-measurable. Take $\bar{a}, \bar{b}, \bar{c}$ to be tuples from M . Let $\text{tp}(\bar{c}_1/\bar{a}\bar{b}), \dots, \text{tp}(\bar{c}_n/\bar{a}\bar{b})$ be the finitely many complete non-forking extensions of $\text{tp}(\bar{c}/\bar{a})$ to $\bar{a}\bar{b}$. Then,*

$$\frac{\mu(\bar{a}\bar{c})\mu(\bar{a}\bar{b})}{\mu(\bar{a})} = \sum_{i=1}^n \mu(\bar{a}\bar{b}\bar{c}_i). \quad (2)$$

We shall use the above corollary in our proof of non-measurability. There is a partial converse to this corollary. In fact if an ω -categorical structure has a function $\mu : S(\emptyset) \rightarrow \mathbb{R}^{>0}$ satisfying equations of the form of (2) and an algebraicity condition, then it is MS-measurable. This can be extracted from the proof of Theorem 2.21. In this theorem we show that in an ω -categorical MS-measurable structure we may choose our dimension to be any definable dimension yielding non-forking independence as its notion of independence.

Theorem 2.21. *Suppose that \mathcal{M} is an ω -categorical MS-measurable structure with dimension-measure $h = (d, \mu)$. Let $D : \text{Def}(\mathcal{M}) \rightarrow \mathbb{N}$ be any definable dimension for \mathcal{M} whose induced notion of independence is non-forking independence. Then, \mathcal{M} has a dimension-measure $h' = (D, \mu')$, where μ' agrees with μ on complete types over finite sets of parameters.*

Proof. By Ryll-Nardzewski, D is invariant and finite. The dimension-measure $h = (d, \mu)$ induces a function on types over finite sets of parameters $\mu^* : S_{\text{fin}}(\mathcal{M}) \rightarrow \mathbb{R}^{>0}$ given by $\mu^*(\bar{a}/\bar{b}) = \mu(\bar{a}/\bar{b})$. We can write any set X definable over \bar{a} as a finite disjoint union of the sets of realisations of complete types p_1, \dots, p_m over \bar{a} . Say that p_1, \dots, p_m have maximal D -dimension. Then we define,

$$\mu'(X) = \sum_{i=1}^m \mu^*(p_i).$$

We need to show that our definition doesn't depend on the choice of parameters over which X is defined. Suppose that X is defined both over \bar{a} and \bar{b} . We may assume without loss of generality that $\bar{a} \subseteq \bar{b}$. Each p_i over \bar{a} has finitely many complete extensions of maximal D -dimension to \bar{b} , say $p'_{i,1}, \dots, p'_{i,m_i}$, which by assumption are the non-forking extensions of p_i to \bar{b} . By the dimension theorem 2.17, these are the extensions of p_i of maximal d -dimension, and so, by MS-measurability we know that

$$\mu(p_i) = \sum_{j=1}^{m_i} \mu(p'_{ij}).$$

But then,

$$\sum_{i=1}^m \mu(p_i) = \sum_{i=1}^m \sum_{j=1}^{m_i} \mu(p'_{ij}),$$

where the equation on the left is the definition of $\mu'(X)$ as defined over \bar{a} and the equation on the right is the definition of μ' for X as defined over \bar{b} . Hence, μ' is well defined. Positivity, algebraicity, additivity and Fubini are easy to prove. Hence, (D, μ') yields an MS-dimension-measure for \mathcal{M} . \square

An important consequence of this theorem is that for MS-measurable ω -categorical structures we may take the dimension to be SU -rank. In a structure of finite SU -rank, SU -rank is a dimension function: it is additive by the Lascar inequalities [13][Prop. 2.5.19]. In an ω -categorical structure, it is definable being invariant. Hence, since MS-measurable structures have finite SU -rank, in ω -categorical MS-measurable structures, SU -rank yields a definable dimension. By definition it induces non-forking independence as its notion of independence. Hence, it satisfies all of the conditions of Theorem 2.21, and we get the following corollary.

Corollary 2.22. *Suppose that the ω -categorical structure \mathcal{M} is MS-measurable with dimension-measure $h = (d, \mu)$. Then, \mathcal{M} is also MS measurable via the dimension-measure $h' = (SU, \mu')$, where μ' is as in Theorem 2.21. Hence, if an ω -categorical structure \mathcal{M} is MS-measurable, there is a dimension-measure on \mathcal{M} where the dimension is given by SU -rank.*

This statement is known to be false outside of an ω -categorical context. In particular, in [2][Remark 3.8] Elwes and Macpherson show that there are MS-measurable structures where SU -rank is not definable. They also give an example of an ω -categorical structure for which we can artificially choose the dimension in the MS-measure and SU -rank to differ. However, our theorem does prove that if there is an MS-measure in the ω -categorical case, there is no harm in taking the dimension to be SU -rank.

Corollary 2.22 is a powerful tool in the context of MS-measurable ω -categorical structures. For example, Evans' proof [5] employs a highly non-trivial theorem showing that in the class of ω -categorical structures he is considering, any dimension function corresponds to a scaled SU -rank. Our result allows us to skip proving such a theorem. Indeed, for other ω -categorical structures such a theorem is false (e.g. the example in [2][Rem. 3.8]).

3 Independence in measure

In this section we obtain another set of equations that hold for MS -measurable ω -categorical structures. The main idea is that in an MS -measurable structure, non-forking independence induces probabilistic independence in the measure. This is shown explicitly in Theorem 3.5, and Corollary 3.6, which yields equations that we will use later in our proof of non- MS -measurability. We thank Ehud Hrushovski for sharing his notes for [10]. In there, he proves a more general version of Theorem 3.5, which implies the results in this section. Working in an ω -categorical MS -measurable context greatly simplifies the tools and the amount of theory required to obtain the probabilistic independence theorem. Hence, we give proofs for these results in this section.

For a structure \mathcal{M} , let $\mu : \text{Def}_{\bar{y}}(M) \rightarrow \mathbb{R}^{\geq 0}$ be an invariant measure and $\phi(\bar{x}_1, \bar{y}), \psi(\bar{x}_2, \bar{y})$ be \mathcal{L} -formulas. We define the relation $R_\alpha(\bar{x}_1, \bar{x}_2)$ on $M^{|\bar{x}_1|} \times M^{|\bar{x}_2|}$ by

$$R_\alpha(\bar{a}, \bar{b}) \text{ if and only if } \mu(\phi(\bar{a}, \bar{y}) \wedge \psi(\bar{b}, \bar{y})) = \alpha.$$

From [8][Prop.2.25] we know that $R_\alpha(\bar{x}_1, \bar{x}_2)$ is stable.

Definition 3.1. Let $\Delta = \{\phi_i(\bar{x}, \bar{z}_i) \mid i \leq n\}$ be a finite set of formulas and A a set of parameters. By $S_\Delta^x(A)$ we mean the set of Δ -types over A . By $FER_\Delta(A)$ we denote the set of A -definable equivalence relations $E(\bar{x}, \bar{y})$ with finitely many classes such that for any \bar{b} , $E(\bar{x}, \bar{b})$ is elementary equivalent to a Boolean combination of formulas from Δ with the \bar{z}_i taking values in A .

In an ω -categorical structure with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$, for $\bar{a}_1 \perp \bar{a}_2$, the relation $R_\alpha(\bar{a}_1, \bar{a}_2)$ only depends on the individual types of \bar{a}_1 and \bar{a}_2 . A common way of using the finite equivalence relation theorem (see [11] or [14] for similar uses) yields this result.

Lemma 3.2. Let $\mu : \text{Def}_{\bar{y}}(M) \rightarrow \mathbb{R}$ be an invariant measure on an ω -categorical \mathcal{L} -structure \mathcal{M} with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$. Let $\phi_i(\bar{x}_i, \bar{y})$ for $i \in \{1, 2\}$ be \mathcal{L} -formulas. Suppose that \bar{a}_1, \bar{a}_2 are tuples from M such that $\bar{a}_1 \perp \bar{a}_2$. Then, the value of $\mu(\phi_1(\bar{a}_1, \bar{y}) \wedge \phi_2(\bar{a}_2, \bar{y}))$ only depends on $\text{tp}(\bar{a}_1)$ and $\text{tp}(\bar{a}_2)$, but not on $\text{tp}(\bar{a}_1 \bar{a}_2)$.

Proof. Let $\alpha_1, \dots, \alpha_m$ be the finitely many possible values for $\mu(\phi_1(\bar{b}_1, \bar{y}) \wedge \phi_2(\bar{b}_2, \bar{y}))$ with $\bar{b}_1 \perp \bar{b}_2$. Let us expand our language by relation symbols $R_i(\bar{x}_1, \bar{x}_2)$, and consider the expansion of \mathcal{M} obtained by interpreting the R_i as

$$\models R_i(\bar{b}_1, \bar{b}_2) \text{ if and only if } \mu(\phi_1(\bar{b}_1, \bar{y}) \wedge \phi_2(\bar{b}_2, \bar{y})) = \alpha_i.$$

Each of these relations is invariant and, as recalled above, stable. Let Δ be the collection of these relation symbols. Let $\bar{a}_1 \equiv \bar{a}'_1, \bar{a}_1 \perp \bar{a}_2$ and $\bar{a}'_1 \perp \bar{a}_2$. Hence, \bar{a}_1 and \bar{a}'_1 have the same Δ -types over \emptyset .

Let $B \subset M$ and suppose that $q_1, q_2 \in S_\Delta \bar{x}(B)$ are non-forking extensions of $\text{tp}_\Delta(\bar{a}_1)$. By stability of the R_i and the finite equivalence relation theorem, $q_1 \neq q_2$ if and only if there is a finite equivalence relation $E(\bar{x}, \bar{y}) \in FER_\Delta(\emptyset)$ such that $q_1(\bar{x}) \cup q_2(\bar{y}) \vdash E(\bar{x}, \bar{y})$. By $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$, $E(\bar{x}, \bar{y})$ is trivial, i.e. tuples with the same types over the emptyset will be in the same equivalence classes. In particular, since for \bar{a} , $E(\bar{x}, \bar{a})$ is a Boolean combination of formulas in Δ , the equivalence class of \bar{a} is entirely determined by $\text{tp}_\Delta(\bar{a})$, and so $q_1 = q_2$. Hence, for any B there is a unique Δ -type over B which is a non-forking extension of $\text{tp}_\Delta(\bar{a}_1)$. In particular, $\text{tp}_\Delta(\bar{a}_1/\bar{a}_2) = \text{tp}_\Delta(\bar{a}'_1/\bar{a}_2)$, which yields

$$\models R_i(\bar{a}_1, \bar{a}_2) \leftrightarrow R_i(\bar{a}'_1, \bar{a}_2).$$

We can apply the same reasoning in the variable \bar{x}_2 to obtain that for \bar{a}'_2 such that $\bar{a}'_1 \perp \bar{a}'_2$,

$$\models R_i(\bar{a}_1, \bar{a}_2) \leftrightarrow R_i(\bar{a}'_1, \bar{a}'_2).$$

Hence, whenever \bar{a}'_1, \bar{a}'_2 are tuples such that $\bar{a}'_i \equiv \bar{a}_i$ for $i = 1, 2$ and $\bar{a}'_1 \perp \bar{a}'_2$ we have the equality $\mu(\phi_1(\bar{a}'_1, \bar{y}) \wedge \phi_2(\bar{a}'_2, \bar{y})) = \mu(\phi_1(\bar{a}_1, \bar{y}) \wedge \phi_2(\bar{a}_2, \bar{y}))$. \square

Let \mathcal{M} be an \mathcal{L} structure. Let \bar{a} be a tuple from M and $B \subseteq M$. We write $\text{loc}(\bar{a}/B)$ for the set of realisations of $\text{tp}(\bar{a}/B)$ in \mathcal{M} .

Lemma 3.3. *Let \mathcal{M}^{eq} be ω -categorical and a_1, a_2 be tuples from M^{eq} . Then, there is an \mathcal{L}^{eq} formula $\psi(x_1, x_2)$ such that*

$$\models \psi(a'_1, a'_2) \text{ if and only if } a'_1 \equiv a_1 \text{ and } a'_2 \equiv a_2 \text{ and } a'_1 \perp a'_2. \quad (3)$$

Moreover, if \mathcal{M} is MS-measurable,

$$\mu(\psi(M, M)) = \mu(a_1)\mu(a_2).$$

Proof. The first part follows simply by Ryll-Nardzewski and invariance of non-forking independence. For the second part we use Fubini. Firstly, since $\psi(x_1, x_2)$ isolates the types of its variables,

$$\models \exists x_1 \psi(x_1, b_2) \text{ if and only if } b_2 \in \text{loc}(a_2).$$

So, $\mu(\exists x_1 \psi(x_1, M)) = \mu(a_2)$. Secondly, we consider $\mu(\psi(M^{x_1}, b_2))$ for b_2 such that $\models \exists x_1 \psi(x_1, b_2)$. By invariance, $\mu(\psi(M, b_2)) = \mu(\psi(M, a_2))$. But now, for $b_1 \equiv a_1$, by the dimension theorem 2.17:

$$\models \psi(b_1, a_2) \text{ if and only if } d(b_1/a_2) = d(b_1).$$

Hence, by additivity, $\mu(a_1) = \mu(\psi(M, a_2))$. Considering Fubini applied to $\psi(x_1, x_2)$ yields the result. \square

Fact 3.4. *If \mathcal{M} is supersimple, ω -categorical with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$, the independence theorem holds over the empty set.*

Proof. By ω -categoricity, over finite sets, Lascar types and strong types coincide. But then by [13][Proposition 3.2.12], the independence theorem holds over $\text{acl}^{eq}(\emptyset)$. By $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$, the independence theorem holds over the empty set. \square

At this point we have the tools to show how in an ω -categorical context, for a triplet of independent pairs, non-forking independence corresponds to probabilistic independence. We state the following result in terms of \mathcal{M}^{eq} recalling that if \mathcal{M} is MS-measurable, then so is \mathcal{M}^{eq} [15].

Theorem 3.5. *Let \mathcal{M}^{eq} be MS-measurable, ω -categorical with $\text{acl}^{eq}(\emptyset) = \text{dcl}^{eq}(\emptyset)$. Suppose that a_1, a_2, a_3 are tuples such that $a_i \perp a_j$ for $1 \leq i < j \leq 3$. Then, $d(\text{tp}(a_3/a_1) \cup \text{tp}(a_3/a_2)) = d(a_3)$ and*

$$\mu(\text{tp}(a_3/a_1) \cup \text{tp}(a_3/a_2)) = \frac{\mu(a_3/a_1)\mu(a_3/a_2)}{\mu(a_3)}. \quad (4)$$

In the theorem, the set defined by $\text{tp}(a_3/a_1) \cup \text{tp}(a_3/a_2)$ is $\text{loc}(a_3/a_1) \cap \text{loc}(a_3/a_2)$. The statement about the dimension corresponds to the independence theorem over \emptyset . But we get a much stronger result: the events " $x \in \text{loc}(a_3/a_1)$ " and " $x \in \text{loc}(a_3/a_2)$ " are probabilistically independent in the measure μ' induced by μ on definable subsets of $\text{loc}(a_3)$.

Proof. We write M instead of M^{eq} in the proof to simplify the notation. For $i \in \{1, 2\}$, let $\phi_{i3}(x_i, x_3)$ isolate $\text{tp}(a_i/a_3)$. Let $\psi(x_1, x_2)$ be the formula satisfying the condition 3 shown to exist in Lemma 3.3. Our proof consists of calculating in different ways the measure of the set S defined by the formula

$$S(x_1, x_2, x_3) := \phi_{13}(x_1, x_3) \wedge \phi_{23}(x_2, x_3) \wedge \psi(x_1, x_2).$$

We apply Fubini, projecting onto the first two coordinates $\pi_{x_1 x_2} : M^3 \rightarrow M^2$. The formula $\exists x_3 S(x_1, x_2, x_3)$ is satisfied by any $b_1 b_2 \models \psi(x_1, x_2)$ by Fact 3.4. Hence,

$$\exists x_3 S(x_1, x_2, x_3) \equiv \psi(x_1, x_2),$$

and so by Lemma 3.3,

$$\mu(\pi_{x_1 x_2}(S)) = \mu(a_1)\mu(a_2).$$

For $b_1 b_2 \in M$ we consider $\pi_{x_1 x_2}^{-1}(b_1 b_2) \cap S$, i.e. the set defined by $S(b_1, b_2, x_3)$. This definable set is non-empty if and only if $b_1 b_2 \models \psi(x_1, x_2)$. Also, by Fact 3.4, for any such pair, $S(b_1, b_2, x_3)$ will have maximal dimension. Finally, by Lemma 3.2,

$$\mu(S(b_1, b_2, M)) = \mu(S(a_1, a_2, M)) = \mu(\text{tp}(a_3/a_1) \cup \text{tp}(a_3/a_2)).$$

Hence, by Fubini with respect to $\pi_{x_1 x_2}$ we obtain:

$$\mu(S) = \mu(\text{tp}(a_3/a_1) \cup \text{tp}(a_3/a_2))\mu(a_1)\mu(a_2). \quad (5)$$

Let us compute the measure with respect to $\pi_{x_3} : M^3 \rightarrow M$. The formula $\exists x_1 x_2 S(x_1, x_2, x_3)$ isolates $\text{tp}(a_3)$ and so $\mu(\exists x_1 x_2 S(x_1, x_2, M)) = \mu(a_3)$. Meanwhile, for $b_3 \in M$, the formula $S(x_1, x_2, b_3)$ is consistent only if $b_3 \equiv a_3$. By invariance we may consider $S(x_1, x_2, a_3)$. To compute the measure of the set defined by this formula we apply Fubini again. Projecting onto x_1 , $\exists x_1 S(x_1, x_2, a_3)$ isolates $\text{tp}(a_2/a_3)$ and so, $\mu(\exists x_1 S(x_1, M, a_3)) = \mu(a_2/a_3)$. Meanwhile, for $b_2 \equiv_{a_3} a_2$ consider $S(x_1, b_2, a_3)$. Again, by invariance this defines a set with the same measure as $S(x_1, a_2, a_3)$. We claim that $\mu(S(M, a_2, a_3)) = \mu(a_1/a_3)$. By Lemma 3.3, there is a formula $\chi(x_1, x_2)$ over a_3 isolating the pairs $b_1 \equiv_{a_3} a_1$ and $b_2 \equiv_{a_3} a_2$ which are independent over a_3 . So, we may consider $S(M, a_2, a_3)$ as the disjoint union

$$(S(M, a_2, a_3) \wedge \chi(M, a_2)) \sqcup (S(M, a_2, a_3) \wedge \neg\chi(M, a_2)).$$

But the definable set on the right of the disjoint union has lower dimension by the dimension theorem, and so, by additivity,

$$\mu(S(M, a_2, a_3)) = \mu(S(M, a_2, a_3) \wedge \chi(M, a_2)).$$

Consider the extension of $\text{tp}(a_1/a_3)$ to a_2 . By the same reasoning,

$$\mu(a_1/a_3) = \mu(\phi_{13}(M, a_3) \wedge \chi(M, a_2)).$$

Noting that $\chi(x_1, x_2)$ implies $\psi(x_1, x_2)$, we get that

$$\mu(S(M, a_2, a_3)) = \mu(a_1/a_3).$$

Hence, our calculations with respect to π_{x_3} yield:

$$\mu(S) = \mu(a_3)\mu(a_2/a_3)\mu(a_1/a_3). \quad (6)$$

Recall that by Fubini $\mu(a_i a_j) = \mu(a_i/a_j)\mu(a_j)$. Using this fact and comparing the equations 5 and 6, we obtain

$$\mu(\text{tp}(a_3/a_1) \cup \text{tp}(a_3/a_2)) = \frac{\mu(a_2 a_3)\mu(a_1 a_3)}{\mu(a_1)\mu(a_2)\mu(a_3)}. \quad (7)$$

Applying Fubini again, we get the desired result. \square

Corollary 3.6. *Let \mathcal{M}^{eq} be MS-measurable, ω -categorical with $\text{acl}^{\text{eq}}(\emptyset) = \text{dcl}^{\text{eq}}(\emptyset)$. Suppose that a_1, a_2, a_3 are tuples such that $a_i \perp a_j$ for $1 \leq i < j \leq 3$. Let $p_{ij}(x_i, x_j) = \text{tp}(a_i, a_j)$. Then,*

$$d(p_{12}(x_1, x_2) \cup p_{23}(x_2, x_3) \cup p_{13}(x_1, x_3)) = \sum_{i \leq 3} d(a_i), \text{ and}$$

$$\mu(p_{12}(x_1, x_2) \cup p_{23}(x_2, x_3) \cup p_{13}(x_1, x_3)) = \frac{\mu(a_1 a_2)\mu(a_1 a_3)\mu(a_2 a_3)}{\mu(a_1)\mu(a_2)\mu(a_3)}.$$

Proof. Considering the set defined by $p_{12}(x_1, x_2) \cup p_{23}(x_2, x_3) \cup p_{13}(x_1, x_3)$, we apply Fubini by projecting onto the first two coordinates. The projection isolates $\text{tp}(a_1 a_2)$, and the fiber of $a_1 a_2$ isolates $\text{tp}(a_3/a_1) \cup \text{tp}(a_3/a_2)$. Hence, by Fubini

$$\mu(p_{12}(x_1, x_2) \cup p_{23}(x_2, x_3) \cup p_{13}(x_1, x_3)) = \mu(\text{tp}(a_3/a_1) \cup \text{tp}(a_3/a_2))\mu(a_1 a_2).$$

Substituting with equation 7, the result follows. \square

4 Hrushovski constructions

In this section we focus on ω -categorical Hrushovski constructions. We begin with subsection 4.1 which gives a brief introduction to these structures following [6] and [23]. Then, in subsection 4.2, we find the form that the equations of Corollary 2.20 would take in the context of ω -categorical Hrushovski constructions if these were MS -measurable.

4.1 Building ω -categorical Hrushovski constructions

We focus on graphs for simplicity of notation, but we can build Hrushovski constructions in relational languages in general [23]. Let $\overline{\mathcal{K}}$ be the class of graphs and \mathcal{K} be the class of finite graphs. Now, for $\alpha \in \mathbb{R}^{>0}$, and A a finite graph, we define the **predimension** of A , $\delta(A)$ to be:

$$\delta(A) = \alpha|A| - |E(A)|,$$

where $|E(A)|$ is the number of edges of A .

For $A \subseteq B, C \in \mathcal{K}$, suppose $g : A \rightarrow B$ and $f : A \rightarrow C$ are embeddings of A in B and C respectively. The **free amalgamation** of B and C over A , is the (unique) graph $B \amalg_A C$ such that there are embeddings $h : B \rightarrow B \amalg_A C$, and $k : C \rightarrow B \amalg_A C$ such that $h(g(A)) = k(f(A))$ and $E(B \amalg_A C) = E(B) \cup E(C)$.

The predimension we introduced satisfies **submodularity**. That is, for $D \in \overline{\mathcal{K}}$, and $B, C \subset D$ finite, we have that

$$\delta(B \cup C) \leq \delta(B) + \delta(C) - \delta(B \cap C),$$

with equality holding if and only B and C are freely amalgamated over $B \cap C$.

Definition 4.1. Let $B \in \overline{\mathcal{K}}$, $A \subseteq B$ finite. We say that A is **d-closed** in B , and write $A \leq B$ if $\delta(A) < \delta(B')$ for any finite B' such that $A \subsetneq B' \subseteq B$. We say that A is **self sufficient** in B and write $A \leq^* B$ if $\delta(A) \leq \delta(B')$ for any finite B' such that $A \subseteq B' \subseteq B$.

Let $f : \mathbb{R}^{\geq 0} \rightarrow \mathbb{R}^{\geq 0}$ be continuous, increasing and unbounded with $f(0) = 0$. We let

$$\mathcal{K}_f := \{A \in \mathcal{K} : \delta(A') \geq f(|A'|) \text{ for any } A' \subseteq A\}$$

Definition 4.2. We say that \mathcal{K}_f has the **free amalgamation property** if given $A_0 \leq A_1, A_2 \in \mathcal{K}_f$ we have that $A_1 \amalg_{A_0} A_2 \in \mathcal{K}_f$.

When \mathcal{K}_f has the free amalgamation property we can build an ω -categorical Hrushovski construction from it [6][Theorems 3.2, & 3.19]:

Theorem 4.3. *Suppose that \mathcal{K}_f has the free amalgamation property. Then, there is a unique countable structure \mathcal{M}_f such that:*

- **Union of Chains:** \mathcal{M}_f is given by the union of a chain $M_1 \leq M_2 \leq \dots$ where $M_i \in \mathcal{K}_f$.
- **Substructures in \mathcal{K}_f :** Every $A \in \mathcal{K}_f$ is isomorphic to some $A' \leq M_f$.
- **Extension Property:** Given $A \leq M_f$ finite and $A \leq B \in \mathcal{K}_f$, there is an embedding of B over A in \mathcal{M}_f such that $B \leq M_f$.

Furthermore, \mathcal{M}_f is ω -categorical.

We say that f is a **good function** if it is piece-wise smooth, $f'(x) \leq 1/x$ and $f'(x)$ is non-decreasing.

Lemma 4.4. *Suppose that $f(x)$ is a good function for $x \geq t$. and that for $A_0 \leq A_1, A_2$, with $|A_i| \leq t$, we have $\delta(A_0 \amalg_{A_1} A_2) \geq f(|A_0 \amalg_{A_1} A_2|)$. Then \mathcal{K}_f has the free amalgamation property.*

Proof. This is substantially the proof of Example 3.20 in [6], which goes back to Hrushovski's original unpublished note [9]. \square

Furthermore, when $\alpha \in \mathbb{Q}$, choosing f with slow enough growth ensures that \mathcal{M}_f is supersimple of finite SU -rank [12][3][Theorem 3.6]. This is explained further in the Appendix.

For $B \in \mathcal{K}_f$ or $B = \mathcal{M}_f$ and finite $A \subseteq B$ there is a unique smallest d -closed subset of B containing A which we call $\text{cl}_B(A)$, i.e, the **closure** of A in B [23][Lemma 2.1].

In the context of \mathcal{M}_f , $\text{cl}_{\mathcal{M}_f}(A)$ corresponds to algebraic closure of A in \mathcal{M}_f and is bounded above by $f^{-1}(\alpha|A|)$ [6][Theorem 3.19]. In general, we shall avoid the subscript when it is clear we are speaking of $\text{cl}_{\mathcal{M}_f}(A)$.

The predimension of a closed set induces a natural notion of **dimension**. For D finite or $D = \mathcal{M}_f$, given $A \subseteq D$, we write $d_D(A)$ for $\delta(\text{cl}_D(A))$. If $\alpha \in \mathbb{N}$, then this dimension is also a natural number, and it may be re-scaled to be a natural number if $\alpha \in \mathbb{Q}$. As above, we omit the subscript and write $d(A)$ for $d_{\mathcal{M}_f}(A)$.

We write $d_D(A/B)$ for $d_D(AB) - d_B(B)$, where we write AB for $A \cup B$. We say that A and B are **independent** over C in D if $d_D(A/BC) = d_D(A/B)$. This implies that $\text{cl}_D(AB) \cap \text{cl}_D(AC) = \text{cl}_D(A)$. When \mathcal{M}_f is supersimple (of finite SU rank) we have that this notion of independence induced by the Hrushovski dimension in \mathcal{M}_f is precisely non-forking independence and the Hrushovski dimension on \mathcal{M}_f is SU -rank.

We have a clear picture of what independence looks like in \mathcal{M}_f [13][Claim 6.2.9]:

Lemma 4.5. *Let $D \in \mathcal{K}_f$ or $D = \mathcal{M}_f$. Suppose $A = B \cap C \leq B, C \leq D$. Then, B and C are independent over A in D if and only if B and C are freely amalgamated over A and $\delta(B \cup C) = \delta(\text{cl}_D(B \cup C))$.*

Remark 4.6. With the notation above, if B and C are independent over A in D , $BC \leq^* \text{cl}_D(B \cup C)$. To see this, suppose by contradiction that there is some F such that $BC \subseteq F \subsetneq \text{cl}_D(B \cup C)$ and $\delta(F) < \delta(BC)$. We may choose F to be maximal so that for $F \subsetneq H \subseteq \text{cl}_D(B \cup C)$, $\delta(H) > \delta(F)$. Then,

$$BC \subseteq F \leq \text{cl}_D(B \cup C) \leq D.$$

But this contradicts the definition of closure in D as the minimal d -closed subset of D containing BC .

4.2 MS-measurable ω -categorical Hrushovski constructions

In this section we consider some equations which would hold in an MS -measurable ω -categorical Hrushovski construction. Suppose $A \subseteq M_f$, and $B = \text{cl}_{\mathcal{M}_f}(A)$. Let $\bar{a}\bar{b}$ be an enumeration of B where \bar{a} is an enumeration of A . By the extension property in Theorem 4.3, the quantifier-free type of $\bar{a}\bar{b}$ determines the type of \bar{a} in \mathcal{M}_f [23][Cor. 2.4]. For $A \leq M_f$, we write $\theta_{\bar{a}}(\bar{x})$ for the formula isolating $\text{tp}(\bar{a})$.

Notation 4.7. Note that by Fubini, any MS -dimension-measure h is invariant under permutation of variables. That is, for \bar{x}_σ a permutation of the variables of \bar{x} , and $\phi_\sigma(\bar{x}) = \phi(\bar{x}_\sigma)$,

$$h(\phi_\sigma(M^{|\bar{x}|})) = h(\phi(M^{|\bar{x}|})).$$

Hence, if \mathcal{M}_f is MS -measurable, for $A \leq M_f$, we may write $\mu(A) := \mu(\theta_{\bar{a}}(\bar{x}))$ without ambiguity. For $A \in \mathcal{K}_f$ we write $\mu(A)$ for $\mu(\theta_{\bar{a}}(\bar{x}))$, where \bar{a} is an enumeration of a copy A' of A in M_f such that $A' \leq M_f$.

By the notation $\text{tp}(A/B)$ we mean $\text{tp}(\bar{a}/B)$ where \bar{a} is an enumeration of A .

Theorem 4.8. *Let \mathcal{M}_f be the ω -categorical Hrushovski construction for the class \mathcal{K}_f . Suppose that \mathcal{M}_f is MS -measurable. Let $A \leq B, C \leq M_f$ be finite. Let $\text{tp}(C_1/B), \dots, \text{tp}(C_n/B)$ be the finitely many non-forking extensions of $\text{tp}(C/A)$ to B . Let D_i be $\text{cl}(BC_i)$. Then, we have that:*

$$\frac{\mu(B)\mu(C)}{\mu(A)} = \sum_{i=1}^n \frac{\mu(D_i)}{|\text{Aut}(D_i/BC_i)|} \quad (8)$$

where $|\text{Aut}(D_i/BC_i)|$ is the number of automorphisms of D_i fixing BC_i pointwise.

Proof. Let $\bar{a}\bar{b}$ be an enumeration of B where \bar{a} is an enumeration of A . Similarly, let $\bar{a}\bar{c}$ be an enumeration of C . Let $\bar{a}\bar{b}\bar{c};\bar{d}_i$ be an enumeration of D_i where, $\bar{a}\bar{b}\bar{c}_i$ is the corresponding enumeration of ABC_i . Note that, by Corollary 2.20,

$$\frac{\mu(B)\mu(C)}{\mu(A)} = \sum_{i=1}^n \mu(\text{tp}(\bar{a}\bar{b}\bar{c}_i)) \quad (9)$$

We know that $\text{tp}(\bar{a}\bar{b}\bar{c}_i)$ is isolated by $\exists \bar{w}\theta_{\bar{a}\bar{b}\bar{c}_i;\bar{d}_i}(\overline{xyz\bar{w}})$. Note that $|\{e \in M_f | \theta_{\bar{a}\bar{b}\bar{c}_i;\bar{d}_i}(\bar{a}\bar{b}\bar{c}_i e)\}|$ is finite and counts the number of automorphisms of D_i fixing BC_i pointwise. Hence, by Fubini,

$$\mu(\text{tp}(\bar{a}\bar{b}\bar{c}_i))|\text{Aut}(D_i/BC_i)| = \mu(D_i).$$

This yields the desired equation. \square

Following Lemma 4.5, we can express the conditions for a graph D_i to appear as $\text{cl}(BC_i)$ in the theorem above in terms of \mathcal{K}_f .

Definition 4.9. Let $A_0 \leq A_1, A_2, A_1 \amalg_{A_0} A_2 \subseteq D \in \mathcal{K}_f$. We say D is an **eventual closure** of $A_1 \amalg_{A_0} A_2$ if A_1 and A_2 are freely amalgamated over A_0 in D , $A_1 \cup A_2 \leq^* D$, $\delta(A_1 \cup A_2) = \delta(D)$, and $A_i \leq D$ for each $i \in \{0, 1, 2\}$.

Remark 4.10. Consider the closures D_i in Theorem 4.8. We have that there is $C_i \leq M_f$ independent from B over A with $\text{tp}(C_i/A) = \text{tp}(C/A)$ and closure D_i if and only if there is $D'_i \in \mathcal{K}_f$, isomorphic to D_i as a graph, which is an eventual closure of $B \amalg_A C$. The left-to-right implication follows by Lemma 4.5 and Remark 4.6. The right-to-left implication follows by the extension property, noting that $B \leq D'_i \in \mathcal{K}_f$.

Hence, we may rephrase Theorem 4.8 as follows:

Corollary 4.11. Let \mathcal{M}_f be an ω -categorical Hrushovski construction with amalgamation class \mathcal{K}_f . Suppose \mathcal{M}_f is MS -measurable. Let $A \leq B, C \in \mathcal{K}_f$. Let D_1, \dots, D_n be the eventual closures of $B \amalg_A C$. Then,

$$\frac{\mu(B)\mu(C)}{\mu(A)} = \sum_{i=1}^n \frac{\mu(D_i)}{|\text{Aut}(D_i/BC)|}.$$

5 A non-measurability proof

In this section we give a class of ω -categorical Hrushovski constructions. These are super-simple of finite SU -rank, as proven in the Appendix 6. However, we will prove in Subsection 5.2 that these are not MS -measurable. In subsection 5.1 we introduce these structures and show how we have control on small enough eventual closures in them.

5.1 The structure \mathcal{M}_f and small eventual closures

We begin by giving the class of structures that we work with.

Construction 5.1. We set $\alpha = 2$. Let $f : \mathbb{R}^{>0} \rightarrow \mathbb{R}^{>0}$ be such that for $t \geq 6$, f is a good function with $f(3 \cdot t) \leq f(t) + 1$ and for $t \leq 6$, f is piece-wise linear with $f(1) = 2, f(4) = 5, f(6) = 6$. An illustration of the initial values of f is given by Figure 1. As proven in the Appendix 6, the class \mathcal{K}_f has the free amalgamation property, and so by Theorem 4.3 we can build an ω -categorical Hrushovski construction \mathcal{M}_f .

An example of a function satisfying the conditions of Construction 5.1 is the function taking the values specified for $t \leq 6$ and corresponding to $\log_3(x) - \log_3(6) + 6$ for $t > 6$. Of course, there are uncountably many functions satisfying the conditions of our construction.

From now on, by \mathcal{M}_f we mean an ω -categorical Hrushovski construction obtained following Construction 5.1.




Figure 1: The initial values of f are represented by the green line. The red dots represent positions $(|X|, \delta(X))$ of the graphs X in \mathcal{K}_f (some red dots for $|X| \geq 5$ lie outside of the figure). Note that $f(1) = 2$ and $f(2) = 3$ are necessary to include vertices and edges in \mathcal{K}_f . $f(3) = 4, f(4) = 5, f(6) = 6$ imply that the smallest cycle in \mathcal{K}_f is a 6-cycle. We also have that vertices and edges are always closed in \mathcal{M}_f .

Lemma 5.2 (Basic properties of \mathcal{M}_f). *The ω -categorical Hrushovski construction \mathcal{M}_f is super-simple of SU-rank 2. Furthermore, it has weak elimination of imaginaries. Finally, we may choose f with growth slow enough that \mathcal{M}_f satisfies independent n amalgamation over finite algebraically closed sets for any $n \geq 3$ or even for all $n \in \mathbb{N}$.*

Proof. See Appendix 6. □

As we shall see, the choice of values of $f(t)$ for $t \leq 6$ ensures that the eventual closures of graphs of small enough dimension are easy to enumerate.

Definition 5.3. Let $A \in \mathcal{K}_f$. Suppose $A \leq^* A' \in \mathcal{K}_f, \delta(A') = \delta(A)$, and $|A' \setminus A| = 1$. Then, we call A' a **one point extension** of A .

Since $\alpha = 2$, a one-point extension A' of A will be obtained by adding to A a single vertex v joined by two edges to A .

Lemma 5.4. *Let $A, B \in \mathcal{K}_f, A \leq^* B, \delta(A) = \delta(B)$, and $|B \setminus A| < 6$. Then, B is obtained from A by iterating one-point extensions.*

Proof. Note that with our choice of f for $X \in \mathcal{K}_f$ such that $|X| < 6, |E(X)| < |X|$. In fact, for $|X| < 6, f(X) > |X|$. Since $X \in \mathcal{K}_f$,

$$\begin{aligned} 2|X| - |E(X)| &= \delta(X) \geq f(|X|) > |X| & (10) \\ \Rightarrow |E(X)| &< |X|. & (11) \end{aligned}$$

Now, consider the graph X induced by the vertices in $B \setminus A$. Since $|X| < 6, |E(X)| < |X|$. Let $E(A; X) := \{\{u, v\} \in E(B) | u \in A, v \in X\}$. Since $\delta(B) = \delta(A)$, we have that:

$$\delta(B) = 2|B| - E(B) = 2|A| + 2|X| - |E(A)| - |E(X)| - |E(A; X)| = 2|A| - |E(A)| = \delta(A),$$

and so since $|E(X)| < |X|$,

$$0 = 2|X| - |E(X)| - |E(A; X)| > |X| - |E(A; X)|$$

Hence, $|X| < |E(A; X)|$. By the pigeonhole principle there must be a vertex v of X connected to A by two or more points. Note that if v were connected by three or more points $\delta(A \cup \{v\}) < \delta(A)$, and so $A \not\leq^* B$. Hence, $A \cup \{v\}$ must be a one-point extension of A . The same argument can be iterated for $A \cup \{v\}$ so that we see B is constructed by iterating one-point extensions. \square

Remark 5.5. Let D be a proper eventual closure of $A_1 \amalg_{A_0} A_2$ with $|D \setminus (A_1 \cup A_2)| < 6$. Then, as noted in Lemma 5.1, D is obtained by iterating one-point extensions D_1, \dots, D_k for $k \leq 5$. One can see that each D_i is also an eventual closure.

We shall call an eventual closure of $A_1 \amalg_{A_0} A_2$ which is also a one-point extension a **one point closure**.

Given Lemma and Corollary 4.11 we get:

Corollary 5.6. *Let $A \leq \mathcal{M}_f$, and $A \leq B, C \in \mathcal{K}_f$. Let $N = \lfloor f^{-1}(\delta(B \amalg_A C)) \rfloor$ so that N is the largest size of a graph of predimension $\delta(B \amalg_A C)$. Suppose that $N - |B \amalg_A C| < 6$ and that the maximal distance between a point in $B \setminus A$ and a point in $C \setminus A$ in $B \amalg_A C$ is ≤ 3 . Then there is no proper eventual closure for $B \amalg_A C$, and so,*

$$\mu(B \amalg_A C) = \frac{\mu(B)\mu(C)}{\mu(A)}.$$

Proof. Since no graph in \mathcal{K}_f contains cycles of length < 6 , $B \amalg_A C$ has no proper eventual closures. In fact, in a one point closure of $B \amalg_A C$ by a point v , v must be attached to a vertex in $B \setminus A$ and a vertex in $C \setminus A$ since B and C are closed in the eventual closure. But then, this one-point closure would contain a k -cycle for $k < 6$. \square

5.2 Proving non-measurability

We are ready to prove that the structures built in Construction 5.1 are not *MS*-measurable. We first find some equations that an *MS*-measure should satisfy and then note that these equations imply that some definable set has measure zero, breaking the positivity condition of *MS*-measures.

Notation 5.7. We still follow the notation of 4.7. However, for clarity, we represent graphs pictorially. For example, for $A \in \mathcal{K}_f$ a path of length two, instead of writing $\mu(A)$, we may write $\mu(\curvearrowright)$, so that it is clear from our notation which graph we are talking about. Note that when we speak of "the measure of A ", we actually mean "the measure of the definable subset of $M_f^{|A|}$ consisting of copies of A which are such that $A \leq M_f$ ". By our choice of f , points and edges form complete types in \mathcal{M}_f .

Proposition 5.8. *Suppose that \mathcal{M}_f is *MS*-measurable with dimension given by *SU*-rank. Assume without loss of generality that the measure of a single vertex is 1. Let λ be the measure of an edge. Then, μ must satisfy the following equations:*

$$\mu(\curvearrowright) = \lambda^2 \tag{12}$$

$$\mu(\cdot \cdot) + \mu(\curvearrowright) = 1 \tag{13}$$

$$\mu(\sqsupset) = \lambda^3 \tag{14}$$

$$\mu(\cdot \dashrightarrow) + \mu(\sqsupset) = \lambda(1 - \lambda^2) \tag{15}$$

$$\mu(\cdot \dashrightarrow) = \frac{\mu(\cdot \dashrightarrow)^2}{(1 - \lambda^2)} \tag{16}$$

$$\mu\left(\begin{array}{c} \vdots \\ \dashrightarrow \end{array}\right) = \lambda^4 \tag{17}$$

$$\mu(\cdot \overset{\cdot}{\cdot}) + \mu\left(\begin{array}{c} \cdot \\ \cdot \\ \cdot \end{array}\right) = (1 - \lambda^2)^2 \lambda^2. \quad (18)$$

Proof. Equation 18 is obtained with a different technique from the rest. The proof for equations 12-17 consists in identifying various free amalgamations in the graphs involved and in the repeated use of Corollary 5.6 or the fact that all eventual closures we will consider are obtained by iterating one-point extensions and using Theorem 4.11. Again, to obtain these equations we consider the eventual closures D_i of $B \amalg_A C$. Now, for clarity, we shall illustrate this through pictures by colouring a copy $B \amalg_A C$ so that the copy of A is in purple, the copy of B in blue and the copy of C in orange. When dealing with one-point extensions we shall colour the additional points in green.

Let us begin with equation 12. Consider a path of length two as a free amalgamation of two paths of length one over a point as in Figure 2. This free amalgamation has no further eventual closures. Since the measure of an edge is λ and the measure of a point is 1, equation 12 follows from Corollary 5.6.



Figure 2: Note that this path has already maximal size for its dimension as $f(3) = 4$ and so there are no further eventual closures.

Now, for equation 13, consider the free amalgamation of two points over the empty set. This does have a one point closure, namely a copy of a path of length two (by joining the one point of the extension to the two distinct points in the amalgamation). There cannot be further extensions since $f(3) = 4$, and so the resulting graph has maximal size for its dimension. Since vertices have measure one in \mathcal{M}_f , equation 13 follows. Note that as a consequence of equations 13 and 12, we obtain that

$$\mu(\cdot \cdot) = (1 - \lambda^2),$$

and we shall use this fact in future equations.

Equations (14)-(17) are explained in Figures 3-6.



Figure 3: There are no further eventual closures since $f(4) = 5$ and so we have maximal size for this dimension. This amalgamation yields equation 14 recalling that the two paths of length two in the amalgamation have measure λ^2 , and that the path over which they are amalgamated has measure λ .



Figure 4: We can obtain a one point closure by joining a point to the points in $B \setminus A$ and $C \setminus A$. It is the unique one point closure since $|B \setminus A| = |C \setminus A| = 1$. There are no larger eventual closures since $f(4) = 5$. Thus, we get equation 15, recalling that the measure of two disjoint points is $(1 - \lambda^2)$.



Figure 5: The free amalgamation has dimension 6. Since $f(6) = 6$ and the points in $B \setminus A$ and $C \setminus A$ are at distance two from each other, from Corollary 5.6 we obtain the desired equation 16.



Figure 6: Equation 17 follows similarly by Corollary 5.6.

We are now ready to obtain equation 18. For it we can use Corollary 3.6 since the structures we are working with have weak elimination of imaginaries, as proven in the appendix 6. Since in \mathcal{M}_f types of finite tuples are determined by the quantifier-free type of their closures, there are only two 2-types over the emptyset, $p_1(x, y)$ and $p_2(x, y)$, such that $p_i(a, b)$ implies $a \perp b$. These are isolated by the formulas $\phi(x, y)$ and $\psi(x, y)$, saying respectively that x and y are at distance two from each other and that they are at distance > 2 . We consider the measure of $\psi(x_1, x_2) \wedge \psi(x_2, x_3) \wedge \phi(x_1, x_3)$. By Corollary 3.6, we have

$$\mu(\psi(x_1, x_2) \wedge \psi(x_2, x_3) \wedge \phi(x_1, x_3)) = \mu(\psi(x_1, x_2))^2 \mu(\phi(x_1, x_3)) = (1 - \lambda^2)^2 \lambda^2.$$

We wish to express $\psi(x_1, x_2) \wedge \psi(x_2, x_3) \wedge \phi(x_1, x_3)$ as a disjoint union of 3-types of maximal dimension in order to expand the left hand side of this equation. Let us look at the complete 3-types $\text{tp}(a_1 a_2 a_3)$ such that $a_1 a_2 a_3 \models \psi(x_1, x_2) \wedge \psi(x_2, x_3) \wedge \phi(x_1, x_3)$ and $a_i \perp a_j a_k$ for $\{i, j, k\} = \{1, 2, 3\}$ distinct. Let A be the graph in \mathcal{K}_f given by the following picture:




Figure 7: The graph A . Note that if $A \leq M_f$, then $M_f \models \psi(v_1, v_2) \wedge \psi(v_2, v_3) \wedge \phi(v_1, v_3)$ and v_i is independent from $v_j v_k$ for $\{i, j, k\} = \{1, 2, 3\}$.

Again, since the types of these triples are determined by the quantifier-free types of their closures, we may focus on the graphs B in \mathcal{K}_f such that, $A \leq^* B$, $\delta(B) = \delta(A) = 6$, and $v_1 v_2, v_1 v_3, v_2 v_3 \leq B$. Note that since $\delta(B) = 6$, B may have at most 6 points and must be obtained by iterating one-point extensions on the graph A .



Figure 8: Note that since $v_1 v_2, v_1 v_3, v_2 v_3 \leq B$, any one-point extension of A by a vertex w must be such that w has an edge with u . Hence, since we avoid squares, the only possible one-point extension is given by joining w to u and v_1 . Call the graph so obtained B' . Note that B' doesn't have any one-point extension since any two points of B' have distance ≤ 3 from each other and M_f avoids cycles for of size < 6 .

Let $A(v_1, v_2, v_3, u)$ be the formula expressing $v_1 v_2 v_3 u \cong A \leq M_f$ and $B'(v_1, v_2, v_3, u, w)$ be the formula expressing $v_1 v_2 v_3 u w \cong B' \leq M_f$, where B' is described in Figure 8. From the argument in Figure 8 we can see that there are only two 3-types $p(x_1, x_2, x_3)$ which are completions of $\psi(x_1, x_2) \wedge \psi(x_2, x_3) \wedge \phi(x_1, x_3)$ and which satisfy our independence requirements, and they are isolated by $\exists u A(x_1, x_2, x_3, u)$ and $\exists u, w B'(x_1, x_2, x_3, u, w)$. Hence,

by additivity,

$$\mu(\exists u A(x_1, x_2, x_3, u)) + \mu(\exists u, w B'(x_1, x_2, x_3, u, w)) = \mu(\psi(x_1, x_2) \wedge \psi(x_2, x_3) \wedge \phi(x_1, x_3)).$$

Consider the projection $\pi : M_f^4 \rightarrow M_f^3$ onto the first three coordinates. The restriction of this map to $A(M^4)$ is injective, and so by Fubini and algebraicity,

$$\mu(\exists u A(x_1, x_2, x_3, u)) = \mu(A(x_1, x_2, x_3, u)) = \mu(\dot{}).$$

Similarly, considering the projection $\pi' : M_f^5 \rightarrow M_f^3$ onto the first three coordinates and $B(M^5)$,

$$\mu(\exists u, w B'(x_1, x_2, x_3, u, w)) = \mu(B'(x_1, x_2, x_3, u, w)) = \mu\left(\begin{array}{c} \dot{} \\ \vdots \\ \dot{} \end{array}\right).$$

This yields the desired equation. \square

Theorem 5.9. *The structures of the form \mathcal{M}_f built following Construction 5.1 are not MS-measurable. Hence, there are ω -categorical supersimple finite SU-rank structures which are not MS-measurable. Indeed, there are supersimple ω -categorical structures of finite SU-rank and with independent n -amalgamation for all n which are not MS-measurable.*

Proof. By Corollary 2.22 we know that if \mathcal{M}_f is MS-measurable, it also has an MS-dimension-measure where the dimension is given by SU-rank. However, the equations from Proposition 5.8 imply that $\lambda = 0$. This contradicts the positivity assumption for the measure in MS-measurable structures. To see this note that equations 14 and 15 imply that

$$\mu(\dot{}) = \lambda - 2\lambda^3.$$

From equations 16, 17 and 18 we then get that

$$\frac{\lambda^2(1 - 2\lambda^2)^2}{(1 - \lambda^2)} + \lambda^4 = (1 - \lambda^2)^2 \lambda^2.$$

Simplifying, we obtain that $\lambda = 0$. \square

While we had a particular choice for α and the initial values of f , it is plausible that similar results should hold for different choices of both. This raises the question of whether in general, non-trivial ω -categorical Hrushovski constructions are not MS-measurable. This would provide further evidence for the conjecture that MS-measurable ω -categorical structures are one-based.

In subsequent work [16], we find more general reasons for which various classes of ω -categorical Hrushovski constructions are not MS-measurable by studying invariant Keisler measures in these structures. In particular, we prove that ω -categorical MS-measurable structures must satisfy a stronger version of the independence theorem and that the equation in Corollary 3.6 holds also when one of the pairs is weakly algebraically independent. We also prove that in the structures \mathcal{M}_f we introduced, the formula asserting that " x has distance two from a " doesn't fork over the empty-set but is universally measure zero. While our results make it implausible, it remains an open question whether any one-based ω -categorical Hrushovski construction is MS-measurable.

6 Appendix: Proving the main properties of our Hrushovski constructions

In Section 5 we introduced a class of ω -categorical supersimple finite SU-rank Hrushovski constructions and proved they are not MS-measurable. However, we omitted from the body of this article the proofs that our choice of f yields such structures. Indeed, the proof

of simplicity requires some effort. We include these technical results in this appendix.

We begin with an abstract discussion of how to prove simplicity in Subsection 6.1. Then, in Subsection 6.2 we move to proving that our choice of f in Construction 5.1 yields ω -categorical Hrushovski constructions and that these are supersimple of finite SU -rank, with weak elimination of imaginaries. We further note that these can be built so that they may satisfy independent n -amalgamation for all n .

6.1 Simplicity of ω -categorical Hrushovski constructions

In this subsection we discuss under which conditions on f ω -categorical Hrushovski constructions are supersimple of finite SU -rank. This was first explored in [12]. We simplify some of the conditions for supersimplicity discussed in [4] and [3]. This will shorten the proofs of supersimplicity of the structures considered in the next section. Much of this material is implicit in the proof of Theorem 3.6 in [3]. Here we make it explicit, and correct a mistake in Remark 3.8 of the same article.

Let us begin by reminding some basic properties of the d-closed and self-sufficient relations [6][Lemma 3.10]:

Lemma 6.1. *Let $C \in \overline{\mathcal{K}}$ and let \leq' stand for either \leq or \leq^* . Then, the following hold:*

1. *Let $A \leq' C$ be finite, and $B \subseteq C$. Then, $A \cap B \leq' B$.*
2. *Let A and B be finite such that $A \leq' B \leq' C$. Then, $A \leq' C$.*
3. *Let A and B be finite with $A, B \leq' C$. Then, $A \cap B \leq' C$.*

Definition 6.2 (Independence Theorem Diagram). Let D be a graph. Suppose that D has subgraphs D_i and D_{ij} for $0 \leq i < j \leq 3$ all contained in \mathcal{K}_f . To simplify our notation, we allow inverting indices, e.g. $D_{12} = D_{21}$. We say that $(D; D_i; D_{ij})$ is an **independence theorem diagram** (ITD) with respect to \mathcal{K}_f if the following hold:

- $D_{0j} = D_j$ for $1 \leq j \leq 3$;
- $D_i \cap D_j = D_0$ for $1 \leq i < j \leq 3$;
- $D_i, D_j \leq D_{ij}$ for $1 \leq i < j \leq 3$;
- $D_{ij} \cap D_{jk} = D_j$ for $\{i, j, k\} = \{1, 2, 3\}$;
- D_i and D_j are independent over D_0 in D_{ij} for $1 \leq i < j \leq 3$;
- Any edge in D is entirely contained within some D_{ij} for $1 \leq i < j \leq 3$.

Figure 9 gives a visual representation of an independence theorem diagram.

We say that $(D; D_i; D_{ij})$ is a **proper ITD** when it satisfies the following additional conditions:

- $D_0 \subsetneq D_i$ for $1 \leq i \leq 3$;
- $D_i \cup D_j \subsetneq D_{ij} = \text{cl}_D(D_i \cup D_j)$ for $1 \leq i < j \leq 3$;

We say that \mathcal{K}_f is **closed under ITDs** if, whenever $(D; D_i; D_{ij})$ is an ITD, we have that $D \in \mathcal{K}_f$.

We note some basic properties of independence theorem diagrams and proper ITDs.

Lemma 6.3. *Suppose $(D; D_i; D_{ij})$ is an ITD for \mathcal{K}_f . Then,*

1. $D_0 \leq D_{ij}$ for $0 \leq i < j \leq 3$;
2. $D_{ij} \leq D$;




Figure 9: A visual representation of an independence theorem diagram. The first picture represents an ITD with its various parts labelled. To distinguish the parts more clearly, the last three pictures show as a shaded area D_0 , D_1 , and D_{12} respectively.

$$3. D_{ij} \leq D_{ij} \cup D_{jk};$$

$$4. D_{ij} \cup D_{jk} \leq^* D;$$

Proof. Part 1 is a direct consequence of Lemma 6.1.1. Parts 2,3, and 4 are from [4][Lemma 1.6]. \square

From [3][Corollary 2.24 & Theorem 3.6] we know that:

Theorem 6.4. *Suppose that $\alpha \in \mathbb{N}$ and \mathcal{K}_f is closed under ITDs. Then, \mathcal{M}_f is supersimple of finite SU-rank.*

Hence, it is sufficient to include all ITDs in \mathcal{K}_f in order to have supersimplicity and finite SU-rank for \mathcal{M}_f . However, this is somehow not easy to check since to see whether D is in \mathcal{K}_f we need to check that any of its subgraphs $B \subset D$ is also in \mathcal{K}_f . The following lemmas help us simplifying this process.

Lemma 6.5. *Let $C \leq D \in \mathcal{K}_f$, where D is an ITD. Then, C is an ITD.*

Proof. Call C_{ij} the intersection of C and D_{ij} . The only conditions that we need to check are that $C_i \leq C_{ij}$ and that C_i and C_j are independent over C_0 in C_{ij} .

Now, the first condition follows by Lemma 6.1.1 since $D_i \leq D_{ij}$ and $C_{ij} \subseteq D_{ij}$ implies $C_i = D_i \cap C_{ij} \leq C_{ij}$.

For the second condition, first note that since D_i and D_j are freely amalgamated over D_0 in D_{ij} , so must be C_i and C_j over C_0 in C_{ij} , being just intersections of these sets. We know that $D_i \cup D_j \leq^* \text{cl}_{D_{ij}}(D_i \cup D_j)$, $\text{cl}_{C_{ij}}(C_i \cup C_j) \cap (D_j \cup D_i) = C_i \cup C_j$, and $\text{cl}_{C_{ij}}(C_i \cup C_j) \cap \text{cl}_{D_{ij}}(D_i \cup D_j) = \text{cl}_{C_{ij}}(C_i \cup C_j)$. Hence, by Lemma 6.1.1, $C_i \cup C_j \leq^* \text{cl}_{C_{ij}}(C_i \cup C_j)$. But then, by Lemma 4.5, we know that C_i and C_j are independent over C_0 in C_{ij} . \square

Hence, in proving supersimplicity, we can reduce the number of graphs for which we check $f(|D|) \leq \delta(D)$ by only focusing on proper ITDs.

Lemma 6.6. *The following are equivalent:*

1. \mathcal{K}_f is closed under ITDs;
2. \mathcal{K}_f is closed under proper ITDs;
3. For each proper ITD D we have that $f(|D|) \leq \delta(D)$.

Proof. (1) \Rightarrow (2) \Rightarrow (3) follows from the definitions of strong ITD and \mathcal{K}_f . Let us focus then on (2) \Rightarrow (1). Let D be an ITD. For it to be in \mathcal{K}_f we need to show that for any $C \subseteq D$, $f(|C|) \leq \delta(C)$. Note that if $f(|C|) > \delta(C)$, also $\text{cl}_D(C)$ satisfies this condition since

$$f(|\text{cl}_D(C)|) \geq f(|C|) > \delta(|C|) \geq \delta(\text{cl}_D(C)),$$

where the first inequality holds since f is increasing and the last by definition of closure. Hence, we may assume that $C \leq D$. We shall show that if C is not already a proper ITD, it must already be in \mathcal{K}_f , or that there is a proper ITD C' , such that if C' is in \mathcal{K}_f , then so is C .

We label the intersections of C with D so that $C_i := D_i \cap C$ and $C_{ij} := D_{ij} \cap C$. By the previous lemma C is an independence theorem diagram.

Suppose that $C_1 = C_0$, then C is in \mathcal{K}_f , being obtained by freely amalgamating C_{12} and C_{23} over C_2 , and then freely amalgamating this structure with C_{13} over C_3 . Hence, if $C_i = C_0$ for any $1 \leq i \leq 3$, $C \in \mathcal{K}_f$.

Suppose that $C_{12} = C_1 \cup C_2$. Then C is the free amalgamation of C_{23} and C_{13} over C_3 . Hence if $C_{ij} = C_i \cup C_j$, then $C \in \mathcal{K}_f$.

Finally, consider the subgraph of C' of C which constitutes the proper independence theorem diagram obtained by taking $C'_{ij} := \text{cl}^C(C_i \cup C_j)$. Now, by construction and definition of closure, we must have $C'_{ij} \leq C_{ij}$. Furthermore, $C'_{ij} \leq C'$. Hence, C may be obtained as a free amalgamation of C' and C_{ij} over C'_{ij} (eventually repeating this operation for the different C'_{ij} 's).

So, we have seen that if all proper ITDs are in \mathcal{K}_f , then so are all ITDs, and so (2) \Rightarrow (1). We prove that (3) \Rightarrow (2) holds by considering a minimal counterexample. Suppose that D is a minimal proper ITD not in \mathcal{K}_f . Since $f(|D|) \leq \delta(D)$, there must be some $C \subseteq D$ such that $f(|C|) > \delta(|C|)$. Again, we may assume that $C \leq D$. But now, by the previous lemma, C is an ITD. Hence, by the above steps either it will be either a free amalgamation of graphs in \mathcal{K}_f , or the free amalgamation of graphs in \mathcal{K}_f and some proper ITD, or it will be a proper ITD again. But any of these cases implies that $C \in \mathcal{K}_f$. So it is sufficient to check the condition $f(|D|) \leq \delta(D)$ in proper ITDs. \square

Lemma 6.7. *Let $\alpha \in \mathbb{N}$, \mathcal{K}_f be a free amalgamation class. Let $n_1 \in \mathbb{N}$. Suppose that for $n_1 \leq t$, $f(3t) \leq f(t) + k$, for fixed $k \in \mathbb{N}$. Let D be an ITD. Without loss of generality, say that D_{12} is the D_{ij} of biggest predimension, and call its predimension d_{12} . Suppose that $f(n_1) \leq d_{12}$. We have that if $\delta(D) \geq d_{12} + k$, then $f(|D|) \leq \delta(D)$.*

Proof. This proof is substantially identical to the final part of the Theorem 3.6 in [3]. However, we repeat the argument for completeness and in order to avoid confusion with Remark 3.8 [3], which follows the theorem and contains a mistake.

Let g be the inverse of f . Making the substitution $g(x) = t$ into $f(3t) \leq f(t) + k$, we get that $3g(x) \leq g(x + k)$ for $f(n_1) \leq x$. Then, for $f(n_1) \leq d_{12}$:

$$|D| \leq \sum_{1 \leq i < j \leq 3} |D_{ij}| \leq \sum_{1 \leq i < j \leq 3} g(\delta(D_{ij})) \leq 3g(d_{12}) \leq g(d_{12} + k) \leq g(\delta(D))$$

Where the second inequality holds since $D_{ij} \in \mathcal{K}_f$ and the fourth holds since $\delta(D) \geq d_{12} + k$ and g is increasing. Note that the resulting inequality is equivalent to $f(|D|) \leq \delta(D)$. \square

Hence, we have a method to obtain supersimple ω -categorical Hrushovski constructions of finite SU -rank. And now we know how to verify this more easily.

6.2 The structures \mathcal{M}_f and their supersimplicity

In this subsection we focus on our choice of structures \mathcal{M}_f built as specified in Construction 5.1. We prove that these structures have the various properties mentioned in the main article.

We begin by noting that \mathcal{K}_f is a free amalgamation class and so the structures \mathcal{M}_f may be built according to Theorem 4.3.

Proposition 6.8. *The class \mathcal{K}_f is a free amalgamation class.*

Proof. Since f is a good function for $t \geq 6$, by Lemma 4.4, we just need to show that for $A_0 \leq A_1, A_2 \in \mathcal{K}_f$, then $A_1 \amalg_{A_0} A_2 \in \mathcal{K}_f$ for $|A_i| \leq 6$. Note that $|A_1 \amalg_{A_0} A_2| = |A_1| +$

$|A_2| - |A_0|$, and $\delta(A_1 \amalg_{A_0} A_2) = \delta(A_1) + \delta(A_2) - \delta(A_0)$. Hence, in the context of Figure 1, it is sufficient to check that given any three red points p, q , and r (possibly with $q = r$) with $p = (p_1, p_2), q = (q_1, q_2), r = (r_1, r_2)$ such that $p_1 < q_1, r_1$ and $p_2 < q_2, r_2$, the fourth vertex of the parallelogram with edges \overline{pq} and \overline{pr} is still above the function f . Since $f(18) \leq 7$ and our function is increasing this can be easily verified. \square

Now we prove that \mathcal{M}_f is supersimple of SU -rank 2. We proceed step by step by proving that for any proper ITD D for \mathcal{K}_f , $f(|D|) \leq \delta(D)$. Hence, the proof will follow by Lemma 6.6 and Theorem 6.4. We shall adopt the notation we already set for proper ITDs. To avoid confusion when speaking of the various element of an ITD, we always write D for the proper ITD, we write D_{ij} only for $i, j \in \{1, 2, 3\}$, and consistently write D_j for D_{0j} . Furthermore, given D , we shall assume without loss of generality that D_{12} has maximal dimension among the D_{ij} . We shall also write d_i for $\delta(D_i)$, d_{ij} for $\delta(D_{ij})$ and d for $\delta(D)$ to simplify our notation.

Lemma 6.9. *For all proper ITDs D for \mathcal{K}_f with $d_{12} \leq 4$, $f(|D|) \leq \delta(D)$. For this proof we only need to assume that $f(1) = 2, f(2) = 3, f(3) = 4$ and $f(6) \leq 6$.*

Proof. For $d_{12} \leq 4$, $|D_{12}| \leq 3$, and so $|D_{ij}| \leq 3$. Since we require $D_i, D_j \neq \emptyset$ and $D_i \cup D_j \subsetneq D_{ij}$ (for $i \neq j \in \{1, 2, 3\}$), we must have that $|D_{ij}| = 3$ for each $i \neq j \in \{1, 2, 3\}$, $|D_i| = 1$ for each $i \in \{1, 2, 3\}$, and $|D_0| = 0$. There is only one graph satisfying these requirements, i.e. a 6-cycle as shown in figure 10.




Figure 10: We color in D_1, D_2 , and D_3 in blue, purple, and pink respectively. Meanwhile, the points and edges in $D_{ij} \setminus D_i \cup D_j$ are in green. We can see that each D_{ij} must be a copy of a path of length 2 since $|D_{ij}| = 3$ and $d_{ij} \leq 4$. Since we know that each D_i is a single point for $i > 0$, the resulting proper ITD is a 6-cycle. \square

Lemma 6.10. *For all proper ITDs D for \mathcal{K}_f with $d_{12} \leq 5$, $f(|D|) \leq \delta(D)$. We only need to assume $f(1) = 2, f(2) = 3, f(3) = 4, f(4) = 5$, and $f(6) \leq 6$, and $f(12) \leq 7$.*

Proof. Given the previous lemma, we only need to prove the condition for $d_{12} = 5$. Note that since $f(3 \cdot 4) = f(12) \leq 7 = f(4) + 2$, by Lemma 6.7 we only need to check the case of $d = 6$.

We have that $d_0 > 0$ since otherwise, the condition that $d - d_{12} = 1$ forces $d_3 = 1$, which is impossible since no graph in \mathcal{K}_f has predimension 1. Hence, we need to check the cases of $d_0 = 2$ and $d_0 = 3$ (note that for $d_0 > 3$ we cannot have $d = 6$ since $d > d_0 + 3$). By definition of an ITD, the following inclusion-exclusion formula holds:

$$|D| = \sum_{1 \leq i < j \leq 3} |D_{ij}| - \sum_{1 \leq i \leq 3} |D_i| + |D_0|. \quad (19)$$

Hence, knowing the upper bounds for the sizes of the D_{ij} and the lower bounds for the sizes of the D_i we can come to an upper bound for D :

$$|D| \leq \sum_{1 \leq i < j \leq 3} \lfloor f^{-1}(d_i + d_j - d_0) \rfloor - \sum_{1 \leq i \leq 3} \min\{|B| \text{ s.t. } \delta(B) = d_i\} + |D_0| := \beta \quad (20)$$

Furthermore, note that $(d_1 - d_0) + (d_2 - d_0) + d_0 = d_{12}$ and that $d_3 - d_0 = d - d_{12} = 1$.

Hence, for $d_0 = 2$, without loss of generality we have $d_1 = 4, d_2 = 3, d_3 = 3$ (since we must have $d_1 - d_0 = 2, d_2 - d_0 = 1, d_3 - d_0 = 1$). Hence, we get that $\beta = 4 \cdot 2 + 3 - 2 \cdot 3 + 1 = 6$, and so $|D| \leq 6$. Since $f(6) \leq 6, f(|D|) \leq 6 = d$.

For $d_0 = 3$, we obtain that $d_1 - d_0 = 1, d_2 - d_0 = 1, d_3 - d_0 = 1$. Since $d_0 = 3$ implies that $|D_0| = 2, |D_i| \geq 3$ so $\beta = 3 \cdot 4 - 3 \cdot 3 + 2 = 5$, and so $f(|D|) \leq f(5) < 6 = d$, as desired. \square

Theorem 6.11. *Let \mathcal{K}_f and \mathcal{M}_f be as in Construction 5.1. Then, \mathcal{M}_f is supersimple of finite SU-rank. In particular, it has SU-rank 2.*

Proof. From Theorem 6.4 and Lemma 6.6 we need to check that for any proper ITD D , $f(|D|) \leq \delta(D)$. We know from the lemmas above for any proper ITD D for \mathcal{K}_f with $d_{12} \leq 5$, $f(|D|) \leq \delta(D)$. Since for $t \geq 6, f(3 \cdot t) \leq f(t) + 1$, by Lemma 6.7 we have that for $d_{12} \geq 6$, if $\delta(D) \geq d_{12} + 1$, then $f(|D|) \leq \delta(D)$. But $\delta(D) > d_{12}$ in any proper ITD. Hence, any proper ITD with $d_{12} \geq 6$ is such that $f(|D|) \leq \delta(D)$ and so the theorem follows. Finally, SU-rank coincides with the Hrushovski dimension, and so $SU(\mathcal{M}_f) = 2$. This follows from the characterisation of non-forking independence in terms of the dimension [3][Corollary 2.21]. \square

In the process of our proof of Theorem 6.11 we have proven that the smallest proper ITD in \mathcal{K}_f has predimension 6. Our conditions in Construction 5.1 put no constraints on how slowly $f(t)$ grows for $t > 6$. Hence, we can see that we may make f slow growing enough to satisfy independent n -amalgamation for any n of our choice. Indeed, by choosing $f(t)$ with growth the inverse of $(n + 1)!$ for $t \geq 6$, we can choose \mathcal{M}_f to have independent n amalgamation for all n [12].

Finally, we note that structures of the form of \mathcal{M}_f have weak elimination of imaginaries.

Proposition 6.12. *Let \mathcal{M}_f be as in Remark 5.2, then it has weak elimination of imaginaries.*

Proof. From Theorem 5.12, we know that \mathcal{M}_f satisfies conditions (P1)-(P5) from [3]. And these conditions are sufficient for weak elimination of imaginaries by Lemma 2.9 and Corollary 2.7 from [3]. \square

References

- [1] Richard Elwes. "Dimension and Measure in Finite First Order Structures". PhD thesis. School of Mathematics, University of Leeds, July 2005.
- [2] Richard Elwes and Dugald Macpherson. "A survey of asymptotic classes and measurable structures". In: *Model Theory with Applications to Algebra and Analysis*. Ed. by Z. Chatzidakis, D. Macpherson, A. Pillay, and A. Wilkie. Vol. 2. London Mathematical Society Lecture Note Series. Cambridge University Press, 2008, pp. 125–160. DOI: 10.1017/CB09780511735219.004.
- [3] David M. Evans. " \aleph_0 -categorical structures with a predimension". In: *Annals of Pure and Applied Logic* 116.1 (2002), pp. 157–186. ISSN: 0168-0072.
- [4] David M. Evans and Mark W. H. Wong. "Some remarks on generic structures". In: *The Journal of Symbolic Logic* 67.4 (2009), pp. 1143–1154. ISSN: 0022-4812.
- [5] David M. Evans. *Higher amalgamation properties in measured structures*. 2022. DOI: 10.48550/ARXIV.2202.10183. URL: <https://arxiv.org/abs/2202.10183>.
- [6] David M. Evans. *Homogeneous Structures ω -categoricity and Amalgamation Constructions*. [Online notes] Available from: http://www.imperial.ac.uk/~dmevans/Bonn2013_DE.pdf [Last Accessed: 12 December 2019]. 2013.
- [7] William Timothy Gowers. "A new proof of Szemerédi's theorem for arithmetic progressions of length four". In: *Geometric & Functional Analysis GAFA* 8.3 (1998), pp. 529–551.
- [8] Ehud Hrushovski. "Stable Group Theory and Approximate Subgroups". In: *Journal of the American Mathematical Society* 25.1 (2012), pp. 189–243. ISSN: 08940347, 10886834. URL: <http://www.jstor.org/stable/23072155>.

- [9] Ehud Hrushovski. *A stable \aleph_0 -categorical pseudo-plane*. Unpublished notes. 1987.
- [10] Ehud Hrushovski. “Approximate Equivalence Relations”. Unpublished notes. 2015.
- [11] Ehud Hrushovski. “Pseudo-finite fields and related structures”. In: *Model Theory and Applications*. Vol. 11. Quaderni di Matematica. Aracne, Rome, 1991, pp. 151–212.
- [12] Ehud Hrushovski. *Simplicity and the Lascar group*. Unpublished notes. 1998.
- [13] Byunghan Kim. *Simplicity theory*. Oxford logic guides ; 53. Oxford, 2014. ISBN: 9780198567387.
- [14] Byunghan Kim and Anand Pillay. “Simple Theories”. In: *Annals of Pure and Applied Logic* 88.2 (1997). Joint AILA-KGS Model Theory Meeting, pp. 149–164. ISSN: 0168-0072. DOI: [https://doi.org/10.1016/S0168-0072\(97\)00019-5](https://doi.org/10.1016/S0168-0072(97)00019-5). URL: <https://www.sciencedirect.com/science/article/pii/S0168007297000195>.
- [15] Dugald Macpherson and Charles Steinhorn. “One-dimensional asymptotic classes of finite structures”. In: *Transactions of the American Mathematical Society* Vol. 360.n.1 (2008), pp. 411–448. ISSN: 0002-9947.
- [16] Paolo Marimon. *Ergodic Keisler measures in ω -categorical Hrushovski constructions*. Unpublished notes. 2022.
- [17] Anand Pillay. *Geometric stability theory*. Oxford logic guides ; 32. Oxford: Clarendon Press, 1996. ISBN: 9780198534372.
- [18] Vojtech Rödl, Brendan Nagle, Jozef Skokan, Mathias Schacht, and Yoshiharu Kohayakawa. “The hypergraph regularity method and its applications”. In: *Proceedings of the National Academy of Sciences* 102.23 (2005), pp. 8109–8113.
- [19] Katrin Tent and Martin Ziegler. *A Course in Model Theory*. Lecture Notes in Logic. Cambridge University Press, 2012. DOI: 10.1017/CB09781139015417.
- [20] Henry Towsner. *A Model Theoretic Proof of Szemerédi’s Theorem*. 2011. arXiv: 1002.4456 [math.LO].
- [21] Henry Towsner. “An analytic approach to sparse hypergraphs: hypergraph removal”. In: *Discrete analysis* (2018). ISSN: 2397-3129.
- [22] Frank O. Wagner. “Dimensional Groups and Fields”. In: *The Journal of Symbolic Logic* 85.3 (2020), pp. 918–936. DOI: 10.1017/jsl.2020.48.
- [23] Frank O. Wagner. “Relational structures and dimensions”. In: *Automorphisms of First-Order Structures*. Ed. by R. Kaye and D. Macpherson. Clarendon Press, 1994, pp. 153–180.