

ZILBER–PINK IN A PRODUCT OF MODULAR CURVES ASSUMING MULTIPLICATIVE DEGENERATION

CHRISTOPHER DAW AND MARTIN ORR

Dedicated to the memory of Susan Daw (1954–2022)

ABSTRACT. We prove the Zilber–Pink conjecture for curves in $Y(1)^n$ whose Zariski closure in $(\mathbb{P}^1)^n$ passes through the point (∞, \dots, ∞) , going beyond the asymmetry condition of Habegger and Pila. Our proof is based on a height bound following André’s G-functions method. The principal novelty is that we exploit relations between evaluations of G-functions at unboundedly many non-archimedean places.

1. INTRODUCTION

One of the central problems of unlikely intersections is the **Zilber–Pink conjecture** for Shimura varieties:

Conjecture 1.1. *Let V be an irreducible subvariety of a Shimura variety S . If V is not contained in a special subvariety of positive codimension, then the intersection of V with the union of all special subvarieties of S of codimension greater than $\dim(V)$ is not Zariski dense in V .*

An immediate corollary of Conjecture 1.1 is the André–Oort conjecture: the special subvarieties of S are characterised among the irreducible subvarieties of S as those containing a Zariski dense set of special points. A proof of the André–Oort conjecture was recently announced by Pila, Shankar and Tsimerman [PST], who, using work of Esnault and Groechenig, give height bounds for special points, thereby completing a new strategy of Binyamini, Schmidt and Yafaev [BSY].

In [HP12], Habegger and Pila made the first progress on Conjecture 1.1 beyond the André–Oort conjecture, proving it for so-called “asymmetric curves” in $Y(1)^n$. Here $Y(1)$ denotes the level-one modular curve, which is isomorphic to the affine line \mathbb{A}^1 . An irreducible curve $C \subset Y(1)^n$ is called **asymmetric** if the restrictions of the coordinate functions on $Y(1)^n$ to C have distinct degrees, up to at most one exceptional pair of coordinates, and ignoring coordinates whose restrictions to C are constant. Habegger and Pila’s theorem was as follows.

Theorem 1.2. [HP12, Theorem 1] *Let C be an irreducible asymmetric curve in $Y(1)^n$ defined over $\overline{\mathbb{Q}}$. If C is not contained in a special subvariety of positive codimension, then the intersection of C with the union of all special subvarieties of $Y(1)^n$ of codimension at least 2 is finite.*

2020 *Mathematics Subject Classification.* 11G18, 11G50, 14G35.

Removing the asymmetry condition in Theorem 1.2 has been a major challenge. The main result of this article is a **proof of Conjecture 1.1 for a different class of curves in $Y(1)^n$** , namely, those that “intersect infinity.” We say that an irreducible curve in $Y(1)^n \cong \mathbb{A}^n$ **intersects infinity** if its Zariski closure in $(\mathbb{P}^1)^n$ contains the point (∞, \dots, ∞) .

Theorem 1.3. *Let C be an irreducible curve in $Y(1)^n$ which intersects infinity and is defined over $\overline{\mathbb{Q}}$. If C is not contained in a special subvariety of positive codimension, then the intersection of C with the union of all special subvarieties of $Y(1)^n$ of codimension at least 2 is finite.*

The class of curves which intersect infinity neither contains nor is contained in the class of asymmetric curves. Examples of curves in $Y(1)^n$ which intersect infinity but are not asymmetric are given by lines in \mathbb{A}^n which are not parallel to any coordinate hyperplane. If a line is not contained in any diagonal hyperplane, then it is not contained in any special subvariety.

For each positive integer M , we denote by $\Phi_M(X, Y) \in \mathbb{Z}[X, Y]$ the classical modular polynomial. The following theorem is an example of a new case of the Zilber–Pink conjecture included in Theorem 1.3.

Theorem 1.4. *Let a, b be distinct, non-zero algebraic numbers. Then there are only finitely many $z \in \overline{\mathbb{Q}}$ for which there exist positive integers M, N such that the following equation holds:*

$$\Phi_M(z, z + a) = \Phi_N(z, z + b) = 0.$$

As well as proving Theorem 1.3, the aim of this article is also to **demonstrate a new method of proving lower bounds for Galois orbits**, the most difficult aspect of the Zilber–Pink conjecture. We use a method of André [And89, Ch. X] for studying fibres with large endomorphism rings in a family of abelian varieties, based on G-functions. Recently, this method has been used to obtain bounds for Galois orbits in other cases of the Zilber–Pink conjecture [BM21], [DO21a], [DO21b]. André’s method has also been applied by Papas in variations of Hodge structure, going beyond families of abelian varieties [Pap].

The key new contribution of this article is to use relations between evaluations of G-functions *at all non-archimedean places* to obtain lower bounds for Galois orbits, and thus new cases of the Zilber–Pink conjecture. Previous applications used only archimedean evaluations of G-functions (except [And95], which exploited only a fixed finite number of non-archimedean places). Ignoring non-archimedean places restricted the large endomorphism rings for which the method could prove Galois orbit bounds, preventing the method being applied to $Y(1)^n$, or to intersections with E^2 curves in the moduli space of abelian surfaces in [DO21b].

This article’s method can be generalised to other cases of the Zilber–Pink conjecture for curves in moduli spaces of abelian varieties, subject to a condition generalising the “intersects infinity” condition in Theorem 1.3. This condition

corresponds to multiplicative degeneration of the corresponding family of abelian varieties as in [And89, X, Theorem 1.3]. The method can also be used to prove versions of Theorem 1.3 with some variations of this degeneration condition, but removing it entirely is challenging.

1.A. Height bounds and bounds for Galois orbits. For a precise description of special subvarieties in $Y(1)^n$, see [HP12, Section 2.1]. Informally, a **special subvariety** is a subvariety of $Y(1)^n$ defined by a system of equations of two types:

- (i) $\Phi_M(x_i, x_j) = 0$ for some indices $i, j \in \{1, \dots, n\}$;
- (ii) $x_i = a$ for some index $i \in \{1, \dots, n\}$ and some CM j -invariant $a \in \overline{\mathbb{Q}}$.

A **strongly special subvariety** of $Y(1)^n$ is a subvariety defined by equations of type (i) only.

In [HP12, Theorem 2], the variant of Theorem 1.2 which dealt with intersections between C and non-strongly special subvarieties was proved, without the asymmetry restriction. Thus in order to prove Theorem 1.3, it suffices to restrict our attention to strongly special subvarieties.

Furthermore, in [HP12], the asymmetry condition was used only in proving the lower bound for Galois orbits [HP12, Lemma 4.2]. Thus, in this article, we only need to prove a new lower bound for Galois orbits of intersections between strongly special subvarieties and a curve which intersects infinity. Theorem 1.3 then follows via the Pila–Zannier method, exactly as in [HP12]. Our lower bound for Galois orbits is as follows.

Theorem 1.5. *Let $C \subset Y(1)^n$ be an irreducible algebraic curve defined over $\overline{\mathbb{Q}}$ that intersects infinity and is not contained in a special subvariety of positive codimension. Let Σ denote the set of points of C with at least two CM coordinates. There exist positive constants c_1 and c_2 with the following property. Let $i_1, i_2, i_3, i_4 \in \{1, \dots, n\}$ with $i_1 \neq i_2$, $i_3 \neq i_4$ and $\{i_1, i_2\} \neq \{i_3, i_4\}$. For all points $s = (s_1, \dots, s_n) \in C(\overline{\mathbb{Q}}) \setminus \Sigma$ satisfying*

$$\Phi_M(s_{i_1}, s_{i_2}) = \Phi_N(s_{i_3}, s_{i_4}) = 0$$

for some positive integers M and N , we have

$$[\mathbb{Q}(s) : \mathbb{Q}] \geq c_1 \max\{M, N\}^{c_2}.$$

The statement of Theorem 1.5 is the same as [HP12, Lemma 4.2], except that the “asymmetric curve” condition has been replaced by “intersects infinity” (and we always impose the condition $\{i_1, i_2\} \neq \{i_3, i_4\}$, which is not really a change from [HP12, Lemma 4.2] because the condition of intersecting infinity implies that no coordinate function is constant on C).

Habegger and Pila deduce [HP12, Lemma 4.2] from isogeny estimates of Masser–Wüstholz type and a height bound: after a suitable relabelling of the coordinates,

$$h(s_{i_1}) \leq c_3 \max\{1, \log(M)\}. \tag{1}$$

This height bound [HP12, (17)] is obtained by comparing heights of coordinates on C and Faltings heights, with a key input being the asymmetry condition.

The majority of this article is dedicated to proving an analogous height bound for curves that intersect infinity. Let $h : \mathbb{A}^n(\overline{\mathbb{Q}}) \rightarrow \mathbb{R}$ denote the (logarithmic) height defined in [HP12, Section 2.2]. We shall prove a variant of the following height bound.

Theorem 1.6. *Let $C \subset \mathbb{A}^n$ and Σ be as in Theorem 1.5. For every $\epsilon > 0$, there exist constants c_4 and c_5 such that, for all s, M, N as in Theorem 1.5, we have*

$$h(s) < c_4 \max\{[\mathbb{Q}(s) : \mathbb{Q}]^{c_5}, M^\epsilon\}.$$

The height bound in Theorem 1.6 is weaker than the analogous bound (1), but it is sufficient to deduce Theorem 1.5 in the same way as [HP12, Lemma 4.2]. We will not prove or use Theorem 1.6 directly, but rather a variant applying to a suitable finite cover of C . Theorem 1.6 can be deduced from Proposition 5.11 in the same way as Theorem 1.5 is deduced from Proposition 5.12 in Section 5.J.

1.B. Comparison with previous applications of André’s method. As in [And89, Ch. X], our proof of Theorem 1.6 is based on studying the relative periods of a family of abelian varieties \mathcal{A} over the curve C . The Taylor series of some of these period functions (the “locally invariant periods”) around a point where \mathcal{A} degenerates to a torus (in the case of Theorem 1.6, this point is (∞, \dots, ∞)) are power series with special Diophantine properties, called G-functions.

At points $s \in C$ where the corresponding abelian variety \mathcal{A}_s has additional endomorphisms, there are polynomial relations between the evaluations at s of these G-functions. Theorem 1.6 follows using a theorem of Bombieri: if there is a polynomial relation between the evaluations of a set of G-functions at a $\overline{\mathbb{Q}}$ -point s , valid at all places for which s lies inside the radius of convergence of the G-functions, and not coming from a “trivial” relation between the G-functions themselves, then the height of s is bounded in terms of the degree of the polynomial relation.

In [And89], and in previous applications of [And89] to Galois orbit bounds for the Zilber–Pink conjecture, only the archimedean evaluations of G-functions were interpreted as periods. In order to apply Bombieri’s theorem without finding relations between non-archimedean evaluations of G-functions, the extra endomorphisms of \mathcal{A}_s were used to ensure that s lies outside the radius of convergence of the G-functions at all non-archimedean places. This approach is only possible when $\text{End}(\mathcal{A}_s)$ does not inject into $\text{End}(\mathbb{G}_m^g) \otimes_{\mathbb{Z}} \mathbb{Q} \cong M_g(\mathbb{Q})$ (where $g = \dim(\mathcal{A}_s)$).

On the other hand, in Theorem 1.3, the exceptional abelian varieties \mathcal{A}_s have endomorphism rings which inject into $M_4(\mathbb{Q})$. For example, a product of non-CM elliptic curves $E_1 \times E_2 \times E_3 \times E_4$, where there are isogenies $E_1 \rightarrow E_2$ and $E_3 \rightarrow E_4$, has endomorphism algebra $M_2(\mathbb{Q}) \times M_2(\mathbb{Q})$. Thus, in order to apply Bombieri’s theorem, we need relations valid at non-archimedean places.

Relations between non-archimedean evaluations of G-functions were used in [And95, Théorème 1]. However in this result, as in [And89], the global relation needed for Bombieri’s theorem was obtained as a product of relations, one for each place. Thus in order to control the degree of this global relation, it was necessary to limit the number of non-archimedean places at which the G-functions could converge (by imposing an integrality condition at other places). Our construction of relations instead directly constructs a polynomial relation simultaneously valid at all non-archimedean places, based on the Tate uniformisation of elliptic curves in a family degenerating to a torus. At archimedean places, we still using the previous approach of taking a product of local relations, one for each place.

1.C. Effectivity. Our height bound Theorem 1.6 and Galois bound Theorem 1.5 can be made effective. The main work required to do this would be to determine the size and global radius of the G-functions F_λ and G_λ in equation (24) and the singularities of the associated differential operator.

As previously mentioned, the Zilber–Pink conjecture in the case of intersections between curves $C \subset Y(1)^n$ and *non-strongly special* subvarieties was already proved by [HP12, Theorem 2], without any condition on the curve C . When the special subvariety is defined by one CM coordinate and one pair of isogenous coordinates, the underlying Galois bound [HP12, Lemma 4.4] was ineffective due to its dependence on an estimate for Siegel zeros. (The remaining case, namely the André–Oort conjecture for a curve in $Y(1)^n$, is known effectively by a different method, due to Kühne [Küh12].) Our proof of Theorem 1.5 could be adapted to apply to intersections with non-special subvarieties, provided that C intersects infinity, thus leading to an effective Galois bound for such cases – the bound on CM discriminants would come from a height bound as in Theorem 1.6 and an estimate of Masser–Wüstholz type.

Thus, if the relevant point-counting step of the Pila–Zannier strategy were to be made effective, that could be combined with Galois bounds using the techniques of this article, to obtain an effective proof of the Zilber–Pink conjecture for curves in $Y(1)^n$ which intersect infinity.

1.D. Structure of the paper. We begin with some preliminary definitions and lemmas in Section 2. In Section 3, we describe two well-known families of elliptic curves: the “ $1/j$ ” family and the Tate family. We also describe periods of elliptic curves in terms of the relation between these families. In Section 4, we construct polynomial relations between the periods of isogenous elliptic curves, both archimedean and non-archimedean. In Section 5, we prove our main theorems: Theorem 1.5 and a variant of Theorem 1.6 by interpreting the relations from Section 4 as relations between evaluations of G-functions.

Acknowledgements. Both authors would like to thank the Engineering and Physical Sciences Research Council for its support via New Investigator Awards [EP/S029613/1 to C.D., EP/T010134/1 to M.O.]. They would also like to thank the organisers of the Mathematisches Forschungsinstitut Oberwolfach workshop 2216, *Diophantische Approximationen*, for the opportunity to speak about this work.

2. PRELIMINARIES

For the definition of a G-function we refer to [And89, I, 1.3]. For the notions regarding relations between G-functions, see [And89, VII, 5.1].

2.A. Embeddings of a number field. Let K be a number field. We define an **embedding** of K to be a homomorphism $K \rightarrow \mathbb{C}$ or \mathbb{C}_p for some prime p . If ι is an embedding of K , we write \mathbb{C}_ι for the complete algebraically closed field which is the target of ι .

If ι is an embedding of K and $s \in K$, then we write $s^\iota = \iota(s) \in \mathbb{C}_\iota$. Similarly, if S is an algebraic object over K (in particular, a polynomial, a power series, an algebraic variety or a morphism of algebraic varieties), then we write S^ι for the base change of S by $\iota: K \rightarrow \mathbb{C}_\iota$. This superscript notation sometimes carries the connotation that S^ι should be interpreted as an analytic, rather than algebraic, object over \mathbb{C}_ι : in particular, if $f \in K[[X]]$ then f^ι denotes the analytic function obtained by evaluating f on its disc of convergence in \mathbb{C}_ι .

We normalise the absolute value on \mathbb{C}_ι so that it extends the standard archimedean or p -adic absolute value on \mathbb{Q} .

2.B. Non-archimedean power series. Let $f(X) = \sum_{n=0}^{\infty} a_n X^n \in K[[X]]$ be a power series, where $(K, |\cdot|)$ is a complete field equipped with an absolute value. We write $R(f)$ for the radius of convergence of f . Recall that

$$R(f) = \frac{1}{\limsup_{n \rightarrow \infty} |a_n|^{1/n}}.$$

When $(K, |\cdot|)$ is non-archimedean, we also define

$$R^\dagger(f) = 1 / \sup_{n \in \mathbb{Z}_{>0}} |a_n|^{1/n}$$

Note that we ignore a_0 in the definition of R^\dagger . Clearly $R^\dagger(f) \leq R(f)$. It is easy to prove that if $R(f) > 0$, then $R^\dagger(f) > 0$.

The quantity $R^\dagger(f)$ is useful for understanding composition of power series over non-archimedean fields. We note the following bound:

Lemma 2.1. *Let $(K, |\cdot|)$ be a non-archimedean complete field equipped with an absolute value. Let $g \in K[[X]]$ and $x \in K$. If $|x| < R^\dagger(g)$, then*

$$|g(x) - g(0)| < R^\dagger(g)^{-1} |x|.$$

Proof. Let $g(X) = \sum_{n=0}^{\infty} b_n X^n$. Note that $R^\dagger(g) = \inf_{n \geq 1} |b_n|^{-1/n}$. Hence, for each $n \geq 1$, $R^\dagger(g)^n \leq |b_n|^{-1}$. Therefore, for each $n \geq 1$,

$$|b_n x^n| < |b_n| |x| R^\dagger(g)^{n-1} \leq R^\dagger(g)^{-1} |x| \cdot |b_n| R^\dagger(g)^n \leq R^\dagger(g)^{-1} |x| \cdot 1$$

Since also $|x| < R(g)$, $|b_n x^n| \rightarrow 0$ as $n \rightarrow \infty$. It follows that $\sup_{n \geq 1} |b_n x^n| < 1$. Because the absolute value is non-archimedean, this implies that $|g(x) - g(0)| < R^\dagger(g)^{-1} |x|$. \square

Lemma 2.2. *Let $(K, |\cdot|)$ be a non-archimedean complete field equipped with an absolute value. Let $f, g \in K[[X]]$ with $R(f) \geq 1$ and $g(0) = 0$. Then $R(f \circ g) \geq R^\dagger(g)$ and, for all $x \in K$ satisfying $|x| < R^\dagger(g)$, we have $f(g(x)) = (f \circ g)(x)$.*

Proof. Let $f(X) = \sum_{n=0}^{\infty} a_n X^n$ and $g(X) = \sum_{n=1}^{\infty} b_n X^n$.

Let $x \in K$ with $|x| < R^\dagger(g)$. Let $M_{|x|}(g) = \sup_{n \geq 1} |b_n x^n|$. As in the proof of Lemma 2.1 and by hypothesis, we have

$$M_{|x|}(g) < 1 \leq R(f).$$

Since also $|x| < R(g)$, the conditions of [Rob00, 6.1.5, Theorem] are satisfied. Therefore $R(f \circ g) > |x|$ and $f(g(x)) = (f \circ g)(x)$. \square

Over \mathbb{C} , composition of power series is described by the following classical result:

Lemma 2.3. *Let $f, g \in \mathbb{C}[[X]]$ with $g(0) = 0$. Let*

$$s = \sup\{r \leq R(g) : |g(x)| < R(f) \text{ for all } x \in K \text{ with } |x| < r\}.$$

Note that if $R(f) > 0$ and $R(g) > 0$, then $s > 0$. Then $R(f \circ g) \geq s$ and, for all $x \in K$ satisfying $|x| < s$, we have $f(g(x)) = (f \circ g)(x)$.

2.C. Notation. If \mathbb{C}_i is a complete field, we write $D(0, r, \mathbb{C}_i)$ for the open disc in \mathbb{C}_i of radius r , centred at 0.

If N is a non-zero integer, we write $d(N)$ for the number of positive divisors of N . We will use the following fact [HW08, Theorem 315]: for every $\epsilon > 0$,

$$d(N) \ll_{\epsilon} N^{\epsilon}. \quad (2)$$

3. FAMILIES OF ELLIPTIC CURVES

3.A. The “1/j” family of elliptic curves. Let $S = \mathbb{A}_{\mathbb{Q}}^1 \setminus \{1/1728\}$ and let $S^* = S \setminus \{0\}$. Let

$$\mathcal{E} = \left\{ ([x : y : z], s) \in \mathbb{P}^2 \times S : \right. \\ \left. y^2 z + xyz = x^3 + \frac{36s}{1 - 1728s} xz^2 + \frac{s}{1 - 1728s} z^3 \right\} \setminus \{([0 : 0 : 1], 0)\}.$$

This is the S -smooth locus of an elliptic scheme $\pi: \mathcal{E} \rightarrow S$. Indeed, the fibres of \mathcal{E} over S^* are elliptic curves and \mathcal{E}_s has j -invariant $1/s$. The fibre \mathcal{E}_0 is the smooth

locus of a nodal cubic. Therefore, the standard Weierstrass group law makes \mathcal{E} into an S -group scheme, with $\mathcal{E}_0 \cong \mathbb{G}_m$.

We obtain an invariant holomorphic differential form on \mathcal{E}/S by setting

$$\omega = \frac{dx}{2y + x} \in \Omega^1(\mathcal{E}/S).$$

3.B. The Tate family. Let $\iota: \mathbb{Q} \rightarrow \mathbb{C}_\iota$ denote an embedding of \mathbb{Q} . Thus $\mathbb{C}_\iota = \mathbb{C}$ or \mathbb{C}_p . Let Δ_ι denote the open unit disc $D(0, 1, \mathbb{C}_\iota)$ and let $\Delta_\iota^* = \Delta_\iota \setminus \{0\}$.

Note that we shall make use of [Sil94, Chapter V] as our main reference for Tate curves, and it works only over finite extensions of \mathbb{Q}_p or over $\overline{\mathbb{Q}_p}$ rather than over \mathbb{C}_p . However, as noted in [Sil94, Remark V.3.1.2], [Sil94, Theorem V.3.1] remains valid over any non-archimedean complete field, in particular over \mathbb{C}_p . We will not require the surjectivity of the Tate uniformisation homomorphism.

The ι -adic Tate family of elliptic curves is defined, as in [Sil94, Theorems V.1.1 and V.3.1] and [Tat95, (13)], by

$$\begin{aligned} \tilde{\mathcal{E}}_\iota = \{([x : y : z], q) \in \mathbb{P}^2(\mathbb{C}_\iota) \times \Delta_\iota : \\ y^2z + xyz = x^3 + \tilde{a}_4^\iota(q)xz^2 + \tilde{a}_6^\iota(q)z^3\} \setminus \{([0 : 0 : 1], 0)\}, \end{aligned} \quad (3)$$

where

$$\begin{aligned} \tilde{a}_4(q) &= -5s_3(q) = -5q - 45q^2 - 140q^3 - \dots \in \mathbb{Z}[[q]], \\ \tilde{a}_6(q) &= -\frac{5s_3(q) + 7s_5(q)}{12} = -q - 23q^2 - 154q^3 - \dots \in \mathbb{Z}[[q]], \\ s_k(q) &= \sum_{n=1}^{\infty} \sigma_k(n)q^n \in \mathbb{Z}[[q]], \\ \sigma_k(n) &= \sum_{d|n, d>0} d^k. \end{aligned}$$

For each $q \in \Delta_\iota^*$, the fibre $\tilde{\mathcal{E}}_{\iota, q}$ is an elliptic curve over \mathbb{C}_ι . The j -invariant of $\tilde{\mathcal{E}}_{\iota, q}$ is given by the ι -adic evaluation of the q -expansion of the standard j -function:

$$j(q) = q^{-1} + 744 + 196884q + \dots \in \mathbb{Q}[[q]][1/q].$$

Similarly to Section 3.A, the fibre $\tilde{\mathcal{E}}_{\iota, 0}$ is the smooth locus of a nodal cubic, so the Weierstrass group law makes $\tilde{\mathcal{E}}_{\iota, 0}$ into a \mathbb{C}_ι -Lie group isomorphic to $\mathbb{G}_m(\mathbb{C}_\iota)$. (In this paper, we will not make use of non-archimedean analytic geometry so the remark that $\tilde{\mathcal{E}}_{\iota, 0}$ is a \mathbb{C}_ι -Lie group may be safely ignored.)

Again by [Sil94, Theorems V.1.1, V.3.1] or by [Tat95, Theorem 1], for each $q \in \Delta_\iota^*$, there is a (surjective homomorphism) $\phi_{\iota, q}: \mathbb{C}_\iota^* \rightarrow \tilde{\mathcal{E}}_{\iota, q}$ with kernel $q^{\mathbb{Z}}$. The morphism $\phi_{\iota, q}$ is defined by explicit series and we can verify that, if we substitute $q = 0$ into these series, we obtain an isomorphism $\phi_{\iota, 0}: \mathbb{C}_\iota^* \rightarrow \tilde{\mathcal{E}}_{\iota, 0}$.

Define an invariant differential form on $\tilde{\mathcal{E}}_\iota/\Delta_\iota$ by

$$\tilde{\omega}_\iota = \frac{dx}{2y+x} \in \Omega^1(\tilde{\mathcal{E}}_\iota/\Delta_\iota).$$

Calculation, or [Tat95, p. 15], shows that, if u denotes the coordinate on \mathbb{C}_ι^* , then

$$\phi_{\iota,q}^*(\tilde{\omega}_{\iota,q}) = \frac{du}{u}. \quad (4)$$

3.C. Map from the Tate family to the “ $1/j$ ” family. For each $q \in \Delta_\iota^*$ such that $j^\iota(q) \in S^\iota(\mathbb{C}_\iota)$, the elliptic curves $\tilde{\mathcal{E}}_{\iota,q}$ and $\mathcal{E}_{1/j^\iota(q)}^\iota$ have the same j -invariant, so they are isomorphic over \mathbb{C}_ι . In fact, these isomorphisms come from an analytic map of the family of elliptic curves, as we shall now show.

Let $\Delta'_\iota = \Delta_\iota$ if ι is non-archimedean and $D(e^{-2\pi}, 0, \mathbb{C}_\iota)$ if \mathbb{C}_ι is archimedean.

Lemma 3.1. *There exists a power series $\alpha \in \mathbb{Z}[[X]]$ and, for each embedding ι of \mathbb{Q} , a morphism of \mathbb{C}_ι -manifolds $\psi_\iota: \tilde{\mathcal{E}}_\iota|_{\Delta'_\iota} \rightarrow \mathcal{E}^\iota(\mathbb{C}_\iota)$, with the following properties:*

(i) *The following diagram commutes:*

$$\begin{array}{ccc} \tilde{\mathcal{E}}_\iota|_{\Delta'_\iota} & \xrightarrow{\psi_\iota} & \mathcal{E}^\iota(\mathbb{C}_\iota) \\ \downarrow & & \downarrow \pi^\iota \\ \Delta'_\iota & \xrightarrow{1/j^\iota} & S(\mathbb{C}_\iota) \end{array}$$

(ii) *For each embedding ι of \mathbb{Q} and each $q \in \Delta'_\iota$, ψ_ι restricts to an isomorphism of \mathbb{C}_ι -algebraic groups $\psi_{\iota,q}: \tilde{\mathcal{E}}_{\iota,q} \rightarrow \mathcal{E}_{1/j^\iota(q)}^\iota$.*

(iii) *For each embedding ι of \mathbb{Q} , α^ι converges on Δ'_ι and, for each $q \in \Delta'_\iota$,*

$$\psi_{\iota,q}^*(\omega_{1/j^\iota(q)}) = \alpha^\iota(q)\tilde{\omega}_{\iota,q}.$$

Proof. Note first that j^ι does not take values 0 or 1728 on Δ'_ι . Hence $1/j^\iota$ maps Δ'_ι into $S(\mathbb{C}_\iota)$ and the bottom row of the diagram in the lemma is valid.

The q -expansions of normalized Eisenstein series $E_4(q) = 1 + 240s_3(q)$ and $E_6(q) = 1 - 504s_5(q)$ each have constant coefficient 1 and all their other coefficients are in $4\mathbb{Z}$. Hence, the same is true of $E_6/E_4 \in \mathbb{Q}[[q]]$ and we can write

$$E_6/E_4 = 1 + 4h,$$

where $h(q) \in \mathbb{Z}[[q]]$ has no constant term.

The power series

$$(1 + 4X)^{1/2} = 1 + \sum_{n=1}^{\infty} (-1)^{n+1} \frac{2}{n} \binom{2n-2}{n-1} X^n$$

has coefficients in \mathbb{Z} because $\frac{1}{n} \binom{2n-2}{n-1}$ is a Catalan number. Hence,

$$\alpha(q) := (1 + 4X)^{1/2} \circ h = 1 + 372q + 127692q^2 + \dots$$

has coefficients in \mathbb{Z} . Furthermore, $\alpha^2 = E_6/E_4$ in the ring $\mathbb{Q}[[q]]$.

Since $\alpha \in \mathbb{Z}[X]$, for every non-archimedean embedding ι of \mathbb{Q} , α^ι converges on Δ_ι . For the archimedean embedding ι of \mathbb{Q} , recall that the modular forms $E_4^\iota(e^{2\pi i\tau})$, $E_6^\iota(e^{2\pi i\tau})$ are holomorphic on \mathcal{H} and that they have no zeros on \mathcal{H} with imaginary part greater than 1. Consequently, $E_6^\iota(e^{2\pi i\tau})/E_4^\iota(e^{2\pi i\tau})$ is holomorphic and non-zero on $\mathcal{S} = \{\tau \in \mathcal{H} : \text{Im}(\tau) > 1\}$, so E_6^ι/E_4^ι is holomorphic and non-zero on the image of \mathcal{S} under $e^{2\pi i-}$, namely Δ'_ι . It follows that E_6^ι/E_4^ι has a holomorphic square root on this disc. If we choose the square root of E_6^ι/E_4^ι with the correct sign, its Taylor series will be α^ι , so α^ι converges on Δ'_ι .

Next we construct the morphism ψ_ι , restricted to $\tilde{\mathcal{E}}_\iota|_{\Delta_\iota^*}$. For each embedding ι of \mathbb{Q} and each $q \in \Delta_\iota^*$, since $\tilde{\mathcal{E}}_{\iota,q}$ and $\mathcal{E}_{1/j^\iota(q)}^\iota$ have the same j -invariant, there is an isomorphism of elliptic curves $\psi_{\iota,q}: \tilde{\mathcal{E}}_{\iota,q} \rightarrow \mathcal{E}_{1/j^\iota(q)}^\iota$. By [Sil86a, Prop. III.3.1(b)], this isomorphism must have the form

$$\psi_{\iota,q}(\tilde{x}, \tilde{y}) = (u(q)^2\tilde{x} + r(q), u(q)^3\tilde{y} + u(q)^2s(q)\tilde{x} + t(q)) \quad (5)$$

for some $u(q) \in \mathbb{C}_\iota^\times$ and $r(q), s(q), t(q) \in \mathbb{C}_\iota$.

The short Weierstrass coefficients for $\mathcal{E}_{1/j^\iota(q)}^\iota$ and for $\tilde{\mathcal{E}}_{\iota,q}$ are

$$\begin{aligned} c_4(1/j^\iota(q)) &= \frac{j^\iota(q)}{j^\iota(q) - 1728}, & \tilde{c}_4(q) &= E_4^\iota(q), \\ c_6(1/j^\iota(q)) &= -\frac{j^\iota(q)}{j^\iota(q) - 1728}, & \tilde{c}_6(q) &= -E_6^\iota(q). \end{aligned}$$

From [Sil86a, Table 3.1], these short Weierstrass coefficients are related by

$$u(q)^4\tilde{c}_4(q) = c_4(1/j^\iota(q)), \quad u(q)^6\tilde{c}_6(q) = c_6(1/j^\iota(q)).$$

Therefore,

$$u(q)^2 = \frac{\tilde{c}_4(q)c_6(1/j^\iota(q))}{\tilde{c}_6(q)c_4(1/j^\iota(q))} = \frac{E_4^\iota(q)}{E_6^\iota(q)} = \alpha^\iota(q)^{-2}.$$

(Note that, as discussed above, j^ι does not take the values 0 or 1728 on Δ'_ι , so $c_4(1/j^\iota(q))$ and $c_6(1/j^\iota(q))$ are finite and non-zero on Δ'_ι for any embedding ι . Since u is non-zero on Δ'_ι , it follows that $\tilde{c}_4(q)$ and $\tilde{c}_6(q)$ are non-zero on Δ'_ι .)

Thus $u(q) = \pm\alpha^\iota(q)^{-1}$. If $u(q) = -\alpha^\iota(q)^{-1}$, then we can replace the isomorphism ψ_q by $[-1] \circ \psi_q$. This replacement multiplies u by -1 . Thus we obtain $u(q) = \alpha^\iota(q)^{-1}$ for all $q \in \Delta_\iota^*$.

We can calculate $r(q), s(q), t(q)$ using the first three lines of [Sil86a, Table 3.1] and the following coefficients of the long Weierstrass equations for \mathcal{E} and $\tilde{\mathcal{E}}_\iota$: $a_1 = \tilde{a}_1 = 1$ and $a_2 = \tilde{a}_2 = a_3 = \tilde{a}_3 = 0$. We obtain that $r(q), s(q), t(q)$ are polynomials in $\alpha^\iota(q)^{-1}$, thus, they are ι -adic analytic functions on Δ_ι^* . Thus, (5) not only defines fibre-wise isomorphisms $\psi_{\iota,q}$, but these fit together into an analytic map $\psi_\iota: \tilde{\mathcal{E}}_\iota|_{\Delta_\iota^*} \rightarrow \mathcal{E}^\iota|_{S^*(\mathbb{C}_\iota)}$.

Again from [Sil86a, Table 3.1], we read off that

$$u(q)^{-1}\tilde{\omega}_{\iota,q} = \psi_{\iota,q}^*(\omega_{1/j^\iota(q)}^{\iota}),$$

proving (iii) for all $q \neq 0$.

In order to extend ψ_ι to a morphism $\tilde{\mathcal{E}}_\iota|_{\Delta'_\iota} \rightarrow \mathcal{E}^\iota$, we define $\psi_{\iota,0}$ by the formula (5). We have $u(0) = \alpha(0) = 1$ and can calculate $r(0) = s(0) = t(0) = 0$. Thus $\psi_{\iota,0}$ is the identity morphism. Since $\tilde{\mathcal{E}}_{\iota,0} = \mathcal{E}_0^\iota$, we see that $\psi_{\iota,0}$ is indeed an isomorphism $\tilde{\mathcal{E}}_{\iota,0} \rightarrow \mathcal{E}_0^\iota$. It satisfies

$$\psi_{\iota,0}^*(\omega_0) = \tilde{\omega}_{\iota,0} = \alpha^\iota(0)\tilde{\omega}_{\iota,0}. \quad \square$$

3.D. Periods. In order to define a power series F which will play a central role throughout the paper, we modify α from Lemma 3.1 so that its ι -adic evaluations become functions of $s = 1/j^\iota(q) \in S^\iota(\mathbb{C}_\iota)$ instead of functions of $q \in \Delta'_\iota$. To accomplish this, note first that $1/j$ has no constant term and its coefficient of degree 1 is a unit in \mathbb{Z} . Hence $1/j$ has a compositional inverse in $\mathbb{Z}[[X]]$, which we denote by

$$\theta = q + 744q^2 + 750420q^3 + \cdots \in \mathbb{Z}[[X]].$$

Define

$$F = \alpha \circ \theta \in \mathbb{Z}[[X]].$$

Since $\theta, \alpha \in \mathbb{Z}[[X]]$, by Lemma 2.2, for all non-archimedean embeddings ι of \mathbb{Q} , $R(F^\iota) \geq 1$ and for all $s \in \Delta_\iota$,

$$F^\iota(s) = \alpha^\iota(\theta^\iota(s)). \quad (6)$$

Note that $\Delta_\iota \subset S^\iota(\mathbb{C}_\iota)$ since $1/1728$ is a p -adic integer for every prime p .

For the archimedean embedding ι of \mathbb{Q} , let Δ_S denote an open disc in \mathbb{C} centred at 0 with the following properties:

- (1) the radius of Δ_S is at most $1/1728$ (so $\Delta_S \subset S^\iota(\mathbb{C})$);
- (2) θ^ι converges on Δ_S ;
- (3) $\theta^\iota(\Delta_S) \subset D(0, e^{-2\pi i}, \mathbb{C})$.

Then the disc of convergence of F^ι contains Δ_S and (6) holds when ι is the archimedean embedding for all $s \in \Delta_S$.

We can interpret the evaluation of F at the archimedean embedding $\iota: \mathbb{Q} \rightarrow \mathbb{C}$ as giving periods of elliptic curves over \mathbb{C} . Indeed, let $\tilde{\gamma}$ be the generator of $H_1(\mathbb{C}^\times, \mathbb{Z}) \cong \mathbb{Z}$ such that $\int_{\tilde{\gamma}} du/u = 2\pi i$ (this fixes an orientation of \mathbb{C}). Then, for $s \in \Delta_S$ and $q = \theta^\iota(s) \in \Delta'_\iota$,

$$\gamma_s := \psi_{\iota,q^*} \phi_{\iota,q^*}(\tilde{\gamma}) \in H_1(\mathcal{E}_s^\iota(\mathbb{C}), \mathbb{Z})$$

are the values of a section $\gamma \in (R_1\pi_*\mathbb{Q})(\Delta_S)$. In particular, for $s \in \Delta_S^*$, γ_s is locally invariant (that is, invariant under the monodromy action of $\pi_1(\Delta_S^*, s)$). Using

Lemma 3.1(iii), (4) and (6), we have

$$\frac{1}{2\pi i} \int_{\gamma_s} \omega_s^\iota = \frac{1}{2\pi i} \int_{\tilde{\gamma}} \phi_{\iota,q}^* \psi_{\iota,q}^* (\omega_s^\iota) = \frac{1}{2\pi i} \int_{\tilde{\gamma}} \alpha^\iota(q) \frac{du}{u} = \alpha^\iota(q) = F^\iota(s). \quad (7)$$

Now we define three additional functions which, together with F , make up the full period matrix of the family \mathcal{E} under the archimedean embedding $\iota: \mathbb{Q} \rightarrow \mathbb{C}$. The first of these, G , is the archimedean evaluation of a power series with coefficients in \mathbb{Q} ; this matters because we need G to be a G-function, but we will not make use of non-archimedean evaluations of G . The other two functions, F^* and G^* , are defined only as holomorphic functions on a subset of Δ_S^* and are not evaluations of power series or even Laurent series at all (they are not meromorphic at 0).

Let $\eta \in H_{DR}^1(\mathcal{E}|_{S^*/S^*})$ denote the de Rham cohomology class which, on each fibre \mathcal{E}_s , is represented by the differential form of the second kind

$$(x + \frac{1}{12}) \frac{dx}{2y + x}.$$

For each $s \in S^*(K)$ (where K is any field of characteristic 0), the de Rham classes $[\omega_s]$ and $[\eta_s]$ form a symplectic basis for $H_{DR}^1(\mathcal{E}_s/K)$ with respect to the residue pairing. Define a holomorphic function $G^\iota: \Delta_S \rightarrow \mathbb{C}$ by

$$G^\iota(s) = \frac{1}{2\pi i} \int_{\gamma_s} \eta_s^\iota. \quad (8)$$

Lemma 3.2. G^ι is the complex evaluation of a power series $G \in \mathbb{Q}[[X]]$.

Proof. Let ∇ denote the Gauss–Manin connection on $H_{DR}^1(\mathcal{E}|_{S^*/S^*})$. Then ω and $\nabla_{d/ds}(\omega)$ form a $\mathbb{Q}(S)$ -basis for $H_{DR}^1(\mathcal{E}_{\mathbb{Q}(S)}/\mathbb{Q}(S))$. So $\eta = a\omega + b\nabla_{d/ds}(\omega)$ for some $a, b \in \mathbb{Q}(S)$. It follows that G^ι is the ι -adic evaluation of

$$G = aF + b \frac{dF}{dX} \in \mathbb{Q}[[X]]. \quad \square$$

Let Δ_S^\dagger be a simply connected open subset of $\Delta_S \setminus \{0\}$ which contains $S^*(\overline{\mathbb{Q}}) \cap \Delta_S$. For example, we can choose Δ_S^\dagger to be the complement in $\Delta_S \setminus \{0\}$ of a ray from 0 which misses all $\overline{\mathbb{Q}}$ -points.

Since Δ_S^\dagger is simply connected, the local system $R_1\pi_*\mathbb{Q}|_{\Delta_S^\dagger}$ is trivial. This local system has rank 2, so we can choose a section δ of $R_1\pi_*\mathbb{Q}|_{\Delta_S^\dagger}$ such that γ and δ form an oriented basis for $(R_1\pi_*\mathbb{Q})(\Delta_S^\dagger)$. (“Oriented” means that $\gamma \cdot \delta = +1$, where \cdot denotes the intersection pairing on the fibres.) Define

$$F^*(s) = \frac{1}{2\pi i} \int_{\delta_s} \omega_s^\iota, \quad G^*(s) = \frac{1}{2\pi i} \int_{\delta_s} \eta_s^\iota.$$

These are holomorphic functions on Δ_S^\dagger , but they do not extend to holomorphic functions on Δ_S^* because δ has non-trivial monodromy around 0.

By the Legendre period relation for elliptic curves, for every $s \in \Delta_S^\dagger$,

$$\det \begin{pmatrix} F^\iota(s) & F^*(s) \\ G^\iota(s) & G^*(s) \end{pmatrix} = \frac{1}{(2\pi i)^2} \det \begin{pmatrix} \int_{\gamma_s} \omega_s^\iota & \int_{\delta_s} \omega_s^\iota \\ \int_{\gamma_s} \eta_s^\iota & \int_{\delta_s} \eta_s^\iota \end{pmatrix} = \frac{1}{2\pi i}. \quad (9)$$

Remark 3.3. This will not be used in this paper, but we can also interpret the p -adic evaluation of F in terms of the p -adic period pairing

$$\langle -, - \rangle : T_p(G) \times H_{DR}^1(G/\mathbb{C}_p) \rightarrow B_{DR},$$

defined for G a commutative algebraic group over \mathbb{C}_p . Indeed, let ι denote the embedding $\mathbb{Q} \rightarrow \mathbb{C}_p$. Choose a generator $\tilde{\gamma}_\iota$ for $T_p(\mathbb{G}_m) = \mathbb{Z}_p(1)$ (a “ p -adic orientation”). Then the “ p -adic $2\pi i$ ”

$$t_p := \langle \tilde{\gamma}_\iota, du/u \rangle$$

is a uniformiser of B_{DR}^+ , and $\psi_{\iota, q^*} \phi_{\iota, q^*}(\tilde{\gamma}_\iota)$ is a locally invariant p -adic cycle on $\mathcal{E}^\iota(\mathbb{C}_\iota)|_{\Delta_\iota}$. The same calculation as (7) shows that, for all $s \in \Delta_\iota$ with $q = \theta^\iota(s)$,

$$\frac{1}{t_p} \langle \psi_{\iota, q^*} \phi_{\iota, q^*}(\tilde{\gamma}_\iota), \omega_s^\iota \rangle = \frac{1}{t_p} \langle \tilde{\gamma}_\iota, \psi_{\iota, q}^* \phi_{\iota, q}^*(\omega_s^\iota) \rangle = \frac{1}{t_p} \langle \tilde{\gamma}_\iota, F^\iota(s) du/u \rangle = F^\iota(s).$$

4. PERIOD RELATIONS FOR ISOGENOUS POINTS

In this section, we show how isogenies between elliptic curves over a number field \hat{K} lead to polynomial relations between corresponding evaluations of F and G . We construct these relations separately for archimedean and non-archimedean embeddings of \hat{K} . The conclusion can be summed up as follows.

Theorem 4.1. *Let \hat{K} be a number field. Let M, N be positive integers. Let $s_1, s_2, s_3, s_4 \in S^*(\hat{K})$ be points such that:*

- (i) *there is an isogeny $\mathcal{E}_{s_1} \rightarrow \mathcal{E}_{s_2}$ of degree M defined over \hat{K} ;*
- (ii) *there is an isogeny $\mathcal{E}_{s_3} \rightarrow \mathcal{E}_{s_4}$ of degree N defined over \hat{K} .*

Then:

- (1) *There exists a non-zero polynomial $P_\infty \in \hat{K}[Y_1, Z_1, \dots, Y_4, Z_4]$ of degree at most $2[\hat{K} : \mathbb{Q}]$ such that, for every archimedean embedding ι of \hat{K} satisfying $s_i^\iota \in \Delta_S$ for all $i = 1, 2, 3, 4$, we have*

$$P_\infty^\iota(F^\iota(s_1^\iota), G^\iota(s_1^\iota), \dots, F^\iota(s_4^\iota), G^\iota(s_4^\iota)) = 0.$$

Furthermore, P_∞ is not in the ideal $\langle Y_1 - Y_3, Z_1 - Z_3 \rangle$.

- (2) *There exists a non-zero polynomial $P_{fin} \in \hat{K}[Y_1, Y_2]$ of degree at most $2d(M)$ such that, for every non-archimedean embedding ι of \hat{K} satisfying $|s_i^\iota| < 1$ for $i = 1, 2$, we have*

$$P_{fin}^\iota(F^\iota(s_1^\iota), F^\iota(s_2^\iota)) = 0.$$

Proof. (1) Let P_∞ be the product of the polynomials P_ι given by Proposition 4.4, as ι runs through the archimedean embeddings ι of \hat{K} .

(2) Let P_{fin} be the product of the polynomials P_m from Proposition 4.7 (applied to s_1 and s_2), as m runs through the divisors (positive and negative) of M . \square

Remark 4.2. For each archimedean embedding ι , $s^\iota \in \Delta_S$ implies that $|s^\iota| < \min(R(F^\iota), R(G^\iota))$. Hence the evaluations in Theorem 4.1(1) make sense.

Note that the relations at non-archimedean embeddings involve only F , not G . This enables us to construct the relation without knowing a geometric interpretation of the p -adic evaluations of G . It also saves us the effort of estimating $R(G^\iota)$ which would otherwise be needed to control the set on which Theorem 4.1(2) makes sense.

The non-archimedean relations involve only s_1, s_2 , not s_3, s_4 . This is not important, except that it avoids the need to assert that $P_{fin} \notin \langle Y_1 - Y_3, Z_1 - Z_3 \rangle$.

We may need to apply Theorem 4.1 with $s_1 = s_3$. The assertion that $P_\infty \notin \langle Y_1 - Y_3, Z_1 - Z_3 \rangle$ in Theorem 4.1 is needed to ensure that P_∞ does not collapse to 0 upon making the substitution $Y_1 = Y_3, Z_1 = Z_3$.

Remark 4.3. The polynomial P_∞ is obtained by finding a relation for each archimedean embedding of \hat{K} (Proposition 4.4), then multiplying these together. There are $[\hat{K} : \mathbb{Q}]$ archimedean embeddings, leading to a bound for $\deg(P_\infty)$.

The polynomial P_{fin} is similarly constructed as a product, where one of the factors gives a relation between evaluations of F at each non-archimedean embedding of \hat{K} (Proposition 4.7). However, there are infinitely many non-archimedean embeddings, so it is important that the relations at non-archimedean embeddings are not independent of each other: instead Proposition 4.7 gives a finite collection of polynomials (of controlled size) whose members give relations between evaluations of F at every non-archimedean embedding where s_1^ι and s_2^ι are small enough.

4.A. Complex period relations. To obtain a period relation at an archimedean embedding of \hat{K} , we require two isogenies $\mathcal{E}_{s_1} \rightarrow \mathcal{E}_{s_2}$ and $\mathcal{E}_{s_3} \rightarrow \mathcal{E}_{s_4}$. A single isogeny gives a relation whose coefficients involve $2\pi i$, not just algebraic numbers. A second isogeny gives a second such relation, allowing us to eliminate $2\pi i$.

Proposition 4.4. *Let \hat{K} be a number field. Let M, N be positive integers. Let $s_1, s_2, s_3, s_4 \in S^*(\hat{K})$ with the following properties:*

- (i) *there is an isogeny $\mathcal{E}_{s_1} \rightarrow \mathcal{E}_{s_2}$ of degree M defined over \hat{K} ;*
- (ii) *there is an isogeny $\mathcal{E}_{s_3} \rightarrow \mathcal{E}_{s_4}$ of degree N defined over \hat{K} .*

Then, for each archimedean embedding ι of \hat{K} satisfying $s_i^\iota \in \Delta_S$ for all $i = 1, 2, 3, 4$, there exists a non-zero polynomial $P_\iota \in \hat{K}[Y_1, Z_1, \dots, Y_4, Z_4]$ of degree 2 such that

$$P_\iota(F^\iota(s_1^\iota), G^\iota(s_1^\iota), \dots, F^\iota(s_4^\iota), G^\iota(s_4^\iota)) = 0.$$

Furthermore, P_ι is not in the ideal $\langle Y_1 - Y_3, Z_1 - Z_3 \rangle$.

Proof. Let ι be an embedding $\hat{K} \rightarrow \mathbb{C}$ satisfying $s_i^t \in \Delta_S$ for all $i = 1, 2, 3, 4$. To reduce notation, we omit mention of ι for the rest of this proof.

We consider first the pair s_1, s_2 , obtaining an inhomogeneous relation of degree at most 2 between the ι -adic evaluations of $F(s_1), G(s_1), F(s_2), G(s_2)$ whose value is a rational multiple of $1/2\pi i$.

Let $f: \mathcal{E}_{s_1} \rightarrow \mathcal{E}_{s_2}$ be an isogeny of degree M . Note that $f^*(\omega_{s_2}) \in \Omega^1(\mathcal{E}_{s_1}/\hat{K})$, which is a 1-dimensional \hat{K} -vector space, so there is an element $a \in \hat{K}$ such that

$$f^*(\omega_{s_2}) = a\omega_{s_1}. \quad (10)$$

There are also $b, d \in \hat{K}$ such that

$$[f^*(\eta_{s_2})] = b[\omega_{s_1}] + d[\eta_{s_1}]$$

where $[\cdot]$ denotes the class of a differential form in $H_{DR}^1(\mathcal{E}_{s_1}/\hat{K})$. Since $[\omega_s], [\eta_s]$ form a symplectic basis of $H_{DR}^1(\mathcal{E}_s/\hat{K})$ for each s , we have

$$ad = \det \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} = M.$$

Looking at homology, there are $p, q, r, s \in \mathbb{Z}$ such that

$$\begin{aligned} f_*(\gamma_{s_1}) &= p\gamma_{s_2} + r\delta_{s_2}, \\ f_*(\delta_{s_1}) &= q\gamma_{s_2} + s\delta_{s_2}. \end{aligned}$$

Here $\det \begin{pmatrix} p & q \\ r & s \end{pmatrix} = M$.

Since $s_1, s_2, s_3, s_4 \in \Delta_S$, we can apply (7) and (8): the evaluations of power series $F(s_1), G(s_1)$ are equal to periods of \mathcal{E}_{s_1} , and so on for s_2, s_3, s_4 . Hence, using equations such as

$$\int_{\gamma_{s_1}} f^*(\omega_{s_2}) = \int_{f_*(\gamma_{s_1})} \omega_{s_2},$$

we obtain the period relations

$$\begin{pmatrix} a & 0 \\ b & d \end{pmatrix} \begin{pmatrix} F(s_1) & F^*(s_1) \\ G(s_1) & G^*(s_1) \end{pmatrix} = \begin{pmatrix} F(s_2) & F^*(s_2) \\ G(s_2) & G^*(s_2) \end{pmatrix} \begin{pmatrix} p & q \\ r & s \end{pmatrix}. \quad (11)$$

Multiplying on the left by the row vector $(-G(s_2), F(s_2))$ and using (9), we obtain

$$\begin{aligned} & (-aG(s_2) + bF(s_2), dF(s_2)) \begin{pmatrix} F(s_1) & F^*(s_1) \\ G(s_1) & G^*(s_1) \end{pmatrix} \\ &= (-G(s_2)F(s_2) + F(s_2)G(s_2), -G(s_2)F^*(s_2) + F(s_2)G^*(s_2)) \begin{pmatrix} p & q \\ r & s \end{pmatrix} \\ &= (0, 1/(2\pi i)) \begin{pmatrix} p & q \\ r & s \end{pmatrix}. \end{aligned}$$

Multiplying on the right by the column vector $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$, we obtain

$$-aG(s_2)F(s_1) + bF(s_2)F(s_1) + dF(s_2)G(s_1) = \frac{r}{2\pi i}. \quad (12)$$

If $r = 0$, then (12) is a relation of degree 2 with coefficients in \hat{K} between the ι -adic evaluations of power series $F(s_1)$, $F(s_2)$, $G(s_1)$, $G(s_2)$. Note that this relation is non-zero because $ad = M \neq 0$.

If $r \neq 0$, then the $2\pi i$ on the right hand side means that (12) does not directly give us a relation between evaluations of F and G . Instead, we apply the same argument as above to s_3 and s_4 , obtaining a relation

$$-a'G(s_4)F(s_3) + b'F(s_4)F(s_3) + d'F(s_4)G(s_3) = \frac{r'}{2\pi i} \quad (13)$$

where $a', b', d' \in \hat{K}$ and $r' \in \mathbb{Z}$. Combining (12) and (13), we obtain

$$\begin{aligned} r'(-aG(s_2)F(s_1) + bF(s_2)F(s_1) + dF(s_2)G(s_1)) \\ + r(a'G(s_4)F(s_3) - b'F(s_4)F(s_3) - d'F(s_4)G(s_3)) = 0. \end{aligned} \quad (14)$$

In other words, the following polynomial in $\hat{K}[Y_1, Z_1, \dots, Y_4, Z_4]$ vanishes at $(F(s_1), G(s_1), \dots, F(s_4), G(s_4))$:

$$P_i = r'(-aZ_2Y_1 + bY_2Y_1 + dY_2Z_1) + r(a'Z_4Y_3 - b'Y_4Y_3 - d'Y_4Z_3).$$

Each of the terms in the above expression involves a distinct monomial. Since we are assuming that $r \neq 0$ and $a'd' = N \neq 0$, the term $ra'Z_4Y_3$ is non-zero. Hence, this is a non-zero polynomial.

Furthermore, the monomials in the above expression remain distinct after the substitution $Y_1 = Y_3$, $Z_1 = Z_3$, so $P_i \notin \langle Y_1 - Y_3, Z_1 - Z_3 \rangle$. \square

Remark 4.5. If $r = 0$, then we can obtain a relation of degree 1 directly from the top left corner of (11), namely, $aF(s_1) = pF(s_2)$. Thus $r = 0$ corresponds to Case 2 in [And89, 2.4], while $r \neq 0$ corresponds to Case 3, leading to a quadratic relation and requiring two inhomogeneous relations to eliminate a multiple of $1/2\pi i$.

4.B. Non-archimedean period relations. In order to prove Theorem 4.1(2), we shall use the following result about isogenies and the Tate uniformisation.

Proposition 4.6. *Let p be a prime number and let ι be the embedding $\mathbb{Q} \rightarrow \mathbb{C}_p$. Let $q_1, q_2 \in \Delta_\iota$ be such that there is an isogeny $\tilde{f}: \tilde{\mathcal{E}}_{\iota, q_1} \rightarrow \tilde{\mathcal{E}}_{\iota, q_2}$ of degree M . There is an integer m dividing M such that the following diagram commutes:*

$$\begin{array}{ccc} \mathbb{G}_m(\overline{\mathbb{Q}}_p) & \xrightarrow{[m]} & \mathbb{G}_m(\overline{\mathbb{Q}}_p) \\ \downarrow \phi_{\iota, q_1} & & \downarrow \phi_{\iota, q_2} \\ \tilde{\mathcal{E}}_{\iota, q_1} & \xrightarrow{\tilde{f}} & \tilde{\mathcal{E}}_{\iota, q_2} \end{array}$$

Proof. According to [Tat95, “Isogenies”, Theorem], every isogeny between p -adic Tate curves is of the form $\alpha_{m,n}: \tilde{\mathcal{E}}_{\iota,q_1} \rightarrow \tilde{\mathcal{E}}_{\iota,q_2}$ for some integers m and n such that $q_1^m = q_2^n$, where $\alpha_{n,m}$ fits into the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{1 \mapsto q_1} & \overline{\mathbb{Q}}_p^\times & \xrightarrow{\phi_{\iota,q_1}} & \tilde{\mathcal{E}}_{\iota,q_1} & \longrightarrow & 0 \\ & & \downarrow [n] & & \downarrow [m] & & \downarrow \alpha_{n,m} & & \\ 0 & \longrightarrow & \mathbb{Z} & \xrightarrow{1 \mapsto q_2} & \overline{\mathbb{Q}}_p^\times & \xrightarrow{\phi_{\iota,q_2}} & \tilde{\mathcal{E}}_{\iota,q_2} & \longrightarrow & 0 \end{array}$$

(note that our m and n are the opposite way round to those in [Tat95]). Furthermore, the theorem in [Tat95] also tells us that $mn = M$ so m divides M . \square

Proposition 4.7. *Let \hat{K} be a number field. Let $s_1, s_2 \in S^*(\hat{K})$ be such that there exists an isogeny $\mathcal{E}_{s_1} \rightarrow \mathcal{E}_{s_2}$ of degree M defined over \hat{K} . Then there exists a set of non-zero linear polynomials $P_m \in \hat{K}[Y_1, Y_2]$, indexed by the positive and negative integers m dividing M , such that, for every non-archimedean embedding ι of \hat{K} satisfying $|s_i^\iota| < 1$ for $i = 1, 2$, there is some divisor m of M for which*

$$P_m^\iota(F^\iota(s_1^\iota), F^\iota(s_2^\iota)) = 0.$$

Proof. Let $f: \mathcal{E}_{s_1} \rightarrow \mathcal{E}_{s_2}$ be an isogeny of degree M . As in (10), let $a \in \hat{K}$ be the scalar such that $f^*(\omega_{s_2}) = a\omega_{s_1}$. Note that a is independent of the embedding ι .

For each divisor m of M in \mathbb{Z} , let

$$P_m(Y_1, Y_2) = aY_1 - mY_2 \in \hat{K}[Y_1, Y_2]. \quad (15)$$

These are the polynomials we need for the conclusion of Proposition 4.7.

Let p be a rational prime and let $\iota: \hat{K} \rightarrow \overline{\mathbb{Q}}_p$ be an embedding such that $s_1^\iota, s_2^\iota \in \Delta_\iota$. Let $q_1 = \theta^\iota(s_1), q_2 = \theta^\iota(s_2) \in \Delta_\iota$.

Since $\psi_{\iota,q}$ is an isomorphism of elliptic curves, there is a unique isogeny $\tilde{f}_\iota: \tilde{\mathcal{E}}_{\iota,q_1} \rightarrow \tilde{\mathcal{E}}_{\iota,q_2}$ of elliptic curves over $\overline{\mathbb{Q}}_p$ such that the lower square of the following diagram commutes. Applying Proposition 4.6 to \tilde{f}_ι , we obtain an integer m dividing M such that the upper square of the diagram commutes.

$$\begin{array}{ccc} \mathbb{G}_m(\overline{\mathbb{Q}}_p) & \xrightarrow{\dots \dots [m] \dots \dots} & \mathbb{G}_m(\overline{\mathbb{Q}}_p) \\ \downarrow \phi_{\iota,q_1} & & \downarrow \phi_{\iota,q_2} \\ \tilde{\mathcal{E}}_{\iota,q_1} & \xrightarrow{\dots \dots \tilde{f}_\iota \dots \dots} & \tilde{\mathcal{E}}_{\iota,q_2} \\ \downarrow \psi_{\iota,q_1} & & \downarrow \psi_{\iota,q_2} \\ \mathcal{E}_{s_1}(\overline{\mathbb{Q}}_p) & \xrightarrow{f} & \mathcal{E}_{s_2}(\overline{\mathbb{Q}}_p) \end{array} \quad (16)$$

Pulling back $\tilde{\omega}_{\iota,q_2}$ around the upper square of diagram (16), we obtain

$$\phi_{\iota,q_1}^* \tilde{f}_\iota^* (\tilde{\omega}_{\iota,q_2}) = [m]^* \phi_{\iota,q_2}^* (\tilde{\omega}_{\iota,q_2}) = m \frac{du}{u} = m \phi_{\iota,q_1}^* (\tilde{\omega}_{\iota,q_1}).$$

Since ϕ_{ι, q_1}^* is an isomorphism $\Omega^1(\tilde{\mathcal{E}}_{\iota, q_1}/\overline{\mathbb{Q}}_p) \rightarrow \Omega^1(\mathbb{G}_m/\overline{\mathbb{Q}}_p)$, it follows that

$$\tilde{f}_\iota^*(\tilde{\omega}_{\iota, q_2}) = m \tilde{\omega}_{\iota, q_1}.$$

Using this together with (6), (10) and the lower square of (16), we can calculate

$$\begin{aligned} \tilde{f}_\iota^* \psi_{\iota, q_2} * (\omega_{s_2}^\iota) &= \tilde{f}_\iota^* (F^\iota(s_2^\iota) \tilde{\omega}_{\iota, q_2}) = F^\iota(s_2^\iota) m \tilde{\omega}_{\iota, q_1} \\ &= \psi_{\iota, q_1}^* \tilde{f}_\iota^* (\omega_{s_2}^\iota) = \psi_{\iota, q_1}^* (a \omega_{s_1}^\iota) = a F^\iota(s_1^\iota) \tilde{\omega}_{\iota, q_1}. \end{aligned}$$

Since $\tilde{\omega}_{\iota, q_1} \neq 0$, we conclude that

$$a F^\iota(s_1^\iota) = m F^\iota(s_2^\iota). \quad (17)$$

In other words, (15) gives a relation between the ι -adic evaluations of $F(s_1)$ and $F(s_2)$, as required. \square

5. PROOF OF LOWER BOUND FOR GALOIS ORBITS

In this section we prove the height bound Theorem 1.6 and deduce the Galois orbits bound Theorem 1.5. The proof follows the strategy of [And89, Ch. X], using the polynomial relations between periods of fibres with exceptional isogenies obtained in Theorem 4.1. In particular, the work of finding relations between non-archimedean evaluations of F was already carried out in Theorem 4.1(2). Thus the only new work in this section needed to make use of non-archimedean evaluations of G-functions is to pay attention to non-archimedean convergence of power series.

In order to be confident that the argument is valid with non-archimedean evaluations, we have written the full argument in detail. This includes expanding the remark on [And89, p. 202] about finite ramification at s_0 when choosing the local parameter, via Lemma 5.1. To help the reader follow the essential structure of the argument, we have included in Section 5.A a summary for a special case in which many of the technical difficulties do not arise.

We found Papas's paper [Pap] very helpful when writing this section, especially for describing the trivial relations using André's Normal Monodromy Theorem [And92], which is simpler than the approach in [And89, X, Lemma 3.3].

5.A. A special case. In order to illustrate our strategy for the proof of Theorem 1.6, we consider a special case in which \overline{C} (the Zariski closure of C in $(\mathbb{P}^1)^n$) is isomorphic to \mathbb{P}^1 . This demonstrates the central idea of how we exploit the global relations given by Theorem 4.1, while avoiding technical complications which appear in the general case. We shall not give full details of all steps in this special case, because full details are given in the general case.

Let $s_0 = (\infty, \dots, \infty) \in \overline{C}$. Choose an isomorphism $x: \overline{C} \rightarrow \mathbb{P}^1$ such that $x(s_0) = 0$. For each $i = 1, \dots, n$, let $\rho_i: \overline{C} \rightarrow \mathbb{P}^1$ denote the reciprocal of the projection onto the i -th coordinate (note that $\rho_i(s_0) = 0$ for all i). Let ξ_i denote

the Taylor series of ρ_i in terms of the local parameter x around s_0 . Define power series $F_i, G_i \in K[[X]]$ by

$$F_i = F \circ \xi_i, \quad G_i = G \circ \xi_i.$$

For every embedding ι of K , on a suitable neighbourhood of s'_0 in $\overline{C}^\iota(\mathbb{C}_\iota)$, we have

$$F_i^\iota(x^\iota(s)) = F^\iota(\rho_i^\iota(s)),$$

and similarly for G (diagram (25)).

Restricting our attention to $C_0 = \bigcap_i \rho_i^{-1}(S) \subset C$ (throwing away only finitely many points), we may consider the semiabelian schemes $\rho_i^* \mathcal{E}$ over C_0 . We can check that, for each archimedean embedding ι of K and on a suitable disc around s'_0 in $\overline{C}^\iota(\mathbb{C})$, the ι -adic evaluations of F_i and G_i give locally invariant periods of $(\rho_i^* \mathcal{E})^\iota$ (equation (27)). It follows that F_i and G_i are G-functions (Lemma 5.6).

Furthermore, since C is Hodge generic, the semiabelian schemes $\rho_i^* \mathcal{E}$ are not geometrically generically isogenous. This implies that there are no non-zero trivial relations between the G-functions $\{F_i, G_i : 1 \leq i \leq n\}$ (Proposition 5.10).

Let $s \in C(\overline{\mathbb{Q}})$ satisfy the hypothesis of Theorem 1.6, and let $\hat{K} = K(s)$. Since $\Phi_M(s_{i_1}, s_{i_2}) = 0$, there is an isogeny $\mathcal{E}_{\rho_{i_1}(s)} \rightarrow \mathcal{E}_{\rho_{i_2}(s)}$ of degree M . Similarly, there is an isogeny $\mathcal{E}_{\rho_{i_3}(s)} \rightarrow \mathcal{E}_{\rho_{i_4}(s)}$ of degree N .

Applying Theorem 4.1, we obtain polynomials P_∞, P_{fin} such that, for all embeddings $\hat{\iota}$ of \hat{K} in which $s^{\hat{\iota}}$ is close to $s_0^{\hat{\iota}}$, we have (if $\hat{\iota}$ is archimedean)

$$\begin{aligned} 0 &= P_\infty^\iota(F^\iota(\rho_{i_1}(s)^{\hat{\iota}}), G^\iota(\rho_{i_1}(s)^{\hat{\iota}}), \dots, F^\iota(\rho_{i_4}(s)^{\hat{\iota}}), G^\iota(\rho_{i_4}(s)^{\hat{\iota}})) \\ &= P_\infty^\iota(F_{i_1}^\iota(x(s)^{\hat{\iota}}), G_{i_1}^\iota(x(s)^{\hat{\iota}}), \dots, F_{i_4}^\iota(x(s)^{\hat{\iota}}), G_{i_4}^\iota(x(s)^{\hat{\iota}})) \end{aligned}$$

and (if $\hat{\iota}$ is non-archimedean)

$$0 = P_{fin}^\iota(F^\iota(\rho_{i_1}(s)^{\hat{\iota}}), F^\iota(\rho_{i_2}(s)^{\hat{\iota}})) = P_{fin}^\iota(F_{i_1}^\iota(x(s)^{\hat{\iota}}), F_{i_2}^\iota(x(s)^{\hat{\iota}})).$$

Define a polynomial $Q \in \hat{K}[Y_1, Z_1, \dots, Y_n, Z_n]$ as follows:

$$Q(\underline{Y}, \underline{Z}) = P_\infty(Y_{i_1}, Z_{i_1}, \dots, Y_{i_4}, Z_{i_4}) \cdot P_{fin}(Y_{i_1}, Y_{i_2}).$$

Then

$$Q^\iota(F_i^\iota(x(s)^{\hat{\iota}}), G_i^\iota(x(s)^{\hat{\iota}})) : 1 \leq i \leq n = 0$$

for every embedding $\hat{\iota}$ at which $x^\iota(s^{\hat{\iota}})$ is small (since the only zero of x in \overline{C} is at s_0 , this is equivalent to $s^{\hat{\iota}}$ being close to $s_0^{\hat{\iota}}$).

By paying careful attention to the radii within which the above manipulations of power series are valid, we can conclude that Q is a global relation between the evaluations at $x(s)$ of the G-functions $F_1, G_1, \dots, F_n, G_n$, along with a rational function which we add to the set of G-functions in order to control the radii of convergence in the definition of global relation (Lemma 5.7).

By consideration of the possible coincidences between the indices i_1, i_2, i_3, i_4 , and using the facts that $P_\infty \notin \langle Y_1 - Y_3, Z_1 - Z_3 \rangle$ and $P_{fin} \neq 0$, we can ensure that $Q \neq 0$, and thus Q is not a trivial relation.

From the degree bounds in Theorem 4.1, we have $\deg(Q) \leq 2[\hat{K} : \mathbb{Q}] + 2d(M) \ll \max\{[K(s) : K], M^\epsilon\}$. We can thus complete the proof of this special case of Theorem 1.6 by applying [And89, Theorem E].

5.B. Outline of technical issues in the general case. The proof of a special case of Theorem 1.6 in Section 5.A made use of two properties of the morphism $x: \overline{C} \rightarrow \mathbb{P}^1$: the only zero of x was at $s_0 = (\infty, \dots, \infty)$ and x was a local parameter at s_0 . A morphism with these properties must have degree 1, so can only exist when \overline{C} is a rational curve.

In order to prove Theorem 1.6 when \overline{C} is not rational, we will replace \overline{C} by another curve C_4 , equipped with a surjective morphism $C_4 \rightarrow \overline{C}$ and with a finite set of points $s_1, \dots, s_\ell \in C_4$ mapping to $s_0 \in \overline{C}$. We will choose C_4 so that there is a morphism $x: C_4 \rightarrow \mathbb{P}^1$ which vanishes only at s_1, \dots, s_ℓ and which is a local parameter at each of s_1, \dots, s_ℓ . This comes as close as we can to the properties of x used in Section 5.A, at the cost of having finitely many degeneration points s_1, \dots, s_ℓ instead of the single point s_0 .

If $s \in C_4(\hat{K})$ and $x(s)^\ell$ is small for some embedding $\hat{\iota}$ of \hat{K} , then s^ℓ is close to one of the s_k^ℓ . However, even for a fixed \hat{K} -point s , s^ℓ might be close to s_k^ℓ for different k depending on the embedding $\hat{\iota}$. We therefore have to look for relations between locally invariant periods of the semiabelian schemes $(\rho_i^* \mathcal{E})^\ell$ near each of $s_1^\ell, \dots, s_\ell^\ell$ (where ρ_i denotes the reciprocal of a coordinate on \overline{C} , composed with the map $C_4 \rightarrow \overline{C}$).

Working with periods around different degeneration points is awkward, especially when it comes to understanding trivial relations between the period functions. We therefore impose an extra condition on C_4 : for every $k = 1, \dots, \ell$, C_4 has an automorphism such that $\sigma_k(s_1) = s_k$. We can then work with the semiabelian scheme $\sigma_k^* \rho_i^* \mathcal{E}$ in a neighbourhood of s_1 instead of $\rho_i^* \mathcal{E}$ in a neighbourhood of s_k .

There may exist isogenies between the semiabelian schemes $\sigma_k^* \rho_i^* \mathcal{E}$ (over a geometric generic point of the base), leading to trivial relations between the associated period functions. We therefore choose one semiabelian scheme from each isogeny class among the $\sigma_k^* \rho_i^* \mathcal{E}$. These are the semiabelian schemes which we will call \mathcal{A}_λ . Their locally invariant periods around s_1 give us a collection of G-functions with no trivial relations.

After the technical setup summarised above, we will finally be able to complete the proof by finding global relations between values of these G-functions at exceptional points along the lines explained in Section 5.A.

5.C. Curves and local parameters. As explained in Section 5.B, we need to replace \overline{C} by a curve C_4 with a suitable rational function x on C_4 which we will use as a local parameter at each of finitely many degeneration points. The following lemma constructs C_4 and x with the required properties.

Lemma 5.1. *Let C be an irreducible projective algebraic curve over a field K of characteristic zero. Let s_0 be a closed point of C . Then, after replacing K by a finite extension, there exists an irreducible smooth projective algebraic curve C_4 over K , a non-constant morphism $\nu: C_4 \rightarrow C$ and a non-constant rational function $x \in K(C_4)$ such that*

- (i) *every point $s \in C_4(\overline{K})$ where $x(s) = 0$ is a simple zero of x ;*
- (ii) *every point $s \in C_4(\overline{K})$ where $x(s) = 0$ satisfies $\nu(s) = s_0$;*
- (iii) *$x: C_4 \rightarrow \mathbb{P}^1$ is Galois, that is, $\text{Aut}_x(C_4) := \{\sigma \in \text{Aut}(C_4) : x \circ \sigma = x\}$ acts transitively on each \overline{K} -fibre of x .*

Proof. Let $\nu_1: C_1 \rightarrow C$ denote the normalisation of C . Let t be a point of C_1 such that $\nu_1(t) = s_0$. (There may be more than one such point; choose one of them.)

By the Riemann–Roch theorem, there exists a rational function on C_1 which has a pole at t and nowhere else. Taking the reciprocal of this function, we obtain a rational function $x_1 \in K(C_1)$ with a zero at t and nowhere else. Let $d \in \mathbb{N}$ denote the order of vanishing of x_1 at t .

Let C_2 denote an irreducible component of the fibre product of $x_1: C_1 \rightarrow \mathbb{P}^1$ with the d -th power map $[d]: \mathbb{P}^1 \rightarrow \mathbb{P}^1$. Let $\nu_2: C_3 \rightarrow C_2$ denote the normalisation of C_2 and let x_3 and κ be the morphisms indicated in the following commutative diagram:

$$\begin{array}{ccccccc} C_3 & \xrightarrow{\nu_2} & C_2 & \xrightarrow{\kappa} & C_1 & \xrightarrow{\nu_1} & C \\ & \searrow x_3 & \downarrow & & \downarrow x_1 & & \\ & & \mathbb{P}^1 & \xrightarrow{[d]} & \mathbb{P}^1 & & \end{array}$$

After replacing K by a finite extension, we may construct a smooth projective curve C_4 over K with a morphism $\nu_3: C_4 \rightarrow C_3$ which is the Galois closure of $x_3: C_3 \rightarrow \mathbb{P}^1$. Let ν and x be the morphisms indicated in the following diagram:

$$\begin{array}{ccccccc} & & & & \nu & & \\ & & & & \curvearrowright & & \\ C_4 & \xrightarrow{\nu_3} & C_3 & \xrightarrow{\nu_2} & C_2 & \xrightarrow{\kappa} & C_1 & \xrightarrow{\nu_1} & C \\ & \searrow x & \searrow & \downarrow & \downarrow x_1 & & & & \\ & & & \mathbb{P}^1 & \xrightarrow{[d]} & \mathbb{P}^1 & & & \end{array} \quad (18)$$

Consider a point $s \in C_4(\overline{K})$ such that $x(s) = 0$. Then $x_1 \kappa \nu_2 \nu_3(s) = [d]x(s) = 0$. Hence by the choice of x_1 , $\kappa \nu_2 \nu_3(s) = t$ so $\nu(s) = \nu_1(t) = s_0$, establishing (ii). Furthermore (iii) holds because we chose C_4 to be the Galois closure of $x_3: C_3 \rightarrow \mathbb{P}^1$.

Since t is the only zero of x_1 in \mathbb{P}^1 , and since the ramification degree of $x_1: C_1 \rightarrow \mathbb{P}^1$ at t is equal to the ramification degree of $[d]: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ at 0, by Abhyankar’s lemma [Sta, Tag 0BRM], $x_3: C_3 \rightarrow \mathbb{P}^1$ is unramified at every zero of x_3 in C_3 . (Note that the function field $K(C_3) = K(C_2)$ is a direct factor of $K(C_1) \otimes_{K(\mathbb{P}^1), [d]} K(\mathbb{P}^1)$.) Passing to the Galois closure, it follows that $x: C_4 \rightarrow \mathbb{P}^1$ is unramified at every zero of x in C_4 , proving (i). \square

In the setting of Theorem 1.6, $C \subset \mathbb{A}^n$ is a geometrically irreducible Hodge generic curve defined over a number field K that intersects infinity. Let \overline{C} denote the Zariski closure of C in $(\mathbb{P}^1)^n$ and let $s_0 = (\infty, \dots, \infty) \in \overline{C}$.

Applying Lemma 5.1 to \overline{C} , after replacing K by a finite extension, we obtain a smooth projective algebraic curve C_4 with a morphism $\nu: C_4 \rightarrow \overline{C}$ and a rational function $x \in K(C_4)$. Replace the field K by a further finite extension so that all zeros of x in C_4 are defined over K . Label the zeros of x in $C_4(K)$ as s_1, \dots, s_ℓ . Since these zeros are simple, x is a local parameter for C_4 around s_k (which is to say, x is a uniformiser in the local ring \mathcal{O}_{C_4, s_k}) for all $k = 1, \dots, \ell$.

Let $\zeta_1, \dots, \zeta_n: \overline{C} \rightarrow \mathbb{P}^1$ denote the reciprocals of the projections onto the \mathbb{P}^1 factors. For each $i = 1, \dots, n$, let $\rho_i = \zeta_i \circ \nu: C_4 \rightarrow \mathbb{P}^1$.

Remark 5.2. By Lemma 5.1(iii), for each $k = 1, \dots, \ell$, there is an automorphism $\sigma_k \in \text{Aut}_x(C_4)$ such that $\sigma_k(s_1) = s_k$.

Since x is unramified at s_1, \dots, s_ℓ , we have $\deg(x) = t$. Since x is Galois, $\#\text{Aut}_x(C_4) = \deg(x)$. It follows that $\sigma_1, \dots, \sigma_k$ are the only elements of $\text{Aut}_x(C_4)$.

5.D. Semiabelian schemes. We wish to pull back the $1/j$ -family of elliptic curves \mathcal{E} to C_4 via ρ_i . Note that \mathcal{E} is defined only over $S = \mathbb{A}^1 \setminus \{1/1728\}$, not over \mathbb{P}^1 , so we must replace C_4 by a Zariski open subset in order to be able to do this. Define

$$C_0 = \{s \in C_4 : \rho_i(\sigma_k(s)) \in S \text{ for all } k = 1, \dots, \ell, i = 1, \dots, n\}.$$

Similarly define

$$C_0^* = \{s \in C_4 : \rho_i(\sigma_k(s)) \in S^* \text{ for all } k = 1, \dots, \ell, i = 1, \dots, n\}.$$

Since the morphisms $\rho_i \circ \sigma_k$ are non-constant, each of C_0 and C_0^* is the complement of finitely many points in C_4 , so they are non-empty Zariski open subsets of C_4 . Note also that σ_k restricts to an automorphism of C_0 for each $k = 1, \dots, \ell$.

By the definition of C_0 , we have $\rho_i(C_0) \subset S$. We can therefore define $\mathcal{A}_i = (\rho_i|_{C_0})^* \mathcal{E}$. In other words, \mathcal{A}_i is a semiabelian scheme over C_0 which fits into the following commutative diagram:

$$\begin{array}{ccc} \mathcal{A}_i & \xrightarrow{\theta_i} & \mathcal{E} \\ \downarrow \pi_i & & \downarrow \pi \\ C_0 & \xrightarrow{\rho_i|_{C_0}} & S \\ \cap & & \cap \\ C_4 & \xrightarrow{\nu} \overline{C} \xrightarrow{\zeta_i} & \mathbb{P}^1 \end{array} \quad (19)$$

Since $\rho_i(C_0^*) \subset S^*$, the fibres of \mathcal{A}_i over C_0^* are elliptic curves. Meanwhile the fibres over the points $s_1, \dots, s_\ell \in C_0$ are isomorphic to \mathbb{G}_m , because $\rho_i(s_k) = 0$ for all i and k .

There may exist points $s \in C_0 \setminus C_0^* \setminus \{s_1, \dots, s_\ell\}$. For such points, the fibres $\mathcal{A}_{i,s}$ may be elliptic curves for some i and isomorphic to \mathbb{G}_m for other i . Such points will not appear further in our analysis.

As explained in Section 5.B, we will need to analyse periods of \mathcal{A}_i near the points s_1, \dots, s_ℓ (where “near” is interpreted with respect to an embedding of K). It is easier to compare periods of a larger number of semiabelian schemes over a neighbourhood of a single point, so we will instead consider the semiabelian schemes $\sigma_k^* \mathcal{A}_i$ ($1 \leq i \leq n$, $1 \leq k \leq \ell$), always near the point s_1 . The periods of $\sigma_k^* \mathcal{A}_i$ near s_1 are the same as the periods of \mathcal{A}_i near s_k . In order to avoid trivial relations between the periods of the schemes $\sigma_k^* \mathcal{A}_i$, we pick out one of these schemes in each generic isogeny class.

More precisely, let $\bar{\eta}$ denote a geometric generic point of C_0 . Define an equivalence relation \sim on $\{1, \dots, \ell\} \times \{1, \dots, n\}$ by

$$(k, i) \sim (k', i') \text{ if there exists an isogeny } \sigma_k^* \mathcal{A}_{i, \bar{\eta}} \rightarrow \sigma_{k'}^* \mathcal{A}_{i', \bar{\eta}}.$$

Let Λ denote a set of representatives of the equivalence classes for \sim .

We define some convenient notation:

- Given $(k, i) \in \{1, \dots, \ell\} \times \{1, \dots, n\}$, let $[k, i]$ be the unique element of Λ such that $[k, i] \sim (k, i)$.
- If $\lambda = (k, i) \in \Lambda$, we shall write $\mathcal{A}_\lambda = \sigma_k^* \mathcal{A}_i$ and $\rho_\lambda = \rho_i \circ \sigma_k: C_0 \rightarrow \mathbb{P}^1$.

Note that combining these two pieces of notation leads to $\mathcal{A}_{[k, i]} = \sigma_{k'}^* \mathcal{A}_{i'}$ where $[k, i] = (k', i')$. Thus if $(k, i) \notin \Lambda$, $\mathcal{A}_{[k, i]}$ need not be equal to $\sigma_k^* \mathcal{A}_i$. However $\mathcal{A}_{[k, i], \bar{\eta}}$ is always isogenous to $\sigma_k^* \mathcal{A}_{i, \bar{\eta}}$. Thanks to this isogeny, periods of $\mathcal{A}_{[k, i]}$ near s_1 are \bar{K} -linear combinations of periods of \mathcal{A}_i near s_k .

Define T to be a positive integer such that, for every $k = 1, \dots, \ell$ and $i = 1, \dots, n$, there exists an isogeny $\mathcal{A}_{[k, i], \bar{\eta}} \rightarrow \sigma_k^* \mathcal{A}_{i, \bar{\eta}}$ of degree dividing T .

Remark 5.3. Since C is Hodge generic in \mathbb{A}^n , there are no isogenies $\mathcal{A}_{i, \bar{\eta}} \rightarrow \mathcal{A}_{i', \bar{\eta}}$ for distinct i, i' . It follows that $(k, i) \not\sim (k, i')$ for all k and all $i \neq i'$.

5.E. Neighbourhoods. It is well-known that, if x is a local parameter at a point s_0 on an algebraic curve C over a number field K , then for every embedding $\iota: K \rightarrow \mathbb{C}_\iota$, x^ι restricts to an analytic isomorphism between a sufficiently small neighbourhood of s_0^ι in $C^\iota(\mathbb{C}_\iota)$ and an open disc in \mathbb{C}_ι . The following lemma is a formal statement of the properties we shall require of these analytic isomorphisms. A key point for us is that the discs can be taken to have radius at least 1 for almost all embeddings ι ; since this fact is perhaps not so well-known as the rest, we have written out a proof of the lemma.

In the subsequent sections of the paper, we will only use the neighbourhood $U_{\iota, 1}$. However, the full collection of open neighbourhoods $U_{\iota, 2}, \dots, U_{\iota, \ell}$ are needed in the proof of the properties of $U_{\iota, 1}$. In particular, (iii) is used in the proof that $x^\iota|_{U_{\iota, 1}}$ is injective, and this in turn is used in the proof of (v).

Lemma 5.4. *Let C_0 be a smooth algebraic curve over a number field K . Let $x \in K(C_0)$ be a rational function of degree ℓ , such that x has ℓ distinct unramified zeros $s_1, \dots, s_\ell \in C_0(K)$. Then for each embedding ι of K , there exists a real number $r_\iota > 0$ and open sets $U_{\iota,1}, \dots, U_{\iota,\ell} \subset C_0^\iota(\mathbb{C}_\iota)$ with the following properties:*

- (i) $r_\iota \geq 1$ for almost all embeddings ι of K ;
- (ii) $s_k^\iota \in U_{\iota,k}$;
- (iii) for each ι , the sets $U_{\iota,1}, \dots, U_{\iota,\ell}$ are pairwise disjoint and $U_{\iota,1} \cup \dots \cup U_{\iota,\ell}$ is equal to the preimage under x^ι of $D(0, r_\iota, \mathbb{C}_\iota)$;
- (iv) for each ι and k , x^ι restricts to a bijection from $U_{\iota,k}$ to the open disc $D(0, r_\iota, \mathbb{C}_\iota)$;
- (v) for every rational function $f \in K(C_0)$ which is regular at s_k , if $\hat{f} \in K[[X]]$ denotes the Taylor series of f around s_k in terms of the local parameter x (defined algebraically), then, for all $s \in U_{\iota,k}$ satisfying $|x^\iota(s)| < R(\hat{f}^\iota)$, we have $\hat{f}^\iota(x^\iota(s)) = f^\iota(s)$.

Proof. For an archimedean embedding ι , the existence of r_ι and $U_{\iota,1}, \dots, U_{\iota,\ell}$ satisfying (ii)–(v) is classical. We can ignore finitely many embeddings in (i), so for the rest of this proof we consider only non-archimedean embeddings.

Let $\overline{C_0}$ be the smooth projective model of C_0 . Embed $\overline{C_0}$ in some projective space \mathbb{P}^m . After applying a linear automorphism of \mathbb{P}^m , we may assume that $s_1, \dots, s_\ell \in \mathbb{A}^m$ (regarded as an open subset of \mathbb{P}^m). Replacing C_0 by a Zariski open subset if necessary, we may assume that $C_0 \subset \mathbb{A}^m$. Let $y_1, \dots, y_m: C_0 \rightarrow \mathbb{A}^1$ denote the coordinate functions.

Let $P, Q \in K[Y_1, \dots, Y_m]$ be polynomials such that

$$x = \frac{P(y_1, \dots, y_m)}{Q(y_1, \dots, y_m)}. \quad (20)$$

Let $\eta_{k,j} \in K[[X]]$ denote the Taylor series of y_j around s_k in terms of the local parameter x (defined algebraically).

For each non-archimedean embedding ι , let

$$d_\iota = \min\{1, |y_j(s_k)^\iota - y_j(s_{k'})^\iota| : 1 \leq k, k' \leq \ell, 1 \leq j \leq m, y_j(s_k) \neq y_j(s_{k'})\},$$

$$r_\iota = \min\{R^\dagger(\eta_{k,j}^\iota) d_\iota : 1 \leq k \leq \ell, 1 \leq j \leq m\}.$$

For each j, k, k' , $y_j(s_k) - y_j(s_{k'}) \in K$, so if $y_j(s_k) \neq y_j(s_{k'})$, then $|y_j(s_k)^\iota - y_j(s_{k'})^\iota| > 0$ for all ι and $|y_j(s_k)^\iota - y_j(s_{k'})^\iota| \geq 1$ for almost all ι . Hence $d_\iota > 0$ for all ι and $d_\iota \geq 1$ for almost all ι .

The field of rational functions $K(C_0)$ has transcendence degree 1, so each y_j is algebraic over $K(x)$. It follows that the power series $\eta_{k,j}$ are algebraic (over $K(X)$). Hence, by Eisenstein's theorem, the $\eta_{k,j}$ are globally bounded, so $R^\dagger(\eta_{k,j}^\iota) > 0$ for all ι and $R^\dagger(\eta_{k,j}^\iota) \geq 1$ for almost all ι . Thus $r_\iota > 0$ for all ι and $r_\iota \geq 1$ for almost all ι , proving (i).

Let

$$U_{\iota,k} = \{s \in C^{\dagger\iota}(\mathbb{C}_\iota) : |x^\iota(s)| < r_\iota, |y_j^\iota(s) - y_j(s_k)^\iota| < d_\iota \text{ for all } j = 1, \dots, m\}.$$

Clearly $s_k \in U_{\iota,k}$, so (ii) is satisfied.

Next we prove (iii) and (iv). Let $\underline{\eta}_k^\iota = (\eta_{k,1}^\iota, \dots, \eta_{k,m}^\iota): D(0, r_\iota, \mathbb{C}_\iota) \rightarrow \mathbb{A}^m(\mathbb{C}_\iota)$. We shall show that $\underline{\eta}_k^\iota$ is the inverse of $x^\iota|_{U_{\iota,k}}$.

Firstly, suppose that $s \in U_{\iota,k} \cap U_{\iota,k'}$ where $k \neq k'$. There exists some j such that $y_j(s_k) \neq y_j(s_{k'})$. Then from the definitions of $U_{\iota,k}$ and d_ι , we have

$$\begin{aligned} |y_j^\iota(s) - y_j(s_k)^\iota| &< d_\iota \leq |y_j(s_k)^\iota - y_j(s_{k'})^\iota|, \\ |y_j^\iota(s) - y_j(s_{k'})^\iota| &< d_\iota \leq |y_j(s_k)^\iota - y_j(s_{k'})^\iota|. \end{aligned}$$

This contradicts the non-archimedean triangle inequality. Thus $U_{\iota,1}, \dots, U_{\iota,\ell}$ are pairwise disjoint.

Consider $z \in D(0, r_\iota, \mathbb{C}_\iota)$. For every f in the ideal of $\mathcal{I}(C_0) \subset K[Y_1, \dots, Y_m]$, we have $f(y_1, \dots, y_m) = 0$ in $K[C_0]$, hence $f(\eta_{k,1}, \dots, \eta_{k,m}) = 0$ in $K[[X]]$, hence $f^\iota(\eta_{k,1}^\iota(z), \dots, \eta_{k,m}^\iota(z)) = 0$ in \mathbb{C}_ι (because taking Taylor series and evaluating ι -adically are both K -algebra homomorphisms). Thus, $\underline{\eta}_k^\iota(z) \in C_0(\mathbb{C}_\iota)$. A similar argument shows that

$$XQ(\eta_{k,1}, \dots, \eta_{k,m}) = P(\eta_{k,1}, \dots, \eta_{k,m}) \text{ in } K[[X]]$$

so

$$zQ^\iota(\eta_{k,1}^\iota(z), \dots, \eta_{k,m}^\iota(z)) = P^\iota(\eta_{k,1}^\iota(z), \dots, \eta_{k,m}^\iota(z)).$$

Hence,

$$x^\iota(\underline{\eta}_k^\iota(z)) = \frac{P^\iota(\underline{\eta}_k^\iota(z))}{Q^\iota(\underline{\eta}_k^\iota(z))} = z. \quad (21)$$

Since $|z| < r_\iota \leq R^\dagger(\eta_{k,j}^\iota)$, using Lemma 2.1, we have

$$|y_j^\iota(\underline{\eta}_k^\iota(z)) - y_j(s_k)^\iota| = |\eta_{k,j}^\iota(z) - \eta_{k,j}^\iota(0)| \leq R^\dagger(\eta_{k,j}^\iota)^{-1}|z| < R^\dagger(\eta_{k,j}^\iota)r_\iota \leq d_\iota$$

for all j . Thus $\underline{\eta}_k^\iota(z) \in U_{\iota,k}$.

By (21), the points $\underline{\eta}_1^\iota(z) \in U_{\iota,1}, \dots, \underline{\eta}_\ell^\iota(z) \in U_{\iota,\ell}$ are preimages of z in $C^\dagger(\mathbb{C}_\iota)$ under x^ι . These points are distinct because $U_{\iota,1}, \dots, U_{\iota,\ell}$ are pairwise disjoint. Since $\deg(x) = t$, these are the only preimages of z under x^ι in $C_0(\mathbb{C}_\iota)$.

This establishes that $U_{\iota,1} \cup \dots \cup U_{\iota,\ell} = (x^\iota)^{-1}(D(0, r_\iota, \mathbb{C}_\iota))$ and that each $z \in D(0, r_\iota, \mathbb{C}_\iota)$ has at most one preimage in each $U_{\iota,k}$ under x^ι . Thanks to (21), this implies that $x^\iota|_{U_{\iota,k}}$ is injective. It is immediate from (21) that $x^\iota|_{U_{\iota,k}}$ is surjective onto $D(0, r_\iota, \mathbb{C}_\iota)$. This completes the proof of (iii) and (iv).

Let $f \in K(C_0)$ be a rational function which is regular at s_k . Let $\hat{f} \in K[[X]]$ be the Taylor series of f around s_k in terms of x . Write $f = R/S$, where $R, S \in K[Y_1, \dots, Y_m]$ and S^ι has no zeros in $U_{\iota,k}$ except at poles of f^ι .

Suppose that $s \in U_{\iota,k}$ with $|x^\iota(s)| < R(\hat{f}^\iota)$. Let $z = x^\iota(s)$. By the same argument as for (21), we obtain

$$\hat{f}^\iota(z) = \frac{R^\iota(\eta_{k,1}^\iota(z), \dots, \eta_{k,m}^\iota(z))}{S^\iota(\eta_{k,1}^\iota(z), \dots, \eta_{k,m}^\iota(z))}. \quad (22)$$

Since x^ι restricts to a bijection $U_{\iota,k} \rightarrow D(0, r_\iota, \mathbb{C}_\iota)$ with inverse $\underline{\eta}_k^\iota$, we have $\underline{\eta}_k^\iota(z) = s$. Substituting this in (22) completes the proof of (v). \square

Remark 5.5. If $s \in C_0^\iota(\mathbb{C}_\iota)$ and $|x^\iota(s)| < r_\iota$, then there exists k such that $(\sigma_k^\iota)^{-1}(s) \in U_{\iota,1}$. Indeed, by Lemma 5.4(iv), there is a unique point $s' \in U_{\iota,1}$ such that $x^\iota(s') = x^\iota(s)$. By Lemma 5.1(iii), $x: C_4 \rightarrow \mathbb{P}^1$ is Galois, so there exists $\sigma \in \text{Aut}_x(C_4)$ such that $\sigma^\iota(s') = s$. By Remark 5.2, $\sigma = \sigma_k$ for some k .

5.F. Power series. We now define power series $F_\lambda, G_\lambda \in K[[X]]$ for each $\lambda \in \Lambda$. As we will show in the next subsection, the archimedean evaluations of these power series can be interpreted as locally invariant periods of \mathcal{A}_λ near s_1 . These will be the G-functions to which we apply Bombieri's theorem.

Let ξ_λ denote the Taylor series of $\rho_\lambda: C_0 \rightarrow S \subset \mathbb{A}^1$ in terms of the local parameter x near s_1 . (Note that, since $x \circ \sigma_{k'} = x$, this is the same as the Taylor series of ρ_i in terms of x near s_k , where $\lambda = (k, i)$.) These are power series in $K[[X]]$ and, since $\rho_\lambda(s_1) = \rho_i(s_k) = 0$, they have no constant term.

Shrinking r_ι and $U_{\iota,1}$ if necessary, we may assume that

$$r_\iota \leq R^\dagger(\xi_\lambda^\iota)$$

for all non-archimedean ι and all $(k, i) \in \Lambda$. Note that the power series ξ_λ are algebraic, hence globally bounded. Hence, we may perform this shrinking while preserving the property that

$$r_\iota > 0 \text{ for all } \iota \text{ and } r_\iota \geq 1 \text{ for almost all } \iota. \quad (23)$$

Similarly, for archimedean embeddings ι , shrinking r_ι and $U_{\iota,1}$ if necessary, we may assume that

$$r_\iota \leq R(\xi_\lambda^\iota)$$

and that ξ_λ^ι maps $D(0, r_\iota, \mathbb{C})$ into the disc Δ_S from Section 3.

Let F and G denote the power series defined in Section 3. For each $\lambda \in \Lambda$, let

$$F_\lambda = F \circ \xi_\lambda, \quad G_\lambda = G \circ \xi_\lambda \in K[[X]]. \quad (24)$$

Since $F \in \mathbb{Z}[[X]]$, $R(F^\iota) \geq 1$ for all non-archimedean embeddings ι of K . Hence, by Lemma 2.2,

$$R(F_\lambda^\iota) \geq R^\dagger(\xi_\lambda^\iota) \geq r_\iota.$$

Using Lemma 2.2 together with Lemma 5.4(v), the following diagram of ι -adic evaluations commutes:

$$\begin{array}{ccccc}
 U_{\iota,1} & & & & \\
 x^\iota \downarrow & \searrow^{\rho_\lambda^\iota} & & & \\
 D(0, r_\iota, \mathbb{C}_\iota) & \xrightarrow{\xi_\lambda^\iota} & D(0, R(F^\iota), \mathbb{C}_\iota) & \xrightarrow{F^\iota} & \mathbb{P}^1(\mathbb{C}_\iota) \\
 & & \searrow_{F_\lambda^\iota} & &
 \end{array} \tag{25}$$

We have not estimated $R(G^\iota)$ for non-archimedean ι , so we do not know whether this diagram makes sense with G in place of F . This is not a problem because we will not need to reason about non-archimedean evaluations of G_λ .

For archimedean embeddings ι , ξ_λ^ι maps $D(0, r_\iota, \mathbb{C}_\iota)$ to Δ_S , which is contained in $D(0, R(F^\iota), \mathbb{C}_\iota)$ and in $D(0, R(G^\iota), \mathbb{C}_\iota)$. Hence, all evaluations of power series in the above diagram make sense. Over \mathbb{C} , this is sufficient to guarantee that the diagram commutes, as well as the analogous diagram for G .

5.G. Periods and G-functions. Let $\lambda = (k, i) \in \Lambda$ and let ι be an archimedean embedding of \hat{K} . We shall interpret the ι -adic evaluation of the power series F_λ and G_λ as periods of the semiabelian scheme \mathcal{A}_λ near s_1 . This interpretation has two applications: showing that F_λ and G_λ are G-functions, and determining the trivial relations between them.

Write $\pi_\lambda: \mathcal{A}_\lambda \rightarrow C_0$ for the structure morphism of the semiabelian scheme \mathcal{A}_λ . Let θ_λ be the morphism which completes the following fibre product diagram:

$$\begin{array}{ccc}
 \mathcal{A}_\lambda & \xrightarrow{\theta_\lambda} & \mathcal{E} \\
 \downarrow \pi_\lambda & & \downarrow \pi \\
 C_0 & \xrightarrow{\rho_\lambda|_{C_0}} & S
 \end{array} \tag{26}$$

In Section 3.D, we defined a non-zero locally invariant cycle $\gamma \in R_1\pi_*^\iota(\mathbb{Q})(\Delta_S)$. For each $s \in C_0^{\iota*}(\mathbb{C})$, $\theta_{\lambda,s}^\iota$ is an isomorphism $\mathcal{A}_{\lambda,s}^\iota \rightarrow \mathcal{E}_{\rho_\lambda^\iota(s)}^\iota$ and hence induces an isomorphism $\theta_{\lambda,s*}^\iota: H_1(\mathcal{A}_{\lambda,s}^\iota, \mathbb{Q}) \rightarrow H_1(\mathcal{E}_{\rho_\lambda^\iota(s)}^\iota, \mathbb{Q})$. Consequently, for each $s \in U_{\iota,1}$, there is a unique $\gamma_{\iota,\lambda,s} \in H_1(\mathcal{A}_{\lambda,s}^\iota, \mathbb{Q})$ such that

$$\theta_{\lambda,s*}^\iota(\gamma_{\iota,\lambda,s}) = \gamma_{\rho_\lambda^\iota(s)}.$$

These cycles $\gamma_{\iota,\lambda,s}$ are locally constant as s varies in $U_{\iota,1}^*$. Since γ is a section in $\pi_*^\iota(\mathbb{Q})(\Delta_S^*)$, it follows that $\gamma_{\iota,\lambda,s}$ are the values of a section in $R_1\pi_{\lambda*}^\iota(\mathbb{Q})(U_{\iota,1}^*)$. By [And89, IX, 4.3, (4.3.2)], this extends uniquely to a section $\gamma_{\iota,\lambda} \in R_1\pi_{\lambda*}^\iota(\mathbb{Q})(U_{\iota,1})$.

Let $[\omega], [\eta] \in H_{DR}^1(\mathcal{E}/S)$ denote the de Rham cohomology classes of the differential forms ω, η defined in sections 3.A and 3.D respectively. Base-changing these via the fibre product diagram (26), we obtain

$$\omega_\lambda = \rho_\lambda^*\omega, \quad \eta_\lambda = \rho_\lambda^*\eta \in H_{DR}^1(\mathcal{A}_\lambda/C_0).$$

From (7) and (25), for every $s \in U_{\iota,1}$,

$$\frac{1}{2\pi i} \int_{\gamma_{\iota,\lambda,s}} \omega'_{\lambda,s} = \frac{1}{2\pi i} \int_{\gamma_{\rho'_\lambda(s)}} \omega'_{\rho'_\lambda(s)} = F^\iota(\rho'_\lambda(s)) = F'_\lambda(x^\iota(s)) \quad (27)$$

and, similarly,

$$\frac{1}{2\pi i} \int_{\gamma_{\iota,\lambda,s}} \eta'_{\lambda,s} = G^\iota_\lambda(x^\iota(s)).$$

Lemma 5.6. F_λ and G_λ are G -functions for all $\lambda \in \Lambda$.

Proof. Since $\pi_\lambda|_{C_0^*}$ is an abelian scheme with multiplicative reduction at s_1 , we can apply [And89, X, 4.2, Theorem 1] to deduce that there is some basis ω', η' for $H_{DR}^1(\mathcal{A}_\lambda/C_0)$ for which the Taylor expansions in x of the locally invariant periods $\frac{1}{2\pi i} \int_{\gamma_{\iota,\lambda,s}} \omega'_{\lambda,s}$, $\frac{1}{2\pi i} \int_{\gamma_{\iota,\lambda,s}} \eta'_{\lambda,s}$ are G -functions. Now ω and η are $K(C_0)$ -linear combinations of ω' and η' . Hence F_λ and G_λ are $K(X)$ -linear combinations of G -functions. Thus, by [And89, Theorem D], F_λ and G_λ are G -functions themselves. \square

5.H. Global relations. Let \mathcal{G} denote the set of G -functions defined in (24):

$$\mathcal{G} = \{F_\lambda, G_\lambda : \lambda \in \Lambda\}.$$

By definition, a **global relation** between the elements of \mathcal{G} at a point $z \in \hat{K}$ (a finite extension of K) is a polynomial $Q \in \hat{K}[Y_\lambda, Z_\lambda : \lambda \in \Lambda]$ which satisfies

$$Q^\iota(F_\lambda^\iota(t^\iota), G_\lambda^\iota(t^\iota) : \lambda \in \Lambda) = 0$$

for every embedding $\hat{\iota}$ of \hat{K} satisfying $|z^{\hat{\iota}}| < \min\{1, R(F_\lambda^\iota), R(G_\lambda^\iota) : \lambda \in \Lambda\}$.

In order to control the radii within which global relations have to hold, we shall add an additional G -function H to our set. Choose $\zeta \in K^\times$ such that $|\zeta^\iota| \leq r_\iota$ for all embeddings ι of K at which $r_\iota < 1$. (By (23), we have only constrained $|\zeta^\iota|$ at finitely many embeddings, so such a ζ exists.) Let H denote the G -function $\zeta/(\zeta - X) = \sum_{k=0}^{\infty} (X/\zeta)^k$ and let $\mathcal{G}' = \mathcal{G} \cup \{H\}$. This additional G -function H will not appear directly in our relations, but because it reduces the number of embeddings $\hat{\iota}$ at which a given parameter value z satisfies the radius bound, some polynomials in $\hat{K}[F_\lambda, G_\lambda]$ may be global relations between the elements of \mathcal{G}' at a given point without being global relations between the elements of \mathcal{G} .

Let $R'_\iota = \min\{1, R(H^\iota), R(F_\lambda^\iota), R(G_\lambda^\iota) : \lambda \in \Lambda\}$. Thus, for every embedding ι ,

$$R'_\iota \leq \min(1, R(H^\iota)) = \min(1, |\zeta^\iota|) \leq r_\iota.$$

Lemma 5.7. Let M, N be positive integers. Let $s \in C_0^*(\overline{\mathbb{Q}})$ be a point such that there exist $i_1, i_2, i_3, i_4 \in \{1, \dots, n\}$ such that $i_1 \neq i_2$, $i_3 \neq i_4$, $\{i_1, i_2\} \neq \{i_3, i_4\}$ and:

- (i) there exists an isogeny $\mathcal{A}_{i_1,s} \rightarrow \mathcal{A}_{i_2,s}$ of degree M ;
- (ii) there exists an isogeny $\mathcal{A}_{i_3,s} \rightarrow \mathcal{A}_{i_4,s}$ of degree N .

Then there exists a non-zero global relation between the evaluations of the G -functions \mathcal{G}' at $x(s)$ of degree at most $288\ell[K(s) : \mathbb{Q}] + 2\ell d(MT^2)$. This relation does not involve H .

Proof. The proof has two parts: first we construct a polynomial Q , then we prove that Q is a global relation.

Let i_1, i_2, i_3, i_4 be the indices appearing in the statement of the lemma. The hypotheses $i_1 \neq i_2$, $i_3 \neq i_4$, $\{i_1, i_2\} \neq \{i_3, i_4\}$ imply that either i_1, i_2, i_3, i_4 are distinct, or that one of i_1, i_2 is equal to one of i_3, i_4 and there are no other equalities between the indices. In the latter case, assume without loss of generality that $i_1 = i_3$.

The strategy is as follows. For each embedding $\hat{\iota}$ of $K(s)$, if $|x^\iota(s^\hat{\iota})|$ is small, then for some $k \in \{1, \dots, \ell\}$, the point $\sigma_k^{-1}(s)^\hat{\iota}$ is close to s_1 . Consequently $\hat{\iota}$ -adic locally invariant periods of the elliptic curves $\mathcal{A}_{[k, i_j], \sigma_k^{-1}(s)^\hat{\iota}}$ can be interpreted as evaluations of the G -functions $F_{[k, i_j]}$ and $G_{[k, i_j]}$ at $x(s)^\hat{\iota}$. The isogeny $\mathcal{A}_{i_1, s} \rightarrow \mathcal{A}_{i_2, s}$ from hypothesis (i) gives rise to an isogeny $\mathcal{A}_{[k, i_1], \sigma_k^{-1}(s)} \rightarrow \mathcal{A}_{[k, i_2], \sigma_k^{-1}(s)}$, and similarly for i_3, i_4 (after replacing $K(s)$ by a controlled extension \hat{K}). Hence, via Theorem 4.1, we get a relation between the $\hat{\iota}$ -adic evaluations of $F_{[k, i_j]}$ and $G_{[k, i_j]}$ at $x(s)$.

In the above outline of the strategy, the index k depends on the embedding $\hat{\iota}$. Therefore, we need to apply Theorem 4.1 once for each k and multiply the resulting relations together to obtain a global relation.

For each $k = 1, \dots, \ell$ and $j = 1, 2, 3, 4$, write $z_{kj} = \rho_{[k, i_j]}(\sigma_k^{-1}(s)) \in K(s)$.

By the definition of $\mathcal{A}_{[k, i_1]}$ and of T , there exists an isogeny $\mathcal{A}_{[k, i_1], \bar{\eta}} \rightarrow \sigma_k^* \mathcal{A}_{i_1, \bar{\eta}}$ of degree dividing T . Specialising this isogeny to the fibre over $\sigma_k^{-1}(s)$ (which requires extending scalars to \bar{K}) and composing with isomorphisms coming from fibre product diagrams (in particular, $\mathcal{A}_{[k, i_1]} \cong \rho_{[k, i_1]}^* \mathcal{E}$), we obtain

$$\mathcal{E}_{z_{k1}, \bar{K}} \cong \mathcal{A}_{[k, i_1], \sigma_k^{-1}(s), \bar{K}} \rightarrow (\sigma_k^* \mathcal{A}_{i_1})_{\sigma_k^{-1}(s), \bar{K}} \cong \mathcal{A}_{i_1, s, \bar{K}}.$$

We similarly obtain an isogeny $\mathcal{E}_{z_{k2}, \bar{K}} \rightarrow \mathcal{A}_{i_2, s, \bar{K}}$ of degree dividing T . By hypothesis, we have an isogeny $\mathcal{A}_{i_1, s} \rightarrow \mathcal{A}_{i_2, s}$ of degree M . We thus obtain a chain of isogenies

$$\mathcal{E}_{z_{k1}, \bar{K}} \rightarrow \mathcal{A}_{i_1, s, \bar{K}} \rightarrow \mathcal{A}_{i_2, s, \bar{K}} \rightarrow \mathcal{E}_{z_{k2}, \bar{K}}$$

whose composition ϕ_{12} has degree dividing MT^2 .

Similarly, we obtain an isogeny $\phi_{34}: \mathcal{E}_{z_{k3}, \bar{K}} \rightarrow \mathcal{E}_{z_{k4}, \bar{K}}$ of degree dividing NT^2 .

By [MW90, Lemma 6.1], ϕ_{12} and ϕ_{34} are each defined over extensions of $K(s)$ of degree at most 12. Thus we can choose an extension $\hat{K}/K(s)$ of degree at most 144 such that ϕ_{12} and ϕ_{34} are both defined over \hat{K} . For the remainder of this proof, we shall take this field \hat{K} as our base field.

Applying Theorem 4.1 to the isogenies ϕ_{12} and ϕ_{34} , we obtain non-zero polynomials $P_{\infty, k} \in \hat{K}[Y_1, Z_1, \dots, Y_4, Z_4]$ and $P_{fin, k} \in \hat{K}[Y_1, Y_2]$ such that

$$P_{\infty, k}(F^\iota(z_{k1}^\hat{\iota}), G^\iota(z_{k1}^\hat{\iota}), \dots, F^\iota(z_{k4}^\hat{\iota}), G^\iota(z_{k4}^\hat{\iota})) = 0 \quad (28)$$

for every archimedean embedding $\hat{\iota}$ of \hat{K} satisfying $z_{k1}^{\hat{\iota}}, \dots, z_{k4}^{\hat{\iota}} \in \Delta_S$ and

$$P_{fin,k}(F^\nu(z_{k1}^{\hat{\iota}}), F^\nu(z_{k2}^{\hat{\iota}})) = 0 \quad (29)$$

at every non-archimedean embedding $\hat{\iota}$ of \hat{K} satisfying $|z_{k1}^{\hat{\iota}}|, |z_{k2}^{\hat{\iota}}| < 1$.

Define a new polynomial $Q \in \hat{K}[Y_\lambda, Z_\lambda : \lambda \in \Lambda]$ by

$$Q(\underline{Y}, \underline{Z}) = \prod_{k=1}^{\ell} P_{\infty,k}(Y_{[k,i_1]}, Z_{[k,i_1]}, Y_{[k,i_2]}, Z_{[k,i_2]}, Y_{[k,i_3]}, Z_{[k,i_3]}, Y_{[k,i_4]}, Z_{[k,i_4]}) \\ \cdot P_{fin,k}(Y_{[k,i_1]}, Y_{[k,i_2]}).$$

We claim that $Q \neq 0$. Indeed, if i_1, i_2, i_3, i_4 are distinct, then, by Remark 5.3, for each k , $[k, i_1], [k, i_2], [k, i_3]$ and $[k, i_4]$ are distinct, so $Q \neq 0$ is immediate from the facts that $P_{\infty,k} \neq 0$ and $P_{fin,k} \neq 0$ for all k . If $i_1 = i_3$ and i_1, i_2, i_4 are distinct, then, similarly, for each k , $[k, i_1] = [k, i_3]$ while $[k, i_1], [k, i_2]$ and $[k, i_4]$ are distinct so the facts that $P_{\infty,k} \notin \langle Y_1 - Y_3, Z_1 - Z_3 \rangle$, $P_{fin,k} \neq 0$ suffice to establish that each factor of Q is non-zero.

From Theorem 4.1, we have $\deg(P_{\infty,k}) \leq 2[\hat{K} : \mathbb{Q}]$ and $\deg(P_{fin,k}) \leq d(MT^2)$ for all k . Hence

$$\deg(Q) \leq 2\ell[\hat{K} : \mathbb{Q}] + 2\ell d(MT^2) \leq 288\ell[K(s) : \mathbb{Q}] + 2\ell d(MT^2).$$

It remains to prove that Q is a global relation between the elements of \mathcal{G}' at the point $x(s)$. To that end, let $\hat{\iota}$ denote an embedding of \hat{K} and let ι denote its restriction to K . Suppose that $|x^\iota(s^\iota)| < R'_\iota$. In particular, $|x^\iota(s^\iota)| < r_\iota$.

By Remark 5.5, there exists some $k \in \{1, \dots, \ell\}$ such that $\sigma_k^{-1}(s)^\iota \in U_{\iota,1}$. (Note that this k depends on $\hat{\iota}$.)

From diagram (25), noting that $x \circ \sigma_k^{-1} = x$, we have

$$F^\nu(z_{k1}^{\hat{\iota}}) = F^\nu(\rho_{[k,i_1]}^\iota(\sigma_k^{-1}(s)^\iota)) = F_{[k,i_1]}^\nu(x^\iota(\sigma_k^{-1}(s)^\iota)) = F_{[k,i_1]}^\nu(x^\iota(s^\iota)). \quad (30)$$

Similar equalities hold for i_2, i_3 and i_4 and, at archimedean embeddings, for G in place of F .

If $\hat{\iota}$ is archimedean, then, since $\sigma_k^{-1}(s)^\iota$ and by our choice of r_ι , $\rho_{[k,i_1]}^\iota(s^\iota) \in \Delta_S$ and so on for i_2, i_3, i_4 . Hence we can substitute (30) (and analogous equations) into (28) and obtain

$$P_\infty(F_{[k,i_1]}^\nu(x(s)^\iota), G_{[k,i_1]}^\nu(x(s)^\iota), \dots, F_{[k,i_4]}^\nu(x(s)^\iota), G_{[k,i_4]}^\nu(x(s)^\iota)) = 0.$$

If $\hat{\iota}$ is non-archimedean, then, since $|x^\iota(s^\iota)| < r_\iota \leq R^\dagger(\xi_{k,i_1}^\iota)$, by Lemma 2.1, we have $|\rho_{[k,i_1]}^\iota(s^\iota)| < 1$ and similarly for i_2 . Hence, we can substitute (30) into (29) and obtain

$$P_{fin}(F_{[k,i_1]}^\nu(x(s)^\iota), F_{[k,i_2]}^\nu(x(s)^\iota)) = 0.$$

In either case, the factor of Q indexed by k vanishes at the $\hat{\iota}$ -adic evaluations of \mathcal{G}' at $x(s)$. Hence, Q itself vanishes at these evaluations. \square

5.I. Trivial relations. Our aim in this section is to prove that there are no non-zero trivial relations between the G-functions $\{F_\lambda, G_\lambda : \lambda \in \Lambda\}$. For this, it suffices to consider the evaluations of F_λ and G_λ at a single archimedean embedding ι .

Let \mathcal{A} denote the fibre product over C_0 of the semiabelian schemes \mathcal{A}_λ for $\lambda \in \Lambda$. Let $\mathcal{A}^* = \mathcal{A}|_{C_0^*}$ and let $\pi_{\mathcal{A}} : \mathcal{A}^* \rightarrow C_0^*$ denote the structure morphism. Because the elements of Λ lie in distinct classes for the equivalence relation \sim , the factors of $\mathcal{A}_{\bar{\eta}}$ are pairwise non-isogenous elliptic curves, so the generic Mumford–Tate group of $\mathcal{A}^{*\iota}$ is

$$\{(g_\lambda) \in \mathbf{GL}_2^\Lambda : \det(g_\lambda) \text{ is the same for all } \lambda \in \Lambda\}.$$

Let V denote a simply connected dense open subset of $U_{\iota,1}^* \subset C_0^{*\iota}(\mathbb{C})$. The local system $R_1\pi_{\lambda^*}^\iota(\mathbb{Q})|_V$ is trivial and has rank 2, so we may choose a section $\delta_{\iota,\lambda} \in R_1\pi_{\lambda^*}^\iota(\mathbb{Q})(V)$ such that $\gamma_{\iota,\lambda}$ and $\delta_{\iota,\lambda}$ form an oriented basis for $R_1\pi_{\lambda^*}^\iota(\mathbb{Q})|_V$.

We define holomorphic functions on V by

$$\begin{aligned} \mathcal{P}_{\lambda,1,1}(s) &= \frac{1}{2\pi i} \int_{\gamma_{\iota,\lambda,s}} \omega_{\lambda,s}^\iota, & \mathcal{P}_{\lambda,1,2}(s) &= \frac{1}{2\pi i} \int_{\delta_{\iota,\lambda,s}} \omega_{\lambda,s}^\iota, \\ \mathcal{P}_{\lambda,2,1}(s) &= \frac{1}{2\pi i} \int_{\gamma_{\iota,\lambda,s}} \eta_{\lambda,s}^\iota, & \mathcal{P}_{\lambda,2,2}(s) &= \frac{1}{2\pi i} \int_{\delta_{\iota,\lambda,s}} \eta_{\lambda,s}^\iota, \end{aligned}$$

forming the full period matrix of the elliptic curve $\mathcal{A}_{\lambda,s}^\iota$ with respect to the bases chosen above.

From (27), for all $s \in V$, we have

$$\mathcal{P}_{\lambda,1,1}(s) = F_\lambda^\iota(x^\iota(s)). \quad (31)$$

Similarly

$$\mathcal{P}_{\lambda,2,1}(s) = G_\lambda^\iota(x^\iota(s)).$$

Putting together all of these period functions $\mathcal{P}_{\lambda,\cdot,\cdot}$ yields a holomorphic function $\mathcal{P} : V \rightarrow \mathbf{M}_2(\mathbb{C})^\Lambda$, which gives the (block diagonal) period matrix of \mathcal{A}_s^ι . We denote its graph in $V \times \mathbf{M}_2(\mathbb{C})^\Lambda$ by Γ . The following lemma and its proof is inspired by [Pap, 7.2.1].

Lemma 5.8. *Let Θ denote the subvariety of $\mathbf{M}_{2,\mathbb{C}}$ defined by the equation $\det = (2\pi i)^{-1}$. The \mathbb{C} -Zariski closure of Γ in $C_0^{*\iota} \times \mathbf{M}_2^\Lambda$ is $C_0^{*\iota} \times \Theta^\Lambda$.*

Proof. The Legendre period relation states that Γ is contained in $C_0^{*\iota} \times \Theta^m$ (cf. equation (9)). To deduce the reverse inclusion let $s \in V$ be a Hodge generic point and let $\rho : \pi_1(C_0^{*\iota}(\mathbb{C}), s) \rightarrow \mathbf{SL}_2(\mathbb{Q})^\Lambda$ denote the monodromy representation on $R_1\pi_{\mathcal{A}^*}^\iota(\mathbb{Q}) = \bigoplus_{\lambda \in \Lambda} R_1\pi_{\lambda^*}^\iota(\mathbb{Q})$ at s (recall that the monodromy action preserves the polarization). Let \mathbf{H} denote the \mathbb{Q} -Zariski closure of the image of ρ . By André’s Normal Monodromy Theorem [And92, Theorem 1], the neutral component \mathbf{H}° is a normal subgroup of \mathbf{SL}_2^Λ . The connected normal subgroups of \mathbf{SL}_2 are $\{1\}$ and \mathbf{SL}_2 itself. Since the ρ_i are non-constant, \mathcal{A}_λ^* are not isotrivial and so the projections $\mathbf{H}^\circ \rightarrow \mathbf{SL}_2$ cannot be trivial. It follows immediately that $\mathbf{H}^\circ = \mathbf{SL}_2^\Lambda$. Since $\mathbf{H} \subset \mathbf{SL}_2^\Lambda$, it follows that $\mathbf{H} = \mathbf{H}^\circ$. We now argue exactly as in [Pap, 7.2.1]. \square

Let $\mathcal{P}' : V \rightarrow \mathbb{A}^{2 \times \Lambda}(\mathbb{C})$ denote the composition of \mathcal{P} with the projection $\mathbf{M}_2(\mathbb{C})^\Lambda \rightarrow \mathbb{A}^{2 \times \Lambda}(\mathbb{C})$ on to the first column in each factor. The following is immediate from Lemma 5.8:

Corollary 5.9. *The graph Γ' of \mathcal{P}' is \mathbb{C} -Zariski dense (and hence $\overline{\mathbb{Q}}$ -Zariski dense) in $C'_0 \times \mathbb{A}^{2 \times m}$.*

Proposition 5.10. *There are no non-zero trivial relations between the G -functions $\{F_\lambda, G_\lambda : \lambda \in \Lambda\}$. In other words, there is no non-zero polynomial $P \in \overline{\mathbb{Q}}[X][Y_\lambda, Z_\lambda : \lambda \in \Lambda]$ such that $P(X)(F_\lambda(X), G_\lambda(X) : \lambda \in \Lambda) = 0$ in $\overline{\mathbb{Q}}[[X]]$.*

Proof. Let C'_0 denote the complement in C_0 of the poles of x .

By (31), we have the following commutative diagram:

$$\begin{array}{ccc} V & \xrightarrow{\mathcal{P}'=(\mathcal{P}_{\lambda,j,1})} & \mathbb{A}^{2 \times \Lambda}(\mathbb{C}) \\ x^\iota \downarrow & \nearrow G=(g_{\lambda,j}) & \\ D(0, r_\iota, \mathbb{C}) & & \end{array}$$

where λ runs through Λ , j runs through $\{1, 2\}$, and $g_{\lambda,1} = F'_\lambda$, $g_{\lambda,2} = G'_\lambda$. Let P be as in the statement of the proposition (that is, $P(X)(g_{\lambda,j} : \lambda \in \Lambda, j \in \{1, 2\}) = 0$). Then the graph of G is contained in the \mathbb{C} -points of the subvariety $V(P)$ of $\mathbb{A}^1 \times \mathbb{A}^{2 \times \Lambda}$ defined by P . It follows that Γ' is contained in the subvariety of $C'_0 \times \mathbb{A}^{2 \times \Lambda}$ defined by $\tilde{P} = P \circ (x \times \text{id}_{\mathbb{A}^{2 \times \Lambda}}) \in \overline{\mathbb{Q}}[C'_0 \times \mathbb{A}^{2 \times \Lambda}]$. We conclude by Corollary 5.9 that $\tilde{P} = 0$. Since x is dominant, it follows that $P = 0$. \square

5.J. Proof of main theorems.

Proposition 5.11. *For every $\epsilon > 0$, there exist constants c_6 and c_7 such that the following holds: Let M, N be positive integers. Let $s \in C_0^*(\mathbb{Q})$ be a point such that there exist $i_1, i_2, i_3, i_4 \in \{1, \dots, n\}$ such that $i_1 \neq i_2$, $i_3 \neq i_4$, $\{i_1, i_2\} \neq \{i_3, i_4\}$, and:*

- (i) *there exists an isogeny $\mathcal{A}_{i_1,s} \rightarrow \mathcal{A}_{i_2,s}$ of degree M ;*
- (ii) *there exists an isogeny $\mathcal{A}_{i_3,s} \rightarrow \mathcal{A}_{i_4,s}$ of degree N .*

Then

$$h(x(s)) < c_6 \max\{[K(s) : K]^{c_7}, M^\epsilon\}.$$

Proof. Let s be a point satisfying the hypotheses of the proposition. By Lemma 5.7, there is a non-zero global relation Q between the elements of \mathcal{G}' at $x(s)$. Since this global relation does not involve the G -function H , Proposition 5.10 guarantees that it is non-trivial.

By [And89, Theorem E] (essentially due to Bombieri), we have

$$h(x(s)) \leq c_8 \deg(Q)^{c_9}$$

where the constants c_8, c_9 do not depend on s or ϵ . From Lemma 5.7 and (2), we have

$$\deg(Q) \leq 288\ell[K(s) : \mathbb{Q}] + 2\ell d(MT^2) \leq c_{10} \max\{[K(s) : K], M^{\epsilon/c_9}\}$$

where the constant c_{10} depends on ϵ but not on s . Putting these together proves the proposition. \square

Proposition 5.12. *There exist constants c_{11} and c_{12} such that the following holds: Let M, N be positive integers. Let $s \in C_0^*(\overline{\mathbb{Q}})$ be a point such that there exist $i_1, i_2, i_3, i_4 \in \{1, \dots, n\}$ such that $i_1 \neq i_2, i_3 \neq i_4, \{i_1, i_2\} \neq \{i_3, i_4\}$, and:*

- (i) *there exists a cyclic isogeny $\mathcal{A}_{i_1,s} \rightarrow \mathcal{A}_{i_2,s}$ of degree M ;*
- (ii) *there exists a cyclic isogeny $\mathcal{A}_{i_3,s} \rightarrow \mathcal{A}_{i_4,s}$ of degree N ;*
- (iii) *$\mathcal{A}_{i_1,s}$ and $\mathcal{A}_{i_3,s}$ do not have CM.*

Then

$$[K(s) : K] > c_{11} \max\{M, N\}^{c_{12}}.$$

Proof. This argument is essentially the one appearing in the proof of [HP12, Lemma 4.2]. Throughout this proof, the notation c_k will denote suitably chosen positive constants which depend on the curve C but not on M, N or s .

By swapping i_1, i_2 with i_3, i_4 if necessary, we may assume without loss of generality that $M \geq N$.

By [GR14, Theorem 1.4] (an effective version of the main theorem of [MW90]), there exists an isogeny $\mathcal{A}_{i_1,s} \rightarrow \mathcal{A}_{i_2,s}$ of degree

$$M' \leq c_{13} \max\{[K(s) : K], h_F(\mathcal{A}_{i_1,s})\}^{c_{14}},$$

where $h_F(\cdot)$ denotes the semistable Faltings height. By Silverman's comparison estimate between the Faltings and modular heights of elliptic curves [Sil86b, Prop. 2.1], we have

$$|h(1/\rho_{i_1}(s)) - 12h_F(\mathcal{A}_{i_1,s})| \leq c_{15} \log \max\{2, h(1/\rho_{i_1}(s))\}.$$

By the standard height machine [HS00, Theorem B.3.2], bearing in mind that $\text{NS}(\overline{C}) \cong \mathbb{Z}$, we have

$$|h(x(s)) - c_{16}h(1/\rho_{i_1}(s))| \leq c_{17}h(x(s)).$$

Combining the above inequalities, we obtain

$$M' \leq c_{18} \max\{[K(s) : K], h(x(s))\}^{c_{19}}.$$

Applying Proposition 5.11 with $\epsilon = 1/2c_{19}$, we get

$$M' \leq c_{20} \max\{[K(s) : K]^{c_{21}}, M^{1/2}\}.$$

Since $\mathcal{A}_{i_1,s}$ does not have CM, the \mathbb{Z} -module of isogenies $\mathcal{A}_{i_1,s} \rightarrow \mathcal{A}_{i_2,s}$ is generated by any cyclic isogeny. Therefore, $M \leq M'$ and the proposition follows. \square

To deduce Theorem 1.5, it suffices to note that if $s \in C(\overline{\mathbb{Q}}) \setminus \Sigma$ satisfies $\Phi_M(s_{i_1}, s_{i_2}) = \Phi_N(s_{i_3}, s_{i_4}) = 0$, then we can choose $\tilde{s} \in C_4(\overline{\mathbb{Q}})$ with $\nu(\tilde{s}) = s$. After removing finitely many points from C , we may assume that $\tilde{s} \in C_0(\overline{\mathbb{Q}})$. Now $\Phi_M(s_{i_1}, s_{i_2})$ implies that there is a cyclic isogeny $\mathcal{E}_{1/s_{i_1}} \rightarrow \mathcal{E}_{1/s_{i_2}}$ of degree M . We have

$$\mathcal{A}_{i_1, \tilde{s}} \cong \mathcal{E}_{\rho_{i_1}(\tilde{s})} = \mathcal{E}_{1/s_{i_1}}$$

and similarly for i_2 and for i_3, i_4 . If $\mathcal{A}_{i_1, \tilde{s}}$ has CM, then so does $\mathcal{A}_{i_2, \tilde{s}}$ (because they are isogenous), contradicting the assumption that $s \notin \Sigma$. Similarly $\mathcal{A}_{i_3, \tilde{s}}$ does not have CM. Thus $\tilde{s}, i_1, i_2, i_3, i_4$ satisfy the conditions of Proposition 5.12. Since $[\mathbb{Q}(s) : \mathbb{Q}]/[K(\tilde{s}) : K]$ is bounded below by a positive constant independent of s , the bound in Proposition 5.12 implies the bound in Theorem 1.5.

REFERENCES

- [And89] Y. André, *G-functions and geometry*, Aspects of Mathematics, E13, Friedr. Vieweg & Sohn, Braunschweig, 1989.
- [And92] Y. André, *Mumford–Tate groups of mixed Hodge structures and the theorem of the fixed part*, *Compositio Math.* **82** (1992), no. 1, 1–24.
- [And95] Y. André, *Théorie des motifs et interprétation géométrique des valeurs p -adiques de G -fonctions (une introduction)*, Number theory (Paris, 1992–1993), London Math. Soc. Lecture Note Ser., vol. 215, Cambridge Univ. Press, Cambridge, 1995, pp. 37–60.
- [BM21] G. Binyamini and D. Masser, *Effective André–Oort for non-compact curves in Hilbert modular varieties*, *Comptes Rendus. Mathématique* **359** (2021), no. 3, 313–321.
- [BSY] G. Binyamini, H. Schmidt, and A. Yafaev, *Lower bounds for Galois orbits of special points on Shimura varieties: a point-counting approach*, to appear in *Math. Ann.*, available at arXiv:2104.05842.
- [DO21a] C. Daw and M. Orr, *Unlikely intersections with $E \times \text{CM}$ curves in \mathcal{A}_2* , *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **22** (2021), no. 4, 1705–1745.
- [DO21b] ———, *Quantitative reduction theory and unlikely intersections*, *International Mathematics Research Notices* (2021), rnab173.
- [GR14] É. Gaudron and G. Rémond, *Théorème des périodes et degrés minimaux d’isogénies*, *Comment. Math. Helv.* **89** (2014), no. 2, 343–403.
- [HP12] P. Habegger and J. Pila, *Some unlikely intersections beyond André–Oort*, *Compos. Math.* **148** (2012), no. 1, 1–27.
- [HS00] M. Hindry and J. H. Silverman, *Diophantine geometry*, Graduate Texts in Mathematics, vol. 201, Springer-Verlag, New York, 2000, An introduction.
- [HW08] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, sixth ed., Oxford University Press, Oxford, 2008, Revised by D. R. Heath-Brown and J. H. Silverman, With a foreword by Andrew Wiles.
- [Küh12] L. Kühne, *An effective result of André–Oort type*, *Ann. of Math. (2)* **176** (2012), no. 1, 651–671.
- [MW90] D. Masser and G. Wüstholz, *Estimating isogenies on elliptic curves*, *Invent. Math.* **100** (1990), no. 1, 1–24.
- [Pap] G. Papas, *Height bounds for certain exceptional points in some variations of Hodge structures*, preprint, available at arXiv:2201.11240.
- [PST] J. Pila, A. Shankar, and J. Tsimerman., *Canonical Heights on Shimura Varieties and the André–Oort Conjecture*, preprint, available at arXiv:2109.08788.

- [Rob00] A. M. Robert, *A course in p -adic analysis*, Graduate Texts in Mathematics, vol. 198, Springer-Verlag, New York, 2000.
- [Sil86a] J. H. Silverman, *The arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 106, Springer-Verlag, New York, 1986.
- [Sil86b] ———, *Heights and elliptic curves*, Arithmetic geometry (Storrs, Conn., 1984), Springer, New York, 1986, pp. 253–265.
- [Sil94] ———, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 151, Springer-Verlag, New York, 1994.
- [Sta] The Stacks Project Authors, *Stacks Project*, <https://stacks.math.columbia.edu>.
- [Tat95] J. Tate, *A review of non-Archimedean elliptic functions*, Elliptic curves, modular forms, & Fermat’s last theorem (Hong Kong, 1993), Ser. Number Theory, I, Int. Press, Cambridge, MA, 1995, pp. 162–184.

DAW: DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF READING, WHITE-KNIGHTS, PO BOX 220, READING, BERKSHIRE RG6 6AX, UNITED KINGDOM

Email address: `chris.daw@reading.ac.uk`

ORR: DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF MANCHESTER, ALAN TURING BUILDING, OXFORD ROAD, MANCHESTER M13 9PL, UNITED KINGDOM

Email address: `martin.orr@manchester.ac.uk`