

A NOTE ON BISHOP-PHELPS-BOLLOBÁS PROPERTY FOR OPERATORS

T. GRANDO AND M. L. LOURENÇO

ABSTRACT. The main purpose of this paper is to study Bishop-Phelps-Bollobás property for operators on c_0 -sums of euclidean spaces. We show that the pair $(c_0(\bigoplus_{k=1}^{\infty} \ell_2^k), Y)$ has the Bishop-Phelps-Bollobás property for operators (in short BPBp for operators) whenever Y is a uniformly convex Banach space.

1. INTRODUCTION

In 1961, Bishop and Phelps [4], proved that, for any Banach space, the subset of norm attaining functionals is dense in the topological dual space. This result is known as Bishop-Phelps theorem. These authors posed the problem of possible extensions of such result to operators. A lot of attention has been devoted to extending Bishop-Phelps theorem to operators and interesting results have been obtained about this topic. In 1970, Bollobás [5], proved a refinement of the Bishop-Phelps theorem, which states that, every norm one functional and its almost norming points can be approximated by a norm attaining functional and its norm attaining point. In 2008, Acosta, Aron, Garcia and Maestre [1], defined a new notion for a pair of Banach spaces, which is called the Bishop-Phelps-Bollobás property for operators, and provided many notable results. Many important reference works in the field have appeared, among others, [1], [2], [6], [7], [8], [9], [11].

In [1] they showed, that (ℓ_{∞}^n, Y) satisfies the Bishop-Phelps-Bollobás property for operators for every $n \in \mathbb{N}$, whenever Y is a uniformly convex Banach space. They also left the open question if (c_0, Y) satisfies the BPBp for operators, whenever Y is a uniformly convex Banach space. In this sense, S. K. Kim [11], answered the question in a positive way. More generally, G. Choi and S. W. Kim proved, in [6], that $(c_0(\bigoplus_{k=1}^{\infty} X), Y)$ has BPBp for operators, if X is a uniformly convex

2010 *Mathematics Subject Classification*. Primary Primary 46B04; Secondary 46B07, 46B20.

Key words and phrases. Banach space, Bishop-Phelps-Bollobás theorem, Norm attaining operator,

Banach space and Y is \mathbb{C} -uniformly convex Banach space. Every uniformly convex complex space is \mathbb{C} -uniformly convex and the converse is not true.

The purpose of this work is to show that $(c_0(\bigoplus_{n=1}^{\infty} \ell_2^n), Y)$ satisfies the BPBp for operators whenever Y is uniformly convex complex Banach space. Considering the real case, when Y is strictly convex, we prove that if $(c_0(\bigoplus_{k=1}^{\infty} \ell_2^n), Y)$ satisfies BPBp for operators then Y is uniformly convex. In this sense, this is an improvement of the result presented in [6], since the c_0 -sum is performed with different spaces.

Notice that, the Banach space $c_0(\bigoplus_{n=1}^{\infty} \ell_2^n)$, known as c_0 -sums of the Euclidean n -space, is not isometric to c_0 . The importance of such space is due to the fact that the result presented by C. Stegall, in [14], where he showed that $\ell_{\infty}(\bigoplus_{n=1}^{\infty} \ell_2^n)$ does not have Dunford-Pettis property, but its predual, $\ell_1(\bigoplus_{n=1}^{\infty} \ell_2^n)$, has it.

Each $x \in c_0(\bigoplus_{n=1}^{\infty} \ell_2^n)$ can be represented by $x = \sum_{n=1}^{\infty} \sum_{k \in I(n)} x_k e_k$, where for every $n \in \mathbb{N}$, $I(n) = \{l \in \mathbb{N} : s(n-1)+1 \leq l \leq s(n)\}$ with $s : \mathbb{N}_0 \rightarrow \mathbb{N}_0$ the auxiliary function defined by $s(n) = 0$ if $n = 0$ and $s(n) = 1 + 2 + \dots + n$ if $n \neq 0$, and (e_k) is the standard basis of $c_0(\bigoplus_{n=1}^{\infty} \ell_2^n)$. The norm of x is given by the formula $\|x\| := \sup_{n \in \mathbb{N}} \left(\sum_{k \in I(n)} |x_k|^2 \right)^{1/2}$.

2. RESULTS

It will be convenient to recall the following notation. Let X and Y be Banach spaces. We denote by S_X , B_X , X^* , and $\mathcal{L}(X, Y)$, the unit sphere, the closed unit ball, the topological dual space of X and the space of all bounded linear operators from X into Y , respectively. An operator $T \in \mathcal{L}(X, Y)$ is said to attain its norm at $x_0 \in S_X$, if $\|T\| = \|T(x_0)\|$. We denote by $NA(X, Y)$ the subset of $\mathcal{L}(X, Y)$ of all norm attaining operators between X and Y . Now, we recall a few definitions.

Definition 2.1. *Let X and Y be Banach spaces. We say that the pair (X, Y) has the Bishop-Phelps-Bollobás property for operators (in shortly BPBp for operators) if given $\varepsilon > 0$, there is $\eta(\varepsilon) > 0$ such that whenever $T \in S_{\mathcal{L}(X, Y)}$ and $x_0 \in S_X$ satisfy that $\|Tx_0\| > 1 - \eta(\varepsilon)$, then there exist a point $u_0 \in S_X$ and an operator $S \in S_{\mathcal{L}(X, Y)}$ satisfying the following conditions*

$$\|Su_0\| = 1, \quad \|u_0 - x_0\| < \varepsilon, \quad \text{and} \quad \|S - T\| < \varepsilon.$$

A Banach space X is *uniformly convex* if for every $\epsilon > 0$ there is a $0 < \delta < 1$ such that, if $\left\| \frac{x+y}{2} \right\| > 1 - \delta$ for every $x, y \in B_X$, then $\|x - y\| < \epsilon$. In this case, the modulus of convexity is given by

$\delta(\epsilon) = \inf \left\{ 1 - \left\| \frac{x+y}{2} \right\| : x, y \in B_X, \|x - y\| \geq \epsilon \right\}$. A Banach space X is *strictly convex* if $\left\| \frac{x+y}{2} \right\| < 1$ whenever $x, y \in S_X$ and $x \neq y$. We remark that uniformly convexity implies strictly convexity, but the converse is not true.

We observe that in the following results, we will use similar techniques found in [1], [6] and [11].

Lemma 2.2. *Let Y be a strictly convex Banach space and let $T \in \mathcal{L}(c_0(\bigoplus_{k=1}^{\infty} \ell_2^k), Y)$. If $\|T(x)\| = \|T\|$ for some $x = \sum_{n=1}^{\infty} \sum_{k \in I(n)} x_k e_k \in S_{c_0(\bigoplus_{k=1}^{\infty} \ell_2^k)}$, so*

$$T(e_k) = 0 \quad \text{for all } k \in \left\{ j \in \mathbb{N} : \left(\sum_{i \in I(j)} |x_i|^2 \right)^{1/2} < 1 \right\}.$$

Proof. Let $x = \sum_{n=1}^{\infty} \sum_{k \in I(n)} x_k e_k$ be an element of $S_{c_0(\bigoplus_{k=1}^{\infty} \ell_2^k)}$ such that $\|T(x)\| = \|T\|$. The set $\left\{ j \in \mathbb{N} : \left(\sum_{i \in I(j)} |x_i|^2 \right)^{1/2} < 1 \right\}$ is nonempty, since $\|x\| = \sup_{n \in \mathbb{N}} \left(\sum_{i \in I(n)} |x_i|^2 \right)^{1/2} = 1$.

Now, we will show by a contradiction, and we assume that there exists $k_0 \in \left\{ j \in \mathbb{N} : \left(\sum_{i \in I(j)} |x_i|^2 \right)^{1/2} < 1 \right\}$ such that $T(e_{k_0}) \neq 0$. If we consider $I(k_0) = \{k, \dots, l\}$, where $k, \dots, l \in \mathbb{N}$, then the element $v := \left(x_k \pm \left(1 - \left(\sum_{i \in I(k_0)} |x_i|^2 \right)^{1/2} \right), x_{k+1}, \dots, x_l \right) \in \ell_2^{k_0}$ and $\|v\| \leq 1$. It implies that

$$\left\| x \pm \left(1 - \left(\sum_{i \in I(k_0)} |x_i|^2 \right)^{1/2} \right) e_k \right\| = \sup_{n \in \mathbb{N}} \left\{ \left(\sum_{i \in I(n) \setminus \{k_0\}} |x_i|^2 \right)^{1/2}, \|v\|_2 \right\} \leq 1.$$

By assumption $\|T(x)\| = \|T\|$, we have naturally $\|T(2x)\| = 2\|T\|$,

$$\begin{aligned} 2\|T\| &\leq \left\| T \left(x + \left(1 - \left(\sum_{i \in I(k_0)} |x_i|^2 \right)^{1/2} \right) e_k \right) \right\| \\ &\quad + \left\| T \left(x - \left(1 - \left(\sum_{i \in I(k_0)} |x_i|^2 \right)^{1/2} \right) e_k \right) \right\| \\ &\leq 2\|T\|. \end{aligned}$$

So, $\left\| T \left(x \pm \left(1 - \left(\sum_{i \in I(k_0)} |x_i|^2 \right)^{1/2} \right) e_k \right) \right\| = \|T\|$, and $\frac{T \left(x \pm \left(1 - \left(\sum_{i \in I(k_0)} |x_i|^2 \right)^{1/2} \right) e_k \right)}{\|T\|} \in S_Y$. Finally,

$$\left\| \frac{\frac{T \left(x + \left(1 - \left(\sum_{i \in I(k_0)} |x_i|^2 \right)^{1/2} \right) e_k \right)}{\|T\|} + \frac{T \left(x - \left(1 - \left(\sum_{i \in I(k_0)} |x_i|^2 \right)^{1/2} \right) e_k \right)}{\|T\|}}{2} \right\| = \left\| \frac{2T(x)}{2\|T\|} \right\| = 1.$$

Contradicting the fact that Y is a strictly convex Banach space, then $T(e_{k_0}) = 0$. \blacksquare

As a consequence of the Lemma 2.2 we get the following theorem.

Theorem 2.3. *Let $c_0 \left(\bigoplus_{k=1}^{\infty} \ell_2^k \right)$ a real Banach space and Y a strictly convex real Banach space. If $(c_0 \left(\bigoplus_{k=1}^{\infty} \ell_2^k \right), Y)$ satisfies the BPBp, then Y is an uniformly convex Banach space.*

Proof. Suppose that Y is not an uniformly convex Banach space. Then there exists $\epsilon > 0$ and sequences $(y_k), (z_k) \subset S_Y$ such that

$$(1) \quad \lim_{k \rightarrow \infty} \left\| \frac{y_k + z_k}{2} \right\| = 1 \quad \text{and} \quad \|y_k - z_k\| > \epsilon, \quad \forall k.$$

For each positive integer $i \in \mathbb{N}$, we define $T_i : c_0 \left(\bigoplus_{k=1}^{\infty} \ell_2^k \right) \rightarrow Y$ by

$$T_i(x) = \left(\frac{x_1 + x_2}{2} \right) y_i + \left(\frac{x_1 - x_2}{2} \right) z_i, \quad x = (x_k) \in X.$$

For each $i \in \mathbb{N}$ and each $x \in S_{c_0 \left(\bigoplus_{k=1}^{\infty} \ell_2^k \right)}$ we have that

$$\|T_i(x)\| \leq \frac{1}{2} \{ |x_1 + x_2| + |x_1 - x_2| \} \leq 1.$$

As $\|T_i(e_1 + e_2)\| = 1$, it follows that $\|T_i\| = 1$, for each $i \in \mathbb{N}$. We observe that, for each $i \in \mathbb{N}$, $\|T_i(e_1)\| = \left\| \frac{y_i + z_i}{2} \right\|$, by $\|T_i(e_1)\|$ converges to 1 when $i \rightarrow \infty$. Then there is $i_0 \in \mathbb{N}$ such that $\|T_{i_0}(e_1)\| > 1 - \eta \left(\frac{\epsilon}{2} \right)$. As the pair $(c_0 \left(\bigoplus_{k=1}^{\infty} \ell_2^k \right), Y)$ has the BPBp, there are an operator $R \in S_{\mathcal{L}(c_0 \left(\bigoplus_{k=1}^{\infty} \ell_2^k \right), Y)}$ and a point $u \in S_{c_0 \left(\bigoplus_{k=1}^{\infty} \ell_2^k \right)}$ such that

$$(2) \quad \|R(u)\| = 1, \quad \|R - T_{i_0}\| < \frac{\epsilon}{2}, \quad \|u - e_1\| < \beta \left(\frac{\epsilon}{2} \right) < 1.$$

Then $\left(\sum_{i \in I(k)} |u_i|^2\right)^{1/2} < 1$, for all $k \in \mathbb{N} \setminus \{1\}$, and by Lemma 2.2,

$$R(e_k) = 0, \quad \text{for all } k \in \mathbb{N} \setminus \{1\}.$$

Therefore, it can be assumed that $u = e_1$, so $R(e_1) = R(e_1 + e_2) = R(e_1 - e_2)$, it implies that

$$\begin{aligned} \|y_{i_0} - z_{i_0}\| &= \|T_{i_0}(e_1 + e_2) - T_{i_0}(e_1 - e_2)\| \\ &= \|T_{i_0}(e_1 + e_2) - R(e_1 + e_2) + R(e_1 - e_2) - T_{i_0}(e_1 - e_2)\| \\ &\leq \|T_{i_0} - R\| \|e_1 + e_2\| + \|R - T_{i_0}\| \|e_1 - e_2\| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned}$$

It is a contradiction, so Y is a uniformly convex Banach space. ■

We need the next Lemma to show the main result.

Lemma 2.4. *Let $F \subset \mathbb{N}$ and $A = \bigcup_{i \in F} I(i)$. Suppose that $0 < \epsilon < 1$ and Y an uniformly convex Banach space with modulus of convexity $\delta(\epsilon)$. If $T \in S_{\mathcal{L}(c_0(\oplus_{k=1}^{\infty} \ell_2^k), Y)}$ satisfies that $\|TP_A\| > 1 - \delta(\epsilon)$, then $\|T(I - P_A)\| \leq \epsilon$.*

Proof. Let $0 < \epsilon < 1$ and $T \in S_{\mathcal{L}(c_0(\oplus_{k=1}^{\infty} \ell_2^k), Y)}$ an operator such that $\|TP_A\| > 1 - \delta(\epsilon)$. Then there exists $x \in S_{c_0(\oplus_{k=1}^{\infty} \ell_2^k)} \cap P_A(c_0(\oplus_{k=1}^{\infty} \ell_2^k))$ such that $\|TP_A(x)\| > 1 - \delta(\epsilon)$. Fix an element $y = \sum_{n=1}^{\infty} \sum_{k \in I(n)} y_k e_k \in B_{c_0(\oplus_{k=1}^{\infty} \ell_2^k)}$ with $\text{supp } y \subset \mathbb{N} \setminus A$, then

$$\begin{aligned} \|x \pm y\| &= \sup_{j \in \mathbb{N}} \left\{ \left(\sum_{i \in I(j), j \in F} |x_i|^2 \right)^{1/2}, \left(\sum_{i \in I(j), j \in \mathbb{N} \setminus F} |y_i|^2 \right)^{1/2} \right\} \\ &\leq 1, \end{aligned}$$

it implies that, $\|T(x \pm y)\| \leq 1$, for every $y \in B_{c_0(\oplus_{k=1}^{\infty} \ell_2^k)}$ with $\text{supp } y \subset \mathbb{N} \setminus A$. Notice that, for every $z \in B_{c_0(\oplus_{k=1}^{\infty} \ell_2^k)}$, the support of the vector $(I - P_A)(z)$ is a subset of $\mathbb{N} \setminus A$ and then, $\|T(x) \pm T(I - P_A)(z)\| \leq 1$. So

$$\begin{aligned} \left\| \frac{T(x + (I - P_A)(z)) + T(x - (I - P_A)(z))}{2} \right\| &= \|TP_A(x)\| \\ &> 1 - \delta(\epsilon). \end{aligned}$$

As Y is an uniformly convex Banach space, it implies that

$$\|T(x + (I - P_A)(z)) - T(x - (I - P_A)(z))\| < \epsilon.$$

That is, $\|T(I - P_A)(z)\| < \frac{\epsilon}{2} < \epsilon$, whenever $z \in B_X$. This shows that $\|T(I - P_A)\| < \epsilon$. \blacksquare

In [1] the authors proved that the pair (X, Y) has the BPBp for operators when X and Y are both finite-dimensional. In the next proposition, we show that for specifically finite dimensional space X and Y is any uniformly convex space. The next proposition is similar the result in the Remark 2.5 in [6].

Proposition 2.5. *If Y is an uniformly convex Banach space, then the pair $(\ell_\infty(\bigoplus_{k=1}^n \ell_2^k), Y)$ satisfies the Bishop-Phelps-Bollobás for operators.*

Proof. The proof can be performed analogously to the Theorem. 2.4 in [6] with some changes, for example we need to use the convexity modulus equal the maximal in the convexity modulus of each space involved in the finite sum. \blacksquare

Theorem 2.6. *If Y is an uniformly convex Banach space, then the pair $(c_0(\bigoplus_{k=1}^\infty \ell_2^k), Y)$ satisfies the Bishop-Phelps-Bollobás for operators.*

Proof. Let $0 < \epsilon < 1$. As $(\ell_\infty(\bigoplus_{k=1}^n \ell_2^k), Y)$ has the BPBp for operators, itthen that there exist $0 < \eta(\epsilon) < \epsilon$ and $\beta(\epsilon) > 0$, with $\lim_{\epsilon \rightarrow 0} \beta(\epsilon) = 0$, such that, for every $Q \in S_{\mathcal{L}(\ell_\infty(\bigoplus_{k=1}^n \ell_2^k), Y)}$ and $z \in S_{\ell_\infty(\bigoplus_{k=1}^n \ell_2^k)}$ which satisfies that $\|Q(z)\| > 1 - \eta(\epsilon)$ there exist $\tilde{Q} \in S_{\mathcal{L}(\ell_\infty(\bigoplus_{k=1}^n \ell_2^k), Y)}$ and $z_0 \in S_{\ell_\infty(\bigoplus_{k=1}^n \ell_2^k)}$ such that

$$(3) \quad \|\tilde{Q}(z_0)\| = 1, \quad \|z - z_0\| < \beta(\epsilon), \quad \|\tilde{Q} - Q\| < \epsilon.$$

Now, let $T \in S_{\mathcal{L}(c_0(\bigoplus_{k=1}^\infty \ell_2^k), Y)}$ and $x = \sum_{n=1}^\infty \sum_{k \in I(n)} x_k e_k \in S_{c_0(\bigoplus_{k=1}^\infty \ell_2^k)}$ such that

$$\|T(x)\| > 1 - \eta(\epsilon) + \gamma(\epsilon),$$

and

$$\|T(x)\| > 1 - \delta(\epsilon) + \gamma(\epsilon),$$

where $\delta(\epsilon) > 0$ is the modulus of convexity of Y , $\gamma(\epsilon) > 0$ and $\lim_{\epsilon \rightarrow 0} \gamma(\epsilon) = 0$. Since c_{00} is a dense subspace of $c_0(\bigoplus_{k=1}^\infty \ell_2^k)$, we can be choose a vector $u \in S_{c_0(\bigoplus_{k=1}^\infty \ell_2^k)}$, with finite support, such that $\|x - u\| < \gamma(\epsilon)$. Then

$$(4) \quad \begin{aligned} \|T(u)\| &= \|T(x - x + u)\| \geq \|T(x)\| - \|T(x - u)\| \\ &> 1 - \eta(\epsilon). \end{aligned}$$

Similarly, $\|T(u)\| > 1 - \delta(\epsilon)$. Let $n = \min\{k \in \mathbb{N} : \text{supp } u \subset \bigcup_{j=1}^k I(j)\}$, and $A = \bigcup_{k=1}^n I(k)$. Then

$$\|TP_A\| \geq \|TP_A(u)\| = \|T(u)\| > 1 - \delta(\epsilon),$$

and

$$(5) \quad \|TP_A\| > 1 - \eta(\epsilon).$$

By Lemma 2.4,

$$(6) \quad \|T(I - P_A)\| \leq \epsilon.$$

Let $J : \ell_\infty \left(\bigoplus_{k=1}^n \ell_2^k \right) \rightarrow c_0 \left(\bigoplus_{k=1}^\infty \ell_2^k \right)$ be the map defined by

$$J(w) = \begin{cases} w_i, & \text{se } i \in A \\ 0, & \text{se } i \in \mathbb{N} \setminus A. \end{cases}$$

We observe that $\|J(w)\| = \max_{1 \leq j \leq n} \left(\sum_{i \in I(j)} |w_i|^2 \right)^{1/2} = \|w\|$, for every $w \in \ell_\infty \left(\bigoplus_{i=1}^n \ell_2^k \right)$. Let $Q : \ell_\infty \left(\bigoplus_{k=1}^n \ell_2^k \right) \rightarrow Y$ be the bounded linear operator defined by $Q(w) = \frac{TP_A J}{\|TP_A J\|}(w)$ and the element $z = (z_i) \in \ell_\infty \left(\bigoplus_{k=1}^n \ell_2^k \right)$ given by $z_i = u_i$, if $i \in A$ and $z_i = 0$ if $i \in \mathbb{N} \setminus A$. It is easy to check that $\|Q\| = \|z\| = 1$, and as $\|TP_A J\| \leq 1$, we get

$$(7) \quad \|Q(z)\| = \left\| \frac{TP_A J}{\|TP_A J\|}(z) \right\| \geq \|TP_A(u)\| = \|T(u)\|,$$

and by (4) we have that $\|Q(z)\| > 1 - \eta(\epsilon)$. By Proposition 2.5 the pair $(\ell_\infty \left(\bigoplus_{k=1}^n \ell_2^k \right), Y)$ has the BPBp for operators, so there exist $\tilde{R} \in S_{\mathcal{L}(\ell_\infty \left(\bigoplus_{k=1}^n \ell_2^k \right), Y)}$ and $\tilde{u} \in S_{\ell_\infty \left(\bigoplus_{k=1}^n \ell_2^k \right)}$, such that

$$(8) \quad \|\tilde{R}(\tilde{u})\| = 1, \quad \|\tilde{R} - Q\| < \epsilon, \quad \|z - \tilde{u}\| < \beta(\epsilon).$$

Now, we consider $(e_k), (f_k)$ be the canonical basis of $c_0 \left(\bigoplus_{k=1}^\infty \ell_2^k \right)$ and $\ell_\infty \left(\bigoplus_{k=1}^n \ell_2^k \right)$, respectively, and $R : c_0 \left(\bigoplus_{k=1}^\infty \ell_2^k \right) \rightarrow Y$ be the bounded linear operator given by

$$R(y) = \sum_{j=1}^{\infty} \sum_{i \in I(j)} y_i R(e_i),$$

where

$$R(e_i) = \begin{cases} \tilde{R}(f_i), & \text{if } i \in A \\ 0, & \text{if } i \in \mathbb{N} \setminus A, \end{cases}$$

and the vector $v = (v_i)_i \in c_0 \left(\bigoplus_{k=1}^\infty \ell_2^k \right)$ defined by

$$v_i = \begin{cases} \tilde{u}_i, & i \in A \\ x_i, & i \in \mathbb{N} \setminus A. \end{cases}$$

So $R \in S_{\mathcal{L}(c_0(\oplus_{k=1}^{\infty} \ell_2^k), Y)}$, $v \in S_{c_0(\oplus_{k=1}^{\infty} \ell_2^k)}$ and $\|R(v)\| = \|\tilde{R}(\tilde{u})\| = 1$.

It follows that R attain its norm in v . Next we will show that $\|R - T\| < \epsilon$.

$$\begin{aligned}
\|R - T\| &\leq \left\| R - \frac{TP_A}{\|TP_A\|} \right\| + \left\| \frac{TP_A}{\|TP_A\|} - TP_A \right\| + \|TP_A - T\| \\
&= \left\| \tilde{R} - \frac{TP_A J}{\|TP_A J\|} \right\| + \left\| \frac{TP_A}{\|TP_A\|} - TP_A \right\| + \|TP_A - T\| \\
&= \left\| \tilde{R} - \frac{TP_A J}{\|TP_A J\|} \right\| + \|TP_A\| \left| \frac{1}{\|TP_A\|} - 1 \right| + \|TP_A - T\| \\
&= \left\| \tilde{R} - Q \right\| + |1 - \|TP_A\|| + \|TP_A - T\| \\
&< \epsilon + 1 - 1 + \eta(\epsilon) + \epsilon < 3\epsilon.
\end{aligned}$$

Finally, we show that the vectors v and x are close,

$$\begin{aligned}
\|v - x\| = \|P_A(v - x)\| &= \max_{1 \leq j \leq n} \left(\sum_{i \in I(j)} |v_i - x_i|^2 \right)^{1/2} \\
&= \max_{1 \leq j \leq n} \left(\sum_{i \in I(j)} |\tilde{u}_i - x_i|^2 \right)^{1/2} \\
(9) \quad &\leq \max_{1 \leq j \leq n} \left(\sum_{i \in I(j)} |\tilde{u}_i - u_i|^2 \right)^{1/2} \\
&\quad + \max_{1 \leq j \leq n} \left(\sum_{i \in I(j)} |u_i - x_i|^2 \right)^{1/2} \\
&\leq \|\tilde{u} - z\| + \|u - x\| < \beta(\epsilon) + \gamma(\epsilon),
\end{aligned}$$

where $\lim_{\epsilon \rightarrow 0} \beta(\epsilon) + \gamma(\epsilon) = 0$. Therefore $(c_0(\oplus_{k=1}^{\infty} \ell_2^k), Y)$ satisfies the *BPBp* for operators. \blacksquare

REFERENCES

- [1] M.D. Acosta, R.M. Aron, D. García and M. Maestre, *The Bishop-Phelps-Bollobás theorem for operators*, J. Funct. Anal. **254** (11) (2008), 2780–2799.
- [2] M Acosta, J. Becerra-Guerrero, Y.S. Choi, M. Ciesielski S. K. Kim, H.. Lee, M.L. Lourenço and M. Martín *The Bishop-Phelps-Bollobás property for operators between spaces of continuous functions*, Nonlinear Analysis, Theory, Methods and Applications **95**(2014),323-332.

- [3] R.M. Aron, Y.S. Choi, D. García and M. Maestre, *The Bishop-Phelps-Bollobás theorem for $\mathcal{L}(L_1(\mu), L_\infty[0, 1])$* , Adv. Math. **228** (1) (2011), 617-628.
- [4] E. Bishop and R.R. Phelps, *A proof that every Banach space is subreflexive*, Bull. Amer. Math. Soc. (N.S.) **67** (1961), 97-98.
- [5] B. Bollobás, *An extension to the theorem of Bishop and Phelps*, Bull. London Math. Soc. **2** (1970), 181-182.
- [6] G. Choi and S. K. Kim, *The Bishop-Phelps-Bollobás property on the space of c_0 -sum*, Mediterr. J. Math. **19**, 72(2022), 16pp.
- [7] Y.S. Choi and S.K. Kim, *The Bishop-Phelps-Bollobás theorem for operators from $L_1(\mu)$ to Banach spaces with the Radon-Nikodým property*, J. Funct. Anal. **261**(6) (2011), 1446-1456.
- [8] Y.S. Choi and S.K. Kim, *The Bishop-Phelps-Bollobás property and lush spaces*, J. Math. Anal. Appl. **390** (2012), 549-555.
- [9] Y.S. Choi, S.K. Kim, H.J. Lee and M. Martín, *The Bishop-Phelps-Bollobás theorem for operators on $L_1(\mu)$* , J. Funct. Anal. **267**(91)(2014),214-242.
- [10] Y. S. Choi, S. K. Kim, H. J. Lee and M. Martín, *On Banach spaces with the approximate hyperplane series property*, Banach J. Math. Anal. **9** (4)(2015), 243-258.
- [11] S.K. Kim, *The Bishop-Phelps-Bollobás theorem for operators from c_0 to uniformly convex spaces*, Israel J. Math. **197**(1) (2013), 425-435.
- [12] S.K. Kim and H.J. Lee, *Uniform convexity and Bishop-Phelps-Bollobás property*, Canadian J. Math. **66**(2)(2014),372-386.
- [13] J. Lindenstrauss, *On operators which attain their norm*, Israel J. Math. **1** (1963), 139-148.
- [14] C. Stegall, *Duals of certain spaces with the Dunford-Pettis property*, Notices Amer. Math. Soc. **19** (1972), 799.

THIAGO GRANDO - DEMAT- UNIVERSIDADE ESTADUAL DO CENTRO-OESTE , ALAMEDA ÉLIO ANTONIO DALLA VECCHIA, VILA CARLI, 05508090 - GUARAPUAVA, PR - BRAZIL,EMAIL: TGRANDO@UNICENTRO.BR

MARY LILIAN LOURENÇO - DEPARTAMENTO DE MATEMÁTICA- INSTITUTO DE MATEMÁTICA E ESTATÍSTICA-UNIVERSIDADE DE SÃO PAULO CAIXA POSTAL 66281 - CEP : 05315-970 - SÃO PAULO, BRAZIL. E.MAIL : MLLOUREN@IME.USP.BR