

SOME CONGRUENCES FROM THE KARLSSON-MINTON SUMMATION FORMULA

CHEN WANG, JUNHANG LI, AND YEZHENYANG TANG

ABSTRACT. Let p be an odd prime. In this paper, by using the well-known Karlsson-Minton summation formula, we mainly prove two supercongruences as variants of a supercongruence of Deines-Fuselier-Long-Swisher-Tu, which confirm some recent conjectures of V.J.W. Guo.

1. INTRODUCTION

For any $n \in \mathbb{N} = \{0, 1, 2, \dots\}$, let $(x)_n = x(x+1)\cdots(x+n-1)$ denote the Pochhammer symbol. For $n, r \in \mathbb{N}$ and $a_0, \dots, a_r, b_1, \dots, b_r, z \in \mathbb{C}$ with $(b_1)_n, \dots, (b_r)_n$ being nonzero, the truncated hypergeometric series ${}_{r+1}F_r$ are defined as

$${}_{r+1}F_r \left[\begin{matrix} a_0 & a_1 & \cdots & a_r \\ b_1 & \cdots & b_r \end{matrix} \middle| z \right]_n = \sum_{k=0}^n \frac{(a_0)_k \cdots (a_r)_k}{(b_1)_k \cdots (b_r)_k} \cdot \frac{z^k}{k!}.$$

Clearly, they are partial sums of the classical hypergeometric series. Let p be an odd prime. For any integer $n \geq 1$, the p -adic Gamma function introduced by Morita (cf. [7, 11]) is defined as

$$\Gamma_p(n) = (-1)^n \prod_{1 \leq k < n} \frac{1}{p^{\lfloor k/p \rfloor}}.$$

Moreover, set $\Gamma_p(0) = 1$, and for any p -adic integer x set

$$\Gamma_p(x) = \lim_{n \rightarrow x} \Gamma_p(n),$$

where n runs through any sequence of positive integers p -adically approaching x .

Rodriguez-Villegas [12] investigated hypergeometric families of Calabi-Yau manifolds, and discovered (numerically) a number of possible supercongruences. Some of them have been proved in [8, 9] where Mortenson, with the help of the Gross-Koblitz formula, determined ${}_2F_1 \left[\begin{matrix} \alpha & 1-\alpha \\ 1 \end{matrix} \middle| 1 \right]_{p-1}$ modulo p^2 for $\alpha \in \{1/2, 1/3, 1/4, 1/6\}$. For instance, he showed that

$${}_2F_1 \left[\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 \end{matrix} \middle| 1 \right]_{p-1} \equiv (-1)^{(p-1)/2} \pmod{p^2} \quad (1.1)$$

for any prime $p \geq 5$. Using the Legendre relation of p -adic Gamma function (cf. [11, p. 370]), we may replace the right-hand side of (1.1) with $-\Gamma_p(1/2)^2$. Later, Sun [13] extended

2020 *Mathematics Subject Classification*. Primary 33C20; Secondary 05A10, 11B65, 11A07, 33E50.

Key words and phrases. truncated hypergeometric series, supercongruences, Karlsson-Minton summation formula.

Mortenson's result to the general p -adic integer α . Let \mathbb{Z}_p denote the ring of all p -adic integers and $\mathbb{Z}_p^\times := \{x \in \mathbb{Z}_p : p \nmid x\}$. Z.-H. Sun proved that for each odd prime p and $\alpha \in \mathbb{Z}_p^\times$,

$${}_2F_1 \left[\begin{matrix} \alpha & 1 - \alpha \\ 1 \end{matrix} \middle| 1 \right]_{p-1} \equiv (-1)^{\langle -\alpha \rangle_p} \pmod{p^2}, \quad (1.2)$$

where $\langle x \rangle_p$ is the least nonnegative residue of x modulo p , i.e., $0 \leq \langle x \rangle_p \leq p-1$ and $x \equiv \langle x \rangle_p \pmod{p}$. On the other hand, Deines et al. [1] obtained the following generalization of (1.1): for any integer $d > 1$ and prime $p \equiv 1 \pmod{d}$,

$${}_dF_{d-1} \left[\begin{matrix} 1 - \frac{1}{d} & 1 - \frac{1}{d} & \cdots & 1 - \frac{1}{d} \\ 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1} \equiv -\Gamma_p \left(\frac{1}{d} \right)^d \pmod{p^2}. \quad (1.3)$$

In fact, Deines et al. also conjectured that for any integer $d \geq 3$ and prime $p \equiv 1 \pmod{d}$, (1.3) holds modulo p^3 , and this conjecture was later confirmed by the first author and Pan [14].

In the past decade, many mathematicians studied q -analogues of (1.1) and its generalizations; among these q -congruences, the first one was obtained by Guo and Zeng [4]. Recently, via the so-called 'creative microscoping' method introduced by Guo and Zudilin [5], Guo [3] established a q -analogue of (1.3). Meanwhile, Guo obtained a variant of (1.3) as follows: for any integer $d > 1$ and prime $p \equiv 1 \pmod{d}$,

$$\sum_{k=0}^{p-1} \frac{k \left(\frac{d-1}{d} \right)_k^d}{k!^d} \equiv \frac{(d-1) \Gamma_p \left(\frac{1}{d} \right)^d}{2d} \pmod{p^2}.$$

The main purpose of this paper is to prove the following variants of (1.3) which confirm two conjectures of Guo [3, (5.4) and (5.5)].

Theorem 1.1. (i) *Let $d \geq 4$ be an even integer. Then, for any prime $p \equiv -1 \pmod{d}$ with $p \geq 2d-1$,*

$${}_dF_{d-1} \left[\begin{matrix} \frac{1}{d} - 1 & 1 + \frac{1}{d} & 1 + \frac{1}{d} & \cdots & 1 + \frac{1}{d} \\ 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1} \equiv \frac{d-1}{d^2} \Gamma_p \left(-\frac{1}{d} \right)^d \pmod{p^2}. \quad (1.4)$$

(ii) *Let $d \geq 3$ be an odd integer. Then, for any prime $p \equiv -1 \pmod{d}$,*

$${}_dF_{d-1} \left[\begin{matrix} \frac{1}{d} & \frac{1}{d} & 1 + \frac{1}{d} & 1 + \frac{1}{d} & \cdots & 1 + \frac{1}{d} \\ 1 & 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1} \equiv -\frac{1}{d^2} \Gamma_p \left(-\frac{1}{d} \right)^d \pmod{p^2}. \quad (1.5)$$

The rest of this paper is organized as follows. In the next section, we list some necessary lemmas which play key roles in the whole paper. Section 3 is devoted to the proof of Theorem 1.1. In Section 4, we shall pose a conjecture for further research.

2. SOME NECESSARY LEMMAS

The first key ingredient of our proofs is the following Karlsson-Minton summation formula (cf. [2, p. 19]).

Lemma 2.1. *Let m_1, m_2, \dots, m_n be nonnegative integers. Then*

$$\begin{aligned} & {}_{n+1}F_n \left[\begin{matrix} -(m_1 + \dots + m_n) & b_1 + m_1 & \dots & b_n + m_n \\ & b_1 & \dots & b_n \end{matrix} \middle| 1 \right] \\ &= (-1)^{m_1 + \dots + m_n} \cdot \frac{(m_1 + \dots + m_n)!}{(b_1)_{m_1} \dots (b_n)_{m_n}}. \end{aligned} \quad (2.1)$$

Our proofs also rely on some properties of the p -adic Gamma functions.

Lemma 2.2. [11, p. 369] *Let p be an odd prime and $x \in \mathbb{Z}_p$. Then*

$$\frac{\Gamma_p(x+1)}{\Gamma_p(x)} = \begin{cases} -x, & \text{if } p \nmid x, \\ -1, & \text{if } p \mid x, \end{cases} \quad (2.2)$$

$$\Gamma_p(x)\Gamma_p(1-x) = (-1)^{\langle -x \rangle_{p-1}}. \quad (2.3)$$

Remark 2.1. (a) By (2.2), it is easy to see that for any positive integer $n \leq p$,

$$\Gamma_p(n) = (-1)^n \Gamma(n). \quad (2.4)$$

(b) (2.3) is a p -adic analogue of the following Legendre relation of the classical Gamma function:

$$\Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin \pi x}.$$

The next lemma concerns a p -adic approximation to Γ_p -quotients.

Lemma 2.3. [6, Theorem 14] *For any prime $p \geq 5$ and $x \in \mathbb{Z}_p$, there exists $G_1(x) \in \mathbb{Z}_p$ such that for any $t \in \mathbb{Z}_p$,*

$$\Gamma_p(x+tp) \equiv \Gamma_p(x)(1+G_1(x)tp) \pmod{p^2}. \quad (2.5)$$

For the properties of $G_1(x)$, the reader may consult [10]. The following lemma lists two identities involving the derivatives of $(1+\alpha+x)_k$, which can be verified directly.

Lemma 2.4. *For any integer $k \geq 0$ and $\alpha, \beta \in \mathbb{R}$,*

$$\begin{aligned} \frac{d}{dx}(1+\alpha+x)_k &= (1+\alpha+x)_k \sum_{j=1}^k \frac{1}{j+\alpha+x}, \\ \frac{d}{dx} \left(\frac{1}{(1+\beta+x)_k} \right) &= - \frac{1}{(1+\beta+x)_k} \sum_{j=1}^k \frac{1}{j+\beta+x}. \end{aligned}$$

3. PROOF OF THEOREM 1.1

Throughout this section, we set $m = (p+1)/d$. We first prove (1.4). To show (1.4), we need the following preliminary result.

Lemma 3.1. *Under the assumptions of Theorem 1.1 (i), modulo p , we have*

$$\sum_{k=0}^{p-1} \frac{(m-1)_k (m+1)_k^{d-1}}{(1)_k^d} \left(\sum_{j=0}^{k-1} \frac{1}{m-1+j} - \sum_{j=0}^{k-1} \frac{1}{m+1+j} \right) \equiv \frac{(p-1)!}{(1)_{m-2} (1)_m^{d-1}} \left(\frac{1}{m} + \frac{1}{m-1} \right).$$

Proof. For $x, y \in (-1, +\infty)$, set

$$\Psi(x, y) = {}_{d+1}F_d \left[\begin{matrix} 1-p & m-1+x & m+1+y & m+1 & \cdots & m+1 \\ & 1+x & 1+y & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1}. \quad (3.1)$$

Clearly, $\Psi(x, y)$ is smooth on $(-1, +\infty) \times (-1, +\infty)$. Since $(1-p)_k = 0$ for all $k \geq p$, we have

$$\Psi(x, y) = {}_{d+1}F_d \left[\begin{matrix} 1-p & m-1+x & m+1+y & m+1 & \cdots & m+1 \\ & 1+x & 1+y & 1 & \cdots & 1 \end{matrix} \middle| 1 \right].$$

As

$$m-2 + (d-1)m = m-2 + p+1 - m = p-1,$$

by Lemma 2.1,

$$\Psi(x, y) = \frac{(p-1)!}{(1+x)_{m-2} (1+y)_m (1)_m^{d-2}}. \quad (3.2)$$

Now we calculate $\Psi_x(0, 0) - \Psi_y(0, 0)$ in two different ways, where $\Psi_x(0, 0)$ and $\Psi_y(0, 0)$ stand for the partial derivatives of Ψ at $(0, 0)$ with respect to x and y . By (3.1) and Lemma 2.4, we obtain

$$\begin{aligned} \Psi_x(0, 0) &= \sum_{k=0}^{p-1} \frac{(1-p)_k (m-1)_k (m+1)_k^{d-1}}{(1)_k^{d+1}} \left(\sum_{j=0}^{k-1} \frac{1}{m-1+j} - H_k \right), \\ \Psi_y(0, 0) &= \sum_{k=0}^{p-1} \frac{(1-p)_k (m-1)_k (m+1)_k^{d-1}}{(1)_k^{d+1}} \left(\sum_{j=0}^{k-1} \frac{1}{m+1+j} - H_k \right), \end{aligned}$$

where $H_k = \sum_{j=1}^k 1/j$ denotes the harmonic number. Therefore,

$$\Psi_x(0, 0) - \Psi_y(0, 0) = \sum_{k=0}^{p-1} \frac{(1-p)_k (m-1)_k (m+1)_k^{d-1}}{(1)_k^{d+1}} \left(\sum_{j=0}^{k-1} \frac{1}{m-1+j} - \sum_{j=0}^{k-1} \frac{1}{m+1+j} \right). \quad (3.3)$$

On the other hand, by (3.2) and Lemma 2.4,

$$\Psi_x(0, 0) - \Psi_y(0, 0) = \frac{(p-1)!}{(1)_{m-2} (1)_m^{d-1}} \left(\frac{1}{m} + \frac{1}{m-1} \right). \quad (3.4)$$

Note that for k among $0, 1, \dots, p-1$, $(1-p)_k \equiv (1)_k \pmod{p}$ and

$$\frac{(m-1)_k(m+1)_k^{d-1}}{(1)_k^{d+1}} \left(\sum_{j=0}^{k-1} \frac{1}{m-1+j} - \sum_{j=0}^{k-1} \frac{1}{m+1+j} \right) \in \mathbb{Z}_p.$$

This, together with (3.3) and (3.4), gives the desired result. \square

Proof of (1.4). For any $x, y \in \mathbb{Z}_p$, let

$$\Phi(x, y) = {}_dF_{d-1} \left[\begin{matrix} m-1+x & m+1+y & \cdots & m+1+y \\ & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1}.$$

Obviously, for any $s, t \in \mathbb{Z}_p$,

$$\Phi(sp, tp) \equiv \Phi(0, 0) + sp\Phi_x(0, 0) + tp\Phi_y(0, 0) \pmod{p^2}. \quad (3.5)$$

In particular,

$$\Phi(-p, 0) \equiv \Phi(0, 0) - p\Phi_x(0, 0) \pmod{p^2}. \quad (3.6)$$

Substituting (3.6) into (3.5), we get

$$\Phi(sp, tp) \equiv \Phi(-p, 0) + (s+1)p\Phi_x(0, 0) + tp\Phi_y(0, 0) \pmod{p^2}.$$

Taking $s = t = -1/d$, in view of Lemmas 2.4 and 3.1,

$$\Phi\left(-\frac{p}{d}, -\frac{p}{d}\right) \quad (3.7)$$

$$= {}_dF_{d-1} \left[\begin{matrix} \frac{1}{d}-1 & 1+\frac{1}{d} & 1+\frac{1}{d} & \cdots & 1+\frac{1}{d} \\ & 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1}$$

$$\equiv \Phi(-p, 0) + \left(1 - \frac{1}{d}\right) p\Phi_x(0, 0) - \frac{p}{d}\Phi_y(0, 0)$$

$$\equiv \Phi(-p, 0) + \left(1 - \frac{1}{d}\right) p \sum_{k=0}^{p-1} \frac{(m-1)_k(m+1)_k^{d-1}}{(1)_k^d} \left(\sum_{j=0}^{k-1} \frac{1}{m-1+j} - \sum_{j=0}^{k-1} \frac{1}{m+1+j} \right)$$

$$\equiv \Phi(-p, 0) + \left(1 - \frac{1}{d}\right) p \cdot \frac{(p-1)!}{(1)_{m-2}(1)_m^{d-1}} \left(\frac{1}{m} + \frac{1}{m-1} \right) \pmod{p^2}. \quad (3.8)$$

Now we evaluate $\Phi(-p, 0)$ modulo p^2 . Since $m(d-1) = p+1-m$, by Lemma 2.1,

$$\Phi(-p, 0) = (-1)^m \frac{(p+1-m)!}{(1)_m^{d-1}}.$$

With the help of (2.2), we obtain

$$\begin{aligned} \Phi(-p, 0) &= (-1)^m \frac{\Gamma(p-m+2)}{\Gamma(m+1)^{d-1}} = (-1)^{p+2-(m+1)(d-1)} \frac{\Gamma_p(p-m+2)}{\Gamma_p(m+1)^{d-1}} \\ &= (-1)^{1-(m+1)(d-1)} (p-m)(p-m+1) \frac{\Gamma_p(p-m)}{\Gamma_p(m+1)^{d-1}} \end{aligned}$$

$$= (-1)^m (p-m)(p-m+1) \frac{\Gamma_p(p-m)}{\Gamma_p(m+1)^{d-1}}, \quad (3.9)$$

where in the last step we have used the fact that d is even. In light of (2.3) and Lemma 2.3,

$$\begin{aligned} \frac{\Gamma_p(p-m)}{\Gamma_p(m+1)^{d-1}} &= (-1)^{(m-1)(d-1)} \Gamma_p(p-m) \Gamma_p(-m)^{d-1} \\ &\equiv (-1)^{m-1} \Gamma_p \left(-\frac{1}{d} \right)^d \left(1 + \left(1 - \frac{1}{d} \right) p G_1 \left(-\frac{1}{d} \right) \right) \left(1 - \frac{p}{d} G_1 \left(-\frac{1}{d} \right) \right)^{d-1} \\ &\equiv (-1)^{m-1} \Gamma_p \left(-\frac{1}{d} \right)^d \pmod{p^2}. \end{aligned} \quad (3.10)$$

Moreover,

$$\begin{aligned} (p-m)(p-m+1) &= \left(-\frac{1}{d} + \left(1 - \frac{1}{d} \right) p \right) \left(1 - \frac{1}{d} + \left(1 - \frac{1}{d} \right) p \right) \\ &\equiv \frac{1-d}{d^2} + \frac{(d-1)(d-2)p}{d^2} \pmod{p^2}. \end{aligned}$$

Combining this with (3.9) and (3.10), we obtain

$$\Phi(-p, 0) \equiv \left(\frac{d-1}{d^2} - \frac{(d-1)(d-2)p}{d^2} \right) \Gamma_p \left(-\frac{1}{d} \right)^d \pmod{p^2}. \quad (3.11)$$

Similarly, it is routine to verify that

$$\left(1 - \frac{1}{d} \right) p \cdot \frac{(p-1)!}{(1)_{m-2} (1)_m^{d-1}} \left(\frac{1}{m} + \frac{1}{m-1} \right) \equiv \frac{(d-1)(d-2)p}{d^2} \Gamma_p \left(-\frac{1}{d} \right)^d \pmod{p^2}. \quad (3.12)$$

Substituting (3.11) and (3.12) into (3.7), we immediately get (1.4). \square

In order to show (1.5), we also need an auxiliary lemma.

Lemma 3.2. *Under the assumptions of Theorem 1.1 (ii), we have*

$$\sum_{k=0}^{p-1} \frac{(m)_k^2 (m+1)_k^{d-2}}{(1)_k^d} \left(\sum_{j=0}^{k-1} \frac{1}{m+j} - \sum_{j=0}^{k-1} \frac{1}{m+1+j} \right) \equiv \frac{(p-1)!}{(1)_{m-1} (1)_m^{d-1}} \pmod{p}.$$

Proof. For $x, y \in (-1, +\infty)$, set

$$\Upsilon(x, y) = {}_{d+1}F_d \left[\begin{matrix} 1-p & m+x & m & m+1+y & m+1 & \cdots & m+1 \\ & 1+x & 1 & 1+y & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1}.$$

Similarly as in the proof of Lemma 3.1, we are led to the desired result by considering $\Upsilon_x(0, 0) - \Upsilon_y(0, 0)$. \square

Proof of (1.5). For any $x, y \in \mathbb{Z}_p$, set

$$\Omega(x, y) = {}_dF_{d-1} \left[\begin{matrix} m+x & m+y & m+1+y & \cdots & m+1+y \\ & 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1}.$$

Similarly as before, by Lemmas 2.1, 2.4 and 3.2,

$$\Omega\left(-\frac{p}{d}, -\frac{p}{d}\right) \tag{3.13}$$

$$= {}_dF_{d-1} \left[\begin{matrix} \frac{1}{d} & \frac{1}{d} & 1+\frac{1}{d} & 1+\frac{1}{d} & \cdots & 1+\frac{1}{d} \\ & 1 & 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1}$$

$$\equiv \Omega(-p, 0) + \left(1 - \frac{1}{d}\right) p \Omega_x(0, 0) - \frac{p}{d} \Omega_y(0, 0)$$

$$\equiv \Omega(-p, 0) + \left(1 - \frac{2}{d}\right) p \sum_{k=0}^{p-1} \frac{(m)_k^2 (m+1)_k^{d-2}}{(1)_k^d} \left(\sum_{j=0}^{k-1} \frac{1}{m+j} - \sum_{j=0}^{k-1} \frac{1}{m+1+j} \right)$$

$$\equiv \Omega(-p, 0) + \left(1 - \frac{2}{d}\right) p \cdot \frac{(p-1)!}{(1)_{m-1} (1)_m^{d-1}} \pmod{p^2}. \tag{3.14}$$

Then, in view of Lemmas 2.1–2.3, we have

$$\Omega(-p, 0) \equiv \left(-\frac{1}{d^2} + \frac{(d-2)p}{d^2}\right) \Gamma_p \left(-\frac{1}{d}\right)^d \pmod{p^2}$$

and

$$\left(1 - \frac{2}{d}\right) p \cdot \frac{(p-1)!}{(1)_{m-1} (1)_m^{d-1}} \equiv \frac{(2-d)p}{d^2} \cdot \Gamma_p \left(-\frac{1}{d}\right)^d \pmod{p^2}.$$

The proof of (1.5) follows by combining the above. \square

4. CONCLUDING REMARKS

It is routine to check that

$$\begin{aligned} & {}_dF_{d-1} \left[\begin{matrix} \frac{1}{d} - 1 & 1 + \frac{1}{d} & \cdots & 1 + \frac{1}{d} \\ & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1} + (d-1) {}_dF_{d-1} \left[\begin{matrix} \frac{1}{d} & \frac{1}{d} & 1 + \frac{1}{d} & \cdots & 1 + \frac{1}{d} \\ & 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1} \\ &= d \cdot {}_dF_{d-1} \left[\begin{matrix} \frac{1}{d} - 1 & \frac{1}{d} & 1 + \frac{1}{d} & \cdots & 1 + \frac{1}{d} \\ & 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1}. \end{aligned} \tag{4.1}$$

Note that Guo [3, Corollaries 4.2 and 4.4] proved that (1.4) and (1.5) also hold for odd integers $d \geq 3$ and even integers $d \geq 4$, respectively. This, together with (1.4), (1.5) and (4.1), gives that

$${}_dF_{d-1} \left[\begin{matrix} \frac{1}{d} - 1 & \frac{1}{d} & 1 + \frac{1}{d} & 1 + \frac{1}{d} & \cdots & 1 + \frac{1}{d} \\ & 1 & 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{p-1} \equiv 0 \pmod{p^2} \tag{4.2}$$

for any integer $d \geq 3$ and prime $p \equiv -1 \pmod{d}$ with $p \neq d - 1$. In fact, (4.2) can also be proved independently by using the method we used to prove Theorem 1.1 and the following Karlsson-Minton summation formula:

$${}_{n+1}F_n \left[\begin{matrix} a & b_1 + m_1 & \cdots & b_n + m_n \\ & b_1 & \cdots & b_n \end{matrix} \middle| 1 \right] = 0$$

provided that m_1, \dots, m_n are nonnegative integers and $\Re(-a) > m_1 + \cdots + m_n$.

Motivated by (4.2) and based on some numerical calculations, we made the following conjecture for further study.

Conjecture 4.1. *Let $d \geq 2$ be an integer. Let n be a positive integer with $n \equiv -1 \pmod{d}$ and $n < d - 1$. Then*

$$\frac{(n-1)! d^{dn-d}}{n^2} \cdot {}_dF_{d-1} \left[\begin{matrix} \frac{1}{d} - 1 & \frac{1}{d} & 1 + \frac{1}{d} & 1 + \frac{1}{d} & \cdots & 1 + \frac{1}{d} \\ & 1 & 1 & 1 & \cdots & 1 \end{matrix} \middle| 1 \right]_{n-1} \in \mathbb{Z}. \quad (4.3)$$

We think it is possible that the ‘creative microscoping’ could be used to prove Conjecture 4.1. We hope that an interested reader will make some progress on it.

Acknowledgment. This work is supported by the Jiangsu Province Student Innovation Training Program (grant no. 202210298128Y).

REFERENCES

- [1] A. Deines, J.G. Fuselier, L. Long, H. Swisher and F.-T. Tu, *Hypergeometric series, truncated hypergeometric series, and Gaussian hypergeometric functions*, Directions in number theory, 125–159, Assoc. Women Math. Ser. 3, Springer, 2016.
- [2] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, 2nd ed., Encyclopedia of Mathematics and its Applications 96, Cambridge University Press, Cambridge, 2004.
- [3] V.J.W. Guo, *Some q -supercongruences from Gasper’s Karlsson-Minton type summation*, Ramanujan J. (2022), in press. <https://doi.org/10.1007/s11139-022-00621-0>
- [4] V.J.W. Guo and J. Zeng, *Some q -analogues of supercongruences of Rodriguez-Villegas*, J. Number Theory **145** (2014), 301–316.
- [5] V.J.W. Guo and W. Zudilin, *A q -microscope for supercongruences*, Adv. Math. **346** (2019), 329–358.
- [6] L. Long and R. Ramakrishna, *Some supercongruences occurring in truncated hypergeometric series*, Adv. Math. **290** (2016), 773–808.
- [7] Y. Morita, *A p -adic analogue of the Γ -function*. J. Fac. Sci. Univ. Tokyo Sect. IA Math. **22** (1975), 255–266.
- [8] E. Mortenson, *A supercongruence conjecture of Rodriguez-Villegas for a certain truncated hypergeometric function*, J. Number Theory **99** (2003), 139–147.
- [9] E. Mortenson, *Supercongruences between truncated ${}_2F_1$ hypergeometric functions and their Gaussian analogs*, Trans. Amer. Math. Soc. **355** (2003), 987–1007.
- [10] H. Pan, R. Tauraso and C. Wang, *A local-global theorem for p -adic supercongruences*, J. reine angew. Math. (2022), in press. <https://doi.org/10.1515/crelle-2022-0032>
- [11] A.M. Robert, *A Course in p -Adic Analysis*, Springer, New York, 2000.
- [12] F. Rodriguez-Villegas, *Hypergeometric families of Calabi-Yau manifolds*, in: Calabi-Yau Varieties and Mirror Symmetry (Toronto, ON, 2001), Fields Inst. Commun., 38, Amer. Math. Soc., Providence, RI, 2003, pp. 223–231.

- [13] Z.-H. Sun, *Generalized Legendre polynomials and related supercongruences*, J. Number Theory **143** (2014) 293–319.
- [14] C. Wang and H. Pan, *Supercongruences concerning truncated hypergeometric series*, Math. Z. **300** (2022), 161–177.

(Chen Wang) DEPARTMENT OF APPLIED MATHEMATICS, NANJING FORESTRY UNIVERSITY, NANJING 210037, PEOPLE'S REPUBLIC OF CHINA

Email address: `cwang@smail.nju.edu.cn`

(Junhang Li) DEPARTMENT OF APPLIED MATHEMATICS, NANJING FORESTRY UNIVERSITY, NANJING 210037, PEOPLE'S REPUBLIC OF CHINA

Email address: `745679122@qq.com`

(Yezhenyang Tang) DEPARTMENT OF APPLIED MATHEMATICS, NANJING FORESTRY UNIVERSITY, NANJING 210037, PEOPLE'S REPUBLIC OF CHINA

Email address: `2742425229@qq.com`