

ON A CONJECTURE OF R. M. MURTY AND V. K. MURTY

YUCHEN DING

ABSTRACT. Let $\omega^*(n)$ be the number of primes p such that $p-1$ divides n . Recently, R. M. Murty and V. K. Murty proved that

$$x(\log \log x)^3 \ll \sum_{n \leq x} \omega^*(n)^2 \ll x \log x.$$

They further conjectured that there is some positive constant C such that

$$\sum_{n \leq x} \omega^*(n)^2 \sim Cx \log x$$

as $x \rightarrow \infty$. In this short note, we give the correct order of the sum by showing that

$$\sum_{n \leq x} \omega^*(n)^2 \asymp x \log x.$$

Let $\omega(n)$ be the number of distinct prime divisors of n . In about one hundred years ago, Hardy and Ramanujan [3] found out that $\omega(n)$ has normal order $\log \log n$ which means that for all most all integers n we have $\omega(n) \sim \log \log n$. Later, Turán [6] provided a quite elegantly simplified proof by establishing

$$\sum_{n \leq x} (\omega(n) - \log \log n)^2 \ll x \log \log x.$$

In 1955, Prachar [5] considered a variant arithmetic function of ω . Let $\omega^*(n)$ be the number of primes p such that $p-1$ divides n . Prachar proved that

$$\sum_{n \leq x} \omega^*(n) = x \log \log x + Bx + O(x/\log x)$$

and

$$\sum_{n \leq x} \omega^*(n)^2 = O(x(\log x)^2),$$

where B is a constant. Motived by Prachar's work, Erdős and Prachar [2] proved that the number of pairs of primes p and q so that the least common multiple $[p-1, q-1] \leq x$ is bounded by $O(x \log \log x)$. Following a remark of Erdős and Prachar, R. M. Murty and V. K. Murty [4] improved this to $O(x)$. By this improvement, they reached the nice bounds

$$x(\log \log x)^3 \ll \sum_{n \leq x} \omega^*(n)^2 \ll x \log x.$$

2010 *Mathematics Subject Classification*. Primary 11A41.

Key words and phrases. Primes; Bombieri–Vinogradov theorem.

With these in hands, R. M. Murty and V. K. Murty conjectured that there is some positive constant C such that

$$\sum_{n \leq x} \omega^*(n)^2 \sim Cx \log x$$

as $x \rightarrow \infty$. In this note, the author shall give a slight improvement of the result due to R. M. Murty and V. K. Murty towards the correct direction of their conjecture.

Theorem 1. *There are two absolute constants a_1 and a_2 such that*

$$a_1 x \log x \leq \sum_{n \leq x} \omega^*(n)^2 \leq a_2 x \log x.$$

Proof. We only need to prove the lower bound as the upper bound is displayed by R. M. Murty and V. K. Murty. Throughout our proof, the number x is sufficiently large. From the paper of R. M. Murty and V. K. Murty [4, equation (4.10)], we have

$$\sum_{n \leq x} \omega^*(n)^2 = x \sum_{d \leq x} \varphi(d) \left(\sum_{\substack{p \leq x \\ p \equiv 1 \pmod{d}}} \frac{1}{p-1} \right)^2 + O(x). \quad (1)$$

Integrating by parts gives

$$\sum_{\substack{p \leq x \\ p \equiv 1 \pmod{d}}} \frac{1}{p} = \frac{\pi(x; d, 1)}{x} + \int_2^x \frac{\pi(t; d, 1)}{t^2} dt \geq \int_{x^{3/4}}^x \frac{\pi(t; d, 1)}{t^2} dt, \quad (2)$$

where $\pi(t; d, 1)$ is the number of primes $p \equiv 1 \pmod{d}$ up to t . Thus, from equations (1) and (2) we obtain

$$\sum_{n \leq x} \omega^*(n)^2 \geq x \sum_{d \leq x^{1/3}} \varphi(d) \left(\int_{x^{3/4}}^x \frac{\pi(t; d, 1)}{t^2} dt \right)^2 + O(x). \quad (3)$$

For any integer $0 \leq j \leq \left\lfloor \frac{\log x}{13 \log 2} \right\rfloor$, let $Q_j = 2^j x^{1/4}$. Then $Q_j < x^{1/3}$ for all integers j . From a weak form of the Bombieri–Vinogradov theorem (see for example [1]), we have

$$\sum_{Q_j < d \leq 2Q_j} \max_{y \leq z} \left| \pi(y; d, 1) - \frac{\text{li } y}{\varphi(d)} \right| \ll \frac{z}{(\log z)^5},$$

for any $0 \leq j \leq \left\lfloor \frac{\log x}{13 \log 2} \right\rfloor$ and $x^{3/4} \leq z \leq x$, where the implied constant is absolute. It follows immediately that

$$\max_{y \leq z} \left| \pi(y; d, 1) - \frac{\text{li } y}{\varphi(d)} \right| < \frac{\text{li } z}{\varphi(d) \log z} \quad (4)$$

hold for all $Q_j < d \leq 2Q_j$ but at most $O(Q_j/(\log x)^2)$ exceptions. From equation (4) we have

$$\pi(y; d, 1) > \frac{\text{li } y}{2\varphi(d)} \quad (z/2 < y \leq z)$$

for all $Q_j < d \leq 2Q_j$ with at most $O(Q_j/(\log x)^2)$ exceptions. A little thought with the dichotomy of z between the interval $[x^{3/4}, x]$ leads to the fact

$$\pi(y; d, 1) > \frac{\text{li } y}{2\varphi(d)} > \frac{y}{3\varphi(d) \log y} \quad (\forall x^{3/4} \leq y \leq x) \quad (5)$$

for all $Q_j < d \leq 2Q_j$ except for $O(Q_j/\log x)$ exceptions. Let S_j be the set of all integers $Q_j < d \leq 2Q_j$ such that equation (5) hold. Thus, from the analysis above and equation (3), (5) we conclude that

$$\begin{aligned} \sum_{n \leq x} \omega^*(n)^2 &\geq \frac{x}{9} \sum_{0 \leq j \leq \lfloor \frac{\log x}{13 \log 2} \rfloor} \sum_{d \in S_j} \varphi(d) \left(\int_{x^{3/4}}^x \frac{1}{\varphi(d)t \log t} dt \right)^2 + O(x) \\ &\gg x \sum_{0 \leq j \leq \lfloor \frac{\log x}{13 \log 2} \rfloor} \sum_{d \in S_j} \frac{1}{\varphi(d)} \gg x \log x. \end{aligned}$$

□

It is worth here mentioning that we have the following corollary

$$\sum_{p, q \leq x} \frac{1}{[p-1, q-1]} \asymp \log x$$

due to (see [4, page 6, last line])

$$\sum_{n \leq x} \omega^*(n)^2 = \sum_{p, q \leq x} \frac{x}{[p-1, q-1]} + O(x).$$

ACKNOWLEDGMENTS

The author is supported by the Natural Science Foundation of Jiangsu Province of China, Grant No. BK20210784, China Postdoctoral Science Foundation, Grant No. 2022M710121. He is also supported by foundation numbers JSSCBS20211023 and YZ-LYJF2020PHD051.

REFERENCES

- [1] H. Davenport, *Multiplicative Number Theory*, Second edition, Graduate Texts in Mathematics 74, Springer-Verlag, New York, 1980.
- [2] P. Erdős, K. Prachar, *Über die Anzahl der Lösungen von $[p-1, q-1] \leq x$* , Monatsh. Math. **59** (1955), 318–319.
- [3] G. H. Hardy, S. Ramanujan, *The normal number of prime factors of a number n* , Quart. J. Math. **48** (1917), 76–92.
- [4] R. M. Murty, V. K. Murty, *A variant of the Hardy–Ramanujan theorem*, Hardy–Ramanujan J. **44** (2021), 32–40.
- [5] K. Prachar, *Über die Anzahl der Teiler einer natürlichen Zahl, welche die Form $p-1$ haben*, Monatsh. Math. **59** (1955), 91–97.
- [6] P. Turán, *On a theorem of Hardy and Ramanujan*, J. London Math. Soc. **9** (1934), 274–276.

(YUCHEN DING) SCHOOL OF MATHEMATICAL SCIENCE, YANGZHOU UNIVERSITY, YANGZHOU 225002, PEOPLE'S REPUBLIC OF CHINA

Email address: ycding@yzu.edu.cn