

Generalized quantum phase spaces for the κ -deformed extended Snyder model

Jerzy Lukierski¹, Stjepan Meljanac², Salvatore Mignemi³, Anna Pachol⁴

¹Institute of Theoretical Physics, Wrocław University, pl. Maxa Borna 9, 50-205 Wrocław, Poland

² Division of Theoretical Physics, Rudjer Bošković Institute, Bijenička c.54, HR-10002 Zagreb, Croatia

³ Dipartimento di Matematica, Università di Cagliari via Ospedale 72, 09124 Cagliari, Italy and INFN, Sezione di Cagliari

⁴ Department of Microsystems, University of South-Eastern Norway, Campus Vestfold, Raveien 215, 3184 Borre, Norway

Abstract

We describe, in an algebraic way, the κ -deformed extended Snyder models, that depend on three parameters β, κ and λ , which in a suitable algebra basis are described by the de Sitter algebras $o(1, N)$. The commutation relations of the algebra contain a parameter λ , which is used for the calculations of perturbative expansions. For such κ -deformed extended Snyder models we consider the Heisenberg double with dual generalized momenta sector, and provide the respective generalized quantum phase space depending on three parameters mentioned above. Further, we study for these models an alternative Heisenberg double, with the algebra of functions on de Sitter group. In both cases we calculate the formulae for the cross commutation relations between generalized coordinate and momenta sectors, at linear order in λ . We demonstrate that in the commutators of quantum space-time coordinates and momenta of the quantum-deformed Heisenberg algebra the terms generated by κ -deformation are dominating over β -dependent ones for small values of λ .

1 Introduction

The non-commutative (NC) quantum space-times, where the coordinates are elements of NC quantum space-time algebra, have been considered as a tool for the description of quantum gravity (QG) (see, e.g. [1], [2]), which unifies two basic theories in physics: general relativity and quantum mechanics. One of the first examples of quantum space-time was proposed by Snyder already in 1947 [3]. Other quantum space-time models were studied in the 1990's (e.g. κ -deformed relativistic theories [4]-[7]). Due to the increasing interest in QG, the framework of quantum symmetries described by the theory of quantum groups was developed. The NC structure of space-time at the Planck scale appeared as a way to describe QG at very short distances and led to the description of quantum-mechanical and quantum-deformed relativistic phase spaces. It appeared that these NC structures of quantum space-times and quantum relativistic symmetries are associated with the formalism of Hopf algebras and the case of quantum phase spaces has been generalized to the Hopf algebroids framework (see, e.g. [8], [9]).

In this paper we find the quantum-deformed relativistic phase space for the κ -deformed extended Snyder model [10]¹ which will be obtained by the Heisenberg double construction. The Heisenberg double, which constitutes a Hopf algebroid, is defined within the Hopf algebraic

¹We use “ κ -extended Snyder model” as a short notation for the λ -dependent κ -deformed extended Snyder model.

framework and can be considered as providing the way to define quantum-deformed phase spaces in the presence of QG effects.

The κ -deformed Snyder model, proposed firstly in [11], [12], unifies the two perhaps best known models which describe the NC relativistic space-time coordinates \hat{x}_i :²

i) the Snyder model:

$$[\hat{x}_i, \hat{x}_j] = i\beta M_{ij} \quad (1)$$

where β is a parameter with dimension of length square and M_{ij} are the generators of Lorentz transformations;

ii) κ -Minkowski quantum space-time:

$$[\hat{x}_i, \hat{x}_j] = i(a_i \hat{x}_j - a_j \hat{x}_i) \quad (2)$$

where a_i is a constant vector multiplied by the parameter κ^{-1} with dimension of length. Both Snyder and κ -deformed Snyder models lead to non-associativity and non-coassociativity [13], [14]. In order to avoid problems with non-associativity, an alternative Snyder model, in which Snyder space is a subspace of a larger non-commutative space was proposed in [14] for the 3 dimensional Euclidean Snyder space, and its generalization to the extended Snyder model was proposed in [15]³. If one uses Latin indices the algebra describing the λ -dependent κ -extended Snyder model looks as follows [15]

$$[\hat{x}_i, \hat{x}_j] = i\lambda\beta\hat{x}_{ij}, \quad [\hat{x}_{ij}, \hat{x}_{kl}] = i\lambda(\eta_{ik}\hat{x}_{jl} - \eta_{il}\hat{x}_{jk} - \eta_{jk}\hat{x}_{il} + \eta_{jl}\hat{x}_{ik}), \quad (4)$$

$$[\hat{x}_{ij}, \hat{x}_k] = i\lambda(\eta_{ik}\hat{x}_j - \eta_{jk}\hat{x}_i), \quad (5)$$

where $\hat{x}_i = \sqrt{\beta}\hat{x}_{iN}$.

Unification of κ -Minkowski and extended Snyder space-times was proposed in [10] generalizing the extended Snyder model [15] by including the κ -Minkowski algebra terms (2) using orthogonal algebra with metric tensor g [16], [17]. The main step for achieving this unification was introducing particular modification of constant metric tensor instead of Minkowski metric.

In this paper we introduce the generalization of quantum-mechanical phase space corresponding to the κ -extended Snyder model. The phase spaces with NC space-time coordinates, have been already considered for Snyder model and for models covariant under the κ -deformed Poincaré algebra: the generalized phase spaces containing the κ -Minkowski NC space-time were considered in [18] - [23] and the phase space for the Snyder model in, e.g. [3], [24], [25] as well as for the extended Snyder model in [26]. Quantum space-times and deformed phase spaces for Snyder and Yang models were discussed in [27], [28] together with their extensions to supersymmetric models, see also [29].

Here we propose the description of the corresponding deformed phase spaces for the κ -extended Snyder model. The unification of the extended Snyder and κ -Minkowski models proposed in [10], [30] is realized in the framework of associative and coassociative Hopf algebra, what permits to apply the Heisenberg double construction that is considered in the present paper. The resulting generalized phase spaces offer interesting insights into the κ -extended Snyder model, which is described by the superposition of two quantum deformations with a further parameter λ .

²N-dimensional Latin indices include zero, which describes time-like dimension, i.e. $i, j = 0, \dots, N - 1$.

³The N -dimensional extended Snyder algebra [14], after introducing the λ -dependence [15], takes the form of an N -dimensional de Sitter algebra given by ($\mu = 0, 1 \dots N$)

$$[\hat{x}_{\mu\nu}, \hat{x}_{\rho\sigma}] = i\lambda(\eta_{\mu\rho}\hat{x}_{\nu\sigma} - \eta_{\nu\rho}\hat{x}_{\mu\sigma} + \eta_{\nu\sigma}\hat{x}_{\mu\rho} - \eta_{\mu\sigma}\hat{x}_{\nu\rho}) \quad (3)$$

where $\hat{x}_{\mu\nu}$ are the standard $o(1, N)$ generators. A specific new feature of such formulation of extended Snyder algebra introduced in [15], is the appearance of a dimensionless parameter λ such that for $\lambda \rightarrow 0$ (classical limit) the extended Snyder algebra reduces to the Abelian algebra. The extended Snyder algebra (3) contains the generators defining Snyder quantum space-time coordinates \hat{x}_i as well as tensorial coordinates \hat{x}_{ij} (see (4,5)).

2 κ -deformed extended Snyder model

Let us recall the algebra corresponding to the κ -extended Snyder model, which was introduced in [10] as the extended unified κ -Minkowski Snyder model. We shall denote it as $o(1, N; g)$. It is defined by the following Lie-algebraic set of commutation relations:

$$[\hat{X}_{\mu\nu}, \hat{X}_{\rho\sigma}] = i\lambda(g_{\mu\rho}\hat{X}_{\nu\sigma} - g_{\nu\rho}\hat{X}_{\mu\sigma} + g_{\nu\sigma}\hat{X}_{\mu\rho} - g_{\mu\sigma}\hat{X}_{\nu\rho}). \quad (6)$$

The metric $g \equiv (g_{\mu\nu})$ has the form:

$$g = \begin{pmatrix} -1 & 0 & \dots & 0 & g_0 \\ 0 & 1 & \dots & 0 & g_1 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & g_{N-1} \\ g_0 & g_1 & \dots & g_{N-1} & g_N \end{pmatrix}$$

with $\det g = -g_0^2 + \sum_{i=1}^{N-1} g_i^2 - g_N$. If we rewrite the generators $\hat{X}_{\mu\nu}$ as \hat{X}_{ij} and $\hat{X}_{kN} = \kappa\hat{X}_k$ where κ is a new mass-like parameter, and rewrite the metric as follows

$$g_{ij} = \eta_{ij}, \quad g_{iN} = \kappa a_i, \quad g_{NN} = g_N = \kappa^2 \beta$$

the algebra (6) splits to the following set of relations

$$[\hat{X}_i, \hat{X}_j] = i\lambda(a_i\hat{X}_j - a_j\hat{X}_i + \beta\hat{X}_{ij}), \quad (7)$$

$$[\hat{X}_{ij}, \hat{X}_k] = i\lambda(\eta_{ik}\hat{X}_j - \eta_{jk}\hat{X}_i + a_j\hat{X}_{ik} - a_i\hat{X}_{jk}), \quad (8)$$

$$[\hat{X}_{ij}, \hat{X}_{kl}] = i\lambda(\eta_{ik}\hat{X}_{jl} - \eta_{jk}\hat{X}_{il} + \eta_{jl}\hat{X}_{ik} - \eta_{il}\hat{X}_{jk}). \quad (9)$$

From the first commutator we see how the Snyder and κ -Minkowski space-time relations are unified. The \hat{X}_i are the NC space-time coordinates, and \hat{X}_{ij} can be interpreted as non-commutative tensorial coordinates.

Moreover, when $g_N = 0$, the relations (6) describe the κ -Minkowski space-time with Lorentz covariance algebra. Alternatively, when $g_0 = \dots = g_{N-1} = 0$ and $g_N = 1$, then the relations (6) reduce to the algebra describing the extended Snyder model. Also note that if $g_{\mu\nu} \rightarrow \eta_{\mu\nu}$ then the algebra (6) becomes a standard orthogonal algebra, which describes the N -dimensional suitably rescaled de Sitter algebra.

The coalgebra sector is a classical one (primitive), i.e.

$$\Delta(\hat{X}_{\mu\nu}) = \Delta_0(\hat{X}_{\mu\nu}), \quad (10)$$

$$\epsilon(\hat{X}_{\mu\nu}) = 0 \text{ and } S(\hat{X}_{\mu\nu}) = -\hat{X}_{\mu\nu} \quad (11)$$

and reduces accordingly to coproducts, counits and antipodes for \hat{X}_k and \hat{X}_{ik} .

If one defines the following change of coordinates:

$$\hat{X}_{\mu\nu} = (O\hat{x}O^T)_{\mu\nu}, \quad g_{\mu\nu} = (O\eta O^T)_{\mu\nu}, \quad (12)$$

with the choice of the matrix

$$O = \begin{pmatrix} 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 1 & 0 \\ -g_0 & g_1 & \dots & g_{N-1} & \rho \end{pmatrix} \quad (13)$$

where $\rho = \sqrt{g_N - g_k g_k} = \sqrt{-\det g}$, one can show that the algebra (6) reduces to (see also ³⁾)

$$[\hat{x}_{\mu\nu}, \hat{x}_{\rho\sigma}] = i\lambda(\eta_{\mu\rho}\hat{x}_{\nu\sigma} - \eta_{\nu\rho}\hat{x}_{\mu\sigma} + \eta_{\nu\sigma}\hat{x}_{\mu\rho} - \eta_{\mu\sigma}\hat{x}_{\nu\rho}), \quad (14)$$

where $\eta_{\mu\nu} = (-1, 1, \dots, 1)$ and $g_N = 1$. This algebra has been discussed in [14], [15], as the extended Snyder algebra. We recall (see ³⁾ and (4), (5)) that the algebra (14) reduces to the N -dimensional Snyder space extended by the Lorentz algebra. We add that Heisenberg double for the extended Snyder model (14) was studied in [26].

Due to the form of the matrix O we have the following relations between two sets of coordinates:

$$\hat{X}_i = \rho\hat{x}_i + a_j\hat{x}_{ij}, \quad \hat{X}_{ij} = \hat{x}_{ij}. \quad (15)$$

Both sets of the coordinates $\hat{X}_{\mu\nu}$ and $\hat{x}_{\mu\nu}$ describe NC extended space-time and are related via relations (15).

In the classical limit ($\lambda \rightarrow 0$) the generators of the algebras (6) and (14) reduce to the commutative (Abelian) ones [30] $\hat{X}_{\mu\nu} \rightarrow X_{\mu\nu}$ with $[X_{\mu\nu}, X_{\rho\sigma}] = 0$; ($\hat{x}_{\mu\nu} \rightarrow x_{\mu\nu}$ with $[x_{\mu\nu}, x_{\rho\sigma}] = 0$). These Abelian coordinates are related with each other via the matrix O given by (13)

$$X_{\mu\nu} = (OxO^T)_{\mu\nu},$$

i.e. explicitly:

$$X_i = \rho x_i + a_j x_{ij}, \quad X_{ij} = x_{ij},$$

where $X_i = \frac{1}{\kappa} X_{iN}$ and $x_i = \frac{1}{\kappa} x_{iN}$. One can say that the algebra (6), with the generators $\hat{X}_{\mu\nu}$ can be seen as the deformation of the underlying commutative space described by $X_{\mu\nu}$, with λ as the deformation parameter.

For the commutative extended spacetime coordinates $X_{\mu\nu}$ and $x_{\mu\nu}$ we can introduce the extended momenta $(P^{\mu\nu}, p^{\mu\nu})$ which can be realized in a standard way as $P^{\mu\nu} = -i\frac{\partial}{\partial X_{\mu\nu}}$ and $p^{\mu\nu} = -i\frac{\partial}{\partial x_{\mu\nu}}$. In this way one can introduce two copies of the generalized Heisenberg algebra as unital, associative algebras generated by the extended coordinates and momenta, with the tensorial coordinates antisymmetric under the exchange $\mu \leftrightarrow \nu$. The following commutation relations are valid (we put $\hbar = 1$)

$$[X_{\mu\nu}, X_{\alpha\beta}] = 0 = [P^{\mu\nu}, P^{\alpha\beta}], \quad [X_{\mu\nu}, P^{\rho\sigma}] = i(\delta_\mu^\rho \delta_\nu^\sigma - \delta_\mu^\sigma \delta_\nu^\rho), \quad (16)$$

and similarly

$$[x_{\mu\nu}, x_{\alpha\beta}] = 0 = [p^{\mu\nu}, p^{\alpha\beta}], \quad [x_{\mu\nu}, p^{\rho\sigma}] = i(\delta_\mu^\rho \delta_\nu^\sigma - \delta_\mu^\sigma \delta_\nu^\rho), \quad (17)$$

where Greek indices are raised and lowered by the metric $g_{\mu\nu}$.

One can introduce, as well, the formulae for the momenta with a single index (i.e $P^i = \kappa P^{iN}$, $p^i = \kappa p^{iN}$) as dual to space-time coordinates X_i and x_i ; Latin indices are raised and lowered by the flat metric η_{ij} . We also have analogous relations between the two types of momenta (16) and (17)

$$p_i = \rho P_i, \quad p_{ij} = P_{ij} - a_i P_j + a_j P_i, \quad (18)$$

with the relation $P^{\mu\nu} = \left((O^{-1})^T p (O^{-1}) \right)^{\mu\nu}$. For more details about these coordinates and momenta we refer the reader to [10], [30] where this model was introduced and studied.

In order to discuss the phase spaces associated with the κ -deformed extended Snyder model (6) one can use the Heisenberg double construction method.

2.1 Commutative momenta for the κ -extended Snyder model and their coproducts

One can introduce the Abelian momenta, dual to the coordinates $\hat{X}_{\mu\nu}$ which describe the algebra $o(1, N; g)$ (see (6)), by deforming the canonically conjugate Abelian momenta $P^{\mu\nu}$. The technique to calculate the coproducts ΔP_i and ΔP_{ij} was proposed in [12], [13]. The commutation relations for $P^i = \kappa P^{iN}$, P^{ij} remain unchanged:

$$[P_i, P_j] = 0, \quad [P_{ik}, P_{jl}] = 0, \quad [P_i, P_{jl}] = 0. \quad (19)$$

while the coalgebraic sector of the momentum generators [30] looks as follows⁴:

$$\begin{aligned} \Delta P_i &= P_i \otimes 1 + 1 \otimes P_i \\ &+ \lambda[-c_1(P_j \otimes P_{ij} + P_j \otimes P_i a_j) \\ &+ (1 - c_1)(P_{ij} \otimes P_j + P_i \otimes P_j a_j) + (2c_1 - 1)P_j \otimes P_j a_i] + O(\lambda^2) \end{aligned} \quad (20)$$

$$\begin{aligned} \Delta P_{ij} &= P_{ij} \otimes 1 + 1 \otimes P_{ij} \\ &+ \frac{\lambda}{2}[-\beta P_i \otimes P_j - (P_{ik} \otimes P_{jk} + a_k(P_i \otimes P_{jk} - P_{jk} \otimes P_i)) \\ &- (2c_1 - 1)(P_k \otimes P_{jk} + P_k \otimes P_j a_k + P_{jk} \otimes P_k + P_j \otimes P_k a_k) a_i - (i \leftrightarrow j)] + O(\lambda^2), \end{aligned} \quad (21)$$

$$S(P_i) = -P_i + \lambda(1 - 2c_1)(P_j P_{ij} + a_k P_k P_i - P_k P_k a_i), \quad (22)$$

$$S(P_{ij}) = -P_{ij} - \lambda(2c_1 - 1)(a_i P_{jk} P_k + a_i P_j a_k P_k - (i \leftrightarrow j)), \quad (23)$$

$$\epsilon(P_i) = \epsilon(P_{ij}) = 0. \quad (24)$$

where a new parameter c_1 depends on the realization (see Sec. 3 in [30]), and coproducts are presented up to the first order in the deformation parameter λ . The coproducts (20), (21) correspond to the so-called generic realization [30], and define the momentum sector of the κ -extended Snyder model as parametrized by λ, β, a_i and c_1 . Such parametrization occurs when the coproducts are written up to the first order in λ ; in higher orders of λ additional parameters appear (see Sec.3 in [30] for more details). For the concrete Weyl realization [10], one should set $c_1 = \frac{1}{2}$ and for the natural realization (i.e. with classical algebra basis) $c_1 = 0$.

We postulate the standard duality relation:

$$\langle P_j, \hat{X}_i \rangle = -i\eta_{ij}, \quad (25)$$

$$\langle P_k, \hat{X}_{ij} \rangle = 0, \quad (26)$$

$$\langle P_{kl}, \hat{X}_i \rangle = 0, \quad (27)$$

$$\langle P_{kl}, \hat{X}_{ij} \rangle = -i(\eta_{ik}\eta_{jl} - \eta_{jk}\eta_{il}). \quad (28)$$

One can check that from the duality relation between the products in the algebra generated by \hat{X} and the coproducts in the coalgebra generated by P , the following compatibility conditions hold

$$\langle b_{(1)}, a \rangle \langle b_{(2)}, a' \rangle = \langle b, a \cdot a' \rangle, \quad (29)$$

$$\langle b, a_{(1)} \rangle \langle b', a_{(2)} \rangle = \langle b \cdot b', a \rangle. \quad (30)$$

⁴Note that there was a typo in sign in term $P_{jk} \otimes P_i a_k$ in the coproduct of ΔP_{ij} in [30].

3 Generalized quantum phase space from the Heisenberg double

We introduce the left Hopf action \triangleright of momenta on coordinates defined by the formula $P \triangleright \hat{X} = \langle P, \hat{X}_{(2)} \rangle \hat{X}_{(1)}$, and recall that we use the Sweedler notation for the coproduct. From (25)-(28) it follows immediately that $P_j \triangleright \hat{X}_i = -i\eta_{ij}$, $P_k \triangleright \hat{X}_{ij} = 0$, $P_{kl} \triangleright \hat{X}_i = 0$, $P_{kl} \triangleright \hat{X}_{ij} = -i(\eta_{ik}\eta_{jl} - \eta_{jk}\eta_{il})$. The corresponding Heisenberg double commutators follow from the cross product construction:

$$\left[P, \hat{X} \right] = \hat{X}_{(1)} \langle P_{(1)}, \hat{X}_{(2)} \rangle P_{(2)} - \hat{X}P. \quad (31)$$

written shortly without the indices. Doing the calculation explicitly and using the coproducts for momenta (20)-(21), case by case, we obtain:

$$\left[P_j, \hat{X}_i \right] = -i\eta_{ji} + i\lambda[c_1(P_{ji} + P_j a_i) - (1 - c_1)\eta_{ij}P_r a_r - (2c_1 - 1)P_i a_j] + O(\lambda^2), \quad (32)$$

$$\left[P_j, \hat{X}_{is} \right] = -i\lambda(1 - c_1)(\eta_{ij}P_s - \eta_{sj}P_i) + O(\lambda^2), \quad (33)$$

$$\begin{aligned} \left[P_{ij}, \hat{X}_k \right] &= \frac{i\lambda}{2}(\beta(\eta_{ik}P_j - \eta_{jk}P_i) + (\eta_{ik}P_{jl} - \eta_{jk}P_{il})a_l \\ &\quad + (2c_1 - 1)[a_i(P_{jk} + P_j a_k + \eta_{jk}P_l a_l) - a_j(P_{ik} + P_i a_k + \eta_{ik}P_l a_l)]) + O(\lambda^2), \end{aligned} \quad (34)$$

$$\begin{aligned} \left[P_{ij}, \hat{X}_{st} \right] &= -i(\eta_{si}\eta_{tj} - \eta_{ti}\eta_{sj}) \\ &\quad + \frac{i\lambda}{2}[\eta_{si}(P_{jt} + P_j a_t - (2c_1 - 1)P_t a_j) - \eta_{ti}(P_{js} + P_j a_s - (2c_1 - 1)P_s a_j) \\ &\quad - \eta_{sj}(P_{it} + P_i a_t - (2c_1 - 1)P_t a_i) + \eta_{tj}(P_{is} + P_i a_s - (2c_1 - 1)P_s a_i)] + O(\lambda^2). \end{aligned} \quad (35)$$

The terms linear in λ in the commutation relations $\left[P_j, \hat{X}_i \right]$, $\left[P_j, \hat{X}_{is} \right]$, and $\left[P_{ij}, \hat{X}_{st} \right]$ do not include any terms proportional to β parameter.

i) Reduction to the κ -Minkowski case

When $g_N = 0$ the cross commutators obtained above are reduced to the particular case of κ -Minkowski relations. Further one can compare these particular relations with the results obtained in the literature (see, e.g. in [21], [22]) where the Heisenberg double construction was investigated for the κ -Minkowski space-time and quantum symmetry described by the κ -Poincaré algebra (in Snyder model the quantum symmetry is linked with the de Sitter algebra).

In linear order of λ the commutator (32) does not depend on the β parameter and the resulting cross commutation relations (32) describing basic quantum-deformed Heisenberg algebra relation contain only the terms generated by κ -deformation. We know however from the standard Snyder model (see [3]) that the term linear in β is bilinear in the momenta. Because terms bilinear in P_i are as well bilinear in λ , the parameter β can contribute only at the second perturbative order in λ . One can state that for small λ the terms generated by κ -deformation dominate over the β -dependent terms, but for more precise statement the perturbative λ^2 order terms should also be calculated for the relation (32).

We add that all of the generalized phase space relations, including the tensorial coordinates and momenta, depend on the c_1 parameter. In particular for $c_1 = 0$, we obtain in place of (32)

$$\left[P_k, \hat{X}_i \right] = -i\eta_{ki}(1 + \lambda a_j P_j) + i\lambda P_i a_k + O(\lambda^2). \quad (36)$$

Such commutator would correspond to the so-called classical basis of κ -Poincaré (named also the natural realization) [31], [32], [33]. For $c_1 = \frac{1}{2}$ we get:

$$\left[P_k, \hat{X}_i \right] = -i\eta_{ki} \left(1 + \frac{1}{2} \lambda a_j P_j \right) - \frac{i}{2} \lambda (P_{ik} - P_k a_i) + O(\lambda^2) \quad (37)$$

which is the result corresponding to the so-called Weyl realization of κ -Poincaré algebra [31].

ii) Reduction to the extended Snyder model

We can reduce the κ -dependent terms in commutation relations (32)-(35) and compare such results with the results obtained from the Heisenberg double of the extended Snyder model [26]. When $g_0 = \dots = g_{N-1} = 0$ and $g_N = 1$, the algebra (6), after the change of variables (15), (18), provides the extended Snyder algebra. The phase space relation calculated in (32)-(35) are reduced to the following ones

$$[p_k, \hat{x}_i] = -i\eta_{ki} + i\lambda c_1 p_{ki} + O(\lambda^2), \quad (38)$$

$$[p_k, \hat{x}_{ij}] = -i\lambda (1 - c_1) (\eta_{ik} p_j - \eta_{jk} p_i) + O(\lambda^2), \quad (39)$$

$$[p_{ij}, \hat{x}_k] = i\frac{\lambda}{2} \beta (\eta_{ik} p_j - \eta_{jk} p_i) + O(\lambda^2), \quad (40)$$

$$[p_{ij}, \hat{x}_{st}] = -i (\eta_{si} \eta_{tj} - \eta_{ti} \eta_{sj}) + i\frac{\lambda}{2} [(\eta_{si} p_{jt} - \eta_{sj} p_{it}) - (\eta_{ti} p_{js} - \eta_{tj} p_{is})] + O(\lambda^2). \quad (41)$$

The cross commutation relations in the generalized phase space obtained by the Heisenberg double method for the extended Snyder model were calculated in [26]. They do agree with the above results if $c_1 = 1/2$.

4 Another Heisenberg double for the κ -extended Snyder model

Using the algebra (6) describing κ -extended Snyder model, also other Heisenberg double construction can be considered. If we introduce the algebra of functions generated by dual de Sitter group matrices $\Lambda_{\alpha\beta}$: $\{\Lambda_{\alpha\beta} : [\Lambda_{\alpha\beta}, \Lambda_{\mu\nu}] = 0 : \Lambda^T g \Lambda = g\}$ we should postulate

$$\Delta(\Lambda_{\rho\sigma}) = \Lambda_{\rho\alpha} \otimes \Lambda_{\alpha\sigma}; \quad \epsilon(\Lambda_{\rho\sigma}) = g_{\rho\sigma} \quad ; S(\Lambda_{\rho\sigma}) = (\Lambda^{-1})_{\rho\sigma} = \Lambda_{\sigma\rho}. \quad (42)$$

One can also introduce the matrices $\tilde{\Lambda}_{\alpha\beta}$ which are related with the above group elements via the map $\Lambda_{\rho\sigma} = \left(O \tilde{\Lambda} O^T \right)_{\rho\sigma}$ ⁵, ($\alpha, \beta = 0, 1, \dots, N$). The basic duality relation is given by:

$$\langle \Lambda_{\rho\sigma}, \hat{X}_{\mu\nu} \rangle = -i\lambda (g_{\rho\mu} g_{\sigma\nu} - g_{\rho\nu} g_{\sigma\mu}). \quad (43)$$

We consider the following left Hopf action \triangleright :

$$\Lambda_{\rho\sigma} \triangleright \hat{X}_{\mu\nu} = \langle \Lambda_{\rho\sigma}, \hat{X}_{\mu\nu(2)} \rangle \hat{X}_{\mu\nu(1)} = g_{\rho\sigma} \hat{X}_{\mu\nu} - i\lambda (g_{\rho\mu} g_{\sigma\nu} - g_{\rho\nu} g_{\sigma\mu}) \quad (44)$$

⁵In this case we have: $\{\tilde{\Lambda}_{\alpha\beta} : [\tilde{\Lambda}_{\alpha\beta}, \tilde{\Lambda}_{\mu\nu}] = 0 : \tilde{\Lambda}^T \eta \tilde{\Lambda} = \eta\}$,

$$\Delta(\tilde{\Lambda}_{\rho\sigma}) = \tilde{\Lambda}_{\rho\alpha} \otimes \tilde{\Lambda}_{\alpha\sigma}; \quad \epsilon(\tilde{\Lambda}_{\rho\sigma}) = \eta_{\rho\sigma} \quad ; S(\tilde{\Lambda}_{\rho\sigma}) = (\tilde{\Lambda}^{-1})_{\rho\sigma} = \tilde{\Lambda}_{\sigma\rho}.$$

Such algebra was also used in [26] in our studies of the Heisenberg double for the extended Snyder model.

and we obtain the cross commutation relations which are defined by the Heisenberg double method

$$\begin{aligned} [\Lambda_{\rho\sigma}, \hat{X}_{\mu\nu}] &= \hat{X}_{\mu\nu(1)} < \Lambda_{\rho\sigma(1)}, \hat{X}_{\mu\nu(2)} > \Lambda_{\rho\sigma(2)} - \hat{X}_{\mu\nu} \Lambda_{\rho\sigma} \\ &= -i\lambda(g_{\rho\mu}\Lambda_{\nu\sigma} - g_{\rho\nu}\Lambda_{\mu\sigma}). \end{aligned} \quad (45)$$

If we recall the relations $\hat{X}_{kN} = \kappa\hat{X}_k$ and $g_{ij} = \eta_{ij}$, $g_{iN} = \kappa a_i$, $g_{NN} = g_N = \kappa^2\beta$, we get from (45) the cross commutation relations between the dual group elements and quantum algebra generators

$$[\Lambda_{jk}, \hat{X}_i] = -\frac{i\lambda}{\kappa}(g_{ji}\Lambda_{Nk} - g_{jN}\Lambda_{ik}) = -\frac{i\lambda}{\kappa}(\eta_{ji}\Lambda_{Nk} - \kappa a_j\Lambda_{ik}), \quad (46)$$

$$[\Lambda_{jN}, \hat{X}_i] = -\frac{i\lambda}{\kappa}(g_{ji}\Lambda_{NN} - g_{jN}\Lambda_{iN}) = -\frac{i\lambda}{\kappa}(\eta_{ji}\Lambda_{NN} - \kappa a_j\Lambda_{iN}), \quad (47)$$

$$[\Lambda_{Nk}, \hat{X}_i] = -i\lambda(a_i\Lambda_{Nk} - \kappa\beta\Lambda_{ik}), \quad (48)$$

$$[\Lambda_{NN}, \hat{X}_i] = -i\lambda(a_i\Lambda_{NN} - \kappa\beta\Lambda_{iN}). \quad (49)$$

For the tensorial coordinates \hat{X}_{ij} we obtain the cross relations:

$$[\Lambda_{lk}, \hat{X}_{ij}] = -i\lambda(\eta_{li}\Lambda_{jk} - \eta_{lj}\Lambda_{ik}), \quad (50)$$

$$[\Lambda_{lN}, \hat{X}_{ij}] = -i\lambda(\eta_{li}\Lambda_{jN} - \eta_{lj}\Lambda_{iN}), \quad (51)$$

$$[\Lambda_{Nk}, \hat{X}_{ij}] = -i\lambda\kappa(a_i\Lambda_{jk} - a_j\Lambda_{ik}), \quad (52)$$

$$[\Lambda_{NN}, \hat{X}_{ij}] = -i\lambda\kappa(a_i\Lambda_{jN} - a_j\Lambda_{iN}). \quad (53)$$

Note that $\Lambda_{ij} = \tilde{\Lambda}_{ij}$ ⁶. One can check explicitly that the duality $< \Lambda_{\rho\sigma}, \hat{X}_{\mu\nu} >$ given above reduces to

$$< \tilde{\Lambda}_{\rho\sigma}, \hat{x}_{\mu\nu} > = -i\lambda(\eta_{\rho\mu}\eta_{\sigma\nu} - \eta_{\rho\nu}\eta_{\sigma\mu}) \quad (56)$$

for the extended Snyder model (i.e. when $g_{Ni} = g_0 = \dots = g_{N-1} = 0$, $g_{ij} = \eta_{ij}$, $g_{NN} = g_N = \kappa^2\beta$). Therefore, all the cross commutation relations (46)-(53) are the same as the ones obtained in [26] where the extended Snyder model was investigated.

In the reduction of the above results to the κ -Minkowski case ($g_N = 0$) only two relations (48), (49) depend on β and we easily see the result of reduction. Further, focusing on the relation (46), after putting $a_j = \frac{1}{\kappa}\delta_j^0$ (i.e. assuming a time-like κ -deformation) we obtain the following covariance relations

$$[\Lambda_{jk}, \hat{X}_i] = -\frac{i}{\kappa}\lambda(\eta_{ji}\Lambda_{Nk} - \delta_j^0\Lambda_{ik}). \quad (57)$$

5 Discussion and outlook

For the κ -extended Snyder model introduced in [10] we constructed the generalized quantum phase space, which depends on three parameters, β , κ and λ . The Hopf algebra describing

⁶Footnote ⁵ can be supplemented by the following relations obtained from (46), (50):

$$[\tilde{\Lambda}_{jk}, \hat{X}_i] = -\frac{i\lambda}{\kappa}(\eta_{ji}\tilde{\Lambda}_{Nk} - \kappa a_j\tilde{\Lambda}_{ik}), \quad (54)$$

$$[\tilde{\Lambda}_{lk}, \hat{X}_{ij}] = -i\lambda(\eta_{li}\tilde{\Lambda}_{jk} - \eta_{lj}\tilde{\Lambda}_{ik}). \quad (55)$$

the quantum symmetries of the model is coassociative, hence one can use the Heisenberg double construction for the correct description of the respective generalized quantum phase space. Since this model contains Snyder ($\beta \neq 0$) and κ -Minkowski sectors ($\kappa \neq 0$), one can discuss how the generalized quantum phase space can be reduced to these special cases. The algebraic relations determining the generalized phase space for the κ -extended Snyder model have been calculated to the first order in the λ parameter, and for most commutation relations to such order we get only the formulae modified by κ -deformation. Our algebra also includes the orthogonal (de Sitter) algebra generators, which are described by tensorial coordinates.

The quantum group describing quantum symmetries of the κ -extended Snyder model is built up from deformed coproducts of momenta [30] and undeformed Lorentz algebra relations. It would be interesting to see what realizations of κ -Poincaré, after the interpretation of β as the inverse square of de Sitter radius ($\beta \sim R^{-2}$), could be obtained by the quantum Inonu-Wigner contraction procedure.

The Heisenberg double for the κ -Poincaré quantum group leads to the κ -deformed phase space that was investigated in different bases for the κ -Poincaré Hopf algebra [18]-[22], [30]. In κ -extended Snyder model with different quantum symmetry group and different coalgebra sector, the comparison between these two cases and the present model involving tensorial coordinates should still be discussed.

Further, one of the tasks which would be interesting to investigate is the Hopf algebroid providing the generalized quantum phase spaces of the κ -extended Snyder model. Hopf algebroids describing the quantum phase spaces with NC space-time coordinates have been studied in the literature recently [19], [34]-[38], and it should be recalled that Heisenberg double construction provides a natural example of the Hopf bialgebroid structure [8].

It can finally be added that the parameters β , κ and λ are not treated here as genuine quantum deformations parameters introducing quantum deformation determined by classical r -matrices, which satisfy classical or modified Yang Baxter equations. These parameters appear in the procedure of changing the $o(1, N)$ basis, which effectively leads to the modified de Sitter relations and noncanonical modified de Sitter metric $g_{\mu\nu}$. The parameters β, κ and λ appear as determining the algebra basis and in this way we introduce physical parameters (β is related with de Sitter radius, κ usually is linked with Planck mass, and the parameter λ is related with the Planck constant). Thus, in order to describe the κ -extended Snyder model we only redefine the standard basis of classical de Sitter algebra, this mathematically rather trivial operation leads to results which might be significant in physical applications.

Acknowledgements

S. Meljanac, S. Mignemi and A. Pachoł would like to acknowledge the support of the European Cooperation in Science and Technology COST Action CA18108. S. Mignemi acknowledges support from GNFM. J. Lukierski and A. Pachoł were supported by Polish NCN grant 2017/27/B/ST2/01902; J. Lukierski would like also to thank A. Borowiec and M. Woronowicz for valuable comments.

References

- [1] S. Doplicher, F. Fredenhagen, J.E. Roberts, *Comm. Math. Phys.* 172, 187 (1995) [arXiv:hep-th/0303037];
- [2] S.Majid, “Foundations of quantum group theory”, Cambridge Univ. Press. (1995);
- [3] H. S. Snyder, *Phys.Rev.* 71, 38, 0031-899 (1947);

- [4] J. Lukierski, A. Nowicki, H. Ruegg, V.N. Tolstoy, Phys.Lett. B264, 331 (1991);
- [5] S. Majid, H. Ruegg, Phys. Lett. B334, 348 (1994) [arXiv:hep-th/9405107];
- [6] J. Madore, “An Introduction to Noncommutative Differential Geometry and its Physical Applications”, CUP (1995);
- [7] P. Podles, S.L. Woronowicz, Comm. Math.Phys. 178, 61 (1996);
- [8] J.H. Lu, Int. J. Math. 7, 47–70 (1996)[arXiv:q-alg/9505024];
- [9] P.Xu, Comm.Math.Phys. 216,539 (2001) [arXiv:math.QA/9965192];
- [10] S. Meljanac, S. Mignemi, Phys. Lett. B 814, 136117 (2021) [arXiv:2101.05275];
- [11] S. Meljanac, D. Meljanac, A. Samsarov, M. Stojić (2009) [arXiv:0909.1706];
- [12] S. Meljanac, D. Meljanac, A. Samsarov, M. Stojić, Phys.Rev. D83, 065009 (2011) [arXiv:1102.1655];
- [13] M. V. Battisti, S. Meljanac, Phys. Rev. D 82, 024028 (2010) [arXiv:1003.2108];
- [14] F. Girelli, E. Livine, JHEP 1103, 132 (2011) [arXiv:1004.0621];
- [15] S. Meljanac, S. Mignemi, Phys. Rev. D 102, 126011 (2020) [arXiv:2007.13498];
- [16] A. Borowiec, A. Pachoł, Phys. Rev. D79, 045012 (2009); [arXiv:0812.0576];
- [17] A. Borowiec, A. Pachoł, Eur. Phys. J. C 74 (2014) 2812 [arXiv:1311.4499];
- [18] P. Kosinski, P. Maslanka, (1994) [arXiv:hep-th/9411033];
- [19] J. Lukierski, Z. Škoda, M. Woronowicz, Phys. Lett. B 750, 401-406 (2015) [arXiv:1507.02612];
- [20] G. Amelino-Camelia, J. Lukierski, A. Nowicki, Phys. Atomic Nuclei, 61, 1811-1815 (1998) [arXiv:hep-th/9706031];
- [21] J. Lukierski, A. Nowicki, Proc. 21st Intl. Colloquium on Group Theoretical Methods in Physics (Group 24: Physical and mathematical aspects of symmetries, V. K. Dobrev, H. D. Doebner, eds.), Heron Press, Sofia , 186-192 (1997) [arXiv:q-alg/9702003];
- [22] A. Borowiec, A. Pachoł, Theor. Math. Phys. 169 (2), 1620-1628 (2011);
- [23] S. Meljanac, R. Štrajn, SIGMA 18, 022 (2022) [arXiv:2112.12038];
- [24] M. V. Battisti, S. Meljanac, Phys. Rev. D 79, 067505 (2009) [arXiv:0812.3755];
- [25] S. Mignemi, Int.J.Mod.Phys.D 24, 6, 1550043 (2015) [arXiv:1308.0673];
- [26] S. Meljanac, A. Pachoł, Symmetry 13, 1055 (2021) [arXiv:2101.02512];
- [27] J. Lukierski, M. Woronowicz, Phys. Lett. B 824, 136783 (2022) [arXiv:2110.13697];
- [28] J. Lukierski, M. Woronowicz, Proc. of CORFU (2021) [arXiv:2204.07787];
- [29] S. Meljanac, S. Mignemi, Phys. Lett. B 833, 137289 (2022) [arXiv:2206.04772];
- [30] S. Meljanac, S. Mignemi, Phys. Rev. D 104, 086006 (2021) [arXiv:2106.08131];

- [31] S. Kresic-Jurić, S. Meljanac, M. Stojić, Eur. Phys. J. C 51, 229-240 (2007) [arXiv:hep-th/0702215];
- [32] A. Borowiec, A. Pachoł, J. Phys. A 43, 045203 (2010) [arXiv:0903.5251];
- [33] T. Jurić, S. Meljanac, D. Pikutić, R. Štrajn, JHEP 07, 055 (2015) [arXiv:1502.02972];
- [34] T. Jurić, S. Meljanac, R. Štrajn, Phys. Lett. A 377, 2472-2476 (2013) [arXiv:1303.0994];
- [35] S. Meljanac, Z. Škoda, M. Stojić, Lett. Math. Phys. 107 no.3, 475-503 (2017) [arXiv:1409.8188];
- [36] A. Borowiec, A. Pachoł, J. Phys. A 50 no.5, 055205 (2017) [arXiv:1603.09280];
- [37] J. Lukierski, D. Meljanac, S. Meljanac, D. Pikutić, M. Woronowicz, Phys. Lett. B 777, 1–7 (2018) [arXiv:1710.09772];
- [38] J. Lukierski, S. Meljanac, M. Woronowicz, Phys. Lett. B 789, 82-87 (2019) [arXiv:1811.07365].