

# All identically conserved gravitational tensors are metric variations of invariant actions

S. Deser

*Walter Burke Institute for Theoretical Physics,  
California Institute of Technology, Pasadena, CA 91125;  
Physics Department, Brandeis University, Waltham, MA 02454  
deser@brandeis.edu*

## Abstract

I prove an old unsolved conjecture, the hard part of its obvious sufficiency, namely that all identically conserved symmetric 2-tensors are necessarily metric variations of invariant actions, thus sparing the world more alternative gravity theories. The proof is reasonably simple.

## 1 Introduction

The possible existence of covariantly conserved gravitational 2-tensors, i.e., those depending only on the metric and curvatures (and their covariant derivatives) that do not stem from an action would seem to have been raised and solved over a century ago, but to my knowledge was only raised recently [1]. Given its physical importance — adding a whole new set of alternate gravity models — this is a bit surprising, to say the least. Here I fill this gap, showing that they are indeed excluded.

## 2 Proof

We divide all conserved tensors into two disjoint categories, those that do not contain terms with one or more leading covariant derivatives, and those that do. The former consists of two types, the Einstein tensor  $G^{\mu\nu}$  and (in  $D > 4$ ) the Lovelock (double density) tensor [2]

$$L^{\mu\nu} = \epsilon^{\mu\dots} \epsilon^{\nu\dots} g_{..} g_{..} R_{...} R_{...} \quad (1)$$

Its covariant conservation is manifest, by the Bianchi identities, but what is most relevant is that it is the metric variation of  $\int 1/\sqrt{-g} L^{\mu\nu}$  because the metric variation of the curvatures vanishes by Bianchi as well. For Einstein, I need not belabor the same result — here the variation of the curvature's contribution vanishes since it consists of two covariant derivatives  $\sim DD\delta g$ , to which

the explicit metric in  $\int \sqrt{-g}g^{\mu\nu}R_{\mu\nu}$  is transparent. It was proved in [2] that those are the only two conserved tensors containing second derivatives (but no higher) of the metric. This means they have NO explicit  $D$  since they can only act on curvatures, thus making them of higher than 2 derivative order, and subject to part 2. That takes care of part 1.

Intermediate between case 1, whose  $T$  consisted entirely of a  $Z$  (in (2)) and case 2 below where both parts are present, lies a somewhat unusual  $T$ , consisting entirely of the projector part with no  $Z$ , because its action has no explicit metric-dependence. This is the famous Chern-Simons gravity model in  $D = 3$ ; details are to be found in [3]. It is of third derivative order — between the 2nd order Einstein and Lovelock and all other, fourth order or higher. Its  $T$  is the Cotton tensor (the Weyl tensor in  $D = 3$ ) whose form is  $\sim DR$ , that is a projector with one  $D$  only (a curl). This is a curiosity, but nicely stands between the 2 main cases.

The second, disjoint, category consists of tensors some of whose terms begin with covariant derivatives, the simplest being of the form

$$T^{\mu\nu} \sim (D^\mu D^\nu - g^{\mu\nu} D^2)X + Z^{\mu\nu}, \quad (2)$$

where  $X$  is a scalar function of the curvatures their derivatives, and metrics and the  $DD$  term is the variation of  $R$ . The remaining,  $Z$ , terms are corrections to ensure conservation. In terms of the underlying actions, the  $DD$  terms always come from varying the curvatures, while the  $Z$  comes from varying the explicit metrics. To be concrete, in the simplest case where  $X \sim R^n$ , the action is just

$$I = \int \sqrt{-g}R^{n+1} \quad (3)$$

and

$$T^{\mu\nu} = (n+1)(DD..)R^n + 1/2g^{\mu\nu}R^{n+1} - nR^n R^{\mu\nu} \quad (4)$$

since  $R$  varies into the above  $DD$  form times  $\delta g^{\mu\nu}$  while the explicit metric variations yield the  $Z$ . The case  $n = 0$  manifestly reduces to Einstein. Terms containing the Ricci or full curvature tensors and their derivatives are just more complicated algebraically but obviously fit the same pattern: their  $DD$  parts are well-known, hence the reconstruction of the action always replaces them there by the respective curvatures (or derivatives thereof). This is essential: given that the leading,  $DD$ , term must start with a transverse projector (see below), the story is identical to our simplest example, except for purely algebraic complications. That is, one might ask whether the second category could have other types of leading  $DD$  terms<sup>1</sup>: the answer is no, as can be seen at linearized order where they must necessarily reduce to transverse projectors such as  $(\partial\partial - \eta\partial^2)$  or four-index ones such as the operator  $\sim (\partial\partial\eta..)^{\mu\nu}$  acting on  $h_{\alpha\beta}$  to yield  $G^{\mu\nu}(\text{Lin})$ , or the even longer ones for linearized Riemann Bianchi. Let me be more pedantic: Any conserved candidate must be a sum of the above three types of projectors; that is only necessary, however; that is, there must be  $Z^{\mu\nu}$  correction terms to insure that conservation is really present. This is not automatic, and obvious counter-examples are easy to construct. The existence of  $Z$ s is only ensured by underlying actions. Indeed, the “good”  $Z$  are only those that come from an action, whose variation is the allowed sum

---

<sup>1</sup>There is an infinite variety from the basic forms, for example just insert powers of d’Alembertians in the middle of any actions etc. The universal point is that all  $T$  must start as projectors by the linearized argument and as can also be seen from the action’s variations — varying curvatures, even if there are many derivatives on them, the outermost factor is always the projector. The apparent exceptions, Einstein and Lovelock, are not — it’s just that there the projectors annihilate their subjects.

of projectors and corrections. The simplest example of a bad combination is as follows. Consider the action  $\sim \int RG^2_{\mu\nu}$ , which yields a  $T$  of the exact form  $T = P(R)G^2 + 2P(G)RG$ , where the  $P$  are the respective projectors; indices are implicit. Unless the relative coefficient is precisely 2, hence from the above action, this  $T$  is not conserved, nor can any  $Z$  save it, as is easily seen.<sup>2</sup> To summarize, the algorithmic procedure to test  $DT = 0$  candidates is first two check if the projector sum is compatible with an underlying action's. If not, no  $Z$  can help, as illustrated by the  $\int RG^2$ . If yes, then success is guaranteed simply because it comes from that action; hence conservation and being the variation of an action are synonymous.

This completes our proof: the second category, the one with explicit leading  $D$ -terms, so of higher derivative order, necessarily begins with what would be transverse projectors at linearized level plus corrections as in our simplest example, and also must stem from an action or else there are no possible  $Z$ -correction terms.

### 3 Conclusion

Trees have been saved by the proof that all viable-conserved, candidate gravity tensors are necessary derivable from actions, as has always been tacitly assumed to date.

### Acknowledgements

I thank my old collaborator, Y. Pang for useful correspondence and J. Franklin for tech help. This work was supported by the U.S. Department of Energy, Office of Science, Office of High Energy Physics, Award Number de-sc0011632.

### References

- [1] S. Deser and Y. Pang Phys Lett B790, 533(2019), arXiv 1811.03124; S.Deser ibid B795,482(2019), arXiv 1905.12831.
- [2] D. Lovelock, J Math Phys 12, 498(1971).
- [3] S. Deser, R. Jackiw and S.Templeton, Ann Phys 140, 372(1982).

---

<sup>2</sup>We dispose of one last hurdle: If the  $X$  term in (2) contained derivatives, say d'Alembertians, before the  $P^{\mu\nu}$  — but not of course powers of curvatures, which would spoil the linearized limit — this would not alter this limit's validity. However, no  $Z$  could cure the extra  $(D, D) = R$  terms raised by taking the divergence past those derivatives on its way to  $P$ : This is obvious by dimensionality —  $Z$  contains fewer derivatives than the  $X$  term, each fewer pair replaced by a curvature. So its divergence cannot match the  $X$  part's. Of course, as stated in text,  $T$ s coming from an action always have  $P$  standing to their extreme left from varying curvatures, even if they carry arbitrary derivatives.