

Positivity of the exterior power of the tangent bundles

Kiwamu Watanabe

ABSTRACT. Let X be a complex smooth projective variety such that the exterior power of the tangent bundle $\bigwedge^r T_X$ is nef for some $1 \leq r < \dim X$. We prove that, up to an étale cover, X is a Fano fiber space over an Abelian variety. This gives generalizations of the structure theorem of varieties with nef tangent bundle by Demailly, Peternell and Schneider [5] and that of varieties with nef $\bigwedge^2 T_X$ by the author [19]. Our result also gives an answer to a question raised by Li, Ou and Yang [14] for varieties with strictly nef $\bigwedge^r T_X$ when $r < \dim X$.

1. Introduction

Positivity for vector bundles such as ampleness and nefness has left its mark on the study of algebraic geometry. Let X be a complex smooth projective variety of dimension n ; we focus on the positivity of the tangent bundle T_X , which reflects the global geometry of X . As a generalization of the Hartshorne-Frankel conjecture solved by Mori [16] (see also [17] by Siu and Yau), Campana and Peternell [1] studied the structure of smooth projective varieties with nef tangent bundle, paying special attention to 3-folds. In higher dimensional case, Demailly, Peternell and Schneider obtained the following structure theorem:

THEOREM 1.1 ([5, Main Theorem]). *If T_X is nef, then there exists a finite étale cover $X' \rightarrow X$ such that X' is a locally trivial fibration $\varphi : X' \rightarrow \text{Alb}(X')$ whose fibers are a Fano variety.*

Some years ago, Cao and Höring extended Theorem 1.1 to a more general setting:

THEOREM 1.2 ([3, Theorem 1.3]). *If the anticanonical divisor $-K_X$ is nef, then there exists a finite étale cover $X' \rightarrow X$ such that $X' \cong Y \times Z$ where K_Y is trivial and Z is a locally trivial fibration $\varphi : Z \rightarrow \text{Alb}(Z)$ with a rationally connected fiber.*

In general a fiber of φ in Theorem 1.2 is not a Fano variety, because there exists a lot of rationally connected projective varieties with nef anticanonical divisor which is not Fano (for instance, consider weak Fano varieties). The main result of this paper is a generalization of Theorem 1.1:

2010 *Mathematics Subject Classification.* 14J40, 14J45, 14M22.

The author is partially supported by JSPS KAKENHI Grant Number 21K03170 and the Sumitomo Foundation Grant Number 190170.

THEOREM 1.3. *Let X be a smooth projective variety of dimension n . Assume that $\bigwedge^r T_X$ is nef for some $1 \leq r < n$. Then if we take a suitable finite étale cover $\tilde{X} \rightarrow X$, there exists a locally trivial fibration $\varphi : \tilde{X} \rightarrow A$ such that the fiber F is a Fano variety and A is an Abelian variety. Moreover, if $\dim A \geq r - 1$, then T_X is nef; otherwise $\bigwedge^{r-\dim A} T_{\tilde{X}/A}$ is nef.*

This theorem reduces the study of smooth projective varieties with nef $\bigwedge^r T_X$ ($r < \dim X$) to that of Fano varieties. For $r = 1$, Theorem 1.3 is nothing but Theorem 1.1; for $r = 2$ this was obtained in [19, Theorem 1.5]. The proof of [19, Theorem 1.5] involves the deformation theory of rational curves and some complicated arguments. On the other hand, in this short paper, we give a really simple proof of Theorem 1.3. Our proof relies on two key ingredients; one is Theorem 1.1; the other is a recent result of Gachet [4]. In [4, Theorem 1.2], she proved that for a smooth projective variety X of dimension n if $\bigwedge^{n-1} T_X$ is strictly nef, then X is a Fano variety. Her proof works if we replace the assumption that $\bigwedge^{n-1} T_X$ is strictly nef by the assumption that X is rationally connected and $\bigwedge^{n-1} T_X$ is nef. Moreover by using a result by Laytimi and Nahm [11], we see that if $\bigwedge^r T_X$ is nef for some $r < n$, then so is $\bigwedge^{n-1} T_X$. Thus we have the following:

PROPOSITION 1.4. *Let X be a smooth projective variety of dimension n . Assume that X is rationally connected and $\bigwedge^r T_X$ is nef for some $1 \leq r < n$. Then X is a Fano variety.*

Remark that, combining with [14, Theorem 1.2], Proposition 1.4 gives an affirmative answer to the following question by Li, Ou and Yang when $r < \dim X$:

QUESTION 1.5 ([14, Remark 5.3], [15, Conjecture 4.9], [4, Question in Section 1]). Assume that $\bigwedge^r T_X$ is strictly nef for some $1 \leq r \leq n$. Then is X a Fano variety?

Finally, Theorem 1.3 follows from Theorem 1.2, Proposition 1.4 and standard arguments.

2. Preliminaries

2.1. Notation and Conventions. We will use the basic notation and definitions in [8], [9], [12], [13] and [10]. Along this paper, we work over the complex number field.

- A *curve* means a projective variety of dimension one.
- Let X be a smooth projective variety. A line bundle L on X is said to be *strictly nef* (resp. *nef*) if the intersection number $L \cdot C$ is positive (resp. non-negative) for any curve $C \subset X$. In general, we say that a vector bundle \mathcal{E} is *strictly nef* (resp. *nef*) if the tautological line bundle $\mathcal{O}_{\mathbb{P}(\mathcal{E})}(1)$ is strictly nef (resp. nef) on $\mathbb{P}(\mathcal{E})$.
- For a non-constant morphism $f : \mathbb{P}^1 \rightarrow X$ from a projective line \mathbb{P}^1 to a smooth projective variety X , f is said to be *free* if f^*T_X is nef.

Throughout this section, we always assume the following:

ASSUMPTION 2.1. Assume X is a smooth projective variety of dimension n such that the exterior power $\bigwedge^r T_X$ is nef for some $1 \leq r < n$.

PROPOSITION 2.2. *The following hold:*

- (i) *The anticanonical divisor $-K_X$ is nef.*
- (ii) *If the Kodaira dimension $\kappa(X) = 0$, then there exists a finite étale cover $f : \tilde{X} \rightarrow X$ such that \tilde{X} is an Abelian variety.*

PROOF. The first part follows from $\det(\wedge^r T_X) \cong \mathcal{O}_X \left(\binom{n-1}{r-1} (-K_X) \right)$. The second part follows from [21, Theorem 1.1] (see also [2, Proposition 1.2]). \blacksquare

LEMMA 2.3 ([2, Lemma 1.3], [20, Lemma 2.9]). *Let $f : \mathbb{P}^1 \rightarrow X$ be a non-free rational curve, that is, f^*T_X is not nef. Then we have $-K_X \cdot f_*(\mathbb{P}^1) \geq n - r + 1$.*

PROOF. Assume that the splitting type of f^*T_X is (a_1, a_2, \dots, a_n) , that is,

$$f^*T_X \cong \bigoplus_{i=1}^n \mathcal{O}_{\mathbb{P}^1}(a_i) \quad (a_1 \geq a_2 \geq \dots \geq a_n, a_1 \geq 2).$$

The r -th exterior power

$$\bigwedge^r f^*T_X \cong \bigoplus_{1 \leq i_1 < i_2 < \dots < i_r \leq n} \mathcal{O}_{\mathbb{P}^1}(a_{i_1} + a_{i_2} + \dots + a_{i_r})$$

is nef; this yields

$$a_{n-r+1} + a_{n-r+2} + \dots + a_n \geq 0.$$

Since f is not free, a_n is negative. These imply that

$$(r-1)a_{n-r+1} \geq a_{n-r+1} + a_{n-r+2} + \dots + a_{n-1} \geq -a_n \geq 1.$$

Thus a_{n-r+1} is positive. As a consequence, we have the inequality

$$-K_X \cdot f_*(\mathbb{P}^1) = a_1 + (a_2 + \dots + a_{n-r}) + (a_{n-r+1} + \dots + a_n) \geq 2 + (n-r-1) + 0 = n-r+1. \quad \blacksquare$$

PROPOSITION 2.4 ([18, Proposition 3.3]). *Let $\varphi : X \rightarrow A$ be a smooth morphism onto an Abelian variety with irreducible fibers. Then the following hold:*

- (i) *If $\dim A \geq r - 1$, then T_X is nef.*
- (ii) *If $\dim A < r - 1$, then $\bigwedge^{r-\dim A} T_{X/A}$ is nef.*

PROOF. We have an exact sequence

$$(1) \quad 0 \rightarrow T_{X/A} \rightarrow T_X \rightarrow \varphi^*T_A \rightarrow 0.$$

By [8, Chapter II, Exercise 5.16 (d)], we have a filtration of $\bigwedge^r T_X$:

$$\bigwedge^r T_X = E^0 \supset E^1 \supset E^2 \supset \dots \supset E^{r+1} = 0$$

such that

$$E^p/E^{p+1} \cong \left(\bigwedge^p T_{X/A} \right) \otimes \left(\bigwedge^{r-p} \varphi^*T_A \right)$$

for any p . In particular, we have the following exact sequences:

$$(2) \quad 0 \rightarrow E^1 \rightarrow \bigwedge^r T_X \rightarrow \bigwedge^r \varphi^*T_A \rightarrow 0$$

$$(3) \quad 0 \rightarrow E^2 \rightarrow E^1 \rightarrow T_{X/A} \otimes \left(\bigwedge^{r-1} \varphi^*T_A \right) \rightarrow 0$$

To prove (i), assume $\dim A \geq r - 1$. Remark that $T_A \cong \mathcal{O}_A^{\oplus \dim A}$. We claim that E^1 is nef. If $\dim A \geq r$, then it follows from the sequence (2) and [1, Proposition 1.2 (8)] that E^1 is nef. If $\dim A = r - 1$, then the sequence (2) yields $E^1 \cong \bigwedge^r T_X$; this implies that E^1 is nef. By the sequence (3), $T_{X/A} \otimes (\bigwedge^{r-1} \varphi^* T_A)$ is nef. Since $\bigwedge^{r-1} (\varphi^* T_A)$ is trivial bundle, we conclude that the relative tangent bundle $T_{X/A}$ is nef. Finally our assertion follows from the sequence (1).

To prove (ii), assume $\dim A < r - 1$. Since $\bigwedge^p \varphi^* T_A = 0$ for any $p > \dim A$, we have

$$\bigwedge^r T_X = E^0 = E^1 = \dots = E^{r-\dim A}.$$

Thus we have a surjection $\bigwedge^r T_X = E^{r-\dim A} \rightarrow \bigwedge^{r-\dim A} T_{X/A}$; this implies that $\bigwedge^{r-\dim A} T_{X/A}$ is nef. \blacksquare

3. Proof of the Main Theorem

The following is due to Gachet:

PROPOSITION 3.1 ([4, Theorem 1.2]). *Let X be a smooth projective variety of dimension n . Assume that X is rationally connected and $\bigwedge^{n-1} T_X$ is nef. Then $-K_X$ is ample, that is, X is a Fano variety.*

PROOF. This follows from the same argument as in [4, Lemma 3.1, Lemma 3.3]. Actually the proof works if we replace the assumption that $\bigwedge^{n-1} T_X$ is strictly nef by the assumption that X is rationally connected and $\bigwedge^{n-1} T_X$ is nef; this yields that $-K_X$ is nef and big. Then we may conclude that $-K_X$ is ample by the same argument as in [4, Lemma 3.3] and Lemma 2.3. \blacksquare

REMARK 3.2. Although Proposition 3.1 is not written explicitly in [4], Gachet introduced this statement holds at Algebraic Geometry seminar of the University of Tokyo (see Acknowledgements below).

THEOREM 3.3 ([11, Theorem 3.3], [6]). *Let X be a smooth projective variety of dimension n . For a vector bundle E of rank r , assume that its exterior power $\bigwedge^m E$ is nef for some positive integer m . Then the vector bundle $\bigwedge^{m+k} E$ is also nef for any $0 \leq k \leq n - m$.*

REMARK 3.4. In general, if a vector bundle E is strictly nef, it is not necessarily that its exterior power $\bigwedge^r E$ is strictly nef. For instance, see [7, Section 10 in Chapter I] and [15, Example 2.1]). This means that an analogue of Theorem 3.3 does not hold if we replace nefness of $\bigwedge^m E$ by strictly nefness.

PROOF OF PROPOSITION 1.4. Assume that X is rationally connected and $\bigwedge^r T_X$ is nef for some $1 \leq r < n$. Then Theorem 3.3 implies that $\bigwedge^{n-1} T_X$ is nef. Applying Proposition 2.4, we see that X is a Fano variety. \blacksquare

PROOF OF THEOREM 1.3. By Proposition 2.2 (i), $-K_X$ is nef; according to Theorem 1.2, this turns out that there exists a finite étale cover $X' \rightarrow X$ such that $X' \cong Y \times Z$ where K_Y is trivial and Z is a locally trivial fibration $Z \rightarrow \text{Alb}(Z)$ with a rationally connected fiber. Since we have $X' \rightarrow X$ is étale, $\bigwedge^r T_{X'}$ is also nef; then by Theorem 3.3, $\bigwedge^{n-1} T_{X'}$ is also nef. Let $p_1 : X' \rightarrow Y$ (resp. $p_2 : X' \rightarrow Z$)

be the first projection (resp. the second projection). We denote by ℓ the dimension of Y . Since we have

$$\bigwedge^{n-1} T_{X'} \cong \left[p_1^* \left(\bigwedge^\ell T_Y \right) \otimes p_2^* \left(\bigwedge^{n-\ell-1} T_Z \right) \right] \oplus \left[p_1^* \left(\bigwedge^{\ell-1} T_Y \right) \otimes p_2^* \left(\bigwedge^{n-\ell} T_Z \right) \right],$$

the direct summand $p_1^* \left(\bigwedge^{\ell-1} T_Y \right) \otimes p_2^* \left(\bigwedge^{n-\ell} T_Z \right)$ is nef; restricting this bundle to a fiber of the projection p_2 , we see that $\bigwedge^{\ell-1} T_Y$ is also nef provided that $\ell > 0$. If $\ell = 1$, then Y is an elliptic curve. Furthermore if $\ell > 1$, then Proposition 2.2 (ii) implies that Y is a finite étale quotient of an Abelian variety \tilde{Y} . Hence, in any case, there exists a finite étale cover $\tilde{X} \rightarrow X'$ such that \tilde{X} is a locally trivial fibration $\varphi : \tilde{X} \rightarrow A$ onto an Abelian variety A with a rationally connected fiber. Then our assertion follows from Proposition 2.4 and Proposition 1.4. \blacksquare

Acknowledgements

On August the 1st 2022, Cécile Gachet gave a talk on the result of [4] in Algebraic Geometry seminar of the University of Tokyo; the author knew her excellent result [4, Theorem 1.2] at the time. The author would like to thank Cécile Gachet for sending him her preprint [4] and answering his questions. The author would like to thank Kento Fujita for sending him his private note [6], which states that for a vector bundle E of rank r if $\bigwedge^\ell E$ is nef, then so is $\bigwedge^m E$ for any $\ell \leq m \leq r$. On the other hand, while preparing this paper, the author noticed the result of [6] was contained in [11].

References

- [1] Frédéric Campana and Thomas Peternell. Projective manifolds whose tangent bundles are numerically effective. *Math. Ann.*, 289(1):169–187, 1991.
- [2] Frédéric Campana and Thomas Peternell. On the second exterior power of tangent bundles of threefolds. *Compositio Math.*, 83(3):329–346, 1992.
- [3] Junyan Cao and Andreas Höring. A decomposition theorem for projective manifolds with nef anticanonical bundle. *J. Algebraic Geom.*, 28(3):567–597, 2019.
- [4] Gachet Cécile. Positivity of higher exterior powers of the tangent bundle. Preprint arXiv:2207.10854, 2022.
- [5] Jean-Pierre Demailly, Thomas Peternell, and Michael Schneider. Compact complex manifolds with numerically effective tangent bundles. *J. Algebraic Geom.*, 3(2):295–345, 1994.
- [6] Kento Fujita. Remarks on ampleness of wedge vector bundles. private note.
- [7] Robin Hartshorne. *Ample subvarieties of algebraic varieties*. Lecture Notes in Mathematics, Vol. 156. Springer-Verlag, Berlin, 1970. Notes written in collaboration with C. Musili.
- [8] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [9] János Kollár. *Rational curves on algebraic varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1996.
- [10] János Kollár and Shigefumi Mori. *Birational geometry of algebraic varieties*, volume 134 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 1998. With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- [11] F. Laytimi and W. Nahm. A vanishing theorem. *Nagoya Math. J.*, 180:35–43, 2005.
- [12] Robert Lazarsfeld. *Positivity in algebraic geometry. I*, volume 48 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2004. Classical setting: line bundles and linear series.

- [13] Robert Lazarsfeld. *Positivity in algebraic geometry. II*, volume 49 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2004. Positivity for vector bundles, and multiplier ideals.
- [14] Duo Li, Wenhao Ou, and Xiaokui Yang. On projective varieties with strictly nef tangent bundles. *J. Math. Pures Appl. (9)*, 128:140–151, 2019.
- [15] Jie Liu, Wenhao Ou, and Xiaokui Yang. Strictly nef vector bundles and characterizations of \mathbb{P}^n . *Complex Manifolds*, 8(1):148–159, 2021.
- [16] Shigefumi Mori. Projective manifolds with ample tangent bundles. *Ann. of Math. (2)*, 110(3):593–606, 1979.
- [17] Yum Tong Siu and Shing Tung Yau. Compact Kähler manifolds of positive bisectional curvature. *Invent. Math.*, 59(2):189–204, 1980.
- [18] Kiwamu Watanabe. Fano manifolds of coindex three admitting nef tangent bundle. *Geom. Dedicata*, 210:165–178, 2021.
- [19] Kiwamu Watanabe. Positivity of the second exterior power of the tangent bundles. *Adv. Math.*, 385:Paper No. 107757, 27, 2021.
- [20] Kazunori Yasutake. On the second exterior power of tangent bundles of Fano fourfolds with picard number $\rho_x \geq 2$. Preprint arXiv:1212.0685, 2012.
- [21] Kazunori Yasutake. On the second and third exterior power of tangent bundles of Fano manifolds with birational contractions. Preprint arXiv:1403.5304, 2014.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ENGINEERING, CHUO UNIVERSITY.
1-13-27 KASUGA, BUNKYO-KU, TOKYO 112-8551, JAPAN
Email address: watanabe@math.chuo-u.ac.jp