

DISTRIBUTIONS OF RESONANCES OF SUPERCRITICAL QUASI-PERIODIC OPERATORS

WENCAI LIU

ABSTRACT. We discover that the distribution of (frequency and phase) resonances plays a role in determining the spectral type of supercritical quasi-periodic Schrödinger operators. In particular, we disprove the second spectral transition line conjecture of Jitomirskaya in the early 1990s.

1. INTRODUCTION

In this paper, we study the quasi-periodic Schrödinger operator $H = H_{\lambda v, \alpha, \theta}$ defined on $\ell^2(\mathbb{Z})$:

$$(1) \quad (H_{\lambda v, \alpha, \theta} u)(n) = u(n+1) + u(n-1) + \lambda v(\theta + n\alpha)u(n), \quad n \in \mathbb{Z}$$

where λ is the coupling constant, $v : \mathbb{T} = \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$ is the (analytic) potential, $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ is the frequency, and $\theta \in \mathbb{R}$ is the phase. Letting $v = 2 \cos 2\pi\theta$ in (1), we obtain the almost Mathieu operator

$$(2) \quad (H_{\lambda, \alpha, \theta} u)(n) = u(n+1) + u(n-1) + 2\lambda \cos 2\pi(\theta + n\alpha)u(n).$$

The almost Mathieu operator is a popular model in both mathematics and physics. We refer readers to [20, 21, 27, 38, 40, 41] for the history and recent development of quasi-periodic Schrödinger operators.

By Avila's global theory [5], the analytic quasi-periodic Schrödinger operator (1) can be categorized into three cases based on the behavior of the complexified Lyapunov exponent of corresponding Schrödinger cocycles: subcritical, critical and supercritical. For the almost Mathieu operator, the Lyapunov exponent on the spectrum only depends on λ [15], so the three regimes are characterized by the coupling constant λ ,

1. subcritical: $|\lambda| < 1$;
2. critical: $|\lambda| = 1$;

Key words and phrases. Almost Mathieu operator, supercritical quasiperiodic Schrödinger operator, Lyapunov exponent, second spectral transition line, frequency resonance, phase resonance, Anderson localization.

2020 Mathematics Subject Classification. Primary: 47A10. Secondary: 81Q10, 47B39.

3. supercritical: $|\lambda| > 1$.

We remark that the three regimes of the almost Mathieu operator are not surprising, which has already been seen from the prediction of Aubry and André [1]:

1. if $|\lambda| < 1$, $H_{\lambda,\alpha,\theta}$ has purely absolutely continuous spectrum for all $\alpha \in \mathbb{R}$ and $\theta \in \mathbb{R}$;
2. if $|\lambda| = 1$, $H_{\lambda,\alpha,\theta}$ has purely singular continuous spectrum for all $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $\theta \in \mathbb{R}$.
3. if $|\lambda| > 1$, $H_{\lambda,\alpha,\theta}$ has only pure point spectrum for all $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $\theta \in \mathbb{R}$.

For the subcritical (namely $|\lambda| < 1$) almost Mathieu operator, after many earlier works [6, 8, 19, 30], Avila finally proved that the spectrum is purely absolutely continuous [3] for any $(\alpha, \theta) \in \mathbb{R}^2$. From the almost reducibility conjecture [2, 4], subcritical analytic quasi-periodic Schrödinger operators (for small λ , the operator (1) is always subcritical) always have purely absolutely continuous spectrum.

It is expected that the critical quasi-periodic Schrödinger operator only supports the singular continuous spectrum. This has been recently proved for the almost Mathieu operator [9, 22]. We should mention that critical analytic quasi-periodic Schrödinger operators do not typically happen [5].

The supercritical analytic quasi-periodic operator (for large λ , the operator (1) is always supercritical) has Anderson localization (only pure point spectrum with exponentially decaying eigenfunctions) for almost every $(\alpha, \theta) \in \mathbb{R}^2$ [13, 14, 30], but not for all $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $\theta \in \mathbb{R}$ since the arithmetics of frequencies α and phases θ play roles [12, 18, 28, 39]. This in particular implies that Part 3 of Aubry-Andre's prediction does not hold all $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ and $\theta \in \mathbb{R}$.

From the discussion above, one can see that it is particularly interesting to study supercritical quasi-periodic operators, which is exactly the main focus of this paper.

By Kotani theory, the supercritical analytic quasi-periodic operator does not have any absolutely continuous spectrum [31, 32]. It is believed that whether the spectrum is pure point or singular continuous depends on resonances arising from the small denominators and the Lyapunov exponent.

There are two types of resonances appearing naturally. The first one, the frequency resonance, was first observed in [12, 18], which is

quantified by

$$(3) \quad \beta(\alpha) = \limsup_{|k| \rightarrow \infty} -\frac{\ln \|k\alpha\|_{\mathbb{R}/\mathbb{Z}}}{|k|},$$

where $\|x\|_{\mathbb{R}/\mathbb{Z}} = \text{dist}(x, \mathbb{Z})$.

Another one, the phase resonance, happens for even functions [28] and is quantified by

$$(4) \quad \delta(\alpha, \theta) = \limsup_{|k| \rightarrow \infty} -\frac{\ln \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}}}{|k|}.$$

A celebrated result of Jitomirskaya show that when $\beta(\alpha) = \delta(\alpha, \theta) = 0$, $H_{\lambda, \alpha, \theta}$ has Anderson localization for $|\lambda| > 1$ [24, 30]. In [29], Jitomirskaya conjectured that for the supercritical almost Mathieu operator, the spectral type (pure point spectrum/localization or singular continuous spectrum) completely determined by competitions between the Lyapunov exponent and frequency/phase resonances quantified by (3) and (4). More precisely,

Conjecture 1

- the almost Mathieu operator $H_{\lambda, \alpha, \theta}$ satisfies Anderson localization if $|\lambda| > e^{\beta(\alpha) + \delta(\alpha, \theta)}$;
- the almost Mathieu operator $H_{\lambda, \alpha, \theta}$ has purely singular continuous spectrum if $1 < |\lambda| < e^{\beta(\alpha) + \delta(\alpha, \theta)}$.

Conjecture 1 is referred to as the second spectral transition line conjecture.

Significant progress in establishing the second spectral transition line has been made in the past 15 years [7, 8, 10, 11, 16, 17, 23, 25, 26, 33, 35–37]. Here is a summary.

Second spectral transition line

1. When $|\lambda| > e^{\beta(\alpha)}$ and $\delta(\alpha, \theta) = 0$, $H_{\lambda, \alpha, \theta}$ has Anderson localization [25].
2. When $|\lambda| > e^{\delta(\alpha, \theta)}$ and $\beta(\alpha) = 0$, $H_{\lambda, \alpha, \theta}$ has Anderson localization [26].
3. When $|\lambda| > e^{2\beta(\alpha)}$, $H_{\lambda, \alpha, \theta}$ satisfies Anderson localization for completely resonant phase θ , namely, $2\theta \in \alpha\mathbb{Z} + \mathbb{Z}$ [34].
4. When $1 < |\lambda| < e^{\beta(\alpha)}$, $H_{\lambda, \alpha, \theta}$ has purely singular continuous spectrum [11].
5. When $1 < |\lambda| < e^{\delta(\alpha, \theta)}$, $H_{\lambda, \alpha, \theta}$ has purely singular continuous spectrum [26].

It is worthwhile to mention that Part 3 solves a conjecture of Avila and Jitomirskaya [7]. It is easy to see that for completely resonant phases

θ , namely $2\theta \in \alpha\mathbb{Z} + \mathbb{Z}$, $\delta(\alpha, \theta) = \beta(\alpha)$, so Part 3 is a particular case of Conjecture 1. All previous results give positive answers to Conjecture 1.

In this paper, we discover that not just strength of frequency and phase resonances plays a role in the supercritical regime, distributions of resonances are also crucial to the spectral types. In particular, we disprove Conjecture 1.

We postpone our main theorem to Section 2 and list some corollaries first.

We denote the continued fraction expansion of α by

$$(5) \quad \alpha = [a_0, a_1, a_2, \dots] = a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \frac{1}{\ddots}}}}.$$

Corollary 1.1. *Assume that α satisfies $0 < \beta(\alpha) < \infty$ and $\liminf_{n \rightarrow \infty} a_n \geq 10^{20}$. Then there exists θ with $0 < \delta(\alpha, \theta) < \infty$ such that*

1. for $|\lambda| > e^{\delta(\alpha, \theta)}$, $H_{\lambda, \alpha, \theta}$ has Anderson localization;
2. for $1 \leq |\lambda| < e^{\delta(\alpha, \theta)}$, $H_{\lambda, \alpha, \theta}$ has purely singular continuous spectrum;
3. for $|\lambda| < 1$, $H_{\lambda, \alpha, \theta}$ has purely absolutely continuous spectrum.

Corollary 1.2. *For any $0 < \mu < \infty$, there exist α with $\beta(\alpha) = \mu$ and θ with $0 < \delta(\alpha, \theta) < \infty$ such that*

1. for $|\lambda| > e^{\delta(\alpha, \theta)}$, $H_{\lambda, \alpha, \theta}$ has Anderson localization;
2. for $1 \leq |\lambda| < e^{\delta(\alpha, \theta)}$, $H_{\lambda, \alpha, \theta}$ has purely singular continuous spectrum;
3. for $|\lambda| < 1$, $H_{\lambda, \alpha, \theta}$ has purely absolutely continuous spectrum.

Remark 1.3. • The second spectral transition line for (α, θ) in Corollaries 1.1 and 1.2 is $|\lambda| = e^{\delta(\alpha, \theta)}$ differing from $|\lambda| = e^{\beta(\alpha) + \delta(\alpha, \theta)}$, which disproves Conjecture 1.

- Parts 2 and 3 in Corollaries 1.1 and 1.2 are known for all possible α and θ [3, 22, 26]. They are included here for completeness.
- Part 1 in Corollaries 1.1 and 1.2 follows from our main Theorem 2.1 below and some simple facts about continued fraction expansions.

The proof of our main results is constructive. As we mentioned earlier, our key observation is that the distribution of resonances plays a role in determining the spectral type. The main idea behind our construction is that when locations of phase resonances and frequency resonances are far away, the Lyapunov exponent with a smaller magnitude (predicted by Conjecture 1) is sufficient to beat both frequency

and phase resonances. Our analysis in the present paper gives us the indication that Conjecture 1 is only true for the two extreme cases ($\beta(\alpha) = 0$ or $\delta(\alpha, \theta) = 0$). So, we conjecture that

Conjecture 2

For any α with $0 < \beta(\alpha) < \infty$ and $0 < \kappa < \infty$, there exists θ with $\delta(\alpha, \theta) = \kappa$ such that the following is not true. $H_{\lambda, \alpha, \theta}$ has Anderson localization for $|\lambda| > e^{\kappa + \beta(\alpha)}$ and has purely singular continuous spectrum for $1 < |\lambda| < e^{\kappa + \beta(\alpha)}$.

We list two simpler versions of Conjecture 2 here.

Conjecture 3

For any α with $0 < \beta(\alpha) < \infty$, there exists θ with $0 < \delta(\alpha, \theta) < \infty$ such that the following is not true. $H_{\lambda, \alpha, \theta}$ has Anderson localization for $|\lambda| > e^{\delta(\alpha, \theta) + \beta(\alpha)}$ and has purely singular continuous spectrum for $1 < |\lambda| < e^{\delta(\alpha, \theta) + \beta(\alpha)}$.

Conjecture 4

For any $0 < \mu < \infty$ and $0 < \kappa < \infty$, there exist α and θ with $\beta(\alpha) = \mu$ and $\delta(\alpha, \theta) = \kappa$ such that the following is not true. $H_{\lambda, \alpha, \theta}$ has Anderson localization for $|\lambda| > e^{\kappa + \mu}$ and has purely singular continuous spectrum for $1 < |\lambda| < e^{\kappa + \mu}$.

As a corollary of Theorem 2.1 below, we prove Conjecture 4 in the regime $\kappa \in [100\mu, \infty)$ and $0 < \mu < \infty$, namely,

Corollary 1.4. *For any $0 < \mu < \infty$ and any $\kappa \in [100\mu, \infty)$, there exist α and θ with $\beta(\alpha) = \mu$ and $\delta(\alpha, \theta) = \kappa$ such that*

1. for $|\lambda| > e^{\delta(\alpha, \theta)}$, $H_{\lambda, \alpha, \theta}$ has Anderson localization;
2. for $1 \leq |\lambda| < e^{\delta(\alpha, \theta)}$, $H_{\lambda, \alpha, \theta}$ has purely singular continuous spectrum;
3. for $|\lambda| < 1$, $H_{\lambda, \alpha, \theta}$ has purely absolutely continuous spectrum.

2. CONSTRUCTIONS OF θ

Without loss of generality, assume $\alpha \in [0, 1] \setminus \mathbb{Q}$. Recall (5) and for $n \geq 1$, let

$$(6) \quad \frac{p_n}{q_n} = [a_1, a_2, \dots, a_n] = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \frac{1}{\ddots + \frac{1}{a_n}}}}},$$

where $(p_n, q_n) = 1$. By setting $q_0 = 1$ and $q_{-1} = 0$, one has that for any $n \geq 1$,

$$(7) \quad q_n = a_n q_{n-1} + q_{n-2}.$$

By basic facts of the continued fraction expansion, we have that for any $1 \leq k < q_{n+1}$,

$$(8) \quad \|k\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \|q_n\alpha\|_{\mathbb{R}/\mathbb{Z}},$$

$$(9) \quad \frac{1}{q_n + q_{n+1}} \leq \|q_n\alpha\|_{\mathbb{R}/\mathbb{Z}} = |q_n\alpha - p_n| \leq \frac{1}{q_{n+1}}.$$

and for even n ,

$$(10) \quad \frac{p_n}{q_n} < \alpha < \frac{p_{n+1}}{q_{n+1}}.$$

By (3) and (7)-(9), one has that

$$(11) \quad \beta(\alpha) = \limsup_{n \rightarrow \infty} \frac{\ln q_{n+1}}{q_n}.$$

We are in a position to construct θ .

Fix any η with $0 < \eta \leq 10^{-2}$. Assume that α satisfies

$$(12) \quad \liminf_{n \rightarrow \infty} a_n \geq 10^{18}\eta^{-1}.$$

By the assumption (12), there exists large enough \tilde{j}_0 such that for any $j \geq \tilde{j}_0$,

$$(13) \quad \eta q_{j+1} \geq 10^{17} q_j.$$

Let k_0 be a positive integer (the existence is guaranteed by the ergodic theory) so that

$$(14) \quad (-k_0\alpha) \pmod{\mathbb{Z}} \in \left[\frac{1}{10}, \frac{1}{5} \right].$$

Case 1: $\limsup_{j \rightarrow \infty} \frac{\ln q_{2j}}{q_{2j-1}} = \beta(\alpha)$.

Let $k_{\tilde{j}_0} = k_0$. For any $j \geq \tilde{j}_0$, define

$$(15) \quad k_{j+1} = k_j + \left\lfloor \eta \frac{q_{2j+1}}{q_{2j}} \right\rfloor q_{2j},$$

where $\lfloor x \rfloor$ is the largest integer that is less than or equal to x . For any $j \geq \tilde{j}_0$, let

$$(16) \quad I_j = \left\{ \theta \in [0, 1/2] : -\frac{10\eta}{q_{2j}} \leq 2\theta - ((-k_j\alpha) \pmod{\mathbb{Z}}) \leq -\frac{\eta}{10q_{2j}} \right\}.$$

By (9), (10), (13), (14), (15) and inductions, one has that for any $j \geq \tilde{j}_0$,

$$\begin{aligned} ((-k_{j+1}\alpha) \bmod \mathbb{Z}) - ((-k_j\alpha) \bmod \mathbb{Z}) &= - \left\lfloor \eta \frac{q_{2j+1}}{q_{2j}} \right\rfloor (q_{2j}\alpha - p_{2j}) \\ &\geq -\eta \frac{q_{2j+1}}{q_{2j}} |q_{2j}\alpha - p_{2j}| \\ &\geq -\frac{\eta}{q_{2j}}, \end{aligned}$$

$$\begin{aligned} ((-k_{j+1}\alpha) \bmod \mathbb{Z}) - ((-k_j\alpha) \bmod \mathbb{Z}) &\leq -\eta \frac{q_{2j+1}}{2q_{2j}} |q_{2j}\alpha - p_{2j}| \\ &\leq -\frac{\eta}{4q_{2j}}, \end{aligned}$$

and

$$(-k_j\alpha) \bmod \mathbb{Z} \in \left[\frac{1}{20}, \frac{1}{5} \right].$$

Therefore, we have that I_j (a closed interval), $j \geq \tilde{j}_0$, is monotone, namely $I_{j+1} \subset I_j$. This implies there exists θ such that

$$(17) \quad \theta = \bigcap_{j=\tilde{j}_0}^{\infty} I_j.$$

Case 2: $\limsup_{j \rightarrow \infty} \frac{\ln q_{2j+1}}{q_{2j}} = \beta(\alpha)$.

For any $j \geq \tilde{j}_0$, define

$$(18) \quad k_{j+1} = k_j + \left\lfloor \eta \frac{q_{2j}}{q_{2j-1}} \right\rfloor q_{2j-1}.$$

For any $j \geq \tilde{j}_0$, let

$$(19) \quad I_j = \left\{ \theta \in [0, 1/2] : \frac{\eta}{10q_{2j-1}} \leq 2\theta - ((-k_j\alpha) \bmod \mathbb{Z}) \leq \frac{10\eta}{q_{2j-1}} \right\}.$$

By (9), (10), (13), (14), (18) and inductions, one has that for any $j \geq \tilde{j}_0$,

$$\begin{aligned} ((-k_{j+1}\alpha) \bmod \mathbb{Z}) - ((-k_j\alpha) \bmod \mathbb{Z}) &= - \left\lfloor \eta \frac{q_{2j}}{q_{2j-1}} \right\rfloor (q_{2j-1}\alpha - p_{2j-1}) \\ &\leq \eta \frac{q_{2j}}{q_{2j-1}} |q_{2j-1}\alpha - p_{2j-1}| \\ &\leq \frac{\eta}{q_{2j-1}}, \end{aligned}$$

$$\begin{aligned} ((-k_{j+1}\alpha) \bmod \mathbb{Z}) - ((-k_j\alpha) \bmod \mathbb{Z}) &\geq \eta \frac{q_{2j}}{2q_{2j-1}} |q_{2j-1}\alpha - p_{2j-1}| \\ &\geq \frac{\eta}{4q_{2j-1}}, \end{aligned}$$

and

$$(-k_j\alpha) \bmod \mathbb{Z} \in \left[\frac{1}{10}, \frac{2}{5} \right].$$

Therefore, we have that I_j (a closed interval), $j \geq \tilde{j}_0$, is monotone, namely $I_{j+1} \subset I_j$. This implies there exists θ such that

$$(20) \quad \theta = \bigcap_{j=\tilde{j}_0}^{\infty} I_j.$$

Theorem 2.1. *Fix any η with $0 < \eta \leq 10^{-2}$. Assume that α satisfies $0 < \beta(\alpha) < \infty$ and $\liminf_{n \rightarrow \infty} a_n \geq 10^{18}\eta^{-1}$. Let θ be constructed by (17) or (20). Then for any $|\lambda| > e^{\delta(\alpha, \theta)}$, $H_{\lambda, \alpha, \theta}$ has Anderson localization.*

In the following sections, we prove Theorem 2.1. In order to avoid repetitions, we only prove the Case 1: $\limsup_{j \rightarrow \infty} \frac{\ln q_{2j}}{q_{2j-1}} = \beta(\alpha)$. The proof of Case 2 is similar. We list the definitions and standard facts in Section 3. In Section 4, we provide several technical lemmas. Sections 5 and 6 are devoted to treating non-resonant and resonant sites respectively. In Section 7, we prove corollaries and three claims which are used in Sections 5 and 6. Proofs of those claims are quite standard if readers are familiar with the small denominator arguments related to almost Mathieu operators.

3. BASICS

All basics in this section are taken from [33]. We refer readers to [33] and references therein for details. Let H be an operator on $\ell^2(\mathbb{Z})$. We say ϕ is a generalized eigenfunction corresponding to the generalized eigenvalue E if

$$(21) \quad H\phi = E\phi, \text{ and } |\phi(k)| \leq \hat{C}(1 + |k|).$$

By Shnol's theorem, in order to prove Anderson localization of H , we only need to show that every generalized eigenfunction is in fact an exponentially decaying eigenfunction, namely, there exists some constant $c > 0$ such that

$$|\phi(k)| \leq e^{-c|k|} \text{ for large } k.$$

For simplicity, we assume $\hat{C} = 1$ in (21).

From now on, we always assume ϕ is a generalized eigenfunction of $H_{\lambda,\alpha,\theta}$ and E is the corresponding generalized eigenvalue. Without loss of generality assume $\phi(0) = 1$. It is well known that every generalized eigenvalue must be in the spectrum, namely $E \in \Sigma_{\lambda,\alpha}$, where $\Sigma_{\lambda,\alpha}$ is the spectrum of $H_{\lambda,\alpha,\theta}$ (the spectrum does not depend on θ).

For any $x_1, x_2 \in \mathbb{Z}$ with $x_1 < x_2$, denote by

$$P_{[x_1, x_2]}(\lambda, \alpha, \theta) = \det(R_{[x_1, x_2]}(H_{\lambda,\alpha,\theta} - E)R_{[x_1, x_2]}),$$

where $R_{[x_1, x_2]}$ is the restriction on $[x_1, x_2]$. Let us denote

$$P_k(\lambda, \alpha, \theta) = \det(R_{[0, k-1]}(H_{\lambda,\alpha,\theta} - E)R_{[0, k-1]}).$$

When there is no ambiguity, we drop the dependence on parameters E, λ, α or θ . Clearly,

$$(22) \quad P_{[x_1, x_2]}(\theta) = P_k(\theta + x_1\alpha),$$

where $k = x_2 - x_1 + 1$.

Let

$$(23) \quad A_k(\theta) = \prod_{j=k-1}^0 A(\theta + j\alpha) = A(\theta + (k-1)\alpha)A(\theta + (k-2)\alpha) \cdots A(\theta)$$

and

$$(24) \quad A_{-k}(\theta) = A_k^{-1}(\theta - k\alpha)$$

for $k \geq 1$, where $A(\theta) = \begin{pmatrix} E - 2\lambda \cos 2\pi\theta & -1 \\ 1 & 0 \end{pmatrix}$. A_k is called the (k -step) transfer matrix.

By the definition, for any $k \in \mathbb{Z}_+, m \in \mathbb{Z}$, one has that

$$(25) \quad \begin{pmatrix} \phi(k+m) \\ \phi(k+m-1) \end{pmatrix} = A_k(\theta + m\alpha) \begin{pmatrix} \phi(m) \\ \phi(m-1) \end{pmatrix}.$$

It is easy to check that for $k \in \mathbb{Z}_+$,

$$(26) \quad A_k(\theta) = \begin{pmatrix} P_k(\theta) & -P_{k-1}(\theta + \alpha) \\ P_{k-1}(\theta) & -P_{k-2}(\theta + \alpha) \end{pmatrix}.$$

The Lyapunov exponent is given by

$$(27) \quad L(E) = \lim_{k \rightarrow \infty} \frac{1}{k} \int_{\mathbb{R}/\mathbb{Z}} \ln \|A_k(\theta)\| d\theta.$$

The Lyapunov exponent can be computed precisely for E in the spectrum of $H_{\lambda,\alpha,\theta}$.

Lemma 3.1. [15] For $E \in \Sigma_{\lambda,\alpha}$ and $|\lambda| > 1$, we have $L(E) = \ln |\lambda|$.

In the following, let $L := \ln |\lambda|$ be the Lyapunov exponent of the almost Mathieu operator for energies in the spectrum.

By upper semicontinuity and unique ergodicity, one has

$$(28) \quad L = \limsup_{k \rightarrow \infty} \sup_{\theta \in \mathbb{R}} \frac{1}{k} \ln \|A_k(\theta)\|.$$

Therefore, for any $\varepsilon > 0$,

$$(29) \quad \|A_k(\theta)\| \leq e^{(L+\varepsilon)k},$$

when k is large enough (independent of θ).

By (26) and (29), one has that for large k ,

$$(30) \quad |P_k(\theta)| \leq e^{(L+\varepsilon)k},$$

and hence for large $|x_2 - x_1|$,

$$(31) \quad |P_{[x_1, x_2]}(\theta)| \leq e^{(L+\varepsilon)|x_2 - x_1|}.$$

By (25) and (29), one has that for large $|k_1 - k_2|$,

$$(32) \quad \left\| \begin{pmatrix} \phi(k_1 + 1) \\ \phi(k_1) \end{pmatrix} \right\| \leq e^{(L+\varepsilon)|k_1 - k_2|} \left\| \begin{pmatrix} \phi(k_2 + 1) \\ \phi(k_2) \end{pmatrix} \right\|.$$

For any $x_1, x_2 \in \mathbb{Z}$ with $x_1 < x_2$, let $G_{[x_1, x_2]}$ be the Green's function:

$$G_{[x_1, x_2]} = (R_{[x_1, x_2]}(H_{\lambda, \alpha, \theta} - E)R_{[x_1, x_2]})^{-1}.$$

By Cramer's rule, for any $y \in [x_1, x_2]$, one has

$$(33) \quad |G_{[x_1, x_2]}(x_1, y)| = \left| \frac{P_{[y+1, x_2]}}{P_{[x_1, x_2]}} \right|,$$

$$(34) \quad |G_{[x_1, x_2]}(y, x_2)| = \left| \frac{P_{[x_1, y-1]}}{P_{[x_1, x_2]}} \right|.$$

It is easy to check that

$$(35) \quad \phi(y) = -G_{[x_1, x_2]}(x_1, y)\phi(x_1 - 1) - G_{[x_1, x_2]}(y, x_2)\phi(x_2 + 1).$$

Denote by $x'_1 = x_1 - 1$ and $x'_2 = x_2 + 1$.

By (33), (34) and (35), one has that for any $y \in [x_1, x_2]$,

$$(36) \quad |\phi(y)| \leq |P_{[x_1, x_2]}|^{-1} |P_{[x_1, y-1]}| |\phi(x'_2)| + |P_{[x_1, x_2]}|^{-1} |P_{[y+1, x_2]}| |\phi(x'_1)|.$$

Given a set $\{\theta_1, \dots, \theta_{k+1}\}$, the Lagrange interpolation terms Lag_m , $m = 1, 2, \dots, k+1$, are defined by

$$(37) \quad \text{Lag}_m = \ln \max_{x \in [-1, 1]} \prod_{j=1, j \neq m}^{k+1} \frac{|x - \cos 2\pi\theta_j|}{|\cos 2\pi\theta_m - \cos 2\pi\theta_j|}.$$

Lemma 3.2. *Given a set $\{\theta_1, \dots, \theta_{k+1}\}$, there exists some θ_m in set $\{\theta_1, \dots, \theta_{k+1}\}$ such that*

$$\left| P_k \left(\theta_m - \frac{k-1}{2} \alpha \right) \right| \geq \frac{e^{kL - \text{Lag}_m}}{k+1}.$$

Assume k is odd. Set $I = [j_0 - \frac{k+1}{2} + 1, j_0 + \frac{k+1}{2} - 1] = [x_1, x_2]$. Assume that $|P_{[x_1, x_2]}| \geq e^{kL - \mu}$. By (31) and (36), for any $y \in [x_1, x_2]$ with large enough $|y - x_1|$ and $|y - x_2|$, one has that

$$\begin{aligned} |\phi(y)| &\leq \sum_{i=1,2} e^{(L+\varepsilon)(k-|y-x_i|) - kL + \mu} |\phi(x'_i)| \\ (38) \quad &\leq e^\mu \sum_{i=1,2} |\phi(x'_i)| e^{\varepsilon k - L|y-x_i|}. \end{aligned}$$

4. TECHNICAL PREPARATIONS

The estimates in the following proposition will be constantly used in the proof.

Proposition 4.1. *Let θ be given by (17). When j is large enough, we have that*

$$(1) \text{ for any } k = k_j + l_1 q_{2j-1} \text{ with } 1 \leq |l_1| \leq \frac{q_{2j}}{10},$$

$$(39) \quad \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_{2j}};$$

$$(2) \text{ for any } k = k_j + l_1 q_{2j-1} + l_2 \text{ with } |l_1| \leq \frac{q_{2j}}{10q_{2j-1}} \text{ and } 1 \leq l_2 < q_{2j-1},$$

$$(40) \quad \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_{2j-1}};$$

$$(3) \text{ for any } k = k_j + l_1 q_{2j} \text{ with } |l_1| \leq \frac{\eta q_{2j+1}}{30q_{2j}},$$

$$(41) \quad \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{\eta}{20q_{2j}};$$

$$(4) \text{ for any } k = k_j + l_1 q_{2j} + l_2 \text{ with } |l_1| \leq \frac{10\eta q_{2j+1}}{q_{2j}} \text{ and } 1 \leq l_2 < q_{2j},$$

$$(42) \quad \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_{2j}};$$

$$(5) \text{ for any } k = l_1 q_n + l_2 \text{ with } |l_1| \leq \frac{q_{n+1}}{10q_n} \text{ and } 1 \leq l_2 < q_n,$$

$$(43) \quad \|k\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n}.$$

Proof. The proof is based on (8), (9) and (16). By direct computations, we have that in (1),

$$\begin{aligned}
\|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} &= \|2\theta + k_j\alpha + l_1q_{2j-1}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq |l_1| \|q_{2j-1}\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|2\theta + k_j\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \frac{|l_1|}{2q_{2j}} - \frac{10\eta}{q_{2j}} \\
&\geq \frac{1}{4q_{2j}},
\end{aligned}$$

in (2)

$$\begin{aligned}
\|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} &= \|2\theta + k_j\alpha + l_1q_{2j-1}\alpha + l_2\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \|l_2\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|2\theta + k_j\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|l_1q_{2j-1}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \frac{1}{2q_{2j-1}} - \frac{10\eta}{q_{2j}} - \frac{|l_1|}{q_{2j}} \\
&\geq \frac{1}{2q_{2j-1}} - \frac{10\eta}{q_{2j}} - \frac{1}{10q_{2j-1}} \\
&\geq \frac{1}{4q_{2j-1}},
\end{aligned}$$

in (3),

$$\begin{aligned}
\|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} &= \|2\theta + k_j\alpha + l_1q_{2j}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \|2\theta + k_j\alpha\|_{\mathbb{R}/\mathbb{Z}} - |l_1| \|q_{2j}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \frac{\eta}{10q_{2j}} - \frac{|l_1|}{q_{2j+1}} \\
&\geq \frac{\eta}{20q_{2j}},
\end{aligned}$$

in (4)

$$\begin{aligned}
\|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} &= \|2\theta + k_j\alpha + l_1q_{2j}\alpha + l_2\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \|l_2\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|2\theta + k_j\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|l_1q_{2j}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \frac{1}{2q_{2j}} - \frac{10\eta}{q_{2j}} - \frac{|l_1|}{q_{2j+1}} \\
&\geq \frac{1}{2q_{2j}} - \frac{20\eta}{q_{2j}} \\
&\geq \frac{1}{4q_{2j}},
\end{aligned}$$

and in (5),

$$\begin{aligned}
 \|k\alpha\|_{\mathbb{R}/\mathbb{Z}} &= \|l_1 q_n \alpha + l_2 \alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 &\geq \|l_2 \alpha\|_{\mathbb{R}/\mathbb{Z}} - \|l_1 q_n \alpha\|_{\mathbb{R}/\mathbb{Z}} \\
 &\geq \frac{1}{2q_n} - \frac{|l_1|}{q_{n+1}} \\
 &\geq \frac{1}{4q_n}.
 \end{aligned}$$

□

Lemma 4.2. *Let θ be defined by (17). Then*

$$(44) \quad \delta(\alpha, \theta) = \limsup_{j \rightarrow \infty} -\frac{\ln \|2\theta + k_j \alpha\|_{\mathbb{R}/\mathbb{Z}}}{k_j} = \limsup_{j \rightarrow \infty} \frac{\ln q_{2j}}{k_j}$$

and

$$(45) \quad (1 - 10^{-14}) \frac{\beta}{\eta} \leq \delta(\alpha, \theta) \leq (1 + 10^{-14}) \frac{\beta}{\eta}.$$

Proof. By (15), one has that for $j \geq j_0$,

$$(46) \quad -q_{2j} \leq k_{j+1} - k_j - \eta q_{2j+1} \leq 0.$$

By (46) and inductions, we have that for large j ,

$$(47) \quad |k_j - \eta q_{2j-1}| \leq 100q_{2j-2} \leq 10^{-15} \eta q_{2j-1}.$$

This implies that

$$(48) \quad \begin{aligned}
 \delta(\alpha, \theta) &= \limsup_{k \rightarrow \infty} -\frac{\ln \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}}}{|k|} \\
 &\geq \limsup_{j \rightarrow \infty} -\frac{\ln \|2\theta + k_j \alpha\|_{\mathbb{R}/\mathbb{Z}}}{k_j}
 \end{aligned}$$

$$(49) \quad \geq (1 - 10^{-14}) \frac{\beta}{\eta}.$$

Assume k satisfies $k_j < |k| \leq k_{j+1}$.

Case 1: $|k| \geq 10^{-3} \eta q_{2j+1}$ and $k \neq k_{j+1}$

In this case, by (8), (9) and (16), one has

$$\begin{aligned}
\|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} &= \|2\theta + k_{j+1}\alpha + (k - k_{j+1})\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \|(k - k_{j+1})\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|2\theta + k_{j+1}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \|q_{2j}\alpha\|_{\mathbb{R}/\mathbb{Z}} - \|2\theta + k_{j+1}\alpha\|_{\mathbb{R}/\mathbb{Z}} \\
&\geq \frac{1}{2q_{2j+1}} - \frac{10\eta}{q_{2j+2}} \\
&\geq \frac{1}{4q_{2j+1}}.
\end{aligned}$$

Case 2: $k_j < |k| < 10^{-3}\eta q_{2j+1}$

Write k as $k = k_j + l_1 q_{2j} + l_2$, where $l_1 \in \mathbb{Z}$ with $|l_1| \leq \frac{10^{-3}\eta q_{2j+1}}{q_{2j}} + 2$ and $0 \leq l_2 < q_{2j}$.

By (41) and (42), one has that

$$\|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{\eta}{20q_{2j}}.$$

Putting both cases together, we have that

$$\begin{aligned}
\delta(\alpha, \theta) &= \limsup_{k \rightarrow \infty} \frac{\ln \|2\theta + k\alpha\|_{\mathbb{R}/\mathbb{Z}}}{|k|} \\
(50) \quad &\leq \limsup_{j \rightarrow \infty} \frac{\ln q_{2j}}{k_j}
\end{aligned}$$

$$(51) \quad \leq (1 + 10^{-14}) \frac{\beta}{\eta}.$$

We conclude that (44) follows from (48) and (50), and (45) follows from (49) and (51). \square

Lemma 4.3. *Let θ be defined by (17). Assume $\liminf a_n = \infty$. Then*

$$(52) \quad \delta(\alpha, \theta) = \frac{\beta}{\eta}.$$

Proof. By the first inequality of (47) and the fact that $\liminf a_n = \infty$, one has that

$$(53) \quad \lim_{j \rightarrow \infty} \frac{k_j}{q_{2j-1}} = \eta.$$

Now (52) follows from (44) and (53). \square

5. NON-RESONANT SITES

In the following, we always assume

- θ is given by (17),

- $L = \ln |\lambda| > \delta(\alpha, \theta)$,
- $\varepsilon > 0$ is an arbitrarily small constant.
- C is a large constant (depends on λ and α) and it may change even in the same equation,
- n is large enough which depends on all parameters and constants.

Let $b_n = 10^{-7}\eta q_n$. For any $\ell \in \mathbb{Z}$, let

$$r_\ell^{\varepsilon, n} = \sup_{|r| \leq 10\varepsilon} |\phi(\ell q_n + r q_n)|,$$

and for $n = 2j - 1$,

$$r_{\ell+\eta}^{\varepsilon, n} = \sup_{|r| \leq 10\varepsilon} |\phi(\ell q_n + k_j + r q_n)|.$$

From (47), one can see that $k_j \approx \eta q_n$. So $r_{\ell+\eta}^{\varepsilon, n}$ is essentially the value of $|\phi(y)|$ with $y \approx (\ell + \eta)q_n$.

Lemma 5.1. *Assume that $|\lambda| > 1$. Let ℓ be such that $0 \leq |\ell| \leq 50 \frac{b_{n+1}}{q_n}$. Then for sufficiently large n , the following statements hold:*

- when $n = 2j - 1$ (namely, n is odd), we have that for any $y \in [\ell q_n + 10\varepsilon q_n, \ell q_n + k_j - 10\varepsilon q_n]$,

(54)

$$|\phi(y)| \leq r_\ell^{\varepsilon, n} \exp\{-(L-\varepsilon)(|y-\ell q_n|-3\varepsilon q_n)\} + r_{\ell+\eta}^{\varepsilon, n} \exp\{-(L-\varepsilon)(|\ell q_n+k_j-y|-3\varepsilon q_n)\},$$

and for any $y \in [\ell q_n + k_j + 10\varepsilon q_n, \ell q_n + q_n - 10\varepsilon q_n]$,

(55)

$$|\phi(y)| \leq r_{\ell+\eta}^{\varepsilon, n} \exp\{-(L-\varepsilon)(|y-\ell q_n-k_j|-3\varepsilon q_n)\} + r_{\ell+1}^{\varepsilon, n} \exp\{-(L-\varepsilon)(|\ell q_n+q_n-y|-3\varepsilon q_n)\};$$

- when $n = 2j$ (namely, n is even), we have that for any $y \in [\ell q_n + 10\varepsilon q_n, (\ell + 1)q_n - 10\varepsilon q_n]$,

(56)

$$|\phi(y)| \leq r_\ell^{\varepsilon, n} \exp\{-(L-\varepsilon)(|y-\ell q_n|-3\varepsilon q_n)\} + r_{\ell+1}^{\varepsilon, n} \exp\{-(L-\varepsilon)(|\ell q_n+q_n-y|-3\varepsilon q_n)\}.$$

Proof. Assume $n = 2j - 1$ first. For any $p \in [\ell q_n, \ell q_n + q_n]$ satisfying $|p - \ell q_n| \geq \varepsilon q_n$, $|p - \ell q_n - q_n| \geq \varepsilon q_n$ and $|p - \ell q_n - k_j| \geq \varepsilon q_n$, let

$$d_p = \frac{1}{100} \min\{|p - \ell q_n|, |p - \ell q_n - q_n|, |p - \ell q_n - k_j|, 10^{-5}\eta q_n\}$$

Let n_0 be the smallest integer such that

$$2q_{n-n_0} \leq d_p,$$

and let s be the largest positive integer such that $2sq_{n-n_0} \leq d_p$. Notice that $2(s+1)q_{n-n_0} \geq d_p$, one has that

$$(57) \quad sq_{n-n_0} \geq \frac{1}{4}d_p \geq \frac{1}{400}\varepsilon q_n.$$

Case 1: $p \in [\ell q_n + \varepsilon q_n, \ell q_n + k_j - \varepsilon q_n]$.

If $p \leq \ell q_n + \frac{k_j}{2}$, we construct intervals

$$I_1 = [-2sq_{n-n_0}, 2sq_{n-n_0} - 1], I_2 = [p - 2sq_{n-n_0}, p - 1].$$

If $p > \ell q_n + \frac{k_j}{2}$, we construct intervals

$$I_1 = [-2sq_{n-n_0}, 2sq_{n-n_0} - 1], I_2 = [p + 1, p + 2sq_{n-n_0}].$$

Case 2: $p \in [\ell q_n + k_j + \varepsilon q_n, \ell q_n + q_n - \varepsilon q_n]$.

If $p \leq \ell q_n + \frac{k_j}{2} + \frac{1}{2}q_n$, we construct intervals

$$I_1 = [-2sq_{n-n_0}, 2sq_{n-n_0} - 1], I_2 = [p - 2sq_{n-n_0}, p - 1].$$

If $p > \ell q_n + \frac{k_j}{2} + \frac{1}{2}q_n$, we construct intervals

$$I_1 = [-2sq_{n-n_0}, 2sq_{n-n_0} - 1], I_2 = [p + 1, p + 2sq_{n-n_0}].$$

By the construction of I_1 and I_2 , one has that for any $j_1, j_2 \in I_1 \cup I_2$ with $j_1 \neq j_2$, there exist l_1 and l_2 with $|l_1| \leq 100\frac{b_{n+1}}{q_n} + 4$ and $1 \leq l_2 < q_n$, and l'_1 and l'_2 with $|l'_1| \leq 100\frac{b_{n+1}}{q_n} + 4$ and $1 \leq l'_2 < q_n$ such that

$$j_1 - j_2 = l_1 q_n + l_2,$$

and

$$j_1 + j_2 = k_j + l'_1 q_n + l'_2.$$

Therefore, by (40) and (43), we have that that for any $j_1, j_2 \in I_1 \cup I_2$ with $j_1 \neq j_2$,

$$(58) \quad \|(j_1 - j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n}$$

and

$$(59) \quad \|2\theta + (j_1 + j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n}.$$

Assume $n = 2j$.

For any $p \in [\ell q_n, \ell q_n + q_n]$ satisfying $|p - \ell q_n| \geq \varepsilon q_n$ and $|p - \ell q_n - q_n| \geq \varepsilon q_n$, let

$$d_p = \frac{1}{100} \min\{|p - \ell q_n|, |p - \ell q_n - q_n|, 10^{-5}\eta q_n\}$$

Let n_0 be the smallest integer such that

$$2q_{n-n_0} \leq d_p,$$

and let s be the largest positive integer such that $2sq_{n-n_0} \leq d_p$. Notice that $2(s+1)q_{n-n_0} > d_p$, one has that

$$(60) \quad sq_{n-n_0} \geq \frac{1}{4}d_p \geq \frac{1}{400}\varepsilon q_n.$$

We construct intervals

$$I_1 = [-sq_{n-n_0}, sq_{n-n_0} - 1], I_2 = [p - sq_{n-n_0}, p + sq_{n-n_0} - 1].$$

In this case, one has that for any $j_1, j_2 \in I_1 \cup I_2$ and $j_1 \neq j_2$,

$$j_1 - j_2 = l_1 q_n + l_2,$$

where $|l_1| \leq 100 \frac{b_{n+1}}{q_n} + 4$ and $1 \leq l_2 < q_n$. Therefore, by (43), we have that

$$(61) \quad \|(j_1 - j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n}.$$

and by (41) and (42), we have that

$$(62) \quad \|2\theta + (j_1 + j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{\eta}{20q_n}.$$

In all cases, let $\theta_m = \theta + m\alpha$ for $m \in I_1 \cup I_2$. By the small divisor conditions (58), (59), (61) and (62), and modifying the proof of [7, Lemma 9.9] (or Appendices in [25] and [33]), we can prove that for any $m \in I_1 \cup I_2$, one has that $\text{Lag}_m \leq \varepsilon q_n$. Now Lemma 5.1 follows from standard block expansions (e.g., [25, Lemma 4.1] and [26, Lemma 3.4]). \square

6. RESONANT SITES AND PROOF OF THEOREM 2.1

We are going to deal with the resonances $\ell q_n + k_j$ and ℓq_n first. For simplicity, we drop the dependence of n and ε in notations $r_\ell^{\varepsilon, n}$ and $r_{\ell+\eta}^{\varepsilon, n}$. Denote by $\beta_n = \frac{\ln q_{n+1}}{q_n}$. Clearly, for large n , $\beta_n \leq \beta + \varepsilon$. Since $L > \delta$, by Lemma 4.2, one has that for large j ,

$$(63) \quad L > \frac{\beta_{2j-1} q_{2j-1}}{k_j},$$

and for large n

$$(64) \quad L > (1 - 10^{-14}) \frac{\beta}{\eta} > (1 - 10^{-13}) \frac{\beta_n}{\eta}.$$

Theorem 6.1. *Assume that $n = 2j - 1$ and $|\ell| \leq 20 \frac{b_{n+1}}{q_n}$. Then*

$$(65) \quad r_{\ell+\eta} \leq e^{C\varepsilon q_n + \beta_n q_n} (e^{-Lk_j r_\ell} + e^{-Lq_n + Lk_j r_{\ell+1}}).$$

Proof. Take any p with $|p - (\ell q_n + k_j)| \leq 10\varepsilon q_n$ into consideration. Let n_0 be the least positive integer such that

$$q_{n-n_0} \leq \frac{\varepsilon}{2} \left(\frac{k_j}{2} - 2\varepsilon q_n \right).$$

Let s be the largest positive integer such that $sq_{n-n_0} \leq \frac{k_j}{2} - 2\varepsilon q_n$. By the fact that $(s+1)q_{n-n_0} \geq \frac{k_j}{2} - 2\varepsilon q_n$, one has

$$s \geq \frac{1}{\varepsilon}$$

and

$$(66) \quad \frac{k_j}{2} - 3\varepsilon q_n \leq sq_{n-n_0} \leq \frac{k_j}{2} - 2\varepsilon q_n.$$

Construct intervals

$$I_1 = [-sq_{n-n_0}, sq_{n-n_0} - 1], I_2 = [\ell q_n + k_j - sq_{n-n_0}, \ell q_n + k_j + sq_{n-n_0} - 1].$$

Let $\theta_m = \theta + m\alpha$ for $m \in I_1 \cup I_2$. The set $\{\theta_m\}_{m \in I_1 \cup I_2}$ consists of $4sq_{n-n_0}$ elements. Let $k = 4sq_{n-n_0} - 1$. By modifying the proof of [7, Lemma 9.9] and [36, Lemma 4.1] (or Appendices in [25] and [33]), we can prove the claim (Claim 1): for any $m \in I_1 \cup I_2$, one has $\text{Lag}_m \leq (\beta_n + \varepsilon)q_n$. For convenience, we include a proof in Section 7.

Applying Lemma 3.2, there exists some j_0 with $j_0 \in I_1 \cup I_2$ such that

$$(67) \quad \left| P_k \left(\theta_{j_0} - \frac{k-1}{2}\alpha \right) \right| \geq e^{kL - (\beta_n + \varepsilon)q_n}.$$

First, assume $j_0 \in I_2$.

Set $I = [j_0 - 2sq_{n-n_0} + 1, j_0 + 2sq_{n-n_0} - 1] = [x_1, x_2]$. By (38),

$$(68) \quad |\phi(p)| \leq \sum_{i=1,2} e^{\beta_n q_n + C\varepsilon q_n} |\phi(x'_i)| e^{-L|p-x_i|},$$

where $x'_1 = x_1 - 1$ and $x'_2 = x_2 + 1$.

For simplicity, we are not going to make the difference between $a \in \mathbb{Z}$ and $\tilde{a} \in \mathbb{Z}$ if $|a - \tilde{a}| \leq 100\varepsilon q_n$. Clearly,

$$(69) \quad x'_1 \in \left[\ell q_n - \frac{1}{2}k_j, \ell q_n + \frac{1}{2}k_j \right], x'_2 \in \left[\ell q_n + \frac{3}{2}k_j, \ell q_n + \frac{5}{2}k_j \right].$$

By Lemma 5.1, one has that

$$(70) \quad |\phi(x'_2)| \leq r_{\ell+\eta} e^{C\varepsilon q_n - L|x_2 - \ell q_n - k_j|} + r_{\ell+1} e^{C\varepsilon q_n - L|\ell q_n + q_n - x_2|}.$$

By Lemma 5.1 again, one has that for any $x'_1 \in [\ell q_n, \ell q_n + \frac{1}{2}k_j]$,

$$(71) \quad |\phi(x'_1)| \leq r_{\ell+\eta} e^{C\varepsilon q_n - L|\ell q_n + k_j - x_1|} + r_{\ell} e^{C\varepsilon q_n - L|x_1 - \ell q_n|},$$

and for any $x'_1 \in [\ell q_n - \frac{1}{2}k_j, \ell q_n]$,

$$(72) \quad |\phi(x'_1)| \leq r_{\ell-1+\eta} e^{C\varepsilon q_n - L|x_1 - \ell q_n + q_n - k_j|} + r_{\ell} e^{C\varepsilon q_n - L|\ell q_n - x_1|}.$$

By (68)-(72), one has

$$(73) \quad \begin{aligned} |\phi(p)| &\leq e^{C\varepsilon q_n - Lk_j + \beta_n q_n} (r_{\ell+\eta} + r_{\ell}) + e^{C\varepsilon q_n - Lq_n + \beta_n q_n} r_{\ell-1+\eta} \\ &\quad + e^{C\varepsilon q_n - Lq_n + \beta_n q_n + Lk_j} r_{\ell+1}. \end{aligned}$$

Therefore, we have

$$(74) \quad r_{\ell+\eta} \leq e^{C\varepsilon q_n - Lk_j + \beta_n q_n} (r_{\ell+\eta} + r_\ell) + e^{C\varepsilon q_n - Lq_n + \beta_n q_n} r_{\ell-1+\eta} + e^{C\varepsilon q_n - Lq_n + \beta_n q_n + Lk_j} r_{\ell+1}.$$

Since $L > \delta(\alpha, \theta) > \frac{\beta_n q_n}{k_j}$ (by (63)), one has

$$(75) \quad e^{C\varepsilon q_n - Lk_j + \beta_n q_n} \leq \frac{1}{2}.$$

By (74) and (75), we have that

$$(76) \quad r_{\ell+\eta} \leq e^{C\varepsilon q_n - Lk_j + \beta_n q_n} r_\ell + e^{C\varepsilon q_n - Lq_n + \beta_n q_n} r_{\ell-1+\eta} + e^{C\varepsilon q_n - Lq_n + \beta_n q_n + Lk_j} r_{\ell+1}$$

By (32), one has

$$(77) \quad r_{\ell-1+\eta} \leq e^{C\varepsilon q_n} e^{Lq_n - Lk_j} r_\ell.$$

By (76) and (77), one has that

$$(78) \quad r_{\ell+\eta} \leq e^{C\varepsilon q_n + \beta_n q_n} (e^{-Lk_j} r_\ell + e^{-Lq_n + Lk_j} r_{\ell+1}).$$

Thus in order to prove the theorem, it suffices to exclude the case $j_0 \in I_1$.

Suppose $j_0 \in I_1$. Following the proof of (73), we obtain (move $-\ell q_n - k_j$ units in (73))

$$|\phi(0)| \leq \frac{1}{2}.$$

This contradicts $\phi(0) = 1$. \square

Theorem 6.2. *Assume that $n = 2j - 1$ and $0 < |\ell| \leq 20 \frac{b_{n+1}}{q_n}$. Then*

$$(79) \quad r_\ell \leq e^{C\varepsilon q_n + 3\beta_n q_n} (e^{-Lq_n} r_{\ell+1} + e^{-Lq_n + Lk_j} r_{\ell-1+\eta}).$$

Proof. Take any p with $|p - \ell q_n| \leq 10\varepsilon q_n$ into consideration.

Construct intervals

$$I_1 = \left[-\left\lfloor \frac{q_n}{2} \right\rfloor, q_n - \left\lfloor \frac{q_n}{2} \right\rfloor - 1 \right],$$

and

$$I_2 = \left[\ell q_n - \left\lfloor \frac{q_n}{2} \right\rfloor, (\ell + 1)q_n - \left\lfloor \frac{q_n}{2} \right\rfloor - 1 \right].$$

Let $\theta_m = \theta + m\alpha$ for $m \in I_1 \cup I_2$. The set $\{\theta_m\}_{m \in I_1 \cup I_2}$ consists of $2q_n$ elements. Let $k = 2q_n - 1$. In this case, similar to Claim 1, we can prove the claim (Claim 2): for any $m \in I_1 \cup I_2$, one has $\text{Lag}_m \leq (3\beta_n + \varepsilon)q_n$. For convenience, we include a proof in Section 7.

Applying Lemma 3.2, there exists some j_0 with $j_0 \in I_1 \cup I_2$ such that

$$(80) \quad \left| P_k \left(\theta_{j_0} - \frac{k-1}{2} \alpha \right) \right| \geq e^{kL - (3\beta_n + \varepsilon)q_n}.$$

First, assume $j_0 \in I_2$.

Set $I = [j_0 - q_n + 1, j_0 + q_n - 1] = [x_1, x_2]$. By (38), one has

$$(81) \quad |\phi(p)| \leq \sum_{i=1,2} e^{3\beta_n q_n + C\varepsilon q_n} |\phi(x'_i)| e^{-L|p-x_i|},$$

where $x'_1 = x_1 - 1$ and $x'_2 = x_2 + 1$.

Clearly,

$$(82) \quad x'_1 \in \left[\ell q_n - \frac{3}{2}q_n, \ell q_n - \frac{1}{2}q_n \right], x'_2 \in \left[\ell q_n + \frac{1}{2}q_n, \ell q_n + \frac{3}{2}q_n \right].$$

By Lemma 5.1,

- for $x'_2 \in [\ell q_n + \frac{1}{2}q_n, \ell q_n + q_n]$,

$$(83) \quad |\phi(x'_2)| \leq r_{\ell+\eta} e^{C\varepsilon q_n - L|x_2 - \ell q_n - k_j|} + r_{\ell+1} e^{C\varepsilon q_n - L|\ell q_n + q_n - x_2|},$$

- for $x'_2 \in [\ell q_n + q_n, \ell q_n + q_n + k_j]$,

$$(84) \quad |\phi(x'_2)| \leq r_{\ell+1+\eta} e^{C\varepsilon q_n - L|\ell q_n + q_n + k_j - x_2|} + r_{\ell+1} e^{C\varepsilon q_n - L|x_2 - q_n - \ell q_n|}.$$

- for $x'_2 \in [\ell q_n + q_n + k_j, \ell q_n + \frac{3}{2}q_n]$,

$$(85) \quad |\phi(x'_2)| \leq r_{\ell+1+\eta} e^{C\varepsilon q_n - L|x_2 - \ell q_n - q_n - k_j|} + r_{\ell+2} e^{C\varepsilon q_n - L|\ell q_n + 2q_n - x_2|}.$$

By Lemma 5.1 again, one has

- for $x'_1 \in [\ell q_n - q_n + k_j, \ell q_n - \frac{1}{2}q_n]$,

$$(86) \quad |\phi(x'_1)| \leq r_{\ell-1+\eta} e^{C\varepsilon q_n - L|x_1 - \ell q_n + q_n - k_j|} + r_{\ell} e^{C\varepsilon q_n - L|\ell q_n - x_1|},$$

- for $x'_1 \in [\ell q_n - q_n, \ell q_n - q_n + k_j]$, one has

$$(87) \quad |\phi(x'_1)| \leq r_{\ell-1+\eta} e^{C\varepsilon q_n - L|\ell q_n - q_n + k_j - x_1|} + r_{\ell-1} e^{C\varepsilon q_n - L|x_1 - \ell q_n + q_n|}.$$

- for $x'_1 \in [\ell q_n - \frac{3}{2}q_n, \ell q_n - q_n]$ one has

$$(88) \quad |\phi(x'_1)| \leq r_{\ell-2+\eta} e^{C\varepsilon q_n - L|x_1 - \ell q_n + 2q_n - k_j|} + r_{\ell-1} e^{C\varepsilon q_n - L|\ell q_n - q_n - x_1|}.$$

By (81)-(88), one has

$$(89) \quad \begin{aligned} |\phi(p)| &\leq e^{C\varepsilon q_n - Lq_n + 3\beta_n q_n} r_{\ell} + e^{C\varepsilon q_n - Lq_n + Lk_j + 3\beta_n q_n} r_{\ell+\eta} \\ &\quad + e^{C\varepsilon q_n - Lq_n - Lk_j + 3\beta_n q_n} r_{\ell+1+\eta} + e^{C\varepsilon q_n - Lq_n + 3\beta_n q_n} r_{\ell+1} \\ &\quad + e^{C\varepsilon q_n - 2Lq_n + 3\beta_n q_n} r_{\ell+2} + e^{C\varepsilon q_n - Lq_n + Lk_j + 3\beta_n q_n} r_{\ell-1+\eta} \\ &\quad + e^{C\varepsilon q_n - Lq_n + 3\beta_n q_n} r_{\ell-1} + e^{C\varepsilon q_n - 2Lq_n + Lk_j + 3\beta_n q_n} r_{\ell-2+\eta}. \end{aligned}$$

Therefore, we have

$$(90) \quad \begin{aligned} r_{\ell} &\leq e^{C\varepsilon q_n - Lq_n + 3\beta_n q_n} r_{\ell} + e^{C\varepsilon q_n - Lq_n + Lk_j + 3\beta_n q_n} r_{\ell+\eta} \\ &\quad + e^{C\varepsilon q_n - Lq_n - Lk_j + 3\beta_n q_n} r_{\ell+1+\eta} + e^{C\varepsilon q_n - Lq_n + 3\beta_n q_n} r_{\ell+1} \\ &\quad + e^{C\varepsilon q_n - 2Lq_n + 3\beta_n q_n} r_{\ell+2} + e^{C\varepsilon q_n - Lq_n + Lk_j + 3\beta_n q_n} r_{\ell-1+\eta} \\ &\quad + e^{C\varepsilon q_n - Lq_n + 3\beta_n q_n} r_{\ell-1} + e^{C\varepsilon q_n - 2Lq_n + Lk_j + 3\beta_n q_n} r_{\ell-2+\eta}. \end{aligned}$$

By (32), one has

$$(91) \quad r_{\ell+\eta} \leq e^{C\varepsilon q_n} e^{Lk_j} r_\ell, r_{\ell+1+\eta} \leq e^{C\varepsilon q_n} e^{Lk_j} r_{\ell+1}, r_{\ell+2} \leq e^{C\varepsilon q_n} e^{Lq_n} r_{\ell+1},$$

and

$$(92) \quad r_{\ell-1} \leq e^{C\varepsilon q_n} e^{Lk_j} r_{\ell-1+\eta}, r_{\ell-2+\eta} \leq e^{C\varepsilon q_n} e^{Lq_n} r_{\ell-1+\eta},$$

By (90), (91) and (92), we have

$$(93) \quad \begin{aligned} r_\ell &\leq e^{C\varepsilon q_n - Lq_n + 2Lk_j + 3\beta_n q_n} r_\ell + e^{C\varepsilon q_n - Lq_n + 3\beta_n q_n} r_{\ell+1} \\ &\quad + e^{C\varepsilon q_n - Lq_n + Lk_j + 3\beta_n q_n} r_{\ell-1+\eta}. \end{aligned}$$

By (64), one has

$$(94) \quad e^{C\varepsilon q_n - Lq_n + 2Lk_j + 3\beta_n q_n} \leq \frac{1}{2}.$$

By (93) and (94), one has

$$(95) \quad r_\ell \leq e^{C\varepsilon q_n - Lq_n + 3\beta_n q_n} r_{\ell+1} + e^{C\varepsilon q_n - Lq_n + Lk_j + 3\beta_n q_n} r_{\ell-1+\eta}.$$

Thus to prove the theorem, it suffices to exclude the case $j_0 \in I_1$. Suppose $j_0 \in I_1$. Following the proof of (95), we have

$$|\phi(0)| \leq \frac{1}{2}.$$

This contradicts $\phi(0) = 1$. □

Theorem 6.3. *Assume that $n = 2j$ is even and $0 < |\ell| \leq 20 \frac{b_{n+1}}{q_n}$. Then*

$$(96) \quad r_\ell \leq e^{C\varepsilon q_n - Lq_n + \beta_n q_n} (r_{\ell-1} + r_{\ell+1}).$$

Proof. Take any p with $|p - \ell q_n| \leq 10\varepsilon q_n$ into consideration.

Construct intervals

$$I_1 = \left[-\left\lfloor \frac{q_n}{2} \right\rfloor, q_n - \left\lfloor \frac{q_n}{2} \right\rfloor - 1 \right],$$

and

$$I_2 = \left[\ell q_n - \left\lfloor \frac{q_n}{2} \right\rfloor, (\ell + 1)q_n - \left\lfloor \frac{q_n}{2} \right\rfloor - 1 \right].$$

Let $\theta_m = \theta + m\alpha$ for $m \in I_1 \cup I_2$. The set $\{\theta_m\}_{m \in I_1 \cup I_2}$ consists of $2q_n$ elements. Let $k = 2q_n - 1$. In this case, similar to Claims 1 and 2, we can prove the claim (Claim 3): for any $m \in I_1 \cup I_2$, one has $\text{Lag}_m \leq (\beta_n + \varepsilon)q_n$. For convenience, we include a proof in Section 7.

Applying Lemma 3.2, there exists some j_0 with $j_0 \in I_1 \cup I_2$ such that

$$(97) \quad \left| P_k \left(\theta_{j_0} - \frac{k-1}{2} \alpha \right) \right| \geq e^{kL - (\beta_n + \varepsilon)q_n}.$$

First assume $j_0 \in I_2$.

Set $I = [j_0 - q_n + 1, j_0 + q_n - 1] = [x_1, x_2]$. By (38), one has that

$$(98) \quad |\phi(p)| \leq \sum_{i=1,2} e^{\beta_n q_n + C\varepsilon q_n} |\phi(x'_i)| e^{-L|p-x_i|},$$

where $x'_1 = x_1 - 1$ and $x'_2 = x_2 + 1$.

Clearly,

$$(99) \quad x'_1 \in \left[\ell q_n - \frac{3}{2}q_n, \ell q_n - \frac{1}{2}q_n \right], x'_2 \in \left[\ell q_n + \frac{1}{2}q_n, \ell q_n + \frac{3}{2}q_n \right].$$

By Lemma 5.1, one has that

- for $x'_2 \in [\ell q_n + \frac{1}{2}q_n, \ell q_n + q_n]$,

$$(100) \quad |\phi(x'_2)| \leq r_\ell e^{C\varepsilon q_n - L|x_2 - \ell q_n|} + r_{\ell+1} e^{C\varepsilon q_n - L|\ell q_n + q_n - x_2|},$$

- for $x'_2 \in [\ell q_n + q_n, \ell q_n + \frac{3}{2}q_n]$ one has

$$(101) \quad |\phi(x'_2)| \leq r_{\ell+1} e^{C\varepsilon q_n - L|x_2 - \ell q_n - q_n|} + r_{\ell+2} e^{C\varepsilon q_n - L|\ell q_n + 2q_n - x_2|}.$$

By Lemma 5.1 again, one has that

- for $x'_1 \in [\ell q_n - q_n, \ell q_n - \frac{1}{2}q_n]$,

$$(102) \quad |\phi(x'_1)| \leq r_{\ell-1} e^{C\varepsilon q_n - L|x_1 - \ell q_n + q_n|} + r_\ell e^{C\varepsilon q_n - L|\ell q_n - x_1|},$$

- for $x'_1 \in [\ell q_n - \frac{3}{2}q_n, \ell q_n - q_n]$,

$$(103) \quad |\phi(x'_1)| \leq r_{\ell-2} e^{C\varepsilon q_n - L|x_1 - \ell q_n + 2q_n|} + r_{\ell-1} e^{C\varepsilon q_n - L|\ell q_n - q_n - x_1|}.$$

By (98)-(103), one has that

$$(104) \quad |\phi(p)| \leq e^{C\varepsilon q_n - Lq_n + \beta_n q_n} (r_\ell + r_{\ell+1} + r_{\ell-1}) + e^{C\varepsilon q_n - 2Lq_n + \beta_n q_n} (r_{\ell+2} + r_{\ell-2}).$$

Therefore, we have that

$$(105) \quad r_\ell \leq e^{C\varepsilon q_n - Lq_n + \beta_n q_n} (r_\ell + r_{\ell+1} + r_{\ell-1}) + e^{C\varepsilon q_n - 2Lq_n + \beta_n q_n} (r_{\ell+2} + r_{\ell-2}).$$

Since $L > \frac{\beta}{2\eta}$, one has that

$$e^{C\varepsilon q_n - Lq_n + \beta_n q_n} \leq \frac{1}{2}.$$

By (105), we have that

$$(106) \quad r_\ell \leq e^{C\varepsilon q_n - Lq_n + \beta_n q_n} (r_{\ell+1} + r_{\ell-1}) + e^{C\varepsilon q_n - 2Lq_n + \beta_n q_n} (r_{\ell+2} + r_{\ell-2}).$$

By (32), one has

$$(107) \quad r_{\ell+2} \leq e^{C\varepsilon q_n} e^{Lq_n} r_{\ell+1}, r_{\ell-2} \leq e^{C\varepsilon q_n} e^{Lq_n} r_{\ell-1}.$$

By (106) and (107), one has that

$$(108) \quad r_\ell \leq e^{C\varepsilon q_n - Lq_n + \beta_n q_n} (r_{\ell+1} + r_{\ell-1}).$$

Thus in order to prove the theorem, it suffices to exclude the case $j_0 \in I_1$.

Suppose $j_0 \in I_1$. Following the proof of (104), we have that

$$|\phi(0)| \leq \frac{1}{2}.$$

This contradicts $\phi(0) = 1$. \square

Proof of Theorem 2.1. Once we have Theorems 6.1, 6.2, 6.3 and Lemma 5.1 at hand, Theorem 2.1 follows from standard iterations. See [25, 33, 34] for example. \square

7. PROOF OF COROLLARIES AND CLAIMS 1-3

Proof of Corollary 1.1. It follows from Remark 1.3, (45) and Theorem 2.1 with $\eta = 10^{-2}$. \square

Proof of Corollary 1.2. Let $a_0 = 0$, $a_1 = \frac{2}{\mu} + 10$, $q_{-1} = 0$ and $q_0 = 1$. For $n \geq 1$, define a_n and q_n inductively by

$$(109) \quad q_n = a_n q_{n-1} + q_{n-2}.$$

and

$$(110) \quad a_{n+1} = \lfloor e^{\mu q_n} \rfloor.$$

Let $\alpha = [a_1, a_2, \dots, a_n, \dots]$. By some simple facts of the continued fraction expansion, one has that $\beta(\alpha) = \mu$ (see Section 2 for more details). Now Corollary 1.2 follows from Corollary 1.1. \square

Proof of Corollary 1.4. It follows from Theorem 2.1, Lemma 4.3 and the construction of α in Corollary 1.2. \square

Proof of Claim 1. By the construction of I_1 and I_2 in Claim 1, we have the following estimates:

- for any $j_1, j_2 \in I_1 \cup I_2$ with $j_1 \neq j_2$, one has that

$$(111) \quad j_1 - j_2 = l_1 q_n + l_2,$$

where $1 \leq l_2 < q_n$ and $|l_1| \leq 40 \frac{b_{n+1}}{q_n} + 4$. Therefore, by (43),

$$(112) \quad \|(j_1 - j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n}.$$

- for any $j_1, j_2 \in I_1 \cup I_2$ with $j_1 \neq j_2$, one has

$$(113) \quad (j_1 + j_2) = k_j + l_1 q_n + l_2,$$

where $0 \leq l_2 < q_n$ and $|l_1| \leq 40\frac{b_{n+1}}{q_n} + 4$. Therefore by (40), for $l_2 \geq 1$,

$$(114) \quad \|2\theta + (j_1 + j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n},$$

and by (39) and (16), for $l_2 = 0$

$$(115) \quad \|2\theta + (j_1 + j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{\eta}{10q_{n+1}} \geq e^{-(\beta_n + \varepsilon)q_n}.$$

Moreover, for $j_1 \in I_1 \cup I_2$, there is at most one $j_2 \in I_1 \cup I_2$ such that the l_2 in (113) is 0. By the standard arguments (e.g. Appendices in [25, 33]), we have that for any $m \in I_1 \cup I_2$,

$$\text{Lag}_m \leq \beta_n q_n + \varepsilon q_n.$$

□

Proof of Claim 2. By the construction of I_1 and I_2 in Claim 2, and Proposition 4.1, we have the following estimates:

- for any $j_1, j_2 \in I_1 \cup I_2$ with $j_1 \neq j_2$, one has that

$$(116) \quad j_1 - j_2 = l_1 q_n + l_2,$$

where $0 \leq l_2 < q_n$ and $|l_1| \leq 40\frac{b_{n+1}}{q_n} + 4$. Therefore, for $l_2 \neq 0$

$$(117) \quad \|(j_1 - j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n},$$

and for $l_2 = 0$,

$$(118) \quad \|(j_1 - j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{2q_{n+1}} \geq e^{-(\beta_n + \varepsilon)q_n}.$$

- for any $j_1, j_2 \in I_1 \cup I_2$ with $j_1 \neq j_2$, one has

$$(119) \quad (j_1 + j_2) = k_j + l_1 q_n + l_2,$$

where $0 \leq l_2 < q_n$ and $|l_1| \leq 40\frac{b_{n+1}}{q_n} + 4$. Therefore, for $l_2 \neq 0$

$$(120) \quad \|2\theta + (j_1 + j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n},$$

and for $l_2 = 0$

$$(121) \quad \|2\theta + (j_1 + j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{\eta}{10q_{n+1}} \geq e^{-(\beta_n + \varepsilon)q_n}.$$

Moreover, for any $j_1 \in I_1 \cup I_2$, there is at most one $j_2 \in I_1 \cup I_2$ with $j_2 \neq j_1$ such that the l_2 in (116) is 0. For any $j_1 \in I_1 \cup I_2$, there is at most two $j_2 \in I_1 \cup I_2$ with $j_2 \neq j_1$ such that the l_2 in (119) is 0. Following the discussion in Claim 1, we have that for any $m \in I_1 \cup I_2$,

$$\text{Lag}_m \leq 3\beta_n q_n + \varepsilon q_n.$$

□

Proof of Claim 3. By the construction of I_1 and I_2 in Claim 3, and Proposition 4.1, we have the following estimates:

- for any $j_1, j_2 \in I_1 \cup I_2$ with $j_1 \neq j_2$, one has that

$$(122) \quad j_1 - j_2 = l_1 q_n + l_2,$$

where $0 \leq l_2 < q_n$ and $|l_1| \leq 40 \frac{b_{n+1}}{q_n} + 4$. Therefore, for $l_2 \neq 0$

$$(123) \quad \|(j_1 - j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{4q_n},$$

and for $l_2 = 0$

$$(124) \quad \|(j_1 - j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{1}{2q_{n+1}} \geq e^{-(\beta_n + \varepsilon)q_n}.$$

- for any $j_1, j_2 \in I_1 \cup I_2$,

$$(125) \quad \|2\theta + (j_1 + j_2)\alpha\|_{\mathbb{R}/\mathbb{Z}} \geq \frac{\eta}{20q_n}.$$

Moreover, for $j_1 \in I_1 \cup I_2$, there is at most one $j_2 \in I_1 \cup I_2$ with $j_2 \neq j_1$ such that the l_2 in (122) is 0. Following the discussion in Claim 1, we have that for any $m \in I_1 \cup I_2$,

$$\text{Lag}_m \leq \beta_n q_n + \varepsilon q_n.$$

□

ACKNOWLEDGMENTS

W. L. was supported by NSF DMS-2000345 and DMS-2052572.

REFERENCES

- [1] S. Aubry and G. André. Analyticity breaking and Anderson localization in incommensurate lattices. *Ann. Israel Phys. Soc.*, 3(133):18, 1980.
- [2] A. Avila. KAM, Lyapunov exponents and the spectral dichotomy for one-frequency Schrödinger operators. *In preparation*.
- [3] A. Avila. The absolutely continuous spectrum of the almost Mathieu operator. *arXiv preprint arXiv:0810.2965*, 2008.
- [4] A. Avila. Almost reducibility and absolute continuity I. *arXiv preprint arXiv:1006.0704*, 2010.
- [5] A. Avila. Global theory of one-frequency Schrödinger operators. *Acta Math.*, 215(1):1–54, 2015.

- [6] A. Avila and D. Damanik. Absolute continuity of the integrated density of states for the almost Mathieu operator with non-critical coupling. *Invent. Math.*, 172(2):439–453, 2008.
- [7] A. Avila and S. Jitomirskaya. The Ten Martini Problem. *Ann. of Math. (2)*, 170(1):303–342, 2009.
- [8] A. Avila and S. Jitomirskaya. Almost localization and almost reducibility. *J. Eur. Math. Soc. (JEMS)*, 12(1):93–131, 2010.
- [9] A. Avila, S. Jitomirskaya, and C. A. Marx. Spectral theory of extended Harper’s model and a question by Erdős and Szekeres. *Invent. Math.*, 210(1):283–339, 2017.
- [10] A. Avila, S. Jitomirskaya, and Q. Zhou. Second phase transition line. *Math. Ann.*, 370(1-2):271–285, 2018.
- [11] A. Avila, J. You, and Q. Zhou. Sharp phase transitions for the almost Mathieu operator. *Duke Math. J.*, 166(14):2697–2718, 2017.
- [12] J. Avron and B. Simon. Singular continuous spectrum for a class of almost periodic Jacobi matrices. *Bull. Amer. Math. Soc. (N.S.)*, 6(1):81–85, 1982.
- [13] J. Bourgain. *Green’s function estimates for lattice Schrödinger operators and applications*, volume 158 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2005.
- [14] J. Bourgain and M. Goldstein. On nonperturbative localization with quasi-periodic potential. *Ann. of Math. (2)*, 152(3):835–879, 2000.
- [15] J. Bourgain and S. Jitomirskaya. Continuity of the Lyapunov exponent for quasiperiodic operators with analytic potential. volume 108, pages 1203–1218. 2002. Dedicated to David Ruelle and Yasha Sinai on the occasion of their 65th birthdays.
- [16] L. Ge and J. You. Arithmetic version of Anderson localization via reducibility. *Geom. Funct. Anal.*, 30(5):1370–1401, 2020.
- [17] L. Ge, J. You, and X. Zhao. The arithmetic version of the frequency transition conjecture: New proof and generalization. *Peking Mathematical Journal to appear*.
- [18] A. Y. Gordon. The point spectrum of the one-dimensional Schrödinger operator. *Uspekhi Matematicheskikh Nauk*, 31(4):257–258, 1976.
- [19] A. Y. Gordon, S. Jitomirskaya, Y. Last, and B. Simon. Duality and singular continuous spectrum in the almost Mathieu equation. *Acta Math.*, 178(2):169–183, 1997.
- [20] S. Jitomirskaya. One-dimensional quasiperiodic operators: global theory, duality, and sharp analysis of small denominators. *Proceedings, 2022 ICM*.

- [21] S. Jitomirskaya. Critical phenomena, arithmetic phase transitions, and universality: some recent results on the almost mathieu operator, 2019.
- [22] S. Jitomirskaya. On point spectrum of critical almost Mathieu operators. *Adv. Math.*, 392:Paper No. 107997, 6, 2021.
- [23] S. Jitomirskaya and I. Kachkovskiy. L^2 -reducibility and localization for quasiperiodic operators. *Math. Res. Lett.*, 23(2):431–444, 2016.
- [24] S. Jitomirskaya, D. A. Koslover, and M. S. Schulteis. Localization for a family of one-dimensional quasiperiodic operators of magnetic origin. *Ann. Henri Poincaré*, 6(1):103–124, 2005.
- [25] S. Jitomirskaya and W. Liu. Universal hierarchical structure of quasiperiodic eigenfunctions. *Ann. of Math. (2)*, 187(3):721–776, 2018.
- [26] S. Jitomirskaya and W. Liu. Universal reflective-hierarchical structure of quasiperiodic eigenfunctions and sharp spectral transition in phase. *arXiv preprint arXiv:1802.00781*, 2018.
- [27] S. Jitomirskaya, W. Liu, and S. Zhang. Arithmetic spectral transitions: a competition between hyperbolicity and the arithmetics of small denominators. *Harmonic Analysis and Applications*, 27:35, 2020.
- [28] S. Jitomirskaya and B. Simon. Operators with singular continuous spectrum. III. Almost periodic Schrödinger operators. *Comm. Math. Phys.*, 165(1):201–205, 1994.
- [29] S. Y. Jitomirskaya. Anderson localization for the almost Mathieu equation: a nonperturbative proof. *Comm. Math. Phys.*, 165(1):49–57, 1994.
- [30] S. Y. Jitomirskaya. Metal-insulator transition for the almost Mathieu operator. *Ann. of Math. (2)*, 150(3):1159–1175, 1999.
- [31] S. Kotani. Ljapunov indices determine absolutely continuous spectra of stationary random one-dimensional Schrödinger operators. In *Stochastic analysis (Katata/Kyoto, 1982)*, volume 32 of *North-Holland Math. Library*, pages 225–247. North-Holland, Amsterdam, 1984.
- [32] Y. Last and B. Simon. Eigenfunctions, transfer matrices, and absolutely continuous spectrum of one-dimensional Schrödinger operators. *Invent. Math.*, 135(2):329–367, 1999.
- [33] W. Liu. Almost Mathieu operators with completely resonant phases. *Ergodic Theory Dynam. Systems*, 40(7):1875–1893, 2020.
- [34] W. Liu. Small denominators and large numerators of quasiperiodic Schrödinger operators. *arXiv preprint arXiv:2205.04648*, 2022.

- [35] W. Liu and X. Yuan. Anderson localization for the almost Mathieu operator in the exponential regime. *J. Spectr. Theory*, 5(1):89–112, 2015.
- [36] W. Liu and X. Yuan. Anderson localization for the completely resonant phases. *J. Funct. Anal.*, 268(3):732–747, 2015.
- [37] W. Liu and X. Yuan. Spectral gaps of almost Mathieu operators in the exponential regime. *J. Fractal Geom.*, 2(1):1–51, 2015.
- [38] C. A. Marx and S. Jitomirskaya. Dynamics and spectral theory of quasi-periodic Schrödinger-type operators. *Ergodic Theory Dynam. Systems*, 37(8):2353–2393, 2017.
- [39] P. Sarnak. Spectral behavior of quasiperiodic potentials. *Comm. Math. Phys.*, 84(3):377–401, 1982.
- [40] W. Schlag. An introduction to multiscale techniques in the theory of Anderson localization, Part I. *Nonlinear Anal.*, 220:Paper No. 112869, 55, 2022.
- [41] J. You. Quantitative almost reducibility and its applications. In *Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. III. Invited lectures*, pages 2113–2135. World Sci. Publ., Hackensack, NJ, 2018.

(W. Liu) DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA

Email address: liuwencai1226@gmail.com; wencail@tamu.edu