

Tau functions and KP solutions on families of algebraic curves

Takashi Ichikawa

Department of Mathematics, Faculty of Science and Engineering, Saga University, Saga 840-8502, Japan. E-mail: ichikawn@cc.saga-u.ac.jp

Abstract: Using abelian differentials and periods of the universal Mumford curve, we study the universal expression and asymptotic behavior of tau functions defined for stably degenerating families of algebraic curves with additional data. Furthermore, we apply this study to constructing solutions to the KP hierarchy which are expressed by nonarchimedean theta functions and by mixtures of quasi-periodic solutions with solitons containing real solutions defined on families of M-curves.

Mathematics Subject Classification 2020. 11G20, 14H10, 14H15, 14D15, 14H40, 14P99, 32G20, 35C80, 37K10.

1. Introduction

Tau functions were introduced by Sato, Jimbo, Miwa and others in the study of integrable systems. Especially, Riemann surfaces with additional data such as marked points and line bundles give rise to tau functions which play important roles in the theory of soliton equations (cf. [23, 34]) and of conformal field theory in operator formalism (cf. [3, 19, 20, 35]). The aim of this paper is to study the tau functions defined for degenerating families of algebraic curves using their explicit formula in terms of theta functions from the viewpoint of formal arithmetic geometry. We have universal expressions of the tau functions and the associated solutions of the KP hierarchy (KP solutions for short) related with soliton solutions which are shown to be meaningful in the fields of complex geometry, real algebraic geometry, nonarchimedean geometry and tropical geometry.

The organization of this paper is as follows.

In Sections 2–5, as a main tool to showing our results, we review results given in [16, 17] on the universal Mumford curve with abelian differentials and periods, and its application to their variational formulas partially obtained in [8, 4, 12].

In Section 6, we give universal and computable formulas of tau functions and the associated KP solutions for families of Schottky uniformized Riemann surfaces. By the arithmeticity of the formulas, extending results in [13] we can express tau functions for all Mumford curves over nonarchimedean complete valuation fields of characteristic 0 in terms of the associated nonarchimedean theta functions (cf. [6, 9]).

In Section 7, we study the asymptotic behavior of tau functions under degeneration of considering Riemann surfaces in order to extend the above conformal field theory for degenerated objects. From the variational formulas stated in Section 5, we can obtain modified tau functions for degenerate objects which give also KP solutions given by

mixtures of quasi-periodic solutions and soliton ones. Similar results were obtained in [29]–[31] and [5] for hyperelliptic curves using the theory of Sato Grassmannian [32] and of sigma functions. Furthermore, generalizing results in [11, 28] we have modified tau functions expressed by rational functions which give rise to soliton solutions as a special case.

In Section 8, we show that the universal Mumford curve give rise to degenerating families of M-curves, i.e., real algebraic curves with maximal number of components. Furthermore, we give a similar statement to results in [1] by showing that the tau functions associated with these M-curves are real, i.e., real-valued for real variables, and tending to soliton KP solutions. This result seems also related to theta functions of tropical curves and their limits which were discussed in [2].

2. Generalized Tate curve

2.1. Schottky uniformization. A Schottky group Γ of rank g is defined as a free group with generators $\gamma_i \in PGL_2(\mathbb{C})$ ($1 \leq i \leq g$) which map Jordan curves $C_{-i} \subset \mathbb{P}_{\mathbb{C}}^1 = \mathbb{C} \cup \{\infty\}$ to other Jordan curves $C_i \subset \mathbb{P}_{\mathbb{C}}^1$ with orientation reversed, where $C_{\pm 1}, \dots, C_{\pm g}$ with their interiors are mutually disjoint. Each element $\gamma \in \Gamma - \{1\}$ is conjugate to an element of $PGL_2(\mathbb{C})$ sending z to $\beta_\gamma z$ for some $\beta_\gamma \in \mathbb{C}^\times$ with $|\beta_\gamma| < 1$ which is called the *multiplier* of γ . Therefore, one has

$$\frac{\gamma(z) - \alpha_\gamma}{z - \alpha_\gamma} = \beta_\gamma \frac{\gamma(z) - \alpha'_\gamma}{z - \alpha'_\gamma}$$

for some element $\alpha_\gamma, \alpha'_\gamma$ of $\mathbb{P}_{\mathbb{C}}^1$ called the *attractive, repulsive* fixed points of γ respectively. Then the discontinuity set $\Omega_\Gamma \subset \mathbb{P}_{\mathbb{C}}^1$ under the action of Γ has a fundamental domain D_Γ which is given by the complement of the union of $C_{\pm 1}, \dots, C_{\pm g}$ with their interiors. The quotient space $R_\Gamma = \Omega_\Gamma / \Gamma$ is a (compact) Riemann surface of genus g which is called *Schottky uniformized* by Γ (cf. [33]). Furthermore, by a result of Koebe, every Riemann surface of genus g can be represented in this manner.

2.2. Generalized Tate curve. A (marked) curve is called *degenerate* if it is a stable (marked) curve and the normalization of its irreducible components are all projective (marked) lines. Then the dual graph $\Delta = (V, E, T)$ of a stable marked curve is a collection of 3 finite sets V of vertices, E of edges, T of tails and 2 boundary maps

$$b : T \rightarrow V, \quad b : E \rightarrow (V \cup \{\text{unordered pairs of elements of } V\})$$

such that the geometric realization of Δ is connected and that Δ is *stable*, namely its each vertex has at least 3 branches. The number of elements of a finite set X is denoted by $\sharp X$, and a (connected) stable graph $\Delta = (V, E, T)$ is called *of (g, n) -type* if $\text{rank}_{\mathbb{Z}} H_1(\Delta, \mathbb{Z}) = g$, $\sharp T = n$. Then under fixing a bijection $\nu : T \xrightarrow{\sim} \{1, \dots, n\}$, which we call a numbering of T , $\Delta = (V, E, T)$ becomes the dual graph of a degenerate curve of genus g with n points such that each tail $h \in T$ corresponds to the $\nu(h)$ th marked point.

In particular, a stable graph without tail is the dual graph of a degenerate (unmarked) curve by this correspondence. If Δ is trivalent, i.e. any vertex of Δ has just 3 branches, then a degenerate $\sharp T$ -marked curve with dual graph Δ is maximally degenerate. An *orientation* of a stable graph $\Delta = (V, E, T)$ means giving an orientation of each $e \in E$. Under an orientation of Δ , denote by $\pm E = \{e, -e \mid e \in E\}$ the set of oriented edges, and by v_h the terminal vertex of $h \in \pm E$ (resp. the boundary vertex of $h \in T$). For each $h \in \pm E$, denote by $|h| \in E$ be the edge h without orientation.

Let $\Delta = (V, E, T)$ be a stable graph. Fix an orientation of Δ , and take a subset \mathcal{E} of $\pm E \cup T$ whose complement \mathcal{E}_∞ satisfies the condition that

$$\pm E \cap \mathcal{E}_\infty \cap \{-h \mid h \in \mathcal{E}_\infty\} = \emptyset,$$

and that $v_h \neq v_{h'}$ for any distinct $h, h' \in \mathcal{E}_\infty$. We attach variables x_h for $h \in \mathcal{E}$ and $y_e = y_{-e}$ for $e \in E$. Let R_Δ be the \mathbb{Z} -algebra generated by x_h ($h \in \mathcal{E}$), $1/(x_e - x_{-e})$ ($e, -e \in \mathcal{E}$) and $1/(x_h - x_{h'})$ ($h, h' \in \mathcal{E}$ with $h \neq h'$ and $v_h = v_{h'}$), and let

$$A_\Delta = R_\Delta[[y_e \ (e \in E)]], \quad B_\Delta = A_\Delta \left[\prod_{e \in E} y_e^{-1} \right].$$

According to [14, Section 2], we construct the universal Schottky group Γ_Δ associated with oriented Δ and \mathcal{E} as follows. For $h \in \pm E$, put

$$\begin{aligned} \phi_h &= \begin{pmatrix} x_h & x_{-h} \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & y_h \end{pmatrix} \begin{pmatrix} x_h & x_{-h} \\ 1 & 1 \end{pmatrix}^{-1} \\ &= \frac{1}{x_h - x_{-h}} \left\{ \begin{pmatrix} x_h & -x_h x_{-h} \\ 1 & -x_{-h} \end{pmatrix} - \begin{pmatrix} x_{-h} & -x_h x_{-h} \\ 1 & -x_h \end{pmatrix} y_h \right\}, \end{aligned}$$

where x_h (resp. x_{-h}) means ∞ if h (resp. $-h$) belongs to \mathcal{E}_∞ . This gives an element of $PGL_2(B_\Delta) = GL_2(B_\Delta)/B_\Delta^\times$ which we denote by the same symbol, and satisfies

$$\frac{\phi_h(z) - x_h}{z - x_h} = y_h \frac{\phi_h(z) - x_{-h}}{z - x_{-h}} \quad (z \in \mathbb{P}^1), \quad (2.2.1)$$

where PGL_2 acts on \mathbb{P}^1 by linear fractional transformation.

For any reduced path $\rho = h(1) \cdot h(2) \cdots h(l)$ which is the product of oriented edges $h(1), \dots, h(l)$ such that $v_{h(i)} = v_{-h(i+1)}$, one can associate an element ρ^* of $PGL_2(B_\Delta)$ having reduced expression $\phi_{h(l)} \phi_{h(l-1)} \cdots \phi_{h(1)}$. Fix a base vertex v_b of V , and consider the fundamental group $\pi_1(\Delta, v_b)$ which is a free group of rank $g = \text{rank}_{\mathbb{Z}} H_1(\Delta, \mathbb{Z})$. Then the correspondence $\rho \mapsto \rho^*$ gives an injective anti-homomorphism $\pi_1(\Delta, v_b) \rightarrow PGL_2(B_\Delta)$ whose image is denoted by Γ_Δ .

It is shown in [14, Section 3] and [15, 1.4] (see also [18, Section 2] when Δ is trivalent and has no loop) that for any stable graph Δ , there exists a stable marked curve \mathcal{C}_Δ of genus g over A_Δ which satisfies the following properties:

- (P1) The closed fiber $\mathcal{C}_\Delta \otimes_{A_\Delta} R_\Delta$ of \mathcal{C}_Δ obtained by substituting $y_e = 0$ ($e \in E$) becomes the degenerate marked curve over R_Δ with dual graph Δ which is obtained from the collection of $P_v := \mathbb{P}_{R_\Delta}^1$ ($v \in V$) by identifying the points $x_e \in P_{v_e}$ and $x_{-e} \in P_{v_{-e}}$ ($e \in E$), where x_h denotes ∞ if $h \in \mathcal{E}_\infty$.
- (P2) \mathcal{C}_Δ gives rise to a universal deformation of degenerate marked curves with dual graph Δ . More precisely, \mathcal{C}_Δ satisfies the following: For a noetherian and normal complete local ring R with residue field k , let C be a marked Mumford curve over R , namely a stable marked curve over R with nonsingular generic fiber such that the closed fiber $C \otimes_R k$ is a degenerate marked curve with dual graph Δ , in which all double points and marked points are k -rational. Then there exists a ring homomorphism $A_\Delta \rightarrow R$ giving $\mathcal{C}_\Delta \otimes_{A_\Delta} R \cong C$.
- (P3) $\mathcal{C}_\Delta \otimes_{A_\Delta} B_\Delta$ is smooth over B_Δ and is Mumford uniformized (cf. [26]) by Γ_Δ .
- (P4) Take x_h ($h \in \mathcal{E}$) as complex numbers such that $x_e \neq x_{-e}$ and that $x_h \neq x_{h'}$ if $h \neq h'$ and $v_h = v_{h'}$, and take y_e ($e \in E$) as sufficiently small nonzero complex numbers. Then \mathcal{C}_Δ gives rise to a family of Riemann surfaces denoted by \mathcal{R}_Δ with marked points which are uniformized by Schottky groups over \mathbb{C} obtained from Γ_Δ .

We review the construction of \mathcal{C}_Δ given in [14, Theorem 3.5]. Let T_Δ be the tree obtained as the universal cover of Δ , and denote by \mathcal{P}_{T_Δ} be the formal scheme as the union of $\mathbb{P}_{A_\Delta}^1$'s indexed by vertices of T_Δ under the B_Δ -isomorphism by ϕ_e ($e \in E$). Then it is shown in [14, Theorem 3.5] that \mathcal{C}_Δ is the formal scheme theoretic quotient of \mathcal{P}_{T_Δ} by Γ_Δ .

3. Universal abelian differential

3.1. *Universal abelian differential.* Let $\Delta = (V, E, T)$ be a stable graph with

$$\text{rank}_{\mathbb{Z}} H_1(\Delta, \mathbb{Z}) = g,$$

and the notation be as in 2.2.

Proposition 3.1. *Let ϕ be a product $\phi_{h(1)} \cdots \phi_{h(l)}$ with $v_{-h(i)} = v_{h(i+1)}$ ($1 \leq i \leq l-1$) which is reduced in the sense that $h(i) \neq -h(i+1)$ ($1 \leq i \leq l-1$), and put $y_\phi = y_{h(1)} \cdots y_{h(l)}$.*

$$(1) \text{ One has } \phi(z) - x_{h(1)} \in y_{h(1)} \left(R_\Delta \left[z, \prod_{h \in \pm E} (z - x_h)^{-1} \right] [[y_e \ (e \in E)]] \right).$$

(2) *If $a \in A_\Delta$ satisfies $a - x_{-h(l)} \in A_\Delta^\times$, then $\phi(a) - x_{h(1)} \in I$. Furthermore, if $a' - x_{-h(l)} \in A_\Delta^\times$, then $\phi(a) - \phi(a') \in (a - a')y_\phi A_\Delta^\times$.*

$$(3) \text{ One has } \frac{d\phi(z)}{dz} \in y_\phi \left(R_\Delta \left[\prod_{h \in \pm E} (z - x_h)^{-1} \right] [[y_e \ (e \in E)]] \right).$$

Proof. Since the assertion (2) is proved in [14, Lemma 1.2], we will prove (1) and (3). Put

$$\phi = \begin{pmatrix} a_\phi & b_\phi \\ c_\phi & d_\phi \end{pmatrix}.$$

Since

$$\begin{pmatrix} \alpha & -\alpha\beta \\ 1 & -\beta \end{pmatrix} \begin{pmatrix} \gamma & -\gamma\delta \\ 1 & -\delta \end{pmatrix} = (\gamma - \beta) \begin{pmatrix} \alpha & -\alpha\beta \\ 1 & -\delta \end{pmatrix},$$

a_ϕ, b_ϕ, c_ϕ and d_ϕ are elements of A_Δ whose constant terms are $x_{h(1)}t, -x_{h(1)}x_{-h(1)}t, t$ and $-x_{-h(1)}t$ respectively, where

$$t = \frac{\prod_{s=2}^l (x_{h(s)} - x_{-h(s-1)})}{\prod_{s=1}^l (x_{h(s)} - x_{-h(s)})} \in A_\Delta^\times.$$

Then $c_\phi z + d_\phi = t(z - x_{-h(1)}) + \cdots$, and hence

$$\phi(z) - x_{h(1)} \in R_\Delta \left[z, \prod_{h \in \pm E} (z - x_h)^{-1} \right] [[y_e \ (e \in E)]].$$

In order to prove (1), we may assume that $l = 1$, and then $\phi(z) = x_{h(1)}$ under $y_{h(1)} = 0$. Therefore, the assertion (1) holds. The assertion (3) follows from

$$\frac{d\phi(z)}{dz} = \frac{\det(\phi)}{(c_\phi z + d_\phi)^2} = \frac{\prod_{s=1}^l \det(\phi_{h(s)})}{(c_\phi z + d_\phi)^2} = \frac{\prod_{s=1}^l y_{h(s)}}{(c_\phi z + d_\phi)^2},$$

and the above calculation. \square

For a stable graph $\Delta = (V, E, T)$, we define universal abelian differentials on a generalized Tate curve \mathcal{C}_Δ . Let $\Gamma_\Delta = \text{Im}(\pi_1(\Delta, v_b) \rightarrow PGL_2(B_\Delta))$ be the universal Schottky group as above. Then it is shown in [14, Lemma 1.3] that each $\gamma \in \Gamma_\Delta - \{1\}$ has its attractive (resp. repulsive) fixed points α (resp. α') in $\mathbb{P}_{B_\Delta}^1$ and its multiplier $\beta \in \sum_{e \in E} A_\Delta \cdot y_e$ which satisfy

$$\frac{\gamma(z) - \alpha}{z - \alpha} = \beta \frac{\gamma(z) - \alpha'}{z - \alpha'}.$$

Fix a set $\{\gamma_1, \dots, \gamma_g\}$ of generators of Γ_Δ , and for each γ_i , denote by α_i (resp. α_{-i}) its attractive (resp. repulsive) fixed points, and by β_i its multiplier. Then under the assumption that there is no element of $\pm E \cap \mathcal{E}_\infty$ with terminal vertex v_b , for each $1 \leq i \leq g$, we define the associated *universal abelian differential of the first kind* as

$$\omega_i = \sum_{\gamma \in \Gamma_\Delta / \langle \gamma_i \rangle} \left(\frac{1}{z - \gamma(\alpha_i)} - \frac{1}{z - \gamma(\alpha_{-i})} \right) dz.$$

Assume that

$$\{h \in \pm E \cap \mathcal{E}_\infty \mid v_h = v_b\} = \emptyset, \quad \{t \in T \mid v_t = v_b\} \neq \emptyset.$$

Then for each $t \in T$ with $v_t = v_b$ and $k > 1$, we define the associated *universal abelian differential of the second kind* as

$$\omega_{t,k} = \begin{cases} \sum_{\gamma \in \Gamma_\Delta} \frac{d\gamma(z)}{(\gamma(z) - x_t)^k} & (x_t \neq \infty), \\ \sum_{\gamma \in \Gamma_\Delta} \gamma(z)^{k-2} d\gamma(z) & (x_t = \infty). \end{cases}$$

Furthermore, put $T_\infty = \{t \in T \cap \mathcal{E}_\infty \mid v_t = v_b\}$ whose cardinality is 0 or 1, and take a maximal subtree \mathcal{T}_Δ of Δ , and for each $t \in T$, take the unique path $\rho_t = h(1) \cdots h(l)$ in \mathcal{T}_Δ from v_t to v_b , and put $\phi_t = \phi_{h(l)} \cdots \phi_{h(1)}$. Then for each $t_1, t_2 \in T$ with $t_1 \neq t_2$, we define the associated *universal abelian differential of the third kind* as

$$\omega_{t_1, t_2} = \sum_{\gamma \in \Gamma_\Delta} \left(\frac{d\gamma(z)}{\gamma(z) - \phi_{t_1}(x_{t_1})} - \frac{d\gamma(z)}{\gamma(z) - \phi_{t_2}(x_{t_2})} \right),$$

where $\phi_{t_i}(x_{t_i}) = \infty$ if $t_i \in T_\infty$.

Theorem 3.2.

- (1) For each $1 \leq i \leq g$, ω_i is a regular differential on $\mathcal{C}_\Delta \otimes_{A_\Delta} B_\Delta$.
- (2) For each $t \in T$ with $v_t = v_b$ and $k > 1$, $\omega_{t,k}$ is a meromorphic differential on $\mathcal{C}_\Delta \otimes_{A_\Delta} B_\Delta$ which has only pole (of order k) at the point p_t corresponding to t .
- (3) For each $t_1, t_2 \in T$ such that $t_1 \neq t_2$, ω_{t_1, t_2} is a meromorphic differential on $\mathcal{C}_\Delta \otimes_{A_\Delta} B_\Delta$ which has only (simple) poles at the points p_{t_1} (resp. p_{t_2}) corresponding to t_1 (resp. t_2) with residue 1 (resp. -1).
- (4) Take x_h ($h \in \pm E \cup T$) and y_e ($e \in E$) be complex numbers as in 2.2 (P4). Then $\omega_i, \omega_{t,k}, \omega_{t_1, t_2}$ are relative abelian differentials on the family \mathcal{R}_Δ of Riemann surfaces.

Proof. By Proposition 3.1, ω_i are differentials on \mathcal{P}_{T_Δ} , and for any $\delta \in \Gamma_\Delta$,

$$\begin{aligned} \omega_i(\delta(z)) &= \sum_{\gamma \in \Gamma_\Delta / \langle \gamma_i \rangle} \left(\frac{1}{\delta(z) - \gamma(\alpha_i)} - \frac{1}{\delta(z) - \gamma(\alpha_{-i})} \right) d\delta(z) \\ &= \sum_{\gamma \in \Gamma_\Delta / \langle \gamma_i \rangle} \left(\frac{\gamma(\alpha_i) - \gamma(\alpha_{-i})}{(\delta(z) - \gamma(\alpha_i))(\delta(z) - \gamma(\alpha_{-i}))} \right) d\delta(z) \\ &= \sum_{\gamma \in \Gamma_\Delta / \langle \gamma_i \rangle} \left(\frac{(\delta^{-1}\gamma)(\alpha_i) - (\delta^{-1}\gamma)(\alpha_{-i})}{(z - (\delta^{-1}\gamma)(\alpha_i))(z - (\delta^{-1}\gamma)(\alpha_{-i}))} \right) dz \end{aligned}$$

$$\begin{aligned}
&= \sum_{\gamma \in \Gamma_{\Delta}/\langle \gamma_i \rangle} \left(\frac{1}{z - (\delta^{-1}\gamma)(\alpha_i)} - \frac{1}{z - (\delta^{-1}\gamma)(\alpha_{-i})} \right) dz \\
&= \omega_i(z).
\end{aligned}$$

Therefore, by the construction of \mathcal{C}_{Δ} reviewed in 2.2, ω_i give rise to differentials on \mathcal{C}_{Δ} which are regular outside $\bigcup_{e \in E} \{y_e = 0\}$, and hence the assertions (1) follows. One can prove the assertions (2), (3) similarly, and hence it is enough to prove (4). As is stated in 2.1, R_{Γ} is given by the quotient space Ω_{Γ}/Γ . Under the assumption on complex numbers x_h and y_e , it is shown in [33] that $\sum_{\gamma \in \Gamma} |\gamma'(z)|$ is uniformly convergent on any compact subset in $\Omega_{\Gamma} - \bigcup_{\gamma \in \Gamma} \gamma(\infty)$, and hence the assertion (4) holds for $\omega_{t,k}$, ω_{t_1,t_2} . If $a \in \Omega_{\Gamma} - \bigcup_{\gamma \in \Gamma} \gamma(\infty)$, then $\lim_{n \rightarrow \infty} \gamma_i^{\pm n}(a) = \alpha_{\pm i}$, and hence

$$\begin{aligned}
d \left(\int_a^{\gamma_i(a)} \sum_{\gamma \in \Gamma} \frac{d\gamma(\zeta)}{\gamma(\zeta) - z} \right) &= \sum_{\gamma \in \Gamma} \left(\frac{1}{z - (\gamma\gamma_i)(a)} - \frac{1}{z - \gamma(a)} \right) dz \\
&= \sum_{\gamma \in \Gamma/\langle \gamma_i \rangle} \lim_{n \rightarrow \infty} \left(\frac{1}{z - (\gamma\gamma_i^n)(a)} - \frac{1}{z - (\gamma\gamma_i^{-n})(a)} \right) dz \\
&= \omega_i(z).
\end{aligned}$$

Therefore, ω_i is absolutely and uniformly convergent on any compact subset in Ω_{Γ} , and hence is an abelian differential on Ω_{Γ}/Γ . \square

3.2. Stability of universal abelian differentials. For a vertex $v \in V$, denote by C_v the corresponding irreducible component of $\mathcal{C}_{\Delta} \otimes_{A_{\Delta}} R_{\Delta}$. Then $P_v = \mathbb{P}_{R_{\Delta}}^1$ is the normalization of C_v .

Theorem 3.3.

(1) For each $1 \leq i \leq g$, let $\phi_{h_i(1)} \cdots \phi_{h_i(l_i)}$ ($h_i(j) \in \pm E$) be the unique reduced product such that $v_{-h_i(j)} = v_{h_i(j+1)}$ and $h_i(1) \neq -h_i(l_i)$ which is conjugate to γ_i . Then for each $v \in V$, the pullback $(\omega_i|_{C_v})^*$ of $\omega_i|_{C_v}$ to P_v is given by

$$\left(\sum_{v_{h_i(j)}=v} \frac{1}{z - x_{h_i(j)}} - \sum_{v_{-h_i(k)}=v} \frac{1}{z - x_{-h_i(k)}} \right) dz.$$

(2) For each $v \in V$, $(\omega_{t,k}|_{C_v})^*$ is given by $\frac{dz}{(z - x_t)^k}$ if $v = v_t$, and is 0 otherwise.

(3) Denote by $\rho_{t_j} = h_j(1) \cdots h_j(l_j)$ the unique path from v_{t_j} ($t_j \in T$) to v_b in \mathcal{T}_{Δ} . Then for each $v \in V$, $(\omega_{t_1,t_2}|_{C_v})^*$ is given by

$$\left(\sum_{v_h=v} \frac{1}{z - x_h} - \sum_{v_{-k}=v} \frac{1}{z - x_{-k}} \right) dz,$$

where h, k runs through $\{t_1, h_1(1), \dots, h_1(l_1), -t_2, -h_2(1), \dots, -h_2(l_2)\}$.

Proof. For the proof of (1), we may assume that $\gamma_i = \phi_{h_i(1)} \cdots \phi_{h_i(l_i)}$. Let γ be an element of Γ_Δ . Then by Proposition 3.1 (2), putting $y_e = 0$ ($e \in E$),

$$\frac{1}{z - \gamma(\alpha_i)} - \frac{1}{z - \gamma(\alpha_{-i})} = \frac{\gamma(\alpha_i) - \gamma(\alpha_{-i})}{(z - \gamma(\alpha_i))(z - \gamma(\alpha_{-i}))}$$

becomes

$$\frac{1}{z - x_{h_i(j)}} - \frac{1}{z - x_{-h_i(j-1)}}$$

if $j \in \{1, \dots, l_i\}$ ($h_i(j-1) := h_i(l_i)$ when $j = 1$), $v(h_i(j)) = v$, $\phi_{h_i(j)} \phi_{h_i(j+1)} \cdots \phi_{h_i(l_i)}$ belongs to $\gamma(\gamma_i)$, and becomes 0 otherwise. Therefore, the assertion follows from the definition of ω_i .

The assertion (2) follows from Proposition 3.1 (1) and (3), and the assertion (3) can be shown in the same way as above. \square

A (regular or meromorphic) global section of the dualizing sheaf on a stable curve is called a *stable differential* (cf. [7]).

Theorem 3.4.

(1) For each $1 \leq i \leq g$, ω_i is a regular stable differential on $\mathcal{C}_\Delta/A_\Delta$, namely an element of the space $H^0(\mathcal{C}_\Delta, \omega_{\mathcal{C}_\Delta/A_\Delta})$ of global sections of the dualizing sheaf $\omega_{\mathcal{C}_\Delta/A_\Delta}$ on $\mathcal{C}_\Delta/A_\Delta$. Furthermore, $\{\omega_i\}_{1 \leq i \leq g}$ gives a basis of $H^0(\mathcal{C}_\Delta, \omega_{\mathcal{C}_\Delta/A_\Delta})$.

(2) For each $t \in T$ with $v_t = v_b$ and $k > 1$, $\omega_{t,k}$ is a meromorphic stable differential on $\mathcal{C}_\Delta/A_\Delta$ which has only pole (of order k) at the point p_t corresponding to t .

(3) For each $t_1, t_2 \in T$ with $t_1 \neq t_2$, ω_{t_1, t_2} is a meromorphic stable differential on $\mathcal{C}_\Delta/A_\Delta$ which has only (simple) poles at the points p_{t_1} (resp. p_{t_2}) corresponding to t_1 (resp. t_2) with residue 1 (resp. -1).

Proof. We only show that the latter assertion in (1) since the remains follow from Theorems 3.2 and 3.3. For each i , take the product $\phi_{h_i(1)} \cdots \phi_{h_i(l_i)}$ of ϕ_h ($h \in \pm E$) as in Theorem 3.3 (1). Then by this theorem, $\omega_i|_{\mathcal{C}_{v_{h_i(1)}}}$ has simple pole at $x_{h_i(1)}$ with residue 1, and hence $\{\omega_i\}_{1 \leq i \leq g}$ gives a basis of $H^0(\mathcal{C}_\Delta, \omega_{\mathcal{C}_\Delta/A_\Delta})$ since $\{\gamma_i\}$ is a basis of $H_1(\Delta, \mathbb{Z}) \cong \Gamma_\Delta/[\Gamma_\Delta, \Gamma_\Delta]$. \square

Remark 3.5. For a subset E' of the set E of edges in the stable graph Δ , let $\Delta' = (V', E', T')$ be the associated graph obtained from Δ , and denote by $\mathcal{C}_{\Delta'}$ the stable curve given by putting $y_e = 0$ ($e \in E'$). Let \mathcal{R}_Δ be a family of Schottky uniformized Riemann surfaces as in 2.2 (P4), and let c_e be a cycle corresponding to $e \in E - E'$ which is oriented by the right-hand rule for γ_i . Then by Theorem 3.3 (1), the restriction of ω_i to each irreducible component of $\mathcal{R}_{\Delta'}$ is characterized analytically as follows:

- its integrals along c_e ($e \in E - E'$) are $2\pi\sqrt{-1}$ if $e = |h_i(j)|$ for some integer $1 \leq j \leq l_i$, and are 0 otherwise, and its residues at $x_{h_i(j)}$ (resp. $x_{h_i(k)}$) are 1 (resp. -1) for some integers $1 \leq j, k \leq l_i$, and are 0 otherwise.

By results in Section 5 below, this characterization holds for any deformation of a stable complex curve whose irreducible components are not necessarily projective lines. This modification gives a more explicit formula than [12, Corollary 4.6], and one has similar statements on $\omega_{t,k}$ and ω_{t_1,t_2} .

3.3. Period.

Theorem 3.6.

(1) *The Jacobian of \mathcal{C}_Δ becomes the Mumford abelian scheme [27] whose multiplicative periods is given in [14, 3.10] as $P_{ij} = \prod_\gamma \psi_{ij}(\gamma)$ ($1 \leq i, j \leq g$), where γ runs through all representatives of $\langle \gamma_i \rangle \backslash \Gamma_\Delta / \langle \gamma_j \rangle$ and*

$$\psi_{ij}(\gamma) = \begin{cases} \beta_i & (i = j \text{ and } \gamma \in \langle \gamma_i \rangle), \\ \frac{(\alpha_i - \gamma(\alpha_j))(\alpha_{-i} - \gamma(\alpha_{-j}))}{(\alpha_{-i} - \gamma(\alpha_j))(\alpha_i - \gamma(\alpha_{-j}))} & (\text{otherwise}). \end{cases}$$

(2) *Let R be a Riemann surface obtained from \mathcal{C}_Δ as in 2.2 (P4), and take counterclockwise oriented small closed paths a_i ($1 \leq i \leq g$) in R surrounding α_i , and closed paths b_i ($1 \leq i \leq g$) in R corresponding to γ_i . Denote by the same symbols the abelian differentials of the first kind and the multiplicative periods for R given by the above ω_j ($1 \leq j \leq g$) and P_{ij} ($1 \leq i, j \leq g$) respectively. Then*

$$\int_{a_i} \omega_j = 2\pi\sqrt{-1}\delta_{ij}, \quad \exp\left(\int_{b_i} \omega_j\right) = P_{ij},$$

where δ_{ij} denotes the Kronecker delta.

Proof. Since the assertion (1) is shown in [14, Theorem 3.13] and the first formula of the assertion (2) follows from Remark 3.5, we will prove the second formula of (2) (cf. [33] and [25, Theorem 2]). Let Γ be a Schottky group over \mathbb{C} obtained from Γ_Δ as in 2.2 (P4). Then as is seen in 2.1, $R = R_\Gamma$ is obtained as the quotient space of the discontinuity set Ω_Γ under the action of Γ . Take a point a on Ω_Γ . Then under the condition that $i = j$ and $\gamma \in \langle \gamma_i \rangle$,

$$\frac{(\gamma_i(a) - \gamma(\alpha_i))(a - \gamma(\alpha_{-i}))}{(a - \gamma(\alpha_i))(\gamma_i(a) - \gamma(\alpha_{-i}))} = \beta_i = \psi_{ii}(\gamma).$$

Since $\lim_{n \rightarrow \infty} \gamma_i^{\pm n}(a) = \alpha_{\pm i}$, if $i \neq j$ or $\gamma \notin \langle \gamma_i \rangle$, then

$$\begin{aligned} & \prod_{n \in \mathbb{Z}} \left(\frac{(\gamma_i(a) - \gamma_i^{-n}\gamma(\alpha_j))(a - \gamma_i^{-n}\gamma(\alpha_{-j}))}{(a - \gamma_i^{-n}\gamma(\alpha_j))(\gamma_i(a) - \gamma_i^{-n}\gamma(\alpha_{-j}))} \right) \\ &= \prod_{n \in \mathbb{Z}} \left(\frac{(\gamma_i^{n+1}(a) - \gamma(\alpha_j))(\gamma_i^n(a) - \gamma(\alpha_{-j}))}{(\gamma_i^n(a) - \gamma(\alpha_j))(\gamma_i^{n+1}(a) - \gamma(\alpha_{-j}))} \right) \\ &= \frac{(\alpha_i - \gamma(\alpha_j))(\alpha_{-i} - \gamma(\alpha_{-j}))}{(\alpha_{-i} - \gamma(\alpha_j))(\alpha_i - \gamma(\alpha_{-j}))} \\ &= \psi_{ij}(\gamma). \end{aligned}$$

Therefore,

$$\exp\left(\int_{b_i} \omega_j\right) = \prod_{\gamma \in \langle \gamma_i \rangle \backslash \Gamma / \langle \gamma_j \rangle} \psi_{ij}(\gamma) = P_{ij}$$

which completes the proof. \square

4. Universal Mumford curve, differentials and periods

4.1. Universal Mumford curve. Let $\Delta = (V, E, T)$ be a stable graph which is not trivalent. Then there exists a vertex $v_0 \in V$ which has at least 4 branches. Take two elements h_1, h_2 of \mathcal{E} such that $h_1 \neq h_2$ and $v_{h_1} = v_{h_2} = v_0$, and let $\Delta' = (V', E', T')$ be a stable graph obtained from Δ by replacing v_0 with an oriented (nonloop) edge h_0 such that $v_{h_1} = v_{h_2} = v_{h_0}$ and that $v_h = v_{-h_0}$ for any $h \in \pm E \cup T - \{h_1, h_2\}$ with $v_h = v_0$. Put $e_i = |h_i|$ for $i = 0, 1, 2$. Then we have the following identifications:

$$V = V' - \{v_{-h_0}\} \text{ (in which } v_0 = v_{h_0}), \quad E = E' - \{e_0\}, \quad T = T'.$$

Theorem 4.1

(1) *The generalized Tate curves \mathcal{C}_Δ and $\mathcal{C}_{\Delta'}$ associated with Δ and Δ' respectively are isomorphic over $R_{\Delta'}[[s_e (e \in E' - \{e_0\})]][s_{e_0}^{-1}]$, where*

$$\frac{x_{h_1} - x_{h_2}}{s_{e_0}}, \quad \frac{y_{e_i}}{s_{e_0} s_{e_i}} \text{ (} i = 1, 2 \text{ with } h_i \notin T), \quad \frac{y_e}{s_e} \text{ (} e \in E - \{e_1, e_2\})$$

belong to $(A_{\Delta'})^\times$ if $h_1 \neq -h_2$, and

$$\frac{x_{h_1} - x_{h_2}}{s_{e_0}}, \quad \frac{y_e}{s_e} \text{ (} e \in E)$$

belong to $(A_{\Delta'})^\times$ if $h_1 = -h_2$.

(2) *The assertion (1) holds in the category of complex geometry when $x_{h_1} - x_{h_2}, y_e$ and s_e are taken to be sufficiently small complex numbers with $x_{h_1} \neq x_{h_2}, s_{e_0} \neq 0$.*

Proof. First, we prove the assertion (1). Denote by t_h the moduli parameters of $\mathcal{C}_{\Delta'}$ corresponding to $h \in \pm E' \cup T'$. Then by [13, Lemma 1.2], $\phi_{-h_0}(t_{h_1}) - \phi_{-h_0}(t_{h_2})$ belongs to $s_{e_0} \cdot (A_{\Delta'})^\times$, and hence

$$\mathcal{C}_{\Delta'} \otimes_{A_{\Delta'}} (R_{\Delta'}[[s_e]][s_{e_0}^{-1}])$$

gives a universal deformation of a universal degenerate curve with dual graph Δ . Then by the universality of generalized Tate curves, there exists an injective homomorphism $A_\Delta \hookrightarrow R_{\Delta'}[[s_e]][s_{e_0}^{-1}]$ which gives rise to an isomorphism $\mathcal{C}_\Delta \cong \mathcal{C}_{\Delta'}$. Under this homomorphism,

$$\begin{aligned} & (P_{v_{-h_0}}; \phi_{-h_0}(t_{h_1}), \phi_{-h_0}(t_{h_2}), t_h \text{ (} v_h = v_{-h_0}, h \neq h_0)) \\ & \cong (P_{v_0}; x_{h_1}, x_{h_2}, x_h \text{ (} v_h = v_0, h \neq h_1, h_2)), \end{aligned}$$

and hence $x_{h_1} - x_{h_2} \in s_{e_0} \cdot (A_{\Delta'})^\times$. Furthermore, when $h_1 \neq -h_2$, the deformation parameters of

$$\left(P_{v_{-h_0}}; \phi_{-h_0}(t_{h_1}), \phi_{-h_0}(t_{h_2}), t_h \ (v_h = v_{-h_0}, h \neq h_0) \right)$$

corresponding to $h_i \cdot (-h_0)$ ($i = 1, 2$) are y_{h_i} , and hence by [13, Proposition 1.3], $y_{h_i} \in (s_{h_0} \cdot s_{h_i}) \cdot (A_{\Delta'})^\times$. When $h_1 = -h_2$, the deformation parameters of

$$\left(P_{v_{-h_0}}; \phi_{-h_0}(t_{h_1}), \phi_{-h_0}(t_{h_2}), t_h \ (v_h = v_{-h_0}, h \neq h_0) \right)$$

corresponding to $h_0 \cdot h_1 \cdot (-h_0)$ is y_{h_1} , and hence $y_{h_1} \in s_{h_1} \cdot (A_{\Delta'})^\times$.

The assertion (2) follows from 2.2 (P4). \square

Remark 4.2 Let Γ_Δ and $\Gamma_{\Delta'}$ denote the universal Schottky groups associated with Δ and Δ' respectively. Since $\mathcal{C}_\Delta \cong \mathcal{C}_{\Delta'}$, by [26, Corollary 4.11], there exists a conjugation isomorphism $\varphi : \Gamma_\Delta \xrightarrow{\sim} \Gamma_{\Delta'}$, and hence the multiplier of an element of Γ_Δ and the cross-ratio of the fixed points of four elements of Γ_Δ are invariant under φ . Therefore, one can calculate the precise formula connecting the moduli and deformation parameters for \mathcal{C}_Δ and $\mathcal{C}_{\Delta'}$. For the detail, see the proof of [14, Theorem 1].

For nonnegative integers g, n such that $2g - 2 + n > 0$, denote by $\overline{\mathcal{M}}_{g,n}$ the moduli stack over \mathbb{Z} of stable n -marked curves of genus g (cf. [7, 21, 22]). Then by definition, there exist the universal stable marked curve $\mathcal{C}_{g,n}$ over $\overline{\mathcal{M}}_{g,n}$, and the associated curve \mathcal{C}_g obtained by forgetting marked points on $\mathcal{C}_{g,n}$.

Theorem 4.3. *There exists a deformation space $\mathcal{A}_{g,n}$ of all n -marked degenerate curves of genus g , and an n -marked stable curve of genus g over $\mathcal{A}_{g,n}$ whose fiber by the canonical morphism $\text{Spec}(A_\Delta) \rightarrow \mathcal{A}_{g,n}$ becomes the generalized Tate curve \mathcal{C}_Δ for each stable graph Δ of (g, n) -type.*

Proof. Let $\Delta = (V, E, T)$ be a stable graph of (g, n) -type, and take a system of coordinates on $P_v = \mathbb{P}_{R_\Delta}^1$ ($v \in V$) such that $x_h = \infty$ ($h \in \mathcal{E}_\infty$) and that $\{0, 1\} \subset P_v$ is contained in the set of points given by x_h ($h \in \mathcal{E}$ with $v_h = v$). Under this system of coordinates, one has the generalized Tate curve \mathcal{C}_Δ whose closed fiber $\mathcal{C}_\Delta \otimes_{A_\Delta} R_\Delta$ gives a family of degenerate curves over the open subspace of

$$S_\Delta = \left\{ (p_h \in P_{v_h})_{h \in \pm E \cup T} \mid p_h \neq p_{h'} \ (h \neq h', v_h = v_{h'}) \right\}$$

defined as $p_e \neq p_{-e}$ for nonloop edges $e \in E$. Therefore, taking another system of coordinates on P_v obtained by mutual changes of $0, 1, \infty$ and comparing the associated generalized Tate curves with the original \mathcal{C}_Δ as in Theorem 4.1, \mathcal{C}_Δ can be extended over the deformation space of all marked degenerate curves with dual graph Δ . Since two stable graphs of (g, n) -type can be translated by a combination of replacements $\Delta \leftrightarrow \Delta'$ given in 3.1, one can define a scheme $\mathcal{A}_{g,n}$ obtained by gluing $\text{Spec}(A_\Delta)$ (Δ :

stable graphs of (g, n) -type) along the isomorphism given in Theorem 4.1. Then $\mathcal{A}_{g,n}$ is regarded as the deformation space of all n -marked degenerate curves of genus g over which there exists an n -marked stable curve of genus g obtained by gluing \mathcal{C}_Δ . \square

Definition 4.4. We call the above n -marked stable curve of genus g over $\mathcal{A}_{g,n}$ the n -marked *universal Mumford curve* of genus g which is the fiber of $\mathcal{C}_{g,n}$ by the canonical morphism $\mathcal{A}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$. By 2.2 (P2) and (P4), one can see that this universal Mumford curve gives rise to all n -marked Mumford curves of genus g , and to all n -marked Riemann surfaces of genus g close to degenerate curves.

Remark 4.5. Gerritzen-Herrlich [10] introduced the extended Schottky space $\overline{\mathcal{S}}_g$ of genus $g > 1$ as the fine moduli space of stable complex curves of genus g with Schottky structure. For integers g, n as above, one can consider the extended Schottky space $\overline{\mathcal{S}}_{g,n}$ for stable n -marked complex curves of genus g with Schottky structure. Then by the result of Koebe referred in 2.1, $\overline{\mathcal{S}}_{g,n}/\text{Out}(F_g)$ becomes a covering space of the moduli space of n -marked stable complex curves of genus g which is also the complex analytic space $\overline{\mathcal{M}}_{g,n}^{\text{an}}$ associated with $\overline{\mathcal{M}}_{g,n}$. Furthermore, the n -marked universal Mumford curve of genus g can be analytically extended to the universal family of marked stable complex curves over $\overline{\mathcal{S}}_{g,n}/\text{Out}(F_g)$.

4.2. Differentials and periods of the universal Mumford curve. Denote by $\Delta_0 = (V_0, E_0, T_0)$ the stable graph of (g, n) -type consisting of one vertex and g loops, and fix generators ρ_1, \dots, ρ_g of $\pi_1(\Delta_0)$. For each stable graph $\Delta = (V, E, T)$ of (g, n) -type, Δ_0 is obtained from Δ by contracting some nonloop edges in E . Then there exists a unique \mathbb{Z} -basis $\{b_i\}_{1 \leq i \leq g}$ of $H_1(\Delta, \mathbb{Z})$ corresponding to

$$\{[\rho_i]\}_{1 \leq i \leq g} \in H_1(\Delta_0, \mathbb{Z}),$$

and T is identified with T_0 . Therefore, by Theorem 3.4, there exist associated stable differentials

$$\omega_i \ (i = 1, \dots, g), \quad \omega_{t,k} \ (t \in T_0, k > 1), \quad \omega_{t_1, t_2} \ (t_i \in T_0, t_1 \neq t_2)$$

on \mathcal{C}_Δ of the first, second, third kind respectively. Furthermore, b_i ($i = 1, \dots, g$) give rise to homology cycles on members of \mathcal{R}_Δ given in 2.2 (P4), and on the universal family of stable complex curves over the extended Schottky space $\overline{\mathcal{S}}_{g,n}$ by the analytic extension. We also denote by b_i these homology cycles.

Theorem 4.6.

(1) *The differentials $\omega_i, \omega_{t,k}, \omega_{t_1, t_2}$ on \mathcal{C}_Δ are glued to stable differentials on $\mathcal{C}_g/\mathcal{A}_{g,n}$ which we call the universal differentials of the first kind, second kind, third kind, and denote by $\overline{\omega}_i, \overline{\omega}_{t,k}, \overline{\omega}_{t_1, t_2}$ respectively.*

(2) *The universal differentials $\overline{\omega}_i$ ($i = 1, \dots, g$) on $\mathcal{C}_g/\mathcal{A}_{g,n}$ of the first kind make a basis of the sheaf $\mathcal{H}^0(\omega_{\mathcal{C}_g/\mathcal{A}_{g,n}})$ which consists of sections of the relative stable differentials on $\mathcal{C}_g/\mathcal{A}_{g,n}$. Furthermore, $\overline{\omega}_i$ are analytically extended to stable differentials on*

the universal family of stable complex curves over $\overline{S}_{g,n}$ which we denote by the same symbols.

Proof. First, we prove (1). As is stated in Remark 4.2, the isomorphism $\mathcal{C}_\Delta \cong \mathcal{C}_{\Delta_0}$ over B_Δ considered in Theorem 3.1 corresponds uniquely to an isomorphism $\Gamma_\Delta \cong \Gamma_{\Delta_0}$. Therefore, under this isomorphism $\mathcal{C}_\Delta \cong \mathcal{C}_{\Delta_0}$, the differentials $\omega_i, \omega_{t,k}, \omega_{t_1,t_2}$ on \mathcal{C}_Δ are mapped to those on \mathcal{C}_{Δ_0} , and hence can be glued to differentials on $\mathcal{C}_g/\mathcal{A}_{g,n}$.

Second, we prove the latter assertion of (2) since the former one follows from Theorem 4.6 (1). For a stable n -marked complex curve C of genus g with Schottky structure, as is stated in [10], one can take a cut system $\{a_1, \dots, a_g\}$ consisting of disjoint oriented simple loops in C such that the intersection numbers $a_i \cdot b_j$ are the Kronecker delta δ_{ij} . Then there exist uniquely regular stable differentials $\omega_{C,i}$ ($i = 1, \dots, g$) on C such that

$$\int_{a_j} \omega_{C,i} = 2\pi\sqrt{-1}\delta_{ij}.$$

Therefore, moving C on $\overline{S}_{g,n}$, $\omega_{C,i}$ form the analytic extension of $\overline{\omega}_i$. \square

Denote by $\mathcal{B}_{g,n}$ the maximal open subscheme of $\mathcal{A}_{g,n}$ over which the associated marked curves are smooth, and by $S_{g,n}$ the (ordinary) Schottky space which is defined as the open subspace of $\overline{S}_{g,n}$ classifying n -marked Riemann surfaces of genus g with Schottky structure.

Theorem 4.7. *There exist $\mathcal{P}_{ij} \in \mathcal{O}_{\mathcal{B}_{g,n}}^\times$ ($1 \leq i, j \leq g$) which give the multiplicative periods of Mumford curves over p -adic fields obtained from $\mathcal{C}_g/\mathcal{B}_{g,n}$ as in 2.2 (P2). Furthermore, \mathcal{P}_{ij} can be analytically continued to regular functions on $S_{g,n}$ which become $\exp\left(\int_{b_i} \overline{\omega}_j\right)$, where $\overline{\omega}_j$ are given in Theorem 4.6. We call \mathcal{P}_{ij} the universal periods.*

Proof. For a stable graph Δ of (g, n) -type, take generators $\gamma_1, \dots, \gamma_g$ of Γ_Δ such that $[\gamma_i] = b_i$ ($i = 1, \dots, g$), and for each γ_i , denote by α_i (resp. α_{-i}) its attractive (resp. repulsive) fixed points and by β_i its multiplier. Then by [13, Theorem 3.13], the multiplicative periods P_{ij} ($1 \leq i, j \leq g$) of \mathcal{C}_Δ are elements of B_Δ^\times defined as $P_{ij} = \prod_\gamma \psi_{ij}(\gamma)$, where γ runs through all representatives of $\langle \gamma_i \rangle \backslash \Gamma_\Delta / \langle \gamma_j \rangle$ and

$$\psi_{ij}(\gamma) = \begin{cases} \beta_i & (i = j, \gamma \in \langle \gamma_i \rangle), \\ \frac{(\alpha_i - \gamma(\alpha_j))(\alpha_{-i} - \gamma(\alpha_{-j}))}{(\alpha_{-i} - \gamma(\alpha_j))(\alpha_i - \gamma(\alpha_{-j}))} & (\text{otherwise}). \end{cases}$$

This implies that P_{ij} depend only on $\{b_i\}_{1 \leq i \leq g}$, and they give rise to the multiplicative periods $\langle [\gamma_i], [\gamma_j] \rangle$ of Mumford curves over p -adic fields by [25, Theorem 2], and to $\exp\left(\int_{b_i} \omega_j\right)$ by the definition of ω_j (see also [24, Theorem 7.5]). Therefore, as in

the proof of Theorem 5.1, P_{ij} are seen to be glued to regular functions \mathcal{P}_{ij} on $\mathcal{B}_{g,n}$ satisfying the required properties. \square

5. Degeneration of abelian differentials

5.1. Irreducible degeneration. We apply results in Section 3 to the study of abelian differentials on a general family of Riemann surfaces which stably degenerates to an irreducible singular complex curve.

Let \mathcal{R} be a family of Riemann surfaces of genus $g > 0$ with symplectic bases $\{a_i, b_i\}_{1 \leq i \leq g}$ (of the first Betti homology groups of members in \mathcal{R}) and local coordinates u at marked points p which is defined over a small disk around $y = 0$ in \mathbb{C} . Here u are centered at p that is $u(p) = 0$, and $\{a_i, b_i\}$ are taken as homotopy classes of closed paths in members in \mathcal{R} such that $(a_i, b_j) = \delta_{ij}$, $(a_i, a_j) = (b_i, b_j) = 0$. Assume that letting $y \rightarrow 0$ corresponds to pinching a_g to a point, and then \mathcal{R} degenerates to a singular complex curve obtained from a Riemann surface R' of genus $g - 1$ by identifying two points p_1, p_2 on R' , where b_g becomes (up to homotopy) to a path from p_1 to p_2 in $R'/(p_1 = p_2)$. Denote by $(\{a'_i, b'_i\}_{1 \leq i \leq g-1}, p, u)$ a symplectic basis and a local coordinate at a marked point of R' obtained from $(\{a_i, b_i\}_{1 \leq i \leq g-1}, p, u)$ by putting $y = 0$.

Let $\{\omega_1, \dots, \omega_g\}$ be a unique basis of the space of relative abelian differentials of the first kind on \mathcal{R} which is normalized for $\{a_i\}_{1 \leq i \leq g}$ in the sense that

$$\int_{a_i} \omega_j = 2\pi\sqrt{-1}\delta_{ij} \quad (1 \leq i, j \leq g).$$

By the theorem of Riemann-Roch, for each positive integer n , there exists a unique relative abelian differential $\omega^{(n)}$ of the second kind with order $n + 1$ on \mathcal{R} which is normalized for $(\{a_i\}_{1 \leq i \leq g}, p, u)$ in the sense that

- $\omega^{(n)}$ is holomorphic outside p ,
- $\omega^{(n)} - \frac{1}{u^{n+1}}du$ is holomorphic at p ,
- $\int_{a_i} \omega^{(n)} = 0$ for any $i = 1, \dots, g$.

Theorem 5.1.

(1) Under $y \rightarrow 0$, $\{\omega_1, \dots, \omega_{g-1}\}$ becomes to the basis of the space of abelian differentials of the first kind on R' which is normalized for $\{a'_i\}_{1 \leq i \leq g-1}$, and ω_g becomes to the unique abelian differential ω' of the third kind on R' which is normalized for $(\{a'_i\}_{1 \leq i \leq g-1}, p_1, p_2)$ in the sense that ω' has only simple poles at p_1, p_2 with residue $1, -1$ respectively and satisfies

$$\int_{a'_i} \omega' = 0 \quad (1 \leq i \leq g - 1).$$

(2) Under $y \rightarrow 0$, $\omega^{(n)}$ becomes to the unique abelian differential of the second kind with order $n + 1$ on R' normalized for $(\{a'_i\}_{1 \leq i \leq g-1}, p, u)$.

Proof. First, we prove the assertions (1) and (2) in the case when \mathcal{R} is obtained from a generalized Tate curve as in 2.2 (P4). Let $\Delta = (V, E, T)$ be a stable graph such that $\text{rank}_{\mathbb{Z}} H_1(\Delta, \mathbb{Z}) = g$, $\#V = \#T = 1$ and that there is an oriented loop in $\pm E$ which we denote by l , where Δ is obtained from a stable graph Δ' with three tails t, t_1, t_2 by connecting t_1, t_2 with l . There exist generators $\gamma_1, \dots, \gamma_g$ of Γ_{Δ} such that $\gamma_1, \dots, \gamma_{g-1}$ are generators of $\Gamma_{\Delta'}$ and that γ_g corresponds to l . We take a symplectic basis $\{a_i, b_i\}_{1 \leq i \leq g}$ of a family \mathcal{R}_{Δ} of Riemann surfaces obtained from \mathcal{C}_{Δ} such that b_i corresponds to γ_i as in 2.2 (P4). Then by Theorem 3.4, the universal abelian differentials of the first kind

$$\omega_i = \sum_{\gamma \in \Gamma_{\Delta} / \langle \gamma_i \rangle} \left(\frac{1}{z - \gamma(\alpha_i)} - \frac{1}{z - \gamma(\alpha_{-i})} \right) dz \quad (1 \leq i \leq g),$$

on \mathcal{C}_{Δ} give the basis of the space of relative abelian differentials of the first kind on \mathcal{R}_{Δ} normalized for $\{a_i\}_{1 \leq i \leq g}$. By Proposition 3.1 (2),

$$\left(\frac{1}{z - \gamma(\alpha_i)} - \frac{1}{z - \gamma(\alpha_{-i})} \right) \Big|_{y_g=0} = \frac{\gamma(\alpha_i) - \gamma(\alpha_{-i})}{(z - \gamma(\alpha_i))(z - \gamma(\alpha_{-i}))} \Big|_{y_g=0} = 0$$

if $i \neq g$, $\gamma \in \Gamma_{\Delta} - \Gamma_{\Delta'}$ or $i = g$, γ has a reduced product $\gamma' \gamma_g^n$, where $\gamma' \in \Gamma_{\Delta} - \Gamma_{\Delta'}$ and $n \in \mathbb{Z}$. Therefore,

$$\omega_i|_{y_g=0} = \sum_{\gamma \in \Gamma_{\Delta'} / \langle \gamma_i \rangle} \left(\frac{1}{z - \gamma(\alpha_i)} - \frac{1}{z - \gamma(\alpha_{-i})} \right) dz \quad (1 \leq i \leq g-1),$$

and hence $\omega_i|_{y_g=0}$ ($1 \leq i \leq g-1$) are the universal abelian differentials of the first kind on $\mathcal{C}_{\Delta'}$ which give the basis of the space of relative abelian differentials of the first kind on $\mathcal{R}_{\Delta'}$ normalized for $\{a_i\}_{1 \leq i \leq g-1}$. Furthermore,

$$\begin{aligned} \omega_g|_{y_g=0} &= \sum_{\gamma \in \Gamma_{\Delta'}} \left(\frac{1}{z - \gamma(\alpha_g)} - \frac{1}{z - \gamma(\alpha_{-g})} \right) dz \\ &= \sum_{\gamma \in \Gamma_{\Delta'}} \left(\frac{d\gamma(z)}{\gamma(z) - \alpha_g} - \frac{d\gamma(z)}{\gamma(z) - \alpha_{-g}} \right), \end{aligned}$$

and hence $\omega_g|_{y_g=0}$ is the universal abelian differential of the third kind on $\mathcal{C}_{\Delta'}$ which gives the unique relative abelian differential of the third kind on $\mathcal{R}_{\Delta'}$ normalized for $(\{a_i\}, \alpha_g, \alpha_{-g})$. Therefore, the assertion (1) holds for $(\mathcal{R}_{\Delta}, \{a_i, b_i\})$.

For an integer $k > 1$ and the unique tail t of Δ , by Theorem 3.4, the universal abelian differential of the second kind

$$\omega_{t,k} = \sum_{\gamma \in \Gamma_{\Delta}} \frac{d\gamma(z)}{(\gamma(z) - x_t)^k}$$

on \mathcal{C}_Δ gives the unique relative abelian differential of the second kind with order k on \mathcal{R}_Δ normalized for $(\{a_i\}_{1 \leq i \leq g}, p_t, z - p_t)$. By Proposition 3.1 (3), $d\gamma(z)|_{y_g=0} = 0$ if $\gamma \in \Gamma_\Delta - \Gamma_{\Delta'}$. Therefore,

$$\omega_{t,k}|_{y_g=0} = \sum_{\gamma \in \Gamma_{\Delta'}} \frac{d\gamma(z)}{(\gamma(z) - x_t)^k},$$

and hence $\omega_{t,k}|_{y_g=0}$ is the universal abelian differential of the second kind on $\mathcal{C}_{\Delta'}$ which gives the unique relative abelian differential of the second kind with order k on $\mathcal{R}_{\Delta'}$ normalized for $(\{a_i\}_{1 \leq i \leq g-1}, p_t, z - x_t)$. Therefore, the assertion (2) holds for $(\mathcal{R}_\Delta, \{a_i, b_i\}, p_t)$, where $z - p_t$ can be replaced with general local coordinates at p_t .

To prove the assertions generally, for integers $g, n \geq 0$ such that $2g - g + n > 0$, we consider the moduli space $\mathcal{H}_{g,n}$ of $(R, \{a_i, b_i\}_{1 \leq i \leq g}, p_1, \dots, p_n)$, where R are Riemann surfaces, $\{a_i, b_i\}$ are symplectic bases of R and p_1, \dots, p_n are points on R which are different each other. Then $\mathcal{H}_{g,n}$ is a complex manifold as a covering space of the moduli space of Riemann surfaces of genus g with n marked points (cf. [22, 21]), and is seen to be connected by the Teichmüller theory. Furthermore, the above $(\mathcal{R}_{\Delta'}, \{a_i, b_i\}_{1 \leq i \leq g-1}, p_{t_1}, p_{t_2}, z)$ make an open subset of $\mathcal{H}_{g-1,3}$, and $(\mathcal{R}_\Delta, \{a_i, b_i\}, z)$ give deformations of $(\mathcal{R}_{\Delta'}/(p_{t_1} = p_{t_2}), \{a_i, b_i\}, z)$ for which the assertion (1) holds. Take a member $(R', \{a'_i, b'_i\}, p_1, p_2, z)$ of $\mathcal{H}_{g-1,3}$ with deformation $(\mathcal{R}, \{a_i, b_i\}, z)$ of $(R'/(p_1 = p_2), \{a'_i, b'_i\}, z)$ by a complex parameter y as above. Let $\{\omega_1, \dots, \omega_g\}$ be the basis of the space of relative abelian differentials of the first kind on \mathcal{R} normalized for $\{a_i\}$, $\{\omega'_1, \dots, \omega'_{g-1}\}$ be the basis of the space of abelian differentials of the first kind on R' normalized for $\{a'_i\}$, and ω'_g be the abelian differential of the third kind on R' normalized for $(\{a'_i\}, p_1, p_2)$. Then for each $1 \leq i \leq g$, values at $z \in R'$ of

$$\lim_{y \rightarrow 0} \omega_i - \omega'_i$$

are holomorphic on the moduli space $\mathcal{H}_{g-1,3}$ of $(R', \{a'_i, b'_i\}, p_1, p_2, z)$ and are 0 on its nonempty open subset $\{(\mathcal{R}_{\Delta'}, \{a_i, b_i\}, p_{t_1}, p_{t_2}, z)\}$. Therefore, by the identity theorem, these values are 0 on the whole $\mathcal{H}_{g-1,3}$, and hence the assertion (1) holds generally.

In order to show the assertion (2), for each positive integer k , we consider the moduli space $\mathcal{H}_{g-1,2,k}$ of $(R', \{a'_i, b'_i\}, p, u, z)$, where $(R', \{a'_i, b'_i\}, p, z)$ are members of $\mathcal{H}_{g-1,2}$ and u are k -jets of local coordinates at p . Then $\mathcal{H}_{g-1,2,k}$ is a connected complex manifold which contains $\{(\mathcal{R}_{\Delta'}, \{a_i, b_i\}, p_t, u, z)\}$ as its open subset, where u are k -jets of local coordinates at p_t . Therefore, a similar argument as above shows the assertion (2). \square

5.2. Reducible degeneration. We consider abelian differentials on a general family of Riemann surfaces stably degenerating to a reducible singular complex curve which is a union of two Riemann surfaces.

Let \mathcal{R} be a family of Riemann surfaces of genus $g > 0$ with symplectic bases $\{a_i, b_i\}_{1 \leq i \leq g}$ and local coordinates u at marked points p which is defined over a small

disk centered at $y = 0$ in \mathbb{C} . Take integers $g_1, g_2 \geq 0$ satisfying $g_1 + g_2 = g$, and assume that under $y \rightarrow 0$, \mathcal{R} degenerates to a union of two Riemann surfaces R_1, R_2 of genus g_1, g_2 respectively by identifying $p_1 \in R_1$ and $p_2 \in R_2$, and $\{a_i, b_i\}_{1 \leq i \leq g}$ becomes, as a homology basis, to the union of symplectic bases

$$\{a_{1,i}, b_{1,i}\}_{1 \leq i \leq g_1}, \{a_{2,i}, b_{2,i}\}_{g_1+1 \leq i \leq g_1+g_2}$$

of R_1, R_2 respectively. Let $\{\omega_1, \dots, \omega_g\}$ be the basis of the space of relative abelian differentials of the first kind on \mathcal{R} normalized for $\{a_i\}_{1 \leq i \leq g}$, and $\omega^{(n)}$ be the unique relative abelian differential of the second kind with order $n + 1$ on \mathcal{R} normalized for $(\{a_i\}_{1 \leq i \leq g}, p, u)$.

Theorem 5.2.

(1) Under $y \rightarrow 0$, $\{\omega_1, \dots, \omega_g\}$ becomes

$$\{\omega_{1,1}, \dots, \omega_{1,g_1}, \omega_{2,g_1+1}, \dots, \omega_{2,g_1+g_2}\},$$

where $\{\omega_{1,1}, \dots, \omega_{1,g_1}\}$ (resp. $\{\omega_{2,g_1+1}, \dots, \omega_{2,g_1+g_2}\}$) are the bases of the spaces of abelian differentials of the first kind on R_1 (resp. R_2) normalized for $\{a_{1,i}\}_{1 \leq i \leq g_1}$ (resp. $\{a_{2,i}\}_{g_1+1 \leq i \leq g_1+g_2}$).

(2) Assume that under $y \rightarrow 0$, (p, u) becomes a point on R_1 with local coordinate which we denote by the same symbol. Then $\lim_{y \rightarrow 0} \omega^{(n)}$ is the unique abelian differential of the second kind with order $n + 1$ on R_1 normalized for $(\{a_{1,i}\}_{1 \leq i \leq g_1}, p, u)$.

Proof. As is shown in the proof of Theorem 5.1, it is enough to prove the assertions (1) and (2) in the case when \mathcal{R} is obtained from a generalized Tate curve as in 2.2 (P4). Let $\Delta_i = (V_i, E_i, T_i)$ ($i = 1, 2$) be two stable graphs such that $\text{rank}_{\mathbb{Z}} H_1(\Delta_i, \mathbb{Z}) = g_i$, $V_i = \{v_i\}$ and $T_1 = \{t, t_1\}$, $T_2 = \{t_2\}$, and Δ be a stable graph containing a nonloop oriented edge e which is obtained from Δ_1, Δ_2 by connecting t_1, t_2 with e , where $v_e = v_1, v_{-e} = v_2$. Take generators $\gamma_{1,1}, \dots, \gamma_{1,g_1}$ (resp. $\gamma_{2,g_1+1}, \dots, \gamma_{2,g_1+g_2}$) of $\Gamma_{\Delta_1} \cong \pi_1(\Delta_1, v_1)$ (resp. $\Gamma_{\Delta_2} \cong \pi_1(\Delta_2, v_2)$), and symplectic bases $\{a_{1,i}, b_{1,i}\}_{1 \leq i \leq g_1}$ (resp. $\{a_{2,i}, b_{2,i}\}_{g_1+1 \leq i \leq g_1+g_2}$) of families \mathcal{R}_{Δ_1} (resp. \mathcal{R}_{Δ_2}) of Riemann surfaces obtained from \mathcal{C}_{Δ_1} (resp. \mathcal{C}_{Δ_2}) such that $b_{1,i}$ (resp. $b_{2,i}$) correspond to $\gamma_{1,i}$ (resp. $\gamma_{2,i}$). If ϕ_e is defined as in 2.2, then

$$\gamma_i = \begin{cases} \gamma_{1,i} & (1 \leq i \leq g_1), \\ \phi_e \cdot \gamma_{2,i} \cdot \phi_e^{-1} & (g_1 + 1 \leq i \leq g_1 + g_2) \end{cases}$$

are generators of $\Gamma_{\Delta} \cong \pi_1(\Delta, v_1)$. Therefore, by Theorem 3.4, the universal abelian differentials of the first kind

$$\omega_i = \sum_{\gamma \in \Gamma_{\Delta} / \langle \gamma_i \rangle} \left(\frac{1}{z - \gamma(\alpha_i)} - \frac{1}{z - \gamma(\alpha_{-i})} \right) dz \quad (1 \leq i \leq g)$$

on \mathcal{C}_Δ give the basis of the space of relative abelian differentials of the first kind on \mathcal{R}_Δ normalized for

$$\{a_1, \dots, a_g\} = \{a_{1,1}, \dots, a_{1,g_1}, a_{2,g_1+1}, \dots, a_{2,g_1+g_2}\}.$$

By Proposition 3.1 (2),

$$\omega_i|_{y_e=0} = \begin{cases} \sum_{\gamma \in \Gamma_{\Delta_1}/\langle \gamma_i \rangle} \left(\frac{1}{z - \gamma(\alpha_i)} - \frac{1}{z - \gamma(\alpha_{-i})} \right) dz & \text{on } \mathcal{C}_{\Delta_1} \text{ for } 1 \leq i \leq g_1, \\ 0 & \text{on } \mathcal{C}_{\Delta_2} \text{ for } g_1 + 1 \leq i \leq g_1 + g_2, \end{cases}$$

and hence $\{\omega_i|_{y_e=0}\}_{1 \leq i \leq g_1}$ gives the basis of the space of relative abelian differentials of the first kind on \mathcal{R}_{Δ_1} normalized for $\{a_{1,i}\}_{1 \leq i \leq g_1}$. Similarly, $\{\omega_i|_{y_e=0}\}_{g_1+1 \leq i \leq g_1+g_2}$ gives the basis of the space of relative abelian differentials of the first kind on \mathcal{R}_{Δ_2} normalized for $\{a_{2,i}\}_{g_1+1 \leq i \leq g_1+g_2}$. For an integer $k > 1$ and the unique tail t of Δ , by Theorem 3.4, the universal abelian differential of the second kind

$$\omega_{t,k} = \sum_{\gamma \in \Gamma_\Delta} \frac{d\gamma(z)}{(\gamma(z) - x_t)^k}$$

on \mathcal{C}_Δ gives the unique relative abelian differential of the second kind with order k on \mathcal{R}_Δ normalized for $(\{a_i\}_{1 \leq i \leq g}, p_t, z - p_t)$. By Proposition 3.1 (3),

$$\omega_{t,k}|_{y_e=0} = \begin{cases} \sum_{\gamma \in \Gamma_{\Delta_1}} \frac{d\gamma(z)}{(\gamma(z) - x_t)^k} & \text{on } \mathcal{C}_{\Delta_1}, \\ 0 & \text{on } \mathcal{C}_{\Delta_2}, \end{cases}$$

and hence $\omega_{t,k}|_{y_e=0}$ gives the unique relative abelian differential of the second kind with order k on \mathcal{R}_{Δ_1} normalized for $(\{a_i\}_{1 \leq i \leq g-1}, p_t, z - x_t)$. Therefore, the assertions (1) and (2) hold for $(\mathcal{R}_\Delta, \{a_i, b_i\}, p_t)$, where $z - p_t$ is replaced with general local coordinates at p_t . This completes the proof. \square

6. Tau functions and quasi-periodic KP solutions

6.1. Tau function and quasi-periodic KP solution. We fix a Riemann surface R of genus $g > 0$ with symplectic basis $\{a_i, b_i\}_{1 \leq i \leq g}$, a point $p \in R$ and a local coordinate u at p , and denote the whole data by $X = (R, \{a_i, b_i\}_{1 \leq i \leq g}, p, u)$. Then there exists a unique basis $\{\omega_1, \dots, \omega_g\}$ of the space of abelian differentials of the first kind on R normalized for $\{a_i\}$, namely $\int_{a_i} \omega_j = 2\pi\sqrt{-1}\delta_{ij}$ ($1 \leq i, j \leq g$), and

$$Z(X) = (Z_{i,j}(X))_{1 \leq i, j \leq g} = \left(\frac{1}{2\pi\sqrt{-1}} \int_{b_i} \omega_j \right)_{1 \leq i, j \leq g}$$

is the period matrix of $(R, \{a_i, b_i\}_{1 \leq i \leq g})$. Take $r_{j,m}(X) \in \mathbb{C}$ such that

$$\omega_j = \sum_{m=1}^{\infty} r_{j,m}(X) u^{m-1} du \quad \text{at } p,$$

and put $r_m(X) = (r_{1,m}(X), \dots, r_{g,m}(X))$. For each positive integer n , there exists a unique abelian differential $\omega^{(n)}$ of the second kind on R satisfying

- $\omega^{(n)}$ is holomorphic outside p ,
- $\omega^{(n)} = \left(\frac{1}{u^{n+1}} + \sum_{m=1}^{\infty} \frac{q_{n,m}(X)}{n} u^{m-1} \right) du$ at p for some $q_{n,m}(X) \in \mathbb{C}$,
- $\int_{a_i} \omega^{(n)} = 0$ for any $i = 1, \dots, g$.

For $z = (z_i)_{1 \leq i \leq g} \in \mathbb{C}^g$, we denote the Riemann theta function of $(R, \{a_i, b_i\}_{1 \leq i \leq g})$ by

$$\Theta_{(R, \{a_i, b_i\})}(z) = \sum_{v \in \mathbb{Z}^g} \left\{ \prod_{i,j=1}^g \exp(\pi \sqrt{-1} Z_{i,j}(X))^{v_i v_j} \prod_{i=1}^g \exp(z_i)^{v_i} \right\}, \quad (6.1.1)$$

where $v = (v_i)_{1 \leq i \leq g}$. Let t_m ($m = 1, 2, \dots$) be indeterminates, and put $t = (t_1, t_2, \dots)$. Following [20, Definition 5.5] (cf. [3, (14)], [19], [35, (24)]), for the addressed Riemann surface X with line bundle of degree 0 corresponding to $c = (c_1, \dots, c_g) \in \mathbb{C}^g$, the associated *tau function* $\tau(t, X_c)$ is defined as

$$\tau(t, X_c) = \exp \left(\frac{1}{2} \sum_{n,m=1}^{\infty} q_{n,m}(X) t_n t_m \right) \cdot \Theta_{(R, \{a_i, b_i\})} \left(c + \sum_{m=1}^{\infty} r_m(X) t_m \right). \quad (6.1.2)$$

Then $\tau(t, X_c)$ is regarded as an element of $\mathbb{C}[[t]] = \mathbb{C}[[t_1, t_2, \dots]]$ by using the expression

$$\prod_{i=1}^g \exp \left(\sum_{m=1}^{\infty} r_{i,m}(X) t_m \right)^{v_i} = \sum_{n=0}^{\infty} \frac{1}{n!} \left(\sum_{m=1}^{\infty} \left(\sum_{i=1}^g v_i \cdot r_{i,m}(X) \right) t_m \right)^n.$$

If $\Theta_{(R, \{a_i, b_i\})}(c) \neq 0$, then we put

$$\frac{\tau(t - [\alpha], X_c)}{\tau(t, X_c)} = 1 + \sum_{k=1}^{\infty} w_k(t, X_c) \alpha^k, \quad (6.1.3)$$

where $[\alpha] = (\alpha, \alpha^2/2, \alpha^3/3, \dots)$, and define two micro-differential operators

$$W(t, X_c) = 1 + \sum_{k=1}^{\infty} w_k(t, X_c) \partial_x^{-k} \quad (6.1.4)$$

and

$$L(t, X_c) = W(t+x, X_c) \cdot \partial_x \cdot W(t+x, X_c)^{-1} \quad (6.1.5)$$

with coefficients in $\mathbb{C}[[x, t]]$, where $t+x = (t_1+x, t_2, t_3, \dots)$. Then it is known (cf. [23, 34]) that $L(t, X_c)$ satisfies the KP hierarchy

$$\frac{\partial L}{\partial t_n} = [(L^n)_+, L] \quad (n = 1, 2, \dots).$$

In particular,

$$u_1(x, t_2, t_3) = \frac{\partial^2}{\partial x^2} \log \Theta_{(R, \{a_i, b_i\})} (c + xr_1(X) + t_2 r_2(X) + t_3 r_3(X)) + q_{1,1}(X)$$

satisfies the KP equation

$$\frac{3}{4} \frac{\partial^2 u_1}{\partial t_2^2} - \frac{\partial}{\partial x} \left(\frac{\partial u_1}{\partial t_3} - \frac{1}{4} \frac{\partial^3 u_1}{\partial x^3} - 3u_1 \frac{\partial u_1}{\partial x} \right) = 0.$$

Since $L(t, X_c)$ and $u_1(x, t_2, t_3)$ are expressed by theta functions with quasi-periodicity, they are called *quasi-periodic KP solutions*.

6.2. Universal tau function. Let $\Delta = (V, E, T)$ be a stable graph of $(g, 1)$ -type with $T = \{t_0\}$, and fix generators ρ_1, \dots, ρ_g of $\pi_1(\Delta)$. Then we denote by \mathcal{X}_Δ the family \mathcal{R}_Δ of Riemann surfaces associated with \mathcal{C}_Δ with symplectic basis $\{a_i, b_i\}_{1 \leq i \leq g}$, marked point $x_{t_0} \in P_{v_{t_0}}$ and local coordinate $z - x_{t_0}$, where b_i correspond to $[\rho_i] \in H_1(\Delta, \mathbb{Z})$ ($i = 1, \dots, g$).

Theorem 6.1.

(1) Denote by $\mathbb{Q}[\gamma_1^{\pm 1}, \dots, \gamma_g^{\pm 1}]$ the \mathbb{Q} -algebra of Laurent polynomials in variables $\gamma_1, \dots, \gamma_g$. Then there exists the universal tau function $\tau(t, (\mathcal{C}_\Delta)_\gamma)$ as an element of

$$\mathcal{A}_\Delta = A_\Delta \left[\sqrt{P_{ii}} \ (i = 1, \dots, g) \right] \widehat{\otimes}_{\mathbb{Z}} \mathbb{Q}[\gamma_1^{\pm 1}, \dots, \gamma_g^{\pm 1}][[t]]$$

which becomes the tau functions of members of $(\mathcal{X}_\Delta)_c$, where $\exp(c_i) = \gamma_i$.

(2) The universal tau function $\tau(t, (\mathcal{C}_\Delta)_\gamma)$ gives rise to a KP solution as in 6.1.

(3) By 2.2 (P2), $\tau(t, (\mathcal{C}_\Delta)_\gamma)$ gives tau functions of all marked Mumford curves with dual graph Δ over nonarchimedean complete valuation fields of characteristic 0, and these tau functions give rise to KP solutions as in 6.1.

Proof. The assertion (1) follows from Theorems 3.4 and 3.6. By [14, Proposition 3.11], P_{ii} ($i = 1, \dots, g$) belong to the ideal of A_Δ generated by y_e ($e \in E$), and $\tau(t, (\mathcal{C}_\Delta)_\gamma) - 1$ belongs to the ideal of \mathcal{A}_Δ generated by $\sqrt{P_{ii}}$ ($i = 1, \dots, g$), and hence $\tau(t, (\mathcal{C}_\Delta)_\gamma)$ defines a micro-differential operator $L(t, (\mathcal{C}_\Delta)_\gamma)$ with coefficients in $\mathcal{A}_\Delta[[x]]$. Then the assertion (2) follows from (1), and (3) follows from (2). \square

Remark 6.2. By the results in Section 4, one can see that $\tau(t, (\mathcal{C}_\Delta)_\gamma)$ (Δ : stable graphs of $(g, 1)$ -type) gives rise to tau functions over the family of all degenerating Riemann surfaces, and that of all Mumford curves.

7. Behavior of tau functions

7.1. Variation of tau functions. Let the notation be as in Section 6, and put

$$c = 2\pi\sqrt{-1}(\alpha + Z(X)\beta),$$

where $\alpha = (\alpha_i)_{1 \leq i \leq g} \in \mathbb{C}^g, \beta = (\beta_i)_{1 \leq i \leq g} \in \mathbb{R}^g$. Then we study the asymptotic behavior of the tau function (6.1.2) under degenerations of X as in Section 6, and define the associated modified tau function as its regularized limit.

7.2. Irreducible degeneration. Let the notation be as in 5.1, especially

$$\mathcal{X} = (\mathcal{R}, \{a_i, b_i\}_{1 \leq i \leq g}, p, u), \quad X' = (R', \{a'_i, b'_i\}_{1 \leq i \leq g-1}, p, u),$$

and \mathcal{R} degenerates to R' by $y \rightarrow 0$. Then we study the relationship between the the tau functions of \mathcal{X} and X' .

Theorem 7.1. *Assume that $\beta_g \notin \mathbb{Z} + 1/2$, and take a unique integer $\bar{\beta}$ such that $|\bar{\beta} + \beta_g| < 1/2$.*

(1) *Under $y \rightarrow 0$,*

$$\exp(-\pi\sqrt{-1}(\bar{\beta} + 2\beta_g)\bar{\beta}Z_{g,g}(\mathcal{X})) \cdot \Theta_{(\mathcal{R}, \{a_i, b_i\})} \left(c + \sum_{m=1}^{\infty} r_m(\mathcal{X})t_m \right)$$

tends to

$$\exp \left(2\pi\sqrt{-1} \left(\alpha_g + \sum_{i=1}^{g-1} \beta_i \bar{Z}_{i,g} \right) + \sum_{m=1}^{\infty} \bar{r}_{g,m} t_m \right)^{\bar{\beta}} \cdot \Theta_{(R', \{a'_i, b'_i\})} \left(\bar{c} + \sum_{m=1}^{\infty} \bar{r}_m t_m \right),$$

where

$$\begin{aligned} \bar{Z}_{i,g} &= \lim_{y \rightarrow 0} Z_{i,g}(\mathcal{X}) \quad (1 \leq i \leq g-1), \\ \bar{c} &= (\alpha_i + \bar{Z}_{i,g}(\bar{\beta} + \beta_g))_{1 \leq i \leq g-1}, \\ \bar{r}_m &= (r_{i,m}(X'))_{1 \leq i \leq g-1}, \end{aligned}$$

and $\sum_{m=1}^{\infty} \bar{r}_{g,m} u^{m-1} du$ is the expansion of ω' given in Theorem 5.1 (1).

(2) *Under $y \rightarrow 0$,*

$$\exp(-\pi\sqrt{-1}(\bar{\beta} + 2\beta_g)\bar{\beta}Z_{g,g}(\mathcal{X})) \cdot \tau(t, \mathcal{X}_c)$$

tends to

$$\exp \left(2\pi\sqrt{-1} \left(\alpha_g + \sum_{i=1}^{g-1} \beta_i \bar{Z}_{i,g} \right) + \sum_{m=1}^{\infty} \bar{r}_{g,m} t_m \right)^{\bar{\beta}}$$

which we call the associated modified tau function.

Proof. By (6.1.1), the assertion (1) follows from the formula

$$\begin{aligned} & \Theta \left(c + \sum_{m=1}^{\infty} r_m t_m \right) \\ &= \sum_{v \in \mathbb{Z}^g} \left\{ \prod_{i,j=1}^g \exp(\pi\sqrt{-1} v_i Z_{i,j} v_j) \prod_{i=1}^g \exp(2\pi\sqrt{-1} \alpha_i)^{v_i} \prod_{i,j=1}^g \exp(2\pi\sqrt{-1} v_i Z_{i,j} \beta_j) \right. \\ & \quad \left. \times \prod_{i=1}^g \exp \left(\sum_{m=1}^{\infty} r_{i,m} t_m \right)^{v_i} \right\} \\ &= \prod_{i,j=1}^g \exp(-\pi\sqrt{-1} \beta_i Z_{i,j} \beta_j) \sum_{v \in \mathbb{Z}^g} \left\{ \prod_{i,j=1}^g \exp(\pi\sqrt{-1} (v_i + \beta_i) Z_{i,j} (v_j + \beta_j)) \right. \\ & \quad \left. \times \prod_{i=1}^g \exp(2\pi\sqrt{-1} \alpha_i)^{v_i} \sum_{n=0}^{\infty} \frac{1}{n!} \left(\sum_{m=1}^{\infty} \left(\sum_{i=1}^g v_i r_{i,m} \right) t_m \right)^n \right\}, \end{aligned}$$

and the assertion (2) follows from (1). \square

Theorem 7.2. Assume that $\beta_g \in \mathbb{Z} + 1/2$, and put $\bar{\beta}_1 = -\beta_g - 1/2$, $\bar{\beta}_2 = -\beta_g + 1/2$ which are the nearest integers to β_g .

(1) Under $y \rightarrow 0$,

$$\exp(\pi\sqrt{-1} (\beta_g^2 - 1/4) Z_{g,g}(\mathcal{X})) \cdot \Theta_{(\mathcal{R}, \{a_i, b_i\})} \left(c + \sum_{m=1}^{\infty} r_m(X) t_m \right)$$

tends to

$$\begin{aligned} & \exp \left(2\pi\sqrt{-1} \left(\alpha_g + \sum_{i=1}^{g-1} \beta_i \bar{Z}_{i,g} \right) + \sum_{m=1}^{\infty} \bar{r}_{g,m} t_m \right)^{\bar{\beta}_1} \\ & \quad \times \Theta_{(R', \{a'_i, b'_i\})} \left(\bar{c}_1 + \sum_{m=1}^{\infty} \bar{r}_m t_m \right) \\ & + \exp \left(2\pi\sqrt{-1} \left(\alpha_g + \sum_{i=1}^{g-1} \beta_i \bar{Z}_{i,g} \right) + \sum_{m=1}^{\infty} \bar{r}_{g,m} t_m \right)^{\bar{\beta}_2} \\ & \quad \times \Theta_{(R', \{a'_i, b'_i\})} \left(\bar{c}_2 + \sum_{m=1}^{\infty} \bar{r}_m t_m \right), \end{aligned}$$

where

$$\begin{aligned}\bar{Z}_{i,g} &= \lim_{y \rightarrow 0} Z_{i,g}(\mathcal{X}) \quad (1 \leq i \leq g-1), \\ \bar{c}_k &= (\alpha_i + \bar{Z}_{i,g}(\bar{\beta}_k + \beta_g))_{1 \leq i \leq g-1} \quad (k = 1, 2), \\ \bar{r}_m &= (r_{i,m}(X'))_{1 \leq i \leq g-1},\end{aligned}$$

and $\sum_{m=1}^{\infty} \bar{r}_{g,m} u^{m-1} du$ is the expansion of ω' given in Theorem 5.1 (1).

(2) Under $y \rightarrow 0$,

$$\exp(\pi\sqrt{-1}(\beta_g^2 - 1/4)Z_{g,g}(\mathcal{X})) \cdot \tau(t, \mathcal{X}_c)$$

tends to

$$\begin{aligned}& \exp\left(2\pi\sqrt{-1}\left(\alpha_g + \sum_{i=1}^{g-1} \beta_i \bar{Z}_{i,g}\right) + \sum_{m=1}^{\infty} \bar{r}_{g,m} t_m\right)^{\bar{\beta}_1} \cdot \tau(t, X'_{\bar{c}_1}) \\ & + \exp\left(2\pi\sqrt{-1}\left(\alpha_g + \sum_{i=1}^{g-1} \beta_i \bar{Z}_{i,g}\right) + \sum_{m=1}^{\infty} \bar{r}_{g,m} t_m\right)^{\bar{\beta}_2} \cdot \tau(t, X'_{\bar{c}_2})\end{aligned}$$

which we call the associated modified tau function.

Proof. One can prove the assertion as in Theorem 7.1. \square

7.3. Reducible degeneration. Let the notation and assumption be as in 5.2, especially

$$\mathcal{X} = (\mathcal{R}, \{a_i, b_i\}_{1 \leq i \leq g}, p, u), \quad X_1 = (R_1, \{a_{1,i}, b_{1,i}\}_{1 \leq i \leq g_1}, p, u),$$

and \mathcal{R} degenerates to the union of R_1, R_2 by $y \rightarrow 0$.

Theorem 7.3. *Let $c = (c_1, \dots, c_g)$ be an element of \mathbb{C}^g .*

(1) Under $y \rightarrow 0$, $\Theta_{(\mathcal{R}, \{a_i, b_i\})}(c + \sum_{m=1}^{\infty} r_m(\mathcal{X})t_m)$ tends to

$$\Theta_{(R_1, \{a_{1,i}, b_{1,i}\})}\left(c' + \sum_{m=1}^{\infty} r_m(X_1)t_m\right) \cdot \Theta_{(R_2, \{a_{2,i}, b_{2,i}\})}(c''),$$

where $c' = (c_1, \dots, c_{g_1})$ and $c'' = (c_{g_1+1}, \dots, c_{g_1+g_2})$.

(2) Under $y \rightarrow 0$, $\tau(t, \mathcal{X}_c)$ tends to

$$\tau(t, (X_1)_{c'}) \cdot \Theta_{(R_2, \{a_{2,i}, b_{2,i}\})}(c'')$$

which we call the associated modified tau function.

Proof. By Theorem 5.2,

$$\lim_{y \rightarrow 0} Z_{i,j} = \begin{cases} 0 & (1 \leq i \leq g_1, g_1 + 1 \leq j \leq g_1 + g_2), \\ 0 & (g_1 + 1 \leq i \leq g_1 + g_2, 1 \leq j \leq g_1), \\ (i, j) \text{ period of } (R_1, \{a_{1,i}, b_{1,i}\}) & (1 \leq i, j \leq g_1), \\ (i, j) \text{ period of } (R_2, \{a_{2,i}, b_{2,i}\}) & (g_1 + 1 \leq i, j \leq g_1 + g_2). \end{cases}$$

Therefore, the assertions (1) and (2) follow from Theorem 5.2. \square

7.4. KP solutions and generalized soliton. By using Theorems 7.1, 7.2 and 7.3 repeatedly, for degenerations to stable complex curves with suitably chosen symplectic bases, we can express the associated modified tau functions as mixtures of quasi-periodic solutions and solitons. Then we have the following assertion.

Theorem 7.4. *The modified tau functions give rise to KP solutions as in 6.1.*

Proof. We only prove the assertion in the case of Theorem 7.1 since other cases can be shown similarly. By (6.1.3),

$$\tau(t, \mathcal{X}_c), \exp(-\pi\sqrt{-1}(\bar{\beta} + 2\beta_g)\bar{\beta}Z_{g,g}(\mathcal{X})) \cdot \tau(t, \mathcal{X}_c)$$

gives the same micro-differential operators in (6.1.4) and (6.1.5), and hence their limits under $y \rightarrow 0$ is associated with the modified tau function, and gives rise to a KP solution. \square

In particular, we have solutions expressed by rational functions containing soliton solutions as follows. Let $\Delta_0 = (V_0, E_0, T_0)$ be the stable graph of $(g, 1)$ -type consisting of one vertex v_0 , and identify E_0, T_0 with $\{1, \dots, g\}, \{t_0\}$ respectively. Then under $y_i \rightarrow 0$ ($i = 1, \dots, g$), \mathcal{X}_{Δ_0} degenerates to the irreducible complex curve $\mathbb{P}^1/(x_i = x_{-i})$ which is obtained from $\mathbb{P}^1 = P_{v_0}$ identifying $2g$ points $x_{\pm i}$ ($i = 1, \dots, g$) in pairs x_i, x_{-i} with local coordinate $u = z - x_{t_0}$ at x_{t_0} .

Theorem 7.5. *Let $\beta = (n_i - 1/2)_{1 \leq i \leq g}$ be an element of $(\mathbb{Z} + 1/2)^g$, and put*

$$c = 2\pi\sqrt{-1}(\alpha + Z(\mathcal{X}_{\Delta_0})\beta)$$

as in 7.1, Then there exists $\alpha' = (\alpha'_i)_{1 \leq i \leq g} \in \mathbb{C}^g$ such that the associated modified tau function

$$\lim_{y_1, \dots, y_g \rightarrow 0} \left(\prod_{i=1}^g \exp(\pi\sqrt{-1}(n_i^2 - n_i)Z_{ii}(\mathcal{X}_{\Delta_0})) \cdot \tau(t, (\mathcal{X}_{\Delta_0})_c) \right)$$

is expressed as

$$\sum_{u \in \{0,1\}^g} \left\{ \prod_{1 \leq i < j \leq g} \left(\frac{(x_i - x_j)(x_{-i} - x_{-j})}{(x_{-i} - x_j)(x_i - x_{-j})} \right)^{(u_i - n_i)(u_j - n_j)} \right\}$$

$$\times \prod_{i=1}^g \exp \left(\alpha'_i - \sum_{m=1}^{\infty} \left(\frac{1}{(x_i - x_{t_0})^m} - \frac{1}{(x_{-i} - x_{t_0})^m} \right) t_m \right)^{u_i - n_i} \Bigg\}$$

which gives rise to KP solutions as in 6.1. When $n_i = 0$ ($i = 1, \dots, g$), these solutions are the soliton solutions.

Proof. By using Theorem 7.2 repeatedly, the modified tau function becomes the sum over $u \in \{0, 1\}^g$ of

$$\prod_{1 \leq i < j \leq g} \exp(2\pi\sqrt{-1}(u_i - n_i)\bar{Z}_{i,j}(u_j - n_j)) \cdot \prod_{i=1}^g \exp \left((u_i - n_i) \left(\alpha'_i + \sum_{m=1}^{\infty} \bar{r}_{i,m} t_m \right) \right).$$

Denote by P_{ij} ($1 \leq i, j \leq g$) the multiplicative periods of \mathcal{C}_{Δ_0} as in 3.3. By Theorem 3.6 (1), if $i \neq j$, then

$$P_{ij} = \frac{(x_i - x_j)(x_{-i} - x_{-j})}{(x_{-i} - x_j)(x_i - x_{-j})} \cdot \prod_{\gamma} \left(1 + \frac{(x_i - x_{-i})(\gamma(x_j) - \gamma(x_{-j}))}{(x_{-i} - \gamma(x_j))(x_i - \gamma(x_{-j}))} \right),$$

where γ runs through $\langle \gamma_i \rangle \setminus (\Gamma_{\Delta_0} - \langle \gamma_i \rangle \langle \gamma_i \rangle) / \langle \gamma_j \rangle$, and hence by Proposition 3.1 (2),

$$\exp(2\pi\sqrt{-1}\bar{Z}_{i,j}) = \lim_{y_1, \dots, y_g \rightarrow 0} P_{ij} = \frac{(x_i - x_j)(x_{-i} - x_{-j})}{(x_{-i} - x_j)(x_i - x_{-j})}.$$

Furthermore, by Proposition 3.1 (2), the abelian differential ω_i ($i = 1, \dots, g$) on \mathcal{C}_{Δ_0} given in 3.1 satisfy

$$\lim_{y_1, \dots, y_g \rightarrow 0} \omega_i = - \left(\frac{1}{(x_i - x_{t_0}) - u} - \frac{1}{(x_{-i} - x_{t_0}) - u} \right) du,$$

and hence

$$\bar{r}_{i,m} = - \left(\frac{1}{(x_i - x_{t_0})^m} - \frac{1}{(x_{-i} - x_{t_0})^m} \right).$$

Then we have the above expression of the modified tau function. The remaining assertion follows from Theorem 7.4 and results in [11, 28]. \square

8. Family of M-curves and real tau functions

8.1. Family of M-curves. Let $\Delta_0 = (V_0, E_0, T_0)$ be the stable graph of (g, n) -type consisting of one vertex v_0 and g oriented loops ρ_1, \dots, ρ_g , and identify E_0 with $\{1, \dots, g\}$. Take $x_{\pm i}$ ($i = 1, \dots, g$), x_t ($t \in T_0$) as real numbers distinct to each other satisfying

$$x_{-1} < x_1 < x_{-2} < x_2 < \dots < x_{-g} < x_g$$

which are regarded as cyclically ordered real points on $P_{v_0} = \mathbb{P}^1$, y_i ($i = 1, \dots, g$) as sufficiently small positive real numbers. Then by the relation (2.2.1), the associated

Schottky uniformized Riemann surfaces are M-curves, namely real algebraic curves with maximal number of components. Furthermore, under comparing \mathcal{C}_Δ with $\mathcal{C}_{\Delta'}$ as in Theorem 4.1, if the parameters x_h ($h \in \pm E \cup T$), y_e ($e \in E$) for \mathcal{C}_Δ are given as real numbers associated with M-curves, then we can construct M-curves from $\mathcal{C}_{\Delta'}$ by taking the parameters s_e ($e \in E'$) as real numbers such that the ratios in Theorem 4.1 are positive. Therefore, by applying the construction of the universal Mumford curve, we have a family $\mathbb{M}_{g,n}$ of n -marked M-curves of genus g .

8.2. Real tau functions.

Theorem 8.1. *Let $\Delta_0 = (V_0, E_0, T_0)$ be the stable graph of $(g, 1)$ -type. Then for each $\gamma \in (\mathbb{R}^\times)^g$, the universal tau function $\tau(t, (\mathcal{C}_{\Delta_0})_\gamma)$ given in Theorem 6.1 (1) gives rise to tau functions defined for all members in $\mathbb{M}_{g,1}$ which are real, namely real-valued for real variables t_m ($m = 1, 2, \dots$).*

Proof. For the above M-curves defined by Δ_0 , Theorem 3.6 implies that

$$\exp(2\pi\sqrt{-1}Z_{i,j}) \in \mathbb{R}^\times, \quad 0 < \exp(\pi\sqrt{-1}Z_{i,i}) = \exp\left(\frac{1}{2} \int_{[\rho_i]} \omega_i\right) < 1$$

since $\int_{[\rho_i]}$ can be taken as the integral along a real line in \mathcal{X}_{Δ_0} . Therefore, the associated tau functions are real, and hence they are extended to real functions defined for all members in $\mathbb{M}_{g,1}$. \square

Remark 8.2. Using results in Section 7, especially Theorem 7.5, one can similarly construct real quasi-periodic KP solutions tending to soliton solutions studied in [1] which are also obtained from theta functions of tropical curves as their tropical limits in [2].

Acknowledgments

This work is partially supported by the JSPS Grant-in-Aid for Scientific Research No. 20K03516.

References

- [1] Abenda, S., Grinevich, P. G.: Rational degeneration of M-curves, totally positive Grassmannians and KP2-solitons. *Comm. Math. Phys.* **361**, 1029–1081 (2018)
- [2] Agostini, D., Fevola, C., Mandelshtamb, Y., Sturmfels, B.: KP solitons from tropical limits. *J. Symb. Comput.* **114**, 282–301 (2023)
- [3] Alvalex-Gaumé, L., Gomez, C., Reina, C.: Loop groups, grassmanians and string theory. *Phys. Lett.* **190B**, 55–62 (1987)

- [4] Bainbridge, B., Möller, M.: The Deligne-Mumford compactification of the real multiplication locus and Teichmüller curves in genus 3. *Acta Math.* **208**, 1–92 (2012)
- [5] Bernatska, J., Enolski, V., Nakayashiki, A.: Sato Grassmannian and degenerate sigma function. to appear in *Comm. Math. Phys.*, arXiv:1810.01224
- [6] Bosch, S., Lütkebohmert, W.: Degenerating abelian varieties. *Topology* **30**, 653–698 (1991)
- [7] Deligne, P., Mumford, D.: The irreducibility of the space of curves of given genus. *Publ. Math. IHES* **36**, 75–109 (1969)
- [8] Fay, J.: Theta functions on Riemann surfaces. *Lecture Notes in Math.* **352**, Berlin-New York, Springer-Verlag, 1973
- [9] Foster, T., Rabinoff, J., Shokrieh, F., Soto, A.: Non-Archimedean and tropical theta functions. *Math. Ann.* **372**, 891–914 (2018)
- [10] Gerritzen, L., Herrlich, F.: The extended Schottky space. *J. reine angew. Math.* **389**, 190–208 (1988)
- [11] Gonzalez-diez, G.: Theta functions on the boundary of moduli space. in R. Donagi (ed.) *Curves, Jacobians, and Abelian Varieties*, *Contemp. Math.* **136**, Providence RI: Amer. Math. Soc., 1992, pp. 185–208.
- [12] Hu, X., Norton, C.: General variational formulas for abelian differentials. *Int. Math. Res. Not.* **2020-12**, 3540–3581 (2020)
- [13] Ichikawa, T.: P -adic theta functions and solutions of the KP hierarchy. *Comm. Math. Phys.* **176**, 383–399 (1996)
- [14] Ichikawa, T.: Generalized Tate curve and integral Teichmüller modular forms. *Amer. J. Math.* **122**, 1139–1174 (2000)
- [15] Ichikawa, T.: Teichmüller groupoids and Galois action. *J. reine angew. Math.* **559**, 95–114 (2003)
- [16] Ichikawa, T.: The universal Mumford curve and its periods in arithmetic formal geometry. arXiv:2010.11517
- [17] Ichikawa, T.: Stable degeneration of abelian differentials and of quasi-periodic solutions of the KP hierarchy. Preprint
- [18] Ihara, Y., Nakamura, H.: On deformation of maximally degenerate stable marked curves and Oda’s problem. *J. reine angew. Math.* **487**, 125–151 (1997)

- [19] Ishibashi, N., Matsuo, Y., Ooguri, H.: Soliton equations and free fermions on Riemann surfaces. *Mod. Phys. Lett.* **A2**, 119 (1987)
- [20] Kawamoto, N., Namikawa, Y., Tsuchiya, A., Yamada, Y.: Geometric realization of conformal field theory on Riemann surfaces. *Comm. Math. Phys.* **116**, 247–308 (1988)
- [21] Knudsen, F. F.: The projectivity of the moduli space of stable curves II, III. *Math. Scand.* **52**, 161–199, 200–212 (1983)
- [22] Knudsen, F. F., Mumford, D.: The projectivity of the moduli space of stable curves I. *Math. Scand.* **39**, 19–55 (1976)
- [23] Krichever, I. M.: Methods of algebraic geometry in the theory of nonlinear equations. *Russ. Math. Surv.* **32**, 185–214 (1977)
- [24] Manin, Yu. I.: Three-dimensional hyperbolic geometry as 1-adic Arakelov geometry. *Invent. Math.* **104**, 223–244 (1991)
- [25] Manin, Yu., Drinfeld, V.: Periods of p -adic Schottky groups. *J. Reine Angew. Math.* **262/263**, 239–247 (1972)
- [26] Mumford, D.: An analytic construction of degenerating curves over complete local rings. *Compos. Math.* **24**, 129–174 (1972)
- [27] Mumford, D.: An analytic construction of degenerating abelian varieties over complete rings. *Compos. Math.* **24**, 239–272 (1972)
- [28] Mumford, D.: *Tata lectures on theta II*. *Progress in Math.* **43**, Basel: Birkhauser, 1983
- [29] Nakayashiki, A.: Degeneration of trigonal curves and solutions of the KP hierarchy. *Nonlinearity* **31**, 3567–3590 (2018)
- [30] Nakayashiki, A.: On reducible degeneration of hyperelliptic curves and soliton solutions. *SIGMA* **15** (2019), arXiv:1808.06748
- [31] Nakayashiki, A.: One step degeneration of trigonal curves and mixing of solitons and quasi-periodic solutions of the KP equation, arXiv:1911.06524
- [32] Sato, M., Sato, Y.: Soliton equations as dynamical systems on infinite dimensional Grassmann manifold. in P. D. Lax (ed.) *Nonlinear Partial Differential Equations in Applied Sciences*, *Lect. Notes Num. Appl. Anal.* **5**, Amsterdam: North-Holland and Tokyo: Kinokuniya, 1982, pp. 259–271
- [33] Schottky, F.: Über eine specielle Function, welche bei einer bestimmten linearen Transformation ihres Arguments unverändert bleibt. *J. reine angew. Math.* **101**, 227–272 (1887)

- [34] Segal, G., Wilson, G.: Loop groups and equations of KdV type. *Publ. Math. IHES* **61**, 5–65 (1985)
- [35] Vafa, C.: Operator formulation on Riemann surfaces. *Phys. Lett.* **190B**, 47–54 (1987)