

DIRICHLET PROBLEM FOR HARMONIC MAPS FROM STRONGLY RECTIFIABLE SPACES INTO REGULAR BALLS IN CAT(1) SPACE

YOHEI SAKURAI

ABSTRACT. In this note, we study the Dirichlet problem for harmonic maps from strongly rectifiable spaces into regular balls in CAT(1) space. Under the setting, we prove that the Korevaar-Schoen energy admits a unique minimizer.

1. INTRODUCTION

1.1. CAT(0) **target**. The study of harmonic maps between singular spaces is one of the central topics in geometric analysis since the pioneering work by Gromov-Schoen [25]. In an earlier stage, the theory has been developed by Korevaar-Schoen [31], Jost [27], [28] and Lin [33], independently. Korevaar-Schoen [31] have introduced an energy for L^2 -maps from a Riemannian domain into a complete metric space, and derived its basic properties; for instance, the existence of energy density, the consistency of Sobolev functions, and its lower semicontinuity. Furthermore, they have proven that if the target space is a CAT(0) space (i.e., non-positively curved space in the sense of A. D. Alexandrov), then the energy admits a unique minimizer for the Dirichlet problem.

One of the research directions is to generalize the theory for non-smooth source spaces. Such attempts have been done by Gregori [17], Eells-Fuglede [11], Kuwae-Shioya [32], and so on. Gregori [17] and Eells-Fuglede [11] have dealt with a domain of a Lipschitz manifold and that of a Riemannian polyhedron, respectively. Kuwae-Shioya [32] have examined a metric measure space satisfying the so-called strongly measure contraction property of Bishop-Gromov type, called SMCPBG space. A typical example of SMCPBG space is an Alexandrov space, which is a metric space equipped with the notion of a lower sectional curvature bound.

In recent years, the theory of metric measure spaces equipped with the notion of a lower Ricci curvature bound has been vastly developed. In the literature, the main research object is the so-called RCD(K, N) space, which has been introduced by Ambrosio-Gigli-Savaré [1], Erbar-Kuwada-Sturm [12] and Gigli [15]. It is natural to ask whether the Korevaar-Schoen theory can be extended to RCD(K, N) spaces. Here we notice that an RCD(K, N) space is not necessarily a SMCPBG space (especially, in the collapsing case) since the Bishop type inequality is required in the SMCPBG condition, and hence the framework of Kuwae-Shioya [32] does not cover RCD(K, N) spaces. To overcome this issue, Gigli-Tyulenev [23] have established a more general framework, which can be applied to RCD(K, N) spaces. In their framework, source spaces are assumed to be a locally uniformly doubling, strongly rectifiable space satisfying a Poincaré inequality (more precisely, see Subsection 2.2). Under this setting, they have formulated a Korevaar-Schoen type energy, and deduced some fundamental results such as the existence of energy density, and its lower semicontinuity. They have concluded the solvability of the Dirichlet problem when the target space is a CAT(0) space (see [23, Theorem 6.4]).

Date: August 16, 2022.

2020 Mathematics Subject Classification. Primary 53E23; Secondly 58E20.

Key words and phrases. Harmonic map; Korevaar-Schoen energy; Strongly rectifiable space; RCD space; CAT(1) space; Regular ball.

Theorem 1.1 ([23]). *Let $(X, \mathbf{d}, \mathbf{m})$ be a locally uniformly doubling, strongly rectifiable space satisfying a Poincaré inequality, and let Ω be a bounded open subset of X with $\mathbf{m}(X \setminus \Omega) > 0$. Let $Y_o = (Y, \mathbf{d}_Y, o)$ be a pointed CAT(0) space. Let $\bar{u} \in \mathbf{KS}^{1,2}(\Omega, Y_o)$ be a Korevaar-Schoen type Sobolev map which determines a boundary value. Then the Korevaar-Schoen type energy $\mathbf{E}_{2, \bar{u}}^\Omega : L^2(\Omega, Y_o) \rightarrow [0, +\infty]$ admits a unique minimizer.*

Thanks to Theorem 1.1, one can introduce the notion of harmonic map from an $\text{RCD}(K, N)$ space to a CAT(0) space. Very recently, Gigli [17] has shown a quantitative Lipschitz estimate for such harmonic maps, and produced a Cheng type Liouville theorem ([7]) based on [10], [18], [22] (see also [41], [42]). Mondino-Semola [36] have also obtained a similar result, independently.

1.2. CAT(1) target. The purpose of this note is to yield an analogue of Theorem 1.1 for the case where the target space is a regular ball in a CAT(1) space (i.e., geodesic ball whose radius is strictly less than $\pi/2$). In the case where the source space is a Riemannian domain, the solvability of the Dirichlet problem has been established by Serbinowski [38]. Similarly to the non-positively curved case, the result in [38] has been extended to non-smooth source spaces (see Eells-Fuglede [11], Fuglede [13], [14] for Riemannian polyhedra, and Huang-Zhang [26] for Alexandrov spaces). We now aim to generalize it for $\text{RCD}(K, N)$ spaces. Our main result is the following:

Theorem 1.2. *Let $(X, \mathbf{d}, \mathbf{m})$ be an infinitesimally Hilbertian, locally uniformly doubling, strongly rectifiable space satisfying a Poincaré inequality, and let Ω be a bounded open subset of X with $\mathbf{m}(X \setminus \Omega) > 0$. Let $Y_o = (Y, \mathbf{d}_Y, o)$ be a pointed CAT(1) space, and let $\bar{B}_\rho(o)$ be a regular ball (i.e., a closed ball of radius $\rho \in (0, \pi/2)$ centered at o). Let $\bar{u} \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$ be a Korevaar-Schoen type Sobolev map which determines a boundary value. Then the Korevaar-Schoen type energy $\mathbf{E}_{2, \bar{u}}^\Omega : L^2(\Omega, \bar{B}_\rho(o)) \rightarrow [0, +\infty]$ admits a unique minimizer.*

Theorem 1.2 enables us to introduce the notion of harmonic map from an $\text{RCD}(K, N)$ space into a regular ball in a CAT(1) space. When the source space is a Riemannian domain, Zhang-Zhong-Zhu [42] have obtained a quantitative Lipschitz estimate and a Choi type Liouville theorem ([8]).

2. PRELIMINARIES

Let $(X, \mathbf{d}, \mathbf{m})$ denote a metric measure space; namely, (X, \mathbf{d}) is a complete separable metric space, and \mathbf{m} is a non-negative Borel measure, which is finite on bounded sets. This section is devoted to preliminaries for metric measure spaces.

2.1. Sobolev functions. We briefly recall the non-smooth differential calculus on metric measure spaces. The readers can refer to [16], [19] for the details.

Let $(X, \mathbf{d}, \mathbf{m})$ be a metric measure space. Let $C([0, 1], X)$ be the set of all curves in X defined on $[0, 1]$ with the uniform topology. For $t \in [0, 1]$ the *evaluation map* $e_t : C([0, 1], X) \rightarrow X$ is defined as $e_t(\gamma) := \gamma_t$. A curve $\gamma \in C([0, 1], X)$ is said to be *absolutely continuous* if there is $f \in L^1(0, 1)$ such that

$$(2.1) \quad \mathbf{d}(\gamma_t, \gamma_s) \leq \int_s^t f(r) \, dr$$

for all $t, s \in [0, 1]$ with $s < t$. For an absolutely continuous γ , the minimal f satisfying (2.1) in the \mathbf{m} -a.e. sense is called the *metric speed*, which is written as $|\dot{\gamma}_t|$. A Borel probability measure π on $C([0, 1], X)$ is said to be a *test plan* if

$$\int_0^1 \int |\dot{\gamma}_t|^2 \, d\pi(\gamma) \, dt < +\infty, \quad (e_t)_\# \pi \leq C \mathbf{m}$$

for some $C > 0$. The *Sobolev class* $S^2(X)$ is the set of all Borel functions $f : X \rightarrow \mathbb{R}$ such that there exists a non-negative $G \in L^2(\mathfrak{m})$ such that

$$(2.2) \quad \int |f(\gamma_1) - f(\gamma_0)| d\pi(\gamma) \leq \int \int_0^1 G(\gamma_t) |\dot{\gamma}_t| dt d\pi(\gamma)$$

for all test plans π . For $f \in S^2(X)$, a non-negative function $G \in L^2(\mathfrak{m})$ satisfying (2.2) is called a *weak upper gradient*, and the minimal one in the \mathfrak{m} -a.e. sense is called the *minimal weak upper gradient*, which is written by $|Df|$. The space

$$W^{1,2}(X) := L^2(\mathfrak{m}) \cap S^2(X)$$

equipped with the norm

$$\|f\|_{W^{1,2}(X)}^2 := \|f\|_{L^2(\mathfrak{m})}^2 + \||Df|\|_{L^2(\mathfrak{m})}^2$$

is called the *Sobolev space*, which is a Banach space.

It is well-known that there exists a unique couple $(L^2(T^*X), d)$ of an $L^2(\mathfrak{m})$ -normed $L^\infty(\mathfrak{m})$ -module and a linear operator $d : S^2(X) \rightarrow L^2(T^*X)$ such that the following hold (see e.g., [19, Theorem 4.1.1]):

- (1) $L^2(T^*X)$ is generated by $\{df \mid f \in S^2(X)\}$;
- (2) for every $f \in S^2(X)$, it holds that

$$|df| = |Df| \quad \mathfrak{m}\text{-a.e.}$$

The space $L^2(T^*X)$ and the operator d are called the *cotangent module* and the *differential*, respectively. The *tangent module* $L^2(TX)$ is defined as the module dual of $L^2(T^*X)$. For $f \in S^2(X)$, let $\text{Grad}(f)$ be the set of all $v \in L^2(TX)$ such that

$$df(v) = |df|^2 = |v|^2 \quad \mathfrak{m}\text{-a.e.}$$

Note that $\text{Grad}(f)$ is non-empty (see [19, Remark 4.2.10]). The following chain rule holds (see e.g., [19, Theorem 4.2.15]):

Theorem 2.1 ([19]). *Let $f \in S^2(X)$ and $v \in \text{Grad}(f)$. If $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is Lipschitz, then $\varphi' \circ f v \in \text{Grad}(\varphi \circ f)$, where $\varphi'(f)$ is arbitrarily defined on the inverse image of the non-differentiability points of φ .*

A metric measure space (X, d, \mathfrak{m}) is said to be *infinitesimally strictly convex* when for every $f \in S^2(X)$ the set $\text{Grad}(f)$ is a singleton; moreover, the unique element is denoted by ∇f . Also, (X, d, \mathfrak{m}) is said to be *infinitesimally Hilbertian* if $W^{1,2}(X)$ is a Hilbert space.

We have the following characterization (see e.g., [19, Theorem 4.3.3]):

Theorem 2.2 ([19]). *The following are equivalent:*

- (1) (X, d, \mathfrak{m}) is infinitesimally Hilbertian;
- (2) $L^2(T^*X)$ and $L^2(TX)$ are Hilbert modules;
- (3) (X, d, \mathfrak{m}) is infinitesimally strictly convex, and for all $f, g \in W^{1,2}(X)$,

$$\nabla(f + g) = \nabla f + \nabla g \quad \mathfrak{m}\text{-a.e.};$$

- (4) (X, d, \mathfrak{m}) is infinitesimally strictly convex, and for all $f, g \in W^{1,2}(X) \cap L^\infty(\mathfrak{m})$,

$$\nabla(fg) = f\nabla g + g\nabla f \quad \mathfrak{m}\text{-a.e.}$$

In the case where (X, d, \mathfrak{m}) is infinitesimally Hilbertian, we denote by $\langle \cdot, \cdot \rangle$ the pointwise scalar product on $L^2(TX)$.

2.2. Doubling, Poincaré and strongly rectifiability. Here we review a volume doubling property, a Poincaré inequality and the strongly rectifiability on metric measure spaces.

Let $(X, \mathbf{d}, \mathbf{m})$ be a metric measure space. For $r > 0$ and $x \in X$, we denote by $B_r(x)$ the open ball of radius r centered at x . A metric measure space $(X, \mathbf{d}, \mathbf{m})$ is said to be *uniformly locally doubling* if for any $R > 0$ there exists $C_R > 0$ such that

$$\mathbf{m}(B_{2r}(x)) \leq C_R \mathbf{m}(B_r(x))$$

for all $x \in X$ and $r \in (0, R)$. We also say that $(X, \mathbf{d}, \mathbf{m})$ satisfies the *Poincaré inequality* if for every $R > 0$ there are $C_R, \lambda = \lambda_R > 0$ such that for any Lipschitz function $f : X \rightarrow \mathbb{R}$,

$$\int_{B_r(x)} |f - f_{B_r(x)}| \, \mathbf{d}\mathbf{m} \leq C_R r \int_{B_{\lambda r}(x)} \text{lip}(f) \, \mathbf{d}\mathbf{m}$$

for all $x \in X$ and $r \in (0, R)$, where $f_{B_r(x)} := \int_{B_r(x)} f \, \mathbf{d}\mathbf{m}$, and $\text{lip}(f)$ is the pointwise Lipschitz constant.

We say that a metric measure space $(X, \mathbf{d}, \mathbf{m})$ is a *d-dimensional strongly rectifiable space* if for every $\varepsilon > 0$ there exists an ε -atlas $\mathcal{A}^\varepsilon := \{(U_i^\varepsilon, \varphi_i^\varepsilon)\}_i$ such that the following hold:

- (1) Borel subsets $\{U_i^\varepsilon\}_i$ of X forms a partition of X except for an \mathbf{m} -negligible set;
- (2) φ_i^ε is a $(1 + \varepsilon)$ -bi-Lipschitz map from U_i^ε to $\varphi_i^\varepsilon(U_i^\varepsilon) \subset \mathbb{R}^d$;
- (3) for some $c_i > 0$, we have

$$c_i \mathcal{L}^d|_{\varphi_i^\varepsilon(U_i^\varepsilon)} \leq (\varphi_i^\varepsilon)_\# \mathbf{m}|_{U_i^\varepsilon} \leq (1 + \varepsilon) c_i \mathcal{L}^d|_{\varphi_i^\varepsilon(U_i^\varepsilon)}.$$

Remark 2.3. For $K \in \mathbb{R}$ and $N \in [1, \infty)$, every $\text{RCD}(K, N)$ space is a locally uniformly doubling, strongly rectifiable space satisfying the Poincaré inequality. Actually, the doubling property is due to [34], [39], [40], the Poincaré inequality is due to [37]. Moreover, as stated in [23, Theorem 2.19], the strongly rectifiability is a consequence of the results by Brué-Semola [6], Gigli-Pasqualetto [20], Kell-Mondino [29], Mondino-Naber [35], De Philippis-Marchese-Rindler [9].

Remark 2.4. Let $(X, \mathbf{d}, \mathbf{m})$ be a strongly rectifiable space, and let $\{\varepsilon_n\}_n$ be a sequence with $\varepsilon_n \downarrow 0$. A family $\{\mathcal{A}^{\varepsilon_n}\}_n$ of atlases is said to be *aligned* if for all n, m and $(U_i^{\varepsilon_n}, \varphi_i^{\varepsilon_n}) \in \mathcal{A}^{\varepsilon_n}$, $(U_j^{\varepsilon_m}, \varphi_j^{\varepsilon_m}) \in \mathcal{A}^{\varepsilon_m}$, the map $\varphi_i^{\varepsilon_n} - \varphi_j^{\varepsilon_m}$ is $(\varepsilon_n + \varepsilon_m)$ -Lipschitz on $U_i^{\varepsilon_n} \cap U_j^{\varepsilon_m}$. It has been observed in [21, Theorem 3.9], [23, Subsection 2.4] that for any sequence $\{\varepsilon_n\}_n$, an aligned family of atlases $\{\mathcal{A}^{\varepsilon_n}\}_n$ exists.

2.3. Approximate metric differentiability. We further review the notion of approximate metric differentiability of maps introduced by Kirchheim [30] and Gigli-Tyulenev [23].

Let $(X, \mathbf{d}, \mathbf{m})$ be a metric measure space, and let (Y, \mathbf{d}_Y) be a metric space. Let $u : X \rightarrow Y$ be a Borel map. The *pointwise Lipschitz constant of u at $x \in X$* is defined as

$$\text{lip}(u)(x) := \overline{\lim}_{y \rightarrow x} \frac{\mathbf{d}_Y(u(x), u(y))}{\mathbf{d}(x, y)}$$

if x is not an isolated point, and $\text{lip}(u)(x) := 0$ otherwise. For a Borel subset $U \subset X$, a point $x \in U$ is said to be a *density point of U* if

$$\lim_{r \downarrow 0} \frac{\mathbf{m}(B_r(x) \cap U)}{\mathbf{m}(B_r(x))} = 1.$$

The *approximate Lipschitz constant* is defined by

$$\text{ap-lip}(u)(x) := \text{ap-}\overline{\lim}_{y \rightarrow x} \frac{\mathbf{d}_Y(u(x), u(y))}{\mathbf{d}(x, y)},$$

where the right hand side means the approximate upper limit (see e.g., [23, Subsection 2.1] for the precise definition). We possess the following (see [23, Proposition 2.5]):

Proposition 2.5 ([23]). *Let $(X, \mathbf{d}, \mathbf{m})$ be a uniformly locally doubling space, and let (Y, \mathbf{d}_Y) be a complete metric space. Let U be a Borel subset of X , and let $u : U \rightarrow Y$ be Lipschitz. Then for every density point $x \in X$ of U we have*

$$\text{lip}(u)(x) = \text{ap-lip}(u)(x).$$

Let $(X, \mathbf{d}, \mathbf{m})$ be a d -dimensional strongly rectifiable space, and let $\{\mathcal{A}^{\varepsilon_n}\}_n$ be an aligned family of atlases for a sequence $\{\varepsilon_n\}_n$ with $\varepsilon_n \downarrow 0$. Let (Y, \mathbf{d}_Y) be a metric space. We denote by sn^d the set of all semi-norms on \mathbb{R}^d equipped with the complete separable metric

$$D(\mathbf{n}_1, \mathbf{n}_2) := \sup_z |\mathbf{n}_1(z) - \mathbf{n}_2(z)|,$$

where the supremum is taken over all $z \in \mathbb{R}^d$ with $|z| \leq 1$. We write $\|\mathbf{n}\| := D(\mathbf{n}, 0)$. We say that a Borel map $u : X \rightarrow Y$ is *approximately metrically differentiable at $x \in X$ relatively to $\{\mathcal{A}^{\varepsilon_n}\}_n$* if the following hold:

- (1) For every n , there exists $i = i_{x,n}$ such that x belongs to $U_i^{\varepsilon_n}$, it is a density point of $U_i^{\varepsilon_n}$ and $\varphi_i^{\varepsilon_n}(x)$ is a density point of $\varphi_i^{\varepsilon_n}(U_i^{\varepsilon_n})$;
- (2) there exists $\text{md}_x(u) \in \text{sn}^d$, called the *metric differential of u at x* , such that

$$\overline{\lim}_{n \rightarrow \infty} \text{ap-}\overline{\lim}_{\substack{y \rightarrow x \\ y \in U_i^{\varepsilon_n}}} \frac{|\mathbf{d}_Y(u(y), u(x)) - \text{md}_x(u)(\varphi_i^{\varepsilon_n}(y) - \varphi_i^{\varepsilon_n}(x))|}{\mathbf{d}(y, x)} = 0.$$

Gigli-Tyulenev [23] has shown the following (see [23, Lemma 3.4]):

Lemma 2.6 ([23]). *Let $(X, \mathbf{d}, \mathbf{m})$ be a strongly rectifiable space, and let $\{\mathcal{A}^{\varepsilon_n}\}_n$ be an aligned family of atlases for a sequence $\{\varepsilon_n\}_n$ with $\varepsilon_n \downarrow 0$. Let (Y, \mathbf{d}_Y) be a complete metric space. If $u : X \rightarrow Y$ is approximately metrically differentiable at $x \in X$ relatively to $\{\mathcal{A}^{\varepsilon_n}\}_n$, then*

$$\text{ap-lip}(u)(x) = \|\text{md}_x(u)\|.$$

Gigli-Tyulenev [23] have further concluded the a.e. approximately metrically differentiability of Borel maps satisfying the Lusin-Lipschitz property (see [23, Proposition 3.6]).

2.4. Korevaar-Schoen space. In this section, we recall the formulation and basic results of the Korevaar-Schoen type energy introduced by Gigli-Tyulenev [23]. Let $(X, \mathbf{d}, \mathbf{m})$ be a metric measure space, and let Ω be an open subset of X . Let $Y_o = (Y, \mathbf{d}_Y, o)$ be a pointed complete metric space.

We denote by $L^0(\Omega, Y)$ the set of all Borel maps (up to \mathbf{m} -a.e. equality) from Ω to Y with separable range. Let $L^2(\Omega, Y_o)$ be the set of all $u \in L^0(\Omega, Y)$ such that

$$\int_{\Omega} \mathbf{d}_Y^2(u(x), o) \, \mathbf{d}\mathbf{m}(x) < +\infty,$$

which is endowed with the distance function

$$\mathbf{d}_{L^2}(u, v) := \left| \int_{\Omega} \mathbf{d}_Y^2(u(x), v(x)) \, \mathbf{d}\mathbf{m}(x) \right|^{1/2}.$$

Since Y is complete, so is $L^2(\Omega, Y_o)$.

Let $u \in L^2(\Omega, Y_o)$. For $r > 0$ and $x \in \Omega$, the *approximating energy* is defined by

$$\text{ks}_{2,r}[u, \Omega](x) := \begin{cases} \left| \int_{B_r(x)} \frac{\mathbf{d}_Y^2(u(x), u(y))}{r^2} \, \mathbf{d}\mathbf{m}(y) \right|^{1/2} & \text{if } B_r(x) \subset \Omega, \\ 0 & \text{otherwise.} \end{cases}$$

The *Korevaar-Schoen space* $\text{KS}^{1,2}(\Omega, Y_o)$ is defined by the set of all $u \in L^2(\Omega, Y_o)$ such that

$$(2.3) \quad \mathbf{E}_2^\Omega(u) := \varliminf_{r \downarrow 0} \int_{\Omega} \text{ks}_{2,r}^2[u, \Omega] \, \mathbf{d}\mathbf{m} < +\infty.$$

Remark 2.7. Let $W^{1,2}(\Omega)$ denote the Sobolev space provided by means of [23, Definition 5.2]. One can also introduce the associated Sobolev space $W^{1,2}(\Omega, Y_o)$ as the set of all $u \in L^2(\Omega, Y_o)$ such that there is $G \in L^2(\Omega)$ such that for all 1-Lipschitz functions $f : Y \rightarrow \mathbb{R}$ we have $f \circ u \in W^{1,2}(\Omega)$ and $|D(f \circ u)| \leq G$ \mathbf{m} -a.e. on Ω (see [23, Definition 5.3]).

We now present basic properties of the Korevaar-Schoen space (see [23, Theorem 5.7]):

Theorem 2.8 ([23]). *Let $(X, \mathbf{d}, \mathbf{m})$ be a locally uniformly doubling, strongly rectifiable space satisfying the Poincaré inequality, and let Ω be an open subset. Let $Y_o = (Y, \mathbf{d}_Y, \mathbf{o})$ be a pointed complete metric space. Then the following hold:*

- (1) $\mathbf{KS}^{1,2}(\Omega, Y_o) = W^{1,2}(\Omega, Y_o)$;
- (2) for every $u \in \mathbf{KS}^{1,2}(\Omega, Y_o)$, there exists a function $e_2[u] \in L^2(\Omega)$, called the energy density, such that

$$\mathbf{ks}_{2,r}[u, \Omega] \rightarrow e_2[u] \quad \mathbf{m}\text{-a.e. on } \Omega \text{ and in } L^2(\Omega) \text{ as } r \downarrow 0.$$

In particular, $\underline{\lim}$ in (2.3) can be limit, and $E_2^\Omega(u)$ can be written as

$$E_2^\Omega(u) = \begin{cases} \int_{\Omega} e_2^2[u] \, d\mathbf{m} & \text{if } u \in \mathbf{KS}^{1,2}(\Omega, Y_o), \\ +\infty & \text{otherwise;} \end{cases}$$

- (3) $E_2^\Omega : L^2(\Omega, Y_o) \rightarrow [0, +\infty]$ is lower semicontinuous;
- (4) any $u \in \mathbf{KS}^{1,2}(\Omega, Y_o)$ is approximately metrically differentiable \mathbf{m} -a.e. in Ω , where we extend u to the whole X by setting it to be constant outside of Ω .

We also need the following (see [23, Propositions 4.6 and 4.19]):

Proposition 2.9 ([23]). *Let $(X, \mathbf{d}, \mathbf{m})$ be a d -dimensional, locally uniformly doubling, strongly rectifiable space satisfying the Poincaré inequality, and let Ω be an open subset. Then there exists $c_d > 0$ depending only on d such that for every $u \in \mathbf{KS}^{1,2}(\Omega, \mathbb{R})$,*

$$(2.4) \quad e_2[u] = c_d |Du| \quad \mathbf{m}\text{-a.e. on } \Omega.$$

Let $W_0^{1,2}(\Omega)$ be the $W^{1,2}(\Omega)$ -closure of the set of all functions in $W^{1,2}(\Omega)$ whose support is contained in Ω . For a fixed $\bar{u} \in \mathbf{KS}^{1,2}(\Omega, Y_o)$, we define

$$\mathbf{KS}_{\bar{u}}^{1,2}(\Omega, Y_o) := \{u \in \mathbf{KS}^{1,2}(\Omega, Y_o) \mid \mathbf{d}_Y(u, \bar{u}) \in W_0^{1,2}(\Omega)\},$$

and the associated energy functional $E_{2,\bar{u}}^\Omega : L^2(\Omega, Y_o) \rightarrow [0, +\infty]$ as

$$E_{2,\bar{u}}^\Omega(u) := \begin{cases} E_2^\Omega(u) & \text{if } u \in \mathbf{KS}_{\bar{u}}^{1,2}(\Omega, Y_o), \\ +\infty & \text{otherwise.} \end{cases}$$

We close this section with the following (see [23, Proposition 5.10]):

Proposition 2.10 ([23]). *Let $(X, \mathbf{d}, \mathbf{m})$ be a locally uniformly doubling, strongly rectifiable space satisfying the Poincaré inequality, and let Ω be an open subset. Let $Y_o = (Y, \mathbf{d}_Y, \mathbf{o})$ be a pointed complete metric space. Then for a fixed $\bar{u} \in \mathbf{KS}^{1,2}(\Omega, Y_o)$, the following hold:*

- (1) $E_{2,\bar{u}}^\Omega$ is lower semicontinuous;
- (2) for any $u, v \in \mathbf{KS}_{\bar{u}}^{1,2}(\Omega, Y_o)$ we have $\mathbf{d}_Y(u, v) \in W_0^{1,2}(\Omega)$.

3. DIRICHLET PROBLEM

In the present section, we give a proof of Theorem 1.2. We proceed along the line of the proof of [38, Theorem 1.16], [23, Theorem 6.4].

3.1. Technical lemma. We begin with a technical lemma. Let $(X, \mathbf{d}, \mathbf{m})$ be a metric measure space, and let Ω be an open subset of X . Let $Y_o = (Y, \mathbf{d}_Y, \mathbf{o})$ be a pointed complete metric space. Let $u, v, w \in L^2(\Omega, Y_o)$. For $\alpha > 0$ and $x \in \Omega$, we define Borel subsets by

$$(3.1) \quad \begin{aligned} \Omega_{v,\alpha,x} &:= \{y \in \Omega \mid \mathbf{d}_Y(v(x), v(y)) \geq \alpha\}, \\ \Omega_{w,\alpha,x} &:= \{y \in \Omega \mid \mathbf{d}_Y(w(x), w(y)) \geq \alpha\}, \\ \Omega_{v,w,\alpha,x} &:= \Omega_{v,\alpha,x} \cup \Omega_{w,\alpha,x}. \end{aligned}$$

We also define a *modified approximating energy* by

$$(3.2) \quad \mathbf{ks}_{2,r}[u, \Omega; v, w, \alpha](x) := \begin{cases} \left| \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \setminus \Omega_{v,w,\alpha,x}} \frac{\mathbf{d}_Y^2(u(x), u(y))}{r^2} \mathbf{d}\mathbf{m}(y) \right|^{1/2} & \text{if } B_r(x) \subset \Omega, \\ 0 & \text{otherwise.} \end{cases}$$

Let us show the following (cf. [38, Lemma 1.12], [26, Lemma 3.1]):

Lemma 3.1. *Let $(X, \mathbf{d}, \mathbf{m})$ be a locally uniformly doubling, strongly rectifiable space satisfying the Poincaré inequality, and let Ω be an open subset. Let $Y_o = (Y, \mathbf{d}_Y, \mathbf{o})$ be a pointed complete metric space. Let $u, v, w \in \mathbf{KS}^{1,2}(\Omega, Y_o)$. We fix $\alpha > 0$. Then we have*

$$(3.3) \quad \mathbf{ks}_{2,r}[u, \Omega; v, w, \alpha] \rightarrow \mathbf{e}_2[u] \quad \mathbf{m}\text{-a.e. on } \Omega \text{ and in } L^2(\Omega) \text{ as } r \downarrow 0.$$

Proof. We first prove the \mathbf{m} -a.e. convergence. Let us consider a point $x \in \Omega$ satisfying the following properties:

- (1) $\mathbf{ks}_{2,r}[u, \Omega](x) \rightarrow \mathbf{e}_2[u](x)$, $\mathbf{ks}_{2,r}[v, \Omega](x) \rightarrow \mathbf{e}_2[v](x)$, $\mathbf{ks}_{2,r}[w, \Omega](x) \rightarrow \mathbf{e}_2[w](x)$ as $r \downarrow 0$;
- (2) $\text{lip } u(x) < +\infty$.

Notice that \mathbf{m} -a.e. $x \in \Omega$ satisfies these properties. Indeed, the first property is a consequence of Theorem 2.8. Concerning the second one, Theorem 2.8 implies that u is approximately metrically differentiable \mathbf{m} -a.e. $x \in \Omega$ relatively to an aligned family $\{\mathcal{A}^{\varepsilon_n}\}_n$ of atlases. For such a $x \in \Omega$, and for each n , there is $i = i_{x,n}$ such that x belongs to a chart $U_i^{\varepsilon_n}$ of $\mathcal{A}^{\varepsilon_n}$, and it is a density point of $U_i^{\varepsilon_n}$. Thanks to [23, (2.15), Corollary 3.10], we may assume that $u|_{U_i^{\varepsilon_n}}$ is Lipschitz. By Proposition 2.5 and Lemma 2.6, $\text{lip } u(x)$ coincides with $\|\mathbf{m}\mathbf{d}_x(u)\|$, which is finite.

We fix a point $x \in \Omega$ discussed in the above paragraph. We verify (3.3) at x . Since

$$\mathbf{ks}_{2,r}^2[u, \Omega; v, w, \alpha](x) = \mathbf{ks}_{2,r}^2[u, \Omega](x) - \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \cap \Omega_{v,w,\alpha,x}} \frac{\mathbf{d}_Y^2(u(x), u(y))}{r^2} \mathbf{d}\mathbf{m}(y),$$

it suffices to show that

$$(3.4) \quad \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \cap \Omega_{v,w,\alpha,x}} \frac{\mathbf{d}_Y^2(u(x), u(y))}{r^2} \mathbf{d}\mathbf{m}(y) \rightarrow 0$$

as $r \downarrow 0$. We have

$$\begin{aligned} & \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \cap \Omega_{v,w,\alpha,x}} \frac{\mathbf{d}_Y^2(u(x), u(y))}{r^2} \mathbf{d}\mathbf{m}(y) \\ & \leq \frac{\mathbf{m}(B_r(x) \cap \Omega_{v,w,\alpha,x})}{\mathbf{m}(B_r(x))} \sup_{y \in B_r(x) \setminus \{x\}} \frac{\mathbf{d}_Y^2(u(x), u(y))}{\mathbf{d}^2(x, y)} \\ & \leq \left(\frac{\mathbf{m}(B_r(x) \cap \Omega_{v,\alpha,x})}{\mathbf{m}(B_r(x))} + \frac{\mathbf{m}(B_r(x) \cap \Omega_{w,\alpha,x})}{\mathbf{m}(B_r(x))} \right) \sup_{y \in B_r(x) \setminus \{x\}} \frac{\mathbf{d}_Y^2(u(x), u(y))}{\mathbf{d}^2(x, y)} \\ & \leq \frac{r^2}{\alpha^2} (\mathbf{ks}_{2,r}^2[u, \Omega](x) + \mathbf{ks}_{2,r}^2[v, \Omega](x)) \sup_{y \in B_r(x) \setminus \{x\}} \frac{\mathbf{d}_Y^2(u(x), u(y))}{\mathbf{d}^2(x, y)}, \end{aligned}$$

where the last inequality follows from

$$\begin{aligned} \frac{\alpha^2 \mathbf{m}(B_r(x) \cap \Omega_{v,\alpha,x})}{r^2 \mathbf{m}(B_r(x))} &\leq \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \cap \Omega_{v,\alpha,x}} \frac{d_Y^2(v(x), v(y))}{r^2} \, d\mathbf{m}(y) \leq \mathbf{ks}_{2,r}^2[v, \Omega](x), \\ \frac{\alpha^2 \mathbf{m}(B_r(x) \cap \Omega_{w,\alpha,x})}{r^2 \mathbf{m}(B_r(x))} &\leq \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \cap \Omega_{w,\alpha,x}} \frac{d_Y^2(w(x), w(y))}{r^2} \, d\mathbf{m}(y) \leq \mathbf{ks}_{2,r}^2[w, \Omega](x). \end{aligned}$$

By the choices of x , and by letting $r \downarrow 0$, we obtain (3.4). We arrive at the \mathbf{m} -a.e. convergence. The L^2 -convergence follows from $\mathbf{ks}_{2,r}^2[u, \Omega; v, w, \alpha] \leq \mathbf{ks}_{2,r}^2[u, \Omega]$ and [23, Lemma 3.14]. \square

3.2. CAT(κ) space. In this subsection, we review basics of CAT(κ) spaces. We refer to [3], [5]. For $\kappa \in \mathbb{R}$, let M_κ be the 2-dimensional space form of constant curvature κ . Let D_κ be the diameter of M_κ ; more precisely,

$$D_\kappa := \begin{cases} \pi/\sqrt{\kappa} & \text{if } \kappa > 0, \\ +\infty & \text{otherwise.} \end{cases}$$

Let (Y, d_Y) be a metric space. A curve $\gamma : [0, 1] \rightarrow Y$ is said to be a *geodesic* if

$$d_Y(\gamma_t, \gamma_s) = |s - t| d_Y(\gamma_0, \gamma_1)$$

for all $t, s \in [0, 1]$. Further, (Y, d) is called a *geodesic space* if for any pair of points $x, y \in Y$, there is a geodesic $\gamma : [0, 1] \rightarrow Y$ with $\gamma_0 = x$ and $\gamma_1 = y$. For $\kappa \in \mathbb{R}$, a complete geodesic space (Y, d_Y) is said to be a *CAT(κ)-space* if it satisfies the so-called κ -triangle comparison principle: For every geodesic triangle Δ_{pqr} in Y whose perimeter is less than $2D_\kappa$, and for every point x on the segment between q and r , it holds that

$$d_Y(p, x) \leq d_\kappa(\bar{p}, \bar{x}),$$

where d_κ is the Riemannian distance on M_κ , and \bar{p} and \bar{x} are comparison points in M_κ .

Let us recall the following basic properties (see e.g., [5, Proposition 1.4]):

Proposition 3.2 ([5]). *For $\kappa \in \mathbb{R}$, let (Y, d_Y) be a CAT(κ) space. Then we have:*

- (1) *For each pair of points $x, y \in Y$ with $d_Y(x, y) < D_\kappa$, there exists a unique geodesic from x to y ; moreover, it varies continuously with its end points;*
- (2) *any balls in Y whose radius is less than $D_\kappa/2$ are convex; namely, for each pair of points in such a ball, the unique geodesic joining them are contained the ball.*

Remark 3.3. In [5], Proposition 3.2 has been deduced from the following: Let $\gamma, \eta : [0, 1] \rightarrow Y$ be geodesics such that

$$\gamma_0 = \eta_0, \quad d_Y(\gamma_0, \gamma_1) + d_Y(\gamma_1, \eta_1) + d_Y(\eta_1, \eta_0) < 2D_\kappa.$$

For $l \in (0, D_\kappa)$, we assume $d_Y(\gamma_0, \gamma_1) \leq l$ and $d_Y(\eta_0, \eta_1) \leq l$. Then there is a constant $c_{\kappa,l} > 0$ depending only on κ, l such that for all $t \in [0, 1]$,

$$(3.5) \quad d_Y(\gamma_t, \eta_t) \leq c_{\kappa,l} d_Y(\gamma_1, \eta_1).$$

We now collect some useful estimates on CAT(1) spaces, which have been stated in [38] without proof. The detailed proofs can be found in [4, Appendix A]. The first one is the following (see [38, p. 11, ESTIMATE I], and also [4, Lemma A.2]):

Lemma 3.4 ([38]). *Let (Y, d_Y) be a CAT(1) space, and let $\gamma, \eta : [0, 1] \rightarrow Y$ be geodesics with*

$$d_Y(\gamma_0, \eta_0) + d_Y(\eta_0, \eta_1) + d_Y(\eta_1, \gamma_1) + d_Y(\gamma_1, \gamma_0) < 2\pi.$$

Then we have

$$\begin{aligned} & \cos^2 \frac{d_Y(\gamma_0, \gamma_1)}{2} d_Y^2(\gamma_{1/2}, \eta_{1/2}) + \frac{1}{4} (d_Y(\eta_0, \eta_1) - d_Y(\gamma_0, \gamma_1))^2 \\ & \leq \frac{1}{2} (d_Y^2(\gamma_0, \eta_0) + d_Y^2(\gamma_1, \eta_1)) \\ & \quad + \text{Cub}(d_Y(\gamma_0, \eta_0), d_Y(\gamma_1, \eta_1), d_Y(\eta_0, \eta_1) - d_Y(\gamma_0, \gamma_1), d_Y(\gamma_{1/2}, \eta_{1/2})), \end{aligned}$$

where Cub denotes the cubic terms in the indicated variables.

The second one is the following (see [38, p. 13, ESTIMATE II], and also [4, Lemma A.4]):

Lemma 3.5 ([38]). *Let (Y, d_Y) be a CAT(1) space, and let $\gamma, \eta : [0, 1] \rightarrow Y$ be geodesics with $\gamma_1 = \eta_1$ and*

$$d_Y(\gamma_0, \eta_0) + d_Y(\eta_0, \eta_1) + d_Y(\eta_1, \gamma_0) < 2\pi.$$

Then for every $t, s \in [0, 1]$ we have

$$\begin{aligned} d_Y^2(\gamma_t, \eta_s) & \leq \frac{\sin^2(1-t)d_Y(\gamma_0, \gamma_1)}{\sin^2 d_Y(\gamma_0, \gamma_1)} (d_Y^2(\gamma_0, \eta_0) - (d_Y(\eta_0, \eta_1) - d_Y(\gamma_0, \gamma_1))^2) \\ & \quad + (1-t)^2 (d_Y(\eta_0, \eta_1) - d_Y(\gamma_0, \gamma_1))^2 + d_Y^2(\gamma_0, \gamma_1)(s-t)^2 \\ & \quad - 2(1-t)(s-t)d_Y(\gamma_0, \gamma_1)(d_Y(\eta_0, \eta_1) - d_Y(\gamma_0, \gamma_1)) \\ & \quad + \text{Cub}(s-t, d_Y(\gamma_0, \eta_0), d_Y(\eta_0, \eta_1) - d_Y(\gamma_0, \gamma_1), d_Y(\gamma_t, \eta_s)). \end{aligned}$$

Remark 3.6. In Lemmas 3.4 and 3.5, we further see that in each term of Cub, at least one of the variables has a exponent which is greater than or equal to two.

3.3. Energy estimates. We here provide several energy estimates. Let (X, d, \mathbf{m}) be a locally uniformly doubling, strongly rectifiable space satisfying the Poincaré inequality, and let Ω be an open subset of X . Let $Y_o = (Y, d_Y, o)$ be a pointed CAT(1) space. For $\rho \in (0, \pi/2)$, we denote by $\bar{B}_\rho(o)$ the closed ball in Y of radius ρ centered at o . In view of Proposition 3.2, $\bar{B}_\rho(o) = (\bar{B}_\rho(o), d_Y, o)$ is a pointed CAT(1) space itself. We call $\bar{B}_\rho(o)$ a *regular ball*.

For $u, v \in L^0(\Omega, \bar{B}_\rho(o))$ and $t \in [0, 1]$, we define a map $\mathbf{G}_t^{u,v} : \Omega \rightarrow \bar{B}_\rho(o)$ by $x \mapsto \mathbf{G}_t^{u(x), v(x)}$, where $\mathbf{G}^{u(x), v(x)}$ denotes the unique geodesic from $u(x)$ to $v(x)$. By virtue of Proposition 3.2, $\mathbf{G}_t^{u,v}$ belongs to $L^0(\Omega, \bar{B}_\rho(o))$. By the same argument as in [23, Section 6], we see that if $u, v \in L^2(\Omega, \bar{B}_\rho(o))$, then $\mathbf{G}_t^{u,v} \in L^2(\Omega, \bar{B}_\rho(o))$, and it is a unique geodesic from u to v .

We begin with the following energy density estimate (cf. [38, Lemma 1.13]):

Lemma 3.7. *For $u, v \in \text{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$, we define $m := \mathbf{G}_{1/2}^{u,v}$, $d := d_Y(u, v)$. Then we have $m \in \text{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$, $d \in \text{KS}^{1,2}(\Omega, \mathbb{R})$, and*

$$(3.6) \quad \cos^2 \frac{d}{2} e_2^2[m] + \frac{1}{4} e_2^2[d] \leq \frac{1}{2} (e_2^2[u] + e_2^2[v]) \quad \mathbf{m}\text{-a.e. on } \Omega.$$

Proof. We check $m \in \text{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$, $d \in \text{KS}^{1,2}(\Omega, \mathbb{R})$. The claim for d immediately follows from the triangle inequality. We examine m . Put $\alpha := 2(\pi - 2\rho) > 0$. For $x \in \Omega$, we define $\Omega_{u,\alpha,x}, \Omega_{v,\alpha,x}, \Omega_{u,v,\alpha,x}$ by (3.1). For all $y \in \Omega \setminus \Omega_{u,v,\alpha,x}$, we see

$$\begin{aligned} d_Y(u(x), v(x)) + d_Y(v(x), u(y)) + d_Y(u(y), u(x)) & < 4\rho + \alpha = 2\pi, \\ d_Y(v(y), u(y)) + d_Y(u(y), v(x)) + d_Y(v(x), v(y)) & < 4\rho + \alpha = 2\pi. \end{aligned}$$

Therefore, by using (3.5) for $l = 2\rho$ twice, for all $y \in \Omega \setminus \Omega_{u,v,\alpha,x}$,

$$d_Y(m(x), m(y)) \leq c_\rho (d_Y(u(x), u(y)) + d_Y(v(x), v(y)))$$

for some constant $c_\rho > 0$ depending only on ρ . This implies

$$\begin{aligned}
(3.7) \quad \mathbf{ks}_{2,r}^2[m, \Omega](x) &\leq \frac{2c_\rho^2}{\mathbf{m}(B_r(x))} \int_{B_r(x) \setminus \Omega_{u,v,\alpha,x}} \left(\frac{\mathbf{d}_Y^2(u(x), u(y))}{r^2} + \frac{\mathbf{d}_Y^2(v(x), v(y))}{r^2} \right) \mathbf{d}\mathbf{m}(y) \\
&\quad + \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \cap \Omega_{u,v,\alpha,x}} \frac{\mathbf{d}_Y^2(m(x), m(y))}{r^2} \mathbf{d}\mathbf{m}(y) \\
&\leq 2c_\rho^2 (\mathbf{ks}_{2,r}^2[u, \Omega](x) + \mathbf{ks}_{2,r}^2[v, \Omega](x)) + \frac{4\rho^2}{r^2} \frac{\mathbf{m}(B_r(x) \cap \Omega_{u,v,\alpha,x})}{\mathbf{m}(B_r(x))} \\
&\leq \left(2c_\rho^2 + \frac{4\rho^2}{\alpha^2} \right) (\mathbf{ks}_{2,r}^2[u, \Omega](x) + \mathbf{ks}_{2,r}^2[v, \Omega](x)),
\end{aligned}$$

where we used

$$\begin{aligned}
\frac{\alpha^2}{r^2} \frac{\mathbf{m}(B_r(x) \cap \Omega_{u,v,\alpha,x})}{\mathbf{m}(B_r(x))} &\leq \frac{\alpha^2}{r^2} \left(\frac{\mathbf{m}(B_r(x) \cap \Omega_{u,\alpha,x})}{\mathbf{m}(B_r(x))} + \frac{\mathbf{m}(B_r(x) \cap \Omega_{v,\alpha,x})}{\mathbf{m}(B_r(x))} \right) \\
&\leq \mathbf{ks}_{2,r}^2[u, \Omega](x) + \mathbf{ks}_{2,r}^2[v, \Omega](x).
\end{aligned}$$

Integrating (3.7) over Ω , and letting $r \downarrow 0$, we arrive at $m \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$.

We prove (3.6). Put $\beta := \pi - 2\rho > 0$. For $x \in \Omega$, we define $\Omega_{u,\beta,x}, \Omega_{v,\beta,x}, \Omega_{u,v,\beta,x}$ as (3.1) again. For all $y \in \Omega \setminus \Omega_{u,v,\beta,x}$, it holds that

$$\mathbf{d}_Y(u(x), v(x)) + \mathbf{d}_Y(v(x), v(y)) + \mathbf{d}_Y(v(y), u(y)) + \mathbf{d}_Y(u(y), u(x)) < 2\beta + 4\rho = 2\pi.$$

Lemma 3.4 tells us that for all $y \in \Omega \setminus \Omega_{u,v,\beta,x}$,

$$\begin{aligned}
&\cos^2 \frac{\mathbf{d}(x)}{2} \mathbf{d}_Y^2(m(x), m(y)) + \frac{1}{4}(\mathbf{d}(x) - \mathbf{d}(y))^2 \\
&\leq \frac{1}{2} (\mathbf{d}_Y^2(u(x), u(y)) + \mathbf{d}_Y^2(v(x), v(y))) \\
&\quad + \text{Cub}(\mathbf{d}_Y(u(x), u(y)), \mathbf{d}_Y(v(x), v(y)), \mathbf{d}(x) - \mathbf{d}(y), \mathbf{d}_Y(m(x), m(y))).
\end{aligned}$$

We write the last term as $\text{Cub}(x, y)$ for short. It follows that

$$\begin{aligned}
(3.8) \quad &\cos^2 \frac{\mathbf{d}(x)}{2} \mathbf{ks}_{2,r}^2[m, \Omega; u, v, \beta](x) + \frac{1}{4} \mathbf{ks}_{2,r}^2[\mathbf{d}, \Omega; u, v, \beta](x) \\
&\leq \frac{1}{2} (\mathbf{ks}_{2,r}^2[u, \Omega](x) + \mathbf{ks}_{2,r}^2[v, \Omega](x)) + \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \setminus \Omega_{u,v,\beta,x}} \frac{\text{Cub}(x, y)}{r^2} \mathbf{d}\mathbf{m}(y),
\end{aligned}$$

where $\mathbf{ks}_{2,r}^2[m, \Omega; u, v, \beta](x)$ and $\mathbf{ks}_{2,r}^2[\mathbf{d}, \Omega; u, v, \beta](x)$ are defined as (3.2).

By Theorem 2.8 and Lemma 3.1, and by the same argument in the proof of Lemma 3.1, \mathbf{m} -a.e. $x \in \Omega$ satisfies the following properties:

- (1) $\mathbf{ks}_{2,r}[m, \Omega; u, v, \beta](x) \rightarrow \mathbf{e}_2[m](x)$, $\mathbf{ks}_{2,r}[\mathbf{d}, \Omega; u, v, \beta](x) \rightarrow \mathbf{e}_2[\mathbf{d}](x)$ as $r \downarrow 0$;
- (2) $\mathbf{ks}_{2,r}[u, \Omega](x) \rightarrow \mathbf{e}_2[u](x)$, $\mathbf{ks}_{2,r}[v, \Omega](x) \rightarrow \mathbf{e}_2[v](x)$ as $r \downarrow 0$;
- (3) $\text{lip } u(x), \text{lip } v(x), \text{lip } \mathbf{d}(x), \text{lip } m(x) < +\infty$.

We now fix such a point $x \in \Omega$. Let us verify

$$(3.9) \quad \frac{1}{\mathbf{m}(B_r(x))} \int_{B_r(x) \setminus \Omega_{u,v,\beta,x}} \frac{\text{Cub}(x, y)}{r^2} \mathbf{d}\mathbf{m}(y) \rightarrow 0$$

as $r \downarrow 0$. Each term in (3.9) can be written as

$$(3.10) \quad \frac{f(x)}{\mathbf{m}(B_r(x))} \int_{B_r(x) \setminus \Omega_{u,v,\beta,x}} \frac{\mathbf{d}_Y^{\theta_1}(u(x), u(y)) \mathbf{d}_Y^{\theta_2}(v(x), v(y)) (\mathbf{d}(x) - \mathbf{d}(y))^{\theta_3} \mathbf{d}_Y^{\theta_4}(m(x), m(y))}{r^2} \mathbf{d}\mathbf{m}(y)$$

for $\theta_1, \theta_2, \theta_3, \theta_4 \geq 0$ such that $\theta_1 + \theta_2 + \theta_3 + \theta_4 \geq 3$ and at least one of $i = 1, \dots, 4$ has $\theta_i \geq 2$ (see Remark 3.6). We consider the case of $\theta_3 \geq 2$. In this case, the absolute value of (3.10) is bounded from above by

$$(3.11) \quad \begin{aligned} & \frac{|f(x)|}{\mathbf{m}(B_r(x))} \int_{B_r(x) \setminus \Omega_{u,v,\beta,x}} \frac{\mathbf{d}_Y^{\theta_1}(u(x), u(y)) \mathbf{d}_Y^{\theta_2}(v(x), v(y)) |\mathbf{d}(x) - \mathbf{d}(y)|^{\theta_3} \mathbf{d}_Y^{\theta_4}(m(x), m(y))}{r^2} \mathbf{d}\mathbf{m}(y) \\ & \leq |f(x)| r^{\theta_1 + \theta_2 + \theta_3 + \theta_4 - 2} \mathbf{ks}_{2,r}^2[\mathbf{d}, \Omega; u, v, \beta](x) I, \end{aligned}$$

where

$$I := \sup_{y \in B_r(x) \setminus \{x\}} \left(\frac{\mathbf{d}_Y^{\theta_1}(u(x), u(y))}{\mathbf{d}^{\theta_1}(x, y)} \frac{\mathbf{d}_Y^{\theta_2}(v(x), v(y))}{\mathbf{d}^{\theta_2}(x, y)} \frac{|\mathbf{d}(x) - \mathbf{d}(y)|^{\theta_3 - 2}}{\mathbf{d}^{\theta_3 - 2}(x, y)} \frac{\mathbf{d}_Y^{\theta_4}(m(x), m(y))}{\mathbf{d}^{\theta_4}(x, y)} \right).$$

By $\theta_1 + \theta_2 + \theta_3 + \theta_4 - 2 \geq 1$, and by the choices of x , the right hand side of (3.11) tends to 0 as $r \downarrow 0$. For other cases, we can apply the same argument, and conclude (3.9). Thus, Theorem 2.8, Lemma 3.1, (3.9), and letting $r \downarrow 0$ in (3.8) lead to the desired assertion. \square

We next prove the following (cf. [38, Lemma 1.14]):

Lemma 3.8. *Assume that $(X, \mathbf{d}, \mathbf{m})$ is infinitesimally Hilbertian. For $u \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$, $\eta \in \mathbf{KS}^{1,2}(\Omega, [0, 1])$, we define $u_\eta : \Omega \rightarrow \bar{B}_\rho(o)$ by $u_\eta := \mathbf{G}_\eta^{u,o}$. We also define $\mathbf{d} := \mathbf{d}_Y(u, o)$. Then we have $u_\eta \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$, $\mathbf{d} \in \mathbf{KS}^{1,2}(\Omega, \mathbb{R})$, and*

$$e_2^2[u_\eta] \leq \frac{\sin^2(1 - \eta)\mathbf{d}}{\sin^2 \mathbf{d}} (e_2^2[u] - e_2^2[\mathbf{d}]) + e_2^2[(1 - \eta)\mathbf{d}] \quad \mathbf{m}\text{-a.e. on } \Omega.$$

Proof. Let us verify $u_\eta \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$, $\mathbf{d} \in \mathbf{KS}^{1,2}(\Omega, \mathbb{R})$. Similarly to the proof of Lemma 3.7, the assertion for \mathbf{d} can be derived from the triangle inequality. In virtue of (3.5) for $l = \rho$, for all $x, y \in \Omega$,

$$\begin{aligned} \mathbf{d}_Y(u_\eta(x), u_\eta(y)) & \leq c_\rho \mathbf{d}_Y(u(x), u(y)) + |\eta(x) - \eta(y)| \mathbf{d}_Y(u(y), o) \\ & \leq c_\rho \mathbf{d}_Y(u(x), u(y)) + \rho |\eta(x) - \eta(y)| \end{aligned}$$

for some constant $c_\rho > 0$ depending only on ρ . Therefore,

$$\mathbf{ks}_{2,r}^2[u_\eta, \Omega](x) \leq 2c_\rho^2 \mathbf{ks}_{2,r}^2[u, \Omega](x) + 2\rho^2 \mathbf{ks}_{2,r}^2[\eta, \Omega](x).$$

Integrating this estimate over Ω , and letting $r \downarrow 0$, we conclude $u_\eta \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$.

Using Lemma 3.5, for all $x, y \in \Omega$ we have

$$\begin{aligned} \mathbf{d}_Y^2(u_\eta(x), u_\eta(y)) & \leq \frac{\sin^2(1 - \eta(x))\mathbf{d}(x)}{\sin^2 \mathbf{d}(x)} (\mathbf{d}_Y^2(u(x), u(y)) - (\mathbf{d}(y) - \mathbf{d}(x))^2) \\ & \quad + (1 - \eta(x))^2 (\mathbf{d}(y) - \mathbf{d}(x))^2 + \mathbf{d}^2(x) (\eta(y) - \eta(x))^2 \\ & \quad - 2(1 - \eta(x))\mathbf{d}(x) (\eta(y) - \eta(x)) (\mathbf{d}(y) - \mathbf{d}(x)) \\ & \quad + \text{Cub}(\eta(y) - \eta(x), \mathbf{d}_Y(u(x), u(y)), \mathbf{d}(y) - \mathbf{d}(x), \mathbf{d}_Y(u_\eta(x), u_\eta(y))). \end{aligned}$$

Setting $\xi := 1 - \eta$, we possess

(3.12)

$$\begin{aligned}
d_Y^2(u_\eta(x), u_\eta(y)) &\leq \frac{\sin^2 \xi(x) \mathbf{d}(x)}{\sin^2 \mathbf{d}(x)} (\mathbf{d}_Y^2(u(x), u(y)) - (\mathbf{d}(y) - \mathbf{d}(x))^2) \\
&\quad + \xi^2(x) (\mathbf{d}(y) - \mathbf{d}(x))^2 + \mathbf{d}^2(x) (\xi(y) - \xi(x))^2 \\
&\quad + 2\xi(x) \mathbf{d}(x) (\xi(y) - \xi(x)) (\mathbf{d}(y) - \mathbf{d}(x)) \\
&\quad + \text{Cub}(\xi(x) - \xi(y), \mathbf{d}_Y(u(x), u(y)), \mathbf{d}(y) - \mathbf{d}(x), \mathbf{d}_Y(u_\eta(x), u_\eta(y))) \\
&= \frac{\sin^2 \xi(x) \mathbf{d}(x)}{\sin^2 \mathbf{d}(x)} (\mathbf{d}_Y^2(u(x), u(y)) - (\mathbf{d}(y) - \mathbf{d}(x))^2) \\
&\quad + \xi^2(x) (\mathbf{d}(y) - \mathbf{d}(x))^2 + \mathbf{d}^2(x) (\xi(y) - \xi(x))^2 \\
&\quad + \frac{\xi(x) \mathbf{d}(x)}{2} [\{(\xi + \mathbf{d})(x) - (\xi + \mathbf{d})(y)\}^2 - \{(\xi - \mathbf{d})(x) - (\xi - \mathbf{d})(y)\}^2] \\
&\quad + \text{Cub}(\xi(x) - \xi(y), \mathbf{d}_Y(u(x), u(y)), \mathbf{d}(y) - \mathbf{d}(x), \mathbf{d}_Y(u_\eta(x), u_\eta(y))).
\end{aligned}$$

Dividing it by r^2 , and integrating in y over $B_r(x)$ imply

$$\begin{aligned}
\mathbf{ks}_{2,r}^2[u_\eta, \Omega](x) &\leq \frac{\sin^2 \xi(x) \mathbf{d}(x)}{\sin^2 \mathbf{d}(x)} (\mathbf{ks}_{2,r}^2[u, \Omega](x) - \mathbf{ks}_{2,r}^2[\mathbf{d}, \Omega](x)) \\
&\quad + \xi^2(x) \mathbf{ks}_{2,r}^2[\mathbf{d}, \Omega](x) + \mathbf{d}^2(x) \mathbf{ks}_{2,r}^2[\xi, \Omega](x) \\
&\quad + \frac{\xi(x) \mathbf{d}(x)}{2} (\mathbf{ks}_{2,r}^2[\xi + \mathbf{d}, \Omega](x) - \mathbf{ks}_{2,r}^2[\xi - \mathbf{d}, \Omega](x)) + \int_{B_r(x)} \frac{\text{Cub}(x, y)}{r^2} \mathbf{d}\mathbf{m}(y),
\end{aligned}$$

where $\text{Cub}(x, y)$ denotes the cubic terms in (3.12). In the same manner as in the proof of Lemma 3.7, we see

$$\int_{B_r(x)} \frac{\text{Cub}(x, y)}{r^2} \mathbf{d}\mathbf{m}(y) \rightarrow 0 \quad \mathbf{m}\text{-a.e. } x \in \Omega$$

as $r \downarrow 0$. By virtue of Theorem 2.8, we obtain

$$e_2^2[u_\eta] \leq \frac{\sin^2 \xi \mathbf{d}}{\sin^2 \mathbf{d}} (e_2^2[u] - e_2^2[\mathbf{d}]) + \xi^2 e_2^2[\mathbf{d}] + \mathbf{d}^2 e_2^2[\xi] + \frac{\xi \mathbf{d}}{2} (e_2^2[\xi + \mathbf{d}] - e_2^2[\xi - \mathbf{d}]) \quad \mathbf{m}\text{-a.e. on } \Omega.$$

Now, it suffices to prove that

$$\xi^2 e_2^2[\mathbf{d}] + \mathbf{d}^2 e_2^2[\xi] + \frac{\xi \mathbf{d}}{2} (e_2^2[\xi + \mathbf{d}] - e_2^2[\xi - \mathbf{d}]) = e_2^2[\xi \mathbf{d}] \quad \mathbf{m}\text{-a.e. on } \Omega.$$

Moreover, due to Proposition 2.9, for any $u \in \mathbf{KS}^{1,2}(\Omega, \mathbb{R})$, we possess $|Du| = c_d e_2[u]$ \mathbf{m} -a.e. on Ω , and hence it is enough to show that

$$(3.13) \quad \xi^2 |D\mathbf{d}|^2 + \mathbf{d}^2 |D\xi|^2 + \frac{\xi \mathbf{d}}{2} (|D(\xi + \mathbf{d})|^2 - |D(\xi - \mathbf{d})|^2) = |D(\xi \mathbf{d})|^2 \quad \mathbf{m}\text{-a.e. on } \Omega.$$

Since $(X, \mathbf{d}, \mathbf{m})$ is infinitesimally Hilbertian, Theorem 2.2 tells us that

$$(3.14) \quad |D(\xi \mathbf{d})|^2 = \langle \nabla(\xi \mathbf{d}), \nabla(\xi \mathbf{d}) \rangle = \xi^2 |D\mathbf{d}|^2 + \mathbf{d}^2 |\nabla \xi|^2 + 2\xi \mathbf{d} \langle \nabla \xi, \nabla \mathbf{d} \rangle \quad \mathbf{m}\text{-a.e. on } \Omega,$$

where we used the Leibniz rule. Furthermore, by the linearity of the gradient,

$$(3.15) \quad |D(\xi + \mathbf{d})|^2 - |D(\xi - \mathbf{d})|^2 = \langle \nabla(\xi + \mathbf{d}), \nabla(\xi + \mathbf{d}) \rangle - \langle \nabla(\xi - \mathbf{d}), \nabla(\xi - \mathbf{d}) \rangle = 4 \langle \nabla \xi, \nabla \mathbf{d} \rangle \quad \mathbf{m}\text{-a.e. on } \Omega.$$

Combining (3.14) and (3.15) yields

$$|D(\xi \mathbf{d})|^2 = \xi^2 |D\mathbf{d}|^2 + \mathbf{d}^2 |D\xi|^2 + \frac{\xi \mathbf{d}}{2} (|D(\xi + \mathbf{d})|^2 - |D(\xi - \mathbf{d})|^2) \quad \mathbf{m}\text{-a.e. on } \Omega,$$

and this is nothing but (3.13). We complete the proof. \square

We obtain the following convexity of energy (cf. [23, Proposition 1.15]):

Proposition 3.9. *Assume that $(X, \mathbf{d}, \mathbf{m})$ is infinitesimally Hilbertian. For $u, v \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$, we define $m := \mathbf{G}_{1/2}^{u,v} \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$ and $\mathbf{d} := \mathbf{d}_Y(u, v), \mathfrak{d} := \mathbf{d}_Y(m, o) \in \mathbf{KS}^{1,2}(\Omega, \mathbb{R})$. Let $\eta \in \mathbf{KS}^{1,2}(\Omega, [0, 1])$ be a function determined by*

$$(3.16) \quad \frac{\sin(1-\eta)\mathfrak{d}}{\sin \mathfrak{d}} = \cos \frac{\mathbf{d}}{2},$$

and define $m_\eta := \mathbf{G}_\eta^{m,o} \in \mathbf{KS}^{1,2}(\Omega, \bar{B}_\rho(o))$. Then we have

$$\mathbf{E}_2^\Omega(m_\eta) + \cos^8 \rho \mathbf{E}_2^\Omega \left(\frac{\tan \frac{\mathbf{d}}{2}}{\cos \mathfrak{d}} \right) \leq \frac{1}{2} \mathbf{E}_2^\Omega(u) + \frac{1}{2} \mathbf{E}_2^\Omega(v).$$

Proof. Combining Lemmas 3.7 and 3.8, we see

$$(3.17) \quad \begin{aligned} e_2^2[m_\eta] &\leq \frac{\sin^2(1-\eta)\mathfrak{d}}{\sin^2 \mathfrak{d}} (e_2^2[m] - e_2^2[\mathfrak{d}]) + e_2^2[(1-\eta)\mathfrak{d}] \\ &= \cos^2 \frac{\mathbf{d}}{2} (e_2^2[m] - e_2^2[\mathfrak{d}]) + e_2^2[(1-\eta)\mathfrak{d}] \\ &\leq \frac{1}{2} (e_2^2[u] + e_2^2[v]) - \frac{1}{4} e_2^2[\mathbf{d}] - \cos^2 \frac{\mathbf{d}}{2} e_2^2[\mathfrak{d}] + e_2^2[(1-\eta)\mathfrak{d}] \quad \mathbf{m}\text{-a.e. on } \Omega. \end{aligned}$$

We now observe that the following holds:

$$(3.18) \quad e_2^2[(1-\eta)\mathfrak{d}] + \frac{\cos^4 \frac{\mathbf{d}}{2} \cos^4 \mathfrak{d}}{1 - \cos^2 \frac{\mathbf{d}}{2} \sin^2 \mathfrak{d}} e_2^2 \left[\frac{\tan \frac{\mathbf{d}}{2}}{\cos \mathfrak{d}} \right] = \cos^2 \frac{\mathbf{d}}{2} e_2^2[\mathfrak{d}] + \frac{1}{4} e_2^2[\mathbf{d}] \quad \mathbf{m}\text{-a.e. on } \Omega.$$

In view of Proposition 2.9, this is equivalent to the following:

$$|D((1-\eta)\mathfrak{d})|^2 + \frac{\cos^4 \frac{\mathbf{d}}{2} \cos^4 \mathfrak{d}}{1 - \cos^2 \frac{\mathbf{d}}{2} \sin^2 \mathfrak{d}} \left| D \left(\frac{\tan \frac{\mathbf{d}}{2}}{\cos \mathfrak{d}} \right) \right|^2 = \cos^2 \frac{\mathbf{d}}{2} |D\mathfrak{d}|^2 + \frac{1}{4} |D\mathbf{d}|^2 \quad \mathbf{m}\text{-a.e. on } \Omega.$$

Since $(X, \mathbf{d}, \mathbf{m})$ is infinitesimally Hilbertian, it can be derived from Theorems 2.1, 2.2 (chain rule, Leibniz rule), the definition of η , and a straightforward calculation. Combining (3.17) and (3.18),

$$\begin{aligned} e_2^2[m_\eta] + \cos^8 \rho e_2^2 \left[\frac{\tan \frac{\mathbf{d}}{2}}{\cos \mathfrak{d}} \right] &\leq e_2^2[m_\eta] + \frac{\cos^4 \frac{\mathbf{d}}{2} \cos^4 \mathfrak{d}}{1 - \cos^2 \frac{\mathbf{d}}{2} \sin^2 \mathfrak{d}} e_2^2 \left[\frac{\tan \frac{\mathbf{d}}{2}}{\cos \mathfrak{d}} \right] \\ &\leq \frac{1}{2} (e_2^2[u] + e_2^2[v]) \quad \mathbf{m}\text{-a.e. on } \Omega. \end{aligned}$$

By integrating this inequality over Ω , we obtain the desired one. \square

3.4. Proof of Theorem 1.2.

We are now in a position to prove Theorem 1.2.

Proof of Theorem 1.2. Let $(X, \mathbf{d}, \mathbf{m})$ be an infinitesimally Hilbertian, locally uniformly doubling, strongly rectifiable space satisfying a Poincaré inequality, and let Ω be a bounded open subset of X with $\mathbf{m}(X \setminus \Omega) > 0$. Let $Y_o = (Y, \mathbf{d}_Y, o)$ be a pointed CAT(1) space, and let $\bar{B}_\rho(o)$ be a regular ball. Let $(u_n)_n \subset \mathbf{KS}_\rho^{1,2}(\Omega, \bar{B}_\rho(o))$ stand for a minimizing sequence of $\mathbf{E}_{2,\bar{u}}^\Omega$. By Proposition 2.10, the functional $\mathbf{E}_{2,\bar{u}}^\Omega$ is lower semicontinuous, and hence it is sufficient to show that $(u_n)_n$ is an $L^2(\Omega, \bar{B}_\rho(o))$ -Cauchy sequence.

Set $I := \lim_n \mathbf{E}_{2,\bar{u}}^\Omega(u_n) = \inf \mathbf{E}_{2,\bar{u}}^\Omega$. We define

$$m_{n,m} := \mathbf{G}_{1/2}^{u_n, u_m}, \quad \mathbf{d}_{n,m} := \mathbf{d}_Y(u_n, u_m), \quad \mathfrak{d}_{n,m} := \mathbf{d}_Y(m_{n,m}, o), \quad m_{n,m,\eta} := \mathbf{G}_{\eta_{n,m}}^{m_{n,m}, o},$$

where $\eta_{n,m} \in \mathbf{KS}^{1,2}(\Omega, [0, 1])$ is defined as (3.16). Proposition 3.9 yields

$$\begin{aligned} \cos^8 \rho \mathbf{E}_{2,\bar{u}}^\Omega \left(\frac{\tan \frac{d_{n,m}}{2}}{\cos \mathfrak{d}_{n,m}} \right) &\leq \frac{1}{2} \mathbf{E}_{2,\bar{u}}^\Omega(u_n) + \frac{1}{2} \mathbf{E}_{2,\bar{u}}^\Omega(u_m) - \mathbf{E}_{2,\bar{u}}^\Omega(m_{n,m,\eta}) \\ &\leq \frac{1}{2} \mathbf{E}_{2,\bar{u}}^\Omega(u_n) + \frac{1}{2} \mathbf{E}_{2,\bar{u}}^\Omega(u_m) - I; \end{aligned}$$

in particular,

$$\overline{\lim}_{n,m \rightarrow \infty} \int_\Omega \left| D \left(\frac{\tan \frac{d_{n,m}}{2}}{\cos \mathfrak{d}_{n,m}} \right) \right|^2 \mathrm{d}\mathbf{m} = 0.$$

Furthermore, the Poincaré inequality under the Dirichlet boundary condition (see e.g., [23, Lemma 6.3], [2, Subsection 5.5]) together with Proposition 2.10 leads us to

$$\lim_{n,m \rightarrow \infty} \int_\Omega \left| \frac{\tan \frac{d_{n,m}}{2}}{\cos \mathfrak{d}_{n,m}} \right|^2 \mathrm{d}\mathbf{m} = 0.$$

Thus, $\lim_{n,m \rightarrow \infty} \int_\Omega |\mathfrak{d}_{n,m}|^2 \mathrm{d}\mathbf{m} = 0$, and we finish the proof. \square

Acknowledgements. The author is grateful to Keita Kunikawa for fruitful discussions during this work. The author expresses his gratitude to Shouhei Honda for valuable comments. The author was supported by JSPS KAKENHI (JP21K20315).

REFERENCES

- [1] L. Ambrosio, N. Gigli and G. Savaré, *Metric measure spaces with Riemannian Ricci curvature bounded from below*, Duke Math. J. **163** (2014), no. 7, 1405–1490.
- [2] A. Björn and J. Björn, *Nonlinear potential theory on metric spaces*, EMS Tracts in Mathematics, **17**. European Mathematical Society (EMS), Zürich, 2011.
- [3] D. Burago, Y. Burago and S. Ivanov, *A course in metric geometry*, Graduate Studies in Mathematics, **33**. American Mathematical Society, Providence, RI, 2001.
- [4] C. Breiner, A. Fraser, L.-H. Huang, C. Mese, P. Sargent and Y. Zhang, *Existence of harmonic maps into CAT(1) spaces*, Comm. Anal. Geom. **28** (2020), no. 4, 781–835.
- [5] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 319. Springer-Verlag, Berlin, 1999.
- [6] E. Brué and D. Semola, *Constancy of the dimension for RCD(K, N) spaces via regularity of Lagrangian flows*, Comm. Pure Appl. Math. **73** (2020), no. 6, 1141–1204.
- [7] S. Y. Cheng, *Liouville theorem for harmonic maps*, Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), pp. 147–151, Proc. Sympos. Pure Math., XXXVI, Amer. Math. Soc., Providence, R.I., 1980.
- [8] H. I. Choi, *On the Liouville theorem for harmonic maps*, Proc. Amer. Math. Soc. **85** (1982), no. 1, 91–94.
- [9] G. De Philippis, A. Marchese and F. Rindler, *On a conjecture of Cheeger. Measure theory in non-smooth spaces*, 145–155, Partial Differ. Equ. Meas. Theory, De Gruyter Open, Warsaw, 2017.
- [10] S. Di Marino, N. Gigli, E. Pasqualetto and E. Soultanis, *Infinitesimal Hilbertianity of locally CAT(κ)-spaces*, J. Geom. Anal. **31** (2021), no. 8, 7621–7685.
- [11] J. Eells and B. Fuglede, *Harmonic maps between Riemannian polyhedra*, With a preface by M. Gromov. Cambridge Tracts in Mathematics, **142**. Cambridge University Press, Cambridge, 2001.
- [12] M. Erbar, K. Kuwada and K.-T. Sturm, *On the equivalence of the entropic curvature-dimension condition and Bochner’s inequality on metric measure spaces*, Invent. Math. **201** (2015), no. 3, 993–1071.
- [13] B. Fuglede, *The Dirichlet problem for harmonic maps from Riemannian polyhedra to spaces of upper bounded curvature*, Trans. Amer. Math. Soc. **357** (2005), no. 2, 757–792.
- [14] B. Fuglede, *Harmonic maps from Riemannian polyhedra to geodesic spaces with curvature bounded from above*, Calc. Var. Partial Differential Equations **31** (2008), no. 1, 99–136.
- [15] N. Gigli, *On the differential structure of metric measure spaces and applications*, Mem. Amer. Math. Soc. **236** (2015), no. 1113, vi+91 pp.

- [16] N. Gigli, *Nonsmooth differential geometry—an approach tailored for spaces with Ricci curvature bounded from below*, Mem. Amer. Math. Soc. **251** (2018), no. 1196, v+161 pp.
- [17] N. Gigli, *On the regularity of harmonic maps from $\text{RCD}(K, N)$ to $\text{CAT}(0)$ spaces and related results*, preprint arXiv:2204.04317.
- [18] N. Gigli and F. Nobile, *A differential perspective on gradient flows on $\text{CAT}(k)$ -spaces and applications*, J. Geom. Anal. **31** (2021), no. 12, 11780–11818.
- [19] N. Gigli and E. Pasqualetto, *Lectures on nonsmooth differential geometry*, SISSA Springer Series, 2. Springer, Cham, 2020.
- [20] N. Gigli and E. Pasqualetto, *Behaviour of the reference measure on RCD spaces under charts*, Comm. Anal. Geom. **29** (2021), no. 6, 1391–1414.
- [21] N. Gigli and E. Pasqualetto, *Equivalence of two different notions of tangent bundle on rectifiable metric measure spaces*, preprint arXiv:1611.09645, to appear in Comm. Anal. Geom..
- [22] N. Gigli and A. Tyulenev, *Korevaar-Schoen’s directional energy and Ambrosio’s regular Lagrangian flows*, Math. Z. **298** (2021), no. 3-4, 1221–1261.
- [23] N. Gigli and A. Tyulenev, *Korevaar-Schoen’s energy on strongly rectifiable spaces*, Calc. Var. Partial Differential Equations **60** (2021), no. 6, Paper No. 235, 54 pp.
- [24] G. Gregori, *Sobolev spaces and harmonic maps between singular spaces*, Calc. Var. Partial Differential Equations **7** (1998), no. 1, 1–18.
- [25] M. Gromov and R. Schoen, *Harmonic maps into singular spaces and p -adic superrigidity for lattices in groups of rank one*, Inst. Hautes Études Sci. Publ. Math. No. **76** (1992), 165–246.
- [26] J.-C. Huang and H.-C. Zhang, *Harmonic maps between Alexandrov spaces*, J. Geom. Anal. **27** (2017), no. 2, 1355–1392.
- [27] J. Jost, *Equilibrium maps between metric spaces*, Calc. Var. Partial Differential Equations **2** (1994), no. 2, 173–204.
- [28] J. Jost, *Generalized Dirichlet forms and harmonic maps*, Calc. Var. Partial Differential Equations **5** (1997), no. 1, 1–19.
- [29] M. Kell and A. Mondino, *On the volume measure of non-smooth spaces with Ricci curvature bounded below*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **18** (2018), no. 2, 593–610.
- [30] B. Kirchheim, *Rectifiable metric spaces: local structure and regularity of the Hausdorff measure*, Proc. Amer. Math. Soc. **121** (1994), no. 1, 113–123.
- [31] N. J. Korevaar and R. M. Schoen, *Sobolev spaces and harmonic maps for metric space targets*, Comm. Anal. Geom. **1** (1993), no. 3-4, 561–659.
- [32] K. Kuwae and T. Shioya, *Sobolev and Dirichlet spaces over maps between metric spaces*, J. Reine Angew. Math. **555** (2003), 39–75.
- [33] F. H. Lin, *Analysis on singular spaces*, Collection of papers on geometry, analysis and mathematical physics, 114–126, World Sci. Publ., River Edge, NJ, 1997.
- [34] J. Lott and C. Villani, *Ricci curvature for metric-measure spaces via optimal transport*, Ann. of Math. (2) **169** (2009), no. 3, 903–991.
- [35] A. Mondino and A. Naber, *Structure theory of metric measure spaces with lower Ricci curvature bounds*, J. Eur. Math. Soc. (JEMS) **21** (2019), no. 6, 1809–1854.
- [36] A. Mondino and D. Semola, *Lipschitz continuity and Bochner-Eells-Sampson inequality for harmonic maps from $\text{RCD}(K, N)$ spaces to $\text{CAT}(0)$ spaces*, preprint arXiv:2202.01590.
- [37] T. Rajala, *Local Poincaré inequalities from stable curvature conditions on metric spaces*, Calc. Var. Partial Differential Equations **44** (2012), no. 3-4, 477–494.
- [38] T. Serbinowski, *Harmonic maps into metric spaces with curvature bounded above*, Thesis (Ph.D.)—The University of Utah. 1995. 64 pp.
- [39] K.-T. Sturm, *On the geometry of metric measure spaces. I*, Acta Math. **196** (2006), no. 1, 65–131.
- [40] K.-T. Sturm, *On the geometry of metric measure spaces. II*, Acta Math. **196** (2006), no. 1, 133–177.
- [41] H.-C. Zhang and X.-P. Zhu, *Lipschitz continuity of harmonic maps between Alexandrov spaces*, Invent. Math. **211** (2018), no. 3, 863–934.
- [42] H.-C. Zhang, X. Zhong and X.-P. Zhu, *Quantitative gradient estimates for harmonic maps into singular spaces*, Sci. China Math. **62** (2019), no. 11, 2371–2400.

DEPARTMENT OF MATHEMATICS, SAITAMA UNIVERSITY, 255 SHIMO-OKUBO, SAKURA-KU, SAITAMA-CITY, SAITAMA, 338-8570, JAPAN

Email address: ysakurai@rimath.saitama-u.ac.jp