

Rational quadrilaterals

Ajai Choudhry

Abstract

A quadrilateral is said to be rational if its four sides, the two diagonals and the area are all expressible by rational numbers. The problem of constructing rational quadrilaterals dates back to the seventh century when Brahmagupta gave an elegant solution of the problem. In 1848 Kummer gave a method of generating all rational quadrilaterals. In this paper we present an alternative method of generating all rational quadrilaterals. For rational cyclic quadrilaterals, we obtain a complete parametrization and for noncyclic rational quadrilaterals, we give several parametrizations in terms of quadratic and quartic polynomials. The parametrizations obtained in this paper are simpler than the known parametrizations of rational quadrilaterals. We also describe how further parametrizations of rational quadrilaterals may be obtained.

Mathematics Subject Classification 2020: 11D09

Keywords: Rational quadrilateral; cyclic quadrilateral; Brahmagupta quadrilateral.

1 Introduction

Since ancient times there has been considerable interest in geometric objects such as triangles, quadrilaterals and other polygons such that the lengths of their sides and diagonals, as well as their areas are expressible by rational numbers. This paper is concerned with rational quadrilaterals — a rational quadrilateral being defined as one whose four sides, the two diagonals and the area are given by rational numbers.

In the seventh century Brahmagupta gave an elegant method of constructing rational quadrilaterals. Brahmagupta's method was elaborated further by Bhaskara in the 12th century and by Chasles in the 19th century (as quoted by Dickson [6, pp. 216–217]). In 1848 Kummer [12] gave a method of generating all rational quadrilaterals. The method, however, leads to a cumbersome final condition that requires a parametrized quartic function be

made a perfect square. The complete solution of this final condition is not yet known. In 1921 Dickson [5] presented Kummer's construction in a somewhat simplified manner.

Various related problems concerning rational quadrilaterals have been studied by several mathematicians ([1], [2], [3], [6, pp. 216–221], [9], [10], [13], [16]). Some authors have adopted a relaxed definition of rational quadrilaterals that requires only the four sides and the two diagonals of the quadrilateral to be rational without the additional condition of the area being rational while some others have studied only cyclic quadrilaterals. For instance, Sastri [16, pp. 170–171] has given a parametrization of cyclic quadrilaterals whose sides, diagonals and the area are all expressible by rational numbers. However, regarding the original problem of constructing quadrilaterals with rational sides, diagonals and area, there has been no progress in over a hundred years since Dickson's simplification in 1921 of the method described by Kummer.

In this paper we describe a method, different from that of Kummer, of generating all rational convex quadrilaterals. We first given in Section 2 a necessary and sufficient condition for the existence of rational quadrilaterals. In Section 3 we obtain several examples of rational quadrilaterals whose sides, diagonals and area are given by multivariate polynomials. These parametrizations are more general and simpler than the examples obtained by Kummer's method. In fact, as a special case, we obtain, for all rational cyclic quadrilaterals, a complete parametrization that is simpler than the known parametrizations of such quadrilaterals.

2 A necessary and sufficient condition for rational quadrilaterals

We now prove a theorem that describes a necessary and sufficient condition for the existence of a rational convex quadrilateral.

Theorem 1. *A necessary and sufficient condition for the existence of a rational convex quadrilateral is that there exist rational numbers $a, b, c, d, e, f, x_1, x_2, y_1$ and y_2 satisfying the simultaneous diophantine equations,*

$$x_1^2 + y_1^2 = a^2, \tag{2.1}$$

$$(e - x_1)^2 + y_1^2 = b^2, \tag{2.2}$$

$$(e - x_2)^2 + y_2^2 = c^2, \tag{2.3}$$

$$x_2^2 + y_2^2 = d^2, \tag{2.4}$$

$$(x_1 - x_2)^2 + (y_1 - y_2)^2 = f^2. \tag{2.5}$$

and such that the following rational numbers are all positive: $a, b, c, d, e, f, -y_1y_2, -(y_1 - y_2)(x_1y_2 - x_2y_1)$ and $(y_1 - y_2)(ey_1 - ey_2 + x_1y_2 - x_2y_1)$.




Figure 1: Rational quadrilateral

Proof. Without loss of generality, let $OABC$ be a rational convex quadrilateral with one vertex O at the origin of the x - y plane and with its diagonal OB along the x -axis as shown in Figure 1. Let the lengths of the sides OA, AB, BC and OC of the quadrilateral be a, b, c and d respectively, and let the lengths of the diagonals OB and AC be e and f respectively. Let D and E be the feet of the perpendiculars to the diagonal OB drawn from the vertices A and C respectively. Let the coordinates of the vertices A and C be (x_1, y_1) and (x_2, y_2) respectively. It readily follows from the theorem of Pythagoras that the relations (2.1)–(2.5) must be satisfied.

Since $OABC$ is assumed to be a rational quadrilateral, a, b, c, d, e and f are rational numbers and the area A of the quadrilateral is rational. We will now prove that the four numbers $x_i, y_i, i = 1, 2$, are all rational. Since all the three sides of the two triangles OAB and OCB are rational, the cosines of the angles AOB and COB are rational, and hence the lengths $OA \cos AOB$ and $OC \cos COB$, that is, x_1 and x_2 are rational.

Further, it follows from Heron's formula for the area of a triangle that the areas of the two triangles OAB and OCB may be written as $\sqrt{r_1}$ and $\sqrt{r_2}$ where r_1 and r_2 are rational numbers. Since, by definition, the area A of the quadrilateral $OABC$ is rational, it follows that $\sqrt{r_1} + \sqrt{r_2} = A$, or, $\sqrt{r_1} = A - \sqrt{r_2}$, or, $r_1 = A + r_2 - 2A\sqrt{r_2}$, and hence $\sqrt{r_2}$ is a rational number. Similarly, $\sqrt{r_1}$ is also a rational number. Thus, the areas of the two triangles OAB and OCB are rational. Since we may write these areas as $OB \times AD/2$ and $OB \times CE/2$ respectively, and OB is of rational length, it follows that the lengths of the line segments AD and CE are rational, that is, both y_1 and y_2 are rational numbers.

We will now prove that all the rational numbers mentioned towards the end of the theorem are positive. We first note that a, b, c, d, e, f , being the lengths of the sides and diagonals of the quadrilateral, are necessarily positive rational numbers. Further, since $OABC$ is a convex quadrilateral, the diagonal OB lies wholly within the quadrilateral, and hence the vertices A and C must be on opposite sides of the diagonal OB , and hence the ordinates y_1 and y_2 must be of opposite signs, and thus $-y_1y_2$ has to be positive. Finally, since $OABC$ is a convex quadrilateral, the point of intersection M of the two diagonals must strictly lie somewhere between the two vertices O and B , that is, the abscissa of the point M must be between 0 and e . The coordinates of the point M are readily worked out to be $(-(x_1y_2 - x_2y_1)/(y_1 - y_2), 0)$, hence we must have

$$0 < -(x_1y_2 - x_2y_1)/(y_1 - y_2) < e.$$

It now readily follows that the numbers $-(y_1 - y_2)(x_1y_2 - x_2y_1)$ and $(y_1 - y_2)(ey_1 - ey_2 + x_1y_2 - x_2y_1)$ are both positive.

We have thus proved that if there exists a rational convex quadrilateral, then there exist rational numbers $a, b, c, d, e, f, x_1, x_2, y_1$ and y_2 satisfying the relations (2.1)–(2.5) and such that the numbers $a, b, c, d, e, f, -y_1y_2, -(y_1 - y_2)(x_1y_2 - x_2y_1)$ and $(y_1 - y_2)(ey_1 - ey_2 + x_1y_2 - x_2y_1)$ are all positive. We have thus shown that the condition stated in the theorem is necessary.

Next let there exist rational numbers $a, b, c, d, e, f, x_1, x_2, y_1$ and y_2 satisfying the conditions of the theorem. We will construct a quadrilateral $OABC$ with vertices O, A, B and C at the points $(0, 0), (x_1, y_1), (e, 0)$ and (x_2, y_2) respectively as shown in Figure 1. In view of the relations (2.1)–(2.5), it is clear that the lengths of the four sides of the quadrilateral are a, b, c , and d respectively, and the two diagonals OB and AC have lengths e and f respectively, and hence these lengths are all rational.

Further, the line segments OB, AD and CE have rational lengths, hence both the triangles OAB and OCB have rational areas, and thus the area of the quadrilateral $OABC$ is rational. Finally it is readily seen that the conditions that the numbers $-y_1y_2, -(y_1 - y_2)(x_1y_2 - x_2y_1)$ and $(y_1 - y_2)(ey_1 - ey_2 + x_1y_2 - x_2y_1)$ are all positive, ensure that the quadrilateral is convex. Thus, the condition stated in the theorem is sufficient for the existence of a rational convex quadrilateral. This proves the theorem. \square

It is readily seen that if

$$(a, b, c, d, e, f, x_1, x_2, y_1, y_2) = (a', b', c', d', e', f', x'_1, x'_2, y'_1, y'_2)$$

is a solution of the simultaneous diophantine Eqs. (2.1)–(2.5), then $(\varepsilon_1a', \varepsilon_2b', \varepsilon_3c', \varepsilon_4d', \varepsilon_5e', \varepsilon_6f', \varepsilon_5x'_1, \varepsilon_5x'_2, \varepsilon_7y'_1, \varepsilon_7y'_2)$ is also a solution where the value of each $\varepsilon_i, i = 1, \dots, 7$, is taken either as $+1$ or -1 . Accordingly, if we obtain

a numerical solution of Eqs. (2.1)–(2.5) such that all the conditions of a rational convex quadrilateral are not satisfied, it is sometimes, though not always, possible to choose appropriate values of ε_i and obtain a solution that satisfies all the conditions of a convex quadrilateral.

3 Parametrizations of rational quadrilaterals

We will now solve the simultaneous diophantine Eqs. (2.1)–(2.5). It is not difficult to obtain the complete solution of any four of the five Eqs. (2.1)–(2.5), and in each case the final equation reduces to a parameterized quartic function to be made a perfect square. While rational solutions of this final condition can be obtained, these are generally quite cumbersome. This situation is similar to the one that arises on following Kummer’s method.

In the next section we will solve the four Eqs. (2.1)–(2.4) in a manner such that the final condition arising from Eq. (2.5) has simple solutions.

3.1 Quadrilaterals with four sides, one diagonal and the area expressible by rational numbers

We will now obtain the complete solution of the four simultaneous diophantine Eqs. (2.1)–(2.4). We will repeatedly make use of the fact that the complete solution of the diophantine equation $X_1X_2 = Y_1Y_2$ is given by $X_1 = gm, X_2 = hn, Y_1 = gn, Y_2 = hm$, where g, h, m and n are arbitrary parameters (see [15, p. 69]). The complete solution of Eqs. (2.1)–(2.4) will also yield a parametrization of all quadrilaterals $OABC$ in which the four sides, the diagonal OB and the area of the quadrilateral $OABC$ are rational.

Taking the difference of the two equations, (2.1) and (2.2), we get $-e(e - 2x_1) = (a+b)(a-b)$, hence we may write $-e = 2u_1v_1, e - 2x_1 = 2u_2v_2, a+b = 2u_1v_2, a - b = 2u_2v_1$, when we get the solution,

$$a = u_1v_2 + u_2v_1, \quad b = u_1v_2 - u_2v_1, \quad e = -2u_1v_1, \quad x_1 = -u_1v_1 - u_2v_2, \quad (3.1)$$

where u_1, u_2, v_1, v_2 are arbitrary parameters.

Similarly, the complete solution of the difference of the two equations (2.3) and (2.4) is given by

$$d = u_3v_4 + u_4v_3, \quad c = u_3v_4 - u_4v_3, \quad e = -2u_3v_3, \quad x_2 = -u_3v_3 - u_4v_4, \quad (3.2)$$

where u_3, u_4, v_3, v_4 are arbitrary parameters.

The two solutions (3.1) and (3.2) will be consistent if the values of e given by the two solutions are the same, that is, $u_1v_1 = u_3v_3$, and hence we take

$$u_1 = m_1n_1, \quad v_1 = m_2n_2, \quad u_3 = m_1n_2, \quad v_3 = m_2n_1, \quad (3.3)$$

where m_1, m_2, n_1 and n_2 are arbitrary nonzero rational parameters. On substituting the values of $u_i, v_i, i = 1, 2$ given by (3.3) in (3.1) and (3.2), we get the following relations,

$$\begin{aligned}
a &= m_1 n_1 v_2 + m_2 n_2 u_2, \\
b &= m_1 n_1 v_2 - m_2 n_2 u_2, \\
c &= m_1 n_2 v_4 - m_2 n_1 u_4, \\
d &= m_1 n_2 v_4 + m_2 n_1 u_4, \\
e &= -2m_1 m_2 n_1 n_2, \\
x_1 &= -m_1 m_2 n_1 n_2 - u_2 v_2, \\
x_2 &= -m_1 m_2 n_1 n_2 - u_4 v_4,
\end{aligned} \tag{3.4}$$

where $m_1, m_2, n_1, n_2, u_2, u_4, v_2$ and v_4 are arbitrary parameters.

On substituting the values of a and x_1 given by (3.4) in (2.1), we get $y_1^2 = -(m_2 n_2 - v_2)(m_2 n_2 + v_2)(m_1 n_1 - u_2)(m_1 n_1 + u_2)$. Thus the right-hand side of this last equation must be a nonzero perfect square, and this is possible if and only if there exists a nonzero rational number h_1 such that $h_1^2(m_1 n_1 - u_2)(m_1 n_1 + u_2) = -(m_2 n_2 - v_2)(m_2 n_2 + v_2)$, hence there exist nonzero rational parameters $p_i, q_i, i = 1, 2$, such that

$$\begin{aligned}
h_1(m_1 n_1 - u_2) &= 2p_1 q_2, & h_1(m_1 n_1 + u_2) &= 2p_2 q_1, \\
-(m_2 n_2 - v_2) &= 2p_1 q_1, & m_2 n_2 + v_2 &= 2p_2 q_2.
\end{aligned} \tag{3.5}$$

On solving Eqs. (3.5) for m_1, m_2, u_2 and v_2 , we get,

$$\begin{aligned}
m_1 &= (p_1 q_2 + p_2 q_1)/(h_1 n_1), & m_2 &= -(p_1 q_1 - p_2 q_2)/n_2, \\
u_2 &= -(p_1 q_2 - p_2 q_1)/h_1, & v_2 &= p_1 q_1 + p_2 q_2.
\end{aligned} \tag{3.6}$$

Similarly, on substituting the values of d and x_2 given by (3.4) in (2.4), we get $y_2^2 = -(m_2 n_1 - v_4)(m_2 n_1 + v_4)(m_1 n_2 - u_4)(m_1 n_2 + u_4)$, and proceeding as in the previous paragraph, we get the solution,

$$\begin{aligned}
m_1 &= (r_1 s_2 + r_2 s_1)/(h_2 n_2), & m_2 &= -(r_1 s_1 - r_2 s_2)/n_1, \\
u_4 &= -(r_1 s_2 - r_2 s_1)/h_2, & v_4 &= r_1 s_1 + r_2 s_2,
\end{aligned} \tag{3.7}$$

where h_2, r_1, r_2, s_1, s_2 are arbitrary nonzero rational parameters.

For the solutions (3.6) and (3.7) to be consistent, we must impose the conditions that the respective values of m_1 and m_2 given by these two solutions coincide. These conditions are satisfied if and only if we choose h_1, h_2, n_1 and n_2 as follows:

$$\begin{aligned}
h_1 &= (p_1 q_2 + p_2 q_1)(p_1 q_1 - p_2 q_2), & h_2 &= (r_1 s_2 + r_2 s_1)(r_1 s_1 - r_2 s_2), \\
n_1 &= r_1 s_1 - r_2 s_2, & n_2 &= p_1 q_1 - p_2 q_2.
\end{aligned} \tag{3.8}$$

Using the values of $h_i, n_i, i = 1, 2$, given by (3.8) we can now find the values of m_1, m_2, u_2, v_2, u_4 and v_4 from the relations (3.6) and (3.7), and we can solve Eqs. (2.1) and (2.4) to obtain rational values of y_1, y_2 , and further, using the relations (3.4), we get, on appropriate scaling, the following solution of the simultaneous diophantine Eqs. (2.1)–(2.4):

$$\begin{aligned}
a &= q_1 q_2 (r_1 s_2 + r_2 s_1) (r_1 s_1 - r_2 s_2) (p_1^2 + p_2^2), \\
b &= p_1 p_2 (r_1 s_2 + r_2 s_1) (r_1 s_1 - r_2 s_2) (q_1^2 + q_2^2), \\
c &= r_1 r_2 (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) (s_1^2 + s_2^2), \\
d &= s_1 s_2 (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) (r_1^2 + r_2^2), \\
e &= (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) (r_1 s_2 + r_2 s_1) (r_1 s_1 - r_2 s_2), \quad (3.9) \\
x_1 &= q_1 q_2 (r_1 s_2 + r_2 s_1) (r_1 s_1 - r_2 s_2) (p_1 - p_2) (p_1 + p_2), \\
x_2 &= s_1 s_2 (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) (r_1 - r_2) (r_1 + r_2), \\
y_1 &= 2p_1 p_2 q_1 q_2 (r_1 s_2 + r_2 s_1) (r_1 s_1 - r_2 s_2), \\
y_2 &= -2r_1 r_2 s_1 s_2 (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2).
\end{aligned}$$

where $p_i, q_i, r_i, s_i, i = 1, 2$, are arbitrary rational parameters.

While solving the simultaneous diophantine equations (2.1)–(2.4), we had to solve several intermediate equations and for each of these equations, we obtained the complete solution. Hence, the formulae (3.9) give the complete solution of the simultaneous equations (2.1)–(2.4).

It follows that the formulae (3.9) generate all convex quadrilaterals whose four sides a, b, c, d , one diagonal and the area are given by rational numbers. We must, of course, choose the parameters such that the rational numbers mentioned towards the end of Theorem 1 are all positive. The area of the quadrilateral defined by (3.9) is given by

$$\begin{aligned}
A &= (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) (r_1 s_2 + r_2 s_1) (r_1 s_1 - r_2 s_2) \\
&\quad \times [p_1^2 q_1 q_2 r_1 r_2 s_1 s_2 + \{q_1^2 r_1 r_2 s_1 s_2 + (r_1 s_2 + r_2 s_1) (r_1 s_1 - r_2 s_2) q_1 q_2 \\
&\quad \quad \quad - q_2^2 r_1 r_2 s_1 s_2\} p_1 p_2 - p_2^2 q_1 q_2 r_1 r_2 s_1 s_2]. \quad (3.10)
\end{aligned}$$

3.2 Rational quadrilaterals

We will now obtain parametrizations of rational quadrilaterals by obtaining parametric solutions of the simultaneous diophantine Eqs. (2.1)–(2.5).

On substituting the values of x_1, x_2, y_1 and y_2 given by (3.9) in Eq. (2.5),

we get the condition,

$$\begin{aligned}
& q_1^2 q_2^2 r_1^2 r_2^2 (p_1^2 + p_2^2)^2 s_1^4 - 2p_1 p_2 q_1 q_2 r_1 r_2 \{ (q_1 r_1 + q_1 r_2 + q_2 r_1 - q_2 r_2) p_1 + (q_1 r_1 - q_1 r_2 \\
& - q_2 r_1 - q_2 r_2) p_2 \} \{ (q_1 r_1 - q_1 r_2 - q_2 r_1 - q_2 r_2) p_1 - (q_1 r_1 + q_1 r_2 + q_2 r_1 - q_2 r_2) p_2 \} s_1^3 s_2 \\
& + \{ 2p_1^4 q_1^2 q_2^2 r_1^2 r_2^2 + 8q_1 q_2 r_1 r_2 (q_1 r_2 + q_2 r_1) (q_1 r_1 - q_2 r_2) p_1^3 p_2 + (q_1^4 r_1^4 + 2q_1^4 r_1^2 r_2^2 + q_1^4 r_2^4 \\
& + 8q_1^3 q_2 r_1^3 r_2 - 8q_1^3 q_2 r_1 r_2^3 + 2q_1^2 q_2^2 r_1^4 - 24q_1^2 q_2^2 r_1^2 r_2^2 + 2q_1^2 q_2^2 r_2^4 - 8q_1 q_2^3 r_1^3 r_2 \\
& + 8q_1 q_2^3 r_1 r_2^3 + q_2^4 r_1^4 + 2q_2^4 r_1^2 r_2^2 + q_2^4 r_2^4) p_1^2 p_2^2 - 8q_1 q_2 r_1 r_2 (q_1 r_2 + q_2 r_1) (q_1 r_1 - q_2 r_2) p_1 p_2^3 \\
& + 2p_2^4 q_1^2 q_2^2 r_1^2 r_2^2 \} s_1^2 s_2^2 + 2p_1 p_2 q_1 q_2 r_1 r_2 \{ (q_1 r_1 + q_1 r_2 + q_2 r_1 - q_2 r_2) p_1 + (q_1 r_1 - q_1 r_2 \\
& - q_2 r_1 - q_2 r_2) p_2 \} \{ (q_1 r_1 - q_1 r_2 - q_2 r_1 - q_2 r_2) p_1 - (q_1 r_1 + q_1 r_2 + q_2 r_1 - q_2 r_2) p_2 \} s_1 s_2^3 \\
& + q_1^2 q_2^2 r_1^2 r_2^2 (p_1^2 + p_2^2)^2 s_2^4 = f^2. \quad (3.11)
\end{aligned}$$

We need to solve Eq. (3.11) such that the rational numbers mentioned towards the end of Theorem 1 are positive. Thus the parameters $p_i, q_i, r_i, s_i, i = 1, 2$, must all be nonzero and simple solutions of Eq. (3.11) such as $p_1 = p_2 q_2 / q_1$ or $p_1 = -p_2 q_1 / q_2$ or $r_1 = r_2 s_2 / s_1$ or $r_1 = -r_2 s_1 / s_2$ are also ruled out. We will give four simple solutions of (3.11) that satisfy the conditions of Theorem 1, and thus obtain four parametrizations of rational quadrilaterals. We will also show how infinitely many parametrizations of rational quadrilaterals may be obtained.

3.2.1 Rational cyclic quadrilaterals

We first give a solution of Eq. (3.11) that generates all rational cyclic quadrilaterals. It is readily verified that if we take

$$\begin{aligned}
s_1 &= p_1(q_1 r_2 + q_2 r_1) + p_2(q_1 r_1 - q_2 r_2), \\
s_2 &= p_1(q_1 r_1 - q_2 r_2) - p_2(q_1 r_2 + q_2 r_1),
\end{aligned} \quad (3.12)$$

the left-hand side of Eq. (3.11) becomes a perfect square. This yields a rational quadrilateral whose sides a, b, c, d , the diagonals e, f and the area A may be written, after appropriate scaling, as follows:

$$\begin{aligned}
a &= (p_1^2 + p_2^2)(r_1^2 + r_2^2)q_1 q_2, \\
b &= (q_1^2 + q_2^2)(r_1^2 + r_2^2)p_1 p_2, \\
c &= (q_1^2 + q_2^2)(p_1^2 + p_2^2)r_1 r_2, \\
d &= (p_1 q_1 r_2 + p_1 q_2 r_1 + p_2 q_1 r_1 - p_2 q_2 r_2) \\
&\quad \times (p_1 q_1 r_1 - p_1 q_2 r_2 - p_2 q_1 r_2 - p_2 q_2 r_1), \\
e &= (r_1^2 + r_2^2)(p_1 q_2 + p_2 q_1)(p_1 q_1 - p_2 q_2), \\
f &= (q_1^2 + q_2^2)(p_1 r_2 + p_2 r_1)(p_1 r_1 - p_2 r_2), \\
A &= (p_1 q_1 - p_2 q_2)(p_1 q_2 + p_2 q_1)(p_1 r_1 - p_2 r_2) \\
&\quad \times (p_1 r_2 + p_2 r_1)(q_1 r_1 - q_2 r_2)(q_1 r_2 + q_2 r_1).
\end{aligned} \quad (3.13)$$

Further, the related values of y_1, y_2 , obtained from (3.9) are as follows:

$$\begin{aligned}
x_1 &= (p_1 + p_2)(p_1 - p_2)(r_1^2 + r_2^2)q_1q_2, \\
x_2 &= (r_1 - r_2)(r_1 + r_2)(p_1q_1r_2 + p_1q_2r_1 + p_2q_1r_1 - p_2q_2r_2) \\
&\quad \times (p_1q_1r_1 - p_1q_2r_2 - p_2q_1r_2 - p_2q_2r_1)/(r_1^2 + r_2^2), \\
y_1 &= 2(r_1^2 + r_2^2)p_1p_2q_1q_2, \\
y_2 &= -2r_1r_2(p_1q_1r_2 + p_1q_2r_1 + p_2q_1r_1 - p_2q_2r_2) \\
&\quad \times (p_1q_1r_1 - p_1q_2r_2 - p_2q_1r_2 - p_2q_2r_1)/(r_1^2 + r_2^2),
\end{aligned} \tag{3.14}$$

If we assign positive rational values to all the parameters $p_i, q_i, r_i, i = 1, 2$, such that $q_1 > q_2r_2/r_1$ and $p_1 > p_2(q_1r_2 + q_2r_1)/(q_1r_1 - q_2r_2)$, it readily follows that all conditions mentioned in Theorem 1 are satisfied so that the formulae (3.13) define a rational convex quadrilateral. As a numerical example, when we take $(p_1, p_2, q_1, q_2, r_1, r_2) = (4, 1, 3, 1, 2, 1)$, we get a rational convex quadrilateral whose sides, diagonals and the area, after appropriate scaling, are given by $(a, b, c, d, e, f, A) = (51, 40, 68, 75, 77, 84, 3234)$.

We will now show that the formulae (3.13) give the sides, diagonals and the area of a cyclic quadrilateral. In fact, it is useful to recall here that if a, b, c, d are the consecutive sides of a cyclic quadrilateral, the area K and the lengths e and f of the two diagonals of the quadrilateral are given by the following formulae proved by Brahmagupta [7, p. 187]:

$$K = \sqrt{(s-a)(s-b)(s-c)(s-d)}, \tag{3.15}$$

and

$$\begin{aligned}
e &= \sqrt{(ab+cd)(ac+bd)/(ad+bc)}, \\
f &= \sqrt{(ac+bd)(ad+bc)/(ab+cd)},
\end{aligned} \tag{3.16}$$

where s is the semi-perimeter, that is,

$$s = (a + b + c + d)/2. \tag{3.17}$$

Further, the circumradius R of the cyclic quadrilateral is given by the following formula of Paramesvara [8]:

$$R = \frac{1}{4} \sqrt{\frac{(ab+cd)(ac+bd)(ad+bc)}{(s-a)(s-b)(s-c)(s-d)}}. \tag{3.18}$$

We also note that the area Δ of an arbitrary convex quadrilateral having sides a, b, c, d , and the sum of any one pair of opposite angles equal to $2u$, is given by the following formula [11, p. 82]:

$$\Delta^2 = (s-a)(s-b)(s-c)(s-d) - abcd \cos^2 u, \tag{3.19}$$

where s is the semiperimeter.

It follows from the formulae (3.15) and (3.19) that a quadrilateral with sides a, b, c, d , is cyclic if and only if its area is given by the formula (3.15). It is readily verified that the area A of the quadrilateral defined by (3.13) is exactly the area of a cyclic quadrilateral whose sides a, b, c, d , are given by (3.13). Further, the lengths of the two diagonals, given by the formulae (3.16), coincide with the lengths e and f of the two diagonals given by (3.13). Thus, the quadrilateral defined by (3.13) is a cyclic quadrilateral. We also note that the circumradius of the quadrilateral defined by (3.13) is

$$R = (p_1^2 + p_2^2)(q_1^2 + q_2^2)(r_1^2 + r_2^2)/4. \quad (3.20)$$

It is pertinent to consider at this stage whether there are any other values of the parameters $p_i, q_i, r_i, s_i, i = 1, 2$, such the quadrilateral defined by (3.9) and (3.10) becomes a cyclic quadrilateral. The condition that the area A of this quadrilateral becomes equal to the area of a cyclic quadrilateral with sides a, b, c, d , may be written as follows:

$$\begin{aligned} & p_1 p_2 q_1 q_2 r_1 r_2 s_1 s_2 (r_1 s_2 + r_2 s_1)^2 (r_1 s_1 - r_2 s_2)^2 (p_1 q_2 + p_2 q_1)^2 \\ & (p_1 q_1 - p_2 q_2)^2 \{ (p_1 q_1 r_1 - p_1 q_2 r_2 - p_2 q_1 r_2 - p_2 q_2 r_1) s_1 \\ & - (p_1 q_1 r_2 + p_1 q_2 r_1 + p_2 q_1 r_1 - p_2 q_2 r_2) s_2 \}^2 = 0. \quad (3.21) \end{aligned}$$

Equating to 0 any factor on the left-hand side of (3.21), except the last, does not yield a quadrilateral since at least one of the four sides a, b, c, d , becomes 0, and the last factor, when equated to 0, yields the cyclic quadrilateral given by (3.13).

Even if we consider the possibility that for a certain choice of parameters, the formulae (3.9) may yield negative values for one or more of the numbers a, b, c, d , and we obtain a rational convex quadrilateral by changing the signs of a, b, c, d , as necessary, when we impose the condition that the area of the quadrilateral becomes equal to the area of a cyclic quadrilateral, and proceed as above, we eventually get the same parametrization of a quadrilateral as is given by (3.13). It follows that the formulae (3.13) give the complete parametrization of all cyclic quadrilaterals.

We note that Euler (as quoted by Dickson [6, p. 221]) had given complicated expressions for the sides and diagonals of a cyclic quadrilateral to be rational but the area is not rational. As mentioned in the Introduction, Sastry [16, pp. 170–171] has given a parametrization of cyclic quadrilaterals but the formulae (3.13) are simpler and more symmetric as compared to Sastry's formulae.

3.2.2 A parametrization of noncyclic convex quadrilaterals

A second solution of Eq. (3.11) is obtained if we take

$$s_1 = r_1 r_2 \{2p_1 q_1 q_2 + (q_1^2 - q_2^2) p_2\}, \quad s_2 = q_1 q_2 \{(r_1^2 - r_2^2) p_1 - 2p_2 r_1 r_2\}.$$

This yields a rational quadrilateral whose sides a, b, c, d , the diagonals e, f and the area A are as follows:

$$\begin{aligned} a &= q_1 q_2 (p_1^2 + p_2^2) \{(r_1^2 + r_2^2) p_1 q_1 q_2 + (q_1^2 r_2 - 2q_1 q_2 r_1 - q_2^2 r_2) p_2 r_2\} \\ &\quad \times \{(r_1^2 + r_2^2) p_1 q_1 q_2 + (q_1^2 r_1 + 2q_1 q_2 r_2 - q_2^2 r_1) p_2 r_1\}, \\ b &= p_1 p_2 (q_1^2 + q_2^2) \{(r_1^2 + r_2^2) p_1 q_1 q_2 + (q_1^2 r_2 - 2q_1 q_2 r_1 - q_2^2 r_2) p_2 r_2\} \\ &\quad \times \{(r_1^2 + r_2^2) p_1 q_1 q_2 + (q_1^2 r_1 + 2q_1 q_2 r_2 - q_2^2 r_1) p_2 r_1\}, \\ c &= (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) \{(r_1^2 + r_2^2)^2 p_1^2 q_1^2 q_2^2 + 4(q_1 r_1 + q_2 r_2) q_1 q_2 r_1 r_2 \\ &\quad \times (q_1 r_2 - q_2 r_1) p_1 p_2 + (q_1^2 + q_2^2)^2 p_2^2 r_1^2 r_2^2\}, \\ d &= q_1 q_2 (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) (r_1^2 + r_2^2) \{2p_1 q_1 q_2 + (q_1^2 - q_2^2) p_2\} \\ &\quad \times \{(r_1^2 - r_2^2) p_1 - 2p_2 r_1 r_2\}, \\ e &= (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) \{(r_1^2 + r_2^2) p_1 q_1 q_2 + (q_1^2 r_2 - 2q_1 q_2 r_1 \\ &\quad - q_2^2 r_2) p_2 r_2\} \{(r_1^2 + r_2^2) p_1 q_1 q_2 + (q_1^2 r_1 + 2q_1 q_2 r_2 - q_2^2 r_1) p_2 r_1\}, \\ f &= q_1 q_2 \{(r_1^2 + r_2^2)^2 p_1^4 q_1^2 q_2^2 + 2(q_1^2 r_1^4 + q_1^2 r_2^4 + 2q_1 q_2 r_1^3 r_2 \\ &\quad - 2q_1 q_2 r_1 r_2^3 - q_2^2 r_1^4 - q_2^2 r_2^4) p_1^3 p_2 q_1 q_2 + (q_1^4 r_1^4 - q_1^4 r_1^2 r_2^2 + q_1^4 r_2^4 \\ &\quad - q_1^2 q_2^2 r_1^4 - 8q_1^2 q_2^2 r_1^2 r_2^2 - q_1^2 q_2^2 r_2^4 + q_2^4 r_1^4 - q_2^4 r_1^2 r_2^2 + q_2^4 r_2^4) p_1^2 p_2^2 \\ &\quad - 2(q_1^4 r_1^2 - q_1^4 r_2^2 + 2q_1^3 q_2 r_1 r_2 - 2q_1 q_2^3 r_1 r_2 + q_2^4 r_1^2 - q_2^4 r_2^2) p_1 p_2^3 r_1 r_2 \\ &\quad + (q_1^2 + q_2^2)^2 p_2^4 r_1^2 r_2^2\}, \\ A &= q_1 q_2 (p_1 q_2 + p_2 q_1) (p_1 q_1 - p_2 q_2) \{(r_1^2 + r_2^2) p_1 q_1 q_2 + (q_1^2 r_2 - 2q_1 q_2 r_1 \\ &\quad - q_2^2 r_2) p_2 r_2\} \{(r_1^2 + r_2^2) p_1 q_1 q_2 + (q_1^2 r_1 + 2q_1 q_2 r_2 - q_2^2 r_1) p_2 r_1\} \\ &\quad \times \{2p_1^2 q_1 q_2 r_1 r_2 + (q_1 r_2 + q_2 r_1) (q_1 r_1 - q_2 r_2) p_1 p_2 - 2p_2^2 q_1 q_2 r_1 r_2\} \\ &\quad \times \{(r_1^2 - r_2^2) p_1^2 q_1 q_2 + (r_1^2 - r_2^2) (q_1^2 - q_2^2) p_1 p_2 - (q_1^2 - q_2^2) p_2^2 r_1 r_2\}, \end{aligned} \tag{3.22}$$

where $p_i, q_i, r_i, i = 1, 2$, are arbitrary nonzero rational parameters.

As a numerical example, if we take $(p_1, p_2, q_1, q_2, r_1, r_2) = (3, 1, 2, 1, 3, 1)$, we get a rational convex quadrilateral whose sides, diagonals and the area, after appropriate scaling, are given by $(a, b, c, d, e, f, A) = (748, 561, 615, 1000, 935, 1068, 490314)$.

We note that if, in the formulae (3.22), we take q_1, q_2, r_1, r_2 as numerical constants, the sides and diagonals of the quadrilateral are given by quartic polynomials in the parameters p_1 and p_2 . Similarly, we may consider the formulae for the sides and diagonals of the above quadrilateral as quartic polynomials in the parameters r_1 and r_2 .

3.2.3 A second parametrization of noncyclic convex quadrilaterals

To obtain a second parametrization of rational noncyclic convex quadrilaterals, we note that the left-hand side of Eq. (3.11) becomes a perfect square if we take

$$r_1 = -p_2, \quad r_2 = p_1, \quad s_1 = (p_1 - p_2)q_1 - (p_1 + p_2)q_2, \quad s_2 = -(p_1 + p_2)q_1 - (p_1 - p_2)q_2.$$

We thus get a rational quadrilateral whose sides a, b, c, d , the diagonals e, f and the area A may be written, after replacing the parameters p_2 and q_2 by $-p_2$ and $-q_2$ respectively, as follows:

$$\begin{aligned} a &= (p_1^2 + p_2^2)^2 (q_2^2 - q_1^2) q_1 q_2, \\ b &= p_1 p_2 (p_1^2 + p_2^2) (q_2^4 - q_1^4), \\ c &= 2p_1 p_2 (p_1 q_2 + p_2 q_1) (p_2 q_2 - p_1 q_1) (q_1^2 + q_2^2), \\ d &= (p_1 q_2 + p_2 q_1) (p_2 q_2 - p_1 q_1) (p_1 q_1 - p_1 q_2 - p_2 q_1 - p_2 q_2) \\ &\quad \times (p_1 q_1 + p_1 q_2 + p_2 q_1 - p_2 q_2), \\ e &= (p_1^2 + p_2^2) (p_2 q_2 - p_1 q_1) (p_1 q_2 + p_2 q_1) (q_2^2 - q_1^2), \\ f &= p_1 p_2 (p_2^2 - p_1^2) (q_1^2 + q_2^2)^2, \\ A &= p_1 p_2 (p_2^2 - p_1^2) (q_2^2 - q_1^2) (p_1 q_2 + p_2 q_1) (p_2 q_2 - p_1 q_1) \\ &\quad \times (p_1 q_1^2 - p_1 q_2^2 - 2p_2 q_1 q_2) (2p_1 q_1 q_2 + p_2 q_1^2 - p_2 q_2^2), \end{aligned} \tag{3.23}$$

where $p_i, q_i, i = 1, 2$, are arbitrary nonzero rational parameters.

The values of a, b, c, d, e and f given by (3.23) and the values of x_1, x_2, y_1, y_2 given by

$$\begin{aligned} x_1 &= (p_2^4 - p_1^2) (q_2^2 - q_1^2) q_1 q_2, \\ x_2 &= (p_1 q_1 - p_1 q_2 - p_2 q_1 - p_2 q_2) (p_1 q_1 + p_1 q_2 + p_2 q_1 - p_2 q_2) (p_2^2 - p_1^2) \\ &\quad \times (p_1 q_2 + p_2 q_1) (p_2 q_2 - p_1 q_1) / (p_1^2 + p_2^2), \\ y_1 &= 2(p_1^2 + p_2^2) (q_2^2 - q_1^2) p_1 p_2 q_1 q_2, \\ y_2 &= -2p_1 p_2 (p_1 q_1 - p_1 q_2 - p_2 q_1 - p_2 q_2) (p_1 q_1 + p_1 q_2 + p_2 q_1 - p_2 q_2) \\ &\quad \times (p_1 q_2 + p_2 q_1) (p_2 q_2 - p_1 q_1) / (p_1^2 + p_2^2) \end{aligned} \tag{3.24}$$

satisfy the simultaneous Eqs. (2.1)–(2.5).

It follows from Theorem 1 that if we assign positive rational values to the parameters $p_i, q_i, i = 1, 2$, such that $q_2 > q_1$, and $p_2 > p_1(q_1 + q_2)/(q_2 - q_1)$, the quadrilateral defined by (3.23) is a convex quadrilateral.

As a numerical example, on taking $(p_1, p_2, q_1, q_2) = (1, 2, 1, 5)$, we get a rational convex quadrilateral whose sides, diagonals and the area, after appropriate scaling, are given by $(a, b, c, d, e, f, A) = (125, 260, 273, 84, 315, 169, 26334)$.

3.2.4 Rational quadrilaterals with two equal sides

There are special quadrilaterals such as parallelograms, rhombi and kites in which two of the four sides are equal in addition to other geometrical conditions. Parametrizations of such special rational quadrilaterals may be obtained by considering the problem ab initio rather than by considering it as a special case of the general problem of finding rational quadrilaterals. For instance, the problem of finding a rational parallelogram is equivalent to finding a rational triangle with a rational median, and its complete solution has been given by Dickson [5, pp. 249–250]. We will not be concerned with such special quadrilaterals in this paper. We give below a parametrization of a more general quadrilateral in which two of the four sides of the quadrilateral are necessarily equal.

We note that a solution of Eq. (3.11) may be obtained by taking

$$q_1 = -p_2, \quad q_2 = p_1, \quad s_1 = (p_1 - p_2)r_1 - (p_1 + p_2)r_2, \quad s_2 = -(p_1 + p_2)r_1 - (p_1 - p_2)r_2.$$

This yields a rational quadrilateral whose sides a, b, c, d , the diagonals e, f and the area A are given by the following formulae:

$$\begin{aligned} a &= b = (p_1^2 + p_2^2)(r_1^2 + r_2^2), & c &= 4(p_1^2 + p_2^2)r_1r_2, \\ d &= 2(p_1r_1 + p_1r_2 + p_2r_1 - p_2r_2)(p_1r_1 - p_1r_2 - p_2r_1 - p_2r_2), \\ e &= 2(r_1^2 + r_2^2)(p_2^2 - p_1^2), & f &= (p_1^2 + p_2^2)(r_1^2 + r_2^2), \\ A &= 2(p_2^2 - p_1^2)(2p_1r_1r_2 + p_2r_1^2 - p_2r_2^2)(p_1r_1^2 - p_1r_2^2 - 2p_2r_1r_2), \end{aligned} \quad (3.25)$$

where $p_i, r_i, i = 1, 2$, are arbitrary parameters.

It follows from Theorem 1 that if we assign positive rational values to the parameters $p_i, q_i, i = 1, 2$, such that $r_2 > r_1$ and $p_2 > p_1(r_1 + r_2)/(r_2 - r_1)$, the formulae (3.25) will always generate rational convex quadrilaterals with two equal sides. As a numerical example, on taking $(p_1, p_2, r_1, r_2) = (1, 3, 1, 3)$, we get a rational convex quadrilateral whose sides, diagonals and the area, after appropriate scaling, are given by $(a, b, c, d, e, f, A) = (25, 25, 30, 14, 40, 25, 468)$.

3.2.5 Further parametrizations of rational quadrilaterals

We will now show how additional parametrizations of rational quadrilaterals may be obtained by finding more solutions of the diophantine Eq. (3.11).

On writing $s_1 = s_2X$, $f = s_2^2Y$, Eq. (3.11) reduces to

$$\begin{aligned}
Y^2 = & q_1^2q_2^2r_1^2r_2^2(p_1^2 + p_2^2)^2X^4 - 2p_1p_2q_1q_2r_1r_2\{(q_1r_1 + q_1r_2 + q_2r_1 - q_2r_2)p_1 \\
& + (q_1r_1 - q_1r_2 - q_2r_1 - q_2r_2)p_2\}\{(q_1r_1 - q_1r_2 - q_2r_1 - q_2r_2)p_1 - (q_1r_1 + q_1r_2 \\
& + q_2r_1 - q_2r_2)p_2\}X^3 + \{2p_1^4q_1^2q_2^2r_1^2r_2^2 + 8q_1q_2r_1r_2(q_1r_2 + q_2r_1)(q_1r_1 - q_2r_2)p_1^3p_2 \\
& + (q_1^4r_1^4 + 2q_1^4r_1^2r_2^2 + q_1^4r_2^4 + 8q_1^3q_2r_1^3r_2 - 8q_1^3q_2r_1r_2^3 + 2q_1^2q_2^2r_1^4 - 24q_1^2q_2^2r_1^2r_2^2 \\
& + 2q_1^2q_2^2r_2^4 - 8q_1q_2^3r_1^3r_2 + 8q_1q_2^3r_1r_2^3 + q_2^4r_1^4 + 2q_2^4r_1^2r_2^2 + q_2^4r_2^4)p_1^2p_2^2 \\
& - 8q_1q_2r_1r_2(q_1r_2 + q_2r_1)(q_1r_1 - q_2r_2)p_1p_2^3 + 2p_2^4q_1^2q_2^2r_1^2r_2^2\}X^2 + 2p_1p_2q_1q_2r_1r_2 \\
& \times \{(q_1r_1 + q_1r_2 + q_2r_1 - q_2r_2)p_1 + (q_1r_1 - q_1r_2 - q_2r_1 - q_2r_2)p_2\}\{(q_1r_1 - q_1r_2 - q_2r_1 \\
& - q_2r_2)p_1 - (q_1r_1 + q_1r_2 + q_2r_1 - q_2r_2)p_2\}X + q_1^2q_2^2r_1^2r_2^2(p_1^2 + p_2^2)^2. \quad (3.26)
\end{aligned}$$

The right-hand side of Eq. (3.26) is a quartic polynomial in X , and Eq. (3.26) may be considered as representing the quartic model of an elliptic curve over the function field $\mathbb{Q}(p_1, p_2, q_1, q_2, r_1, r_2)$. A point $P = (X_1, Y_1)$ on the elliptic curve (3.26), derived from the solution (3.12) of Eq. (3.11), is as follows:

$$\begin{aligned}
(X_1, Y_1) = & (\{(q_1r_2 + q_2r_1)p_1 + (q_1r_1 - q_2r_2)p_2\} \\
& \times \{(q_1r_1 - q_2r_2)p_1 + (-q_1r_2 - q_2r_1)p_2\}^{-1}, \\
& (p_1q_2 + p_2q_1)(p_1q_1 - p_2q_2)(p_1r_2 + p_2r_1)(p_1r_1 - p_2r_2)(q_1^2 + q_2^2) \\
& \times (r_1^2 + r_2^2)\{(q_1r_1 - q_2r_2)p_1 + (-q_1r_2 - q_2r_1)p_2\}^{-2}).
\end{aligned}$$

Since the coefficient of X^4 in the quartic polynomial on the right-hand side of (3.26) is a perfect square, we can, by following a well-known procedure (see, for instance, [14, p. 77] or [4, pp. 35–36]), transform the elliptic curve (3.26) to the usual cubic model, and find a point P_1 on the cubic model corresponding to the point P on the quartic model given by (3.26). It is easily established that the point P_1 is not of finite order. Thus, by repeated application of the group law, we can find infinitely many rational points on the cubic model of the curve, and hence also on the curve (3.26). We thus get infinitely many rational solutions of Eq. (3.26), and we can thus find infinitely many parametrizations of rational quadrilaterals.

4 Concluding remarks

In this paper we described a method of generating all quadrilaterals whose sides, diagonals and the area are given by rational numbers. This method is different from the classic construction of rational quadrilaterals given by Kummer in 1848. We were able to obtain a complete parametrization of all rational cyclic quadrilaterals. For general rational quadrilaterals, we obtained several parametrizations in terms of quartic polynomials. When two

sides of the quadrilateral are equal, we obtained a parametrization in terms of quadratic polynomials. The formulae for rational quadrilaterals obtained in this paper are simpler than the known parametrizations of such quadrilaterals.

References

- [1] J. H. J. Almering, Rational quadrilaterals, *Indag. Math.* **25** (1963), 192–199.
- [2] J. H. J. Almering, Rational quadrilaterals. II, *Indag. Math.* **27** (1965), 290–304.
- [3] R. H. Buchholz and J. A. MacDougall, Cyclic polygons with rational sides and area, *J. Number Theory* **128** (2008), 17–48.
- [4] J. W. S. Cassels, *Lectures on elliptic curves*, Cambridge University Press, 1991.
- [5] L. E. Dickson, Rational triangles and quadrilaterals, *Am. Math. Month.* **28** (1921), 244–250.
- [6] L. E. Dickson, *History of the theory of numbers Vol. 2*, reprint, Chelsea Publishing Company, 1992.
- [7] H. Eves, *An introduction to the history of mathematics, Third edition*, (Holt, Rinehart and Winston, New York, 1969).
- [8] R. C. Gupta, Paramesvara’s rule for the circumradius of a cyclic quadrilateral, *Historia Math.* **4** (1977), 67–74.
- [9] M. Hajja and J. Sondow, Newton quadrilaterals, the associated cubic equations, and their rational solutions, *Am. Math. Mon.* **126** (2019), 135–150.
- [10] F. Izadi, F. Khoshnam and D. Moody, Heron quadrilaterals via elliptic curves, *Rocky Mt. J. Math.* **47** (2017), 1227–1258.
- [11] R. A. Johnson, *Advanced Euclidean geometry*, Dover Publications, New York, 1960.
- [12] E. E. Kummer, *Collected papers, Vol. 1*, Springer-Verlag 1975, Über die Vierecke, deren Seiten und Diagonalen rational sind, pp. 253–273, (published originally in *Journal für die reine und angewandte Mathematik* **37** (1848), 1–20).

- [13] L. J. Mordell, Rational quadrilaterals, J. London Math. Soc. **35** (1960), 277–282.
- [14] L. J. Mordell, *Diophantine equations*, Academic Press, London, 1969.
- [15] W. Sierpiński, *Elementary theory of numbers* (edited by A. Schinzel), PWN – Polish Scientific Publishers, Warszawa, 1987.
- [16] K. R. S. Sastry, Brahmagupta quadrilaterals, Forum Geom. **2** (2002), 167–173.

Postal address: Ajai Choudhry, 13/4 A, Clay Square, Lucknow - 226001, India

E-mail: ajaic203@yahoo.com