

LOWER ESTIMATES OF THE KOBAYASHI DISTANCE AND LIMITS OF COMPLEX GEODESICS

LUKASZ KOSIŃSKI AND NIKOLAI NIKOLOV

ABSTRACT. It is proved for a strongly pseudoconvex domain D in \mathbb{C}^d with $\mathcal{C}^{2,\alpha}$ -smooth boundary that any complex geodesic through every two close points of D sufficiently close to ∂D and whose difference is non-tangential to ∂D intersect a compact subset of D that depends only on the rate of non-tangentiality. As an application, a lower bound for the Kobayashi distance is obtained.

1. INTRODUCTION

In [9] Huang proved the localization theorem for infinitesimal extremal mappings near \mathcal{C}^3 -smooth strongly pseudoconvex points. This result had various applications, for example in solving the homogeneous Monge-Ampère equations or in establishing the regularities for CR mappings. Its immediate consequence was also the existence of geodesics with prescribed boundary data, a result previously obtained in [5] within the class of \mathcal{C}^{14} -smooth strictly linearly convex domains. Recently, Huang [10] showed that his result remains true for strongly pseudoconvex point with only $\mathcal{C}^{2,\alpha}$ -regularity, $\alpha \in (0, 1]$.

In this paper we extend the localization theorem to non-infinitesimal settings, while the estimate we are able to achieve applied to infinitesimal case is sharper.

Some part of our paper is devoted to yet another application: we obtain lower bounds for the Kobayashi distance. Estimates that we are going to present being in the spirit of Balogh-Bonk results [2] are sensitive in the non-tangential case.

2. DEFINITIONS AND MAIN RESULTS

Let D be a domain in \mathbb{C}^d $z, w \in D$, and $X \in \mathbb{C}^d$. The Kobayashi distance k_D is the largest pseudodistance not exceeding the Lempert

2010 *Mathematics Subject Classification.* 32F45.

Key words and phrases. Kobayashi distance, strongly pseudoconvex domain, strictly linearly convex domain, extremal mapping, complex geodesic.

The first named author was supported by the NCN grant SONATA BIS no. 2017/26/E/ST1/00723. The second named author was partially supported by the Bulgarian National Science Fund, Ministry of Education and Science of Bulgaria under contract KP-06-N52/3.

function

$$l_D(z, w) = \inf\{\tanh^{-1}|\alpha| : \exists\varphi \in \mathcal{O}(\Delta, D) \text{ with } \varphi(0) = z, \varphi(\alpha) = w\},$$

where Δ is the unit disc and $\tanh^{-1}t = \frac{1}{2} \log \frac{1+t}{1-t}$. This distance is the integrated form of the Kobayashi metric

$$\kappa_D(z; X) = \inf\{|\alpha| : \exists\varphi \in \mathcal{O}(\Delta, D), \varphi(0) = z, \alpha\varphi'(0) = X\}.$$

Set $\delta_D(z) = \text{dist}(z, \partial D)$ and $h_D(z, w) = \delta_D(z)^{1/2}\delta_D(w)^{1/2}$.

Recall that if p is a Dini-smooth boundary point of D , then (see [16, Corollary 8])

$$(1) \quad k_D(z, w) \leq \log \left(1 + \frac{2|z-w|}{h_D(z, w)} \right), \quad z, w \text{ near } p.$$

The example of the unit ball shows a similar lower bound has to include the term $\frac{|z-w|^2}{\delta_D(z)^{1/2}\delta_D(w)^{1/2}}$.

Let us mention two results in this direction. If p is a strongly pseudoconvex boundary point of D , then there is $c > 0$ such that (see [18, Theorems 1.6 and 1.7])

$$(2) \quad k_D(z, w) \geq \log \left(1 + \frac{c|z-w|}{\delta_D(z)^{1/2}} \right) \left(1 + \frac{c|z-w|}{\delta_D(w)^{1/2}} \right), \quad z, w \text{ near } p.$$

For strongly pseudoconvex domain D , Balogh and Bonk [2] introduced a positive function g_D on $D \times D$ which depends on δ_D and the Carnot-Carathéodory metric on ∂D , and they prove that $g_D - k_D$ is bounded on $D \times D$. In particular, there exists $C > 0$ such that

$$(3) \quad \log \left(1 + \frac{|z-w|^2}{\delta_D(z)^{1/2}\delta_D(w)^{1/2}} \right) - C \leq k_D(z, w) \\ \leq \log \left(1 + \frac{|z-w|}{\delta_D(z)^{1/2}\delta_D(w)^{1/2}} \right) + C$$

which also follows from (1) and (2).

The following estimate is more sensitive in the non-tangential case.

Proposition 1. *Let p be a strongly pseudoconvex boundary point of a domain D . Then there exist a neighborhood U of p and a constant $c > 0$ such that*

$$k_D(z, w) \geq \log \left(1 + \frac{c|(z-w)_n|}{\delta_D(z)^{1/2}\delta_D(w)^{1/2}} \right), \quad z, w \in D \cap U.$$

Here X_n denotes the projection of X on the real normal to ∂D at the closest point $p(z) \in \partial D$ to z . Note that $|X_n| = 2|\text{Re}\langle X, g_D(z) \rangle|$, where $g_D(z) =: \bar{\partial}\tilde{\delta}_D(z) = \bar{\partial}\tilde{\delta}_D(p(z))$ and $\tilde{\delta}_D$ is the signed distance to ∂D .

Proposition 1 together with (2) lead to the following more precise estimate:

$$k_D(z, w) \geq \log \left(1 + c \left(\frac{|(z-w)_n| + |z-w|^2}{\delta_D(z)^{1/2} \delta_D(w)^{1/2}} + \frac{|z-w|}{\delta_D(z)^{1/2}} + \frac{|z-w|}{\delta_D(w)^{1/2}} \right) \right).$$

One may expect similar results, replacing the real by the complex normal.

Proposition 2. *Let p be a $\mathcal{C}^{2,\alpha}$ -smooth strongly pseudoconvex boundary point of a domain D . Then for any $\varepsilon \in (0, 1]$ there exist a neighborhood U of p and a constant $c > 0$ such that if $z, w \in D \cap U$ and $|(z-w)_N| \geq \varepsilon|z-w|$, then*

$$k_D(z, w) \geq \log \left(1 + \frac{c|(z-w)_N|}{\delta_D(z)^{1/2} \delta_D(w)^{1/2}} \right).$$

Here and in the sequel $\alpha \in (0, 1]$ and X_N denotes the projection of X on the complex normal to ∂D at the closest point $p(z) \in \partial D$ to z . Note that $|X_N| = 2|\langle X, g_D(z) \rangle|$.

It is natural to ask if this result remains true for $\varepsilon = 0$, that is, without the assumption about $(z-w)_N$. Proposition 8 below gives a partial answer to this question.

The key point in the proof of Proposition 2 is the following result which is also of independent interest. To formulate this let us introduce a class of *F-domains*. It is composed of bounded pseudoconvex domains that either have \mathcal{C}^∞ -smooth boundary, or their closures possess Stein neighborhood bases. The motivation for the name that we are using comes from the exposing type theorem that is essentially due to Fornæss (see [7]). Note that any strongly pseudoconvex domain is an *F-domain*.

Theorem 3. *Let D be an F -domain. Suppose that p is a $\mathcal{C}^{2,\alpha}$ -smooth strongly pseudoconvex boundary point of D . Then there is a neighborhood U of p and a constant $C > 0$ such*

$$|(z-w)_N| \leq C|z-w| \operatorname{diam}(\varphi)$$

for any extremal mapping φ through z and w , where $z \neq w \in D \cap U$.

As we mentioned in the introduction the infinitesimal counterpart of the above result has been obtained in [9, 10]. Note that it can be deduced directly from Theorem 3, as the proof of [8, Theorem 1] implies that any extremal mapping whose diameter is small enough must be a complex geodesic.

The proof presented here is elementary in the sense that it follows directly from Lempert's theory in [13, 14]. In particular, we do not consider any additional Lagrange-Euler equations or dwell on showing any more subtle (e.g. boundary) perturbation of stationary discs. The main point of our argument relies on a careful analysis of the scaling carried out in [13].

Theorem 3 together with [5, Proposition 4] (see also (5) below) lead to the following

Corollary 4. *Let D and p be as in Theorem 3. Then for any $\varepsilon \in (0, 1]$ there exist a neighborhood U of p and a compact subset K of D such that if $z \neq w \in D \cap U$ and $|(z - w)_N| \geq \varepsilon|z - w|$, then any extremal mapping through z and w intersects K .*

The respective definitions, as well as other facts about complex geodesic, are given in section 3.

3. EXTREMAL MAPPINGS AND COMPLEX GEODESICS

Let D be a domain in \mathbb{C}^d , $z \neq w \in D$, and $0 \neq X \in \mathbb{C}^d$.

A mapping $\varphi \in \mathcal{O}(\Delta, D)$ is called:

– extremal for $\kappa_D(z; X) := \alpha$ (in short, $\varphi = \varphi_{z;X}$) if $\varphi(0) = z$ and $\alpha\varphi'(0) = X$.

– extremal for $k_D(z; w) := \alpha$ (in short, $\varphi = \varphi_{z,w}$) if $\varphi(0) = z$, $\varphi(\alpha) = w$ and $k_D(z, w) = \tanh^{-1} \alpha$.

– complex geodesic if $k_D(\varphi(\zeta), \varphi(\eta)) = k_\Delta(\zeta, \eta)$ for any $\zeta, \eta \in \Delta$.

Recall that D is said to be strictly linearly convex if D is bounded and, for any $p \in \partial D$, the restriction of the Hessian of $\tilde{\delta}_D$ on $T_p^{\mathbb{C}}(\partial D)$ is a positive Hermitian form.

According to Lempert's theory, if D is strictly linearly convex, there exist unique extremal mappings for $\kappa_D(z; X)$ and $k_D(z, w)$, and they are complex geodesics. Conversely, after a reparametrization, any complex geodesic φ is extremal for $\kappa_D(\varphi(\zeta); \varphi'(\zeta))$ and $k_D(\varphi(\zeta), \varphi(\eta))$ for any $\zeta, \eta \in \Delta$. It is also known (see [15]) that if D has a $\mathcal{C}^{2,\alpha}$ -smooth boundary, then complex geodesics extend \mathcal{C}^{1,α^-} -smoothly to $\overline{\Delta}$ (that is $\mathcal{C}^{1,\alpha-\varepsilon}$ -smoothly for any $\varepsilon > 0$).

Let us mention the following tangential version of Theorem 3.

Theorem 5. ([8, Theorem 1]) *Let D be an F -domain. Suppose that p is a $\mathcal{C}^{2,\alpha}$ -smooth strongly pseudoconvex boundary point of D . Then for any neighborhood U of p , there exists $\varepsilon > 0$ such that for each extremal mapping $\varphi_{z;X}$ with $|z - p| < \varepsilon$ and $|X_N| < \varepsilon|X|$, one has that $\varphi_{z;X}(\Delta) \subset U$ and $\varphi_{z;X}$ is a complex geodesic.*

([3, Proposition 2.5], see also [12]) *The same remains true for each extremal mapping $\varphi_{z,w}$ with $|z - p| < \varepsilon$, $|w - p| < \varepsilon$ and $|z - w|_N < \varepsilon|z - w|$.*

Let us remark that Theorem 5 is stated in [8] under an additional assumption of \mathcal{C}^3 -smoothness, nevertheless the whole proof works in $\mathcal{C}^{2,\alpha}$ -settings, as well.

The next result can be considered as an infinitesimal version of Theorem 3.

Theorem 6. ([10, Corollary 1.2]; [9, Theorem 2] if $\alpha = 1$) *Let p be a $\mathcal{C}^{2,\alpha}$ -smooth strongly pseudoconvex point of a domain D , where $\alpha \in (0, 1]$. Then there exist a neighborhood U of p and $C > 0$ such that for each extremal mapping $\varphi_{z;X}$ with $\varphi_{z;X}(\Delta) \subset U$ one has that $|X_N| < C \text{diam}^\alpha(\varphi_{z;X}(\Delta))|X|$.*

As we already said one can deduce Theorem 6 directly from Theorem 3 with a bit better estimate $|X_N| < C \text{diam}(\varphi_{z;X}(\Delta))|X|$. Note that this estimate is almost precise. Indeed, if D is a ball, then

$$\text{diam}(\varphi_{z;X}(\Delta)) \sim \delta_D(z)^{1/2} + |X_N|/|X|. \quad ^1$$

This observation can be partially extended in a more general setting.

Proposition 7. *Let D be a strongly pseudoconvex domain with $\mathcal{C}^{2,\alpha}$ -smooth boundary. There exist $C > c > 0$ such that for any complex geodesic $\varphi \in \mathcal{O}(\Delta, D)$ one has that*

$$(4) \quad c \left(\delta_D(\varphi(\zeta))^{1/2} + \frac{|\varphi'(\zeta)|_N}{|\varphi'(\zeta)|} \right) \leq \text{diam}(\varphi), \quad \zeta \in \Delta,$$

$$(5) \quad c \max(\delta_D \circ \varphi)^{1/2} \leq \text{diam}(\varphi) \leq C \max(s \circ \delta_D \circ \varphi),$$

where $s(x) = -x^{1/2} \log x$.

In (4) we may assume that $\text{diam}(\varphi)$ is small enough; therefore $\varphi(\Delta)$ is near some $p \in \partial D$ and hence $|\varphi'(\zeta)|_N$ is well-defined.

One may expect that

$$(6) \quad \text{diam}(\varphi) \sim \max(\delta_D \circ \varphi)^{1/2} \sim \max |\varphi'|$$

for any complex geodesic $\varphi \in \mathcal{O}(\Delta, D)$ which is parameterized such that $\delta_D(\varphi(0)) = \max(\delta_D \circ \varphi)$.

Proposition 8. *Proposition 2 remains true for $\varepsilon = 0$ if (6) holds.*

Since (cf. [11, Theorem 19.4.2])

$$(7) \quad \frac{1}{1 - |\zeta|^2} = \kappa_\Delta(\zeta; e) = \kappa_D(\varphi(\zeta); \varphi'(\zeta)) \sim \frac{|\varphi'(\zeta)|_N}{\delta_D(\varphi(\zeta))} + \frac{|\varphi'(\zeta)|}{\delta_D(\varphi(\zeta))^{1/2}},$$

the inequality (4) can be read as

$$(8) \quad \delta_D(\varphi(\zeta)) \lesssim \text{diam}(\varphi) |\varphi'(\zeta)| \delta_\Delta(\zeta).$$

Note that, by [17, Theorem 20],

$$\frac{\delta_D(f(\varphi(0)))}{\delta_D(f(\varphi(\zeta)))} \lesssim \frac{1}{\delta_\Delta(\zeta)}.$$

On the other hand, if (6) is true, then (8) implies the opposite inequality and hence we may replace \lesssim by \sim in (4) and (8) (also,

¹As usual, $f \lesssim g$ means that $f \leq Cg$ for some constant $C > 0$ depending only on D ; $f \sim g$ means that $f \lesssim g \lesssim f$.

$\max |\varphi'| \sim |\varphi'(\zeta)|$ for any $\zeta \in \Delta$). This will give us a quantitative version of the first part of Theorem 5.

Further, for $X \notin T_p^{\mathbb{C}}(\partial D)$, there exists a unique unimodular λ_X such that $\langle \nu_p, \lambda_X X \rangle > 0$, where $\langle \cdot, \cdot \rangle$ is the standard Hermitian product and ν_p is the outward normal to ∂D at p .

Theorems 3, 5 and 6 imply the following.

Corollary 9. *Let D and p be as in Theorem 3. Fix $\varepsilon \in [0, 1)$. Let $(\varphi_{z_j; X_j})$ and $(\psi_{z_j; w_j})$ be sequences of extremal mappings such that $z_j, w_j \rightarrow p$, $X_j \rightarrow X$, and $\frac{z_j - w_j}{|z_j - w_j|} \rightarrow X$.*

(a) *If $X \in T_p^{\mathbb{C}}(\partial D)$, then $\varphi_j, \psi_j \rightarrow p$ uniformly on $\overline{\Delta}$.*

(b) *If $X \notin T_p^{\mathbb{C}}(\partial D)$, then one may find subsequences of $(\varphi_{z_j; X_j})$ and $(\psi_{z_j; w_j})$ which tend, after reparametrizations, resp. to φ and ψ in the \mathcal{C}^1 -norm on $\overline{\Delta}$, where φ and ψ are complex geodesics with $\varphi(1) = \psi(1) = p$ and $\varphi'(1) = \psi'(1) = \lambda_X X$.*

According to Lempert's theory (see also [5, Theorem 2]), φ and ψ coincide up to an automorphism of Δ .

4. PROOFS OF PROPOSITIONS 1, 2, 7, 8 AND COROLLARY 9

Proof of Proposition 1. It is known that there exists a ball V around p such that for any $z \in D$ near p one may find a composition Φ_z (injective on V) of three transformation, all depending continuously of $p(z)$ and hence of z , namely the translation $\pi(z) \rightarrow 0$, a unitary transformation and a change of the variables of the form $\eta = \eta_z = (\zeta_1 + c_z \zeta_1^2 + P_z(\zeta'), \zeta')$, where $c_z \in \mathbb{R}$ and P_z is a quadratic polynomial, such that $G_z = \Phi_z(D \cap V)$ is a convex domain and $\operatorname{Re} \eta_1 < 0$ is the inner normal to ∂G_z at 0. Set $d_z = \delta_{G_z} \circ \Phi_z$. It is easy to check that near p (see below),

$$(9) \quad |(d_z(z) - d_z(w))| = (1 + o(1))|(z - w)_n| + O(|z - w|^2),$$

$$d_z = (1 + o(1))\delta_D.$$

Let now z, w be near p . Assume that $d_z(z) \leq d_z(w)$. Since $G_z \subset \Pi_z = \{\eta : \operatorname{Re} \eta_1 < 0\}$, then

$$\begin{aligned} k_{D \cap V}(z, w) &\geq k_{\Pi_z}(\Phi_z(z), \Phi_z(w)) \geq \frac{1}{2} \log \frac{\delta_{\Pi_z}(\Phi_z(w))}{\delta_{\Pi_z}(\Phi_z(z))} \\ &\geq \frac{1}{2} \log \frac{d_z(w)}{d_z(z)} \geq \log \left(1 + \frac{d_z(w) - d_z(z)}{2d_z(z)^{1/2}d_z(w)^{1/2}} \right). \end{aligned}$$

Hence

$$(10) \quad k_{D \cap V}(z, w) \geq \log \left(1 + \frac{|d_z(w) - d_z(z)|}{2d_z(z)^{1/2}d_z(w)^{1/2}} \right).$$

The same remains true in the case $d_z(z) \geq d_z(w)$ by considering the supporting hyperplane to ∂G_z at the closest point to $\Phi_z(w)$.

On the other hand, the proofs of [4, Theorem 1.4] and [18, Theorem 1.6] imply that for any neighborhood $U \Subset V$ there exists a constant $C > 0$ such that

$$(11) \quad k_{D \cap V}(z, w) \leq k_D(z, w) + C, \quad k_{D \cap V} \leq Ck_D(z, w), \quad z, w \in D \cap U.$$

Combining these two inequalities with (2), (9) and (10) provides a constant $c_1 \in (0, 1)$ such that

$$k_{D \cap V}(z, w) \geq \log \left(1 + \frac{g_D(z, w)}{d_z(z)^{1/2} d_z(w)^{1/2}} \right),$$

where $g_D(z, w) = \max\{c_1|(z-w)_n| - |z-w|^2/c_1, c_1|z-w|^2\}$. It remains to note that $g_D(z, w) \geq \frac{c_1^3}{c_1^2+1}|(z-w)_n|$.

Subproof of (9). Obviously, any translation and unitary transformation keeps δ_D , $|u-v|$ and $|u-v|_N$. So, it suffices to check (9) for $\Phi_z = \eta_z$.

Observe that $\tilde{\delta}_{G_z} \circ \eta_z$ is a defining function for D with the same gradient as δ_D at $p(z) = 0$. This implies that $d_z = (1 + o(1))\delta_D$ near p .

Further, $z = (z_1, 0')$ with $z_1 < 0$ and then

$$\begin{aligned} d_z(w) - d_z(z) &= 2\operatorname{Re}\langle w - z, \bar{\partial}d_z(z) \rangle + O(|w - z|^2) \\ &= (1 + 2c_z z_1)\operatorname{Re}(z_1 - w_1) + O(|w - z|^2). \end{aligned}$$

The proof is finished.

Proof of Proposition 2. Having in mind that D is convexifiable near p , as well as the localizations (11), we may assume that D is strictly convex with $\mathcal{C}^{2,\alpha}$ -smooth boundary.

Let now $c > 0$ and \mathcal{F}_c be the class of complex geodesics $\varphi \in \mathcal{O}(\Delta, D)$ such that $\operatorname{diam}(\varphi) \geq c$ and $\delta_D(\varphi(0)) = \max(\delta_D \circ \varphi)$ (the last one is always achieved after a possible reparametrization). By [5, Proposition 4]) (see also (5)), one may find $\tilde{c} > 0$ such that $\delta_D(\varphi(0)) \geq \tilde{c}$ for any $\varphi \in \mathcal{F}_c$. Then [1, Corollary 2.1] and [8, Proposition 1] provide $\hat{c} > 0$ such that

$$\delta_\Delta \leq \hat{c}(\delta_D \circ \varphi), \quad \max |\varphi'| \leq \hat{c},$$

and hence $|\varphi(\zeta) - \varphi(\eta)| \leq \hat{c}|\zeta - \eta|$ for any $\varphi \in \mathcal{F}_c$.

Since

$$(12) \quad k_D(\varphi(\zeta), \varphi(\eta)) = k_\Delta(\zeta, \eta) \geq \log \left(1 + \frac{|\zeta - \eta|}{2\delta_\Delta(\zeta)^{1/2}\delta_\Delta(\eta)^{1/2}} \right),$$

it follows that

$$k_D(\varphi(\zeta), \varphi(\eta)) \geq \log \left(1 + \frac{|\varphi(\zeta) - \varphi(\eta)|}{2\hat{c}^2\delta_D(\varphi(\zeta))^{1/2}\delta_D(\varphi(\eta))^{1/2}} \right).$$

To complete the proof, it remains to apply Theorem 3.

Proof of Proposition 7. The inequality $|\varphi'(\zeta)|_N/|\varphi'(\zeta)| \lesssim \operatorname{diam}(\varphi)$ follows from Theorem 6.

To prove (4), it remains to show that $\delta_D(\varphi(\zeta))^{1/2} \lesssim \operatorname{diam}(\varphi)$. We may assume that $\zeta = 0$, $\varphi(0) = 0$ and $|\varphi'(0)|_N/|\varphi'(0)| \lesssim \delta_D(0)^{1/2}$. By

(7), we get that $|\varphi'(0)| \sim \delta_D(0)^{1/2}$. We may assume that $|\varphi'_1(0)| \sim \delta_D(0)^{1/2}$. Then the Schwarz lemma implies that

$$\delta_D(0)^{1/2} \sim |\varphi'_1(0)| \leq \sup_{\Delta} |\varphi_1| < \text{diam}(\varphi).$$

Further, the first inequality in (5) is a consequence of (4).

Finally, it follows by [17, Theorem 8] that there exists $C > 0$ such that the Euclidian length of $\varphi(\gamma)$ does not exceed

$$C \max_{u \in \varphi(\gamma)} \left(\delta_D(u)^{1/2} \log \frac{1}{\delta_D(u)} \right),$$

for any real geodesic γ w.r.t. k_{Δ} . This implies second inequality in (5) even in the \mathcal{C}^2 -smooth case.

Proof of Proposition 8. As in Proposition 2, we may assume that D is strictly convex with $\mathcal{C}^{2,\alpha}$ -smooth boundary.

Let $\varphi \in \mathcal{O}(\Delta, D)$ be a complex geodesic. Since

$$|\varphi(\zeta) - \varphi(\eta)| \leq |\zeta - \eta| \cdot \max |\varphi'|$$

and

$$\frac{1}{2} \log \frac{\delta_D(\varphi(0))}{\delta_D(\varphi(\theta))} \leq k_D(\varphi(0), \varphi(\theta)) = k_{\Delta}(0, \theta) < \frac{1}{2} \log \frac{2}{\delta_{\Delta}(\theta)},$$

it follows by (12) and Theorem 3 that

$$\begin{aligned} k_D(\varphi(\zeta), \varphi(\eta)) &\geq \log \left(1 + \frac{\delta_D(\varphi(0))}{4 \max |\varphi'|} \cdot \frac{|\varphi(\zeta) - \varphi(\eta)|}{\delta_D(\varphi(\zeta))^{1/2} \delta_D(\varphi(\eta))^{1/2}} \right) \\ &\geq \log \left(1 + \frac{\delta_D(\varphi(0))}{4C \max |\varphi'| \cdot \text{diam}(\varphi)} \cdot \frac{|(\varphi(\zeta) - \varphi(\eta))_N|}{\delta_D(\varphi(\zeta))^{1/2} \delta_D(\varphi(\eta))^{1/2}} \right). \end{aligned}$$

To complete the proof, it remains to apply (6).

Proof of Corollary 9. (a) follows from Theorem 5.

Let us prove (b). By Theorems 3 and 6, we may assume that, after possible reparametrizations, $\varphi_{z_j; X_j}, \psi_{z_j; w_j} \in \mathcal{F}_c$ for some $c > 0$, and $\varphi_{z_j; X_j}^{-1}(z_j), \psi_{z_j; w_j}^{-1}(z_j) \rightarrow 1$. The proof of [3, Proposition 2.5] implies that, up to subsequences, $\varphi_{z_j; X_j} \rightarrow \varphi$ and $\psi_{z_j; w_j} \rightarrow \psi$ in the \mathcal{C}^1 -norm on $\overline{\Delta}$, where φ and ψ are complex geodesics. Clearly, $\varphi(1) = \psi(1) = p$, $\varphi'(1) = \lambda X$ and $\psi'(1) = \mu X$ for some $\lambda, \mu \in \mathbb{C}$. The Hopf lemma implies that $\lambda/\lambda_X, \mu/\lambda_X > 0$. After automorphisms of Δ , we may get $\lambda = \mu = \lambda_X$.

5. PROOF OF THEOREM 3

In the proof we can assume that D is strictly convex and φ is a geodesic. Actually, the result is trivial when the diameter of an extremal mapping passing through z and w is big. If, in turn, it is small enough, the extremal mapping lies entirely in a neighborhood of some boundary point p , which can be exposed to a strictly convex one (globally) — see the proof of [6, Theorem 1.1] (based on a Fornæss type embedding

theorem). Then this mapping is a stationary disc (see [13, 14]) and consequently a complex geodesic.

Let us recall here again that the core of the our argument relies on a careful analysis of Lempert's idea from his seminal paper [13].

Take $\tilde{z} \in D$ sufficiently close to ∂D . Let us mention that all transformations and function that will appear below can be chosen to depend continuously on \tilde{z} , and all constants will be uniformly bounded. Along the proof we shall keep the notation from the introduction: if $X \in \mathbb{C}^n$, then $|X_N| = 2|\langle X, g_D(\tilde{z}) \rangle|$.

Translate the domain D so that the closest point to \tilde{z} on ∂D is 0. Using unitary map we can make the Levi form at 0 of a defining function r of ∂D to be a positive diagonal matrix. Replacing the coordinates z_j by $\lambda_j z_j$ we can additionally assume that it is the identity matrix. Using unitary map again we can assume that the tangent space is $\text{Re } z_1 = 0$, i.e. that $r(z) = 2 \text{Re } z_1 + |z|^2 + 2 \text{Re } P(z) + O(|z|^{2+\alpha})$, where P is a quadratic polynomial. All these transformations used so far (denote the composition of all of them by F) satisfy the following estimate:

$$\frac{|(\tilde{z} - w)_N|}{|\tilde{z} - w|} \sim \frac{|(F(\tilde{z}) - F(w))_N|}{|F(\tilde{z}) - F(w)|}.$$

Moreover, the point \tilde{z} is mapped a point of the form $(-s, 0)$, where $s > 0$ (in particular, $(\tilde{z} - w)_N = \tilde{z}_1 - w$ and $(F(\tilde{z}) - F(w))_N = F_1(\tilde{z}) - F_1(w)$). Then composing F with an additional transformation $z_1 \mapsto z_1 + P(z)$, $z' \mapsto z'$ gives a biholomorphic map G in a neighborhood U of 0 such that

$$\frac{|F_1(\tilde{z}) - F_1(w)|}{|F(\tilde{z}) - F(w)|} = \frac{|G_1(\tilde{z}) - G_1(w)|}{|G(\tilde{z}) - G(w)|} + O(|\tilde{z}| + |\tilde{z} - w|)$$

and the boundary of $G(D \cap U)$ near 0 is of the form $\{2 \text{Re } \zeta_1 + |\zeta|^2 + O(|\zeta|^{2+\alpha}) < 0\}$. The translation moves 0 to $e_1 = (1, 0')$ and $G(D \cap U)$ to $\{|\zeta|^2 + O(|\zeta - e_1|^{2+\alpha}) < 1\}$. Then point \tilde{z} is mapped to a point of the form $(1 - s + \gamma s^2, 0')$, where $\gamma \in \mathbb{C}$. Moreover, $d_D(\tilde{z}) = s + O(s^2)$.

Let us denote $\varphi = \varphi_{\tilde{z}, w}$. Since $\text{diam } \varphi \gtrsim |\tilde{z}|, |\tilde{z} - w|$ (the first inequality follows from the fact that geodesics are proper and that $|\tilde{z}| \sim \delta_D(\tilde{z})$, the second is trivial) it suffices to focus on achieving estimates for the term $|G_1(\tilde{z}) - G_1(w)|/|G(\tilde{z}) - G(w)|$.

Let $m_t(\lambda) = \frac{\lambda+t}{1+t\lambda}$, $A_t(z) = \left(m_t(z_1), \sqrt{1-t^2} \frac{z'}{1+t z_1}\right)$, and $r_t(z) = \frac{1+t z_1^2}{1-t^2} r(A_t(z))$, $t \in (0, 1)$. Then r_t converges to $\rho(z) = -1 + |z|^2$ in the $\mathcal{C}^{2+\alpha}$ -topology, as $t \rightarrow 1$, when restricted to $\{\text{Re } z_1 > -1/2\}$ (see [13, p. 468]; see also [12]).

Take t such that an analytic disc $A_t^{-1} \circ \varphi$ lies entirely in $\{\text{Re } z_1 > 0\}$ and its boundary touches $\{\text{Re } z_1 = 0\}$, say at point $\mu \in \mathbb{T}$. This, means that

$$(13) \quad \frac{t}{1+t^2} = \frac{\text{Re } \varphi_1(\eta)}{1 + |\varphi_1(\eta)|^2}.$$

It is also clear that the above discs are stationary maps, and thus geodesics, in $A_t^{-1}(D) \cap \{\operatorname{Re} z_1 > -1/2\}$

Claim. There exists $C_1 > 0$ such that $\operatorname{diam}(A_t^{-1} \circ \varphi) > C_1$ for any $t \in (0, 1)$. In particular, all of $A_t^{-1} \circ \varphi$ intersect a compact subset K of $\mathbb{B} \cap \{\operatorname{Re} z_1 > -1/2\}$ (compare with the proof of [5, Proposition 4]).

To prove the claim note that it suffices to show that $A_t^{-1}(\varphi(\eta))$ and $A_t^{-1}(\varphi(0)) = A_t^{-1}(\tilde{z})$ are far away from each other. If this were not the case, $A_t^{-1}(\varphi(\eta))$ would be close to $(\pm i, 0')$, as $A_t^{-1}(\tilde{z}) = (m_t^{-1}(1 - s + \gamma s^2), 0')$.

Expressing $m_t(a + ib) = 1 - s' + i\gamma's'$, where a, b, s' and γ' are real, we see that then $\gamma' = \frac{(1+t)b}{(1+ta)(1-a)-ib^2}$. In particular, if we write $m_t^{-1}(1 - s + \gamma s^2) = a + ib$, $a, b \in \mathbb{R}$, then b must be small (express $1 - s + \gamma s^2 = 1 - s' + iO(s'^2) \in \mathbb{R} + i\mathbb{R}$, where $s' = s + O(s^2)$). This proves the claim.

It thus follow from this claim and Lempert's results [15] that after a possible reparametrization the geodesic $A_t^{-1} \circ \varphi$ is close to a geodesic in the ball in the $\mathcal{C}^{1+\alpha^-}$ -topology (as t is close to 1). Using just the \mathcal{C}^1 -topology we easily get that there exists $C > 0$ such that

$$\frac{|(A_t^{-1}(\tilde{z}) - A_t^{-1}(w))_N|}{|(A_t^{-1}(\tilde{z}) - A_t^{-1}(w))_T|} < C$$

for any $t \in (0, 1)$, where $X_T = X - X_N$. Denoting $x = A_t^{-1}(\tilde{z})$, $y = A_t^{-1}(w)$, we can easily transform the equality above to $|x_1 - y_1|/|y'| < C$. Let us compute

$$\frac{|(\tilde{z} - w)_N|}{|(\tilde{z} - w)_T|} = \frac{|(A_t(x) - A_t(y))_N|}{|(A_t(x) - A_t(y))_T|} = \frac{\sqrt{1-t^2}|x_1 - y_1|}{|(1+tx_1)y'|} \leq C\sqrt{1-t^2}.$$

It follows from (13) that $\frac{(1-t)^2}{1+t^2} = \frac{|1-\varphi_1(\eta)|^2}{1+|\varphi_1(\eta)|^2}$, so

$$\sqrt{1-t^2} \sim |1-\varphi_1(\eta)| \leq |\varphi(\eta) - e_1| \leq |\varphi(\eta) - \tilde{z}| + |\tilde{z} - e_1| \lesssim \operatorname{diam}(\varphi)$$

(note that $|\tilde{z} - e_1| \sim \delta_D(\tilde{z})$) which completes the proof.

Acknowledgement. The authors would like to thank Pascal J. Thomas for careful reading of the manuscript.

REFERENCES

- [1] M. Abate, *Boundary behavior of invariant distances and complex geodesics*, Atti. Accad. Naz. Lincei Rend. 80 (1986), 100–106.
- [2] Z.M. Balogh, M. Bonk, *Gromov hyperbolicity and the Kobayashi metric on strictly pseudoconvex domains*, Comment. Math. Helv. 75 (2000), 504–533.
- [3] F. Bracci, J.E. Fornæss, E.F. Wold, *Comparison of invariant metrics and distances on strongly pseudoconvex domains and worm domains*, Math. Z. 292 (2019), 879–893.
- [4] F. Bracci, N. Nikolov, P.J. Thomas, *Visibility of Kobayashi geodesics in convex domains and related properties*, Math. Z. 301 (2022), 2011–2035.
- [5] C.-H. Chang, M.C. Hu, H.-P. Lee, *Extremal analytic discs with prescribed boundary data*, Trans. Amer. Math. Soc. 310 (1988), 355–369.

- [6] K. Diederich, J.E. Fornæss, E.F. Wold, Exposing points on the boundary of a strictly pseudoconvex or a locally convexifiable domain of finite 1-type, *J. Geom. Anal.* 24 (2014), no. 4, 2124–2134.
- [7] J.E. Fornæss, *Embedding strictly pseudoconvex domains in convex domains*, *Amer. J. Math.* 98, (1976), 529–569.
- [8] X. Huang, *A preservation principle of extremal mappings near a strongly pseudoconvex point and its applications*, *Illinois J. Math.* 38, 2 (1994), 283–302.
- [9] X. Huang, *A non-degeneracy property of extremal mappings and iterates of holomorphic self-mappings*, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (4) Vol. XXI* (1994), 399–419.
- [10] X. Huang, *Revisiting a non-degeneracy property for extremal mappings*, *Acta Math. Sci.* 41B (2021), 1829–1838.
- [11] M. Jarnicki, P. Pflug, *Invariant distances and metrics in complex analysis – 2nd edition*. de Gruyter Expositions in Mathematics 9, Walter de Gruyter, Berlin/Boston, 2013.
- [12] L. Kosiński, *Comparison of invariant functions and metrics*, *Arch. Math. (Basel)* 102 (2014), 271–281.
- [13] L. Lempert, *La métrique de Kobayashi et la représentation des domaines sur la boule*, *Bull. Soc. Math. France* 109 (1981), 427–474.
- [14] L. Lempert, *Intrinsic distances and holomorphic retracts*, *Complex Analysis and Applications '81*, Sofia (1984), 341–364.
- [15] L. Lempert, *A precise result on the boundary regularity of biholomorphic mappings*, *Math. Z.* 193 (1986), no. 4, 559–579.
- [16] N. Nikolov, L. Andreev, *Estimates of the Kobayashi and quasi-hyperbolic distances*, *Ann. Mat. Pura Appl.* 196 (2017), 43–50.
- [17] N. Nikolov, A.Y. Ökten, *Strongly Goldilocks domains, quantitative visibility, and applications*, arXiv:2206.08344.
- [18] N. Nikolov, P.J. Thomas, *Comparison of the real and the complex Green functions, and sharp estimates of the Kobayashi distance*, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) Vol. XVIII* (2018), 1125–1143.

L. KOSIŃSKI: INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, JAGIELLONIAN UNIVERSITY, ŁOJASIEWICZA 6, 30-348 KRAKÓW, POLAND

Email address: lukasz.kosinski@uj.edu.pl

N. NIKOLOV, INSTITUTE OF MATHEMATICS AND INFORMATICS, BULGARIAN ACADEMY OF SCIENCES, ACAD. G. BONCHEV 8, 1113 SOFIA, BULGARIA
FACULTY OF INFORMATION SCIENCES, STATE UNIVERSITY OF LIBRARY STUDIES AND INFORMATION TECHNOLOGIES, SHIPCHENSKI PROHOD 69A, 1574 SOFIA, BULGARIA

Email address: nik@math.bas.bg