

ARTICLE - ARXIV.ORG

How much does it cost to complete a sticker collection? A simulation study using R

Edson Zangiacomi Martinez^a, Jorge Alberto Achcar^a, Marcos Vinicius de Oliveira Peres^b, Ricardo Puziol de Oliveira^c, Regina Albanese Pose^d

^aFaculdade de Medicina de Ribeirão Preto, Universidade de São Paulo (USP), Ribeirão Preto, Brazil.

^bUniversidade Estadual do Paraná, Campus de Paranavaí, Brazil.

^cUniversidade Estadual de Maringá, Maringá, Brazil.

^dUniversidade Municipal de São Caetano do Sul (USCS), São Caetano do Sul (SP), Brazil.

ARTICLE HISTORY

Compiled September 1, 2022

ABSTRACT

This paper considers some computational issues related to the problem of the coupon collector behavior. In addition, we propose a simulation algorithm to find the solution to the problem, where we present as a numerical illustration an album of stickers related to the Qatar 2022 Official Football World Cup, which is being sold in Brazilian newsstands. In this case, the complete album has $N = 670$ stickers, which are sold in packages containing $n = 5$ stickers each. The simulation was based on a computational code written in R. For this application we obtained as a solution, an average number of 950 packages needed to complete the album, at an average cost of 3,800 reais (Brazilian currency), assuming that the collector does not exchange any of its repetitions with nobody. All computer code used in this article is provided on GitHub. The assumed simulation study could be useful for statistics teachers to motivate their students for lectures in computational statistics and fundamentals of probability.

KEYWORDS

Coupon collector's problem; Probability; Simulation; Statistics education.

1. Introduction

The coupon-collector's problem is a classical probability problem. It considers one person that collects coupons and assumes that there is a finite number of different coupons, labeled by the numbers $1, 2, \dots, N$. In this case, we may be interested to find the probability of completing the collection after acquiring exactly n ($n > N$) coupons or in the estimation of the mean number of coupons needed to complete the collection. It is believed that this problem appeared earlier in the work *De Mensura Sortis* (On the Measurement of Chance), written by Abraham de Moivre in 1711, and further results were obtained by Laplace and Euler (Ferrante and Saltalamacchia, 2014). It has many applications in different areas of science such as ecology and evolution (Zoroa et al., 2014), computer science (Nakata and Kubo, 2006), engineering (Boneh and Hofri,

Corresponding author: Ribeirão Preto Medical School, Av. Bandeirantes 3900, University of São Paulo (USP), Ribeirão Preto, 14049-900, Brazil. E-mail: edson@fmrp.usp.br

1997) and quality engineering (Luko, 2009).

This article revisits the coupon-collector's problem and introduces a simulation program developed in the R language. For this purpose, we consider as a special illustrative example, the official 2022 Qatar soccer World Cup sticker album, released in Brazil in August 2022, which is being sold in Brazilian news-stands. The results obtained in this study, especially the results related to the simulation study, could be very useful for statistics teachers to motivate students in teaching fundamentals of probability and statistics in basic and intermediate courses.

2. Methods

2.1. The sticker-collector's problem

Let us consider a population whose members are of N different types (in our case, all stickers from a sticker book). For $1 \leq j \leq N$, let us denote by p_j the probability that a sticker is of type j , where $p_j > 0$ and $\sum_{j=1}^N p_j = 1$. The simplest case occurs when one takes

$$p_1 = p_2 = \dots = p_N = \frac{1}{N},$$

or say, each sticker is produced in equal numbers and so has an equal chance of being found in the next packet that the collector open. In this case, R_i denotes the additional number of stickers that the collector needs to purchase to pass from $i - 1$ to i different types of stickers in her/his collection. Note that R_i has a geometric law with parameter

$$q_i = \frac{N - (i - 1)}{N}.$$

Therefore, $q_1 = 1$, $q_2 = \frac{N-1}{N}$, $q_3 = \frac{N-2}{N}$, and so on. If R is the number of stickers that the collector has to buy in order to find all N existing different stickers, we have that

$$R = R_1 + R_2 + \dots + R_N,$$

and

$$\begin{aligned} E(R) &= \sum_{i=1}^N E(R_i) = \sum_{i=1}^N \frac{1}{q_i} = \sum_{i=1}^N \frac{N}{N - (i - 1)} \\ &= N \sum_{i=1}^N \frac{1}{N - (i - 1)} = N \sum_{i=1}^N \frac{1}{i}. \end{aligned} \tag{1}$$

An approximation of (1) is given by

$$E(R) \approx N(\ln N + \gamma) + 0.5, \tag{2}$$

where $\gamma \approx 0.5772156649$ is the Euler-Mascheroni constant (Doumas, 2015), defined as the limit of the sequence

$$\gamma = \lim_{N \rightarrow \infty} \left(\sum_{i=1}^N \frac{1}{i} - \ln N \right).$$

2.2. The collector's problem in the case of sampling in groups of constant size

We now formulate the collector's problem as follows: given a collection with N different stickers, how often must we draw subsets of $1 \leq n \leq N$ distinct stickers in order to see each stickers at least once? This subsection is based on Stadje (1990), Leite et al. (1992), Adler and Ross (2001), and Diniz et al. (2016), and considers that the stickers are sold in packets of n stickers each. Let N be the total number of stickers in a collection and T denotes the number of packets necessary to complete the collection. The probability that more than k packets are needed to complete the collection is given by $P(T > k)$. Let us consider the following event:

$A_{i,k}$: sticker i is not selected in the first k packets.

The probability $P(T > k)$ is thus given by

$$P(T > k) = P\left(\bigcup_{i=1}^N A_{i,k}\right). \quad (3)$$

Assuming independence between packets, we have $P(A_{i,k}) = p_1^k$ where

$$p_r = \frac{\binom{N-r}{n}}{\binom{N}{n}}$$

is the probability that no sticker from a subset of r stickers is found in the j -th packet with n stickers each ($k \geq 1$, $n < N$, $N - \min(kn, N) \leq r \leq N - n$). Diniz et al. (2016) observed that p_r neither depends on the particular subset nor the particular packet j and, for distinct elements i_1, i_2, \dots, i_s of the set $\{1, \dots, N\}$, the authors used the notation

$$\begin{aligned} p_1^k &= P(A_{i_1,k}) \\ p_2^k &= P(A_{i_1,k} \cap A_{i_2,k}) \\ &\vdots \\ p_s^k &= P(A_{i_1,k} \cap A_{i_2,k} \cap \dots \cap A_{i_s,k}). \end{aligned} \quad (4)$$

From (3) and (4), we have

$$\begin{aligned}
P(T > k) &= \sum_{i_1=1}^N P(A_{i_1,k}) - \sum_{i_1 \neq i_2} P(A_{i_1,k} \cap A_{i_2,k}) + \dots \\
&\quad + (-1)^{N-1} P(A_{1,k} \cap A_{2,k} \cap \dots \cap A_{N,k}) \\
&= \binom{N}{1} p_1^k - \binom{N}{2} p_2^k + \dots + (-1)^{N-1} \binom{N}{N} p_N^k \\
&= \sum_{i=1}^N (-1)^{i-1} \binom{N}{i} p_i^k,
\end{aligned} \tag{5}$$

and

$$\begin{aligned}
P(T = k) &= P(T > k-1) - P(T > k) \\
&= \sum_{i=1}^N (-1)^{i-1} \binom{N}{i} p_i^{k-1} - \sum_{i=1}^N (-1)^{i-1} \binom{N}{i} p_i^k \\
&= \sum_{i=1}^N (-1)^{i-1} \binom{N}{i} (p_i^{k-1} - p_i^k) \\
&= \sum_{i=1}^N (-1)^{i-1} \binom{N}{i} p_i^{k-1} (1 - p_i).
\end{aligned}$$

The expected number of packets necessary to complete the collection is given by

$$E(T) = \sum_{i=1}^N (-1)^{i-1} \binom{N}{i} \frac{1}{1 - p_i}. \tag{6}$$

2.3. A computational simulation

We use the R software to simulate the number of sticker packets that are required to complete one collection sequentially. If we run this simulation a relatively large number of times, i.e., assuming many collectors are buying stickers simultaneously, we can estimate the average number of packets necessary to fill the albums. Considering the official 2022 Qatar soccer World Cup sticker album, we have $N = 670$ and $n = 5$. We simulate each purchased packet of stickers as a uniform sample of size n drawn from a population of size N with replacement. Given N different stickers, the probability of finding n different stickers in a packet of n stickers is

$$\frac{N(N-1) \cdots (N-n+1)}{N^n} = \prod_{j=1}^{n-1} \left(1 - \frac{j}{N}\right).$$

For $N = 670$ and $n = 5$ we have

$$\prod_{j=1}^4 \left(1 - \frac{j}{670}\right) \simeq 0.985.$$

The steps of the algorithm are as follows.

- (1) Assume that $N = 670$ is the total in the album, $k = 5$ is the number of stickers in a packet, $H = 0$ is the number of stickers that the collector have, and $D = 0$ is the number of duplicate stickers.
- (2) Let `is.missing` be a vector with the labels of the missing stickers. Initially `is.missing` is given by $\{1, 2, 3, \dots, 670\}$.
- (3) Let `stickers.b = 0` be the number of packets that are required to complete the collection.
- (4) Simulate one packet by taking a random sample of size $k = 5$ from the elements of $\{1, 2, 3, \dots, 670\}$.
- (5) Verify that each of the simulated stickers belongs to the vector `is.missing`. If a sticker does not belong to this vector, it is a duplicate sticker. In this case we take $D = D + 1$. Otherwise we exclude the sticker from the vector `is.missing` and take H equal to N minus the number of elements of `is.missing`.
- (6) Take `stickers.b = stickers.b + 1`.
- (7) Repeat the items (4) to (6) while $H < N$.

In order to provide a didactic approach to this simulation, a code in R was developed to help visualize in an animation the steps of the algorithm until an album is filled.

We can use this algorithm to simulate a large B number of collections, and from there we can estimate the average number of packets necessary to fill the albums without using the equations (1), (2), or (6).

2.4. Computational details

All codes and data underlying this work are available in GitHub at <https://github.com/edsonzmartinez/stickers>.

3. Results

According to the expression (1), an average of $E(R) = 4,748$ stickers are required to complete a collection of $N = 670$ stickers. We get the same result for $E(R)$ if we use the approximation given by expression (2). Alternatively, considering $N = 670$ and that the stickers come in packets of $n = 5$, we approximated the expression (6) to be 946.98 using the WolframAlpha calculator (Dimiceli et al., 2010). Rounding up, we get 947 packets and $947 \times 5 = 4,735$ stickers. At a cost of 4 Brazilian reais a packet, this implies a total average cost of $947 \times 4 = 3,788$ Brazilian reais (BRL) per album (as of August 2022, 1 BRL was equivalent to 0.19 US dollars).

Table 1 shows the probability $1 - P(T > k)$ of completing the album given the number k of purchased packets and the corresponding cost in BRL. Values of $P(T > k)$ were obtained using the expression (5). In addition, Figure 1 describes the relationship between $1 - P(T > k)$ and k . Table 1 and Figure 1 show that the probabilities of completing the album by purchasing 826, 918 and 1036 packets are, respectively, near to 0.25, 0.50, 0.75 (the lower quartile, median, and upper quartile). If a collector purchases 1500 packets, he/she will have a chance of about 0.99 to complete the album at a cost of 6,000 BRL.

Table 1. Probability of completing the album given the number k of purchased packets.

k	$1 - P(T > k)$	Cost (BRL)
500	0.00001	2000.00
600	0.00044	2400.00
650	0.00516	2600.00
700	0.02735	2800.00
750	0.08516	3000.00
800	0.18488	3200.00
826	0.24972	3304.00
850	0.31416	3400.00
900	0.45170	3600.00
918	0.49950	3672.00
950	0.57939	3800.00
1000	0.68734	4000.00
1036	0.75114	4144.00
1100	0.83773	4400.00
1150	0.88541	4600.00
1200	0.91974	4800.00
1300	0.96123	5200.00
1500	0.99120	6000.00
1600	0.99583	6400.00
1650	0.99713	6600.00
1700	0.99803	6800.00
1750	0.99864	7000.00
1755	0.99869	7020.00

3.1. Simulation results

Figure 2 shows a grid representing the results of a simulation after the purchase of 30 sticker packets. Continuing this simulation, Figure 3 shows that it was required 824 packets to complete the album, at a cost of 3,296 BRL. Based on this algorithm, we simulated $B = 100.000$ independent collections. The number of required packets ranged from 547 to 2,455, and the corresponding cost ranged from 2,188 to 9,820 BRL. Figure 4 shows a histogram of the distribution of the number of packets required to complete these 100.000 collections. The average number of required packets was 950 at a cost of 3,800 BRL. The lower quartile, median, and upper quartile were 829, 921, and 1,038, respectively, at corresponding costs of 3,316, 3,684, and 4,152 BRL. Therefore, the results of our simulation are close to that showed in Table 1 and Figure 1, based on the expression (5).

4. Discussion

Using a simulation study, we show that it is necessary to purchase on average 950 packets of $n = 5$ stickers, at a cost of 3,800 BRL, to complete a collection of $N = 670$ stickers. Similar results were obtained using expression (6), which does not require the use of computational simulation. However, these results ignore the facility for buying missing stickers and that collectors can swap stickers with other collectors. Intuitively, collectors know that filling a collection is easier when they exchange duplicated stickers with other people. According to Hayes and Hannigan (2006), although statisticians frequently use the theory of the coupon collector’s problem to respond to inquiries




Figure 1. Probability of completing the album according to the number of packets sold.




Figure 2. Results of a simulation after the purchase of 30 sticker packets.

from newspapers about the cost of filling sticker albums, it is possible that the only way to find the true answer is by simulating the trading environment of the school playground (or other social environments). In this way, authors such as Braband et al. (2015), Morgado and Borges (2019), and Rosa and Probst (2021) proposed different




Figure 3. Final results of the simulation, where it was required 824 packets to complete the album.




Figure 4. Distribution of the number of packets required to complete $B = 100.000$ independent collections.

extensions of the coupon-collector's problem considering that collectors trade duplicate stickers with other people. These articles show how trading repeated stickers with other collectors reduces substantially the amount of stickers that the collector needs to complete the album.

Although the coupon-collector's problem contextualized in collections soccer championships' stickers has already been extensively explored in the literature, the present article can contribute to educating students about some fundamentals of probability and computational statistics by providing detailed R codes for implementing the problem. We believe that the simulation study presented in this article can benefit high

school and college students. For example, by showing that the number of sticker packets needed to complete the collection is not a fixed number but an amount that can vary from one collector to another, we exemplify the concept of a random phenomenon. It is possible that one collector will complete the collection earlier than another, but this is determined by chance.

References

- [1] Adler, I. and Ross, S. M. (2001). The coupon subset collection problem, *Journal of Applied Probability*, **38**, 737–746.
- [2] Alistarh, D., and Davies, P. (2021). Collecting coupons is faster with friends, *International Colloquium on Structural Information and Communication Complexity*, 3–12.
- [3] Boneh, A. and Hofri, M. (1997). The coupon-collector problem revisited—a survey of engineering problems and computational methods, *Stochastic Models*, **13**, 39–66.
- [4] Braband, N., Braband, S., and Braband, M. (2015). New results for the Coupon Collector’s problem with swapping and replacement, *arXiv preprint arXiv*, **1504.00296**.
- [5] Dimiceli, V. E., Lang, A. S., and Locke, L. (2010). Teaching calculus with Wolfram Alpha, *International Journal of Mathematical Education in Science and Technology*, **41**, 1061–1071.
- [6] Diniz, M. A., Lopes, D., Polpo, A., and Salasar, L. E. B. (2016). The Sticker Collector’s Problem, *The College Mathematics Journal*, **47**, 255–263.
- [7] Doumas, A. V. (2015). How many trials does it take to collect all different types of a population with probability p ?, *Journal of Applied Mathematics and Bioinformatics*, **5**, 1–14.
- [8] Ferrante, M. and Saltalamacchia, M. (2014). The coupon collector’s problem, *Materials Mathematics*, **2**, 1–35.
- [9] Hayes, K. and Hannigan, A. (2006). Trading coupons: completing the world cup football sticker album, *Significance*, **3**, 142–144.
- [10] Leite, J. G., Pereira, C. A. B., and Rodrigues, F. W. (1992). Waiting time to exhaust lottery numbers, *Communications in Statistics - Theory and Methods*, **22**, 301–310.
- [11] Luko, S. N. (2009). The “Coupon Collector’s Problem” and Quality Control, *Quality Engineering*, **21**, 168–181.
- [12] Morgado, L. B. and Borges L. S. (2019). Sticker albums: an approach via Markov chains, *REMAT*, **5**, 101–113.
- [13] Nakata, T. and Kubo, I. (2006). A coupon collector’s problem with bonuses, *Discrete Mathematics and Theoretical Computer Science Proceedings*, 215–224.
- [14] Rosa, G. M. and Probst, R. W. (2021). Quanto custa completar o álbum de figurinhas da Copa?, *Proceeding Series of the Brazilian Society of Computational and Applied Mathematics*, **8**, 1–7.
- [15] Sardy, S. and Velenik, Y. (2014). Paninimania: sticker rarity and cost-effective strategy, Université de Genève. Available from: <https://www.unige.ch/math/folks/velenik/Vulg/Paninimania.pdf>.
- [16] Stadje, W. (1990). The collector’s problem with group drawings, *Advances in Applied Probability*, **22**, 866–882.
- [17] Zoroa, N., Lesigne, E., Fernández-Sáez, M. J., Zoroa, P., and Casas, J. (2017). The coupon collector urn model with unequal probabilities in ecology and evolution, *Journal of The Royal Society Interface*, **14**, 20160643.