

Background independence and the Adler-Bardeen theorem

Jochen Zahn

Institut für Theoretische Physik, Universität Leipzig
Brüderstr. 16, 04103 Leipzig, Germany
jochen.zahn@itp.uni-leipzig.de

September 6, 2022

Abstract

We prove that for renormalizable Yang-Mills gauge theory with arbitrary compact gauge group (of at most a single abelian factor) and matter coupling, the absence of gauge anomalies can be established at the one-loop level. This proceeds by relating the gauge anomaly to perturbative agreement, which formalizes background independence.

1 Introduction

In its most general form, the Adler-Bardeen theorem states that in renormalizable Yang-Mills gauge theory, gauge anomalies are absent if they are absent at the one-loop, or $\mathcal{O}(\hbar)$, level.¹ It has been argued for at various levels of rigour and generality [5, 6, 7, 8, 9]. With the exception of [9], the Callan-Symanzik equation [10, 11], and the quantum action principle [12, 13] are employed.² However, the arguments put forward are not completely satisfactory: They are based on a particular (vacuum) state, while in generic backgrounds (of gravitational and/or gauge type), there is no preferred state. Furthermore, the generating functionals used in these approaches are formal, as they are integrated over the entire spacetime, leading to potential infrared issues. Finally, the “algebraic” approaches [7, 8] seem to apply only to the non-abelian anomaly, but not to abelian anomalies (which, for example, are most crucial to establish that anomaly freedom essentially fixes the matter representation in the Standard Model [14]).³

It is thus desirable to prove the Adler-Bardeen theorem in the rigorous framework of locally covariant field theory [15, 16, 17], cf. also [18, 19] for recent reviews or monographs. The author’s attempts to adapt the above mentioned approaches by replacing the Callan-Symanzik equation by renormalization group transformations as defined in [20] failed. However, it turns out that there is a close connection between gauge anomalies and anomalies of background independence, which one can exploit to prove the Adler-Bardeen theorem. Background independence is formalised in *perturbative agreement* [21, 22], which essentially states that quadratic terms can be equivalently included in the free or the interacting part of the action, see also [23] for a discussion. Typically, one thinks of these quadratic terms as arising from the free action by

¹The original statement [1], see also [2, 3], did actually not concern the gauge anomaly, but the chiral anomaly in QED. Recently, also a non-perturbative version of this was proven for lattice QED [4].

²By contrast, in [9] a particular, rather non-trivial, regularization scheme is employed in which the anomaly cancellation to all orders is argued to be manifest.

³Other approaches [5, 6, 9] also apply to some of the possible abelian anomalies (those listed in (45) below), but apparently not to general abelian anomalies, see also Remark 4 below.

variations of some of the background geometric data, such as the metric [21] or a background gauge connection [22]. For our purposes, it is the latter notion which is of relevance.

A first indication of a connection between gauge anomalies and perturbative agreement is the fact that the latter actually has the Adler-Bardeen property: Perturbative agreement can be fulfilled to all orders, if a certain obstruction, visible at $\mathcal{O}(\hbar)$, vanishes [21, 22]. Furthermore, the obstruction is, for the case of variations of a background gauge connection, the covariant divergence $T(\bar{\nabla}_\mu J_0^{\mu I})$ of the corresponding renormalized free current. Here $\bar{\nabla}$ indicates the covariant derivative w.r.t. the background gauge connection, $J_0^{\mu I} := \frac{\delta}{\delta \mathcal{A}_\mu^I} S_0$ is the functional derivative of the free part of the action w.r.t. the background gauge connection (with I a Lie algebra index), and T denotes a renormalization scheme (a time-ordered product in the terminology used below). In more formal approaches a non-vanishing $T(\bar{\nabla}_\mu J_0^{\mu I})$ is called a *covariant anomaly*, and it has (formally) a well-understood relation [24] to the *consistent anomaly*, which governs the gauge anomaly. Hence, in practice, when determining violations of perturbative agreement, one computes the well-known anomalies, just in covariant form.

The fact that an Adler-Bardeen type theorem has already been proven for perturbative agreement suggests the following strategy for proving the Adler-Bardeen theorem: When we assume vanishing of the covariant divergence $T(\bar{\nabla}_\mu J_0^{\mu I})$ of the renormalized free current (which is an $\mathcal{O}(\hbar)$ statement), we may also assume fulfillment of perturbative agreement w.r.t. changes in the background connection. These two properties turn out to be sufficient to prove absence of anomalies. To summarize, we prove the following:

Theorem 1. *Assume that, in a perturbatively renormalizable Yang-Mills gauge theory with a compact gauge group of at most a single $U(1)$ factor and arbitrary matter coupling on a globally hyperbolic four-dimensional spacetime, there exists a locally covariant renormalization scheme such that the covariant divergence $T(\bar{\nabla}_\mu J_0^{\mu I})$ of the free current vanishes (on-shell, when anti-fields are set to zero) for generic on-shell background gauge fields and vanishing background matter fields. Then there exists a locally covariant renormalization scheme in which gauge anomalies are absent for generic on-shell background gauge fields and vanishing background matter fields.*

The restriction to at most a single $U(1)$ factor in the gauge group ensures the absence of so-called exceptional anomalies [25]. This assumption can be relaxed when the occurrence of the exceptional anomalies can be excluded otherwise.

We note that the restriction to trivial matter background implies no restrictions on the matter action (except for the absence of a linear “source” term for the matter fields in the action), as the discussion is limited to the algebraic level, i.e., existence of states is not required. In particular, the vanishing matter background need not be stable (as long as it is a solution to the classical field equations). Nevertheless, in view of the Higgs mechanism in the Standard Model, an extension to non-trivial bosonic background matter fields is desirable, and will be sketched after establishing our main result.

Apart from connecting two up to now disparate concepts, our result also has the practical advantage that the non-vanishing of $\bar{\nabla}_\mu J_0^\mu$ is easy to check, as it can be reduced to studying violations of the Hadamard parametrix being a solution to the equation of motion, as in [26] for the case of chiral fermions in background gauge fields.

In the following, we will first recall some results of [17, 23] on gauge anomalies and background independence and prove some elementary lemmata. We will then, in Section 3, prove Theorem 1. In the concluding Section 4 we in particular sketch the generalization to non-trivial bosonic background matter fields.

2 Gauge anomalies and background independence

We consider a compact Lie group G with at most a single abelian factor, i.e., a quotient by a discrete subgroup of a product of simple groups and at most one $U(1)$, and denote its Lie algebra by \mathfrak{g} . On the latter, we choose an invariant negative definite symmetric bilinear form, which on the simple factors is a multiple of the Killing form. We consider principal G bundles P with base manifolds M , and denote by \mathfrak{p} the associated \mathfrak{g} vector bundle (whose sections would be “Lie-algebra valued functions” in physics parlance). On P , we consider connections \mathcal{A} whose curvatures F are sections of $\mathfrak{p} \otimes \Omega^2$ (“Lie-algebra valued two-forms”). Given a representation ρ of G , we consider matter fields ψ as sections of the associated vector bundle.⁴ The starting point of our considerations is then an action of the form

$$S = \frac{1}{4} \int F_{\mu\nu}^I F^{I\mu\nu} \text{vol} + S_m(\psi, \nabla\psi), \quad (1)$$

with vol the volume four-form, indices I labelling an appropriately normalized basis of \mathfrak{g} , and ∇ the covariant derivative w.r.t. the connection \mathcal{A} .

Given a background configuration $\bar{\mathcal{A}}$ of the gauge connection \mathcal{A} , we consider perturbations A , i.e., we write

$$\mathcal{A} = \bar{\mathcal{A}} + A \quad (2)$$

where A is a section of $\mathfrak{p} \otimes \Omega^1$ (a “Lie algebra valued one-form”). We then consider $\bar{\mathcal{A}}$ as a geometric datum on the same footing as the metric g (see [27] for the formulation of local covariance including background gauge connections). An important case is that of $\bar{\mathcal{A}}$ being on-shell, i.e., a configuration extremizing (1) with $\psi = 0$. Expanding S in A , the component S_{-1} which is of first order in A then drops out. We shall from now on restrict to that case and comment on deviations for off-shell backgrounds below. We denote by \mathcal{S} the “manifold” of on-shell backgrounds and by $T\mathcal{S}$ its tangent space, consisting of infinitesimal perturbations \bar{a} of $\bar{\mathcal{A}}$, which are solutions to the linearized equations of motion.⁶

The free part S_0 of the action, which is of second order in (A, ψ) , does not give rise to hyperbolic equations of motion, i.e., a well-posed initial value formulation. To remedy this, we follow the standard procedure (see, for example, [17, 19]) and first introduce Lagrange multipliers B , ghosts C , and anti-ghosts \tilde{C} , all of which are sections of \mathfrak{p} , and define the Batalin-Vilkovisky (BV) operator s by (the commutators are Lie algebra commutators)

$$sA_\mu^I = \bar{\nabla}_\mu C^I + [A_\mu, C]^I, \quad sC^I = -\frac{1}{2}[C, C]^I, \quad s\tilde{C}^I = B^I, \quad sB^I = 0, \quad s\psi = -\rho(C)\psi. \quad (3)$$

We collectively denote these fields by $\Phi = (A_\mu^I, B^I, C^I, \tilde{C}^I, \psi)$ and attribute a ghost number $(0, 0, 1, -1, 0)$ and mass dimension $(1, 2, 0, 2, *)$, with the mass dimension of the matter fields the canonical one (1 for scalars, $\frac{3}{2}$ for spin $\frac{1}{2}$ fermions). Then s increases the ghost number by one, but does not increase the mass dimension. The Grassmann parity of the fields is the same as their ghost number modulo 2, except for fermionic matter fields, which have Grassmann parity 1. The BV differential is nil-potent when graded w.r.t. the Grassmann parity.

⁴When spin $\frac{1}{2}$ matter fields are considered, M should also be equipped with a spin structure.

⁵The difference of any two principal bundle connections $\mathcal{A}, \mathcal{A}'$ can be naturally interpreted as a section of this bundle.

⁶Strictly speaking, the set of solutions to the Yang-Mills equation is a manifold only up to singular points corresponding to solutions possessing certain symmetries [28], i.e., the linearized equations around these special configurations have solutions which do not correspond to infinitesimal variations of the full solution. This does not affect our considerations, as we work locally in configuration space so that we can avoid these singular configurations.

We also introduce anti-fields $\Phi^\dagger = (A^{\mu I\dagger}, B^{I\dagger}, C^{I\dagger}, \tilde{C}^{I\dagger}, \psi^\dagger)$, which we interpret as densities with ghost numbers $(-1, -1, -2, 0, -1)$. We supplement the action with a source term

$$S_{\text{source}} = - \int \sum_i s\Phi^i \Phi_i^\dagger, \quad (4)$$

with i labelling the different fields. With this definition

$$sF = (S, F) \quad (5)$$

for any local functional F not depending on anti-fields, with the anti-bracket of local functionals defined by

$$(F, G) := \int \left\{ \frac{\delta^R F}{\delta\Phi^i(x)} \frac{\delta^L G}{\delta\Phi_i^\dagger(x)} - \frac{\delta^R G}{\delta\Phi_i^\dagger(x)} \frac{\delta^L F}{\delta\Phi^i(x)} \right\}. \quad (6)$$

The anti-bracket fulfills a graded symmetry and a graded Jacobi identity, see [29], for example.

Gauge fixing is performed by choosing a gauge-fixing fermion Ψ , which we take to be of the form

$$\Psi = \int \tilde{C}^I (\bar{\nabla}^\mu A_\mu^I + \frac{1}{2} B^I) \text{vol}, \quad (7)$$

We then supplement the action by an additional gauge fixing term $s\Psi$ and extend the definition of s to functionals involving anti-fields by (5). The new action thus obtained (including the source and the gauge fixing term) is denoted by S and consists of a free part S_0 containing all terms of second order in (anti-) fields, and an interacting part S_{int} containing all terms of higher order in (anti-) fields.

The original action depended on $\bar{\mathcal{A}}$, A only through the combination $\bar{\mathcal{A}} + A$, and was in this sense background independent. Analogously, we may say that a classical local functional F is background independent, if it only depends on this combination, or in other words, if for any $\bar{a} \in T\mathcal{S}$,

$$\mathcal{D}_{\bar{a}} F := \bar{\delta}_{\bar{a}} F - \delta_{\bar{a}} F = 0, \quad (8)$$

where $\bar{\delta}_{\bar{a}}$ denotes the functional derivative w.r.t. the background field $\bar{\mathcal{A}}$ in the direction \bar{a} , defined as

$$\bar{\delta}_{\bar{a}} F[\bar{\mathcal{A}}, \Phi, \Phi^\dagger] := \frac{d}{d\tau} F[\bar{\mathcal{A}} + \tau\bar{a}, \Phi, \Phi^\dagger] \Big|_{\tau=0}, \quad (9)$$

while $\delta_{\bar{a}}$ denotes the functional derivative w.r.t. the dynamical field A , in the same direction. The gauge fixing destroys the background independence of the action, as we have the formal⁷ identity [23]

$$\bar{\delta}_{\bar{a}} S - \delta_{\bar{a}} S_{\text{int}} = s\bar{\delta}_{\bar{a}} \Psi. \quad (10)$$

The appropriate replacement of the condition (8) for a classical local functional F to be background independent is then [23]

$$\hat{\mathcal{D}}_{\bar{a}} F := \mathcal{D}_{\bar{a}} F - (F, \mathcal{D}_{\bar{a}} \Psi) = 0. \quad (11)$$

From now on, we think of \bar{a} not as a vector, i.e., an element of $T\mathcal{S}$, but as a vector field, i.e., a section of $T\mathcal{S}$. The operator $\hat{\mathcal{D}}$ then fulfills

$$\hat{\mathcal{D}}_{\bar{a}}(F_1, F_2) = (\hat{\mathcal{D}}_{\bar{a}} F_1, F_2) + (F_1, \hat{\mathcal{D}}_{\bar{a}} F_2), \quad (12)$$

$$s\hat{\mathcal{D}}_{\bar{a}} F = \hat{\mathcal{D}}_{\bar{a}} sF, \quad (13)$$

$$\hat{\mathcal{D}}_{\bar{a}} \hat{\mathcal{D}}_{\bar{a}'} F - \hat{\mathcal{D}}_{\bar{a}'} \hat{\mathcal{D}}_{\bar{a}} F = \hat{\mathcal{D}}_{[\bar{a}, \bar{a}']} F, \quad (14)$$

⁷This is only formal as the integrals will in general not be convergent. This will be remedied below, in (53), after we introduce an adiabatic cutoff for the interaction.

where $[\cdot, \cdot]$ denotes the Lie bracket of vector fields. In view of the last equality, it is natural to interpret $\hat{\mathcal{D}}_{\bar{a}}F = (\hat{\mathcal{D}}F)(\bar{a})$ as the evaluation of the “background variation one-form” $\hat{\mathcal{D}}F$ in the vector field \bar{a} . We can thus promote $\hat{\mathcal{D}}$ to a differential, which, by (13), anticommutes with s (if the grading of background variation forms is taken into account). Put differently, one treats the perturbation \bar{a} as a new fermionic field variable.⁸ This is discussed in more detail in Section 3 below.

From the free part S_0 of the action, one obtains hyperbolic equations of motions for the fields Φ , giving rise to a causal propagator Δ being the difference of retarded and advanced propagator. One then defines a \star product on functionals⁹ such that

$$\Phi_i(x) \star \Phi_j(x') - (-1)^{\varepsilon_i \varepsilon_j} \Phi_j(x') \star \Phi_i(x) = i\hbar \Delta_{ij}(x, x'), \quad (15)$$

with ε_i the Grassmann parity of Φ_i . The anti-fields (anti-) commute with all other fields w.r.t. the \star product. The \star product preserves the grading $\text{Deg} = 2 \text{deg}_{\hbar} + \text{deg}_{\Phi(\ddagger)}$ with deg_{\hbar} counting the factors of \hbar and $\text{deg}_{\Phi(\ddagger)}$ the total number of fields and anti-fields. The Deg grading can be used to make sense of the series typically occurring in perturbation theory, i.e., equations relating such series can be understood as equations at each order in Deg.¹⁰

As in [16, 17], one defines local and covariant time-ordered products $T(F_1 \otimes \cdots \otimes F_n)$ of local functionals F_j , which can be inductively fixed up to the total diagonal by their defining property of causal factorization, i.e.,

$$T(F_1 \otimes \cdots \otimes F_n) = T(F_1 \otimes \cdots \otimes F_k) \star T(F_{k+1} \otimes \cdots \otimes F_n) \quad (16)$$

whenever $\text{supp } F_i$ does not intersect the causal past of $\text{supp } F_j$ for all $1 \leq i \leq k$ and $k+1 \leq j \leq n$, and which in particular are graded symmetric and fulfill field independence

$$\frac{\delta}{\delta \Phi(x)} T(F_1 \otimes \cdots \otimes F_n) = \sum_{k=1}^n T(F_1 \otimes \cdots \otimes \frac{\delta}{\delta \Phi(x)} F_k \otimes \cdots \otimes F_n). \quad (17)$$

Furthermore, they respect the Deg grading and can be assumed to act trivially on anti-fields.¹¹

$$T(\Phi^\ddagger(x) \mathcal{O}(x) \otimes F_1 \otimes \cdots \otimes F_n) = \Phi^\ddagger(x) T(\mathcal{O}(x) \otimes F_1 \otimes \cdots \otimes F_n). \quad (18)$$

Time-ordered products are not unique, but are subject to renormalization ambiguities encoded in the *main theorem of renormalization*, see for example [20], which is most conveniently formulated in terms of generating functionals

$$T(e^{\frac{i}{\hbar} F}) := 1 + \sum_{n=1}^{\infty} \frac{(\frac{i}{\hbar})^n}{n!} T(F^{\otimes n}). \quad (19)$$

It states that any two time ordered products T, T' are related by

$$T'(e^{\frac{i}{\hbar} F}) = T(e^{\frac{i}{\hbar} (F + D(e^{\frac{i}{\hbar} F})_D)}) \quad (20)$$

where $D(F_1 \otimes \cdots \otimes F_n)$ are local functionals (supported on the total diagonal of the F_k 's), and at least of $\mathcal{O}(\hbar)$. They inherit many of the properties of time-ordered products, such as field

⁸A similar approach was used in [30], see also Rem. 3.5 of [23] for further references.

⁹Technically, one restricts to *microcausal functionals*, which contain the local functionals and close under the \star product [15].

¹⁰This grading also naturally occurs in the context of deformation quantization, cf. [31].

¹¹This assumption is convenient, but not essential, in the proof of Prop. 7 below.

independence, i.e., (17) also holds for T symbols replaced by D symbols. With respect to the Deg grading, they fulfill

$$\text{Deg } D(F_1 \otimes \cdots \otimes F_n) = \sum_{i=1}^n \text{Deg}(F_i) - 2(n-1). \quad (21)$$

It is sometimes also convenient to consider connected time-ordered products, which can be implicitly defined by [32]

$$T(e_{\otimes}^{\frac{i}{\hbar}F}) = \exp_{\star}(\frac{i}{\hbar}T_c(e_{\otimes}^F)), \quad (22)$$

where

$$\exp_{\star}(G) = 1 + \sum_{n=1}^{\infty} \underbrace{G \star \cdots \star G}_{n \text{ factors}}. \quad (23)$$

One advantage of connected time-ordered products is that they are formal power series in \hbar (in contrast to $T(e_{\otimes}^{\frac{i}{\hbar}F})$). With respect to the Deg grading, they fulfill the same relation (21) as the renormalization map D .

The free BV differential s_0 is the linear part of s , i.e., it acts on fields as

$$s_0 A_{\mu}^I = \bar{\nabla}_{\mu} C^I, \quad s_0 C^I = 0, \quad s_0 \tilde{C}^I = B^I, \quad s_0 B^I = 0, \quad s_0 \psi = 0, \quad (24)$$

and more generally on local functionals by $s_0 F = (S_0, F)$. Like s , it is nilpotent. The incompatibility of time-ordered products with s_0 is encoded in the anomaly, through the *anomalous Ward identity* [17]:

Theorem 2. *We have*¹²

$$s_0 T_c(e_{\otimes}^F) = T_c(\{s_0 F + \frac{1}{2}(F, F) + A(e_{\otimes}^F)\} \otimes e_{\otimes}^F). \quad (25)$$

The anomaly A maps tensor products of local functionals linearly to local functionals. It increases the ghost number by one and is at least of $\mathcal{O}(\hbar)$. It is subject to the consistency condition

$$s_0 A(e_{\otimes}^F) + (F, A(e_{\otimes}^F)) + A(\{s_0 F + \frac{1}{2}(F, F) + A(e_{\otimes}^F)\} \otimes e_{\otimes}^F) = 0. \quad (26)$$

In particular, if $A^m(e_{\otimes}^F)$ is the first non-trivial term in the \hbar expansion of $A(e_{\otimes}^F)$, then

$$s_0 A^m(e_{\otimes}^F) + (F, A^m(e_{\otimes}^F)) + A^m(\{s_0 F + \frac{1}{2}(F, F)\} \otimes e_{\otimes}^F) = 0. \quad (27)$$

If T and T' are two renormalization schemes related by (20), the corresponding anomalies A and A' are related by

$$\begin{aligned} s_0 D(e_{\otimes}^F) + (F, D(e_{\otimes}^F)) + \frac{1}{2}(D(e_{\otimes}^F), D(e_{\otimes}^F)) + A(e_{\otimes}^{F+D(e_{\otimes}^F)}) \\ = D(\{s_0 F + \frac{1}{2}(F, F) + A'(e_{\otimes}^F)\} \otimes e_{\otimes}^F) + A'(e_{\otimes}^F). \end{aligned} \quad (28)$$

The anomaly A inherits many of the properties of time-ordered products. In particular, it is field (and anti-field) independent,

$$\frac{\delta}{\delta\Phi(x)} A(F_1 \otimes \cdots \otimes F_n) = \sum_{k=1}^n A(F_1 \otimes \cdots \otimes \frac{\delta}{\delta\Phi(x)} F_k \otimes \cdots \otimes F_n), \quad (29)$$

¹²In [17] the anomaly is defined via time-ordered products instead of connected time-ordered products. These definitions can easily be seen to be equivalent.

vanishes if one of the factors is a linear field [23], and fulfills the same relation (21) to the Deg grading as the renormalization map.

In order to define the algebra of interacting fields for a space-time region \mathcal{R} , one first introduces a compactly supported adiabatic cutoff $\lambda(x)$ with $\lambda|_{\mathcal{R}} = 1$ in the interacting part S_{int} of the action. We do this in such a way that cubic terms of the Lagrangian are multiplied with $\lambda(x)$ while the quartic ones are multiplied with $\lambda^2(x)$. The generating functional for interacting local functionals F supported inside \mathcal{R} is then (the inverse on the r.h.s. is w.r.t. \star)

$$T^{\text{int}}(e^{\frac{i}{\hbar}F}) = T(e^{\frac{i}{\hbar}S_{\text{int}}})^{-1} \star T(e^{\frac{i}{\hbar}F} \otimes e^{\frac{i}{\hbar}S_{\text{int}}}), \quad (30)$$

and these generate the interacting algebra for the space-time region \mathcal{R} . The algebra thus obtained is independent on the adiabatic cutoff λ (up to unitary equivalence) [33].

The interacting fields thus obtained are in general not gauge invariant, i.e., observables. In the classical theory, the gauge invariant, on-shell observables are given by the cohomology of s at vanishing ghost number. In the quantum theory, there are quantum corrections due to anomalies. The *quantum BV differential* [17, 29, 32]

$$qF := sF + A(F \otimes e^{\frac{i}{\hbar}S_{\text{int}}}), \quad (31)$$

turns out to be nil-potent, i.e., a proper differential, in case that

$$\text{supp } A(e^{\frac{i}{\hbar}S_{\text{int}}}) \cap \mathcal{R} = \emptyset, \quad (32)$$

i.e., the *gauge anomaly* $A(e^{\frac{i}{\hbar}S_{\text{int}}})$ vanishes in the region where the interacting observables are considered.¹³ In this case, we say that gauge anomalies are absent. Then the cohomology of q determines the interacting local observables in the quantum theory. It is thus crucial to understand the potential obstructions to achieving (32).

The gauge anomaly $A(e^{\frac{i}{\hbar}S_{\text{int}}})$ is a local functional, which can be expressed as

$$\sum_{n=1}^{\infty} \sum_{\alpha} \int a_{\alpha_1 \dots \alpha_n}^n \nabla^{\alpha_1} \lambda \dots \nabla^{\alpha_n} \lambda \text{vol}, \quad (33)$$

with α_k multiindices and $a_{\alpha_1 \dots \alpha_n}^n$ determined locally and covariantly out of (anti-) fields and background fields and symmetric under interchange $\alpha_i \leftrightarrow \alpha_j$. The above expression can be understood in the sense of a formal power series in λ . However, with our choice of adiabatic cutoff in S_{int} , the expansion in powers of λ is equivalent to an expansion in the Deg grading, as

$$\text{Deg}(a_{\alpha_1 \dots \alpha_n}^n) = n + 2. \quad (34)$$

We have the following:

Lemma 3. *An expression of the form (33) vanishes for all compactly supported λ only if, for all n ,*

$$a_{\emptyset \dots \emptyset}^n = \nabla^{\mu} b_{\mu}^n \quad (35)$$

for some b_{μ}^n . Hence, an expression of the form (33) determines a four-form

$$a = \sum_{n=1}^{\infty} a_{\emptyset \dots \emptyset}^n \text{vol} \quad (36)$$

up to a total derivative, i.e., modulo d , in the following sense: Two expressions of the form (33) coincide for all compactly supported λ if and only if the corresponding four-forms a defined by (36) coincide modulo an exact form.

¹³In [17, 29, 32], it is required that $A(e^{\frac{i}{\hbar}S_{\text{int}}}) = 0$, but it is easy to see that (32) is sufficient.

Proof. If (33) vanishes for all compactly supported λ , then the functional derivative w.r.t. $\lambda(x)$ must vanish, i.e.,

$$\sum_{n=1}^{\infty} \sum_{\alpha} \sum_{i=1}^n (-1)^{|\alpha_i|} \nabla^{\bar{\alpha}_i} (a_{\alpha_1 \dots \alpha_n}^n \nabla^{\alpha_1} \lambda \dots \nabla^{\alpha_i} \lambda \dots \nabla^{\alpha_n} \lambda) (x) = 0, \quad (37)$$

where $\bar{\alpha}_i$ is the multi-index in reverse order. This must hold order by order in the Deg grading, so we can restrict to a fixed n . Choosing $\lambda = 1$ near x , we obtain

$$\sum_k (-1)^k \nabla^{\mu_k \dots \mu_1} a_{(\mu_1 \dots \mu_k) \emptyset \dots \emptyset}^n = 0. \quad (38)$$

All but the first ($k = 0$) term on the l.h.s. are total derivatives, proving the statement. \square

Expressions of the form (33) with coefficients $a_{\alpha_1 \dots \alpha_n}^n$ fulfilling the relation (34), possibly shifted by an additive constant, will frequently occur, not just for the gauge anomaly. Also to such expressions we can associate a four form (up to a total derivative). We denote this association by $\cdot|_{\lambda=1}$, i.e., in the above case

$$\sum_{n=1}^{\infty} \sum_{\alpha} \int a_{\alpha_1 \dots \alpha_n}^n \nabla^{\alpha_1} \lambda \dots \nabla^{\alpha_n} \lambda \text{vol} \Big|_{\lambda=1} = a. \quad (39)$$

We call two expressions F , G , of the form (33) equivalent, $F \sim G$, if $F|_{\lambda=1} = G|_{\lambda=1} \pmod{\mathfrak{d}}$ (this is similar to the equivalence of “generalized Lagrangians” used in [34]). We can thus reformulate the condition (32) as

$$A(e_{\otimes}^{S_{\text{int}}}) \sim 0. \quad (40)$$

Now denote by $A^k(e_{\otimes}^{S_{\text{int}}})$ the $\mathcal{O}(\hbar^k)$ component of $A(e_{\otimes}^{S_{\text{int}}})$ and assume that $A^k(e_{\otimes}^{S_{\text{int}}}) \sim 0$ for all $k < m$, while $A^m(e_{\otimes}^{S_{\text{int}}}) \not\sim 0$, i.e., $A^m(e_{\otimes}^{S_{\text{int}}})$ is the first non-trivial term in the \hbar expansion of $A(e_{\otimes}^{S_{\text{int}}})$. Denote by $a^m := A^m(e_{\otimes}^{S_{\text{int}}})|_{\lambda=1}$ the corresponding four-form according to (36). Then, using that¹⁴

$$s_0 S_{\text{int}} + \frac{1}{2} (S_{\text{int}}, S_{\text{int}}) \sim 0, \quad (41)$$

it follows from (27) that sa^m is s closed modulo \mathfrak{d} , i.e.,

$$sa^m = \mathfrak{d}b^m. \quad (42)$$

On the other hand, if a^m is s exact modulo \mathfrak{d} , i.e., $a^m = sc^m + \mathfrak{d}d^m$, then one can perform a redefinition of time-ordered products as in (20) to obtain $a^m = 0$ [17], see also (54) below. Proceeding inductively in the order of \hbar , one obtains absence of gauge anomaly, in case that each a^m is s exact modulo \mathfrak{d} . Hence, the potential obstructions to achieve (40) are characterized by the cohomology $H_4^1(s|\mathfrak{d})$ of local and covariant four-forms of ghost number 1 and mass dimension 4 (or less, if the matter action contains terms of mass dimension less than 4). The most general such element of $H_4^1(s|\mathfrak{d})$ is a linear combination of the *non-abelian anomaly*

$$\text{Tr}_N \left[C \left(\bar{F}\bar{F} + \frac{1}{2} (\bar{\nabla} A \bar{F} + \bar{F} \bar{\nabla} A) + \frac{1}{6} (A^2 \bar{F} - 2A \bar{F} A + \bar{F} A^2) \right. \right. \\ \left. \left. + \frac{1}{3} \bar{\nabla} A \bar{\nabla} A + \frac{1}{6} (\bar{\nabla} A A^2 - A \bar{\nabla} A A + A^2 \bar{\nabla} A) \right) \right], \quad (43)$$

¹⁴The nilpotency of s and s_0 leads, via the Jacobi identity for the antibracket, to the formal identity $2s_0 S_{\text{int}} + (S_{\text{int}}, S_{\text{int}}) = 0$. Including the adiabatic cutoff in S_{int} remedies the formality, but introduces violation terms involving $\nabla\lambda$, so that the equality is reduced to the \sim equivalence.

with Tr_N an invariant trace over one of the non-abelian factors and $\bar{\nabla}$ the covariant differential¹⁵ defined by $\bar{\nabla}A = \bar{\nabla}_\mu A_\nu dx^\mu dx^\nu$, and an *abelian anomaly*

$$C^A G^A \quad (44)$$

with C^A the abelian ghost and $G^A \in H_4^0(s)$, cf. [25, 35]. For the later considerations, it is important that a representer G^A of a nontrivial cohomology class in $H_4^0(s)$ can be chosen such that if G_i^A is the lowest non-trivial component in a filtration w.r.t. the total (anti-)field number, then G_i^A is a non-trivial element of $H_4^0(s_0)$, so in particular not s_0 exact, see [36], Prop. 5.6. Furthermore, the restriction of G_i^A to the component of vanishing anti-field number does not vanish, i.e. $G_i^A|_{\Phi^\ddagger=0} \neq 0$, [25], Thm 7.1. It follows that $G_i^A \not\approx_0 0$, where \approx_0 denotes equality modulo the free equations of motions (generated by S_0), when anti-fields are set to zero. For if $G_i^A \approx_0 0$, then $G_i^A|_{\Phi^\ddagger=0} = (s_0 H_i)|_{\Phi^\ddagger=0}$ for some H_i . Then $G_i'^A := G_i^A - s_0 H_i$ is in the same equivalence class of $H_4^0(s_0)$ as G_i^A , but has $G_i'^A|_{\Phi^\ddagger=0} = 0$.

Remark 4. *Examples for abelian anomalies are*

$$G^A = F^A F^A, \quad G^A = \text{Tr}_N FF, \quad G^A = \text{Tr}_{so(3,1)} RR, \quad (45)$$

with R the Riemann curvature tensor, interpreted as an $so(3,1)$ valued two-form. These are relevant in particular for the Standard Model (the vanishing of the corresponding coefficient at $\mathcal{O}(\hbar)$ essentially fixes the hypercharges of quark and leptons [14]). Other potential abelian anomalies would be

$$G^A = F_{\mu\nu}^A F^{A\mu\nu} \text{vol}, \quad G^A = \text{Tr}_N F_{\mu\nu} F^{\mu\nu} \text{vol}, \quad G^A = R_{\mu\nu\lambda\rho} R^{\mu\nu\lambda\rho} \text{vol}, \quad G^A = \langle \varphi, \varphi \rangle_V^2 \text{vol}. \quad (46)$$

Here V is the representation space of the bosonic matter field φ and $\langle \cdot, \cdot \rangle_V$ a G invariant hermitean inner product. Obviously, there are many possible variants of the last term, for example involving the fermionic matter fields or derivatives. Such anomalies have, up to now, not occurred in perturbative calculations and are scarcely discussed in the literature (an exception is [37]). However, we are not aware of a general argument excluding such anomalies.¹⁶ We intend to revisit this topic in the future. In any case, the proof of our main theorem given below does apply to all potential abelian anomalies.

In order to formulate the requirement of perturbative agreement, we have to relax the condition that the background is on-shell, so that we are able to consider infinitesimal changes of the background connection which are compactly supported (and not necessarily pure gauge). We can simply do this by extending the action $S_0 + S_{\text{int}}$ to off-shell backgrounds.¹⁷ The free BV differential s_0 will then no longer be nil-potent, except when acting on functionals supported in the region where the background is on-shell. We denote by a a compactly supported section of $\mathfrak{p} \otimes \Omega^1$, representing an infinitesimal variation of the background connection. Perturbative agreement w.r.t. changes in the background gauge now means that, for any such a ,

$$\delta_a^r T(e_{\otimes}^{\frac{i}{\hbar}F}) = T(\frac{i}{\hbar}\{\bar{\delta}_a F + \bar{\delta}_a S_0\} \otimes e_{\otimes}^{\frac{i}{\hbar}F}) - T(\frac{i}{\hbar}\bar{\delta}_a S_0) \star T(e_{\otimes}^{\frac{i}{\hbar}F}). \quad (47)$$

Here δ_a^r is the *retarded variation* w.r.t. the infinitesimal background perturbation a (as it is not used in the proof of our main statement, we refrain from stating the precise definition, which

¹⁵It is not a proper differential, as it squares to the commutator with the curvature two-form \bar{F} of the background connection.

¹⁶Note that the anomalies in (45) have parity opposite to those in (46). If there is a general argument ruling out anomalies of the form (46), this may be expected to be important.

¹⁷Note that this is not the same as the expansion of the original action around off-shell backgrounds, as this would lead to a term linear in the dynamical fields, which we do not consider.

can be found for example in [23]). The quantity $J_0(a) = \bar{\delta}_a S_0$ can be seen as the free current J_0 smeared with the compactly supported Lie-algebra valued one-form a . By induction in the total number of fields, one can show [21, 22] that perturbative agreement can be fulfilled provided that

$$E(a_1, a_2) := \delta_{a_1}^r T(J_0(a_2)) - \delta_{a_2}^r T(J_0(a_1)) + \frac{i}{\hbar} [T(J_0(a_1)), T(J_0(a_2))]_* = 0, \quad (48)$$

a condition which is in fact closely related to the Wess-Zumino consistency condition [38], see [39]. In dimensions $D \leq 4$, this condition can in turn be fulfilled¹⁸ [22] provided that the covariant divergence of the free current vanishes,

$$T(J_0(\bar{\nabla}\Lambda)) \approx_0 0, \quad (49)$$

for any compactly supported section Λ of \mathfrak{p} (a ‘‘Lie algebra valued function’’). Here \approx_0 denotes equality modulo the free equations of motion, when anti-fields are set to zero. This condition can be rather straightforwardly checked in concrete cases, cf. [26] for the case of chiral fermions in gauge backgrounds. The calculation boils down to determining the violation of the Hadamard parametrix to being a solution of the free field equation, a calculation which actually does not involve any loop integrals. For our purposes the following is crucial [22, 23]:

Theorem 5. *If (49) and perturbative agreement w.r.t. changes in the background gauge connection is fulfilled, then*

$$T(\bar{\delta}_{\bar{\nabla}\Lambda} S \otimes e_{\otimes}^{\frac{i}{\hbar} S_{\text{int}}}) \approx_0 0, \quad (50)$$

$$\bar{\delta}_{\bar{a}} A(e_{\otimes}^F) = A(\{\bar{\delta}_{\lambda\bar{a}} S_0 + \bar{\delta}_{\bar{a}} F\} \otimes e_{\otimes}^F). \quad (51)$$

In (50), Λ is supported in the region where $\lambda = 1$ and \approx_0 denotes equality modulo the free equations of motion (derived from S_0), when anti-fields are set to zero. In (51), F is supported in the region where $\lambda = 1$, and \bar{a} is again a solution to the linearized equations of motions, i.e., an infinitesimal variation of an on-shell background connection.

Note that in (51), we cut off the (not necessarily compactly supported) infinitesimal variation \bar{a} of the free action S_0 by multiplication with the adiabatic cut-off λ . This is necessary to have the expression well-defined, but obviously the choice of λ is irrelevant by the locality of the anomaly and the restriction on the support of F .¹⁹ Later, we will apply (51) to the case $F = S_{\text{int}}$, whose support is not restricted to the region where $\lambda = 1$. It follows that there will be supplementary terms supported in $\text{supp } d\lambda$, i.e., the equality in (51) is replaced by the equivalence \sim as defined below Lemma 3.

In the following, we will need to perform redefinitions of time-ordered products, but have to preserve perturbative agreement. For this, the following is essential (we omit the straightforward proof):

Lemma 6. *If time-ordered products T fulfill perturbative agreement, then the time-ordered products T' defined by (20) also fulfill perturbative agreement if and only if*

$$\bar{\delta}_a D(e_{\otimes}^F) = D(\{\bar{\delta}_a F + \bar{\delta}_a S_0\} \otimes e_{\otimes}^F). \quad (52)$$

Finally, we note that in the presence of an adiabatic cutoff λ in the interaction, the identity (10) can be stated as follows:

$$\bar{\delta}_{\lambda\bar{a}} S - \bar{\delta}_{\bar{a}} S_{\text{int}} - (S, \bar{\delta}_{\lambda\bar{a}} \Psi) \sim 0. \quad (53)$$

¹⁸Using the explicit form of δ_a^r given in [23] one can easily check that (18) can be preserved under the necessary redefinitions of time-ordered products.

¹⁹Note that $A(\bar{\delta}_{\lambda\bar{a}} S_0) = 0$, as by the Deg grading rule the result must be a c number, which is not possible at ghost number 1.

Note the presence of the adiabatic cut-off in all functional derivatives w.r.t. background fields. It ensures the well-definedness of these expressions, but also that in all terms the same (affine) linear relation between the Deg grading and the order in λ holds. Hence, the l.h.s. is of the form (33), now with $\text{Deg}(a_{\alpha_1 \dots \alpha_n}^n) = n + 1$ instead of (34), so that Lemma 3 is applicable.

3 Proof of Theorem 1

As discussed above, if the conditions of Theorem 1 are met, we can assume that perturbative agreement holds. We begin by arguing that trivial anomalies can be removed by redefinitions of time-ordered products. This is not completely trivial, as we want to ensure that also the new time-ordered products fulfill perturbative agreement. Usually, one would argue as follows: Assume m is the lowest order in \hbar at which the anomaly a^m is not d exact. If the anomaly is trivial in $H(s|d)$, i.e., $a^m = sb^m + dc^m$, then one performs a redefinition of the form (20) such that

$$D(e_{\otimes}^{S_{\text{int}}})|_{\lambda=1} = \frac{1}{n!} D(S_1^{\otimes n})|_{\lambda=1} = -b^m. \quad (54)$$

Here S_1 is the part of S_{int} which is cubic in (anti-) fields and $n = 2m - 2 + k$, with k the total number of (anti-) fields in b^m (the latter condition ensures that the redefined time-ordered products respect the Deg grading).²⁰ By (28), the anomaly at $\mathcal{O}(\hbar^m)$ is then removed for the modified time-ordered products. If we are to preserve perturbative agreement in the process, we must then also have, by Lemma 6, (all equalities modulo d)

$$-\bar{\delta}_{\bar{a}} b^m = D(\{\bar{\delta}_{\lambda\bar{a}} S_{\text{int}} + \bar{\delta}_{\lambda\bar{a}} S_0\} \otimes e_{\otimes}^{S_{\text{int}}})|_{\lambda=1} = D(\bar{\delta}_{\lambda\bar{a}} S \otimes e_{\otimes}^{S_{\text{int}}})|_{\lambda=1}. \quad (55)$$

On the other hand, from field independence of D and (53), we also have (again modulo d)

$$-\delta_{\bar{a}} b^m = D(\delta_{\bar{a}} S_{\text{int}} \otimes e_{\otimes}^{S_{\text{int}}})|_{\lambda=1} = D(\{\bar{\delta}_{\lambda\bar{a}} S - (S, \bar{\delta}_{\lambda\bar{a}} \Psi)\} \otimes e_{\otimes}^{S_{\text{int}}})|_{\lambda=1}. \quad (56)$$

These two equations imply (modulo d)

$$D((S, \bar{\delta}_{\lambda\bar{a}} \Psi) \otimes e_{\otimes}^{S_{\text{int}}})|_{\lambda=1} = -\mathcal{D}_{\bar{a}} b^m, \quad (57)$$

where we used the notation introduced in (8). If now $(S, \bar{\delta}_{\lambda\bar{a}} \Psi) \sim 0$ (which happens only if \bar{a} is a perturbation purely in the abelian direction, so that $\bar{\delta}_{\lambda\bar{a}} \Psi = 0$), then this would be inconsistent unless also $\mathcal{D}_{\bar{a}} b^m = 0$ modulo d. But this is indeed the case: If the conditions of Thm. 5 are met, then by (51), field independence (29) of the anomaly, and the relation (53), we have that

$$\mathcal{D}_{\bar{a}}^{\lambda} A^m(e_{\otimes}^{S_{\text{int}}}) - A^m((S, \bar{\delta}_{\lambda\bar{a}} \Psi) \otimes e_{\otimes}^{S_{\text{int}}}) \sim 0, \quad (58)$$

where we introduced $\mathcal{D}_{\bar{a}}^{\lambda} := \bar{\delta}_{\lambda\bar{a}} - \delta_{\bar{a}}$. Hence, $(S, \bar{\delta}_{\lambda\bar{a}} \Psi) \sim 0$ implies that $\mathcal{D}_{\bar{a}} a^m = 0$ modulo d. From this, it follows that b^m may be chosen such that $\mathcal{D}_{\bar{a}} b^m = 0$ modulo d, too.²¹ Thus,

²⁰In case b^m is not homogenous in the total number of (anti-) fields, one decomposes it and applies the above to each component.

²¹As noted above, $(S, \bar{\delta}_{\lambda\bar{a}} \Psi) \sim 0$ implies that \bar{a} is purely abelian, so that $\mathcal{D}_{\bar{a}} \Psi = -\int \tilde{C}^I \tilde{\nabla}^{\mu} \bar{a}_{\mu}^I \text{vol}$. Hence, $(F, \mathcal{D}_{\bar{a}} \Psi) = 0$ unless F contains \tilde{C}^{\dagger} . As the interaction S_{int} does not contain \tilde{C}^{\dagger} or B , the anomaly a^m can not contain these either, by (anti-) field independence (29) of the anomaly. Hence, we also have $\hat{\mathcal{D}}_{\bar{a}} a^m = 0$ modulo d. Given a solution b^m to $a^m = sb^m + dc^m$ at a particular background, we may extend b^m via ‘‘parallel transport’’ w.r.t. the flat ‘‘connection’’ \hat{D} to all backgrounds (recall (14)). Then $a^m = sb^m + dc^m$ is fulfilled on all backgrounds and $\hat{\mathcal{D}}_{\bar{a}} b^m = 0$. It remains to argue that the anti-bracket of b^m with $\mathcal{D}_{\bar{a}} \Psi$ vanishes. This is the case unless b^m contains \tilde{C}^{\dagger} , but by power counting and the fact that we are looking for a scalar (not a section of \mathfrak{p}), the only possibility for this would be $\text{Tr}(B\tilde{C}^{\dagger})$ with some invariant trace Tr . But then sb^m would contain B and not be a total derivative, a contradiction to $a^m = sb^m + dc^m$.

preservation of perturbative agreement does not lead to obstructions to the removal of trivial anomalies. Furthermore, by decomposing S_1 into monomials and choosing the combination of these on the l.h.s. of (54) such that total anti-field content matches that of b^m , we can make sure that condition (18) is still valid after the redefinition.

We may thus assume that at the first non-trivial order in \hbar the anomaly is non-trivial, i.e., a linear combination of the non-abelian anomaly (43) and an abelian anomaly of the form (44).

The above already showed that the abelian anomaly is background independent, i.e., $\mathcal{D}_{\bar{a}} a^m = 0$ (modulo d) for an abelian anomaly a^m , cf. (58). This is however not sufficient to rule it out, as there are many possible background independent abelian anomalies, for example the ones in (45) with F the full field strength (of $\mathcal{A} = \bar{\mathcal{A}} + A$). However, we can proceed differently:

Proposition 7. *If (49) and perturbative agreement w.r.t. changes in the background connection holds, and $a^m = A^m(e_{\otimes}^{S_{\text{int}}})|_{\lambda=1}$ is the lowest non-trivial term in an \hbar expansion of the gauge anomaly, then a^m can not contain a contribution from an abelian anomaly of the form (44).*

Proof. If a^m contains an abelian anomaly (44), then its coefficient G^A can be obtained by functional differentiation w.r.t. the abelian ghost,

$$G^A(x) = \frac{\delta}{\delta C^A(x)} A^m(e_{\otimes}^{S_{\text{int}}}) = A^m\left(\frac{\delta}{\delta C^A(x)} S_{\text{int}} \otimes e_{\otimes}^{S_{\text{int}}}\right), \quad (59)$$

where in the second step we used field independence of the anomaly. We may here and in the following restrict to x contained in the region where the cutoff λ equals 1. On the other hand, by the definition of the anomaly, we have

$$\begin{aligned} s_0 T_c\left(\frac{\delta}{\delta C^A(x)} S_{\text{int}} \otimes e_{\otimes}^{S_{\text{int}}}\right) &= T_c\left((S, \frac{\delta}{\delta C^A(x)} S_{\text{int}}) \otimes e_{\otimes}^{S_{\text{int}}}\right) + T_c\left(A\left(\frac{\delta}{\delta C^A(x)} S_{\text{int}} \otimes e_{\otimes}^{S_{\text{int}}}\right) \otimes e_{\otimes}^{S_{\text{int}}}\right) \\ &\quad - T_c\left(\frac{\delta}{\delta C^A(x)} S_{\text{int}} \otimes \left\{s_0 S_{\text{int}} + \frac{1}{2}(S_{\text{int}}, S_{\text{int}}) + A(e_{\otimes}^{S_{\text{int}}})\right\} \otimes e_{\otimes}^{S_{\text{int}}}\right). \end{aligned} \quad (60)$$

We will be interested in considering this equality modulo anti-fields and the free equations of motion. The only terms in S_{int} which depend on the abelian ghost are the source terms for the matter gauge transformations, i.e., schematically of the form $\int \rho(C)\psi\psi^\dagger$. Hence, $\frac{\delta}{\delta C^A(x)} S_{\text{int}}$ is linear in the matter anti-fields, and thus the last term on the r.h.s. of (60) vanishes when anti-fields are set to zero (recall (18), i.e., that anti-fields are not contracted in time-ordered products). By the same argument, the l.h.s. of (60) vanishes on-shell (w.r.t. to the free equations of motion) when anti-fields are set to zero (recall that the action of s_0 on anti-fields generates the free equations of motion). Regarding the first term on the r.h.s. of (60), we notice that, by the restriction of x to the region where λ equals 1,

$$\begin{aligned} (S, \frac{\delta}{\delta C^A(x)} S_{\text{int}}) &= (S_0, \frac{\delta}{\delta C^A(x)} S_{\text{int}}) - \frac{1}{2} \frac{\delta}{\delta C^A(x)} (S_{\text{int}}, S_{\text{int}}) \\ &= (S_0, \frac{\delta}{\delta C^A(x)} S_{\text{int}}) + \frac{\delta}{\delta C^A(x)} (S_0, S_{\text{int}}) \\ &= (\frac{\delta}{\delta C^A(x)} S_0, S_{\text{int}}). \end{aligned} \quad (61)$$

For convenience, we integrate against $\Lambda^A(x)$, i.e., we consider the functional derivative w.r.t. C^A in the direction Λ^A with Λ^A supported within the region where $\lambda = 1$. Then

$$\langle \frac{\delta}{\delta C} S_0, \Lambda \rangle = - \int (A^{\mu A \dagger} + \bar{\nabla}^\mu \bar{C}^A \text{vol}) \bar{\nabla}_\mu \Lambda^A. \quad (62)$$

and thus

$$\langle \langle \frac{\delta}{\delta C} S_0, \Lambda \rangle, S_{\text{int}} \rangle = -\delta_{\bar{\nabla}\Lambda} S_{\text{int}} = -\bar{\delta}_{\bar{\nabla}\Lambda} S, \quad (63)$$

where in the last step, we used (53) and the fact that the gauge fixing fermion does not depend on the abelian background gauge field. From (50), it then follows that the first term on the r.h.s. of (60) vanishes on-shell w.r.t. the free equations of motion, when anti-fields are set to zero. Hence, we have shown that

$$T_c(G^A(x) \otimes e_{\otimes}^{S_{\text{int}}}) \approx_0 0. \quad (64)$$

Now assume that there is a non-trivial abelian anomaly $C^A G^A$ at the leading order \hbar^m . As the connected time-ordered products are formal power series in \hbar , the $\mathcal{O}(\hbar^m)$ component of the l.h.s. can be expressed as $T_c^m(G^A(x) \otimes e_{\otimes}^{S_{\text{int}}})$, where the superscript m denotes the restriction to $\mathcal{O}(\hbar^m)$. As each factor S_{int} increases the total Deg of $T_c^m(G^A(x) \otimes e_{\otimes}^{S_{\text{int}}})$ at least by 1 (this follows from (21) and $\text{Deg } S_{\text{int}} \geq 3$), the lowest non-vanishing term in a filtration of $T_c^m(G^A(x) \otimes e_{\otimes}^{S_{\text{int}}})$ w.r.t. the total (anti-) field number is $G_i^A(x)$ in the notation introduced below (44). Hence, we have shown that $G_i^A \approx_0 0$. But, as argued below (44), this is not possible for a non-trivial abelian anomaly. \square

It remains to rule out the non-abelian anomaly (43). As we already ruled out trivial and abelian anomalies, we can assume that the contribution a^m of lowest non-vanishing order in \hbar is of the form (43), so in particular anti-field independent. Then (58) can be rewritten as

$$\hat{D}_{\bar{a}}^\lambda A^m(e_{\otimes}^{S_{\text{int}}}) - A^m((S, \bar{\delta}_{\lambda\bar{a}}\Psi) \otimes e_{\otimes}^{S_{\text{int}}}) \sim 0, \quad (65)$$

where we used $\hat{D}_{\bar{a}}^\lambda := \bar{\delta}_{\lambda\bar{a}} - \delta_{\bar{a}} - (\cdot, \mathcal{D}_{\bar{a}}^\lambda \Psi)$. Considering this at lower order in \hbar , this in particular implies

$$A^k((S, \bar{\delta}_{\lambda\bar{a}}\Psi) \otimes e_{\otimes}^{S_{\text{int}}}) \sim 0 \quad \forall k < m. \quad (66)$$

We now use the following corollary of the \hbar expanded consistency condition (27):²²

Corollary 8. *Let $A^m(e_{\otimes}^{S_{\text{int}}})$ be the first non-trivial term (w.r.t. \sim) in the \hbar expansion of $A(e_{\otimes}^{S_{\text{int}}})$. If $A^k(F \otimes e_{\otimes}^{S_{\text{int}}})$, $k \leq m$ is the first non-trivial term in the \hbar expansion of $A(F \otimes e_{\otimes}^{S_{\text{int}}})$, then*

$$(S, A^k(F \otimes e_{\otimes}^{S_{\text{int}}})) + (F, A^k(e_{\otimes}^{S_{\text{int}}})) - A^k((S, F) \otimes e_{\otimes}^{S_{\text{int}}}) \sim 0. \quad (67)$$

Also, if $A^m(F_{1/2} \otimes e_{\otimes}^{S_{\text{int}}})$ are the first non-trivial terms in the \hbar expansion of $A(F_{1/2} \otimes e_{\otimes}^{S_{\text{int}}})$, and $A^k(F_1 \otimes F_2 \otimes e_{\otimes}^{S_{\text{int}}})$, $k \leq m$, is the first non-trivial term in the \hbar expansion of $A(F_1 \otimes F_2 \otimes e_{\otimes}^{S_{\text{int}}})$, then

$$(S, A^k(F_1 \otimes F_2 \otimes e_{\otimes}^{S_{\text{int}}})) + (F_1, A^k(F_2 \otimes e_{\otimes}^{S_{\text{int}}})) + (F_2, A^k(F_1 \otimes e_{\otimes}^{S_{\text{int}}})) \\ - A^k((S, F_1) \otimes F_2 \otimes e_{\otimes}^{S_{\text{int}}}) - A^k(F_1 \otimes (S, F_2) \otimes e_{\otimes}^{S_{\text{int}}}) \sim 0. \quad (68)$$

Here F_1, F_2 are assumed to be bosonic (otherwise the signs need to be adjusted).

From (66) and (67), we may thus conclude that $(S, A^k(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes e_{\otimes}^{S_{\text{int}}})) \sim 0$ for $k < m$. Hence, the four-form field $\xi_{\bar{a}}^k := A^k(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes e_{\otimes}^{S_{\text{int}}})|_{\lambda=1}$ (recall the definition below Lemma 3), which is of vanishing ghost number, mass dimension four, and linear in \bar{a} , is s closed modulo d . Furthermore, as Ψ is independent of the abelian part of the background connection, we can restrict to perturbations \bar{a} which are purely in the non-abelian directions. To such fields, the following applies:

Lemma 9. *Fields $\xi_{\bar{a}}$ of mass dimension four which are linear in a perturbation \bar{a} purely in the non-abelian directions are trivial in $H_4^0(s|d)$.*

²²For present purposes, only the result (67) is relevant. Later, also (68) will be used.

Proof. A general element $\xi_{\bar{a}} \in H_4^0(s|\mathfrak{d})$ is of the form $\xi_{\bar{a}} = \text{Tr}_N(\bar{a} \wedge *D)$ with D a section of $\mathfrak{p} \otimes \Omega^1$ of vanishing ghost number and mass dimension 3 (one mass dimension is contributed by \bar{a}) and $*$ the Hodge dual. Furthermore, $\xi_{\bar{a}}$ being s closed modulo \mathfrak{d} implies that $sD = P_{\text{lin}}E$, where P_{lin} is the linearized Yang-Mills operator (which vanishes on \bar{a} , so that $s\xi_{\bar{a}}$ is a total derivative) for some E of ghost number and mass dimension 1. It must thus be a linear combination of $\bar{\nabla}C^I$ and $[A, C]^I$. But as $P_{\text{lin}} \circ \bar{\nabla} = 0$, the first term can always be added to E^I , so unless it is trivial, we can write it as $E^I = c(\bar{\nabla}C + [A, C])^I = csA^I$ with a numerical coefficient c . Again by $P_{\text{lin}}\bar{a} = 0$, $\text{Tr}_N(\bar{a} \wedge *P_{\text{lin}}A)$ is a total derivative, so $\xi'_{\bar{a}} = \text{Tr}_N(\bar{a} \wedge *(D - cP_{\text{lin}}A))$ is in the same cohomology class as $\xi_{\bar{a}}$. However, the corresponding $D' = D - cP_{\text{lin}}A$ is now s closed: $sD' = P_{\text{lin}}(E - E) = 0$. We are thus looking for an s closed Lie algebra valued one-form D of vanishing ghost number and mass dimension 3. Hence, D must at the same time transform covariantly under background gauge transformations and be invariant under a gauge transformation of the dynamical fields. It follows from the well-known characterization of $H^0(s)$ that it must either be trivial, for example $D^I = s\bar{\nabla}\tilde{C}^I$, or a c -number. By power counting, the only possibility is $D^I_{\nu} = \bar{\nabla}^{\mu}\bar{F}_{\mu\nu}$, which however vanishes by our assumption that the background is on-shell. \square

Analogously to the removal of the gauge anomaly, we can thus, by redefinition of time-ordered products $T(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes e_{\otimes}^{\frac{i}{\hbar}S_{\text{int}}})$ (which does not spoil perturbative agreement), achieve that

$$A^k(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes e_{\otimes}^{S_{\text{int}}}) \sim 0 \quad \forall k < m. \quad (69)$$

Now as both $\bar{\delta}_{\lambda\bar{a}}\Psi$ and the anomaly a^m are anti-field independent, it follows from (66), (69), and (67) that

$$(S, A^m(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes e_{\otimes}^{S_{\text{int}}})) \sim A^m((S, \bar{\delta}_{\lambda\bar{a}}\Psi) \otimes e_{\otimes}^{S_{\text{int}}}). \quad (70)$$

In particular, with (65),

$$\hat{\mathcal{D}}_{\bar{a}}^{\lambda}A^m(e_{\otimes}^{S_{\text{int}}}) \sim (S, A^m(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes e_{\otimes}^{S_{\text{int}}}). \quad (71)$$

This implies that there must be fields $b_{\bar{a}}^m, c_{\bar{a}}^m$ such that

$$\hat{\mathcal{D}}_{\bar{a}}a^m = sb_{\bar{a}}^m + dc_{\bar{a}}^m. \quad (72)$$

As $\hat{\mathcal{D}}_{\bar{a}}$ commutes with \mathfrak{d} and s , cf. (13), it is well-defined on the cohomology $H(s|\mathfrak{d})$. The above shows that if the assumptions of Thm. 5 are fulfilled, then $\hat{\mathcal{D}}_{\bar{a}}a^m$ is trivial in $H^1(s|\mathfrak{d})$, i.e., a^m is background independent in cohomology. If the non-abelian anomaly (43) turned out to be not background independent in cohomology, this would already prove Thm. 1. However, as shown in [35], the non-abelian anomaly (43) fulfills (72), with $b_{\bar{a}}^m$ explicitly given by

$$b_{\bar{a}}^m = \text{Tr}_N \left[-\frac{1}{2}A(\bar{a}\bar{F} + \bar{F}\bar{a}) + \frac{1}{6}\bar{a}(A\bar{\nabla}A + \bar{\nabla}AA) + \frac{1}{12}A^2(\bar{a}A - A\bar{a}) \right]. \quad (73)$$

Hence, (72) can not yet be used to rule out the non-abelian anomaly (43). Nevertheless, we have learned something: By comparing (72) and (71) and recalling the definition of $\xi_{\bar{a}}^m$ given below Corollary 8, we have

$$\xi_{\bar{a}}^m = b_{\bar{a}}^m + d_{\bar{a}}^m \quad (74)$$

for some $d_{\bar{a}}^m \in H^0(s|\mathfrak{d})$ (in particular, this includes \mathfrak{d} exact terms). This ambiguity is due to the fact that (72) defines $b_{\bar{a}}^m$ only up to terms in the kernel of s (modulo \mathfrak{d}). Furthermore, a trivial element of $H^0(s|\mathfrak{d})$ can be removed from $\xi_{\bar{a}}^m$ by a redefinition of $T(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes e_{\otimes}^{\frac{i}{\hbar}S_{\text{int}}})$, which can be performed without destroying perturbative agreement. However, by Lemma 9, the relevant cohomology class is trivial, so that we can ignore the ambiguity $d_{\bar{a}}^m$ in (74).

We would now like to compute a further ‘‘background derivative’’, namely of (74). As already indicated below (14), we promote $\hat{\mathcal{D}}$ to a differential on the tangent space $T\mathcal{S}$ of background variations, and interpret (74) as an equality of background variation one-forms. To deal with background variation p -forms, we introduce some notation. A general background variation p -form field H can be expressed as

$$H_{I_1 \dots I_p}^{\mu_1 \dots \mu_p}(x) \hat{\mathcal{D}}\Xi_{\mu_1}^{I_1}(x) \wedge \dots \wedge \hat{\mathcal{D}}\Xi_{\mu_p}^{I_p}(x) \quad (75)$$

with I_k Lie algebra indices and $\hat{\mathcal{D}}\Xi$ dual to a tangent vector $\bar{a}_\mu^I(x)$, meaning that the evaluation of a tangent vector \bar{a} in $\hat{\mathcal{D}}\Xi$ is given by $\hat{\mathcal{D}}\Xi_\mu^I(x)(\bar{a}) = \bar{a}_\mu^I(x)$. On such an expression, the differential $\hat{\mathcal{D}}$ acts as

$$\hat{\mathcal{D}} \left(H_{I_1 \dots I_p}^{\mu_1 \dots \mu_p}(x) \hat{\mathcal{D}}\Xi_{\mu_1}^{I_1}(x) \wedge \dots \wedge \hat{\mathcal{D}}\Xi_{\mu_p}^{I_p}(x) \right) = (-1)^{\varepsilon_H} \hat{\mathcal{D}}_{I_0}^{\mu_0} H_{I_1 \dots I_p}^{\mu_1 \dots \mu_p}(x) \hat{\mathcal{D}}\Xi_{\mu_0}^{I_0}(x) \wedge \dots \wedge \hat{\mathcal{D}}\Xi_{\mu_p}^{I_p}(x), \quad (76)$$

where ε_H is the Grassmann parity of $H_{I_1 \dots I_p}^{\mu_1 \dots \mu_p}(x)$ and $\hat{\mathcal{D}}_I^\mu$ is defined by

$$\hat{\mathcal{D}}_{\bar{a}} H(x) = \hat{\mathcal{D}}_I^\mu H(x) \bar{a}_\mu^I(x). \quad (77)$$

By (14), (13), the differential $\hat{\mathcal{D}}$ is nilpotent and anticommutes with s . On expressions of the form (33), one similarly defines $\hat{\mathcal{D}}^\lambda$ (due to the presence of the cut-off λ this is a differential only w.r.t. the relation \sim).

In the notation just introduced, $A^m(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes e_\otimes^{S_{\text{int}}})$ can be seen as the evaluation of the background variation one form $A^m(\bar{\delta}_I^{\lambda\mu}\Psi \otimes e_\otimes^{S_{\text{int}}})\hat{\mathcal{D}}\Xi_\mu^I$ in \bar{a} (the superscript λ at $\bar{\delta}$ indicates multiplication with the cut-off λ). As the gauge fixing fermion is of second order in the dynamical fields and does not contain anti-fields, and the anomaly vanishes if one of the factors is a linear field, we may replace $\bar{\delta}_I^{\lambda\mu}\Psi$ by $\hat{\mathcal{D}}_I^{\lambda\mu}\Psi$ in this expression. We now compute a $\hat{\mathcal{D}}^\lambda$ differential of this background variation one-form. A further application of field independence (29), background independence (51), and (53) yields, for $k \leq m$,

$$\begin{aligned} \hat{\mathcal{D}}^\lambda \left(A^k(\bar{\delta}_I^{\lambda\mu}\Psi \otimes e_\otimes^{S_{\text{int}}})\hat{\mathcal{D}}\Xi_\mu^I \right) \\ \sim \left(A^k(\hat{\mathcal{D}}_{I_0}^{\lambda\mu_0}\hat{\mathcal{D}}_{I_1}^{\lambda\mu_1}\Psi \otimes e_\otimes^{S_{\text{int}}}) + A^k(\hat{\mathcal{D}}_{I_1}^{\lambda\mu_1}\Psi \otimes (S, \hat{\mathcal{D}}_{I_0}^{\lambda\mu_0}\Psi) \otimes e_\otimes^{S_{\text{int}}}) \right) \hat{\mathcal{D}}\Xi_{\mu_0}^{I_0} \wedge \hat{\mathcal{D}}\Xi_{\mu_1}^{I_1}. \end{aligned} \quad (78)$$

The first term on the r.h.s. vanishes (w.r.t. \sim), by nilpotency of $\hat{\mathcal{D}}^\lambda$. To deal with the second term, we note that, analogously to (66), (69), we have, for all $k < m$,²³

$$A^k(\bar{\delta}_{\lambda\bar{a}}\Psi \otimes \bar{\delta}_{\lambda\bar{a}'}\Psi \otimes e_\otimes^{S_{\text{int}}}) \sim 0, \quad A^k((S, \bar{\delta}_{\lambda\bar{a}}\Psi) \otimes \bar{\delta}_{\lambda\bar{a}'}\Psi \otimes e_\otimes^{S_{\text{int}}}) \sim 0. \quad (79)$$

Then, with (68), we have (recall that Ψ is fermionic)

$$\begin{aligned} (S, A^m(\hat{\mathcal{D}}_{I_1}^{\lambda\mu_1}\Psi \otimes \hat{\mathcal{D}}_{I_0}^{\lambda\mu_0}\Psi \otimes e_\otimes^{S_{\text{int}}})) \\ \sim A^m((S, \hat{\mathcal{D}}_{I_1}^{\lambda\mu_1}\Psi) \otimes \hat{\mathcal{D}}_{I_0}^{\lambda\mu_0}\Psi \otimes e_\otimes^{S_{\text{int}}}) - A^m(\hat{\mathcal{D}}_{I_1}^{\lambda\mu_1}\Psi \otimes (S, \hat{\mathcal{D}}_{I_0}^{\lambda\mu_0}\Psi) \otimes e_\otimes^{S_{\text{int}}}) \\ \sim A^m(\hat{\mathcal{D}}_{I_0}^{\lambda\mu_0}\Psi \otimes (S, \hat{\mathcal{D}}_{I_1}^{\lambda\mu_1}\Psi) \otimes e_\otimes^{S_{\text{int}}}) - A^m(\hat{\mathcal{D}}_{I_1}^{\lambda\mu_1}\Psi \otimes (S, \hat{\mathcal{D}}_{I_0}^{\lambda\mu_0}\Psi) \otimes e_\otimes^{S_{\text{int}}}). \end{aligned} \quad (80)$$

²³The second identity follows directly from (78) and (69). From the second identity, (69), and (68), we conclude that the field corresponding to the l.h.s. of the first identity is s closed (modulo d). But there are no such fields of ghost number -1 , anti-symmetric and linear in \bar{a} , \bar{a}' and of mass dimension four (two of which are already contributed by \bar{a} and \bar{a}'): By power counting, the only possible fields with these properties are $\text{Tr}_N([\bar{a}^\mu, \bar{a}'_\mu]B^\ddagger)$ and $\text{Tr}_N([\bar{a}^\mu, \bar{a}'_\mu]\hat{C})\text{vol}$, but no linear combination of these is s closed modulo d .

Hence,

$$\hat{\mathcal{D}}^\lambda \left(A^m (\bar{\delta}_I^{\lambda\mu} \Psi \otimes e_\otimes^{S_{\text{int}}}) \hat{\mathcal{D}} \Xi^I \right) \sim -\frac{1}{2} (S, A^m (\hat{\mathcal{D}}_{I_1}^{\lambda\mu_1} \Psi \otimes \hat{\mathcal{D}}_{I_0}^{\lambda\mu_0} \Psi \otimes e_\otimes^{S_{\text{int}}})) \hat{\mathcal{D}} \Xi_{\mu_0}^{I_0} \wedge \hat{\mathcal{D}} \Xi_{\mu_1}^{I_1}. \quad (81)$$

In terms of fields, this means that, modulo d exact terms,

$$\hat{\mathcal{D}} b^m = \hat{\mathcal{D}} \xi^m = s \chi^m, \quad (82)$$

with $\chi^m = \frac{1}{2} A^m (\hat{\mathcal{D}}^\lambda \Psi \otimes \hat{\mathcal{D}}^\lambda \Psi \otimes e_\otimes^{S_{\text{int}}})|_{\lambda=1}$.

On the other hand, we can directly compute the l.h.s. of (82) for b^m corresponding to the non-abelian anomaly, i.e., given by (73). We obtain, up to d exact terms,

$$(\hat{\mathcal{D}} b^m)(\bar{a}_1, \bar{a}_2) = \frac{1}{2} \text{Tr}_N [(\bar{a}_1 \bar{a}_2 - \bar{a}_2 \bar{a}_1) \bar{F}]. \quad (83)$$

The r.h.s., being a c-number and not d exact, is not s exact modulo d. Hence, (82) is not fulfilled. Thus, also the non-abelian anomaly can not be present under the assumptions of Theorem 1. This concludes the proof of Theorem 1.

We remark that the r.h.s. of (83) coincides with the ‘‘non-abelian’’ obstruction $E(a_1, a_2)$ to fulfill perturbative agreement w.r.t. changes in the background connection, cf. [39].

4 Conclusion

We have proven under quite general conditions that the absence of gauge anomalies can be determined at the one-loop level. As our proof is based on perturbative agreement, which is a formalization of background independence (but see [23] for a thorough discussion of background independence in gauge theories), one might also phrase our result as stating that background independence w.r.t. changes in the background connection implies absence of gauge anomalies.

A shortcoming of our analysis is that we restricted to vanishing background bosonic matter fields. We briefly sketch how to include these and how to possibly extend our result to that case. At the level of the action, non-trivial background matter fields can be simply introduced by replacing the bosonic matter field φ by $\bar{\Phi} + \varphi$, with $\bar{\Phi}$ the matter background field. The action of the BV operator on φ then has to be modified to $s\varphi = -\rho(C)(\bar{\Phi} + \varphi)$ so that also the free part s_0 acts non-trivially on the bosonic matter fields. In the gauge fixing fermion (7), also a term $\int \tilde{C}^I H^I(\bar{\Phi}, \varphi) \text{vol}$ would be included, allowing for an R_ξ gauge (in which terms with single derivatives are eliminated from the free part of the action).

As for anomalies, the non-abelian anomaly (43) is independent of matter fields (in particular background matter fields), so if it is absent for vanishing matter background fields (as is guaranteed if the assumptions of Thm. 1 are fulfilled), it will also be absent for non-trivial matter background fields. The same is true for the abelian anomalies of the form (45), which are well-known to potentially occur. It remains to rule out abelian anomalies depending on the background matter field. As discussed in Remark 4, one may hope to rule these out on general grounds. Alternatively, one can slightly strengthen the requirements of Thm. 1 such that also perturbative agreement w.r.t. changes in the background matter field holds and rule out the occurrence of anomalies depending explicitly on the background matter field $\bar{\Phi}$ (not just on the combination $\bar{\Phi} + \varphi$ which has already been ruled out by Thm. 1 for $\bar{\Phi} = 0$) by using again the background independence of the anomaly.

Acknowledgements

I would like to thank Stefan Hollands for useful discussions.

References

- [1] S.L. Adler and W.A. Bardeen, Absence of higher order corrections in the anomalous axial vector divergence equation, *Phys. Rev.* 182 (1969) 1517.
- [2] A. Zee, Axial vector anomalies and the scaling property of field theory, *Phys. Rev. Lett.* 29 (1972) 1198.
- [3] J.H. Lowenstein and B. Schroer, Comment on the absence of radiative corrections to the anomaly of the axial-vector current, *Phys. Rev. D* 7 (1973) 1929.
- [4] V. Mastropietro, Emergent Adler-Bardeen theorem, *JHEP* 03 (2020) 095, [arXiv:2001.00183].
- [5] G. Costa et al., Nonabelian Gauge Theories and Triangle Anomalies, *Nuovo Cim.* A38 (1977) 373.
- [6] G. Bandelloni et al., On the cancellation of hard anomalies in gauge field models: A regularization independent proof, *Commun. Math. Phys.* 72 (1980) 239.
- [7] O. Piguet and S.P. Sorella, On the finiteness of the BRS modulo cocycles, *Nucl. Phys.* B381 (1992) 373, [arXiv:hep-th/9302122].
- [8] O. Piguet and S.P. Sorella, Adler-Bardeen theorem and vanishing of the gauge beta function, *Nucl. Phys.* B395 (1993) 661, [arXiv:hep-th/9302123].
- [9] D. Anselmi, Adler-Bardeen theorem and manifest anomaly cancellation to all orders in gauge theories, *Eur. Phys. J. C* 74 (2014) 3083, [arXiv:1402.6453].
- [10] C.G. Callan, Jr., Broken scale invariance in scalar field theory, *Phys. Rev. D* 2 (1970) 1541.
- [11] K. Symanzik, Small distance behavior in field theory and power counting, *Commun. Math. Phys.* 18 (1970) 227.
- [12] J.H. Lowenstein, Differential vertex operations in Lagrangian field theory, *Commun. Math. Phys.* 24 (1971) 1.
- [13] Y.M.P. Lam, Perturbation Lagrangian theory for scalar fields: Ward-Takahasi identity and current algebra, *Phys. Rev. D* 6 (1972) 2145.
- [14] C.Q. Geng and R.E. Marshak, Uniqueness of Quark and Lepton Representations in the Standard Model From the Anomalies Viewpoint, *Phys. Rev. D* 39 (1989) 693.
- [15] S. Hollands and R.M. Wald, Local Wick polynomials and time ordered products of quantum fields in curved space-time, *Commun. Math. Phys.* 223 (2001) 289, [arXiv:gr-qc/0103074].
- [16] S. Hollands and R.M. Wald, Existence of local covariant time ordered products of quantum fields in curved space-time, *Commun. Math. Phys.* 231 (2002) 309, [arXiv:gr-qc/0111108].
- [17] S. Hollands, Renormalized Quantum Yang-Mills Fields in Curved Spacetime, *Rev. Math. Phys.* 20 (2008) 1033, [arXiv:0705.3340].
- [18] S. Hollands and R.M. Wald, Quantum fields in curved spacetime, *Phys. Rept.* 574 (2015) 1, [arXiv:1401.2026].
- [19] K. Rejzner, *Perturbative algebraic quantum field theory* (Springer, Cham, 2016).

- [20] S. Hollands and R.M. Wald, On the Renormalization Group in Curved Spacetime, *Commun. Math. Phys.* 237 (2003) 123, [arXiv:gr-qc/0209029].
- [21] S. Hollands and R.M. Wald, Conservation of the stress tensor in interacting quantum field theory in curved spacetimes, *Rev. Math. Phys.* 17 (2005) 227, [arXiv:gr-qc/0404074].
- [22] J. Zahn, Locally covariant charged fields and background independence, *Rev. Math. Phys.* 27 (2015) 1550017, [arXiv:1311.7661].
- [23] M. Taslimi Tehrani and J. Zahn, Background independence in gauge theories, *Annales Henri Poincaré* 21 (2020) 1135, [arXiv:1804.07640].
- [24] W.A. Bardeen and B. Zumino, Consistent and Covariant Anomalies in Gauge and Gravitational Theories, *Nucl. Phys.* B244 (1984) 421.
- [25] G. Barnich, F. Brandt and M. Henneaux, Local BRST cohomology in gauge theories, *Phys. Rept.* 338 (2000) 439, [arXiv:hep-th/0002245].
- [26] J. Zahn, Locally covariant chiral fermions and anomalies, *Nucl. Phys.* B890 (2015) 1, [arXiv:1407.1994].
- [27] J. Zahn, The renormalized locally covariant Dirac field, *Rev. Math. Phys.* 26 (2014) 1330012, [arXiv:1210.4031].
- [28] J.M. Arms, The structure of the solution set for the yang-mills equations, *Mathematical Proceedings of the Cambridge Philosophical Society* 90 (1981) 361.
- [29] M. Taslimi Tehrani, Quantum BRST charge in gauge theories in curved space-time, *J. Math. Phys.* 60 (2019) 012304, [arXiv:1703.04148].
- [30] H. Kluberg-Stern and J.B. Zuber, Renormalization of Nonabelian Gauge Theories in a Background Field Gauge. 1. Green Functions, *Phys. Rev.* D12 (1975) 482.
- [31] S. Waldmann, *Poisson-Geometrie und Deformationsquantisierung* (Springer, Berlin, 2007).
- [32] M.B. Fröb, Anomalies in Time-Ordered Products and Applications to the BV–BRST Formulation of Quantum Gauge Theories, *Commun. Math. Phys.* 372 (2019) 281, [arXiv:1803.10235].
- [33] R. Brunetti and K. Fredenhagen, Microlocal analysis and interacting quantum field theories: Renormalization on physical backgrounds, *Commun. Math. Phys.* 208 (2000) 623, [arXiv:math-ph/9903028].
- [34] R. Brunetti, M. Dütsch and K. Fredenhagen, Perturbative Algebraic Quantum Field Theory and the Renormalization Groups, *Adv. Theor. Math. Phys.* 13 (2009) 1541, [arXiv:0901.2038].
- [35] J. Manes, R. Stora and B. Zumino, Algebraic Study of Chiral Anomalies, *Commun. Math. Phys.* 102 (1985) 157.
- [36] O. Piguet and S.P. Sorella, *Algebraic Renormalization, Lecture Notes in Physics Vol. 28* (Springer, 1995).
- [37] Y. Nakayama, Realization of impossible anomalies, *Phys. Rev. D* 98 (2018) 085002, [arXiv:1804.02940].

- [38] J. Wess and B. Zumino, Consequences of anomalous Ward identities, *Phys. Lett.* B37 (1971) 95.
- [39] A. Schenkel and J. Zahn, Global anomalies on Lorentzian space-times, *Annales Henri Poincaré* 18 (2017) 2693, [arXiv:1609.06562].