

# Mixed formulation for the computation of Miura surfaces with Dirichlet boundary conditions

Frédéric Marazzato

Department of Mathematics, Louisiana State University, Baton Rouge, LA 70803, USA  
email: marazzato@lsu.edu

## Abstract

Miura surfaces are the solutions of a constrained nonlinear elliptic system of equations. This system is derived by homogenization from the Miura fold, which is a type of origami fold with multiple applications in engineering. A previous inquiry, gave suboptimal conditions for existence of solutions and proposed an  $H^2$ -conformal finite element method to approximate them. In this paper, the existence of Miura surfaces is studied using a mixed formulation. It is also proved that the constraints propagate from the boundary to the interior of the domain for well-chosen boundary conditions. Then, a numerical method based on a least-squares formulation, Taylor–Hood finite elements and a Newton method is introduced to approximate Miura surfaces. The numerical method is proved to converge at order one in space and numerical tests are performed to demonstrate its robustness.

*Keywords:* Origami, nonlinear elliptic equation, Kinematics of deformation.

*AMS Subject Classification:* 35J66, 65N12, 65N15, 65N30 and 74A05.

## 1 Introduction

The ancient art of origami has attracted a lot of attention in recent years. The monograph [14] presents a very complete description of many different types of origami patterns as well as some questions surrounding them. Origami have found many applications in engineering. Origami folds are, for instance, used to ensure that airbags inflate correctly from their folded state [1]. Applications in aerospace engineering are as varied as radiators, panels for radio telescopes [18] or soft robots [22]. More recently, origami have been studied with a view to produce metamaterials [21, 2].

The Miura fold was introduced in [17]. It has later found applications in solar panels for satellites and in biology [13]. Recent applications also include the design of metamaterials [27]. The Miura ori is flat when completely unfolded and can be fully folded in a very compact form, hence its applications in aerospace engineering. However, the Miura fold can also assume different shapes when partially unfolded. [24] and [26] provided a description of these partially unfolded states through a computation of the Poisson ratio of the Miura fold, which happens to be negative for in-plane deformations and positive for out-of-plane bending. Miura folds are therefore globally saddle shaped. A homogenization procedure for origami folds was proposed in [20] and [19] and then applied to the Miura fold in [15]. The authors obtained a constrained nonlinear elliptic equation describing the limit surface, which is called a Miura surface. The constraints are both equality and inequality constraints. In [16], existence and uniqueness of solutions was proved but only for the unconstrained equation and under restrictive assumptions. [16] also proposed an  $H^2$ -conformal Finite Element Method (FEM) to compute solutions to the problem, which is computationally involved.




Figure 1: Miura reference cell.

In this paper, existence and uniqueness of solutions of the constrained equation is proved under adequate boundary conditions. Thereafter, a least-squares formulation is proposed coupled to a Newton method and Taylor–Hood finite elements to approximate the solutions. The robustness of the method is demonstrated on several test cases.

## 2 Continuous equations

### 2.1 Modeling of the Miura fold

The Miura fold is based on the reference cell sketched in Figure 1, in which all edges have unit length. The reference cell is made of four parallelograms and can be folded along the full lines in Figure 1. Following [26, 24], we consider that the cells can also bend along the dashed lines. However, the cells cannot stretch. We restrict the deformations of the Miura cell to folding along crease lines or bending in the facets. A Miura tessellation is based on the continuous juxtaposition of reference cells, dilated by a factor  $r > 0$ . In the spirit of homogenization, [19, 15] have proposed a procedure to compute a surface that is the limit when  $r \rightarrow 0$  of a Miura tessellation. This procedure leads to the constrained PDE presented in the next section.

### 2.2 Strong form equations

Let  $\Omega \subset \mathbb{R}^2$  be a bounded convex polygon that can be perfectly fitted by triangular meshes. Note that, due to the convexity hypothesis, the boundary  $\partial\Omega$  is Lipschitz [11] and thus verifies an exterior sphere condition. Let  $\varphi : \Omega \subset \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a parametrization of the homogenized surface constructed from a Miura tessellation. The coordinates of  $\varphi$  are written as  $\varphi^i$ , for  $i \in \{1, 2, 3\}$ . As proved in [15],  $\varphi$  is a solution of the following strong form equation:

$$p(\varphi_x)\varphi_{xx} + q(\varphi_y)\varphi_{yy} = 0 \in \mathbb{R}^3, \quad (1a)$$

$$p(\varphi_x)q(\varphi_y) = 4, \quad (1b)$$

$$\varphi_x \cdot \varphi_y = 0, \quad (1c)$$

$$0 < |\varphi_x|^2 \leq 3, \quad 1 < |\varphi_y|^2 \leq 4, \quad (1d)$$

where

$$p(\varphi_x) = \frac{4}{4 - |\varphi_x|^2}, \quad q(\varphi_y) = \frac{4}{|\varphi_y|^2},$$

and the subscripts  $x$  and  $y$  stand respectively for  $\partial_x$  and  $\partial_y$ . (1a) describes the fact that the in-plane and out-of-plane Poisson ratios for the bending mode of the Miura fold are equal, as proved in [24, 26]. (1b) and (1c) are equality constraints that stem from the kinematics of the bending mode of the Miura ori, see [15]. Finally, (1d) is enforced to prevent the metric tensor of the surface  $\varphi(\Omega)$  from being singular, see [15]. In practice, that means that, locally, when the bounds are reached, the pattern is fully folded or fully unfolded.

Equation (1a) is nonlinear elliptic and is proved in the following to have solutions for quite general Dirichlet boundary conditions. However, it is not clear that all solutions verify the constraints (1b)-(1d). As stated in [15], these constraints are necessary to construct a Miura surface. In the following, we investigate under what supplementary assumptions the solutions of (1a) verify the equality constraints (1b) and (1c) and the inequality constraints (1d). Note that the constraint  $1 < |\varphi_y|^2$  in (1d) is particularly challenging as it is non-convex.

### 2.3 Main result

We introduce the Hilbert space  $V := (H^2(\Omega))^3$ , equipped with the usual  $(H^2(\Omega))^3$  Sobolev norm. Note that due to Rellich–Kondrachov theorem [4, Theorem 9.16],  $V \subset (C^0(\bar{\Omega}))^3$ . For  $\varphi \in V$ , let  $\mathcal{A}(\varphi) : V \mapsto L^2(\Omega)^3$ , be the operator defined for  $\psi \in V$  as

$$\mathcal{A}(\varphi)\psi := p(\varphi_x)\psi_{xx} + q(\varphi_y)\psi_{yy} \in \mathbb{R}^3. \quad (2)$$

Solving Equation (1a) consists in finding  $\varphi \in V$  such that

$$\mathcal{A}(\varphi)\varphi = 0 \text{ in } \Omega. \quad (3)$$

Note that (3) admits a translation invariance. Therefore, we consider the convex subset  $\mathcal{V} := \{\varphi \in V \mid \int_{\Omega} \varphi = 0\}$ , as the solution space to remove the translation invariance. The operator  $\mathcal{A}(\varphi)$  is not well adapted to obtain a constrained solution of (1a). Equation (3) is thus reformulated into an equation on  $\mathcal{G} \equiv \nabla\varphi$ .

Let  $W := H^1(\Omega)^{3 \times 2}$  equipped with the usual  $H^1(\Omega)$  norm. Let  $\mathfrak{T} : H^1(\Omega)^{3 \times 2} \rightarrow H^{\frac{1}{2}}(\partial\Omega)^{3 \times 2}$  be the trace operator, see [8, Theorem B.52] for instance. Let  $X := \mathfrak{T}\{G \in W \mid G_y^x = G_x^y \text{ a.e. in } \Omega\}$  be a trace space, and  $Y := \{G \in X \mid G \in H^{\frac{3}{2}}(\partial\Omega)^{3 \times 2}\}$ , equipped with the usual  $H^{\frac{3}{2}}(\partial\Omega)$  norm. The condition in the definition of  $X$  translates the fact that  $\mathcal{G}$  should be a gradient and thus verify a generalization of Clairault’s theorem. Let  $G_D \in Y$  be the imposed Dirichlet boundary condition on  $\partial\Omega$ . We consider the convex subset  $W_D := \{G \in W \mid G = G_D \text{ on } \partial\Omega\}$  as the solution space and  $W_0 := H_0^1(\Omega)^{3 \times 2}$  as the associated homogeneous space. Let  $A(\mathcal{G}) : W \mapsto L^2(\Omega)^3$  be the operator such that, for  $G = (G^x, G^y) \in W$ ,

$$A(\mathcal{G})G := p(\mathcal{G}^x)G_x^x + q(\mathcal{G}^y)G_y^y.$$

$\mathcal{G}^x$  and  $\mathcal{G}^y$  are interpreted as the tangent vectors to the computed Miura surface. Equation (1d) indicates that we are not interested in the values of  $\nabla\varphi$  when  $|\varphi_x| > 3$ ,  $|\varphi_y| > 4$  and  $|\varphi_y| < 1$ . We thus define the Lipschitz cut-offs,

$$\bar{p}(\mathcal{G}^x) := \begin{cases} 4 & \text{if } |\mathcal{G}^x|^2 \geq 3 \\ p(\mathcal{G}^x) & \text{otherwise} \end{cases}, \quad \bar{q}(\mathcal{G}^y) := \begin{cases} 4 & \text{if } |\mathcal{G}^y|^2 \leq 1 \\ 1 & \text{if } |\mathcal{G}^y|^2 \geq 4 \\ q(\mathcal{G}^y) & \text{otherwise} \end{cases}.$$

Therefore,  $1 \leq \bar{p}(\mathcal{G}^x) \leq 4$  and  $1 \leq \bar{q}(\mathcal{G}^y) \leq 4$  and  $\bar{p}$  and  $\bar{q}$  are  $K$ -Lipschitz functions, where  $K > 0$ . The operator  $\bar{A}(\mathcal{G})$  is defined for  $G \in W$  as,

$$\bar{A}(\mathcal{G})G := \bar{p}(\mathcal{G}^x)G_x^x + \bar{q}(\mathcal{G}^y)G_y^y. \quad (4)$$

Note that the operator  $\bar{A}(\mathcal{G})$  is uniformly elliptic, whereas  $A(\mathcal{G})$  is not. Therefore,  $\bar{A}(\mathcal{G})$  verifies the Cordès condition, see [25, 10]. We thus propose to find  $\mathcal{G} \in W_D$ ,  $\mathcal{G}_y^x = \mathcal{G}_x^y$  a.e. in  $\Omega$ , such that

$$\bar{A}(\mathcal{G})\mathcal{G} = 0 \text{ in } \Omega. \quad (5)$$

The main result of this paper is Theorem 9 which proves the existence of solutions to (1) under appropriate Dirichlet boundary conditions. Let us describe the boundary conditions that will be used. We write the unit normal to a Miura surface as

$$\mathcal{N} := \frac{\mathcal{G}^x \wedge \mathcal{G}^y}{|\mathcal{G}^x \wedge \mathcal{G}^y|}.$$

As shown in Figure 2, we impose on  $\partial\Omega$ ,  $\mathcal{N}$ ,  $|\mathcal{G}^x|$ ,  $|\mathcal{G}^y|$  and  $\mathcal{G}^x \cdot \mathcal{G}^y = 0$ . A final degree of freedom needs to be fixed in the form of an angle of rotation around  $\mathcal{N}$ , written  $\alpha$ .




Figure 2: Imposed boundary conditions

This translates to  $G_D$  being chosen so that  $|G_D^x|$  and  $|G_D^y|$  verify (1b) and (1d) on  $\partial\Omega$ .  $G_D^x \cdot G_D^y = 0$  is similarly set to verify (1c) on  $\partial\Omega$ . Note that this consists in imposing the first fundamental form [5] associated to  $\varphi$  on  $\partial\Omega$ . Also note that, it is not a priori easy to know if a given  $G_D$ , as in Figure 2, is in  $X$  (i.e. is the trace of a gradient).

## 2.4 Linearized problem

Let  $\mathcal{G} \in W$ . The linearized equation we want to solve is thus, find  $G \in W_D$ ,  $G_y^x = G_x^y$  a.e. in  $\Omega$ ,

$$\bar{A}(\mathcal{G})G = 0 \text{ in } \Omega. \tag{6}$$

To study the well-posedness of (6), we write an appropriate mixed formulation, following [9]. As  $G_D \in Y \subset H^{\frac{3}{2}}(\partial\Omega)^{3 \times 2}$ , there exists  $\hat{G}_D \in H^2(\Omega)^{3 \times 2}$ ,  $\hat{G}_D = G_D$  on  $\partial\Omega$  and  $\hat{G}_{D,y}^x = \hat{G}_{D,x}^y$  a.e. in  $\Omega$ . We define

$$f := -\bar{A}(\mathcal{G})\hat{G}_D \in L^2(\Omega)^3.$$

Let  $R := \{r \in L^2(\Omega)^3 \mid \int_{\Omega} r = 0\}$ , equipped with the usual  $L^2$  norm. Let  $\hat{G} \in W_0$ , we define the bilinear form, for all  $\tilde{G} \in W_0$ ,

$$a(\mathcal{G}; \hat{G}, \tilde{G}) := \int_{\Omega} \bar{A}(\mathcal{G})\hat{G} \cdot \bar{A}(\mathcal{G})\tilde{G},$$

and for all  $\tilde{r} \in R$ ,

$$b(\hat{G}, \tilde{r}) = \int_{\Omega} (\hat{G}_y^x - \hat{G}_x^y) \cdot \tilde{r}.$$

The mixed formulation of (6) consists in seeking  $(\hat{G}, r) \in W_0 \times R$ ,

$$a(\mathcal{G}, \hat{G}, \tilde{G}) + b(\tilde{G}, r) = \int_{\Omega} f \cdot \bar{A}(\mathcal{G})\tilde{G}, \quad \forall \tilde{G} \in W_0, \tag{7a}$$

$$b(\hat{G}, \tilde{r}) = 0, \quad \forall \tilde{r} \in R. \tag{7b}$$

Then,  $G := \hat{G} + \hat{G}_D$  will be a solution of (6).

**Lemma 1** (Coercivity). *Let  $\gamma(\mathcal{G})$  such that*

$$\gamma(\mathcal{G}) := \frac{\bar{p}(\mathcal{G}^x) + \bar{q}(\mathcal{G}^y)}{\bar{p}(\mathcal{G}^x)^2 + \bar{q}(\mathcal{G}^y)^2}.$$

*One has  $\gamma_0 := \frac{1}{16} \leq \gamma(\mathcal{G}) \leq 4 =: \gamma_1$ . There exists  $\varepsilon \in (0, 1)$ , independent of  $\gamma(\mathcal{G})$ , for all  $G \in W_0$ ,*

$$\left( \frac{1 - \sqrt{1 - \varepsilon}}{\gamma_1} \right)^2 \|\nabla G\|_{L^2(\Omega)}^2 \leq a(\mathcal{G}; G, G).$$

*Thus,  $a(\mathcal{G})$  is coercive over  $W_0^2$ , with a constant independent of  $\mathcal{G}$ .*

*Proof.* Following, [10, 25], as  $\bar{A}(\mathcal{G})$  verifies the Cordès condition, one can prove that, there exists  $\varepsilon \in (0, 1)$ , independent of  $\mathcal{G}$ , such that

$$\|\gamma(\mathcal{G})\bar{A}(\mathcal{G})G\|_{L^2(\Omega)} \geq (1 - \sqrt{1 - \varepsilon})\|\operatorname{div}(G)\|_{L^2(\Omega)}.$$

One also has, for all  $G \in W_0$ ,

$$\|\nabla G\|_{L^2(\Omega)}^2 \leq \|G_y^x - G_x^y\|_{L^2(\Omega)}^2 + \|\operatorname{div}(G)\|_{L^2(\Omega)}^2,$$

as shown in [6, Theorem 2.3]. Therefore, for all  $G \in W_0$ , such that  $G_y^x = G_x^y$  a.e. in  $\Omega$ ,

$$\|\gamma(\mathcal{G})\bar{A}(\mathcal{G})G\|_{L^2(\Omega)} \geq (1 - \sqrt{1 - \varepsilon})\|\nabla G\|_{L^2(\Omega)}.$$

Finally, one has

$$(1 - \sqrt{1 - \varepsilon})^2 \|\nabla G\|_{L^2(\Omega)}^2 \leq \gamma_1^2 \|\bar{A}(\mathcal{G})G\|_{L^2(\Omega)}^2.$$

□

**Lemma 2.** *Equation (7) admits a unique solution  $G \in W_D$ . There exists  $C > 0$ , independent from  $\mathcal{G}$ ,*

$$\|G\|_{H^1(\Omega)} \leq C\|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)}.$$

*Proof.* There exists  $C > 0$ , independent from  $\mathcal{G}$ ,

$$\|\hat{G}_D\|_{H^2(\Omega)} \leq C\|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)}.$$

We use the BNB lemma, see [8, Theorem 2.6, p. 85], in the context of saddle point problems [8, Theorem 2.34, p. 100]. As shown in [9], there exists  $\beta > 0$ , independent of  $\mathcal{G}$ ,

$$\beta \leq \inf_{\hat{G} \in W_0 \setminus \{0\}} \sup_{\tilde{r} \in R \setminus \{0\}} \frac{b(\hat{G}, \tilde{r})}{\|\tilde{r}\|_R \|\hat{G}\|_{W_0}}.$$

$a(\mathcal{G})$  was proved to be coercive in Lemma 1, thus fulfilling the two conditions of the BNB theorem. Applying the BNB theorem and a trace inequality gives

$$\|G\|_{H^1(\Omega)} \leq \|\hat{G}_D\|_{H^1(\Omega)} + C\|\bar{A}(\mathcal{G})\hat{G}_D\|_{L^2(\Omega)} \leq C'\|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)},$$

where  $C, C' > 0$  is independent of  $\mathcal{G}$ .

□

**Proposition 3.** *The solution  $G \in W_D$  of Equation (6) benefits from more regularity as  $G \in H^2(\Omega)^{3 \times 2} \subset C^0(\bar{\Omega})^{3 \times 2}$ . Also, one has*

$$\|G\|_{H^2(\Omega)} \leq C \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)},$$

where  $C > 0$  is independent of  $\mathcal{G}$ .

*Proof.* We derive the first line in the right-hand side of (6) with respect to both variables. As  $f \in H^1(\Omega)^6$ , one gets

$$\begin{cases} \bar{p}(\mathcal{G}^x)G_{xx}^x + \bar{q}(\mathcal{G}^y)G_{yy}^x = f_x - (\bar{p}(\mathcal{G}^x))_x G_x^x - (\bar{q}(\mathcal{G}^y))_x G_y^y \in L^2(\Omega)^3, \\ \bar{p}(\mathcal{G}^x)G_{xx}^y + \bar{q}(\mathcal{G}^y)G_{yy}^y = f_y - (\bar{p}(\mathcal{G}^x))_y G_x^x - (\bar{q}(\mathcal{G}^y))_y G_y^y \in L^2(\Omega)^3. \end{cases}$$

These are two independent linear uniformly elliptic equations in  $G^x$  and  $G^y$ . It is proved in [25, 16], that such equations have a unique solution in  $H^2(\Omega)^3 \cap H_0^1(\Omega)^3$ . Therefore  $G \in H^2(\Omega)^{3 \times 2} \subset C^0(\bar{\Omega})^{3 \times 2}$  and

$$\begin{cases} \|G^x\|_{H^2(\Omega)} \leq C \|f_x - \bar{p}_x G_x^x - \bar{q}_x G_y^y\|_{L^2(\Omega)} \leq C \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)}, \\ \|G^y\|_{H^2(\Omega)} \leq C \|f_y - \bar{p}_y G_x^x - \bar{q}_y G_y^y\|_{L^2(\Omega)} \leq C \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)}, \end{cases}$$

where  $C > 0$  is independent from  $\mathcal{G}$ . □

## 2.5 Nonlinear problem

The existence of a solution to (5) is proved through a fixed point method. Let  $T : H^1(\Omega)^{3 \times 2} \ni \mathcal{G} \mapsto G(\mathcal{G}) \in W_D \cap H^2(\Omega)^{3 \times 2}$  be the map that, given a  $\mathcal{G} \in H^1(\Omega)^{3 \times 2}$ , maps to the unique solution of (6). We define the following subset

$$B = \left\{ \tilde{G} \in H^2(\Omega)^{3 \times 2} \cap W_D; \|\tilde{G}\|_{H^2(\Omega)} \leq C \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \right\}, \quad (8)$$

where  $C > 0$  is the constant from Proposition 3.

**Proposition 4.** *The map  $T$  admits a fixed point  $\mathcal{G} \in B$  which verifies*

$$\begin{cases} \bar{A}(\mathcal{G})\mathcal{G} = 0 \text{ in } \Omega, \\ \mathcal{G} = G_D \text{ on } \partial\Omega. \end{cases}$$

*Proof.* Let us show that  $B$  is stable by  $T$ . Let  $\mathcal{G} \in B$ . Letting  $G := T\mathcal{G}$ , one thus has  $G \in B$ , as a consequence of Proposition 3. Also,  $B \subset H^1(\Omega)^{3 \times 2}$  with a compact Sobolev embedding and  $H^1(\Omega)^{3 \times 2}$  equipped with its usual norm is a Banach space. Thus,  $B$  is a closed convex subset of the Banach space  $H^1(\Omega)^{3 \times 2}$  and  $TB$  is precompact in  $H^1(\Omega)^{3 \times 2}$ .

We prove that  $T$  is continuous over  $H^1(\Omega)^{3 \times 2}$ . Let  $(\mathcal{G}_n)_n$  be a sequence of  $B$  such that  $\mathcal{G}_n \xrightarrow{n \rightarrow +\infty} \mathcal{G} \in B$ , for  $\|\cdot\|_{H^1(\Omega)}$ . Let  $G := T\mathcal{G}$  and  $G_n := T\mathcal{G}_n$ , for  $n \in \mathbb{N}$ . We want to prove that  $G_n \xrightarrow{n \rightarrow +\infty} G$  for  $\|\cdot\|_{H^1(\Omega)}$ . As for all  $n \in \mathbb{N}$ ,  $G_n \in B$ ,  $(G_n)_n$  is bounded in  $H^2(\Omega)^{3 \times 2}$ . Therefore, there exists  $\hat{G} \in H^2(\Omega)^{3 \times 2}$ , up to a subsequence,  $G_n \xrightarrow{n \rightarrow +\infty} \hat{G}$ . Let us show that  $\hat{G}$  is a solution of (6). By definition, one has

$$\begin{aligned} \|\bar{A}(\mathcal{G})G_n\|_{L^2(\Omega)} &= \|\bar{A}(\mathcal{G})G_n - \bar{A}(\mathcal{G}_n)G_n\|_{L^2(\Omega)} \\ &\leq K \|\mathcal{G} - \mathcal{G}_n\|_{H^1(\Omega)} \|G_n\|_{H^2(\Omega)} \\ &\leq CK \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \|\mathcal{G} - \mathcal{G}_n\|_{H^1(\Omega)} \xrightarrow{n \rightarrow \infty} 0, \end{aligned}$$

since  $\bar{A}$  is  $K$ -Lipschitz. The left-hand side converges to  $\|\bar{A}(\mathcal{G})\hat{G}\|_{L^2(\Omega)}$  when  $n \rightarrow +\infty$ . As for all  $n \in \mathbb{N}$ ,  $G_n \in W_D$ , then  $\hat{G} \in W_D$  and  $\hat{G} = G_D$  on  $\partial\Omega$ . Therefore,  $\hat{G}$  solves (6). Also, as  $H^2(\Omega)^{3 \times 2} \subset H^1(\Omega)^{3 \times 2}$ , up to a subsequence,

$$\|G_n - \hat{G}\|_{H^1(\Omega)} \leq \|G_n - \hat{G}\|_{H^2(\Omega)} \xrightarrow{n \rightarrow \infty} 0.$$

Using Lemma 2, the solution of (6) is unique in  $W_D$ . Thus the full sequence  $(G_n)_n$  converges towards  $\hat{G} = G = T\mathcal{G}$  in  $H^1(\Omega)^{3 \times 2}$ . We conclude with the Schauder fixed point Theorem, see Corollary 11.2 of [10], that  $T$  admits a fixed point.  $\square$

**Lemma 5** (Uniqueness). *Let  $\delta > 0$  and  $\|G_D\|_{H^{\frac{3}{2}}(\Omega)} \leq \delta$ . For  $\delta$  small enough, there exists a unique  $\mathcal{G} \in W_D$  solution of (5).*

*Proof.* Let  $\mathcal{G}, \tilde{\mathcal{G}} \in W_D$  solutions of (5) and  $\hat{G}_D \in H^2(\Omega)^{3 \times 2}$ ,  $\hat{G}_D = G_D$  on  $\partial\Omega$ . Let  $G := \mathcal{G} - \hat{G}_D \in H_0^1(\Omega)^{3 \times 2}$  and  $\tilde{G} := \tilde{\mathcal{G}} - \hat{G}_D \in H_0^1(\Omega)^{3 \times 2}$ . One thus has,

$$\begin{cases} \bar{A}(\mathcal{G})G = -\bar{A}(\mathcal{G})\hat{G}_D, \\ \bar{A}(\tilde{\mathcal{G}})\tilde{G} = -\bar{A}(\tilde{\mathcal{G}})\hat{G}_D, \end{cases}$$

Therefore,

$$\begin{aligned} \bar{A}(\tilde{\mathcal{G}})\tilde{G} - \bar{A}(\mathcal{G})G &= (\bar{A}(\mathcal{G}) - \bar{A}(\tilde{\mathcal{G}}))\hat{G}_D, \\ \bar{A}(\tilde{\mathcal{G}})\tilde{G} - \bar{A}(\tilde{\mathcal{G}})G + \bar{A}(\tilde{\mathcal{G}})G - \bar{A}(\mathcal{G})G &= (\bar{A}(\mathcal{G}) - \bar{A}(\tilde{\mathcal{G}}))\hat{G}_D, \\ \bar{A}(\tilde{\mathcal{G}})(\tilde{G} - G) &= (\bar{A}(\mathcal{G}) - \bar{A}(\tilde{\mathcal{G}}))(\hat{G}_D + G). \end{aligned}$$

Using the coercivity proved in Lemma 1, one has

$$c\|G - \tilde{G}\|_{H^1(\Omega)}^2 \leq C\|G - \tilde{G}\|_{H^1(\Omega)}\|(\bar{A}(\mathcal{G}) - \bar{A}(\tilde{\mathcal{G}}))(\hat{G}_D + G)\|_{L^2(\Omega)},$$

where  $c, C > 0$  are constants independent of  $\mathcal{G}, \tilde{\mathcal{G}}$ . Therefore,

$$\begin{aligned} \|G - \tilde{G}\|_{H^1(\Omega)} &\leq \frac{C}{c}K\|\mathcal{G} - \tilde{\mathcal{G}}\|\|\hat{G}_D + G\|_{L^2(\Omega)}, \\ &\leq \frac{CK}{c}\|\mathcal{G} - \tilde{\mathcal{G}}\|_{L^p(\Omega)}\left(\|\nabla\hat{G}_D\|_{L^q(\Omega)} + \|\nabla G\|_{L^q(\Omega)}\right), \end{aligned}$$

since  $\bar{A}$  is  $K$ -Lipschitz, and using a Hölder inequality with  $p^{-1} + q^{-1} = 2^{-1}$ ,  $p > 2$  and  $q < 2$ . Thus,

$$\begin{aligned} \|G - \tilde{G}\|_{H^1(\Omega)} &\leq \frac{CK}{c}\|\mathcal{G} - \tilde{\mathcal{G}}\|_{H^1(\Omega)}\left(\|\nabla\hat{G}_D\|_{L^2(\Omega)} + \|\nabla G\|_{L^2(\Omega)}\right), \\ &\leq \frac{CKC'}{c}\|G_D\|_{H^{\frac{3}{2}}(\Omega)}\|\mathcal{G} - \tilde{\mathcal{G}}\|_{H^1(\Omega)}, \\ &\leq \frac{CKC'}{c}\delta\|\mathcal{G} - \tilde{\mathcal{G}}\|_{H^1(\Omega)}, \end{aligned}$$

where  $C' > 0$  is independent from  $\mathcal{G}$  and  $\tilde{\mathcal{G}}$ . Therefore, for  $\delta > 0$  small enough,  $T$  is contracting and the solution of (5) is unique.  $\square$

The following proposition is destined to get solutions of (3) from solutions of (5).

**Proposition 6.** *Let  $\mathcal{G} \in W_D \cap H^2(\Omega)^{3 \times 2}$  be a solution of (5). There exists a unique  $\varphi \in \mathcal{V} \cap H^3(\Omega)^3$ ,*

$$\nabla \varphi = \mathcal{G} \text{ in } \Omega.$$

*There exists  $\alpha \in (0, 1)$ ,  $\varphi \in \mathcal{C}^1(\bar{\Omega})^3 \cap \mathcal{C}^{2,\alpha}(\Omega)$  and  $\varphi$  is a solution of*

$$\bar{p}(\varphi_x)\varphi_{xx} + \bar{q}(\varphi_y)\varphi_{yy} = 0.$$

*Proof.* We use again the BNB lemma. Therefore, there exists a unique  $\varphi \in X$ ,  $\nabla \varphi = \mathcal{G} \in \mathcal{V} \cap H^2(\Omega)^{3 \times 2}$  and thus  $\varphi \in H^3(\Omega)$ . The proof using the BNB lemma is omitted for concision. Using a classical Sobolev embedding, one has  $\varphi \in \mathcal{C}^1(\bar{\Omega})^3$ . Also, one has

$$\bar{A}(\nabla \varphi)\nabla \varphi = 0 \text{ in } \Omega,$$

and therefore

$$\bar{p}(\varphi_x)\varphi_{xx} + \bar{q}(\varphi_y)\varphi_{yy} = 0 \text{ in } \Omega.$$

Note that, as  $\varphi \in \mathcal{C}^0(\bar{\Omega})$ , one can consider  $\varphi|_{\partial\Omega}$ , the restriction of  $\varphi$  to  $\partial\Omega$ , which is continuous on  $\partial\Omega$ , as a Dirichlet boundary condition for

$$\bar{p}(\varphi_x)\psi_{xx} + \bar{q}(\varphi_y)\psi_{yy} = 0 \text{ in } \Omega,$$

which is a linear equation in  $\psi \in H^2(\Omega)^3$  and has for unique solution  $\varphi$ . Note that because there are no cross derivative terms in the expression, the maximum principle can be applied to each individual component. Applying Theorem 6.13 of [10], there exists  $\alpha \in (0, 1)$ ,  $\varphi \in \mathcal{C}^{2,\alpha}(\Omega)^3$ , as done in [16].  $\square$

## 2.6 Constrained problem

The previous section has proved the existence of solutions of (5) under a general Dirichlet boundary condition  $G_D$  imposed on  $\partial\Omega$ . This section gives some restrictions on  $G_D$  so as to prove that (1b) - (1d) are verified in a subset of  $\Omega$ .

**Lemma 7.** *Let  $\varphi \in \mathcal{V}$  be a solution of Equation (5), and let*

$$\begin{cases} u(\varphi) := \varphi_x \cdot \varphi_y, \\ v(\varphi) := \log \left( \left(1 - \frac{1}{4}|\varphi_x|^2\right)|\varphi_y|^2 \right). \end{cases}$$

*We consider  $\bar{u}(\varphi)$  and  $\bar{v}(\varphi)$  as the Lipschitz cut-offs with constant functions of  $u(\varphi)$  and  $v(\varphi)$  when the constraints  $|\varphi_x|^2 \leq 3$  and  $1 < |\varphi_y|^2 \leq 4$  are not verified in  $\Omega$ . Then  $(\bar{u}, \bar{v}) \in H^1(\Omega)^2$  verifies*

$$\begin{cases} \bar{p}\bar{u}_x + 2\bar{v}_y = 0 \\ \bar{q}\bar{u}_y - 2\bar{v}_x = 0, \end{cases}$$

*where  $\bar{p} \equiv \bar{p}(\varphi_x)$  and  $\bar{q} \equiv \bar{q}(\varphi_y)$ .*

*Proof.* Equation (5) is projected onto  $\varphi_x$  and  $\varphi_y$ . One thus has

$$\begin{cases} \bar{p}\varphi_{xx} \cdot \varphi_x + \bar{q}\varphi_{yy} \cdot \varphi_x = 0, \\ \bar{p}\varphi_{xx} \cdot \varphi_y + \bar{q}\varphi_{yy} \cdot \varphi_y = 0. \end{cases}$$

When  $|\varphi_y|^2 \leq 4$  and  $1 < |\varphi_x|^2 \leq 3$ , one has

$$\begin{cases} \bar{u}_x = \varphi_{xx} \cdot \varphi_y + \varphi_x \cdot \varphi_{xy}, \\ \bar{u}_y = \varphi_{xy} \cdot \varphi_y + \varphi_x \cdot \varphi_{yy}, \end{cases}$$



and

$$\begin{cases} \bar{v}_x = -\frac{\bar{p}}{2}\varphi_{xx} \cdot \varphi_x + \frac{\bar{q}}{2}\varphi_y \cdot \varphi_{yx}, \\ \bar{v}_y = -\frac{\bar{p}}{2}\varphi_{xy} \cdot \varphi_x + \frac{\bar{q}}{2}\varphi_y \cdot \varphi_{yy}. \end{cases}$$

Otherwise,  $\bar{u}(\varphi)$  and  $\bar{v}(\varphi)$  are constant and thus  $\nabla\bar{u}(\varphi) = 0$  and  $\nabla\bar{v}(\varphi) = 0$ . Finally,

$$\begin{cases} \bar{p}\bar{u}_x = -2\bar{v}_y, \\ \bar{q}\bar{u}_y = 2\bar{v}_x. \end{cases}$$

□

Lemma 7 proved the existence of a solution to the equation search for  $(u, v) \in H_0^1(\Omega)^2$ ,

$$\begin{cases} \bar{p}u_x + 2v_y = 0 \\ \bar{q}u_y - 2v_x = 0, \end{cases} \quad (9)$$

under the assumption that the  $(\bar{u}, \bar{v})$ , as define in Lemma 7, are zero on  $\partial\Omega$ . The following proposition proves that the solution of (9) is also unique.

**Proposition 8.** *The unique solution in  $H_0^1(\Omega)^2$  of (9) is the pair  $(0, 0)$ .*

*Proof.* We define the solution space  $\mathcal{X} = H_0^1(\Omega)^2$  and the test space  $\mathcal{Y} = H_0^1(\Omega)$ . Let  $\mathcal{C} := \begin{pmatrix} \bar{p} & 0 & 0 & 2 \\ 0 & \bar{q} & -2 & 0 \end{pmatrix}$ ,  $z = (u, v) \in \mathcal{X}$  and  $\tilde{z} \in \mathcal{Y}$ . We define over  $\mathcal{X} \times \mathcal{Y}$  the bilinear form

$$a(z, \tilde{z}) := \int_{\Omega} \mathcal{C} \nabla z \cdot \nabla \tilde{z},$$

where  $\nabla z := \begin{pmatrix} u_x \\ u_y \\ v_x \\ v_y \end{pmatrix}$  and  $a \in \mathcal{L}(\mathcal{X} \times \mathcal{Y})$ . We use the second condition of the BNB lemma, see [8,

Theorem 2.6, p. 85]. Let us assume that

$$\forall z \in \mathcal{X}, \quad a(z, \tilde{z}) = 0.$$

We consider  $z = (\tilde{z}, 0) \in H_0^1(\Omega)^2$ . Therefore,

$$0 = \int_{\Omega} \bar{p}\tilde{z}_x^2 + \bar{q}\tilde{z}_y^2 \geq \|\nabla\tilde{z}\|_{L^2(\Omega)}^2,$$

and thus  $\tilde{z} = 0$ , which proves the second condition of the BNB lemma, which is equivalent to the uniqueness of solutions of (9).

□

The consequence of Lemma 7 and Proposition 8 consists in saying that if  $(\bar{u}, \bar{v}) = 0$  on  $\partial\Omega$ , then  $(\bar{u}, \bar{v}) = 0$  a.e. in  $\Omega$ .

The main result of this section is stated thereafter.

**Theorem 9.** *Let  $G_D \in Y$ , such that a.e. on  $\partial\Omega$ ,*

$$\begin{cases} 0 < |G_D^x|^2 \leq 3, \\ |G_D^y|^2 = \frac{4}{4 - |G_D^x|^2}, \\ G_D^x \cdot G_D^y = 0. \end{cases}$$

Let  $\mathcal{G} \in W_D$  be a solution of (5) and  $\varphi \in \mathcal{V} \cap H^3(\Omega)^3$  such that  $\nabla\varphi = \mathcal{G}$  in  $\Omega$ . There exists  $\Omega' \subset \Omega$ , open set in  $\mathbb{R}^2$  of non-zero Lebesgues measure, such that  $1 < |\varphi_y|^2$  a.e. in  $\Omega'$ . Then,  $\varphi$  is a solution of Equation (1) in  $\Omega'$ .

**Remark 10.**  $\varphi$  is in general not a solution of (1) in all of  $\Omega$ . The physical interpretation is that on the curves where  $1 = |\varphi_y|^2$ , the Miura fold is fully folded. Therefore, the solution computed in the domain where  $1 > |\varphi_y|^2$  is not physical. Section 4.2 below presents a case in which this phenomenon manifests itself.

*Proof.* On  $\partial\Omega$ , let

$$\begin{cases} |\varphi_x|^2 = e, \\ |\varphi_y|^2 = \frac{4}{4-e} > 1, \\ \varphi_x \cdot \varphi_y = 0, \end{cases}$$

with  $0 < e \leq 3$ . As  $\nabla\varphi$  is continuous and  $1 < |\varphi_y|$  on  $\partial\Omega$ , there exists an open domain  $\Omega' \subset \Omega$  such that  $1 < |\varphi_y|$  in  $\Omega'$ . Also,  $\partial\Omega' \supset \partial\Omega$ . Due to the imposed boundary conditions,  $(1 - \frac{1}{4}|\varphi_x|^2)|\varphi_y|^2 = 1$  on  $\partial\Omega$  and  $\bar{u} = 0$  and  $\bar{v} = 0$  on  $\partial\Omega$  in the notations of (9). Using Proposition 8, one has  $(\bar{u}, \bar{v}) = (0, 0)$  a.e. in  $\Omega$  and thus in  $\Omega'$ . Therefore,

$$\varphi_x \cdot \varphi_y = 0 \text{ and } p(\varphi_x)q(\varphi_y) = 4 \text{ a.e. in } \Omega'.$$

Thus, (1b) and (1c) are verified in  $\Omega'$ . Let us now turn to (1d). Similarly as in the proof of Proposition 3,  $\varphi_x$  verifies

$$\bar{p}(\varphi_x)_{xx} + \bar{q}(\varphi_x)_{yy} = -\bar{p}_x\psi_{xx} - \bar{q}_x\psi_{yy} \in L^2(\Omega)^3.$$

Applying the maximum principle, see Theorem 9.1 of [10], to each component of  $\varphi_x$ , one has

$$\max_{\Omega} |\varphi_x| \leq \max_{\partial\Omega} |\varphi_x| = \max_{\partial\Omega} e \leq 3.$$

Similarly,  $\varphi_y$  verifies

$$\bar{p}(\varphi_y)_{xx} + \bar{q}(\varphi_y)_{yy} = -\bar{p}_y\psi_{xx} - \bar{q}_y\psi_{yy} \in L^2(\Omega)^3.$$

Applying the maximum principle, one has

$$\max_{\Omega} |\varphi_y| \leq \max_{\partial\Omega} |\varphi_y| \leq 4.$$

By definition of  $\Omega'$ , one has  $|\varphi_y| > 1$  in  $\Omega'$ . As  $p(\varphi_x)q(\varphi_y) = 4$  a.e. in  $\Omega'$ , then  $|\varphi_x| > 0$  a.e. in  $\Omega'$  and (1d) is verified in  $\Omega'$ .  $\square$

### 3 Numerical method

In [16], an  $H^2$ -conformal finite element method was used to approximate the solutions of (1a). In this paper, Taylor–Hood finite elements are used coupled to a Newton method to approximate (5). The new method is more efficient as much less degrees of freedom (dofs) are required to operate and proved to converge at order one in space.

### 3.1 Discrete Setting

Let  $(\mathcal{T}_h)_h$  be a family of quasi-uniform and shape regular triangulations [8], perfectly fitting  $\Omega$ . For a cell  $c \in \mathcal{T}_h$ , let  $h_c := \text{diam}(c)$  be the diameter of  $c$ . Then, we define  $h := \max_{c \in \mathcal{T}_h} h_c$  as the mesh parameter for a given triangulation  $\mathcal{T}_h$ . Let  $W_h := \mathbb{P}^2(\mathcal{T}_h)^{3 \times 2}$ , where the corresponding interpolator is written as  $\mathcal{I}_h$ , and  $R_h := \{r_h \in \mathbb{P}^1(\mathcal{T}_h)^3 \mid \int_{\Omega} r_h = 0\}$ . We also define the following solution space,

$$W_{hD} := \{G_h \in W_h; G_h = \mathcal{I}_h G_D \text{ on } \partial\Omega\},$$

and its associated homogeneous space

$$W_{h0} := \{G_h \in W_h; G_h = 0 \text{ on } \partial\Omega\}.$$

### 3.2 Discrete problem

We will approximate a solution of (5) by constructing successive approximations of (7). As explained in [9], a direct discretization of (7) would not be coercive. Indeed, in general, if  $G_h \in \mathbb{P}^2(\mathcal{T}_h)^{3 \times 2}$  verifies

$$b(G_h, \tilde{r}_h) = 0, \quad \forall \tilde{r}_h \in R_h,$$

that does not imply that  $G_{h,y}^x = G_{h,x}^y$  a.e. in  $\Omega$ . Therefore, we define the penalty bilinear form  $\mathfrak{p}$  such that for  $G_h, \tilde{G}_h \in W_h$ ,

$$\mathfrak{p}(G_h, \tilde{G}_h) := \eta \int_{\Omega} (G_{h,y}^x - G_{h,x}^y) \cdot (\tilde{G}_{h,y}^x - \tilde{G}_{h,x}^y),$$

where  $\eta > 0$  is a penalty parameter.

**Lemma 11.** *Given  $\mathcal{G}_h \in W_{hD}$ , the equation search for  $(G_h, r_h) \in W_{hD} \times R_h$  such that*

$$a(\mathcal{G}_h; G_h, \tilde{G}_h) + \mathfrak{p}(G_h, \tilde{G}_h) + b(\tilde{G}_h, r_h) = 0, \quad \forall \tilde{G}_h \in W_{h0}, \quad (10a)$$

$$b(G_h, \tilde{r}_h) = 0, \quad \forall \tilde{r}_h \in R_h. \quad (10b)$$

*admits a unique solution. This solution verifies*

$$\|G_h\|_{H^1(\Omega)} \leq C \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)}, \quad (11)$$

where  $C > 0$  is a constant independent of  $\mathcal{G}_h$ .

*Proof.* There exists  $\hat{G}_D \in H^2(\Omega)^{3 \times 2} \subset C^0(\Omega)^{3 \times 2}$  such that  $\hat{G}_D = G_D$  on  $\partial\Omega$ . We define the linear form for  $\tilde{G}_h \in W_{h0}$ ,

$$L(\mathcal{G}_h; \tilde{G}_h) := - \int_{\Omega} \bar{A}(\mathcal{G}_h) \mathcal{I}_h \hat{G}_D \cdot \bar{A}(\mathcal{G}_h) \tilde{G}_h,$$

which is bounded on  $W_{h0}$ , independently of  $\mathcal{G}_h$ . Therefore, we search for  $(\hat{G}_h, r_h) \in W_{h0} \times R_h$ ,

$$\begin{cases} a(\mathcal{G}_h; \hat{G}_h, \tilde{G}_h) + \mathfrak{p}(\hat{G}_h, \tilde{G}_h) + b(\tilde{G}_h, r_h) = L(\mathcal{G}_h; \tilde{G}_h), & \forall \tilde{G}_h \in W_{h0}, \\ b(\hat{G}_h, \tilde{r}_h) = 0, & \forall \tilde{r}_h \in R_h, \end{cases}$$

where  $G_h = \hat{G}_h + \mathcal{I}_h \hat{G}_D$ . Using Lemma 1, one has

$$\begin{aligned} \|\text{div}(\hat{G})\|_{L^2(\Omega)} &\leq (1 - \sqrt{1 - \varepsilon}) \|\text{div}(\hat{G})\|_{L^2(\Omega)} + \sqrt{1 - \varepsilon} \|\text{div}(\hat{G})\|_{L^2(\Omega)}, \\ &\leq \gamma_1 \|\bar{A}(\mathcal{G}) \hat{G}\|_{L^2(\Omega)} + \sqrt{1 - \varepsilon} \|\text{div}(\hat{G})\|_{L^2(\Omega)}, \\ &\leq \gamma_1 \|\bar{A}(\mathcal{G}) \hat{G}\|_{L^2(\Omega)} + \sqrt{1 - \varepsilon} (\|\text{div}(\hat{G})\|_{L^2(\Omega)} + \|\hat{G}_x^y - \hat{G}_y^x\|_{L^2(\Omega)}). \end{aligned}$$

Thus,

$$\frac{1 - \sqrt{1 - \varepsilon}}{\gamma_1} \|\operatorname{div}(\hat{G})\|_{L^2(\Omega)} \leq \|\bar{A}(\mathcal{G})\hat{G}\|_{L^2(\Omega)} + \frac{\sqrt{1 - \varepsilon}}{\gamma_1} \|\hat{G}_x^y - \hat{G}_y^x\|_{L^2(\Omega)}.$$

Therefore, using a Young inequality,

$$\left(\frac{1 - \sqrt{1 - \varepsilon}}{\gamma_1}\right)^2 \|\operatorname{div}(\hat{G})\|_{L^2(\Omega)}^2 \leq (1 + \lambda) \|\bar{A}(\mathcal{G})\hat{G}\|_{L^2(\Omega)}^2 + \left(1 + \frac{1}{\lambda}\right) \frac{1 - \varepsilon}{\gamma_1^2} \|\hat{G}_x^y - \hat{G}_y^x\|_{L^2(\Omega)}^2,$$

where  $\lambda > 0$  is a parameter. Finally,

$$\begin{aligned} \left(\frac{1 - \sqrt{1 - \varepsilon}}{\gamma_1}\right)^2 \|\nabla \hat{G}\|_{L^2(\Omega)}^2 &\leq (1 + \lambda) \|\bar{A}(\mathcal{G})\hat{G}\|_{L^2(\Omega)}^2 \\ &\quad + \left[ \left(\frac{1 - \sqrt{1 - \varepsilon}}{\gamma_1}\right)^2 + \left(1 + \frac{1}{\lambda}\right) \frac{1 - \varepsilon}{\gamma_1^2} \right] \|\hat{G}_x^y - \hat{G}_y^x\|_{L^2(\Omega)}^2. \end{aligned}$$

Thus there exists  $C > 0$ , independent of  $\mathcal{G}_h$ ,

$$C \|\hat{G}_h\|_{H^1(\Omega)}^2 \leq a(\mathcal{G}_h; \hat{G}_h, \hat{G}_h) + \mathfrak{p}(\hat{G}_h, \hat{G}_h),$$

by choosing adequately a value of  $\eta > 0$ , depending on  $\lambda > 0$ .

Regarding  $b$ , we use the result from [9] and the fact that the Taylor–Hood element is conforming to conclude the inf-sup stability of  $b$  over  $W_{h0} \times R_h$ . As a consequence of the BNB lemma, one has

$$\|G_h\|_{H^1(\Omega)} \leq \|\mathcal{I}_h \hat{G}_D\|_{H^1(\Omega)} + \|\hat{G}_h\|_{H^1(\Omega)} \leq C' \|\mathcal{I}_h \hat{G}_D\|_{H^1(\Omega)} \leq C'' \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)},$$

where  $C', C'' > 0$  are independent from  $\mathcal{G}_h$ .  $\square$

**Proposition 12.** *Let  $\delta > 0$ ,  $\|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \leq \delta$ . For  $\delta$  small enough, there exists a unique solution  $(\mathcal{G}_h, r_h) \in W_{hD} \times R_h$  of*

$$a(\mathcal{G}_h; \mathcal{G}_h, \tilde{G}_h) + \mathfrak{p}(\mathcal{G}_h, \tilde{G}_h) + b(\tilde{G}_h, r_h) = 0, \quad \forall \tilde{G}_h \in W_{h0}, \quad (12a)$$

$$b(\mathcal{G}_h, \tilde{r}_h) = 0, \quad \forall \tilde{r}_h \in R_h. \quad (12b)$$

*Proof.* Let us define

$$B_h := \left\{ G_h \in W_{hD}; \|G_h\|_{H^1(\Omega)} \leq C \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \right\},$$

where  $C > 0$  is the constant from Lemma 11. Let  $T_h : W_{hD} \rightarrow W_{hD}$  be such that  $T_h : \mathcal{G}_h \mapsto G_h$ , where  $G_h$  is the unique solution of (10). (11) shows that  $T_h B_h \subset B_h$ . We prove that  $T_h$  is Lipschitz. Let  $\mathcal{G}_h, \hat{\mathcal{G}}_h \in W_{hD}$ ,  $G_h := T_h \mathcal{G}_h$  and  $\hat{G}_h := T_h \hat{\mathcal{G}}_h$  with associated Lagrange multipliers  $r_h, \hat{r}_h \in R_h$ . Using the coercivity of  $a(\mathcal{G}_h) + \mathfrak{p}$ , with coercivity constant  $c > 0$ , one has

$$\begin{aligned} c \|\nabla G_h - \nabla \hat{G}_h\|_{L^2(\Omega)}^2 &\leq a(\mathcal{G}_h; G_h - \hat{G}_h, G_h - \hat{G}_h) + \mathfrak{p}(G_h - \hat{G}_h, G_h - \hat{G}_h), \\ &= a(\mathcal{G}_h; G_h, G_h - \hat{G}_h) + \mathfrak{p}(G_h, G_h - \hat{G}_h) \\ &\quad - a(\mathcal{G}_h; \hat{G}_h, G_h - \hat{G}_h) - \mathfrak{p}(\hat{G}_h, G_h - \hat{G}_h), \\ &= -b(G_h - \hat{G}_h, r_h) - a(\mathcal{G}_h; \hat{G}_h, G_h - \hat{G}_h) - \mathfrak{p}(\hat{G}_h, G_h - \hat{G}_h), \\ &= -b(G_h - \hat{G}_h, \hat{r}_h) - a(\mathcal{G}_h; \hat{G}_h, G_h - \hat{G}_h) - \mathfrak{p}(\hat{G}_h, G_h - \hat{G}_h), \\ &= a(\hat{\mathcal{G}}_h; \hat{G}_h, G_h - \hat{G}_h) - a(\mathcal{G}_h; \hat{G}_h, G_h - \hat{G}_h), \end{aligned}$$

as  $(G_h, r_h)$  and  $(\hat{G}_h, \hat{r}_h)$  are solutions of (12). Therefore, one has

$$\begin{aligned}
 c\|\nabla G_h - \nabla \hat{G}_h\|_{L^2(\Omega)}^2 &\leq \int_{\Omega} \bar{A}(\hat{\mathcal{G}}_h) \hat{G}_h \cdot \bar{A}(\hat{\mathcal{G}}_h) (G_h - \hat{G}_h) - \int_{\Omega} \bar{A}(\mathcal{G}_h) \hat{G}_h \cdot \bar{A}(\mathcal{G}_h) (G_h - \hat{G}_h), \\
 &= \int_{\Omega} (\bar{A}(\hat{\mathcal{G}}_h) - \bar{A}(\mathcal{G}_h)) \hat{G}_h \cdot \bar{A}(\hat{\mathcal{G}}_h) (G_h - \hat{G}_h) \\
 &\quad + \int_{\Omega} \bar{A}(\mathcal{G}_h) \hat{G}_h \cdot (\bar{A}(\hat{\mathcal{G}}_h) - \bar{A}(\mathcal{G}_h)) (G_h - \hat{G}_h), \\
 &\leq \|(\bar{A}(\mathcal{G}_h) - \bar{A}(\hat{\mathcal{G}}_h)) \hat{G}_h\|_{L^2(\Omega)} \|\bar{A}(\mathcal{G}_h) (\hat{G}_h - G_h)\|_{L^2(\Omega)} \\
 &\quad + \|\bar{A}(\mathcal{G}_h) \hat{G}_h\|_{L^2(\Omega)} \|(\bar{A}(\mathcal{G}_h) - \bar{A}(\hat{\mathcal{G}}_h)) (G_h - \hat{G}_h)\|_{L^2(\Omega)}, \\
 &\leq KC' \|\mathcal{G}_h - \hat{\mathcal{G}}_h\|_{L^2(\Omega)} \|\nabla \hat{G}_h\|_{L^2(\Omega)} \|\nabla \hat{G}_h - \nabla G_h\|_{L^2(\Omega)} \\
 &\quad + KC'' \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \|\mathcal{G}_h - \hat{\mathcal{G}}_h\|_{L^2(\Omega)} \|\nabla \hat{G}_h - \nabla G_h\|_{L^2(\Omega)},
 \end{aligned}$$

because  $\bar{A}$  is  $K$ -Lipschitz, and where  $C', C'' > 0$  are constants independent of  $\mathcal{G}_h$  and  $\hat{\mathcal{G}}_h$ . Using a generalized Hölder inequality with  $p > 2$  and  $q < 2$  such that  $p^{-1} + q^{-1} = 2^{-1}$ , one has

$$\begin{aligned}
 c\|\nabla G_h - \nabla \hat{G}_h\|_{L^2(\Omega)}^2 &\leq KC' \|\mathcal{G}_h - \hat{\mathcal{G}}_h\|_{L^p(\Omega)} \|\nabla \hat{G}_h\|_{L^q(\Omega)} \|\nabla \hat{G}_h - \nabla G_h\|_{L^2(\Omega)} \\
 &\quad + KC'' \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \|\mathcal{G}_h - \hat{\mathcal{G}}_h\|_{L^p(\Omega)} \|\nabla \hat{G}_h - \nabla G_h\|_{L^q(\Omega)}, \\
 &\leq KC''' \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \|\mathcal{G}_h - \hat{\mathcal{G}}_h\|_{H^1(\Omega)} \|\nabla \hat{G}_h - \nabla G_h\|_{L^2(\Omega)},
 \end{aligned}$$

where  $C''' > 0$  is a constant independent of  $\mathcal{G}_h$  and  $\hat{\mathcal{G}}_h$ . Therefore,

$$c\|\nabla G_h - \nabla \hat{G}_h\|_{L^2(\Omega)} \leq C''' \delta K \|\mathcal{G}_h - \hat{\mathcal{G}}_h\|_{H^1(\Omega)}.$$

The existence of the fixed point is provided by the Brouwer fixed point theorem, see [4], and the uniqueness by the fact that, for  $\delta$  small enough,  $T_h$  is contracting.  $\square$

### 3.3 Recomputing $\varphi_h$

Following [9], the approximation space is written as

$$V_h := \left\{ \psi_h \in \mathbb{P}^2(\mathcal{T}_h)^3; \int_{\Omega} \psi_h = 0 \right\}.$$

The least-squares method is used. The bilinear form  $\mathbf{a} \in \mathcal{L}(V_h^2)$  is defined as

$$\mathbf{a}(\varphi_h, \tilde{\varphi}_h) := \int_{\Omega} \nabla \varphi_h \cdot \nabla \tilde{\varphi}_h.$$

It is classically coercive over  $V_h$ . The linear form  $\mathbf{b} \in \mathcal{L}(V_h)$  is defined as

$$\mathbf{b}(\tilde{\varphi}_h) := \int_{\Omega} \mathcal{G}_h \cdot \nabla \tilde{\varphi}_h.$$

**Lemma 13.** *There exists a unique  $\varphi_h \in V_h$  solution of*

$$\mathbf{a}(\varphi_h, \tilde{\varphi}_h) = \mathbf{b}(\tilde{\varphi}_h), \quad \forall \tilde{\varphi}_h \in V_h. \tag{13}$$

The proof is omitted for concision, as this is a classical result.

### 3.4 Convergence of the FEM

**Theorem 14.** *Let  $\delta > 0$ ,  $\|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)}$ . For  $\delta$  and  $h$  small enough, the sequence  $(\varphi_h)_h \in V_h^{\mathbb{N}}$  of solutions of (12) converges towards  $\varphi \in \mathcal{V}$ , solution of (1a), with the following convergence estimate,*

$$\|\varphi - \varphi_h\|_{H^1(\Omega)} \leq Ch \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)}, \quad (14)$$

where  $C > 0$  is a constant independent of  $\varphi$ .

*Proof.* Using a poincaré inequality, there exists  $C > 0$ ,

$$\begin{aligned} \|\varphi - \varphi_h\|_{H^1(\Omega)} &\leq C \|\nabla\varphi - \nabla\varphi_h\|_{L^2(\Omega)}, \\ &\leq C \|\nabla\varphi - \mathcal{G}\|_{L^2(\Omega)} + C \|\mathcal{G} - \mathcal{G}_h\|_{L^2(\Omega)} + C \|\mathcal{G}_h - \nabla\varphi_h\|_{L^2(\Omega)}, \\ &= C \|\mathcal{G} - \mathcal{G}_h\|_{L^2(\Omega)}. \end{aligned}$$

Therefore, (14) is proved by getting a convergence rate on  $\mathcal{G}_h \rightarrow \mathcal{G}$ , when  $h \rightarrow 0$ . Using the coercivity of  $a(\mathcal{G}_h) + \mathfrak{p}$  over  $W_{h0}^2$ , there exists  $c > 0$ , independent of  $\mathcal{G}_h$  such that

$$\begin{aligned} c \|\nabla\mathcal{G} - \nabla\mathcal{G}_h\|_{L^2(\Omega)}^2 &\leq a(\mathcal{G}_h; \mathcal{G} - \mathcal{G}_h, \mathcal{G} - \mathcal{G}_h) + \mathfrak{p}(\mathcal{G} - \mathcal{G}_h, \mathcal{G} - \mathcal{G}_h), \\ &\leq a(\mathcal{G}_h; \mathcal{G} - \mathcal{G}_h, \mathcal{G} - \mathcal{I}_h\mathcal{G}) + a(\mathcal{G}_h; \mathcal{G} - \mathcal{G}_h, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + \mathfrak{p}(\mathcal{G} - \mathcal{G}_h, \mathcal{G} - \mathcal{G}_h). \end{aligned}$$

One thus has

$$\begin{aligned} a(\mathcal{G}_h; \mathcal{G} - \mathcal{G}_h, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) &= a(\mathcal{G}_h; \mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + \mathfrak{p}(\mathcal{G}_h, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r_h), \\ &= a(\mathcal{G}_h; \mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) - a(\mathcal{G}; \mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + \mathfrak{p}(\mathcal{G}_h, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r_h - r), \end{aligned}$$

because  $\mathcal{G}_h$  solves (12) and  $\mathcal{G}$  solves (7). Thus,

$$\begin{aligned} a(\mathcal{G}_h; \mathcal{G} - \mathcal{G}_h, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) &= \int_{\Omega} (\bar{A}(\mathcal{G}_h) - \bar{A}(\mathcal{G}))\mathcal{G} \cdot \bar{A}(\mathcal{G}_h)(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + \mathfrak{p}(\mathcal{G}_h - \mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) \\ &\quad + \int_{\Omega} \bar{A}(\mathcal{G})\mathcal{G} \cdot (\bar{A}(\mathcal{G}_h) - \bar{A}(\mathcal{G}))(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r_h - r), \end{aligned}$$

since  $\mathfrak{p}(\mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) = 0$ . Therefore,

$$\begin{aligned} c \|\nabla\mathcal{G} - \nabla\mathcal{G}_h\|_{L^2(\Omega)}^2 &\leq a(\mathcal{G}_h; \mathcal{G} - \mathcal{G}_h, \mathcal{G} - \mathcal{I}_h\mathcal{G}) + \mathfrak{p}(\mathcal{G}_h - \mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}) + b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r_h - r) \\ &\quad + \int_{\Omega} (\bar{A}(\mathcal{G}_h) - \bar{A}(\mathcal{G}))\mathcal{G} \cdot \bar{A}(\mathcal{G}_h)(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + \int_{\Omega} \bar{A}(\mathcal{G})\mathcal{G} \cdot (\bar{A}(\mathcal{G}_h) - \bar{A}(\mathcal{G}))(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h), \end{aligned}$$

Let us write  $T_1$  the first line of the right-hand side of the previous equation and  $T_2$  the second. One has

$$T_1 \leq C \|\nabla\mathcal{G} - \nabla\mathcal{G}_h\|_{L^2(\Omega)} \|\mathcal{G} - \mathcal{I}_h\mathcal{G}\|_{H^1(\Omega)} + |b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r_h - r)|,$$

where  $C > 0$  is a constant independent of  $\mathcal{G}$  or  $\mathcal{G}_h$ . Let us bound the remaining term. One has

$$\begin{aligned} b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r_h - r) &= b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r_h) - b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r), \\ &= -a(\mathcal{G}_h; \mathcal{G}_h, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) - \mathfrak{p}(\mathcal{G}_h - \mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) + a(\mathcal{G}; \mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h), \end{aligned}$$

since  $\mathcal{G}_y^x = \mathcal{G}_x^y$  a.e. in  $\Omega$  and thus  $\mathfrak{p}(\mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) = 0$ . One also has

$$\begin{aligned} a(\mathcal{G}; \mathcal{G}, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) - a(\mathcal{G}_h; \mathcal{G}_h, \mathcal{I}_h\mathcal{G} - \mathcal{G}_h) &= \int_{\Omega} (\bar{A}(\mathcal{G}) - \bar{A}(\mathcal{G}_h))\mathcal{G} \cdot \bar{A}(\mathcal{G})(\mathcal{G}_h - \mathcal{I}_h\mathcal{G}) \\ &+ \int_{\Omega} \bar{A}(\mathcal{G}_h)\mathcal{G} \cdot (\bar{A}(\mathcal{G}) - \bar{A}(\mathcal{G}_h))(\mathcal{G}_h - \mathcal{I}_h\mathcal{G}) + \int_{\Omega} \bar{A}(\mathcal{G}_h)(\mathcal{G} - \mathcal{G}_h) \cdot \bar{A}(\mathcal{G}_h)(\mathcal{G}_h - \mathcal{I}_h\mathcal{G}). \end{aligned}$$

Therefore,

$$\begin{aligned} |b(\mathcal{I}_h\mathcal{G} - \mathcal{G}_h, r_h - r)| &\leq C' \|\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)} \|\mathcal{G} - \mathcal{I}_h\mathcal{G}\|_{H^1(\Omega)} \\ &+ C'' K \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \|\mathcal{G} - \mathcal{G}_h\| \|\nabla\mathcal{G}_h - \nabla\mathcal{I}_h\mathcal{G}\|_{L^2(\Omega)}, \\ &\leq C' \|\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)} \|\mathcal{G} - \mathcal{I}_h\mathcal{G}\|_{H^1(\Omega)} \\ &+ C''' K \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \|\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)} \|\mathcal{G}_h - \mathcal{I}_h\mathcal{G}\|_{H^1(\Omega)}, \end{aligned}$$

since  $\bar{A}$  is  $K$ -Lipschitz,  $C', C'', C''' > 0$  are constants independent of  $h$  and using a computation similar to the proof of Proposition 12. Let us now bound  $T_2$ . One has

$$T_2 \leq KC^\dagger \|G_D\|_{H^{\frac{3}{2}}(\partial\Omega)} \|\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)} \|\mathcal{I}_h\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)},$$

where  $C^\dagger > 0$  is a constant independent of  $h$ . Summing up the previous results, one has

$$\begin{aligned} \|\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)} &\leq \frac{\delta K'}{c} \|\mathcal{I}_h\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)} + \frac{K''}{c} \|\mathcal{I}_h\mathcal{G} - \mathcal{G}\|_{H^1(\Omega)}, \\ &\leq \frac{\delta K'}{c} \left( \|\mathcal{I}_h\mathcal{G} - \mathcal{G}\|_{H^1(\Omega)} + \|\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)} \right) + \frac{K''}{c} \|\mathcal{I}_h\mathcal{G} - \mathcal{G}\|_{H^1(\Omega)}, \end{aligned}$$

with  $K', K'' > 0$  are constants independent of  $h$ . Let  $\iota := \frac{\delta K'}{c}$ . For  $\delta > 0$  small enough,  $\iota < 1$  and thus,

$$\|\mathcal{G} - \mathcal{G}_h\|_{H^1(\Omega)} \leq \frac{1}{1 - \iota} \left( \iota + \frac{K''}{c} \right) \|\mathcal{I}_h\mathcal{G} - \mathcal{G}\|_{H^1(\Omega)}.$$

As  $\mathcal{G} \in (H^2(\Omega))^3$ , one has only the classical interpolation error [3],

$$\|\mathcal{I}_h\mathcal{G} - \mathcal{G}\|_{H^1(\Omega)} \leq K''' h |\mathcal{G}|_{H^2(\Omega)},$$

where  $K''' > 0$  is independent of  $\mathcal{G}$ . Hence the announced result.  $\square$

**Remark 15.** *Theorem 14 proves that the proposed method converges at order one in  $h$ . However, numerically, we observe an improved convergence rate of two. This suggests that (14) could be improved to*

$$\|\varphi - \varphi_h\| \leq Ch^2 |\mathcal{G}|_{H^1(\Omega)}.$$

*Such a result might be proved using an Aubin–Nitsche trick, as  $a(\mathcal{G}_h)$  is symmetric, but in a nonlinear setting. See [7], for instance.*

## 4 Numerical examples

The method uses the automatic differentiation of Firedrake [23] to solve (12) with a Newton method. The boundary condition  $G_h = \mathcal{I}_h G_D$  on  $\partial\Omega$  is imposed strongly. The stopping criterion based on the relative residual in  $H^1$ -norm is set to  $\varepsilon := 10^{-8}$ . The penalty parameter for  $\mathbf{p}$  is taken as  $\eta := 10$ , for all computations. The initial guess  $(\mathcal{G}_{h,0}, r_{h,0}) \in W_{hD} \times R_h$  for the Newton method is computed as the solution of

$$\begin{cases} \int_{\Omega} \nabla \mathcal{G}_{h,0} \cdot \nabla \tilde{G}_h + \mathbf{p}(\mathcal{G}_{h,0}, \tilde{G}_h) + b(\tilde{G}_h, r_{h,0}) = 0, & \forall \tilde{G}_h \in W_{h0}, \\ b(\mathcal{G}_{h,0}, \tilde{r}_h) = 0, & \forall \tilde{r}_h \in R_h. \end{cases}$$

To handle in practice the constraint  $\int_{\Omega} \varphi_h = 0$  in computing the solution of (13), we follow [8, Remark 3.83] and fix the position of one node of the mesh to zero.

### 4.1 Hyperboloid

This test case comes from [15]. The reference solution is

$$\phi(x, y) = (\rho(x) \cos(\alpha y), \rho(x) \sin(\alpha y), z(x))^T, \text{ where } \begin{cases} \rho(x) = \sqrt{4c_0^2 x^2 + 1}, \\ z(x) = 2s_0 x, \\ \alpha = (1 - s_0^2)^{-1/2}, \end{cases}$$

$c_0 = \cos(\frac{\theta}{2})$ ,  $s_0 = \sin(\frac{\theta}{2})$  and  $\theta \in (0, \frac{2\pi}{3})$ . The domain is  $\Omega = (-s_0^*, s_0^*) \times (0, \frac{2\pi}{\alpha})$ , where  $s_0^* = \sin\left(0.5 \cos^{-1}\left(0.5 \cos\left(\frac{\theta}{2}\right)^{-1}\right)\right)$ . Structured meshes, periodic in  $y$ , are used. This translates into the fact that the dofs on the lines of equation  $y = 0$  and  $y = \frac{2\pi}{\alpha}$  of  $\partial\Omega$  are one and the same, but remain unknown.  $\nabla\phi$  is used as Dirichlet boundary condition on the lines of equations  $x = -s_0^*$  and  $x = s_0^*$ . A convergence test is performed for  $\theta = \frac{\pi}{2}$ . Table 1 contains the errors and estimated convergence rate.

$h$	nb dofs	nb iterations	$\ \mathcal{G}_h - \mathcal{G}\ _{H^1}$	rate	$\ \mathcal{G}_h - \mathcal{G}\ _{L^2}$	rate
0.178	34,125	7	1.064e-02	-	2.577e-04	-
0.00889	135,750	7	2.658e-03	2.01	3.191e-05	3.03
0.00444	541,500	7	6.642e-04	2.00	3.977e-06	3.01

Table 1: Hyperboloid: estimated convergence rate and number of Newton iterations.

The convergence rate is estimated using the formula

$$2 \log\left(\frac{e_1}{e_2}\right) \log\left(\frac{\text{card}(\mathcal{T}_{h_1})}{\text{card}(\mathcal{T}_{h_2})}\right)^{-1},$$

where  $e_1$  and  $e_2$  are the errors. The number of Newton iterations is also reported. The results in Table 1 are compatible with the second order rate announced in Remark 15. They are even better as  $\phi$  is more regular than just  $H^3(\Omega)$ .

### 4.2 Annulus

The domain is  $\Omega = (0, \frac{3}{4}) \times (0, 2\pi)$ . A structured mesh, periodic in the  $y$  direction, is used to mesh  $\Omega$ . The mesh has a size  $h = 0.015$  and contains 181,800 dofs. The Dirichlet boundary conditions



are

$$\begin{cases} G_D^x = (ax + 1)e_r, \\ G_D^y = \frac{4}{4 - |G_D^x|^2}e_\theta, \end{cases}$$

where  $(e_r, e_\theta)$  is the polar basis of  $\mathbb{R}^2$  and  $a = 0.675$ . The Newton method converges after 7 iterations. The resulting surface is presented in Figure 3. In Figure 4, one can observe that  $\Omega' \not\subseteq \Omega$ .




Figure 3: Annulus: Computed surface.

Indeed, in a large band in the middle of the domain, one can notice that  $|\mathcal{G}_{h,x}| \simeq 0$ . The violation of the constraints in  $\Omega \setminus \Omega'$  is interpreted as being the result of having Dirichlet boundary conditions on all of  $\partial\Omega$ . Indeed, as stated in Remark 10, the constraints are not verified after the pattern fully folds on a curve, which corresponds to  $|\varphi_x| = 0$ . As Dirichlet boundary conditions are imposed on all of  $\partial\Omega$ , the pattern cannot relax in any part of the domain, as would have been possible with a part of the boundary having stress-free (homogeneous Neumann) boundary conditions. This leads to over-constraining the pattern and thus to a solution that is not physical in some parts of the domain.

### 4.3 Axysymmetric surface

This example comes from [15]. The domain is  $(0, 2\pi) \times (0, 4)$ . A structured mesh, periodic in the  $x$  direction, is used. The mesh has a size  $h = 0.0628$  and contains 541,500 dofs. The reference solution writes

$$\phi(x, y) := (\rho(y) \cos(x), \rho(y) \sin(x), z(x)),$$

where  $\rho$  is a solution of

$$\ddot{\rho} = \frac{4\rho}{(4 - \rho^2)^2}.$$

An explicit Runge–Kutta method of order 5 (see [12]) is used to integrate this ODE with initial conditions  $\rho_0 = 0.1$  and  $\dot{\rho}_0 = 0$ . Because the reference solution is not known analytically, the Dirichlet boundary condition is imposed weakly using a least-squares penalty in order to remove numerical




Figure 4: Annulus: left:  $\Omega'$  (red) and  $\Omega \setminus \Omega'$  (blue), middle:  $|\mathcal{G}_{h,x}|$  and right:  $|\mathcal{G}_{h,y}|$ .

issues. Let  $\mathcal{E}_h$  the set of the edges of  $\mathcal{T}_h$ . The set  $\mathcal{E}_h$  is partitioned as  $\mathcal{E}_h := \mathcal{E}_h^i \cup \mathcal{E}_h^b$ , where for all  $e \in \mathcal{E}_h^b$ ,  $e \subset \partial\Omega$  and  $\cup_{e \in \mathcal{E}_h^b} e = \partial\Omega$ . Let  $h_e := \text{diam}(e)$ , where  $e \in \mathcal{E}_h$ . Therefore, we search for  $(\mathcal{G}_h, r_h) \in W_h \times R_h$ ,

$$\begin{cases} a(\mathcal{G}_h; \mathcal{G}_h, \tilde{G}_h) + \mathfrak{p}(\mathcal{G}_h, \tilde{G}_h) + b(\tilde{G}_h, r_h) + \sum_{e \in \mathcal{E}_h^b} \frac{\eta}{h_e} \int_e \mathcal{G}_h : \tilde{G}_h \\ = \sum_{e \in \mathcal{E}_h^b} \frac{\eta}{h_e} \int_e \mathcal{I}_h G_D : \tilde{G}_h, \quad \forall \tilde{G}_h \in W_{h0}, \\ b(\mathcal{G}_h, \tilde{r}_h) = 0, \quad \forall \tilde{r}_h \in R_h. \end{cases}$$

3 Newton iterations are necessary to reach converge. Figure 5 shows the computed surface, which




Figure 5: Axysymmetric surface: Computed surface.

looks as expected in [15].

#### 4.4 A failing example

The domain is  $\Omega = (-1, 1)^2$ . It is meshed with an unstructured triangular mesh of size  $h = 0.025$ . The Dirichlet boundary conditions for this test case derive from a hyperbolic paraboloid for which

$$f(x, y) = \left( x, y, \frac{x^2}{a^2} - \frac{y^2}{b^2} \right)^T.$$

The chosen Dirichlet boundary conditions are

$$\begin{cases} G_D^x = f_x, \\ G_D^y = \frac{4}{4 - |f_x|^2} \mathfrak{f}, \end{cases}$$

where  $a = 2$ ,  $b = \frac{4}{3}$  and  $\mathfrak{f} := f_y - \frac{f_y f_x}{|f_x|^2} f_x$ . The boundary condition seems to verify all the hypotheses of Theorem 9. However, when we compute the initial guess  $(\mathcal{G}_{h,0}, r_{h,0})$ , the surface is degenerate and the subsequent Newton iterations diverge. Let  $\mathfrak{G}_h \in \{G_h \in P^2(\mathcal{T}_h)^{3 \times 2}; G_h = \mathcal{I}_h G_D \text{ on } \partial\Omega\}$  be the solution of

$$\int_{\Omega} \nabla \mathfrak{G}_h \cdot \nabla \tilde{G}_h = 0, \quad \forall \tilde{G}_h \in W_{h0}.$$

This computation is performed with 203,502 dofs. Figure 6 shows the resulting surface, in terms of  $\varphi_h$ . It is not degenerate. But after careful consideration, one notices that  $\mathfrak{G}_{h,y}^x = \mathfrak{G}_{h,x}^y$  is not verified in  $\Omega$ , as showed in Figure 7. Therefore, the chosen Dirichlet boundary condition is not the trace of a function  $G \in W$  such that  $G_y^x = G_x^y$  a.e. in  $\Omega$ , which explains why a solution cannot be computed.




Figure 6: Saddle shape: Computed surface.




Figure 7: Saddle shape:  $|\mathfrak{G}_{h,x}^y - \mathfrak{G}_{h,y}^x|^2$ .

## 5 Conclusion

In this paper, the existence and uniqueness of solutions of the constrained system of equations describing a Miura surface are proved under specific boundary conditions that still leave some freedom for design choices, specifically  $\alpha$  and  $\mathcal{N}$  (see Figure 2). Then, a numerical method based on a mixed formulation, Taylor–Hood elements, and a Newton method is introduced to approximate Miura surfaces. The method is proved to converge at order one in  $H^1$  norm in the space discretization parameter  $h$  and an order two is observed in practice. Some numerical tests are performed and show the robustness of the method.

Future work include being able to compute the Miura surface “closest” to a given target surface and to be able to produce Miura tessellations with a given pattern size from Miura surfaces.

## Code availability

The code is available at <https://github.com/marazzaf/Miura.git>

## Acknowledgment

The author would like to thank Hussein Nassar from University of Missouri for stimulating discussions that lead to the content of this paper. The author would also like to thank Zhaonan Dong from Inria Paris for fruitful discussions.

## Funding

This work is supported by the US National Science Foundation under grant number OIA-1946231 and the Louisiana Board of Regents for the Louisiana Materials Design Alliance (LAMDA).

## References

- [1] P. Badagavi, V. Pai, and A. Chinta. Use of origami in space science and various other fields of science. In *2017 2nd IEEE International Conference on Recent Trends in Electronics, Information & Communication Technology (RTEICT)*, pages 628–632. IEEE, 2017.
- [2] E. Boatti, N. Vasios, and K. Bertoldi. Origami metamaterials for tunable thermal expansion. *Advanced Materials*, 29(26):1700360, 2017.
- [3] S. Brenner, L. Scott, and L. Scott. *The mathematical theory of finite element methods*, volume 3. Springer, 2008.
- [4] H. Brézis. *Functional analysis, Sobolev spaces and partial differential equations*, volume 2. Springer, 2011.
- [5] P. G. Ciarlet. An introduction to differential geometry with applications to elasticity. *Journal of Elasticity*, 78(1):1–215, 2005.
- [6] M. Costabel and M. Dauge. Maxwell and lamé eigenvalues on polyhedra. *Mathematical methods in the applied sciences*, 22(3):243–258, 1999.
- [7] M. Dobrowolski and R. Rannacher. Finite element methods for nonlinear elliptic systems of second order. *Mathematische Nachrichten*, 94(1):155–172, 1980.

- 
- [8] A. Ern and J.-L. Guermond. *Theory and practice of finite elements*, volume 159. Springer Science & Business Media, 2013.
- [9] D. Gallistl. Variational formulation and numerical analysis of linear elliptic equations in nondivergence form with Cordès coefficients. *SIAM Journal on Numerical Analysis*, 55(2):737–757, 2017.
- [10] D. Gilbarg and N. Trudinger. *Elliptic partial differential equations of second order*, volume 224. Springer, 2015.
- [11] P. Grisvard. *Elliptic problems in nonsmooth domains*. SIAM, 2011.
- [12] E. Hairer, M. Hochbruck, A. Iserles, and C. Lubich. Geometric numerical integration. *Oberwolfach Reports*, 3(1):805–882, 2006.
- [13] R. J. Lang. *Origami 4*. CRC Press, 2009.
- [14] R. J. Lang. *Twists, tilings, and tessellations: mathematical methods for geometric origami*. AK Peters/CRC Press, 2017.
- [15] A. Lebé, L. Monasse, and H. Nassar. Fitting surfaces with the Miura tessellation. In *7th International Meeting on Origami in Science, Mathematics and Education (7OSME)*, volume 4, page 811. Tarquin, 2018.
- [16] F. Marazzato.  $H^2$ -conformal approximation of Miura surfaces. *arXiv.org*, 2022.
- [17] K. Miura. Proposition of pseudo-cylindrical concave polyhedral shells. *ISAS report/Institute of Space and Aeronautical Science, University of Tokyo*, 34(9):141–163, 1969.
- [18] J. Morgan, S. P. Magleby, and L. L. Howell. An approach to designing origami-adapted aerospace mechanisms. *Journal of Mechanical Design*, 138(5), 2016.
- [19] H. Nassar, A. Lebé, and L. Monasse. Curvature, metric and parametrization of origami tessellations: theory and application to the eggbox pattern. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 473(2197):20160705, 2017.
- [20] H. Nassar, A. Lebé, and L. Monasse. Macroscopic deformation modes of origami tessellations and periodic pin-jointed trusses: the case of the eggbox. In *Proceedings of IASS Annual Symposia*, volume 2017, pages 1–9. International Association for Shell and Spatial Structures (IASS), 2017.
- [21] J. Overvelde, T. A. De Jong, Y. Shevchenko, S. A. Begera, G. M. Whitesides, J. C. Weaver, C. Hoberman, and K. Bertoldi. A three-dimensional actuated origami-inspired transformable metamaterial with multiple degrees of freedom. *Nature communications*, 7(1):1–8, 2016.
- [22] A. Rafsanjani, K. Bertoldi, and A. R. Studart. Programming soft robots with flexible mechanical metamaterials. *Science Robotics*, 4(29):eaav7874, 2019.
- [23] F. Rathgeber, D. Ham, L. Mitchell, M. Lange, F. Luporini, A. McRae, G.-T. Bercea, G. Markall, and P. Kelly. Firedrake: automating the finite element method by composing abstractions. *ACM Transactions on Mathematical Software (TOMS)*, 43(3):1–27, 2016.
- [24] M. Schenk and S. Guest. Geometry of Miura-folded metamaterials. *Proceedings of the National Academy of Sciences*, 110(9):3276–3281, 2013.
-

- [25] I. Smears and E. Suli. Discontinuous Galerkin finite element approximation of nondivergence form elliptic equations with Cordès coefficients. *SIAM Journal on Numerical Analysis*, 51(4):2088–2106, 2013.
- [26] Z. Wei, Z. Guo, L. Dudte, H. Liang, and L. Mahadevan. Geometric mechanics of periodic pleated origami. *Physical review letters*, 110(21):215501, 2013.
- [27] A. Wickeler and H. Naguib. Novel origami-inspired metamaterials: Design, mechanical testing and finite element modelling. *Materials & Design*, 186:108242, 2020.