

EXPANSIVITY AND STRUCTURALLY STABILITY OF COMPOSITION OPERATORS ON ORLICZ SPACES

Y. ESTAREMI

ABSTRACT. The notions of expansivity and positively expansivity for composition operators on Orlicz spaces are investigated. We provide some necessary and sufficient conditions for expansivity, positively expansivity and uniformly expansivity of such operators. Moreover, we provide some equivalent conditions for these concepts in the case that the system is dissipative. Our results are essentially based on the results of [16].

1. Introduction and Preliminaries

The dynamics of continuous linear operators on Banach spaces have been extensively studied during the last three decades. Many connections between dynamical systems and other areas of mathematics, such as ergodic theory, number theory and geometry of Banach spaces have been established. Interested readers can refer to [4, 8, 1, 2, 6, 14]. In [11] Eisenberg and Hedlund investigated relationships between expansivity and spectrum of operators on Banach spaces. Also, in [17] the authors established relationships between notions of expansivity and hypercyclicity, supercyclicity, Li–Yorke chaos and shadowing. In the case that the Banach space is c_0 or l_p ($1 \leq p < \infty$), they have given complete characterizations of weighted shifts which satisfy various notions of expansivity. In [16], the authors investigated the two notions of expansivity and strong structural stability for composition operators on $L^p(\mu)$ -spaces, $1 \leq p < \infty$. Necessary and sufficient conditions for composition operators to be expansive have provided, both in the general and the dissipative case. They also have shown that, in the dissipative setting, the shadowing property implies the strong structural stability and they have proved that these two notions are equivalent under the extra hypothesis of positive expansivity. In this paper we investigate the notions of expansivity and positively expansivity for composition operators on Orlicz spaces. Indeed, first we extend the definitions of some concepts like, dissipative system and bounded distortion property to the setting of Orlicz spaces, in way that it coincides to the definition of these concepts in $L^p(\mu)$ -spaces, in special case. Then we provide some necessary and sufficient conditions for expansivity, positively expansivity and uniformly expansivity of such operators. Moreover, we provide some equivalent conditions for these concepts in the case that the system is dissipative.

In the sequel, for the convenience of the reader, we recall some essential facts on Orlicz spaces for later use. For more details on Orlicz spaces, see [15, 18].

2020 *Mathematics Subject Classification.* 47B33, 37B65.

Key words and phrases. Orlicz spaces, Composition operators, Expansivity, structural stability.

A function $\Phi : \mathbb{R} \rightarrow [0, \infty]$ is called a *Young's function* if Φ is convex, $\Phi(-x) = \Phi(x)$, $\Phi(0) = 0$ and $\lim_{x \rightarrow \infty} \Phi(x) = +\infty$. With each *Young's function* Φ , one can associate another convex function $\Psi : \mathbb{R} \rightarrow [0, \infty]$ having similar properties defined by

$$\Psi(y) = \sup\{x \mid y \mid -\Phi(x) : x \geq 0\}, \quad y \in \mathbb{R}.$$

The function Ψ is called the *complementary function* to Φ . It follows from the definition that $\Psi(0) = 0$, $\Psi(-y) = \Psi(y)$ and $\Psi(\cdot)$ is a convex increasing function satisfying $\lim_{y \rightarrow \infty} \Psi(y) = +\infty$.

Let Φ be a *Young's function*. Then we say Φ satisfies the Δ_2 -condition or Φ is Δ_2 -regular, if $\Phi(2x) \leq K\Phi(x)$ ($x \geq x_0$) for some constants $K > 0$ and $x_0 > 0$. Also, Φ satisfies the Δ_2 -condition globally if $\Phi(2x) \leq K\Phi(x)$ ($x \geq 0$) for some $K > 0$. If $x_0 = 0$, these conditions are said to hold globally.

For a given complete σ -finite measure space (X, \mathcal{F}, μ) , let $L^0(\mathcal{F})$ be the linear space of equivalence classes of \mathcal{F} -measurable real-valued functions on X , that is, we identify functions equal μ -almost everywhere on X . The support $S(f)$ of a measurable function f is defined by $S(f) := \{x \in X : f(x) \neq 0\}$.

For a *Young's function* Φ , let $\rho_\Phi : L^\Phi(\mu) \rightarrow \mathbb{R}^+$ such that $\rho_\Phi(f) = \int_X \Phi(f) d\mu$ for all $f \in L^\Phi(\mu)$. Then the space

$$L^\Phi(\mu) = \{f \in L^0(\mathcal{F}) : \exists k > 0, \rho_\Phi(kf) < \infty\}$$

is called an Orlicz space. Define the functional

$$N_\Phi(f) = \inf\{k > 0 : \rho_\Phi\left(\frac{f}{k}\right) \leq 1\}.$$

The functional $N_\Phi(\cdot)$ is a norm on $L^\Phi(\mu)$ and is called *gauge norm* (or Luxemburge norm). Also, $(L^\Phi(\mu), N_\Phi(\cdot))$ is a normed linear space. If almost every where equal functions are identified, then $(L^\Phi(\mu), N_\Phi(\cdot))$ is a Banach space, the basic measure space (X, \mathcal{F}, μ) is unrestricted. Hence every element of $L^\Phi(\mu)$ is a class of measurable functions that are almost every where equal. Also, there is another norm on $L^\Phi(\mu)$, defined as follow:

$$(1.1) \quad \|f\|_\Phi = \sup\left\{\int_X |fg| d\mu : g \in B_\Psi\right\} = \sup\left\{\left|\int_X fg d\mu\right| : g \in B_\Psi\right\},$$

in which $B_\Psi = \{g \in L^\Psi(\mu) : \int_X \Psi(|g|) d\mu \leq 1\}$. The norm $\|\cdot\|_\Phi$ is called *Orlicz norm*. For any $f \in L^\Phi(\mu)$, Φ being a Young function, we have

$$(1.2) \quad N_\Phi(f) \leq \|f\|_\Phi \leq 2N_\Phi(f).$$

For every $F \in \mathcal{F}$ with $0 < \mu(F) < \infty$, we have $N_\Phi(\chi_F) = \frac{1}{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}$. Moreover, if $\Phi \in \Delta_2$, then $(L^\Phi(\mu))^* = L^\Psi(\mu)$, in which Ψ is the complementary Young's function to Φ . Here we recall the next theorem about convergence of sequences in Orlicz spaces.

Theorem 1.1. [18] *Let $\{f_n\}_{n \geq 1}$ be a sequence from $L^\Phi(\mu)$ and $f \in L^\Phi(\mu)$. Then the following assertions hold:*

(a) If $\|f_n - f\|_\Phi \rightarrow 0$ (or equivalently $N_\Phi(f_n - f) \rightarrow 0$), then $\rho_\Phi(f_n) \rightarrow \rho_\Phi(f)$. The converse holds if Φ is Δ_2 -regular.

(b) If Φ is Δ_2 -regular Young function, or if Φ is continuous and concave $\Phi(0) = 0$, $\Phi \nearrow$ as well, $\rho_\Phi(f_n) \rightarrow \rho_\Phi(f)$ as $n \rightarrow \infty$ and $f_n \rightarrow f$ a.e., or in μ -measure, then $f_n \rightarrow f$ in norm.

Throughout this paper, (X, \mathcal{F}, μ) will be a measure space, that is, X is a set, \mathcal{F} is a sigma algebra on X and μ is a positive measure on \mathcal{F} . Also, we assume that $\varphi : X \rightarrow X$ is a non-singular measurable transformation, that is, $\varphi^{-1}(F) \in \mathcal{F}$, for every $F \in \mathcal{F}$ and $\mu(\varphi^{-1}(F)) = 0$, if $\mu(F) = 0$. Moreover, for the non-singular measurable transformation $\varphi : X \rightarrow X$, if there exists a positive constant c for which

$$\mu(\varphi^{-1}(F)) \leq c\mu(F), \quad \text{for every } F \in \mathcal{F},$$

then the linear operator

$$C_\varphi : L^\Phi(\mu) \rightarrow L^\Phi(\mu), \quad C_\varphi(f) = f \circ \varphi,$$

is well-defined and continuous on the Orlicz space $L^\Phi(\mu)$ and is called composition operator. For more details on composition operators on Orlicz spaces one can refer to [9].

2. Expansivity for composition operators

Here we recall some results of [7].

Proposition 2.1. *Let T be an operator on a Banach space X . Then:*

(a) *T is positively expansive if and only if $\sup_{n \in \mathbb{N}} \|T^n x\| = \infty$, for every nonzero $x \in X$.*

(b) *T is uniformly positively expansive if and only if $\lim_{n \rightarrow \infty} \|T^n x\| = \infty$, uniformly on S_X .*

If, in addition, T is invertible, then:

(c) *T is expansive if and only if $\sup_{n \in \mathbb{Z}} \|T^n x\| = \infty$, for every nonzero $x \in X$.*

(d) *T is uniformly expansive if and only if $S_X = A \cup B$, where $\lim_{n \rightarrow \infty} \|T^n x\| = \infty$, uniformly on A and $\lim_{n \rightarrow \infty} \|T^{-n} x\| = \infty$, uniformly on B .*

Let $C_\varphi : L^\Phi(\mu) \rightarrow L^\Phi(\mu)$ be composition operator and $\mathcal{F}^+ = \{F \in \mathcal{F} : 0 < \mu(F) < \infty\}$. Here we rewrite the definition of composition dynamical system, that is used in [10] and similar papers for composition operators on L^p -spaces, for composition operators on Orlicz spaces $L^\Phi(\mu)$, in which $\Phi \in \Delta_2$.

Definition 2.2. A composition dynamical system is a quintuple $(X, \mathcal{F}, \mu, \varphi, C_\varphi)$ where

- (1) (X, \mathcal{F}, μ) is a σ -finite measure space,
- (2) $\varphi : X \rightarrow X$ is an injective bimeasurable transformation, i.e., $\varphi(F) \in \mathcal{F}$ and $\varphi^{-1}(F) \in \mathcal{F}$ for every $F \in \mathcal{F}$,
- (3) there is $c > 0$ such that

$$(2.1) \quad \mu(\varphi^{-1}(A)) \leq c\mu(A) \quad \text{for every } A \in \mathcal{F},$$

- (4) $C_\varphi : L^\Phi(\mu) \rightarrow L^\Phi(\mu)$ is the composition operator induced by φ , that is

$$C_\varphi f = f \circ \varphi.$$

It is known that the condition 2.1 is a necessary and sufficient condition for boundedness of C_φ on Orlicz space $L^\Phi(\mu)$. In addition, if φ is bijective and φ^{-1} satisfies in condition 2.1 then $C_{\varphi^{-1}}$ is a bounded linear operators and $C_{\varphi^{-1}}^{-1} = C_\varphi$. For more details on composition operators one can refer to [19].

Also, by the fact that for every $F \in \mathcal{F}$ with $0 < \mu(F) < \infty$, we have $\|\chi_F\|_p = (\int_X |\chi_F|^p d\mu)^{\frac{1}{p}} = \mu(F)^{\frac{1}{p}}$ and $N_\Phi(\chi_F) = \frac{1}{\Phi^{-1}(\frac{1}{\mu(F)})}$, we can rewrite the definition of wandering set, dissipative system and dissipative system of bounded distortion in Orlicz spaces, that were defined for L_p spaces before in some contexts such as [10].

Definition 2.3. Let (X, \mathcal{F}, μ) be a measure space and $\varphi : X \rightarrow X$ be an invertible non-singular transformation. A measurable set $W \subseteq X$ is called a wandering set for φ if the sets $\{\varphi^{-n}(W)\}_{n \in \mathbb{Z}}$ are pair-wise disjoint.

Definition 2.4. Let (X, \mathcal{F}, μ) be a measure space and $\varphi : X \rightarrow X$ be an invertible non-singular transformation. The quadruple $(X, \mathcal{F}, \mu, \varphi)$ is called

- a dissipative system generated by W , if $X = \dot{\cup}_{k \in \mathbb{Z}} \varphi^k(W)$ for some $W \in \mathcal{F}$ with $0 < \mu(W) < \infty$ (the symbol $\dot{\cup}$ denotes pairwise disjoint union);
- a dissipative system, of bounded distortion, generated by W , if there exists $K > 0$, such that

$$(2.2) \quad \frac{1}{K} \frac{N_\Phi(C_\varphi^k(\chi_W))}{N_\Phi(\chi_W)} \leq \frac{N_\Phi(C_\varphi^k(\chi_F))}{N_\Phi(\chi_F)} \leq K \frac{N_\Phi(C_\varphi^k(\chi_W))}{N_\Phi(\chi_W)},$$

for all $k \in \mathbb{Z}$ and $F \in \mathcal{F}_W = \{F \cap W : F \in \mathcal{F}\}$. If we replace N_Φ by the norm of L_p , then we will have the bounded distortion property in L_p -spaces.(Definition 2.6.4, [10])

Definition 2.5. A composition dynamical system $(X, \mathcal{F}, \mu, \varphi, C_\varphi)$ is called

- dissipative composition dynamical system, generated by W , if $(X, \mathcal{F}, \mu, \varphi)$ is a dissipative system generated by W ;
- dissipative composition dynamical system, of bounded distortion, generated by W , if $(X, \mathcal{F}, \mu, \varphi)$ is a dissipative system of bounded distortion, generated by W .

In the following we recall Proposition 2.4 of [12].

Proposition 2.6. *Let $(X, \mathcal{F}, \mu, \varphi)$ be a dissipative system of bounded distortion, generated by W . Then there exists $H > 0$ such that*

$$(2.3) \quad \frac{1}{H} \frac{N_\Phi(C_\varphi^{t+s}(\chi_W))}{N_\Phi(C_\varphi^s(\chi_W))} \leq \frac{N_\Phi(C_\varphi^{t+s}(\chi_F))}{N_\Phi(C_\varphi^s(\chi_F))} \leq H \frac{N_\Phi(C_\varphi^{t+s}(\chi_W))}{N_\Phi(C_\varphi^s(\chi_W))},$$

for all $F \in \mathcal{F}_W$ with $\mu(F) > 0$ and all $s, t \in \mathbb{Z}$.

Now in the next Theorem we obtain equivalent conditions for composition operators to be expansive, positively expansive and uniformly positively expansive.

Theorem 2.7. *Let $(X, \mathcal{F}, \mu, \varphi, C_\varphi)$ be a composition dynamical system. The followings hold.*

- (1) The composition operator C_φ is positively expansive if and only if for every $A \in \mathcal{F}$, with $0 < \mu(A) < \infty$, we have

$$\inf_{n \in \mathbb{N}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) = 0.$$

- (2) The composition operator C_φ is expansive if and only if for every $A \in \mathcal{F}$, with $0 < \mu(A) < \infty$, we have

$$\inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) = 0.$$

- (3) If Φ is Δ_2 -regular, then the composition operator C_φ is uniformly positively expansive if and only if

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(A)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right)} = \infty,$$

for all $A \in \mathcal{F}$, with $0 < \mu(A) < \infty$.

- (4) If Φ is Δ_2 -regular, then the composition operator is uniformly expansive if and only if \mathcal{F}^+ can be splitted as $\mathcal{F}^+ = \mathcal{F}_A^+ \cup \mathcal{F}_B^+$ such that

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^n(F))}\right)} = \infty, \quad \text{uniformly on } \mathcal{F}_A^+,$$

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} = \infty, \quad \text{uniformly on } \mathcal{F}_B^+.$$

Proof. First we prove (2). Let C_φ be expansive. Then by part (c) of the Proposition 2.1 we have

$$\sup_{n \in \mathbb{Z}} N_\Phi(C_\varphi^n f) = \sup_{n \in \mathbb{Z}} N_\Phi(C_{\varphi^n} f) = \infty, \quad \text{for each } f \in L^\Phi(\mu) \setminus \{0\}.$$

Now if $F \in \mathcal{F}$ with $0 < \mu(F) < \infty$, then by setting $f = \chi_F$ and for every $n \in \mathbb{Z}$ we have

$$N_\Phi(C_\varphi^n f) = N_\Phi(\chi_{\varphi^{-n}(F)}) = \frac{1}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)}.$$

So we have

$$\sup_{n \in \mathbb{Z}} N_\Phi(C_\varphi^n f) = \sup_{n \in \mathbb{Z}} \frac{1}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} = \infty$$

and therefore

$$\inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right) = 0.$$

Now we prove the converse. Suppose that for each $F \in \mathcal{F}$, with $0 < \mu(F) < \infty$, we have $\inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right) = 0$. If $f \in L^\Phi(\mu) \setminus \{0\}$, then there exists a positive constant δ such that the set $F_\delta = \{x \in X : |f(x)| > \delta\}$ has positive measure, i.e,

$\mu(F_\delta) > 0$. Since (X, \mathcal{F}, μ) is σ -finite, then without loss of generality we can assume that $\mu(F_\delta) < \infty$. So for every $n \in \mathbb{Z}$:

$$\begin{aligned} \int_X \Phi\left(\frac{\delta \cdot \chi_{F_\delta} \circ \varphi^n}{N_\Phi(C_\varphi^n)}\right) d\mu &\leq \int_{\varphi^{-n}(F_\delta)} \Phi\left(\frac{f \circ \varphi^n}{N_\Phi(C_\varphi^n)}\right) d\mu \\ &\leq \int_X \Phi\left(\frac{f \circ \varphi^n}{N_\Phi(C_\varphi^n)}\right) d\mu \\ &\leq 1. \end{aligned}$$

Hence $N_\Phi(\delta \cdot \chi_{F_\delta}) \leq N_\Phi(C_\varphi^n f)$ and so

$$\delta \cdot \frac{1}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F_\delta))}\right)} \leq N_\Phi(C_\varphi^n f).$$

This inequality implies that if

$$\inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F_\delta))}\right) = 0,$$

then $\sup_{n \in \mathbb{Z}} N_\Phi(C_\varphi^n f) = \infty$. Since f is chosen arbitrarily, then by part (c) of the Proposition 2.1 we get that C_φ is expansive on $L^\Phi(\mu)$.

For the proof of part (1) it suffices just to replace \mathbb{Z} by \mathbb{N} .

(3). Let C_φ be uniformly positively expansive. Then by Proposition 2.1 we have

$$\lim_{n \rightarrow \infty} N_\Phi(C_\varphi^n f) = \infty, \quad \text{uniformly on } S_{L^\Phi(\mu)},$$

in which $S_{L^\Phi(\mu)} = \{f \in L^\Phi(\mu) : N_\Phi(f) = 1\}$. Let $f = \Phi^{-1}\left(\frac{1}{\mu(F)}\right)\chi_F$, for each $F \in \mathcal{F}^+$. Then we have $N_\Phi(f) = 1$ and also

$$N_\Phi(C_\varphi^n f) = \Phi^{-1}\left(\frac{1}{\mu(F)}\right)N_\Phi(\chi_{\varphi^{-n}(F)}) = \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)}.$$

So by our assumptions we get that

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} = \infty, \quad \text{uniformly on the sets } F \in \mathcal{F}^+.$$

Conversely, since $\Phi \in \Delta_2$, then the simple functions are dens in $L^\Phi(\mu)$. So it suffices to prove that for any simple function $f \in S_{L^\Phi(\mu)}$, the equality

$$\lim_{n \rightarrow \infty} N_\Phi(C_\varphi^n f) = \infty$$

holds. By our assumptions, for each $M > 0$ there exists $N \in \mathbb{N}$ such that for every $F \in \mathcal{F}^+$, we have

$$\frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} > M, \quad \text{for each } n \geq N.$$

By equation 1.1, it is easy to see that for $f, g \in L^\Phi(\mu)$, with $S(f) \cap S(g) \neq \emptyset$, there exists $c > 0$ such that

$$\|f + g\|_\Phi \geq c(\|f\|_\Phi + \|g\|_\Phi).$$

Now for every simple function $f \in S_{L^\Phi(\mu)}$, with $f = \sum_{i=1}^m \alpha_i \chi_{F_i}$, $\alpha_i \in \mathbb{C} \setminus \{0\}$ and $F_i \in \mathcal{F}$, without loss of generality we can assume that measurable sets $\{F_i\}_{i=1}^m$ are

pairwise disjoint. Hence by inequalities 1.2, for any $n \geq N$, we have

$$\begin{aligned}
N_{\Phi}(C_{\varphi}^n f) &= N_{\Phi}(\sum_{i=1}^m \alpha_i \chi_{F_i} \circ \varphi^n) \\
&\geq \frac{c}{2} \sum_{i=1}^m |\alpha_i| N_{\Phi} \chi_{\varphi^{-n}(F_i)} \\
&= \frac{c}{2} \sum_{i=1}^m |\alpha_i| \frac{1}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F_i))}\right)} \\
&\geq \frac{cM}{2} \sum_{i=1}^m |\alpha_i| \frac{1}{\Phi^{-1}\left(\frac{1}{\mu(F_i)}\right)} \\
&= \frac{cM}{2} \sum_{i=1}^m |\alpha_i| N_{\Phi}(\chi_{F_i}) \\
&\geq \frac{cM}{2} N_{\Phi}(f) \\
&= \frac{cM}{2}.
\end{aligned}$$

This implies that $\lim_{n \rightarrow \infty} N_{\Phi}(C_{\varphi}^n f) = \infty$. As is known, for every $f \in S_{L^{\Phi}(\mu)}$, we have a decomposition as $f = f^+ - f^-$, in which f^+, f^- are positive measurable functions. We can find a sequence of simple functions $\{f_k\}$ with $|f_k| \leq 2|f|$, such that converges point-wisely to f and so the sequence $\{\Phi(f_k)\}$ is also convergent point-wisely to $\Phi(f)$. Then by using Lebesgue Dominated Convergence Theorem we have

$$\lim_{k \rightarrow \infty} \rho_{\Phi}(f_k) = \rho_{\Phi}(f).$$

Hence by Theorem 1.1 we get that

$$\lim_{k \rightarrow \infty} N_{\Phi}(f_k) = N_{\Phi}(f) = 1.$$

Therefore by combining the above observations we get that $\lim_{n \rightarrow \infty} N_{\Phi}(C_{\varphi}^n f) = \infty$.

(4) Let C_{φ} be uniformly expansive. Then by part (d) of the Proposition 2.1 we have a decomposition as $S_{L^{\Phi}(\mu)} = A \cup B$, where

$$\lim_{n \rightarrow \infty} N_{\Phi}(C_{\varphi}^n f) = \infty, \quad \text{uniformly on } A$$

and

$$\lim_{n \rightarrow \infty} N_{\Phi}(C_{\varphi}^{-n} f) = \infty, \quad \text{uniformly on } B.$$

If we put $\mathcal{F}^+ = \{F \in \mathcal{F} : 0 < \mu(F) < \infty\}$, then some basic properties of Orlicz spaces $L^{\Phi}(\mu)$ and the fact that simple functions are dense in them give us: $\mathcal{F}^+ = \mathcal{F}_A^+ \cup \mathcal{F}_B^+$ in which

$$\mathcal{F}_A^+ = \{F \in \mathcal{F}^+ : \Phi^{-1}\left(\frac{1}{\mu(F)}\right) \chi_F \in A\} \quad \text{and} \quad \mathcal{F}_B^+ = \{F \in \mathcal{F}^+ : \Phi^{-1}\left(\frac{1}{\mu(F)}\right) \in B\}.$$

Since

$$\Phi^{-1}\left(\frac{1}{\mu(F)}\right) \chi_F \in S_{L^{\Phi}(\mu)} \quad \text{and} \quad C_{\varphi}^n(\Phi^{-1}\left(\frac{1}{\mu(F)}\right) \chi_F) = \Phi^{-1}\left(\frac{1}{\mu(F)}\right) \chi_{\varphi^{-n}(F)},$$

then by our assumptions we have

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^n(F))}\right)} = \infty, \quad \text{on } \mathcal{F}_A^+$$

and

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} = \infty, \quad \text{on } \mathcal{F}_B^+.$$

So we get the result. Now to prove the converse, we use again part (d) of the Proposition 2.1. Hence it suffices to show that $S_{L^\Phi(\mu)} = A \cup B$, such that

$$\lim_{n \rightarrow \infty} N_\Phi(C_\varphi^n f) = \infty, \quad \text{uniformly on } A$$

and

$$\lim_{n \rightarrow \infty} N_\Phi(C_\varphi^{-n} f) = \infty, \quad \text{uniformly on } B.$$

By the assumptions, for every $M > 0$, there exists $N \in \mathbb{N}$ such that for every $n \geq N$,

$$\frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^n(F))}\right)} > M \quad \text{for each } F \in \mathcal{F}_A^+$$

and

$$\frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} > M \quad \text{for each } F \in \mathcal{F}_B^+.$$

Now, let $f \in S_{L^\Phi(\mu)}$ be a simple function with $f = \sum_{i=1}^m \alpha_i \chi_{F_i}$, in which F_i 's are pairwise disjoint measurable sets with positive measure. Since $\mathcal{F}^+ = \mathcal{F}_A^+ \cup \mathcal{F}_B^+$, then we can define

$$f_{\mathcal{F}_A^+} = \sum_{\{1 \leq i \leq m: F_i \in \mathcal{F}_A^+\}} \alpha_i \chi_{F_i}$$

and

$$f_{\mathcal{F}_B^+} = \sum_{\{1 \leq i \leq m: F_i \in \mathcal{F}_B^+\}} \alpha_i \chi_{F_i}.$$

It is clear that $f = f_{\mathcal{F}_A^+} + f_{\mathcal{F}_B^+}$. So by equation 1.1 there exists $c > 0$ such that

$$\|f\|_\Phi \geq c(\|f_{\mathcal{F}_A^+}\|_\Phi + \|f_{\mathcal{F}_B^+}\|_\Phi) \geq c\|f\|_\Phi$$

and since $N_\Phi(f) = 1$, then by inequalities 1.2 we have

$$2 \geq c(\|f_{\mathcal{F}_A^+}\|_\Phi + \|f_{\mathcal{F}_B^+}\|_\Phi) \geq 2c.$$

This means that

$$\|f_{\mathcal{F}_A^+}\|_\Phi \geq 1 \quad \text{or} \quad \|f_{\mathcal{F}_B^+}\|_\Phi \geq 1$$

and so by inequalities 1.2 we have

$$N_\Phi(f_{\mathcal{F}_A^+}) \geq \frac{1}{2} \quad \text{or} \quad N_\Phi(f_{\mathcal{F}_B^+}) \geq \frac{1}{2}.$$

Hence if $N_\Phi(f_{\mathcal{F}_A^+}) \geq \frac{1}{2}$, then for every $n \geq N$

$$\begin{aligned}
N_\Phi(C_\varphi^{-n}f) &\geq \frac{1}{2}\|C_\varphi^{-n}f\|_\Phi \\
&= \frac{1}{2}\|C_\varphi^{-n}(f_{\mathcal{F}_A^+} + f_{\mathcal{F}_B^+})\|_\Phi \\
&\geq \frac{c}{2}(\|C_\varphi^{-n}(f_{\mathcal{F}_A^+}\|_\Phi + \|f_{\mathcal{F}_B^+}\|_\Phi) \\
&\geq \frac{c}{2}\|C_\varphi^{-n}f_{\mathcal{F}_A^+}\|_\Phi \\
&\geq \frac{c^2}{2} \sum_{\{1 \leq i \leq m: F_i \in \mathcal{F}_A^+\}} |\alpha_i| \frac{1}{\Phi^{-1}(\frac{1}{\mu(\varphi^n(F_i))})} \\
&\geq \frac{c^2 M}{2} \sum_{\{1 \leq i \leq m: F_i \in \mathcal{F}_A^+\}} |\alpha_i| \frac{1}{\Phi^{-1}(\frac{1}{\mu(F_i)})} \\
&\geq \frac{c^2 M}{2} N_\Phi(f_{\mathcal{F}_A^+}) \\
&\geq \frac{c^2 M}{4}.
\end{aligned}$$

By a similar way we get that if $N_\Phi(f_{\mathcal{F}_B^+}) \geq \frac{1}{2}$, then we have $N_\Phi(C_\varphi^n f) \geq \frac{c^2 M}{4}$. If we set

$$S_{L^\Phi(\mu)}^0 = \{f \in S_{L^\Phi(\mu)} : f \text{ is a simple function}\},$$

then from the above observations we get that $S_{L^\Phi(\mu)}^0 = A_0 \cup B_0$, in which

$$A_0 = \{f \in S_{L^\Phi(\mu)}^0 : N_\Phi(f_{\mathcal{F}_A^+}) \geq \frac{1}{2}\} \quad \text{and} \quad B_0 = \{f \in S_{L^\Phi(\mu)}^0 : N_\Phi(f_{\mathcal{F}_B^+}) \geq \frac{1}{2}\}.$$

Let $f \in S_{L^\Phi(\mu)}$ be an arbitrary element. Then we can write $f = f^+ - f^-$ in which f^+ and f^- are positive and negative parts of f , respectively. Hence we can find two non-decreasing sequences of simple functions $\{f_k^+\}_{k \in \mathbb{N}}$ and $\{f_k^+ - f_k^-\}_{k \in \mathbb{N}}$ such that converge pointwisely to f^+ and f^- , respectively. If we define $f_k = f_k^+ - f_k^-$, then we get that the sequence $\{f_k\}_{k \in \mathbb{N}}$ converges pointwisely to f . Since $|f_k| = |f_k^+ - f_k^-| \leq 2|f|$ and so $\Phi(|f_k|) \leq \Phi(2|f|)$, then by Lebesgue Dominated Convergence Theorem we have

$$\lim_{k \rightarrow \infty} \rho_\Phi(f_k) = \rho_\Phi(f).$$

Hence by Theorem 1.1 we get that

$$\lim_{k \rightarrow \infty} N_\Phi(f_k) = N_\Phi(f) = 1.$$

So there exists k_0 such that for every $k \geq k_0$, $N_\Phi(f_k) > \frac{1}{2}$. By the above observations we get that

$$N_\Phi(C_\varphi^n(\frac{f_k}{N_\Phi(f_k)}) < 4N_\Phi(C_\varphi^n(f))$$

and also at least one of the following sets must be infinite:

$$I_1(f) = \{k \in \mathbb{N} : \frac{f_k}{N_\Phi(f_k)} \in A_0\}; \quad I_2(f) = \{k \in \mathbb{N} : \frac{f_k}{N_\Phi(f_k)} \in B_0\}.$$

Therefore, if $I_1(f)$ is infinite, then there is a subsequence

$$\left\{ \frac{f_{k_j}}{N_\Phi(f_{k_j})} \right\}_{j \in \mathbb{N}} \subseteq A_0$$

such that for every $M > 0$, there exists $N \in \mathbb{N}$ such that for each $n \geq N$,

$$N_\Phi(C_\varphi^{-n}(\frac{f_{k_j}}{N_\Phi(f_{k_j})})) > \frac{M}{2}.$$

Thus we have

$$N_\Phi(C_\varphi^{-n}(f)) > \frac{M}{8}, \quad \text{for every } n \geq N.$$

Similarly, if $I_2(f) = \infty$, then for each $M > 0$ we can find $N \in \mathbb{N}$, such that for every $n \geq N$, $N_\Phi(C_\varphi^n(f)) > \frac{M}{8}$. Finally if we set

$$A = \{f \in S_{L^\Phi(\mu)} : I_2(f) = \infty\} \quad \text{and} \quad B = \{f \in S_{L^\Phi(\mu)} : I_1(f) = \infty\},$$

then we have $S_{L^\Phi(\mu)} = A \cup B$, such that

$$\lim_{n \rightarrow \infty} N_\Phi(C_\varphi^n f) = \infty, \quad \text{uniformly on } A$$

and

$$\lim_{n \rightarrow \infty} N_\Phi(C_\varphi^{-n} f) = \infty, \quad \text{uniformly on } B.$$

This completes the proof. \square

Theorem 2.8. *Let $(X, \mathcal{F}, \mu, \varphi, C_\varphi)$ be a dissipative composition dynamical system of bounded distortion, generated by W . Then the following statements hold.*

- (1) C_φ is positively expansive if and only if $\inf_{n \in \mathbb{N}} \Phi^{-1}(\frac{1}{\mu(\varphi^{-n}(W))}) = 0$.
- (2) C_φ is expansive if and only if $\inf_{n \in \mathbb{Z}} \Phi^{-1}(\frac{1}{\mu(\varphi^n(W))}) = 0$.
- (3) C_φ is uniformly positively expansive if and only if

$$\lim_{n \rightarrow \infty} \Phi^{-1}(\frac{1}{\mu(\varphi^{-n}(W))}) = 0.$$

- (4) C_φ is uniformly expansive if and only if one of the following conditions hold:

$$(2.4) \quad \lim_{n \rightarrow \infty} \inf_{k \in \mathbb{Z}} \left(\frac{\Phi^{-1}(\frac{1}{\mu(\varphi^k(W))})}{\Phi^{-1}(\frac{1}{\mu(\varphi^{k+n}(W))})} \right) = \infty,$$

$$(2.5) \quad \lim_{n \rightarrow \infty} \inf_{k \in \mathbb{Z}} \left(\frac{\Phi^{-1}(\frac{1}{\mu(\varphi^k(W))})}{\Phi^{-1}(\frac{1}{\mu(\varphi^{k-n}(W))})} \right) = \infty,$$

$$(2.6) \quad \lim_{n \rightarrow \infty} \inf_{k \in \mathbb{Z}} \left(\frac{\Phi^{-1}(\frac{1}{\mu(\varphi^k(W))})}{\Phi^{-1}(\frac{1}{\mu(\varphi^{k+n}(W))})} \right) = \infty \quad \text{and} \quad \lim_{n \rightarrow \infty} \inf_{k \in -\mathbb{N}_0} \left(\frac{\Phi^{-1}(\frac{1}{\mu(\varphi^k(W))})}{\Phi^{-1}(\frac{1}{\mu(\varphi^{k-n}(W))})} \right) = \infty.$$

Proof. First we prove part (2), then for the proof of part (1) it suffices just to replace \mathbb{Z} by \mathbb{N} .

In Theorem 2.7 we can replace \mathcal{F} by \mathcal{F}^+ , i.e., we can write "the composition operator C_φ is expansive if and only if for every $A \in \mathcal{F}^+$,

$$\inf_{n \in \mathbb{Z}} \Phi^{-1}(\frac{1}{\mu(\varphi^{-n}(A))}) = 0.$$

Hence for the proof of (2), it suffices to prove

$$\inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) = 0, \quad \forall A \in \mathcal{F}^+ \Leftrightarrow \inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^n(W))}\right) = 0.$$

The implication

$$\inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) = 0, \quad \forall A \in \mathcal{F}^+ \Rightarrow \inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^n(W))}\right) = 0$$

is clear. So we prove the converse. Let $F \in \mathcal{F}^+$. Since $X = \cup_{n \in \mathbb{Z}} \varphi^n(W)$, in which $\varphi^n(W)$'s are pair-wise disjoint and $\mu(F) > 0$, then there exists $n_0 \in \mathbb{Z}$ such that $\mu(F \cap \varphi^{n_0}(W)) > 0$. On the other hand $\varphi^{-n_0}(F \cap \varphi^{n_0}(W)) = \varphi^{n_0}(F) \cap W$. Hence if we take $A = \varphi^{-n_0}(F \cap \varphi^{n_0}(W))$, then $A \in \mathcal{F}_W$ and $\mu(A) \leq \mu(W)$. Now by using bounded distortion property for we have

$$\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) \leq \frac{KN_\Phi(\chi_W)}{N_\Phi(\chi_A)} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right), \quad \forall n \in \mathbb{Z}$$

and therefore

$$\inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) \leq \frac{KN_\Phi(\chi_W)}{N_\Phi(\chi_A)} \inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right) = 0.$$

Consequently we have

$$\inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right) \leq \inf_{n \in \mathbb{Z}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) = 0.$$

(3). By Theorem 2.7 it suffices to prove that

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} = \infty, \quad \forall F \in \mathcal{F}^+ \Leftrightarrow \lim_{n \rightarrow \infty} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right) = 0.$$

It is obvious that if

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} = \infty, \quad \forall F \in \mathcal{F}^+,$$

then $\lim_{n \rightarrow \infty} \frac{1}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right)} = \infty$ and so $\lim_{n \rightarrow \infty} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right) = 0$. Now we prove the converse, suppose that $\lim_{n \rightarrow \infty} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right) = 0$ and $F \in \mathcal{F}^+$. Similar to pat (2) we can find $n_0 \in \mathbb{Z}$ such that $0 < \mu(F \cap \varphi^{n_0}(W)) < \infty$ and

$$A = \varphi^{-n_0}(F \cap \varphi^{n_0}(W)) = \varphi^{-n_0}(F) \cap W.$$

Hence $0 < \mu(A) < \infty$ and $A \in \mathcal{F}_W$. Now by using bounded distortion property we have

$$\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) \leq \frac{KN_\Phi(\chi_W)}{N_\Phi(\chi_A)} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right), \quad \forall n \in \mathbb{Z}$$

and so

$$\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n-n_0}(F))}\right) \leq \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(A))}\right) \leq \frac{KN_\Phi(\chi_W)}{N_\Phi(\chi_A)} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right)$$

Thus by taking lim we have

$$\lim_{n \rightarrow \infty} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n-n_0}(F))}\right) \leq \frac{KN_\Phi(\chi_W)}{N_\Phi(\chi_A)} \lim_{n \rightarrow \infty} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right) = 0$$

hence $\lim_{n \rightarrow \infty} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n-n_0}(F))}\right) = 0$ and consequently

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} = \infty.$$

By Theorem 2.7, part (3) we get the proof.

(4) Suppose that C_φ is uniformly expansive. By Theorem 2.7 we have that $\mathcal{F}^+ = \mathcal{F}_A^+ \cup \mathcal{F}_B^+$ such that

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^n(F))}\right)} = \infty, \quad \text{uniformly on } \mathcal{F}_A^+,$$

$$\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} = \infty, \quad \text{uniformly on } \mathcal{F}_B^+.$$

If we set

$$I = \{k \in \mathbb{Z} : \varphi^k(W) \in \mathcal{F}_A\} \quad \text{and} \quad J = \{k \in \mathbb{Z} : \varphi^k(W) \in \mathcal{F}_B\},$$

then by our assumptions we get that

$$\lim_{n \rightarrow \infty} \inf_{k \in I} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)} \right) = \infty$$

$$\lim_{n \rightarrow \infty} \inf_{k \in J} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right) = \infty.$$

By these observations, if $J = \emptyset$, then the condition 2.4 holds, if $I = \emptyset$, then the condition 2.5 holds and if I and J are both non-empty, then there exist $i, j \in \mathbb{Z}$ such that $[i, \infty) \cap \mathbb{Z} \subseteq I$ and $(-\infty, j] \cap \mathbb{Z} \subseteq J$, this implies that the condition 2.6 holds.

For the converse, for every $F \in \mathcal{F}^+$, by our assumptions we have $F = \cup_{k \in \mathbb{Z}} F_k$, where $F_k = F \cap \varphi^k(W)$ and F_k 's are pair-wise disjoint and also by Proposition 2.6, for every $n, k \in \mathbb{Z}$ there exists $H > 0$ such that

$$\frac{1}{H} \frac{N_\Phi(C_\varphi^{k+n}(\chi_W))}{N_\Phi(C_\varphi^k(\chi_W))} N_\Phi(\chi_{F_k}) \leq N_\Phi(C_\varphi^n(\chi_{F_k})) \leq H \frac{N_\Phi(C_\varphi^{k+n}(\chi_W))}{N_\Phi(C_\varphi^k(\chi_W))} N_\Phi(\chi_{F_k}).$$

Also, we have $0 < \mu(F) < \infty$. Hence there exists $k \in \mathbb{Z}$ such that $\mu(F_k) > 0$. Therefore, for each $n \in \mathbb{Z}$,

$$\begin{aligned}
N_{\Phi}(C_{\varphi}^n(\chi_F)) &\geq \frac{1}{2} \|C_{\varphi}^n(\chi_F)\|_{\Phi} \\
&= \frac{1}{2} \|\chi_{\varphi^{-n}(F)}\|_{\Phi} \\
&\geq \frac{c}{2} \sum_{k \in \mathbb{Z}} \|\chi_{\varphi^{-n}(F_k)}\|_{\Phi} \\
&\geq \frac{c}{2} \sum_{k \in \mathbb{Z}} N_{\Phi}(\chi_{\varphi^{-n}(F_k)}) \\
&= \frac{c}{2} \sum_{k \in \mathbb{Z}} N_{\Phi}(C_{\varphi}^n(\chi_{F_k})) \\
&\geq \frac{c}{2H} \sum_{k \in \mathbb{Z}} \frac{N_{\Phi}(C_{\varphi}^{k+n}(\chi_W))}{N_{\Phi}(C_{\varphi}^k(\chi_W))} N_{\Phi}(\chi_{F_k}) \\
&\geq \frac{c}{2H} \inf_{k \in \mathbb{Z}} \frac{N_{\Phi}(C_{\varphi}^{k+n}(\chi_W))}{N_{\Phi}(C_{\varphi}^k(\chi_W))} \sum_{k \in \mathbb{Z}} N_{\Phi}(\chi_{F_k}) \\
&\geq \frac{c}{2H} \inf_{k \in \mathbb{Z}} \frac{N_{\Phi}(C_{\varphi}^{k+n}(\chi_W))}{N_{\Phi}(C_{\varphi}^k(\chi_W))} N_{\Phi}(\chi_F).
\end{aligned}$$

By these observations we get that

$$\frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^n(F))}\right)} \geq \frac{c}{2H} \inf_{k \in \mathbb{Z}} \frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}.$$

This inequality implies that if we set

$$\mathcal{F}_A^+ = \{F \cap (\cup_{k \geq 0} \varphi^k(W)) : F \in \mathcal{F}^+\}$$

and

$$\mathcal{F}_B^+ = \{F \cap (\cup_{k < 0} \varphi^k(W)) : F \in \mathcal{F}^+\},$$

then $\mathcal{F}^+ = \mathcal{F}_A^+ \cup \mathcal{F}_B^+$ and also if one of conditions 2.4, 2.5 or 2.6 holds, then we get that

$$\begin{aligned}
\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^n(F))}\right)} &= \infty, \quad \text{uniformly on } \mathcal{F}_A^+, \\
\lim_{n \rightarrow \infty} \frac{\Phi^{-1}\left(\frac{1}{\mu(F)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(F))}\right)} &= \infty, \quad \text{uniformly on } \mathcal{F}_B^+.
\end{aligned}$$

Thus by Theorem 2.7 the composition operator C_{φ} is uniformly expansive. \square

In the next Proposition we obtain some sufficient conditions for composition operator C_{φ} on Orlicz spaces to decline structural stability and even strongly structurally stability.

Proposition 2.9. *Let $(X, \mathcal{F}, \mu, \varphi, C_{\varphi})$ be a dissipative composition dynamical system of bounded distortion, generated by W and $\Phi \in \Delta'$. If*

$$\overline{\lim}_{n \rightarrow \infty} \sup_{k \in \mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} < 1 \quad \text{and} \quad \underline{\lim}_{n \rightarrow \infty} \inf_{k \in -\mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right)^{\frac{1}{n}} > 1,$$

then the composition operator C_φ can not be structurally stable and so can not be even strongly structurally stable.

Proof. Since

$$\liminf_{n \rightarrow \infty} \inf_{k \in -\mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right)^{\frac{1}{n}} > 1,$$

then there exists $n_0 \in \mathbb{N}_0$ and $t > 1$ such that for every $n \geq n_0$,

$$\inf_{k \in -\mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right)^{\frac{1}{n}} \geq t > 1.$$

Hence

$$\frac{\Phi^{-1}\left(\frac{1}{\mu(W)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right)} \geq \inf_{k \in -\mathbb{N}_0} \frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \geq t^n.$$

This implies that $\sup_{n \in \mathbb{N}} \frac{\Phi^{-1}\left(\frac{1}{\mu(W)}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right)} = \infty$ and therefore

$$\inf_{n \in \mathbb{N}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right) = 0.$$

Thus by theorem 2.7, we get that C_φ is positively expansive. Moreover, for every $n \geq n_0$,

$$\sup_{k \in -\mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right)^{\frac{1}{n}} \geq \inf_{k \in -\mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right)^{\frac{1}{n}} \geq t > 1.$$

Hence we get that

$$\overline{\lim}_{n \rightarrow \infty} \sup_{k \in \mathbb{Z}} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right)^{\frac{1}{n}} \geq \overline{\lim}_{n \rightarrow \infty} \sup_{k \in -\mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right)^{\frac{1}{n}} > 1.$$

In addition, since

$$\overline{\lim}_{n \rightarrow \infty} \sup_{k \in \mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} < 1,$$

then there exists $m_0 \in \mathbb{N}_0$ and $0 < t < 1$ such that for every $n \geq m_0$,

$$\sup_{k \in \mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} \leq t < 1.$$

This implies that

$$\inf_{k \in \mathbb{Z}} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} \leq \inf_{k \in \mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} \leq t < 1$$

and consequently

$$\underline{\lim}_{n \rightarrow \infty} \inf_{k \in \mathbb{Z}} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} \leq \underline{\lim}_{n \rightarrow \infty} \inf_{k \in \mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} < 1.$$

By the above observations we get that none of conditions 2.3, 2.4 and 2.5 of the Corollary 2.20 of [12] holds. This means that C_φ doesn't have the shadowing

property and so it is not hyperbolic. Therefore C_φ is positively expansive but not hyperbolic, hence by Theorem 6 of [5] we get that C_φ is not structurally stable and therefore, not even strongly structurally stable. \square

Theorem 2.10. *Let $(X, \mathcal{F}, \mu, \varphi, C_\varphi)$ be a dissipative composition dynamical system of bounded distortion, generated by W and $\Phi \in \Delta'$. If at least one of the following inequalities holds:*

$$(2.7) \quad \overline{\lim}_{n \rightarrow \infty} \sup_{k \in \mathbb{Z}} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} < 1$$

$$(2.8) \quad \underline{\lim}_{n \rightarrow \infty} \inf_{k \in \mathbb{Z}} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)} \right)^{\frac{1}{n}} > 1$$

$$(2.9) \quad \overline{\lim}_{n \rightarrow \infty} \sup_{k \in -\mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k-n}(W))}\right)} \right)^{\frac{1}{n}} < 1 \text{ and } \underline{\lim}_{n \rightarrow \infty} \inf_{k \in \mathbb{N}_0} \left(\frac{\Phi^{-1}\left(\frac{1}{\mu(\varphi^k(W))}\right)}{\Phi^{-1}\left(\frac{1}{\mu(\varphi^{k+n}(W))}\right)} \right)^{\frac{1}{n}} > 1,$$

then the composition operator C_φ is strongly structurally stable.

Proof. If one of the conditions 2.7, 2.8 and 2.9 holds, then Theorem 2.16 of [12] and Theorem 1 of [3] we get that C_φ is strongly structurally stable. \square

Now by using Theorem 2.10 and also Corollary CSC of [12] we have the next corollary.

Corollary 2.11. *Let $(X, \mathcal{F}, \mu, \varphi, C_\varphi)$ be a dissipative composition dynamical system of bounded distortion, generated by W and $\Phi \in \Delta'$. Then the composition operator C_φ is strongly structurally stable when it has shadowing property or equivalently is generalized hyperbolic.*

Theorem 2.12. *Let $(X, \mathcal{F}, \mu, \varphi, C_\varphi)$ be a dissipative composition dynamical system of bounded distortion, generated by W , $\inf_{n \in \mathbb{N}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right) = 0$ and $\Phi \in \Delta'$. Then the composition operator C_φ is strongly structurally stable if and only if it has the shadowing property.*

Proof. If C_φ has the shadowing property, then by Corollary 2.11 we get that C_φ is strongly structurally stable. Conversely, if C_φ is strongly structurally stable, then by the assumption $\inf_{n \in \mathbb{N}} \Phi^{-1}\left(\frac{1}{\mu(\varphi^{-n}(W))}\right) = 0$ and by Theorem 2.8 we have C_φ is positively expansive. Therefore by Theorem 6 of [5] we get the proof. \square

Declarations

Conflict of interest. The authors have not disclosed any competing interests.

Acknowledgement. My manuscript has no associate data.

REFERENCES

- [1] N.C. Bernardes Jr., A. Bonilla, V. Müller, A. Peris, Distributional chaos for linear operators, *J. Funct. Anal.* 265 (9) (2013) 2143–2163.
- [2] N.C. Bernardes Jr., A. Bonilla, V. Müller, A. Peris, Li–Yorke chaos in linear dynamics, *Ergodic Theory Dynam. Systems* 35 (6) (2015) 1723–1745.
- [3] N. C. Bernardes and A. Messaoudi, A generalized Grobman–Hartman theorem, *Proc. Amer. Math. Soc.* 148 (2020), 4351–4360.

- [4] F. Bayart, É. Matheron, Dynamics of Linear Operators, Cambridge University Press, Cambridge, 2009.
- [5] N. BERNARDES, and A. MESSAOUDI, Shadowing and structural stability for operators. Ergodic Theory and Dynamical Systems, 41 (2021), 961-980.
- [6] J. Bès, Q. Menet, A. Peris, Y. Puig, Recurrence properties of hypercyclic operators, Math. Ann. 366 (1) (2016) 545–572.
- [7] N. C. Bernardes Jr, P. R. Cirilo, U. B. Darji, A. Messaoudi and E. R. Pujals, Expansivity and shadowing in linear dynamics, J. Math. Anal. Appl. 461 (2018) 796–816.
- [8] F. Bayart, I.Z. Ruzsa, Difference sets and frequently hypercyclic weighted shifts, Ergodic Theory Dynam. Systems 35 (3) (2015) 691–709.
- [9] Y. Cui, H. Hudzik, R. Kumar and L. Maligranda, Composition operators in Orlicz spaces, J. Aust. Math. Soc. 76 (2004), 189-206.
- [10] E. D’Aniello, U. B. Darji and M. Maiuriello, Generalized hyperbolicity and shadowing in L^p spaces. J. Difer. Equ. 298 (2021), 68–94.
- [11] M. Eisenberg, J.H. Hedlund, Expansive automorphisms of Banach spaces, Pacific J. Math. 34 (3) (1970) 647–656.
- [12] Y. Estaremi and N. Bamerni, Hyperbolicity and shadowing property in Orlicz spaces, arXiv.org.
- [13] K.-G. Grosse-Erdmann, A. Peris Manguillot, Linear Chaos, Springer-Verlag, London, 2011.
- [14] S. Grivaux, É. Matheron, Invariant measures for frequently hypercyclic operators, Adv. Math. 265 (2014) 371–427.
- [15] M. A. Krasnosel’skii, Ya. B. Rutickii, Convex functions and Orlicz spaces, Noordhoff, Netherlands, 1961.
- [16] M. Maiuriello, Expansivity and strong structural stability for composition operators on L^p spaces, Banach J. Math. Anal. (2022). <https://doi.org/10.1007/s43037-022-00196-4>
- [17] C. Nilson Jr. Bernardes, R. C. Patricia, B. Udayan B. D. A. Messaoudi, R. P. Enrique, Expansivity and shadowing in linear dynamics, J. Math. Anal. Appl. 461 (2018) 796–816.
- [18] M. M. Rao, Z.D. Ren, Theory of Orlicz spaces, Marcel Dekker, New York, 1991.
- [19] R.K. Singh, J.S. Manhas, Composition Operators on Function Spaces, North-Holland Mathematics Studies, vol. 179, North-Holland Publishing Co., Amsterdam, 1993.

Y. ESTAREMI

Email address: y.estaremi@gu.ac.ir

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES, GOLESTAN UNIVERSITY, GORGAN, IRAN.