

Complement Avoidance in Binary Words

James Currie¹ Pascal Ochem² Narad Rampersad³ Jeffrey Shallit⁴

September 21, 2022

Abstract

The complement \bar{x} of a binary word x is obtained by changing each 0 in x to 1 and vice versa. We study infinite binary words \mathbf{w} that avoid sufficiently large complementary factors; that is, if x is a factor of \mathbf{w} then \bar{x} is not a factor of \mathbf{w} . In particular, we classify such words according to their critical exponents.

1 Introduction

Let x be a finite nonempty binary word. We write \bar{x} for the complementary word, image of the morphism that maps $0 \rightarrow 1$ and $1 \rightarrow 0$, and we write x^R for the reversal (mirror image) of x . We say y is a *factor* of a (one-sided) infinite word \mathbf{w} if $\mathbf{w} = xyz$ for a finite word x and an infinite word \mathbf{z} . In this paper, we are interested in the construction of and properties of infinite binary words \mathbf{w} *avoiding* complementary factors: that is, if x is a nonempty factor of \mathbf{w} , then \bar{x} is not. This is not a new notion; for example, complement avoidance in De Bruijn words was studied by Sawada et al. [18].

Evidently it is impossible for an infinite word to avoid complementary factors of *all* lengths, except in the trivial cases $0^\omega = 000\dots$ and $1^\omega = 111\dots$. A natural question then poses itself: are there such infinite words if the set of exceptions is restricted in some way, say by length or by cardinality? And what is the repetition threshold of such infinite words? We now turn to repetitions.

We say that a finite word $w = w[1..n]$ has *period* $p \geq 1$ if $w[i] = w[i+p]$ for $1 \leq i \leq n-p$. The smallest period of a word w is called *the* period, and we write it as $\text{per}(w)$. The *exponent* of a finite word w , written $\text{exp}(w)$ is defined to be $|w|/\text{per}(w)$. For a real number α , we say a word (finite or infinite) is α -*free* if the exponent of all its nonempty factors is $> \alpha$.

¹Department of Math/Stats, University of Winnipeg, 515 Portage Ave., Winnipeg, MB, R3B 2E9, Canada; e-mail j.currie@uwinnipeg.ca.

²LIRMM, CNRS, Université de Montpellier, France; e-mail ochem@lirmm.fr.

³Department of Math/Stats, University of Winnipeg, 515 Portage Ave., Winnipeg, MB, R3B 2E9, Canada; e-mail n.rampersad@uwinnipeg.ca.

⁴School of Computer Science, University of Waterloo, Waterloo, ON N2L 3G1, Canada; e-mail shallit@uwaterloo.ca.

We say a word is α^+ -free if the exponent of all its nonempty factors is $\geq \alpha$. The *critical exponent* of a finite or infinite word x is the supremum, over all nonempty finite factors w of x , of $\exp(w)$; it is written $\text{ce}(x)$. The *repetition threshold* for a language L of infinite words is defined to be the infimum, over all $x \in L$, of $\text{ce}(x)$.

The repetition thresholds for various classes of words have been studied extensively. To name just a few classes, Dejean [8] determined the repetition threshold for all words over a 3-letter alphabet, and conjectured its value for larger alphabets. Her conjecture attracted a lot of attention, and was finally resolved by Rao [16] and Currie and Rampersad [7], independently.

Other classes that have been studied include the Sturmian words, studied by Carpi and de Luca [4, Prop. 15]; the palindromes, studied in [20]; the rich words, studied by Currie et al. [5]; the balanced words, studied by Rampersad et al. [15] and Dvořáková et al. [10]; and complementary symmetric Rote words, studied by Dvořáková et al. [9]. Other related works include [12, 1, 11, 17, 14].

In this paper we study the repetition threshold for two classes of infinite words:

- CAL_ℓ , the binary words for which there is no length- ℓ word x such that both x and \bar{x} appear as factors;
- CAN_n , the binary words for which there are at most n distinct words x such that both x and \bar{x} appear as factors.

It turns out that there is an interesting and subtle hierarchy, depending on the values of ℓ and n .

Our work is very similar in flavor to that of [19], which found a similar hierarchy concerning critical exponents and sizes of squares avoided. The hierarchy for complementary factors, as we will see, however, is significantly more complex. In this paper, in Section 2, we investigate the hierarchy based on word length, and in Section 3 we investigate the hierarchy based on the cardinality of the exceptional set. In Section 4 we study finite words avoiding complementary factors and determine under what conditions there are exponentially many such words.

We will need the following famous infinite words.

- The Fibonacci word $\mathbf{f} = 01001010010010100100100 \dots$, fixed point of the morphism $0 \rightarrow 01, 1 \rightarrow 0$. See, for example, [3].
- The word $\mathbf{p} = 0121021010210121010210121 \dots$, fixed point of the morphism $0 \rightarrow 01, 1 \rightarrow 21, 2 \rightarrow 0$. This is sequence [A287072](#) in the OEIS. Its properties were recently studied in [6].

2 The class CAL_ℓ

In this section, we investigate the repetition threshold for the class CAL_ℓ , the binary words \mathbf{w} with the property that if x is a length- ℓ factor of \mathbf{w} , then \bar{x} is not.

Lemma 1. *Suppose \mathbf{x} is an infinite binary word with critical exponent $\leq \frac{7}{3}$. Then \mathbf{x} contains infinitely many (and hence, arbitrarily large) complementary factors.*

Proof. By a result of Karhumäki and Shallit [13], every infinite binary word avoiding e -powers for $e \leq \frac{7}{3}$ contains $\mu^n(0)$ as a factor for all $n \geq 1$. Such a word is of the form $\mu^{n-1}(01) = \mu^{n-1}(0)\mu^{n-1}(1)$, and these two terms are complementary factors of length 2^{n-1} . \square

By Lemma 1, the lower limit on the repetition threshold is $\frac{7}{3}$. For $\ell = 1$ the only such words are of the form $0^\omega, 1^\omega$. For $\ell = 2$ the only such words are of the form $(\epsilon + 0)1^\omega$ and $(\epsilon + 1)0^\omega$. For larger ℓ the situation is as described in the following theorem.

Theorem 2. *There exists an infinite β^+ -free binary word containing no complementary factors of length $\geq \ell$, for the following pairs (ℓ, β) :*

(a) $(3, 2 + \alpha)$

(b) $(5, 3)$

(c) $(7, \frac{8}{3})$

(d) $(8, \frac{5}{2})$

(e) $(11, \gamma')$, where γ' is the critical exponent of the morphic word $\psi(\mathbf{p})$, where ψ is the morphism $0 \rightarrow 011001, 1 \rightarrow 0, 2 \rightarrow 01101$.

(f) $(13, \frac{7}{3})$.

Proof.

(a) $(3, 2 + \alpha)$ is achieved by any Sturmian word \mathbf{x} with slope $[0, 3, 1, 1, 1, 1, \dots] = (5 - \sqrt{5})/10$. As is well-known, a Sturmian word has exactly $n + 1$ factors of length n , and these factors are independent of the intercept of the Sturmian word. For intercept 0, the first 6 symbols are 001000, and so the four factors of \mathbf{x} of length 3 are 001, 010, 100, 000, and no complement of these words appears as a factor of \mathbf{x} .

On the other hand, we know from the proof of Proposition 15 of [4] that the critical exponent of \mathbf{x} is $2 + \alpha$. Thus there are uncountably many binary words in CAL_3 with critical exponent $2 + \alpha$.

This is best possible, as shown in Theorem 5.

(b) $(5, 3)$ is achieved by applying the morphism

$0 \rightarrow 001010001$

$1 \rightarrow 000101001$

$2 \rightarrow 000100101,$

to any ternary squarefree word. This gives uncountably many infinite binary words and exponentially many finite binary words.

For $\ell = 6$ and $e = 3$ the longest word is of length 50 and one example is 00101001001101001001101001101001001101001101001001.

(c) $(7, \frac{8}{3})$ is achieved by applying the morphism

$$\begin{aligned} 0 &\rightarrow 001001010011001010010011001001010011 \\ 1 &\rightarrow 001001010010011001010011001010010011 \\ 2 &\rightarrow 001001010010011001001010011001010011, \end{aligned}$$

to any ternary squarefree word. This gives uncountably many infinite binary words and exponentially many finite binary words.

For $\ell = 7$ and $e = \frac{8}{3}$ the longest word is of length 51 and one example is 001001100100110010100110010011001010011001010010100.

(d) $(8, \frac{5}{2})$ is achieved by applying the morphism ξ , defined as follows

$$\begin{aligned} 0 &\rightarrow 01 \\ 1 &\rightarrow 0110 \\ 2 &\rightarrow 1, \end{aligned}$$

to the infinite word \mathbf{p} mentioned above.

To determine the critical exponent of $\xi(\mathbf{p})$, we need two facts about \mathbf{p} that are proved in [6]: (1) the critical exponent of \mathbf{p} is γ (which is less than 2.481), and (2) \mathbf{p} is 2-balanced. In general, a word w over an alphabet A is C -balanced if, whenever x and y are factors of w of the same length, we have $||x|_a - |y|_a| \leq C$ for every $a \in A$.

First, consider the factor 0110110 of $\xi(\mathbf{p})$. Let us try to write it in the form $u\xi(v)w$, where u (resp., w) is a proper suffix (resp., prefix) of one of the blocks 01, 0110, or 1, and v is a factor of \mathbf{p} . We find that such a representation is unique, namely, $01.1.0110 = \xi(021)$; other possibilities, such as $01.1.01.1.0 = \xi(0202)0$, for example, imply that \mathbf{p} contains a factor such as 202, which is not the case.

Next, note that occurrences of 021 in \mathbf{p} are separated by 01, 0121, and 012101. Thus, the longest factor separating occurrences of $\xi(021)$ in $\xi(\mathbf{p})$ is $\xi(012101) = 01011010110010110$, which has length 17.

Now, suppose that $\xi(\mathbf{p})$ has a $\frac{5}{2}$ -power $X'X''X'X''X'$, where $|X'| = |X''|$. If $|X'| \geq 41$, then $\xi(\mathbf{p})$ has a factor $Y'Y''Y'Y''Y'$, where Y' begins and ends with $\xi(021)$, and $|Y'| + 34 \geq |Y''|$. The exponent of $Y'Y''Y'Y''Y'$ therefore satisfies

$$\exp(Y'Y''Y'Y''Y') \geq 2 + \frac{|Y'|}{2|Y'| + 34}. \quad (1)$$

Consequently, we have a factor $y'y''y'y''y'$ in \mathbf{p} , where $y' = \xi^{-1}(Y')$ and $y'' = \xi^{-1}(Y'')$. If $|y''| \leq |y'|$, then we have a $\frac{5}{2}$ -power in \mathbf{p} , which is a contradiction, so suppose that $|y''| \geq |y'|$, and write $y'' = y'''x$, where $|y'''| = |y'|$.

Since \mathbf{p} is 2-balanced, we have $||\xi(y''')| - |\xi(y')|| \leq 6$: we get the largest differences between these lengths when the difference between the Parikh vectors of y''' and y' is either $(0, 2, -2)$ or $(0, -2, 2)$. Also, by a computer search over all factors of \mathbf{p} of length 17, we can verify that if $|x| \geq 17$, then $|\xi(x)| \geq 43$.

Suppose that $|x| \geq 17$. We have

$$\begin{aligned} \exp(Y'Y''Y'Y''Y') &= \exp(\xi(y'y''y'y''y')) \\ &= \exp(\xi(y'y'''xy'y'''xy')) \\ &= 2 + \frac{|\xi(y')|}{|\xi(y')| + |\xi(y''')| + |\xi(x)|} \\ &\leq 2 + \frac{|\xi(y')|}{|\xi(y')| + |\xi(y')| - 6 + 43} \\ &\leq 2 + \frac{|Y'|}{2|Y'| + 37}, \end{aligned}$$

which contradicts Eq. (1).

We conclude that $|y''| \leq |y'| + 16$. This implies that

$$\begin{aligned} \exp(y'y''y'y''y') &= 2 + \frac{|y'|}{|y'| + |y''|} \\ &\geq 2 + \frac{|y'|}{2|y'| + 16}. \end{aligned}$$

However, if $|y'| \geq 203$, then

$$2 + \frac{|y'|}{2|y'| + 16} > 2.481,$$

which contradicts the fact that the critical exponent of \mathbf{p} is γ .

Completing the proof that the critical exponent of $\xi(\mathbf{p})$ is $\frac{5}{2}^+$ can now be done by a finite search. We want to verify that the only $\frac{5}{2}$ -power $X'X''X'X''X'$ in $\xi(\mathbf{p})$ is 10101. We consider two cases:

Case 1: $|X'| < 41$. In this case we simply enumerate all factors of $\xi(\mathbf{p})$ of length at most $5 \cdot 40 = 200$ and verify the claim.

Case 2: $|X'| \geq 41$. In this case the previous analysis shows that there is a corresponding repetition $y'y''y'y''y'$ in \mathbf{p} , with $|y'| \leq 202$ and $|y''| \leq 218$. Let us write $z = y'y''y'y''y'$. We need to verify that for all factors of \mathbf{p} of the form uzv , where $|u| = |v| = 6$ and $|z| \leq 3(202) + 2(218) = 1042$, the image $\xi(uzv)$ does not contain any $\frac{5}{2}$ -powers,

except for 10101. We can speed up the calculation significantly by using the results in [6] concerning the possible periods of z . In that paper, the authors showed that the periods of the overlaps in \mathbf{p} (such as z) are of the form X_n or $X_n + X_{n-2}$, where the sequence $(X_n)_{n \geq 1}$ is defined by the recurrence

$$X_1 = 1, X_2 = 2, X_3 = 4, X_4 = 7, \text{ and } X_n = X_{n-1} + X_{n-2} + X_{n-4} \text{ for } n \geq 0 .$$

Here are the first few terms of this recurrence:

| | | | | | | | | | | | | | | |
|-------|---|---|---|---|----|----|----|----|-----|-----|-----|-----|------|------|
| n | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 | 14 |
| X_n | 1 | 2 | 4 | 7 | 12 | 21 | 37 | 65 | 114 | 200 | 351 | 616 | 1081 | 1897 |

Table 1: The recurrence X_n

Since $|z| \leq 1042$, the only periods we need to consider for z come from the finite list

$$\text{PERS} = \{1, 2, 4, 5, 7, 9, 12, 16, 21, 28, 37, 49, 65, 86, 114, 151, 200, 265, 351, 465\}.$$

We then build a list of distinct factors of \mathbf{p} of the form uzv as follows. For each period $p \in \text{PERS}$ and each position $i \geq 6$ of \mathbf{p} , we find the longest factor z_i of \mathbf{p} that has period p and starts at position i . If $|z_i|/p > 2$, we add $u_i z_i v_i$ to our list, where u_i and v_i are the factors of length 6 immediately preceding and following z_i . Then for each $u_i z_i v_i$ in our list, we check that $\xi(u_i z_i v_i)$ contains no $\frac{5}{2}$ -power other than 10101. By [6, Corollary 13], every factor of \mathbf{p} of length n occurs at a position $\leq 3.65n$, so we only need to examine the first $3.65(1042 + 12) \leq 3848$ positions when conducting this search.

For $\ell = 10$ and $e = \frac{5}{2}$ the longest word is of length 75 and one example is
0010110100110010110100110110010110100110010110100110110010110100110110011001100.

(e) $(11, \gamma')$ is achieved by $\psi(\mathbf{p})$.

For $\ell = 12$ and $e = \gamma$, the optimality is proved as follows. Let $z = 1001011001$. We can check that $\psi(\mathbf{p})$ avoids z and contains \bar{z} , z^R , and $\overline{z^R}$. Let x be the prefix of length 40 of $\psi(\mathbf{p})$. A computer check shows that there is no infinite $\frac{5}{2}$ -free binary word with $\ell \leq 12$ that avoids simultaneously x , \bar{x} , x^R , and $\overline{x^R}$.

By symmetry, we consider a bi-infinite $\frac{5}{2}$ -free binary word \mathbf{w} with $\ell \leq 12$ that contains x . Let X be the set containing the complements of the factors of length 12 of x . Thus $\ell \leq 12$ means that \mathbf{w} avoids X .

We compute the set S of factors f such that efg is $\frac{5}{2}$ -free and avoids X and $|e| = |f| = |g| = 100$. We compute the set S' of factors of $\psi(\mathbf{p})$ of length 100. We verify that $S = S'$. This means that $\mathbf{w} = \psi(\mathbf{v})$ for some bi-infinite ternary word \mathbf{v} . Moreover, by considering the pre-images by ψ of $\mathbf{w} = \psi(\mathbf{v})$ and $\psi(\mathbf{p})$, this implies that \mathbf{v} and \mathbf{p}

have the same set of factors of length $100/\max(|\psi(0)|, |\psi(1)|, |\psi(2)|) = 16$. In particular, \mathbf{v} avoids the set $\{00, 11, 22, 20, 212, 0101, 02102, 121012, 01021010, 21021012102\}$ mentioned in [6, Theorem 14].

Also, \mathbf{v} is cube-free since \mathbf{w} is $\frac{5}{2}$ -free. By [6, Theorem 14], we know that \mathbf{v} has the same set of factors as \mathbf{p} . Thus \mathbf{w} has the same set of factors as $\psi(\mathbf{p})$. So, the critical exponent of $\psi(\mathbf{p})$ is optimal for $\ell = 12$.

(f) $(13, \frac{7}{3})$ is achieved by applying the morphism

0 \rightarrow 001001100101101001100101100100110100110010110100110010011010011001011
1 \rightarrow 001001100101101001100100110100110010110100110010110010011010011001011
2 \rightarrow 0010011001011010011001001101001100100110100110010011010011001011010011001011

to any ternary squarefree word. This gives uncountably many infinite binary words and exponentially many finite binary words.

By Proposition 1, if $e = \frac{7}{3}$, then there are arbitrarily long complemented words.

□

3 The class CAN_n

One could also try to minimize the total number n of complemented words that appear. Obviously n has to be even. For $n = 0$ the only such infinite words are $0^\omega, 1^\omega$. For $n = 2$ the only such infinite words are 01^ω and 10^ω . For larger n the situation is summed up in the following theorem:

Theorem 3. *There exists an infinite β^+ -free binary word having at most n complemented words, for the following pairs (n, β) :*

(a) $(4, 2 + \alpha)$

(b) $(8, 3)$

(c) $(24, \frac{8}{3})$

(d) $(36, \frac{5}{2})$

(e) $(64, \gamma')$, where γ' is the number defined in Theorem 2

(f) $(90, \frac{7}{3})$.

Proof. For the positive part, we use the same words as in Theorem 2. That is, for every $\beta \in \{2 + \alpha, 3, \frac{8}{3}, \frac{5}{2}, \gamma', \frac{7}{3}\}$, the infinite β^+ -free binary words given in Theorem 2 to achieve the pair (ℓ, β) also achieve the pair (n, β) in Theorem 3. It is not hard to count the complemented factors in such words since their length is less than ℓ . Now let us consider the negative part.

- (a) For $4 \leq t \leq 6$ the best critical exponent is $2 + \alpha$. To see this note that any such word must avoid having both x and \bar{x} as factors if $|x| \geq 4$, since taking non-empty prefixes of x and \bar{x} gives at least 8 complemented factors. Hence, the best critical exponent is $2 + \alpha$ by Theorem 5.
- (b) For $8 \leq t \leq 22$ the best critical exponent is 3^+ . For $t = 22$ and $e = 3$ the longest word is of length 50 and one example is
00101001001101001001101001101001001101001101001001.
- (c) For $24 \leq t \leq 34$ the best critical exponent is $\frac{8}{3}^+$. For $t = 34$ and $e = \frac{8}{3}$ the longest word is of length 51 and one example is
001001100100110010100110010011001010011001010010100.
- (d) For $36 \leq t \leq 62$ the best critical exponent is $\frac{5}{2}^+$. For $t = 62$ and $e = \frac{5}{2}$ the longest word is of length 73 and one example is
0010110100110010110100110110010110100110010110100110110010110100110110010.
- (e) For $64 \leq t \leq 88$ the best critical exponent is γ' . We use the proof of optimality of Theorem 2 (e) and replace the condition $\ell \leq 12$ by $t \leq 88$.
- (f) For $90 \leq t < \infty$, the best critical exponent is $\frac{7}{3}^+$. By Proposition 1, if $e = \frac{7}{3}$, then there are arbitrarily many complemented words.

□

4 Complexity threshold

Notice that we use two different constructions of infinite words in the proof of Theorem 2. When possible, that is in cases (b), (c), and (f), we use morphic images of arbitrary ternary square-free words, thus showing that exponentially many binary words have the considered property. Otherwise, in cases (a), (d), and (e), every bi-infinite binary word with the considered property has the same set of factors as one or a few morphic words.

In this section, we consider the latter case and we relax the constraints on the critical exponent or the complementary factors in order to get exponential factor complexity.

Let h be the morphism that maps $0 \mapsto 0$ and $1 \mapsto 01$.

Lemma 4. *Every bi-infinite binary 4-free word avoiding pairs of complementary factors of length 4 and $\{1001001, 0110110\}$ has the same set of factors as either $h(\mathbf{f})$ or $\overline{h(\mathbf{f})}$.*

Proof. We compute the set S of factors y such that there exists a binary 4-free word xyz avoiding pairs of complementary factors of length 4 and $\{1001001, 0110110\}$ and such that $|x| = |y| = |z| = 100$. We observe that S consists of the (disjoint) union of the factors of length 100 of $h(\mathbf{f})$ and $\overline{h(\mathbf{f})}$. Let \mathbf{w} be a bi-infinite binary 4-free word avoiding pairs of complementary factors of length 4 and $\{1001001, 0110110\}$. By symmetry, we suppose that \mathbf{w} contains 00 , so that \mathbf{w} has the same factors of length 100 as $h(\mathbf{f})$. So $\mathbf{w} \in \{0, 01\}^\omega$, that

is, $\mathbf{w} = h(\mathbf{v})$ for some bi-infinite binary word \mathbf{v} . Since \mathbf{w} is 4-free, so is \mathbf{v} . Moreover, by considering the pre-images by h of $\mathbf{w} = h(\mathbf{v})$ and $h(\mathbf{f})$, this implies that \mathbf{v} and \mathbf{f} have the same set of factors of length $100/\max(|h(0)|, |h(1)|) = 50$. In particular, \mathbf{v} avoids $\{11, 000, 10101\}$. By [2, Thm. 8], \mathbf{v} has the same set of factors as \mathbf{f} . Thus \mathbf{w} has the same set of factors as $h(\mathbf{f})$. \square

Theorem 5. *Every bi-infinite binary $\frac{11}{3}$ -free word that contains no pair of complementary factors of length 4 has the same set of factors as $h(\mathbf{f})$ or the same set of factors as $\overline{h(\mathbf{f})}$.*

Proof. Suppose that such a word \mathbf{w} contains 1001001.

First, suppose that \mathbf{w} contains 11. Since \mathbf{w} avoids $0110 = \overline{1001}$, \mathbf{w} contains 111. Since \mathbf{w} avoids the 4-power 1111, \mathbf{w} contains 01110. Since \mathbf{w} avoids $1101 = \overline{0010}$, \mathbf{w} contains 011100. Symmetrically, since \mathbf{w} avoids $1011 = \overline{0100}$, \mathbf{w} contains 0011100. So \mathbf{w} contains both 0011 and 1100, a contradiction. Thus, \mathbf{w} avoids 11.

Now, suppose that \mathbf{w} contains 101. Since \mathbf{w} avoids 11, \mathbf{w} contains 01010. So \mathbf{w} contains both 0101 and 1010, a contradiction. Thus, \mathbf{w} avoids 101.

Since \mathbf{w} contains 1001001 and avoids 11, \mathbf{w} contains 010010010. Since \mathbf{w} contains 010010010 and avoids 101, \mathbf{w} contains 00100100100. This is a contradiction, since $00100100100 = (001)^{11/3}$.

So \mathbf{w} avoids 1001001. By symmetry, \mathbf{w} avoids 0110110.

By Lemma 4, \mathbf{w} has the same set of factors as either $h(\mathbf{f})$ or $\overline{h(\mathbf{f})}$. \square

By contrast, there exist exponentially many binary $\frac{11}{3}^+$ -free words with no pair of complementary factors of length 4.

Theorem 6. *The image of any ternary squarefree word by the 21-uniform morphism*

$$\begin{aligned} 0 &\rightarrow 0001001001000100001001 \\ 1 &\rightarrow 0001000100100100001001 \\ 2 &\rightarrow 000100010010001001001 \end{aligned}$$

is a $\frac{11}{3}^+$ -free word containing only 0, 1, 01, and 10 as complementary factors.

Let ρ be the morphism defined as follows

$$\begin{aligned} 0 &\rightarrow 01100101101 \\ 1 &\rightarrow 0110010 \\ 2 &\rightarrow 011001. \end{aligned}$$

Using the technique of Theorem 2 (d), we can show that $\rho(\mathbf{p})$ is a $\frac{5}{2}^+$ -free word with no pair of complementary factors of length 8 and exactly 40 complementary factors.

Theorem 7. *Every bi-infinite binary $\frac{29}{11}$ -free word that contains no pair of complementary factors of length 8 has the same set of factors as either $\xi(\mathbf{p})$, $\overline{\xi(\mathbf{p})}$, $\xi(\mathbf{p})^R$, $\overline{\xi(\mathbf{p})^R}$, $\rho(\mathbf{p})$, $\overline{\rho(\mathbf{p})}$, $\rho(\mathbf{p})^R$, or $\overline{\rho(\mathbf{p})^R}$.*

Proof. First, we show the following. If \mathbf{w} is a bi-infinite cube-free binary word and every factor of length 21 of \mathbf{w} is also a factor of $\xi(\mathbf{p})$, then \mathbf{w} has the same set of factors as $\xi(\mathbf{p})$.

We compute the set S of factors f such that there exists $u = efg$ where u is cube-free, every factor of length 21 of u is a factor of $\xi(\mathbf{p})$, and $|e| = |f| = |g| = 500$. We verify that every factor $f \in S$ is a factor of $\xi(\mathbf{p})$.

This means that $\mathbf{w} = \xi(\mathbf{v})$ for some bi-infinite ternary word \mathbf{v} . Moreover, by considering the pre-images by ξ of $\mathbf{w} = \xi(\mathbf{v})$ and $\xi(\mathbf{p})$, this implies that \mathbf{v} and \mathbf{p} have the same set of factors of length $500/\max(|\xi(0)|, |\xi(1)|, |\xi(2)|) = 125$. In particular, \mathbf{v} has the same set of factors as \mathbf{p} by [6, Theorem 14].

Similarly, we show that if \mathbf{w} is a bi-infinite cube-free binary word and every factor of length 63 of \mathbf{w} is also a factor of $\rho(\mathbf{p})$, then \mathbf{w} has the same set of factors as $\rho(\mathbf{p})$.

Finally, we compute the set X of factors f such that there exists $u = efg$ where u is $\frac{29}{11}$ -free, u contains no pair of complementary factors of length 8, $|f| = 200$, and $|e| = |g| = 80$. We verify that every factor $f \in X$ is a factor of either $\xi(\mathbf{p})$, $\xi(\mathbf{p})^R$, $\xi(\mathbf{p})^R$, $\rho(\mathbf{p})$, $\rho(\mathbf{p})^R$, or $\rho(\mathbf{p})^R$. Since $\frac{29}{11}$ -free words are cube-free, Theorem 7 follows by the two previous results and the symmetries by complement and reversal. \square

By contrast, there exist exponentially many binary $\frac{29}{11}^+$ -free words containing no pair of complementary factors of length 8 and exactly 36 complementary factors.

Theorem 8. *The image of any ternary squarefree word by the 42-uniform morphism*

$$\begin{aligned} 0 &\rightarrow 001010010110010011001011001001100101001011 \\ 1 &\rightarrow 001010010110010011001010010110010011001011 \\ 2 &\rightarrow 001001100101001011001010010110010011001011 \end{aligned}$$

is a $\frac{29}{11}^+$ -free word with no pair of complementary factors of length 8.

We also get exponentially many words if we allow complementary factors of length 8.

Theorem 9. *The image of any ternary squarefree word by the 31-uniform morphism*

$$\begin{aligned} 0 &\rightarrow 0010100110010011001010011001011 \\ 1 &\rightarrow 0010100101100101001100101001011 \\ 2 &\rightarrow 0010011001011001010010110010011 \end{aligned}$$

is a $\frac{5}{2}^+$ -free word with no pair of complementary factors of length 9 and exactly 40 complementary factors.

5 Conclusion

It is quite nice that many morphic words, namely $h(\mathbf{f})$, $\xi(\mathbf{p})$, $\rho(\mathbf{p})$, and $\psi(\mathbf{p})$, show up in this study.

To complete the work in this paper, it remains to determine the critical exponent γ' of $\psi(\mathbf{p})$.

Conjecture 10. $\psi(\mathbf{p})$ and \mathbf{p} have the same critical exponent.

Perhaps some reader can prove this conjecture.

References

- [1] G. Badkobeh and M. Crochemore. Finite-repetition threshold for infinite ternary words. In P. Ambrož, Š. Holub, and Z. Masáková, editors, *WORDS 2011*, Vol. 63 of *Elec. Proc. Theor. Comput. Sci.*, pp. 37–43. Open Publishing Association, 2011.
- [2] A. Baranwal, J. D. Currie, L. Mol, P. Ochem, N. Rampersad, and J. Shallit. Antisquares and critical exponents. Arxiv preprint arXiv:2209.09223 [math.CO], available at <https://arxiv.org/abs/2209.09223>, 2022.
- [3] J. Berstel. Fibonacci words—a survey. In G. Rozenberg and A. Salomaa, editors, *The Book of L*, pp. 13–27. Springer-Verlag, 1986.
- [4] A. Carpi and A. de Luca. Special factors, periodicity, and an application to Sturmian words. *Acta Informatica* **36** (2000), 983–1006.
- [5] J. D. Currie, L. Mol, and N. Rampersad. The repetition threshold for binary rich words. *Discrete Math. & Theoret. Comput. Sci.* **22**(1) (2020), DMTCS-22-1-6 (electronic), <https://dmtcs.episciences.org/6082>
- [6] J. D. Currie, P. Ochem, N. Rampersad, and J. Shallit. Properties of a ternary infinite word. Arxiv preprint arXiv:2206.01776 [cs.DM], available at <https://arxiv.org/abs/2206.01776>, 2022.
- [7] J. Currie and N. Rampersad. A proof of Dejean’s conjecture. *Math. Comp.* **80** (2011), 1063–1070.
- [8] F. Dejean. Sur un théorème de Thue. *J. Combin. Theory. Ser. A* **13** (1972), 90–99.
- [9] L. Dvořáková, K. Medková, and E. Pelantová. Complementary symmetric Rote sequences: the critical exponent and the recurrence function. *Discrete Math. & Theoret. Comput. Sci.* **22**(1) (2020), DMTCS-22-1-20 (electronic), <https://dmtcs.episciences.org/6519/>
- [10] L. Dvořáková, D. Opočenská, E. Pelantová, and A. M. Shur. On minimal critical exponent of balanced sequences. Preprint, available at https://papers.ssrn.com/sol3/papers.cfm?abstract_id=4011686, 2022.
- [11] F. Fiorenzi, P. Ochem, and E. Vaslet. Bounds for the generalized repetition threshold. *Theoret. Comput. Sci.* **412** (2011), 2955–2963.

- [12] L. Ilie, P. Ochem, and J. Shallit. A generalization of repetition threshold. *Theoret. Comput. Sci.* **345** (2005), 359–369.
- [13] J. Karhumäki and J. Shallit. Polynomial versus exponential growth in repetition-free binary words. *J. Combin. Theory. Ser. A* **105**(2) (2004), 335–347.
- [14] H. Mousavi and J. Shallit. Repetition avoidance in circular factors. In M.-P. Béal and O. Carton, editors, *DLT 2013*, Vol. 7907 of *Lecture Notes in Computer Science*, pp. 384–395. Springer-Verlag, 2013.
- [15] N. Rampersad, J. Shallit, and É. Vandomme. Critical exponents of infinite balanced words. *Theoret. Comput. Sci.* **777** (2019), 454–463.
- [16] M. Rao. Last cases of Dejean’s conjecture. *Theoret. Comput. Sci.* **412** (2011), 3010–3018.
- [17] A. V. Samsonov and A. M. Shur. On Abelian repetition threshold. *RAIRO Inform. Théor. App.* **46** (2012), 147–163.
- [18] J. Sawada, B. Stevens, and A. Williams. De Bruijn sequences for the binary strings with maximum density. In N. Katoh and A. Kumar, editors, *WALCOM 2011*, Vol. 6552 of *Lecture Notes in Computer Science*, pp. 182–190. Springer-Verlag, 2011.
- [19] J. Shallit. Simultaneous avoidance of large squares and fractional powers in infinite binary words. *Internat. J. Found. Comp. Sci.* **15** (2004), 317–327.
- [20] J. Shallit. Minimal critical exponents for palindromes. ArXiv preprint, arXiv:1612.05320 [cs.FL]. Available at <https://arxiv.org/abs/1612.05320>, 2016.