

Quantum Wasserstein distance based on an optimization over separable states

Géza Tóth^{1,2,3,4,5,*} and József Pitrik^{5,6,7,†}

¹Theoretical Physics, University of the Basque Country UPV/EHU, ES-48080 Bilbao, Spain

²EHU Quantum Center, University of the Basque Country UPV/EHU, Barrio Sarriena s/n, ES-48940 Leioa, Biscay, Spain

³Donostia International Physics Center (DIPC), ES-20080 San Sebastián, Spain

⁴IKERBASQUE, Basque Foundation for Science, ES-48011 Bilbao, Spain

⁵Institute for Solid State Physics and Optics, Wigner Research Centre for Physics, HU-1525 Budapest, Hungary

⁶Alfréd Rényi Institute of Mathematics, Reáltanoda u. 13-15., HU-1053 Budapest, Hungary

⁷Department of Analysis, Institute of Mathematics, Budapest University of Technology and Economics, Műegyetem rkp. 3., HU-1111 Budapest, Hungary

(Dated: October 28, 2022)

We define the quantum Wasserstein distance such that the optimization is carried out over bipartite separable states rather than bipartite quantum states in general, and examine its properties. Surprisingly, we find that its self-distance is related to the quantum Fisher information. We discuss how the quantum Wasserstein distance introduced is connected to criteria detecting quantum entanglement. We define variance-like quantities that can be obtained from the quantum Wasserstein distance by replacing the minimization over quantum states by a maximization. We extend our results to a family of generalized quantum Fisher information.

I. INTRODUCTION

A classical Wasserstein distance is a metric between probability distributions μ and ν , induced by the problem of optimal mass transportation [1, 2]. It reflects the minimal effort that is required in order to morph the mass of μ into the mass distribution of ν . Methods based on the theory of optimal transport and advantageous properties of Wasserstein metrics have achieved great success in several important fields of pure mathematics including probability theory [3, 4], theory of (stochastic) partial differential equations [5, 6], variational problems [7, 8] and geometry of metric spaces [9–12]. In recent years, there have been a lot of results concerning the description of isometries of Wasserstein spaces, too [13–16]. Moreover, optimal transport and Wasserstein metric are also used as tools in applied mathematics. In particular, there are applications to image and signal processing [17–19], medical imaging [20, 21] and machine learning [22–27].

The non-commutative generalization, the so-called quantum optimal transport has been at the center of attention, as it lead to the definition of several new and very useful notions in quantum physics. The first, semi-classical approach of Życzkowski and Slomczynski, has been motivated by applications in quantum chaos [28–30]. The method of Biane and Voiculescu is related to free probability [31], while the one of Carlen, Maas, Datta, and Rouzé [32–36] is based on a dynamical interpretation. Caglioti, Golse, Mouhot, and Paul presented an approach based on a static interpretation [37–42]. Finally De Palma and Trevisan used quantum channels [43], and De Palma, Marvian, Trevisan, and Lloyd defined the quantum earth mover’s distance, i.e., the quantum Wasserstein distance order 1 [44], while Bistrón, Cole, Eckstein, Friedland and Życzkowski formulated a quantum Wasserstein distance based on an antisymmetric cost function [45–47].

One of the key results of quantum optimal transport is the definition of the quantum Wasserstein distance [28–30, 37–44, 48, 49]. It has the often desirable feature that it is not necessarily maximal for two quantum states orthogonal to each other, which is beneficial, for instance, when performing learning on quantum data [50]. Some of the properties of the new quantities are puzzling, yet point to profound relations between seemingly unrelated fields of quantum physics. For instance, the quantum Wasserstein distance order 2 of the quantum state from itself can be nonzero, while in the classical case the self-distance is always zero. In particular, as we have mentioned, the quantum Wasserstein distance has been defined based on a quantum channel formalism [43], and it has been shown that the square of the self-distance is equal to the Wigner-Yanase skew information of the quantum state [51]. At this point the questions arises: Can other fields of quantum physics help to interpret these results? For example, is it possible to relate the findings above to entanglement theory [52–54] such that we obtain new and meaningful relations naturally?

Before presenting our results, let us summarize the definitions of the quantum Wasserstein distance.

Definition 1.—The square of the distance between two quantum states described by the density matrices ϱ and σ is given by De Palma and Trevisan as [55]

$$D_{\text{DPT}}(\varrho, \sigma)^2 = \frac{1}{2} \min_{\varrho_{12}} \sum_{n=1}^N \text{Tr}[(H_n^T \otimes \mathbb{1} - \mathbb{1} \otimes H_n)^2 \varrho_{12}],$$

s. t. $\varrho_{12} \in \mathcal{D}$,
 $\text{Tr}_2(\varrho_{12}) = \varrho^T$,
 $\text{Tr}_1(\varrho_{12}) = \sigma$, (1)

where A^T denotes the matrix transpose of A , and H_1, H_2, \dots, H_N are Hermitian operators, while \mathcal{D} is the set of density matrices, i.e., Hermitian matrices fulfilling

$$\varrho_{12} = \varrho_{12}^\dagger, \text{Tr}(\varrho_{12}) = 1, \varrho_{12} \geq 0. \quad (2)$$

In this approach, there is a bipartite density matrix ϱ_{12} , called coupling, corresponding to any transport map between

* toth@alumni.nd.edu

† pitrik@math.bme.hu

ϱ and σ , and vice versa, there is a transport map corresponding to any coupling [43]. Moreover, it has been shown that for the self-distance of a state [43]

$$D_{\text{DPT}}(\varrho, \varrho)^2 = \sum_{n=1}^N I_{\varrho}(H_n) \quad (3)$$

holds, where the Wigner-Yanase skew information is defined as [51]

$$I_{\varrho}(H) = \text{Tr}(H^2 \varrho) - \text{Tr}(H \sqrt{\varrho} H \sqrt{\varrho}). \quad (4)$$

This profound result connects seemingly two very different notions of quantum physics, as it has been mentioned in the introduction.

The Wasserstein distance has also been defined in a slightly different way.

Definition 2.—Golse, Mouhot, Paul and Caglioti defined the square of the distance as [37–42]

$$\begin{aligned} D_{\text{GMPC}}(\varrho, \sigma)^2 &= \frac{1}{2} \min_{\varrho_{12}} \sum_{n=1}^N \text{Tr}[(H_n \otimes \mathbb{1} - \mathbb{1} \otimes H_n)^2 \varrho_{12}], \\ &\text{s. t. } \varrho_{12} \in \mathcal{D}, \\ &\text{Tr}_2(\varrho_{12}) = \varrho, \\ &\text{Tr}_1(\varrho_{12}) = \sigma. \end{aligned} \quad (5)$$

In this paper, we will obtain new quantities by restricting the optimization to separable states in the above definitions. We will show that, in this case, the square of the self-distance equals the quantum Fisher information times a constant, while in Eq. (3) it was related to the Wigner-Yanase skew information. The quantum Fisher information is a central quantity of quantum metrology, a field that is concerned with metrological tasks in which the quantumness of the system plays an essential role [56–59]. Recent findings show that the Quantum Fisher information is the convex roof of the variance, apart from a constant factor [60, 61], which allowed, for instance, to derive novel uncertainty relations [62, 63], and will also be used in this article.

The paper is organized as follows. In Sec. II, we summarize basic facts connected to quantum metrology. In Sec. III, we summarize entanglement theory. In Sec. IV, we show how to transform the optimization over decompositions of the density matrix into an optimization of an expectation value over separable states. In Sec. V, we show some applications of these ideas for the Wasserstein distance. We define a novel type of Wasserstein distance based on an optimization over separable states. In Sec. VI, we define variance-like quantities from the Wasserstein distance. In Sec. VII, we discuss how such a Wasserstein distance and the variance-like quantity mentioned above can be used to construct entanglement criteria. In Sec. VIII, we introduce further quantities similar to the formulas giving the Wasserstein distance, but they involve the variance of two-body quantities rather than the second moment. In Sec. IX, we extend our ideas to various types of generalized quantum Fisher information.

II. QUANTUM METROLOGY

Before discussing our main results, we review some of the fundamental relations of quantum metrology. A basic metrological task in an interferometer is estimating the small angle θ for a unitary dynamics

$$U_{\theta} = \exp(-iH\theta), \quad (6)$$

where H is the Hamiltonian. The precision is limited by the Cramér-Rao bound as [56–59, 64–70]

$$(\Delta\theta)^2 \geq \frac{1}{\nu \mathcal{F}_Q[\varrho, H]}, \quad (7)$$

where ν is the number of independent repetitions, and the quantum Fisher information, a central quantity in quantum metrology is defined by the formula [64–68]

$$\mathcal{F}_Q[\varrho, H] = 2 \sum_{k,l} \frac{(\lambda_k - \lambda_l)^2}{\lambda_k + \lambda_l} |\langle k|H|l\rangle|^2, \quad (8)$$

where the density matrix has the eigendecomposition

$$\varrho = \sum_k \lambda_k |k\rangle\langle k|. \quad (9)$$

The quantum Fisher information is bounded from above by the variance and from below by the Wigner-Yanase skew information as

$$4I_{\varrho}(H) \leq \mathcal{F}_Q[\varrho, H] \leq 4(\Delta H)_{\varrho}^2, \quad (10)$$

where if ϱ is pure then the equality holds for both inequalities [66]. For clarity, we add that the variance is defined as

$$(\Delta H)_{\varrho}^2 = \langle H^2 \rangle_{\varrho} - \langle H \rangle_{\varrho}^2, \quad (11)$$

where the expectation value is calculated as

$$\langle H \rangle_{\varrho} = \text{Tr}(\varrho H). \quad (12)$$

The quantum Fisher information is, apart from a constant factor, can be defined also as the convex roof of the variance [60, 61]

$$\mathcal{F}_Q[\varrho, H] = 4 \min_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k (\Delta H)_{\Psi_k}^2, \quad (13)$$

where the optimization is over pure state decompositions of the type

$$\varrho = \sum_k p_k |\Psi_k\rangle\langle \Psi_k|, \quad (14)$$

where for the probabilities $p_k \geq 0$ and $\sum_k p_k = 1$ hold, and the pure states $|\Psi_k\rangle$ are not assumed to be orthogonal to each other.

Apart from the quantum Fisher information, the variance can also be given as a roof [60, 61]

$$(\Delta H)_{\varrho}^2 = \max_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k (\Delta H)_{\Psi_k}^2. \quad (15)$$

Finally, note that convex and concave roofs of more complicated expressions can also be computed. For instance,

$$\min_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k \sum_{n=1}^N (\Delta H_n)_{\Psi_k}^2 \geq \frac{1}{4} \sum_{n=1}^N \mathcal{F}_Q[\varrho, H_n], \quad (16)$$

and the expression

$$\max_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k \sum_{n=1}^N (\Delta H_n)_{\Psi_k}^2 \leq \sum_{n=1}^N (\Delta H_n)_{\varrho}^2, \quad (17)$$

are the convex and concave roofs, respectively, of the sum of several variances over the decompositions of ϱ given in Eq. (14). They play a role in the derivation of entanglement conditions [62], and will also appear in our results about the quantum Wasserstein distance. We add that there is an equality in Eq. (17) for $N = 2$ [71, 72].

III. ENTANGLEMENT THEORY

Next, we review entanglement theory [52–54]. A bipartite quantum state is separable if it can be given as [73]

$$\sum_k p_k |\Psi_k\rangle\langle\Psi_k| \otimes |\Phi_k\rangle\langle\Phi_k|, \quad (18)$$

where p_k are for the probabilities, $|\Psi_k\rangle$ and $|\Phi_k\rangle$ are pure quantum states. We will denote the set of separable states by \mathcal{S} . The mixture of two separable states is also separable, thus the \mathcal{S} set is convex. If a quantum state cannot be written as Eq. (18) then it is called entangled.

A relevant subset of separable states are the symmetric separable states, which can be given as [74, 75]

$$\sum_k p_k |\Psi_k\rangle\langle\Psi_k| \otimes |\Psi_k\rangle\langle\Psi_k|. \quad (19)$$

We will denote the set of symmetric separable states by \mathcal{S}' . The mixture of two symmetric separable states is also symmetric separable, thus the

$$\mathcal{S}' \subset \mathcal{S} \quad (20)$$

set is also convex. Symmetry means that for such states for the expectation value

$$\langle P_s \rangle = 1 \quad (21)$$

holds, where P_s is the projector to the symmetric subspace defined by the basis vectors $|nn\rangle$ and $(|nm\rangle + |mn\rangle)/\sqrt{2}$ for $n \neq m$.

The set of quantum states with a positive partial transpose (PPT) \mathcal{P} consists of the states for which

$$\varrho^{T_k} \geq 0 \quad (22)$$

holds, where T_k is a partial transposition according to the k^{th} subsystem. \mathcal{P} is clearly a convex set. Moreover, all separable states given in Eq. (18) fulfill Eq. (22), thus

$$\mathcal{P} \supset \mathcal{S} \quad (23)$$

holds. For two qubits and for a qubit-qutrit unit, i.e., for 2×2 and 2×3 systems the set of PPT states equals the set of separable states [76–78].

It is generally very difficult to decide whether a state is separable or not, while in general it is very simple to decide whether the condition given in Eq. (22) is fulfilled. Such conditions can be even part of semidefinite programs used to solve various optimization problems (e.g., Ref. [79]). Thus, often the set of PPT quantum states, \mathcal{P} is used instead of the separable states. Since for small systems $\mathcal{P} = \mathcal{S}$, optimization problems over separable states can be solved exactly for those cases. For larger systems, by optimizing over states in \mathcal{P} instead of states in \mathcal{S} , we get lower or upper bounds.

Finally, we will define the set of symmetric PPT states \mathcal{P}' for which naturally

$$\mathcal{S}' \subset \mathcal{P}' \subset \mathcal{P} \quad (24)$$

holds, while for 2×2 and 2×3 systems, we have $\mathcal{P}' = \mathcal{S}'$.

Convex roofs and concave roofs play also a central role in entanglement theory. The entanglement of formation is defined as a convex roof over pure components of the von Neumann entropy of the reduced state [80, 81]

$$E_F(\varrho) = \min_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k E_E(\Psi_k), \quad (25)$$

where the optimization is over the decompositions given in Eq. (14), and the entanglement entropy of the pure components is given as

$$E_E(\Psi_k) = S[\text{Tr}_A(\Psi_k)], \quad (26)$$

where S is the von Neumann entropy. $E_F(\varrho)$ is the minimum entanglement needed to create the state. On the other hand, the entanglement of assistance is obtained as a concave roof [82, 83]

$$E_A(\varrho) = \max_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k E_E(\Psi_k). \quad (27)$$

These quantities correspond to the following scenario. Let us assume that the bipartite quantum state living on parties A and B is realized as the reduced state of a pure state living on parties A, B and C. Party C can make various measurements and with that it can produce any decomposition of the bipartite quantum state to pure states. If party C also sends the measurement outcome to A and B, then average entanglement will be the weighted sum of the entanglement of the components

$$\sum_k p_k E_E(\Psi_k). \quad (28)$$

If party C wants to help, then Eq. (28) can reach the concave roof, which is just the entanglement of assistance given in Eq. (27), while it is always larger than or equal to the entanglement of formation given in Eq. (25).

Now, we will present some conditions that can be used to detect entanglement. That is, they are inequalities fulfilled by all separable states. If they are violated by a quantum state, it must be entangled.

Next, we will introduce an entanglement condition based on the sum of several variances [84–87]. Let us consider a full set of traceless observables $\{G_n\}_{n=1}^{d^2-1}$ for d -dimensional systems fulfilling

$$\text{Tr}(G_n G_{n'}) = 2\delta_{nn'}. \quad (29)$$

Any traceless Hermitian observable can be obtained as a linear combination of G_n . In other words, G_n are the $\text{SU}(d)$ generators. It is known that for pure states (see e.g., Ref. [87])

$$\sum_{n=1}^{d^2-1} (\Delta G_n)^2 = 2(d-1) \quad (30)$$

holds. Due to the concavity of the variance it follows that for mixed states we have [87]

$$\sum_{n=1}^{d^2-1} (\Delta G_n)^2 \geq 2(d-1). \quad (31)$$

Let us now consider a $d \times d$ system. For a product state, $|\Psi\rangle \otimes |\Phi\rangle$, we obtain [84, 87]

$$\begin{aligned} & \sum_{n=1}^{d^2-1} [\Delta(G_n^T \otimes \mathbb{1} - \mathbb{1} \otimes G_n)]_{\Psi \otimes \Phi}^2 \\ &= \sum_{n=1}^{d^2-1} (\Delta G_n^T)_{\Psi}^2 + \sum_{n=1}^{d^2-1} (\Delta G_n)_{\Phi}^2 \\ &= 4(d-1), \end{aligned} \quad (32)$$

where in the last inequality we used that for pure states Eq. (30) holds. We also used the fact that if $\{G_n\}_{n=1}^{d^2-1}$ is a full set of observables with the properties mentioned above, then $\{G_n^T\}_{n=1}^{d^2-1}$ is also a full set of such observables. Then, for bipartite separable states given in Eq. (18) [84–87]

$$\sum_{n=1}^{d^2-1} [\Delta(G_n^T \otimes \mathbb{1} - \mathbb{1} \otimes G_n)]^2 \geq 4(d-1) \quad (33)$$

holds due to the concavity of the variance. Any state that violates the inequality in Eq. (33) is entangled. The left-hand side of Eq. (33) is zero for the maximally entangled state

$$|\Psi_{\text{me}}\rangle = \frac{1}{\sqrt{d}} \sum_{k=0}^{d-1} |k\rangle|k\rangle. \quad (34)$$

Thus, we say that the criterion given in Eq. (33) detects entangled states in the vicinity of the state given in Eq. (34).

Let us consider the $d = 2$ case concretely. Equation (33) can be rewritten as

$$\begin{aligned} & [\Delta(\sigma_x \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_x)]^2 \\ & + [\Delta(\sigma_y \otimes \mathbb{1} + \mathbb{1} \otimes \sigma_y)]^2 \\ & + [\Delta(\sigma_z \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_z)]^2 \geq 4, \end{aligned} \quad (35)$$

where σ_l are Pauli spin matrices. The left-hand side of the inequality given in Eq. (35) is zero for the state

$$\frac{1}{\sqrt{2}}(|00\rangle + |11\rangle). \quad (36)$$

We can have a similar construction with only three operators for any system size. Let us consider the usual angular momentum operators j_x, j_y and j_z , living in a d -dimensional system, fulfilling

$$j_x^2 + j_y^2 + j_z^2 = j(j+1)\mathbb{1}, \quad (37)$$

where $d = 2j + 1$. In particular, let us define the matrix [88]

$$j_z = \text{diag}(-j, -j+1, \dots, j-1, j). \quad (38)$$

Then, we need the ladder operators

$$(j_+)_{m,n} = \delta_{m,n+1} \sqrt{j(j+1) - (j-n)(j+1-n)} \quad (39)$$

and $j_- = j_+^\dagger$, where $1 \leq m, n \leq d$ and δ_{kl} is the well-known Kronecker delta. Then, we define the x and y components as

$$j_x = \frac{j_+ + j_-}{2}, \quad j_y = \frac{j_+ - j_-}{2i}. \quad (40)$$

After clarifying the operators used, we need the uncertainty relation

$$(\Delta j_x)^2 + (\Delta j_y)^2 + (\Delta j_z)^2 \geq j. \quad (41)$$

Based on Eq. (41), it can be proved that for separable states we have [87, 89–91]

$$\begin{aligned} & [\Delta(j_x^T \otimes \mathbb{1} - \mathbb{1} \otimes j_x)]^2 \\ & + [\Delta(j_y^T \otimes \mathbb{1} - \mathbb{1} \otimes j_y)]^2 \\ & + [\Delta(j_z^T \otimes \mathbb{1} - \mathbb{1} \otimes j_z)]^2 \geq 2j, \end{aligned} \quad (42)$$

c. f. Eq. (35). The left-hand side of the inequality given in Eq. (42) is zero for the maximally entangled state given in Eq. (34).

We now present a simple expression for which the maximum for general states is larger than the maximum for separable quantum states. We know that for separable states [90, 92]

$$\langle (\sigma_x \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_x)^2 + (\sigma_y \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_y)^2 \rangle \leq 6, \quad (43)$$

while the maximum for quantum states is 8 and it is taken by the singlet state

$$\frac{1}{\sqrt{2}}(|01\rangle - |10\rangle). \quad (44)$$

We add that the product state $|01\rangle_x$ saturates the inequality Eq. (43).

Let us now determine a complementary relation. We will find the minimum for separable states for the left-hand side of Eq. (43). We need to know that for separable states

$$[\Delta(\sigma_x \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_x)]^2 + [\Delta(\sigma_y \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_y)]^2 \geq 2 \quad (45)$$

holds, c. f. Eq. (35). This can be seen as follows [84, 85]. We need to know that for two-qubit quantum states

$$(\Delta\sigma_x)^2 + (\Delta\sigma_y)^2 \geq 2 =: L_{xy} \quad (46)$$

holds. For product states $|\Psi\rangle \otimes |\Phi\rangle$, the left-hand side of Eq. (45) equals the sum of single system variances, which can be bounded from below as

$$(\Delta\sigma_x)_{\Psi}^2 + (\Delta\sigma_y)_{\Psi}^2 + (\Delta\sigma_x)_{\Phi}^2 + (\Delta\sigma_y)_{\Phi}^2 \geq 2L_{xy} = 4. \quad (47)$$

The bound for separable states given in Eq. (18) is the same as the bound for product states, since the variance is a concave function of the state. The left-hand side of Eq. (45) is zero for the state

$$\frac{1}{\sqrt{2}}(|01\rangle + |10\rangle). \quad (48)$$

The product state $|11\rangle_x$ saturates the inequality given in Eq. (45) and for that state

$$\begin{aligned} \langle \sigma_x \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_x \rangle &= 0, \\ \langle \sigma_y \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_y \rangle &= 0 \end{aligned} \quad (49)$$

hold. Thus, for separable states

$$\langle (\sigma_x \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_x)^2 + (\sigma_y \otimes \mathbb{1} - \mathbb{1} \otimes \sigma_y)^2 \rangle \geq 2 \quad (50)$$

holds and it is even a tight inequality for separable states, c. f. Eq. (43).

IV. OPTIMIZATION OVER THE TWO-COPY SPACE

In this section, we first review the formalism that maps the optimization over the decompositions of the density matrix to an optimization of an operator expectation value over a bipartite symmetric separable quantum state with given marginals [55, 93]. Then, we show that the same results is obtained if the optimization is carried out over general separable quantum states rather than symmetric separable states.

We start from writing the variance of a pure state $|\Psi\rangle$ as an operator expectation value acting on two copies as

$$(\Delta H)_{\Psi}^2 = \text{Tr}(\Omega |\Psi\rangle\langle\Psi| \otimes |\Psi\rangle\langle\Psi|), \quad (51)$$

where we define the operator

$$\Omega = H^2 \otimes \mathbb{1} - H \otimes H. \quad (52)$$

Based on Eq. (51), the expression of the quantum Fisher information given in Eq. (13) can be rewritten as [60, 93]

$$\begin{aligned} \mathcal{F}_Q[\varrho, H] &= \\ 4 \min_{\{p_k, |\Psi_k\rangle\}} &\sum_k p_k \text{Tr}[\Omega (|\Psi_k\rangle\langle\Psi_k|)^{\otimes 2}], \\ \text{s. t. } \varrho &= \sum_k p_k |\Psi_k\rangle\langle\Psi_k|. \end{aligned} \quad (53)$$

The sum can be moved into the trace and we obtain

$$\begin{aligned} \mathcal{F}_Q[\varrho, H] &= \\ 4 \min_{\{p_k, |\Psi_k\rangle\}} &\text{Tr} \left\{ \Omega \left[\sum_k p_k (|\Psi_k\rangle\langle\Psi_k|)^{\otimes 2} \right] \right\}, \\ \text{s. t. } \varrho &= \sum_k p_k |\Psi_k\rangle\langle\Psi_k|. \end{aligned} \quad (54)$$

On the right-hand side in Eq. (54) in the square bracket, we can recognize a mixed state living in the two-copy space

$$\varrho_{12} = \sum_k p_k |\Psi_k\rangle\langle\Psi_k| \otimes |\Psi_k\rangle\langle\Psi_k|, \quad (55)$$

States given in Eq. (55) are symmetric separable states, thus $\varrho_{12} \in \mathcal{S}'$. We can formulate the optimization directly over states given in Eq. (55). Hence, further algebra leads to

$$\begin{aligned} \mathcal{F}_Q[\varrho, H] &= \min_{\varrho_{12}} 4\text{Tr}(\Omega \varrho_{12}), \\ \text{s. t. } \varrho_{12} &\in \mathcal{S}', \\ \text{Tr}_2(\varrho_{12}) &= \varrho. \end{aligned} \quad (56)$$

Due to an optimization over symmetric separable states, $\text{Tr}_1(\varrho_{12}) = \varrho$ is fulfilled without adding it as an explicit constraint.

Equation (56) contains an optimization over symmetric separable states, which cannot be computed directly. However, we can consider an optimization over a larger set, the set of symmetric PPT states, which leads to an expression that can be obtained numerically using semidefinite programming [94]

$$\begin{aligned} \mathcal{F}_Q^{(\text{PPT})}[\varrho, H] &= \min_{\varrho_{12}} 4\text{Tr}(\Omega \varrho_{12}), \\ \text{s. t. } \varrho_{12} &\in \mathcal{P}', \\ \text{Tr}_2(\varrho_{12}) &= \varrho, \end{aligned} \quad (57)$$

In general, the relation

$$\mathcal{F}_Q^{(\text{PPT})}[\varrho, H] \leq \mathcal{F}_Q[\varrho, H] \quad (58)$$

holds, while for two qubits we have an equality, since for that system size $\mathcal{S}' = \mathcal{P}'$, as we discussed in Sec. III.

Let us now change the operator to be optimized, making it permutationally symmetric. Equation (56) can be rewritten as

$$\begin{aligned} \mathcal{F}_Q[\varrho, H] &= \min_{\varrho_{12}} 2\text{Tr}[(H \otimes \mathbb{1} - \mathbb{1} \otimes H)^2 \varrho_{12}], \\ \text{s. t. } \varrho_{12} &\in \mathcal{S}', \\ \text{Tr}_2(\varrho_{12}) &= \varrho. \end{aligned} \quad (59)$$

We will now show that the expression in Eq. (59) remains true if we change the set over which we have to optimize to the set of separable states.

Observation 1.—The quantum Fisher information can be obtained as an optimization over separable states as

$$\begin{aligned} \mathcal{F}_Q[\varrho, H] &= \min_{\varrho_{12}} 2\text{Tr}[(H \otimes \mathbb{1} - \mathbb{1} \otimes H)^2 \varrho_{12}], \\ \text{s. t. } \varrho_{12} &\in \mathcal{S}, \\ \text{Tr}_2(\varrho_{12}) &= \varrho, \\ \text{Tr}_1(\varrho_{12}) &= \varrho. \end{aligned} \quad (60)$$

Proof.—Using

$$\begin{aligned} & \text{Tr}[(H \otimes \mathbf{1} - \mathbf{1} \otimes H)^2 \varrho_{12}] \\ &= 2\text{Tr}(H^2 \varrho) - 2\text{Tr}[(H \otimes H)\varrho_{12}], \end{aligned} \quad (61)$$

the right-hand side of Eq. (60) can be reformulated as

$$\begin{aligned} & 4\text{Tr}(H^2 \varrho) - 4 \max_{\varrho_{12}} \text{Tr}[(H \otimes H)\varrho_{12}], \\ & \text{s. t. } \varrho_{12} \in \mathcal{S}, \\ & \text{Tr}_2(\varrho_{12}) = \varrho, \\ & \text{Tr}_1(\varrho_{12}) = \varrho. \end{aligned} \quad (62)$$

Due to the permutational invariance of $H \otimes H$ and that both marginals must be equal to ϱ , if a separable state given in Eq. (18) maximizes the correlation $\langle H \otimes H \rangle$, then the separable state

$$\sum_k p_k |\Phi_k\rangle\langle\Phi_k| \otimes |\Psi_k\rangle\langle\Psi_k|. \quad (63)$$

also maximizes the correlation. Then, the mixture of the above two separable states also maximize the correlation with the given marginals, which can always be written as

$$\varrho_{12} = \sum_{k=1}^M \tilde{p}_k |\tilde{\Psi}_k\rangle\langle\tilde{\Psi}_k| \otimes |\tilde{\Psi}_{\pi_k}\rangle\langle\tilde{\Psi}_{\pi_k}|, \quad (64)$$

where π_k for $k = 1, 2, \dots, M$ is a permutation of $1, 2, \dots, M$ for some M . Based on our arguments, it is sufficient to look for the separable state that maximizes $\langle H \otimes H \rangle$ in the form Eq. (64), and $\tilde{p}_k \geq 0$, $\sum_k \tilde{p}_k = 1$. Then, the correlation can be given as

$$\text{Tr}[(H \otimes H)\varrho_{12}] = \sum_{k=1}^M \tilde{p}_k h_k h_{\pi_k}, \quad (65)$$

where we define the expectation values of H on $|\tilde{\Psi}_k\rangle$ as

$$h_k = \text{Tr}(H|\tilde{\Psi}_k\rangle\langle\tilde{\Psi}_k|). \quad (66)$$

Based on the Cauchy-Schwarz inequality we have

$$\begin{aligned} \sum_k \tilde{p}_k h_k h_{\pi_k} &\leq \sqrt{\sum_k \tilde{p}_k h_k^2 \sum_k \tilde{p}_k h_{\pi_k}^2} \\ &\leq \max \left(\sum_k \tilde{p}_k h_k^2, \sum_k \tilde{p}_k h_{\pi_k}^2 \right). \end{aligned} \quad (67)$$

Thus, when we maximize $\text{Tr}[(H \otimes H)\varrho_{12}]$ over separable states with the constraints for the marginals, the maximum is taken by a symmetric separable state given in Eq. (55). In this case both inequalities are saturated. ■

Based on Eq. (15) and the previous arguments, we can obtain the variance also as a result of an optimization over a two-copy space.

Observation 2.—The variance can also be obtained as an optimization over separable states as

$$\begin{aligned} (\Delta H)^2 &= \max_{\varrho_{12}} \frac{1}{2} \text{Tr}[(H \otimes \mathbf{1} - \mathbf{1} \otimes H)^2 \varrho_{12}], \\ & \text{s. t. } \varrho_{12} \in \mathcal{S}, \\ & \text{Tr}_2(\varrho_{12}) = \varrho, \\ & \text{Tr}_1(\varrho_{12}) = \varrho. \end{aligned} \quad (68)$$

V. QUANTUM WASSERSTEIN DISTANCE BASED ON A SEPARABLE COUPLING

Next, we will connect our result in Observation 1 to results available in the literature mentioned in the introduction. We will consider the quantum Wasserstein distance such that the optimization takes place over separable states rather than over general bipartite quantum states. We will consider such modifications of $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$ and $D_{\text{DPT,sep}}(\varrho, \sigma)^2$, and examine their properties.

Our first finding concerning the GMPC distance is the following.

Definition 3.—Modifying the definition in Eq. (5) we can define a new type of GMPC distance such that we restrict the optimization over separable states as

$$\begin{aligned} & D_{\text{GMPC,sep}}(\varrho, \sigma)^2 \\ &= \frac{1}{2} \min_{\varrho_{12}} \sum_{n=1}^N \text{Tr}[(H_n \otimes \mathbf{1} - \mathbf{1} \otimes H_n)^2 \varrho_{12}], \\ & \text{s. t. } \varrho_{12} \in \mathcal{S}, \\ & \text{Tr}_1(\varrho_{12}) = \varrho, \\ & \text{Tr}_2(\varrho_{12}) = \sigma. \end{aligned} \quad (69)$$

We can see immediately two relevant properties of the newly defined distance. Observation 1 showed that for $N = 1$ for a given H_1

$$D_{\text{GMPC,sep}}(\varrho, \varrho)^2 = \frac{1}{4} \mathcal{F}_Q[\varrho, H_1] \quad (70)$$

holds. Moreover, based on Eqs. (5) and (69), we can immediately see that

$$D_{\text{GMPC}}(\varrho, \sigma)^2 \leq D_{\text{GMPC,sep}}(\varrho, \sigma)^2, \quad (71)$$

since on the left-hand side of Eq. (71) there is a minimization over a larger set of quantum states than on the right-hand side.

Let us now define another distance based on an optimization over separable states.

Definition 4.—Based on Eq. (1), we can also define

$$\begin{aligned} & D_{\text{DPT,sep}}(\varrho, \sigma)^2 = \frac{1}{2} \min_{\varrho_{12}} \sum_{n=1}^N \text{Tr}[(H_n^T \otimes \mathbf{1} - \mathbf{1} \otimes H_n)^2 \varrho_{12}], \\ & \text{s. t. } \varrho_{12} \in \mathcal{S}, \\ & \text{Tr}_2(\varrho_{12}) = \varrho^T, \\ & \text{Tr}_1(\varrho_{12}) = \sigma. \end{aligned} \quad (72)$$

As an important property of $D_{\text{DPT,sep}}(\varrho, \sigma)^2$, we can see that

$$D_{\text{DPT}}(\varrho, \sigma)^2 \leq D_{\text{DPT,sep}}(\varrho, \sigma)^2 \quad (73)$$

holds, since on the left-hand side of Eq. (73) there is a minimization over a larger set of quantum states than on the right-hand side.

Interestingly, the two different types of distance, defined with the transpose and without it, respectively, are equal to each other.

Observation 3.—The two types of quantum Wasserstein distance are equal to each other

$$D_{\text{DPT,sep}}(\varrho, \sigma)^2 = D_{\text{GMPC,sep}}(\varrho, \sigma)^2. \quad (74)$$

Proof.—Using the fact that for two matrices X and Y act on the bipartite system

$$\text{Tr}(XY) = \text{Tr}(X^{T_k} Y^{T_k}) \quad (75)$$

holds for $k = 1, 2$, from Eq. (69) we arrive at

$$\begin{aligned} D_{\text{GMPC,sep}}(\varrho, \sigma)^2 &= \frac{1}{2} \min_{\varrho_{12}} \sum_{n=1}^N \text{Tr}[(H_n^T \otimes \mathbf{1} - \mathbf{1} \otimes H_n)^2 \varrho_{12}^{T_1}], \\ &\text{s. t. } \varrho_{12} \in \mathcal{S}, \\ &\quad \text{Tr}_2(\varrho_{12}) = \varrho, \\ &\quad \text{Tr}_1(\varrho_{12}) = \sigma. \end{aligned} \quad (76)$$

Then, we replace ϱ_{12} by $\varrho_{12}^{T_1}$, which leaves the result of the minimization unchanged. Then, it follows that

$$\begin{aligned} D_{\text{GMPC,sep}}(\varrho, \sigma)^2 &= \frac{1}{2} \min_{\varrho_{12}^{T_1}} \sum_{n=1}^N \text{Tr}[(H_n^T \otimes \mathbf{1} - \mathbf{1} \otimes H_n)^2 \varrho_{12}^{T_1}], \\ &\text{s. t. } \varrho_{12}^{T_1} \in \mathcal{S}, \\ &\quad \text{Tr}_2(\varrho_{12}^{T_1}) = \varrho^T, \\ &\quad \text{Tr}_1(\varrho_{12}^{T_1}) = \sigma. \end{aligned} \quad (77)$$

Finally, we arrive at Eq. (72) by noticing that

$$\varrho_{12}^{T_1} \in \mathcal{S} \quad (78)$$

holds if and only if

$$\varrho_{12} \in \mathcal{S} \quad (79)$$

holds. \blacksquare

It is instructive to obtain a quantum state that maximizes $D_{\text{DPT,sep}}(\varrho, \varrho)^2$ and $D_{\text{DPT}}(\varrho, \varrho)^2$ for $N = 1$ for a given H_1 as follows. We know that

$$I_\varrho(H_1) \leq \mathcal{F}_Q[\varrho, H_1]/4 \leq (\Delta H_1)_\varrho^2 \leq \frac{1}{4}(h_{\max} - h_{\min})^2 \quad (80)$$

holds. The first two inequalities are based on Eq. (10). The third one can be obtained as follows. Simple algebra shows that for the state

$$|\Psi_{\text{opt}}\rangle = \frac{1}{\sqrt{2}}(|h_{\min}\rangle + |h_{\max}\rangle), \quad (81)$$

where $|h_{\min}\rangle$ ($|h_{\max}\rangle$) is the eigenstate of H_1 with the minimal (maximal) eigenvalue of H_1 , the variance $(\Delta H_1)^2$ is maximal, and all inequalities of Eq. (80) are saturated. Hence, the maximal self-distance is achieved by $|\Psi_{\text{opt}}\rangle$

$$\begin{aligned} D_{\text{DPT,sep}}(|\Psi_{\text{opt}}\rangle\langle\Psi_{\text{opt}}|, |\Psi_{\text{opt}}\rangle\langle\Psi_{\text{opt}}|)^2 &= D_{\text{DPT}}(|\Psi_{\text{opt}}\rangle\langle\Psi_{\text{opt}}|, |\Psi_{\text{opt}}\rangle\langle\Psi_{\text{opt}}|)^2 \\ &= \frac{1}{4}(h_{\max} - h_{\min})^2. \end{aligned} \quad (82)$$




FIG. 1. Geometric representation of the quantum Wasserstein distance between a pure state ϱ and a mixed state σ given in Example 1 for $N = 1$ with operator H_1 . In Eq. (84), the Wasserstein distance square is 1/2 times the sum of three terms, corresponding to the two uncertainties and $(\langle H_1 \rangle_\varrho - \langle H_1 \rangle_\sigma)^2$. Thus, not only the AB distance matters, but the variances computed for the two states. The relevant points are $A(\langle H_1 \rangle_\varrho, 0, 0)$, $B(\langle H_1 \rangle_\sigma, 0, 0)$, $A'(\langle H_1 \rangle_\varrho, \sqrt{(\Delta H_1)_\varrho^2}, 0)$ and $B'(\langle H_1 \rangle_\sigma, 0, \sqrt{(\Delta H_1)_\sigma^2})$. The quantum Wasserstein distance equals 1/2 times the usual Euclidean distance between A' and B' .

Let us calculate $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$ and the other types of quantum Wasserstein distance for some concrete examples.

Example 1.—Let us consider the case when $\varrho = |\Psi\rangle\langle\Psi|$ is a pure state of any dimension and σ is an arbitrary density matrix of the same dimension. Then, when computing the various types of Wasserstein distance between ϱ and σ , the state ϱ_{12} in the optimization is constrained to be the tensor product of the two density matrices. Hence, for the distance from a pure state,

$$\begin{aligned} D_{\text{DPT}}(|\Psi\rangle\langle\Psi|, \sigma)^2 &= D_{\text{DPT,sep}}(|\Psi\rangle\langle\Psi|, \sigma)^2 \\ &= D_{\text{GMPC,sep}}(|\Psi\rangle\langle\Psi|, \sigma)^2 \\ &= D_{\text{GMPC}}(|\Psi\rangle\langle\Psi|, \sigma)^2 \\ &= \sum_{n=1}^N \frac{\langle H_n^2 \rangle_\varrho + \langle H_n^2 \rangle_\sigma - \langle H_n \rangle_\varrho \langle H_n \rangle_\sigma}{2} \end{aligned} \quad (83)$$

holds. The last expression in Eq. (83) can be written also as

$$\frac{1}{2} \sum_{n=1}^N [(\Delta H_n)_\varrho^2 + (\Delta H_n)_\sigma^2 + (\langle H_n \rangle_\varrho - \langle H_n \rangle_\sigma)^2], \quad (84)$$

see Fig. 1.

Example 2.—Let us consider the single-qubit states

$$\varrho = \sigma = \frac{1}{2}\mathbf{1}. \quad (85)$$

Let us take $N = 1$ and $H_1 = \sigma_z$, where the Pauli spin matrix is defined as $\sigma_z = \text{diag}(+1, -1)$. Then, when comput-

ing $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$, for the optimum we have the separable state

$$\varrho_{12} = \frac{1}{2}|00\rangle\langle 00| + \frac{1}{2}|11\rangle\langle 11|, \quad (86)$$

and for the self-distance

$$D_{\text{GMPC,sep}}\left(\frac{1}{2}\mathbb{1}, \frac{1}{2}\mathbb{1}\right)^2 = 0 \quad (87)$$

holds. If H_1 is a different operator then the state ϱ_{12} corresponding to the minimum will be different. It can be obtained from the state given in Eq. (86) with local unitaries. For instance, for $H_1 = \sigma_x$, where the σ_x Pauli spin matrix is defined as

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad (88)$$

the optimal bipartite separable state is

$$\varrho_{12} = \frac{1}{2}|00\rangle\langle 00|_x + \frac{1}{2}|11\rangle\langle 11|_x, \quad (89)$$

where $|\cdot\rangle_x$ is a state given in the x -basis. That is, for a qubit, the basis states in the x -basis are

$$\begin{aligned} |0\rangle_x &= \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle), \\ |1\rangle_x &= \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle). \end{aligned} \quad (90)$$

Example 3.—Let us consider the two-qubit states $\varrho = \sigma = D_p$, where the diagonal state is defined as

$$D_p = \begin{pmatrix} p & 0 \\ 0 & 1-p \end{pmatrix}. \quad (91)$$

Let us take $N = 1$ and $H_1 = \sigma_z$. When computing $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$, the optimal bipartite separable state is

$$\varrho_{12} = p|00\rangle\langle 00| + (1-p)|11\rangle\langle 11|. \quad (92)$$

The self-distance is zero for all p

$$D_{\text{GMPC,sep}}(D_p, D_p)^2 = 0. \quad (93)$$

Example 4.—Let us consider the two single-qubit mixed states

$$\varrho = \frac{1}{2}|1\rangle\langle 1| + \frac{1}{2} \cdot \frac{\mathbb{1}}{2}. \quad (94)$$

and

$$\sigma_\phi = e^{-i\frac{\sigma_y}{2}\phi} \varrho e^{+i\frac{\sigma_y}{2}\phi}, \quad (95)$$

where the σ_y Pauli spin matrix is given as

$$\sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad (96)$$

and ϕ is a real parameter. For $N = 1$, $H_1 = \sigma_z$ we plotted $D_{\text{DPT,sep}}(\varrho, \sigma_\phi)^2$ and $D_{\text{DPT}}(\varrho, \sigma_\phi)^2$ in Fig. 2. The details




FIG. 2. (solid) $D_{\text{DPT,sep}}(\varrho, \sigma_\phi)^2$ and (dashed) $D_{\text{DPT}}(\varrho, \sigma_\phi)^2$ for the states given in Eqs. (94) and (95). The two curves coincide with each other on the right-hand side of the vertical dotted line, where $\phi \geq \phi_0$ and ϕ_0 is given in Eq. (102).

of the numerical calculations are in Appendix A. For $\phi = 0$, $\sigma_\phi = \varrho$, hence in this case the two types of distance equals the corresponding types of self-distance of ϱ . That is, based on Eq. (3), we have

$$D_{\text{DPT}}(\varrho, \varrho)^2 = I_\varrho(H_1) = 1 - \frac{\sqrt{3}}{2} \approx 0.1340, \quad (97)$$

and based on Eq. (70) and taking into account Observation 3, we have

$$D_{\text{DPT,sep}}(\varrho, \varrho)^2 = \frac{\mathcal{F}_Q[\varrho, H_1]}{4} = \frac{1}{4}. \quad (98)$$

Here, we used the formula giving the quantum Fisher information with the variance for pure states mixed with white noise as [69, 95–97]

$$\mathcal{F}_Q \left[p|\Psi\rangle\langle\Psi| + (1-p)\frac{\mathbb{1}}{d}, H \right] = \frac{p^2}{p + 2(1-p)d^{-1}} 4(\Delta H)_\Psi^2, \quad (99)$$

where d is the dimension of the system. For $\phi = \pi/2$,

$$\sigma_{\pi/2} = \frac{1}{2}|1\rangle\langle 1| + \frac{1}{2} \cdot \frac{\mathbb{1}}{2}. \quad (100)$$

Numerics show that

$$D_{\text{DPT,sep}}(\varrho, \sigma_\phi)^2 = D_{\text{DPT}}(\varrho, \sigma_\phi)^2 \quad (101)$$

for

$$\phi \geq \phi_0 \approx 0.2946\pi, \quad (102)$$

while for smaller ϕ an entangled ϱ_{12} is cheaper than a separable one.

We can relate $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$ to the quantum Fisher information.

Observation 4.—For the modified GMPC distance defined in Eq. (69) the inequality

$$D_{\text{GMPC,sep}}(\varrho, \sigma)^2 \geq \frac{1}{8} \sum_{n=1}^N \mathcal{F}_Q[\varrho, H_n] + \frac{1}{8} \sum_{n=1}^N \mathcal{F}_Q[\sigma, H_n] \quad (103)$$

holds, and for $\varrho = \sigma$ and for $N = 1$ we have equality in Eq. (103).

Proof.—We can rewrite the optimization problem in Eq. (69) for separable states as

$$\begin{aligned} D_{\text{GMPC,sep}}(\varrho, \sigma)^2 &= \frac{1}{2} \min_{\{p_k, |\Psi_k\rangle, |\Phi_k\rangle\}} \sum_{n=1}^N \sum_k p_k \left[(\Delta H_n)_{\Psi_k}^2 + (\Delta H_n)_{\Phi_k}^2 \right. \\ &\quad \left. + (\langle H_n \rangle_{\Psi_k} - \langle H_n \rangle_{\Phi_k})^2 \right] \\ &\geq \frac{1}{2} \left[\min_{\{p_k, |\Psi_k\rangle\}} \sum_{n=1}^N \sum_k p_k (\Delta H_n)_{\Psi_k}^2 \right. \\ &\quad \left. + \min_{\{q_k, |\Phi_k\rangle\}} \sum_{n=1}^N \sum_k q_k (\Delta H_n)_{\Phi_k}^2 \right] \\ &\geq \frac{1}{2} \left[\sum_{n=1}^N \min_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k (\Delta H_n)_{\Psi_k}^2 \right. \\ &\quad \left. + \sum_{n=1}^N \min_{\{q_k, |\Phi_k\rangle\}} \sum_k q_k (\Delta H_n)_{\Phi_k}^2 \right], \quad (104) \end{aligned}$$

where for the first optimization the decomposition of ϱ_{12} is given as Eq. (18), and we have the conditions

$$\text{Tr}_2(\varrho_{12}) = \varrho, \quad \text{Tr}_1(\varrho_{12}) = \sigma. \quad (105)$$

For the second optimization, the condition for ϱ is given in Eq. (14). For the third optimization, the condition is

$$\sigma = \sum_k q_k |\Phi_k\rangle \langle \Phi_k|, \quad (106)$$

where for the probabilities $q_k \geq 0$ and $\sum_k q_k = 1$ hold. The fourth and fifth optimization are similar to the second and the third one, however, the order of the sum and the minimization is exchanged. From Eq. (104), the statement follows using the formula giving the quantum Fisher information with the convex roof of the variance in Eq. (13). ■

In Eq. (104), in the second and third lines we can see that, when computing $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$, the quantity to be minimized is the a weighted sum containing the variances of H_n for $|\Psi_k\rangle$ and $|\Phi_k\rangle$, and the expression $(\langle H_n \rangle_{\Psi_k} - \langle H_n \rangle_{\Phi_k})^2$. The two variances are zero if $|\Psi_k\rangle$ and $|\Phi_k\rangle$ are the eigenstates of H . Since ϱ_{12} must obey the constraints given in Eq. (105), in this case, ϱ , σ and H must have common eigenstates. The term $\sum_k (\langle H_n \rangle_{\Psi_k} - \langle H_n \rangle_{\Phi_k})^2$ is zero if

$$\langle H_n \rangle_{\Psi_k} = \langle H_n \rangle_{\Phi_k} \quad (107)$$

for all k .

Let us see now the consequences of the above observations for the self-distance. Unlike in the classical case, it can be nonzero. Based on these, for the self-distance for $N = 1$ for a given H_1

$$D_{\text{GMPC,sep}}(\varrho, \varrho)^2 = 0 \quad (108)$$

holds, if and only if

$$[\varrho, H_1] = 0. \quad (109)$$

Let us look for similar observations for the distance between two different states. For this case, for $N = 1$ for a given H_1

$$D_{\text{GMPC,sep}}(\varrho, \sigma)^2 = 0 \quad (110)$$

holds, if

$$[\varrho, H_1] = [\sigma, H_1] = [\varrho, \sigma] = 0. \quad (111)$$

is satisfied and if Eq. (107) holds for all k . In this case, $|\Psi_k\rangle$ are the eigenvectors of ϱ and of H_1 . $|\Phi_k\rangle$ are the eigenvectors of σ and of H_1 . If ϱ, σ and H_1 have a non-degenerate spectrum then $\{|\Psi_k\rangle\}_{k=1}^d$ and $\{|\Phi_k\rangle\}_{k=1}^d$ are the same, apart from a reordering of the labels, where d is the dimension of the subsystems.

Finally, from the first inequality in Eq. (104), it follows that for any ϱ and σ

$$\begin{aligned} D_{\text{GMPC,sep}}(\varrho, \sigma)^2 &\geq \\ &\frac{1}{2} [D_{\text{GMPC,sep}}(\varrho, \varrho)^2 + D_{\text{GMPC,sep}}(\sigma, \sigma)^2] \end{aligned} \quad (112)$$

holds, since the self-distances can be obtained as

$$\begin{aligned} D_{\text{GMPC,sep}}(\varrho, \varrho)^2 &= \min_{\{p_k, |\Psi_k\rangle\}} \sum_{n=1}^N \sum_k p_k (\Delta H_n)_{\Psi_k}^2, \\ D_{\text{GMPC,sep}}(\sigma, \sigma)^2 &= \min_{\{q_k, |\Phi_k\rangle\}} \sum_{n=1}^N \sum_k q_k (\Delta H_n)_{\Phi_k}^2. \end{aligned} \quad (113)$$

In the formulas giving the self-distances in Eq. (113), for the first optimization, the condition for ϱ is given in Eq. (14). For the second optimization, the condition for σ is given in Eq. (106). Based on Observation 3, an analogous relation holds for $D_{\text{DPT,sep}}(\varrho, \sigma)^2$.

For local dimension $d > 2$, the optimization over separable states is difficult to carry out numerically. For this reason, it is reasonable to define $D_{\text{PPT}}(\varrho, \sigma)^2$ and $D_{\text{GMPC,PPT}}(\varrho, \sigma)^2$ that need an optimization over PPT states rather than separable states.

VI. VARIANCE-LIKE QUANTITIES

When we compare the expression in Eq. (60) defining the quantum Fisher information and the other expression in

Eq. (68) defining the variance, we can see that the main difference is that the minimization is replaced by a maximization. We also showed a similar relation between the entanglement of formation given in Eq. (25), and the entanglement of assistance given in Eq. (27). Based on this observation, we can define variance-like quantities from the various forms of quantum Wasserstein distance by replacing the minimization by maximization.

Such a variance-like quantity can be interpreted in the framework of transport problems as follows. The quantum Wasserstein distance determines the smallest cost possible for the transport problem by a minimization. The variance-like quantities presented in this section determine the largest cost possible for the transport problem.

Let us now define the first variance-like quantity.

Definition 5.—From the GMPC distance with an optimization restricted over separable states given in Eq. (69), we obtain the following variance like quantity

$$\begin{aligned} V_{\text{GMPC,sep}}(\varrho, \sigma) &= \frac{1}{2} \max_{\varrho_{12}} \sum_{n=1}^N \text{Tr}[(H_n \otimes \mathbb{1} - \mathbb{1} \otimes H_n)^2 \varrho_{12}], \\ &\text{s. t. } \varrho_{12} \in \mathcal{S}, \\ &\quad \text{Tr}_2(\varrho_{12}) = \varrho, \\ &\quad \text{Tr}_1(\varrho_{12}) = \sigma. \end{aligned} \quad (114)$$

Clearly, the inequality

$$V_{\text{GMPC,sep}}(\varrho, \sigma) \geq D_{\text{GMPC,sep}}(\varrho, \sigma)^2 \quad (115)$$

holds, since since on the left-hand side of Eq. (115) we maximize over a set of quantum states while on the right-hand side we minimize over the same set.

We can define analogously $V_{\text{DPT,sep}}(\varrho, \sigma)$, $V_{\text{GMPC}}(\varrho, \sigma)$, and $V_{\text{DPT}}(\varrho, \sigma)$, modifying the definition of $D_{\text{DPT,sep}}(\varrho, \sigma)^2$, $D_{\text{GMPC}}(\varrho, \sigma)^2$, and $D_{\text{DPT}}(\varrho, \sigma)^2$, respectively. In all these cases, a relation analogous to the one in Eq. (115) can be obtained.

The value of $V_{\text{GMPC,sep}}(\varrho, \sigma)$ is related to the variance.

Observation 5.—The GMPC variance defined in Eq. (114) is bounded from below as

$$V_{\text{GMPC,sep}}(\varrho, \sigma) \geq \frac{1}{2} \sum_{n=1}^N (\Delta H_n)_\varrho^2 + (\Delta H_n)_\sigma^2. \quad (116)$$

Proof.—We can rewrite the expression to be computed for $V_{\text{GMPC,sep}}(\varrho, \sigma)$ as

$$\begin{aligned} &\frac{1}{2} \max_{\varrho_{12} \in \mathcal{S}} \sum_{n=1}^N \text{Tr}[(H_n \otimes \mathbb{1} - \mathbb{1} \otimes H_n)^2 \varrho_{12}] \\ &= \frac{1}{2} \sum_{n=1}^N (\Delta H_n)_\varrho^2 + (\Delta H_n)_\sigma^2 + (\langle H_n \rangle_\varrho - \langle H_n \rangle_\sigma)^2 \\ &\quad - \min_{\varrho_{12} \in \mathcal{S}} C_{\varrho_{12}}, \end{aligned} \quad (117)$$

where ϱ_{12} has marginals ϱ and σ , as given in Eq. (105), and we define

$$C_{\varrho_{12}} = \sum_{n=1}^N \langle H_n \otimes H_n \rangle_{\varrho_{12}} - \langle H_n \rangle_\varrho \langle H_n \rangle_\sigma. \quad (118)$$

Since the product state $\varrho \otimes \sigma$ is separable and fulfills the conditions on the marginals, it is clear that

$$\min_{\varrho_{12} \in \mathcal{S}} C_{\varrho_{12}} \leq C_{\varrho \otimes \sigma} = 0. \quad (119)$$

Substituting Eq. (119) into Eq. (117), using the fact $(\langle H_n \rangle_\varrho - \langle H_n \rangle_\sigma)^2 \geq 0$, we can prove the observation. ■

Let us see some concrete examples.

Example 5.—Interestingly, the quantity

$$\mathcal{V} := V_{\text{GMPC,sep}}(\varrho, \sigma) \quad (120)$$

can be larger or smaller than or equal to

$$\bar{\mathcal{V}} := \frac{1}{2} [V_{\text{GMPC,sep}}(\varrho, \varrho) + V_{\text{GMPC,sep}}(\sigma, \sigma)], \quad (121)$$

c. f. Eq. (112). We consider $N = 1$ and $H_1 = \sigma_z$. The three possibilities above are realized by the following states. For the state

$$\begin{aligned} \varrho &= |0\rangle\langle 0|, \\ \sigma &= \text{diag}(0.25, 0.75), \end{aligned} \quad (122)$$

we have $\mathcal{V} > \bar{\mathcal{V}}$. For the following state

$$\begin{aligned} \varrho &= |0\rangle\langle 0|, \\ \sigma &= \text{diag}(0.75, 0.25), \end{aligned} \quad (123)$$

we have $\mathcal{V} = \bar{\mathcal{V}}$. Finally, for the following state

$$\begin{aligned} \varrho &= \mathbb{1}/2, \\ \sigma &= \begin{pmatrix} 0.75 & 0.40 \\ 0.40 & 0.25 \end{pmatrix}, \end{aligned} \quad (124)$$

we have $\mathcal{V} < \bar{\mathcal{V}}$.

The two different types of variance-like quantities, defined with the transpose and without it, respectively, are equal to each other.

Observation 6.—The two types of quantum Wasserstein variance are equal to each other

$$V_{\text{DPT,sep}}(\varrho, \sigma) = V_{\text{GMPC,sep}}(\varrho, \sigma). \quad (125)$$

Proof.—The proof is analogous to that of Observation 3. ■

Let us calculate $V_{\text{DPT,sep}}(\varrho, \sigma)$ for some concrete examples.

Example 6.—Let us consider the case when $\varrho = |\Psi\rangle\langle\Psi|$ is a pure state of any dimension and σ is an arbitrary density matrix of the same dimension. Then, when computing the various types of quantum Wasserstein variance between ϱ and σ , the state ϱ_{12} in the optimization is constrained to be the

tensor product of the two density matrices. Hence, it follows that

$$\begin{aligned}
V_{\text{DPT}}(|\Psi\rangle\langle\Psi|, \sigma) &= V_{\text{DPT,sep}}(|\Psi\rangle\langle\Psi|, \sigma) \\
&= V_{\text{GMPC,sep}}(|\Psi\rangle\langle\Psi|, \sigma) \\
&= V_{\text{GMPC}}(|\Psi\rangle\langle\Psi|, \sigma) \\
&= \sum_{n=1}^N \frac{\langle H_n^2 \rangle_\varrho + \langle H_n^2 \rangle_\sigma}{2} - \langle H_n \rangle_\varrho \langle H_n \rangle_\sigma
\end{aligned} \tag{126}$$

holds, c. f. Eq. (83).

Example 7.—For $\varrho = \sigma = |\Psi\rangle\langle\Psi|$, and for $N = 1$ for a given H_1 we obtain

$$V_{\text{DPT}}(|\Psi\rangle\langle\Psi|, |\Psi\rangle\langle\Psi|) = (\Delta H_1)_\Psi^2, \tag{127}$$

which motivates further calling V quantum Wasserstein variance.

Example 8.—Let us consider the single-qubit states given in Eq. (85). Let us take $N = 1$ and $H_1 = \sigma_z$. Then, when computing $V_{\text{GMPC,sep}}(\varrho, \sigma)$, for the optimum we have the separable state

$$\varrho_{12} = \frac{1}{2}|01\rangle\langle 01| + \frac{1}{2}|10\rangle\langle 10|, \tag{128}$$

[c. f. Eq. (86)] and we have

$$V_{\text{GMPC,sep}}\left(\frac{1}{2}\mathbb{1}, \frac{1}{2}\mathbb{1}\right) = 2. \tag{129}$$

If H_1 is a different operator then the state ϱ_{12} corresponding to the maximum will be different. It can be obtained from the state given in Eq. (128) with local unitaries. For instance, for $H_1 = \sigma_x$, the optimal bipartite separable state is

$$\varrho_{12} = \frac{1}{2}|01\rangle\langle 01|_x + \frac{1}{2}|10\rangle\langle 10|_x, \tag{130}$$

where $|\cdot\rangle_x$ is a state given in the x -basis.

VII. QUANTUM WASSERSTEIN DISTANCE AND ENTANGLEMENT CRITERIA

Example 4 highlighted that in certain cases the minimum for separable states is larger than the minimum for general states. In this case, entanglement can help to decrease the quantum Wasserstein distance. In this section, we analyze the relation of the quantum Wasserstein distance and entanglement conditions on the optimal ϱ_{12} coupling.

First, we make the following simple observation.

Observation 7.—If

$$D_{\text{DPT,sep}}(\varrho, \sigma)^2 > D_{\text{DPT}}(\varrho, \sigma)^2 \tag{131}$$

holds then the optimal ϱ_{12} state for the given ϱ and σ , when computing $D_{\text{DPT}}(\varrho, \sigma)^2$, is entangled. The situation is analogous if

$$D_{\text{GMPC,sep}}(\varrho, \sigma)^2 > D_{\text{GMPC}}(\varrho, \sigma)^2. \tag{132}$$

Then the optimal ϱ_{12} state for the given ϱ and σ , when computing $D_{\text{GMPC}}(\varrho, \sigma)^2$, is entangled. Thus, we can even use the quantum Wasserstein distance as an entanglement criterion detecting entanglement in the optimal ϱ_{12} state.

We can even consider conditions with the self-distance. Based on Eqs. (3), (70), and (74), we find that for the $N = 1$ case

$$D_{\text{DPT,sep}}(\varrho, \varrho)^2 > D_{\text{DPT}}(\varrho, \varrho)^2 \tag{133}$$

is equivalent to a relation between the quantum Fisher information and the Wigner-Yanase skew information

$$\frac{F_Q[\varrho, H_1]}{4} > I_\varrho(H_1). \tag{134}$$

Thus, the condition in Eq. (134) implies that the optimal ϱ_{12} , when computing $D_{\text{DPT}}(\varrho, \varrho)^2$, is entangled.

Let us now use entanglement criteria to construct relations for the quantum Wasserstein distance, that can verify that the coupling ϱ_{12} is entangled. If the inequality given in Eq. (33) holds for separable states, so does the inequality

$$\sum_{n=1}^{d^2-1} \langle (G_k^T \otimes \mathbb{1} - \mathbb{1} \otimes G_k)^2 \rangle \geq 4(d-1), \tag{135}$$

since the left-hand side of Eq. (135) is never smaller than the left-hand side of Eq. (33). Any state that violates the inequality in Eq. (135) is entangled. It can be shown that Eq. (135) is a tight inequality for separable states as follows. Based on Eq. (32), we see that for pure product states of the form

$$(|\Psi\rangle\langle\Psi|)^T \otimes |\Psi\rangle\langle\Psi|, \tag{136}$$

for the second moments

$$\sum_{n=1}^{d^2-1} \langle (G_k^T \otimes \mathbb{1} - \mathbb{1} \otimes G_k)^2 \rangle = 4(d-1) \tag{137}$$

holds.

Next, we will define a quantum Wasserstein distance related to the entanglement condition in Eq. (135).

Observation 8.—Let us consider d -dimensional systems with

$$H_n = G_n \tag{138}$$

for $n = 1, 2, \dots, d^2 - 1$. If

$$D_{\text{DPT}}^{\{G_1, G_2, \dots, G_{d^2-1}\}}(\varrho, \sigma)^2 < 2(d-1) \tag{139}$$

holds, then the optimal ϱ_{12} is entangled. Here, for clarity, we give explicitly the observables used to define the distance in the superscript.

Clearly, since when calculating $D_{\text{DPT,sep}}(\varrho, \sigma)^2$, we optimize over separable states, we have

$$D_{\text{DPT,sep}}^{\{G_1, G_2, \dots, G_{d^2-1}\}}(\varrho, \sigma)^2 \geq 2(d-1). \tag{140}$$

Thus, independently from what is ϱ and σ , their distance $D_{\text{DPT}}^{\{G_1, G_2, \dots, G_{d^2-1}\}}(\varrho, \sigma)^2$ cannot be smaller than a bound. This is true even if $\varrho = \sigma$.

Let us now use another entanglement condition to construct relations for the quantum Wasserstein distance that can verify that the coupling is entangled. If the inequality given in Eq. (33) holds for separable states, so does the inequality

$$\begin{aligned} & \langle (j_x^T \otimes \mathbb{1} - \mathbb{1} \otimes j_x)^2 \rangle \\ & + \langle (j_y^T \otimes \mathbb{1} - \mathbb{1} \otimes j_y)^2 \rangle \\ & + \langle (j_z^T \otimes \mathbb{1} - \mathbb{1} \otimes j_z)^2 \rangle \geq 2j. \end{aligned} \quad (141)$$

It can be shown that Eq. (141) is a tight inequality for separable states, since the $|+j, +j\rangle$ state saturates it. Based on these, we can obtain the following bounds on the quantum Wasserstein distance.

Observation 9.—Let us choose the set of operators as

$$\{H_n\} = \{j_x, j_y, j_z\}. \quad (142)$$

Then, if the inequality

$$D_{\text{DPT}}^{\{j_x, j_y, j_z\}}(\varrho, \sigma)^2 < j \quad (143)$$

holds, then the optimal ϱ_{12} is entangled.

Clearly, since when calculating $D_{\text{DPT,sep}}(\varrho, \sigma)^2$, we optimize over separable states, we have

$$D_{\text{DPT,sep}}^{\{j_x, j_y, j_z\}}(\varrho, \sigma)^2 \geq j. \quad (144)$$

Thus, again, independently from what ϱ and σ are, their distance $D_{\text{DPT,sep}}^{\{j_x, j_y, j_z\}}(\varrho, \sigma)^2$ cannot be smaller than a bound. This is true even if $\varrho = \sigma$.

So far we studied the relation of the quantum Wasserstein distance to entanglement. Next, let us consider the relation of $V_{\text{DPT}}(\varrho, \sigma)$ and $V_{\text{GMPC}}(\varrho, \sigma)$ to entanglement.

Observation 10.—If

$$V_{\text{DPT,sep}}(\varrho, \sigma) < V_{\text{DPT}}(\varrho, \sigma) \quad (145)$$

holds then the optimal ϱ_{12} state for the given ϱ and σ , when computing $V_{\text{DPT}}(\varrho, \sigma)$, is entangled. The situation is analogous if

$$V_{\text{GMPC,sep}}(\varrho, \sigma) < V_{\text{GMPC}}(\varrho, \sigma). \quad (146)$$

Next, we will determine a set of H_n operators that can be used efficiently to detect entanglement with the quantum Wasserstein variance. For that, we need to know that for separable states the inequality given in Eq. (43) holds.

Observation 11.—Let us consider an example with $d = 2$ and $\{H_n\} = \{\sigma_x, \sigma_y\}$. If the condition

$$V_{\text{GMPC}}^{\{\sigma_x, \sigma_y\}}(\varrho, \sigma) > 3 \quad (147)$$

holds, then the optimal ϱ_{12} is entangled. Clearly, since when calculating $V_{\text{DPT,sep}}(\varrho, \sigma)$, we optimize over separable states, we have

$$V_{\text{GMPC,sep}}^{\{\sigma_x, \sigma_y\}}(\varrho, \sigma) \leq 3. \quad (148)$$

Let us see now a complementary relation for $D_{\text{GMPC}}(\varrho, \sigma)^2$ and $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$. They will use the same H_n operators that appear in Observation 11. We need to know that for separable states the inequality given in Eq. (50) holds.

Observation 12.—Let us consider $d = 2$ and $\{H_n\} = \{\sigma_x, \sigma_y\}$. If

$$D_{\text{GMPC}}^{\{\sigma_x, \sigma_y\}}(\varrho, \sigma)^2 < 1 \quad (149)$$

then the optimal ϱ_{12} is entangled. Clearly, since when calculating $D_{\text{DPT,sep}}(\varrho, \sigma)$, we optimize over separable states, we have

$$D_{\text{GMPC,sep}}^{\{\sigma_x, \sigma_y\}}(\varrho, \sigma)^2 \geq 1. \quad (150)$$

Statements analogous to those of Observation 11 and Observation 12 can be formulated, with identical bounds, for $V_{\text{DPT}}(\varrho, \sigma)^2$, $V_{\text{DPT,sep}}(\varrho, \sigma)^2$, $D_{\text{DPT}}(\varrho, \sigma)^2$, and $D_{\text{DPT,sep}}(\varrho, \sigma)^2$.

VIII. OPTIMIZATION OF THE VARIANCE OVER THE TWO-COPY SPACE

In this section, we examine the quantity that we obtain after replacing the second moment by a variance in the optimization in the definition of $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$ given in Eq. (69) and in the definition of $V_{\text{GMPC,sep}}(\varrho, \sigma)$ in Eq. (114). Analogous ideas work also for the other types of quantum Wasserstein distance and quantum Wasserstein variance defined before. We will show that such quantities have interesting properties.

Definition 6.—After replacing the second moment by a variance in the optimization in the definition of $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$ given in Eq. (69), we define

$$\begin{aligned} & \tilde{D}_{\text{GMPC,sep}}(\varrho, \sigma)^2 \\ & = \frac{1}{2} \min_{\varrho_{12}} \sum_{n=1}^N [\Delta(H_n \otimes \mathbb{1} - \mathbb{1} \otimes H_n)]_{\varrho_{12}}^2, \\ & \text{s. t. } \varrho_{12} \in \mathcal{S}, \\ & \quad \text{Tr}_2(\varrho_{12}) = \varrho, \\ & \quad \text{Tr}_1(\varrho_{12}) = \sigma. \end{aligned} \quad (151)$$

Let us see some properties of the quantity we have just introduced. In general, for mixed states,

$$\begin{aligned} \tilde{D}_{\text{GMPC,sep}}(\varrho, \sigma)^2 & = D_{\text{GMPC,sep}}(\varrho, \sigma)^2 \\ & - \frac{1}{2} \sum_n \langle (H_n)_\varrho - (H_n)_\sigma \rangle^2 \end{aligned} \quad (152)$$

holds, hence clearly

$$\tilde{D}_{\text{GMPC,sep}}(\varrho, \sigma)^2 \leq D_{\text{GMPC,sep}}(\varrho, \sigma)^2. \quad (153)$$

Due to the relation in Eq. (152), for the self-distance we have

$$\tilde{D}_{\text{GMPC,sep}}(\varrho, \varrho)^2 = D_{\text{GMPC,sep}}(\varrho, \varrho)^2, \quad (154)$$

where $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$ is given in Eq. (69).

We can write the expression to be optimized as

$$\begin{aligned} & \sum_{n=1}^N [\Delta(H_n \otimes \mathbb{1} - \mathbb{1} \otimes H_n)]_{\varrho_{12}}^2 \\ &= \sum_{n=1}^N [(\Delta H_n)_{\varrho}^2 + (\Delta H_n)_{\sigma}^2] - 2C_{\varrho_{12}}, \end{aligned} \quad (155)$$

where ϱ_{12} has marginals ϱ and σ , as given in Eq. (105), and we define $C_{\varrho_{12}}$ as in Eq. (118). For pure states, $\varrho = |\Psi\rangle\langle\Psi|$ and $\sigma = |\Phi\rangle\langle\Phi|$, we have $C = 0$, and hence

$$\begin{aligned} & \tilde{D}_{\text{GMPC,sep}}(|\Psi\rangle\langle\Psi|, |\Phi\rangle\langle\Phi|)^2 \\ &= \frac{1}{2} \left[\tilde{D}_{\text{GMPC,sep}}(|\Psi\rangle\langle\Psi|, |\Psi\rangle\langle\Psi|)^2 \right. \\ & \quad \left. + \tilde{D}_{\text{GMPC,sep}}(|\Phi\rangle\langle\Phi|, |\Phi\rangle\langle\Phi|)^2 \right]. \end{aligned} \quad (156)$$

We can present lower bounds on the distance.

Observation 13.—The modified GMPC distance defined in Eq. (151) is bounded from below as

$$\tilde{D}_{\text{GMPC,sep}}(\varrho, \sigma)^2 \geq \frac{1}{8} \sum_{n=1}^N \mathcal{F}_Q[\varrho, H_n] + \frac{1}{8} \sum_{n=1}^N \mathcal{F}_Q[\sigma, H_n], \quad (157)$$

while for $\varrho = \sigma$ we have equality for $N = 1$ in Eq. (157).

Proof.—We can rewrite the optimization problem in Eq. (151) based on Eqs. (104) and (152) as

$$\begin{aligned} & \tilde{D}_{\text{GMPC,sep}}(\varrho, \sigma)^2 \\ &= \frac{1}{2} \min_{\{p_k, |\Psi_k\rangle, |\Phi_k\rangle\}} \sum_{n=1}^N \sum_k p_k \left[(\Delta H_n)_{\Psi_k}^2 + (\Delta H_n)_{\Phi_k}^2 \right. \\ & \quad \left. + (\langle H_n \rangle_{\Psi_k} - \langle H_n \rangle_{\Phi_k})^2 - (\langle H_n \rangle_{\varrho} - \langle H_n \rangle_{\sigma})^2 \right] \\ &\geq \frac{1}{2} \left[\min_{\{p_k, |\Psi_k\rangle\}} \sum_{n=1}^N \sum_k p_k (\Delta H_n)_{\Psi_k}^2 \right. \\ & \quad \left. + \min_{\{q_k, |\Phi_k\rangle\}} \sum_{n=1}^N \sum_k q_k (\Delta H_n)_{\Phi_k}^2 \right], \\ &\geq \frac{1}{2} \left[\sum_{n=1}^N \min_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k (\Delta H_n)_{\Psi_k}^2 \right. \\ & \quad \left. + \sum_{n=1}^N \min_{\{q_k, |\Phi_k\rangle\}} \sum_k q_k (\Delta H_n)_{\Phi_k}^2 \right], \end{aligned} \quad (158)$$

where we used in the first inequality that for real x_k

$$\sum_k p_k x_k^2 \geq \left(\sum_k p_k x_k \right)^2 \quad (159)$$

holds due to the fact that $f(x) = x^2$ is convex. In Eq. (158), in the first optimization the decomposition of ϱ_{12} is given as Eq. (18), and we have the conditions given in Eq. (105). For the second optimization, the condition is Eq. (14). For the

third optimization, the condition is given in Eq. (106), where for the probabilities $q_k \geq 0$ and $\sum_k q_k = 1$ hold. From Eq. (158), the statement follows using Eq. (13), which shows that the quantum Fisher information is the convex roof of the variance times four. The fourth and fifth optimization are similar to second and the third one, however, the order of the sum and the minimization is exchanged. From Eq. (158), the statement follows using the formula that obtains the quantum Fisher information with a convex roof of the variance given in Eq. (13). ■

We can define another variance-like quantity.

Definition 7.—Analogously, we can define the quantity that we obtain after replacing the second moment by a variance in the optimization in the definition of $V_{\text{GMPC,sep}}(\varrho, \sigma)^2$ in Eq. (114), as

$$\begin{aligned} & \tilde{V}_{\text{GMPC,sep}}(\varrho, \sigma) \\ &= \frac{1}{2} \max_{\varrho_{12}} \sum_{n=1}^N [\Delta(H_n \otimes \mathbb{1} - \mathbb{1} \otimes H_n)]_{\varrho_{12}}^2, \\ & \quad \text{s. t. } \varrho_{12} \in \mathcal{S}, \\ & \quad \text{Tr}_2(\varrho_{12}) = \varrho, \\ & \quad \text{Tr}_1(\varrho_{12}) = \sigma. \end{aligned} \quad (160)$$

Let us see now some properties of $\tilde{V}_{\text{GMPC,sep}}(\varrho, \sigma)$. In general,

$$\begin{aligned} & \tilde{V}_{\text{GMPC,sep}}(\varrho, \sigma) = V_{\text{GMPC,sep}}(\varrho, \sigma) \\ & \quad - \frac{1}{2} \sum_n (\langle H_n \rangle_{\varrho} - \langle H_n \rangle_{\sigma})^2 \end{aligned} \quad (161)$$

holds, hence clearly

$$\tilde{V}_{\text{GMPC,sep}}(\varrho, \sigma) \leq V_{\text{GMPC,sep}}(\varrho, \sigma). \quad (162)$$

Analogously to Eq. (156), for pure states

$$\begin{aligned} & \tilde{V}_{\text{GMPC,sep}}(|\Psi\rangle\langle\Psi|, |\Phi\rangle\langle\Phi|) \\ &= \frac{1}{2} \left[\tilde{V}_{\text{GMPC,sep}}(|\Psi\rangle\langle\Psi|, |\Psi\rangle\langle\Psi|) \right. \\ & \quad \left. + \tilde{V}_{\text{GMPC,sep}}(|\Phi\rangle\langle\Phi|, |\Phi\rangle\langle\Phi|) \right] \end{aligned} \quad (163)$$

holds.

Example 9.—Let us consider the case when $\varrho = |\Psi\rangle\langle\Psi|$ is a pure state of any dimension and σ is an arbitrary density matrix of the same dimension. Then, when computing the various types of quantum Wasserstein variance between ϱ and σ , the state ϱ_{12} in the optimization is constrained to be the tensor product of the two density matrices. Hence, based on Eqs. (126) and (161), it follows that

$$\tilde{V}_{\text{GMPC,sep}}(|\Psi\rangle\langle\Psi|, \sigma) = \frac{1}{2} \sum_{n=1}^N (\Delta H_n)_{\Psi}^2 + (\Delta H_n)_{\sigma}^2 \quad (164)$$

holds, c. f. Observation 5.

Finally, one can define similarly $\tilde{D}_{\text{DPT,sep}}(\varrho, \sigma)^2$, $\tilde{D}_{\text{GMPC}}(\varrho, \sigma)^2$, $\tilde{D}_{\text{DPT}}(\varrho, \sigma)^2$, $\tilde{V}_{\text{DPT,sep}}(\varrho, \sigma)$, $\tilde{V}_{\text{GMPC}}(\varrho, \sigma)$, and $\tilde{V}_{\text{DPT}}(\varrho, \sigma)$.

IX. EXTENSION TO FAMILIES OF THE QUANTUM FISHER INFORMATION

In this section, we will extend our statements about the quantum Fisher information to various types of generalized quantum Fisher information.

The basic idea of Refs. [98, 99] is that for each standard matrix monotone function $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, a generalized variance and a corresponding quantum Fisher information are defined. The notion standard means that f must satisfy

$$f(1) = 1, \quad (165a)$$

$$f(t) = tf(t^{-1}). \quad (165b)$$

For a review on generalized variances, different types of quantum Fisher information and covariances see Ref. [100]. Moreover, it is also useful to define the mean based on f as

$$m_f(a, b) = af\left(\frac{b}{a}\right) \quad (166)$$

and use it instead of f . The normalization condition given in Eq. (165a) corresponds to the condition

$$m_f(a, a) = a \quad (167)$$

for the means. The requirement given in Eq. (165b) corresponds to

$$m_f(a, b) = m_f(b, a). \quad (168)$$

A list of quantum Fisher information generated by various well-known means $m_f(a, b)$ can be found in Refs. [98, 99].

Using the normalization suggested by Ref. [60], we arrive at a family of generalized quantum Fisher information $\mathcal{F}_Q^f[\varrho, H]$ such that (i) for pure states, we have

$$\mathcal{F}_Q^f[|\Psi\rangle, H] = 4(\Delta H)_\Psi^2. \quad (169)$$

(ii) For mixed states, $\mathcal{F}_Q^f[\varrho, H]$ is convex in the state.

The generalized variance $\text{var}_\varrho^f(H)$ fulfills the following two requirements. (i) For pure states, the generalized variance equals the usual variance

$$\text{var}_\Psi^f(H) = (\Delta H)_\Psi^2. \quad (170)$$

(ii) For mixed states, $\text{var}_\varrho^f(H)$ is concave in the state.

The generalized quantum Fisher information and generalized variance fulfilling the above requirements are [60]

$$\mathcal{F}_Q^f[\varrho, H] = 2 \sum_{k,l} \frac{m_f(1,0)}{m_f(\lambda_k, \lambda_l)} (\lambda_k - \lambda_l)^2 |H_{kl}|^2, \quad (171a)$$

$$\begin{aligned} \text{var}_\varrho^f(H) &= \frac{1}{2} \sum_{k,l} \frac{m_f(\lambda_k, \lambda_l)}{m_f(1,0)} |H_{kl}|^2 \\ &\quad - \frac{1}{2m_f(1,0)} \left| \sum_k \lambda_k H_{kk} \right|^2, \end{aligned} \quad (171b)$$

where the matrix elements of H in the eigenbasis of ϱ are denoted as

$$H_{kl} = \langle k|H|l\rangle. \quad (172)$$

Note that the definition of the variance in Eq. (171b) requires that $m_f(1,0) \equiv f(0)$ is nonzero. In such cases f is called regular [101].

The usual quantum Fisher information and the usual variance corresponds to

$$f_{\max}(x) = \frac{1+x}{2}, \quad (173)$$

and the arithmetic mean

$$m_{f_{\max}}(a, b) = \frac{a+b}{2}. \quad (174)$$

The Wigner-Yanase skew information corresponds to

$$f_{\text{WY}}(x) = \frac{(\sqrt{x}+1)^2}{4}, \quad (175)$$

and the mean

$$m_{f_{\text{WY}}}(a, b) = \frac{(\sqrt{a} + \sqrt{b})^2}{4}. \quad (176)$$

Note that we get 4 times the usual Wigner-Yanase skew information due to the chosen normalization

$$\mathcal{F}_Q^{f_{\text{WY}}}[\varrho, H] = 4I_\varrho(H). \quad (177)$$

We now show a method to express the various types of generalized quantum Fisher information with each other.

Observation 14.—Let us define for given f the following matrix in the eigenbasis of ϱ

$$(X_f)_{kl} = \langle k|X|l\rangle = \sqrt{\frac{m_f(1,0)}{m_f(\lambda_k, \lambda_l)}}. \quad (178)$$

Then, any generalized quantum Fisher information can be expressed as

$$\mathcal{F}_Q^{f_1}[\varrho, H] = \mathcal{F}_Q^{f_2}[\varrho, Q_{f_1, f_2} \circ H], \quad (179)$$

where "o" denotes element-wise or Hadamard product defined as

$$A \circ B = \sum_{k,l} \langle k|A|l\rangle \langle k|B|l\rangle |k\rangle \langle l|, \quad (180)$$

where $|k\rangle$ and $|l\rangle$ are the eigenvectors of the density matrix, and the coefficient for converting one type of quantum Fisher information into another is given as

$$(Q_{f_1, f_2})_{kl} = \begin{cases} \frac{(X_{f_1})_{kl}}{(X_{f_2})_{kl}}, & \text{if } \lambda_k \neq \lambda_l, \\ 0, & \text{if } \lambda_k = \lambda_l. \end{cases} \quad (181)$$

Proof.—The statement can be verified by direct comparison of the definition of the quantum Fisher information given in Eq. (171a) and Eq. (179). ■

It is interesting to compute the matrix needed for $f_1 = f_{\max}$ and $f_2 = f_{WY}$. We obtain, in the basis of the eigenvectors of ϱ ,

$$(Q_{f_{\max}, f_{WY}})_{kl} = \begin{cases} \frac{(X_{f_{\max}})_{kl}}{(X_{f_{WY}})_{kl}} = \frac{\sqrt{\lambda_k} + \sqrt{\lambda_l}}{\sqrt{\lambda_k + \lambda_l}}, & \text{if } \lambda_k \neq \lambda_l, \\ 0, & \text{if } \lambda_k = \lambda_l. \end{cases} \quad (182)$$

Here note that in Eq. (182), the denominator of the fraction with λ_k and λ_l is positive if $\lambda_k \neq \lambda_l$. With the matrix in Eq. (182), we can use the Wigner-Yanase skew information to obtain a the quantum Fisher information

$$\mathcal{F}_Q[\varrho, H] = 4I_\varrho(Q_{f_{\max}, f_{WY}} \circ H). \quad (183)$$

We can also obtain the skew information with the quantum Fisher information as

$$I_\varrho(H) = \frac{1}{4} \mathcal{F}_Q[\varrho, Q_{f_{WY}, f_{\max}} \circ H], \quad (184)$$

where $Q_{f_{WY}, f_{\max}}$ is given in the basis of the eigenvectors of ϱ as

$$(Q_{f_{WY}, f_{\max}})_{kl} = \begin{cases} \frac{(X_{f_{WY}})_{kl}}{(X_{f_{\max}})_{kl}} = \frac{\sqrt{\lambda_k + \lambda_l}}{\sqrt{\lambda_k} + \sqrt{\lambda_l}}, & \text{if } \lambda_k \neq \lambda_l, \\ 0, & \text{if } \lambda_k = \lambda_l. \end{cases} \quad (185)$$

Next, we show how the various types of generalized quantum Fisher information can be expressed as a convex roof over the decomposition of the density matrix.

Observation 15.—The various types of generalized quantum Fisher information can be expressed as a convex roof as

$$\mathcal{F}_Q^f[\varrho, H] = 4 \min_{\{p_k, |\Psi_k\rangle\}} \sum_k p_k [\Delta(Y_f \circ H)]_{\Psi_k}^2, \quad (186)$$

where the optimization is over pure state decompositions given in Eq. (14), and in the basis of the eigenvectors of ϱ we define

$$(Y_f)_{kl} = \begin{cases} \frac{(X_f)_{kl}}{(X_{f_{\max}})_{kl}} = \sqrt{\frac{m_f(1,0)}{m_f(\lambda_k, \lambda_l)}} (\lambda_k + \lambda_l), & \text{if } \lambda_k \neq \lambda_l, \\ 0, & \text{if } \lambda_k = \lambda_l. \end{cases} \quad (187)$$

Proof.—It follows from Observation 14 defining $(X_f)_{kl}$ and the definition of the quantum Fisher information with the convex roof of the variance given in Eq. (13). ■

Next, we show how the various types of generalized quantum Fisher information can be expressed as an optimization in the two-copy space.

Observation 16.—The generalized quantum Fisher information can be obtained as an optimization over separable states

$$\begin{aligned} \mathcal{F}_Q^f[\varrho, H] &= \min_{\varrho_{12}} 2\text{Tr}[(Y_f \circ H \otimes \mathbb{1} - \mathbb{1} \otimes Y_f \circ H)^2 \varrho_{12}], \\ \text{s. t. } \varrho_{12} &\in \mathcal{S}, \\ \text{Tr}_2(\varrho_{12}) &= \varrho, \text{Tr}_1(\varrho_{12}) = \varrho. \end{aligned} \quad (188)$$

Proof.—It follows from Observation 1 defining the quantum Fisher information with an optimization over bipartite symmetric quantum states and Observation 14 defining $(X_f)_{kl}$. ■

Let us calculate a concrete example. For instance, for obtaining the Wigner-Yanase skew information times four as an optimization over separable states, we should use

$$(Y_{f_{WY}})_{kl} = (Q_{f_{WY}, f_{\max}})_{kl}, \quad (189)$$

where $Q_{f_{WY}, f_{\max}}$ is defined in Eq. (185).

Observation 17.—We can obtain the various types of quantum Fisher information as an optimization over general quantum states, rather than over separable state states, as

$$\begin{aligned} \mathcal{F}_Q^f[\varrho, H](\varrho) &= \frac{1}{2} \min_{\varrho_{12}} \text{Tr}\{[(Z_f \circ H)^T \otimes \mathbb{1} - \mathbb{1} \otimes Z_f \circ H]^2 \varrho_{12}\}, \\ \text{s. t. } \varrho_{12} &\in \mathcal{D}, \\ \text{Tr}_2(\varrho_{12}) &= \varrho^T, \\ \text{Tr}_1(\varrho_{12}) &= \varrho, \end{aligned} \quad (190)$$

where in the basis of the eigenvectors of ϱ we define

$$(Z_f)_{kl} = \begin{cases} \frac{(X_f)_{kl}}{(X_{f_{WY}})_{kl}}, & \text{if } \lambda_k \neq \lambda_l, \\ 0, & \text{if } \lambda_k = \lambda_l. \end{cases} \quad (191)$$

Proof.—We use the definition of $D_{\text{DPT}}(\varrho, \sigma)^2$ given in Eq. (1) and the equation relating it to the Wigner-Yanase skew information given in Eq. (3), together with Eq. (179). ■

Let us again calculate a concrete example. For instance, for obtaining the quantum Fisher information as an optimization over general quantum states, we should use

$$(Z_{f_{\max}})_{kl} = Q_{f_{\max}, f_{WY}}, \quad (192)$$

where $Q_{f_{\max}, f_{WY}}$ is defined in Eq. (182).

X. CONCLUSIONS

We discussed how to define the quantum Wasserstein distance as an optimization over bipartite separable states rather than an optimization over general quantum states. With such a definition, the self-distance becomes related to the quantum Fisher information. We introduced also variance-like quantities in which we replaced the minimization used in the definition of the quantum Wasserstein distance by a maximization, and examined their properties. We discussed the relation of our findings to entanglement criteria. We examined also the quantity obtained after we considered optimizing the variance rather than the second moment in the usual expression of the quantum Wasserstein distance. Finally, we extended our results to the various types of generalized quantum Fisher information. The details of the numerical calculations are discussed in Appendix A.

ACKNOWLEDGMENTS

We thank I. Apellaniz, F. Fröwis, I. L. Egusquiza, C. Klempt, J. Kołodyński, M. Mosonyi, G. Muga, J. Siewert, Sz. Szalay, K. Życzkowski, G. Vitagliano, and D.

Virosztek for discussions. We acknowledge the support of the EU (COST Action CA15220, QuantERA CEBBEC, QuantERA MENTA, QuantERA QuSiED), the Spanish MCIU (Grant No. PCI2018-092896), the Spanish Ministry of Science, Innovation and Universities and the European Regional Development Fund FEDER through Grant No. PGC2018-101355-B-I00 (MCIU/AEI/FEDER, EU) and through Grant No. PID2021-126273NB-I00 funded by MCIN/AEI/10.13039/501100011033 and by "ERDF A way of making Europe", the Basque Government (Grant No. IT986-16, No. IT1470-22), and the National Research, Development and Innovation Office NKFIH (Grant No. K124351, No. K124152, No. KH129601). We thank the "Front-line" Research Excellence Programme of the NKFIH (Grant No. KKP133827). We thank Project no. TKP2021-NVA-04, which has been implemented with the support provided by the Ministry of Innovation and Technology of Hungary from the National Research, Development and Innovation Fund, financed under the TKP2021-NVA funding scheme. We thank the Quantum Information National Laboratory of Hungary. G.T. is thankful for a Bessel Research Award from the Hum-

boldt Foundation.

Appendix A: Details of the numerical calculations

We used MATLAB [102] for numerical calculations. We used the semidefinite solver MOSEK [103] and the front-end YALMIP [104]. We also used the QUBIT4MATLAB package [105, 106]. $D_{\text{DPT}}(\varrho, \sigma)^2$ given in Definition 1 and $D_{\text{GMPC}}(\varrho, \sigma)^2$ given in Definition 2 can be obtained using semidefinite programming. $D_{\text{GMPC,sep}}(\varrho, \sigma)^2$ in Definition 3 and $D_{\text{DPT,sep}}(\varrho, \sigma)^2$ in Definition 4 need an optimization over separable states. The optimization over separable states can be carried out numerically using semidefinite programming for two qubits, since in this case the set of PPT states equals the set of separable states.

We included the routines computing the various types of quantum Wasserstein distance as `wdistsquare_GMPC_ppt.m`, `wdistsquare_GMPC.m`, `wdistsquare_DPT_ppt.m`, and `wdistsquare_DPT.m`. We included the various types of the variance-like quantities as `wvar_GMPC_ppt.m`, `wvar_GMPC.m`, `wvar_DPT_ppt.m`, and `wvar_DPT.m`. The usage of these routines is demonstrated in `example_wdistsquare.m`.

-
- [1] G. Monge, Mémoire sur la théorie des déblais et des remblais, Mémoires de l'Académie Royale de Sciences de Paris **625** (1781).
- [2] L. Kantorovitch, On the translocation of masses, *Management Science* **5**, 1 (1958).
- [3] E. Boissard, T. Le Gouic and J.-M. Loubes, Distributions template estimate with Wasserstein metric, *Bernoulli* **21**, 740 (2015).
- [4] O. Butkovsky, Subgeometric rates of convergence of Markov processes in the Wasserstein metric, *Ann. Appl. Probab.* **24**, 526 (2014).
- [5] M. Hairer, J.-C. Mattingly and M. Scheutzow, Asymptotic coupling and a general form of Harris' theorem with applications to stochastic delay equations, *Probab. Theory Relat. Fields* **149**, 223 (2011).
- [6] M. Hairer and J. Mattingly, Spectral Gaps in Wasserstein Distances and the 2D Stochastic Navier-Stokes Equations, *Ann. Probab.* **36**, 2050 (2008).
- [7] A. Figalli, F. Maggi and A. Pratelli, A mass transportation approach to quantitative isoperimetric inequalities, *Invent. Math.* **182**, 167 (2010).
- [8] A. Figalli and F. Maggi, On the shape of liquid drops and crystals in the small mass regime, *Arch. Ration. Mech. Anal.* **201**, 143 (2011).
- [9] J. Lott and C. Villani, Ricci curvature for metric-measure spaces via optimal transport, *Ann. of Math.* **169** (3), 903 (2009).
- [10] M.-K. von Renesse and K.-T. Sturm, Transport inequalities, gradient estimates, entropy, and Ricci curvature, *Comm. Pure Appl. Math.* **58**, 923 (2005).
- [11] K.-T. Sturm, On the geometry of metric measure spaces I, *Acta Math.* **196**, 65 (2006).
- [12] K.-T. Sturm, On the geometry of metric measure spaces II, *Acta Math.* **196**, 133 (2006).
- [13] B. Kloeckner, A geometric study of Wasserstein spaces: Euclidean spaces, *Annali della Scuola Normale Superiore di Pisa - Classe di Scienze, Scuola Normale Superiore* **2010 IX** (2), 297 (2010).
- [14] Gy. P. Gehér, T. Titkos, D. Virosztek, Isometric study of Wasserstein spaces – the real line, *Trans. Amer. Math. Soc.* **373**, 5855 (2020).
- [15] Gy. P. Gehér, T. Titkos, D. Virosztek, The isometry group of Wasserstein spaces: the Hilbertian case, *arXiv:2102.02037* (2021).
- [16] Gy. P. Gehér, T. Titkos, D. Virosztek, Isometric rigidity of the Wasserstein torus and the Wasserstein sphere, *arXiv:2203.04054* (2022).
- [17] S. Kolouri, S. R. Park and G. K. Rohde, The Radon cumulative distribution transform and its application to image classification, *IEEE Trans. Image Process.* **25**, 920 (2016).
- [18] W. Wang, D. Slepčev, S. Basu, J. A. Ozolek and G. K. Rohde, A linear optimal transportation framework for quantifying and visualizing variations in sets of images, *Int. J. Comput. Vis.* **101**, 254 (2013).
- [19] S. Kolouri, S. Park, M. Thorpe, D. Slepčev, G. K. Rohde, Optimal Mass Transport: Signal processing and machine-learning applications, *IEEE Signal Processing Magazine* **34**, 43 (2017).
- [20] A. Gramfort, G. Peyré and M. Cuturi, Fast Optimal Transport Averaging of Neuroimaging Data, *Information Processing in Medical Imaging, IPMI 2015. Lecture Notes in Computer Science* **9123**, 261 (2015).
- [21] Z. Su, W. Zeng, Y. Wang, Z. L. Lu and X. Gu, Shape classification using Wasserstein distance for brain morphometry analysis, *Information Processing in Medical Imaging, IPMI 2015. Lecture Notes in Computer Science* **24**, 411 (2015).
- [22] M. Arjovsky, S. Chintala and L. Bottou, Wasserstein Genera-

- tive Adversarial Networks, *Proceedings of Machine Learning Research* **70**, 214 (2017).
- [23] T. A. El Moselhy and Y. M. Marzouk, Bayesian inference with optimal maps, *J. Comput. Phys.* **231**, 7815 (2012).
- [24] G. Peyré and M. Cuturi, Computational Optimal Transport, *Found. Trends Machine Learn.* **11**, 355 (2019).
- [25] C. Frogner, C. Zhang, H. Mobahi, M. Araya and T. A. Poggio, Learning with a Wasserstein loss, [arXiv:1506.05439v3](https://arxiv.org/abs/1506.05439v3) (2015).
- [26] A. Ramdas, N.G. Trillos and M. Cuturi, On Wasserstein Two-Sample Testing and Related Families of Nonparametric Tests, *Entropy* **19**, 47. (2017).
- [27] S. Srivastava, C. Li and D.B. Dunson, Scalable Bayes via Barycenter in Wasserstein Space, *J. Mach. Learn. Res.* **19**, 1 (2018).
- [28] K. Życzkowski and W. Słomczynski, The Monge distance between quantum states, *J. Phys. A: Math. Gen.* **31**, 9095 (1998).
- [29] K. Życzkowski and W. Słomczynski, The Monge metric on the sphere and geometry of quantum states, *J. Phys. A: Math. Gen.* **34**, 6689 (2001).
- [30] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States: An Introduction to Quantum Entanglement* (Cambridge University Press, 2006).
- [31] P. Biane and D. Voiculescu, A free probability analogue of the Wasserstein metric on the trace-state space, *GAFN, Geom. Funct. Anal.* **11**, 1125 (2001).
- [32] E. Carlen and J. Maas, An Analog of the 2-Wasserstein Metric in Non-Commutative Probability Under Which the Fermionic Fokker-Planck Equation is Gradient Flow for the Entropy, *Commun. Math. Phys.* **331**, 887 (2014).
- [33] E. A. Carlen and J. Maas, Gradient flow and entropy inequalities for quantum Markov semigroups with detailed balance, *J. Funct. Anal.* **273**, 1810 (2017).
- [34] E. Carlen and J. Maas, Non-commutative calculus, optimal transport and functional inequalities in dissipative quantum systems, *J. Stat. Phys.* **178**, 319 (2020).
- [35] N. Datta and C. Rouzé, Concentration of quantum states from quantum functional and transportation cost inequalities, *J. Math. Phys.* **60**, 012202 (2019).
- [36] N. Datta and C. Rouzé, Relating relative entropy, optimal transport and Fisher information: A quantum HWI inequality, *Ann. Henri Poincaré* **21**, 2115 (2020).
- [37] F. Golse, C. Mouhot, and T. Paul, On the mean field and classical limits of quantum mechanics, *Commun. Math. Phys.* **343**, 165 (2016).
- [38] F. Golse and T. Paul, The Schrödinger equation in the mean-field and semiclassical regime, *Arch. Ration. Mech. Anal.* **223**, 57 (2017).
- [39] F. Golse and T. Paul, Wave packets and the quadratic Monge-Kantorovich distance in quantum mechanics, *Comptes Rendus Math.* **356**, 177 (2018).
- [40] F. Golse, The quantum N -body problem in the mean-field and semiclassical regime, *Phil. Trans. R. Soc. A* **376**, 20170229 (2018).
- [41] E. Caglioti, F. Golse, and T. Paul, Quantum optimal transport is cheaper, *J. Stat. Phys.* **181**, 149 (2020).
- [42] E. Caglioti, F. Golse, and T. Paul, Towards optimal transport for quantum densities, [arXiv:2101.03256](https://arxiv.org/abs/2101.03256) (2021).
- [43] G. De Palma and D. Trevisan, Quantum optimal transport with quantum channels, *Ann. Henri Poincaré* **22**, 3199 (2021).
- [44] G. De Palma, M. Marvian, D. Trevisan, and S. Lloyd, The quantum Wasserstein distance of order 1, *IEEE Trans. Inf. Theory* **67**, 6627 (2021).
- [45] S. Friedland, M. Eckstein, S. Cole, and K. Życzkowski, Quantum Monge–Kantorovich problem and transport distance between density matrices, *Phys. Rev. Lett.* **129**, 110402 (2022).
- [46] S. Friedland, M. Eckstein, S. Cole, and K. Życzkowski, Quantum optimal transport, [arXiv:2105.06922](https://arxiv.org/abs/2105.06922) (2021).
- [47] R. Bistrón, M. Eckstein, and K. Życzkowski, Monotonicity of the quantum 2-Wasserstein distance, [arXiv:2204.07405](https://arxiv.org/abs/2204.07405) (2022).
- [48] G. P. Gehér, J. Pitrik, T. Titkos, and D. Virosztek, Quantum Wasserstein isometries on the qubit state space, [arXiv:2204.14134](https://arxiv.org/abs/2204.14134) (2022).
- [49] L. Li, K. Bu, D. E. Koh, A. Jaffe, and S. Lloyd, Wasserstein complexity of quantum circuits, [arXiv: 2208.06306](https://arxiv.org/abs/2208.06306) (2022).
- [50] B. T. Kiani, G. D. Palma, M. Marvian, Z.-W. Liu, and S. Lloyd, Learning quantum data with the quantum earth mover’s distance, *Quantum Sci. Technol.* **7**, 045002 (2022).
- [51] E. P. Wigner and M. M. Yanase, Information contents of distributions, *Proc. Natl. Acad. Sci. U.S.A.* **49**, 910 (1963).
- [52] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, *Rev. Mod. Phys.* **81**, 865 (2009).
- [53] O. Gühne and G. Tóth, Entanglement detection, *Phys. Rep.* **474**, 1 (2009).
- [54] N. Friis, G. Vitagliano, M. Malik, and M. Huber, Entanglement certification from theory to experiment, *Nat. Rev. Phys.* **1**, 72 (2019).
- [55] We use the convention that in the expression within the trace in Eq. (1), $H_n \otimes \mathbb{1}$ denotes an operator in which H_n acts on subsystem 1 and $\mathbb{1}$ acts on subsystem 2.
- [56] V. Giovannetti, S. Lloyd, and L. Maccone, Quantum-enhanced measurements: Beating the standard quantum limit, *Science* **306**, 1330 (2004).
- [57] M. G. A. Paris, Quantum estimation for quantum technology, *Int. J. Quant. Inf.* **07**, 125 (2009).
- [58] R. Demkowicz-Dobrzanski, M. Jarzyna, and J. Kolodnyski, Chapter four - Quantum limits in optical interferometry, *Prog. Optics* **60**, 345 (2015), [arXiv:1405.7703](https://arxiv.org/abs/1405.7703).
- [59] L. Pezze and A. Smerzi, Quantum theory of phase estimation, in *Atom Interferometry (Proc. Int. School of Physics 'Enrico Fermi', Course 188, Varenna)*, edited by G. Tino and M. Kasevich (IOS Press, Amsterdam, 2014) pp. 691–741, [arXiv:1411.5164](https://arxiv.org/abs/1411.5164).
- [60] G. Tóth and D. Petz, Extremal properties of the variance and the quantum fisher information, *Phys. Rev. A* **87**, 032324 (2013).
- [61] S. Yu, Quantum fisher information as the convex roof of variance, [arXiv:1302.5311](https://arxiv.org/abs/1302.5311) (2013).
- [62] G. Tóth and F. Fröwis, Uncertainty relations with the variance and the quantum fisher information based on convex decompositions of density matrices, *Phys. Rev. Research* **4**, 013075 (2022).
- [63] S.-H. Chiew and M. Gessner, Improving sum uncertainty relations with the quantum Fisher information, *Phys. Rev. Research* **4**, 013076 (2022).
- [64] C. Helstrom, *Quantum Detection and Estimation Theory* (Academic Press, New York, 1976).
- [65] A. Holevo, *Probabilistic and Statistical Aspects of Quantum Theory* (North-Holland, Amsterdam, 1982).
- [66] S. L. Braunstein and C. M. Caves, Statistical distance and the geometry of quantum states, *Phys. Rev. Lett.* **72**, 3439 (1994).
- [67] D. Petz, *Quantum information theory and quantum statistics* (Springer, Berlin, Heilderberg, 2008).
- [68] S. L. Braunstein, C. M. Caves, and G. J. Milburn, Generalized uncertainty relations: Theory, examples, and Lorentz invariance, *Ann. Phys.* **247**, 135 (1996).
- [69] G. Tóth and I. Apellaniz, Quantum metrology from a quantum

- information science perspective, *J. Phys. A: Math. Theor.* **47**, 424006 (2014).
- [70] L. Pezzè, A. Smerzi, M. K. Oberthaler, R. Schmied, and P. Treutlein, Quantum metrology with nonclassical states of atomic ensembles, *Rev. Mod. Phys.* **90**, 035005 (2018).
- [71] Z. Léka and D. Petz, Some decompositions of matrix variances, *Probab. Math. Statist.* **33**, 191 (2013).
- [72] D. Petz and D. Viosztek, A characterization theorem for matrix variances, *Acta Sci. Math. (Szeged)* **80**, 681 (2014).
- [73] R. F. Werner, Quantum states with einstein-podolsky-rosen correlations admitting a hidden-variable model, *Phys. Rev. A* **40**, 4277 (1989).
- [74] K. Eckert, J. Schliemann, D. Bruss, and M. Lewenstein, Quantum correlations in systems of indistinguishable particles, *Ann. Phys.* **299**, 88 (2002).
- [75] T. Ichikawa, T. Sasaki, I. Tsutsui, and N. Yonezawa, Exchange symmetry and multipartite entanglement, *Phys. Rev. A* **78**, 052105 (2008).
- [76] P. Horodecki, Separability criterion and inseparable mixed states with positive partial transposition, *Phys. Lett. A* **232**, 333 (1997).
- [77] A. Peres, Separability criterion for density matrices, *Phys. Rev. Lett.* **77**, 1413 (1996).
- [78] P. Horodecki, M. Horodecki, and R. Horodecki, Bound entanglement can be activated, *Phys. Rev. Lett.* **82**, 1056 (1999).
- [79] G. Tóth and T. Vértesi, Quantum states with a positive partial transpose are useful for metrology, *Phys. Rev. Lett.* **120**, 020506 (2018).
- [80] S. Hill and W. K. Wootters, Entanglement of a pair of quantum bits, *Phys. Rev. Lett.* **78**, 5022 (1997).
- [81] W. K. Wootters, Entanglement of formation of an arbitrary state of two qubits, *Phys. Rev. Lett.* **80**, 2245 (1998).
- [82] D. P. DiVincenzo, C. A. Fuchs, H. Mabuchi, J. A. Smolin, A. Thapliyal, and A. Uhlmann, Entanglement of assistance, *quant-ph/9803033* (1998).
- [83] J. A. Smolin, F. Verstraete, and A. Winter, Entanglement of assistance and multipartite state distillation, *Phys. Rev. A* **72**, 052317 (2005).
- [84] H. F. Hofmann and S. Takeuchi, Violation of local uncertainty relations as a signature of entanglement, *Phys. Rev. A* **68**, 032103 (2003).
- [85] O. Gühne, Characterizing entanglement via uncertainty relations, *Phys. Rev. Lett.* **92**, 117903 (2004).
- [86] O. Gühne, M. Mechler, G. Tóth, and P. Adam, Entanglement criteria based on local uncertainty relations are strictly stronger than the computable cross norm criterion, *Phys. Rev. A* **74**, 010301 (2006).
- [87] G. Vitagliano, P. Hyllus, I. L. Egusquiza, and G. Tóth, Spin squeezing inequalities for arbitrary spin, *Phys. Rev. Lett.* **107**, 240502 (2011).
- [88] A. Edmonds, *Angular Momentum in Quantum Mechanics*, Investigations in Physics Series (Princeton University Press, 1996).
- [89] G. Tóth, Entanglement detection in optical lattices of bosonic atoms with collective measurements, *Phys. Rev. A* **69**, 052327 (2004).
- [90] G. Tóth, C. Knapp, O. Gühne, and H. J. Briegel, Optimal spin squeezing inequalities detect bound entanglement in spin models, *Phys. Rev. Lett.* **99**, 250405 (2007).
- [91] G. Tóth and M. W. Mitchell, Generation of macroscopic singlet states in atomic ensembles, *New J. Phys.* **12**, 053007 (2010).
- [92] G. Tóth, Detection of multipartite entanglement in the vicinity of symmetric Dicke states, *J. Opt. Soc. Am. B* **24**, 275 (2007).
- [93] G. Tóth, T. Moroder, and O. Gühne, Evaluating convex roof entanglement measures, *Phys. Rev. Lett.* **114**, 160501 (2015).
- [94] L. Vandenberghe and S. Boyd, Semidefinite programming, *SIAM Review* **38**, 49 (1996).
- [95] G. Tóth, Multipartite entanglement and high-precision metrology, *Phys. Rev. A* **85**, 022322 (2012).
- [96] P. Hyllus, W. Laskowski, R. Krischek, C. Schwemmer, W. Wieczorek, H. Weinfurter, L. Pezzè, and A. Smerzi, Fisher information and multiparticle entanglement, *Phys. Rev. A* **85**, 022321 (2012).
- [97] G. Tóth, T. Vértesi, P. Horodecki, and R. Horodecki, Activating hidden metrological usefulness, *Phys. Rev. Lett.* **125**, 020402 (2020).
- [98] D. Petz, Covariance and fisher information in quantum mechanics, *J. Phys. A: Math. Gen.* **35**, 929 (2002).
- [99] P. Gibilisco, F. Hiai, and D. Petz, Quantum covariance, quantum fisher information, and the uncertainty relations, *IEEE Trans. Inf. Theory* **55**, 439 (2009).
- [100] D. Petz and C. Ghinea, Introduction to quantum Fisher information, in *QP- PQ: Quantum Probability and White Noise Analysis*, Vol. 27, edited by R. Rebolledo and M. Orszag (World Scientific, 2011) pp. 261–281.
- [101] P. Gibilisco, D. Girolami, and F. Hansen, A unified approach to local quantum uncertainty and interferometric power by metric adjusted skew information, *Entropy* **23**, 263 (2021).
- [102] MATLAB, *9.9.0.1524771(R2020b)* (The MathWorks Inc., Natick, Massachusetts, 2020).
- [103] MOSEK ApS, *The MOSEK optimization toolbox for MATLAB manual. Version 9.0.* (2019).
- [104] J. Löfberg, Yalmip : A toolbox for modeling and optimization in matlab, in *In Proceedings of the CACSD Conference* (Taipei, Taiwan, 2004).
- [105] G. Tóth, QUBIT4MATLAB V3.0: A program package for quantum information science and quantum optics for MATLAB, *Comput. Phys. Commun.* **179**, 430 (2008).
- [106] The package QUBIT4MATLAB V6.1 is available at <https://www.mathworks.com/matlabcentral/fileexchange/8433>. The actual version of the package QUBIT4MATLAB is available at <http://goth.eu/qubit4matlab.html>.