

PRIME GEODESIC THEOREM AND CLOSED GEODESICS FOR LARGE GENUS

YUNHUI WU AND YUHAO XUE

ABSTRACT. Let \mathcal{M}_g be the moduli space of hyperbolic surfaces of genus g endowed with the Weil-Petersson metric. In this paper, we show that for any $\epsilon > 0$, as $g \rightarrow \infty$, for a generic surface in \mathcal{M}_g , the error term in the Prime Geodesic Theorem is bounded from above by $g \cdot t^{\frac{3}{4}+\epsilon}$, up to a uniform constant multiplication. The expected value of the error term in the Prime Geodesic Theorem over \mathcal{M}_g is also studied. As an application, we show that as $g \rightarrow \infty$, on a generic hyperbolic surface in \mathcal{M}_g most closed geodesics of length significantly less than \sqrt{g} are simple and non-separating, and most closed geodesics of length significantly greater than \sqrt{g} are not simple, which confirms a conjecture of Lipnowski-Wright.

A novel effective upper bound for intersection numbers on $\mathcal{M}_{g,n}$ is also established, when certain indices are large compared to $\sqrt{g+n}$.

1. INTRODUCTION

Let X_g be a closed hyperbolic surface of genus g ($g \geq 2$). The spectrum of X is a discrete closed subset in $\mathbb{R}^{\geq 0}$ and consists of eigenvalues with finite multiplicity. We enumerate them, counted with multiplicity, in the following increasing order

$$0 = \lambda_0(X_g) < \lambda_1(X_g) \leq \lambda_2(X_g) \leq \dots \rightarrow \infty.$$

For any $t > 1$, we denote by $\pi_{X_g}(t)$ the number of oriented primitive closed geodesics of length $\leq \ln t$. A remarkable application of Selberg's trace formula [Sel56] is the Prime Geodesic Theorem (PGT) (see *e.g.* [Hub61, Hej76, Ran77]) which states

$$(1) \quad \pi_{X_g}(t) = \text{Li}(t) + \sum_{\frac{3}{4} < s_j < 1} \text{Li}(t^{s_j}) + O_{X_g} \left(\frac{t^{\frac{3}{4}}}{\ln t} \right)$$

where $\text{Li}(t) = \int_2^t \frac{dx}{\ln x} \sim \frac{t}{\ln t}$ as $t \rightarrow \infty$, $s_j = \frac{1}{2} + \sqrt{\frac{1}{4} - \lambda_j(X_g)}$ and the implied constant depends on the geometry of X_g . As an analog of the Riemann hypothesis for closed hyperbolic surfaces, it is a well-known conjecture that *the $\frac{3}{4}$ in (1) can be replaced by $\frac{1}{2} + \epsilon$* . Rewrite (1) as

$$(2) \quad \pi_{X_g}(t) = \text{Li}(t) + \text{Er}_{X_g}(t).$$

So the error term $\text{Er}_{X_g}(t)$ depends on the geometry and first non-zero eigenvalue of X_g . There are many works on the study of the error term $\text{Er}_{X_g}(t)$,

especially for special arithmetic surfaces. See [Sar80, Iwa84, LRS95, Cai02, SY13] and the references therein.

Let \mathcal{M}_g be the moduli space of closed hyperbolic surfaces of genus g endowed with the Weil-Petersson metric. Based on her celebrated thesis works [Mir07a, Mir07b], Mirzakhani in [Mir13] initiated the study of random hyperbolic surfaces for large genus. In this paper, our aim is to study the average behavior of the error terms in (1) and (2) on \mathcal{M}_g for large g . More precisely, let $\text{Prob}_{\text{WP}}^g$ and \mathbb{E}_{WP}^g be the probability measure and expectation on \mathcal{M}_g respectively given by the Weil-Petersson metric. We apply recent results of [Mon22] and [WX22, LW21] to show that as $g \rightarrow \infty$, for a generic $X_g \in \mathcal{M}_g$, the error term $\text{Er}_{X_g}(t)$ in (2) can be bounded from above by $g \cdot \frac{t^{\frac{3}{4}+\epsilon}}{\ln t}$ up to a universal constant multiplication. More precisely,

Theorem 1. *There exists a universal constant $c > 0$ such that for any $\epsilon > 0$ and all $t = t(g) > 2$ which may depend on g ,*

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; |\pi_{X_g}(t) - \text{Li}(t)| \leq c \cdot g \cdot t^{\frac{3}{4}+\epsilon} \right) = 1.$$

Selberg's theory is important in the proof of Theorem 1. One may see [GLMST21, LS20, Mon22, Tho20, WX22, LW21, Hid21, Rud22] and the references therein for related topics on the same spirit. As an analog of the Riemann hypothesis for Weil-Petersson random hyperbolic surfaces, we conjecture

Conjecture. Theorem 1 still holds with replacing $\frac{3}{4}$ by $\frac{1}{2}$.

We also study the expected value $\mathbb{E}_{\text{WP}}^g [|\pi_{X_g}(t) - \text{Li}(t)|]$ for large genus. In light of (1), we need to study the term $\sum_{\frac{3}{4} < s_j < 1} \text{Li}(t^{s_j}) + O_{X_g} \left(\frac{t^{\frac{3}{4}}}{\ln t} \right)$ for large genus. Motivated by recent work [BP22] of Bourque-Petri, we first prove the following result in which the error term $O_{X_g} \left(\frac{t^{\frac{3}{4}}}{\ln t} \right)$ in (1) for a single surface $X_g \in \mathcal{M}_g$ is explicitly given in terms of the geometry of X_g . More precisely,

Theorem 2. *For any $X_g \in \mathcal{M}_g$ and all $t = t(g) > 2$ which may depend on g ,*

$$\begin{aligned} \pi_{X_g}(t) &= \text{Li}(t) + \sum_{0 < \lambda_j(X_g) \leq \frac{1}{4}} \text{Li}(t^{s_j}) \\ &+ O \left(g \frac{t^{\frac{5}{6}}}{\ln t} + g \frac{t^{\frac{2}{3}}}{\ln t} \max \left\{ 0, \ln \left(\frac{1}{\text{sys}(X_g) t^{\frac{1}{6}}} \right) \right\} \right) \end{aligned}$$

where $\text{sys}(X_g)$ is the length of shortest closed geodesic in X_g , and the implied constant is universal.

With the help of our previous work [WX22], we will get an effective upper bound of $\mathbb{E}_{\text{WP}}^g \left[\sum_{0 < \lambda_j(X_g) \leq \frac{1}{4}} \text{Li}(t^{s_j}) \right]$ for large genus. Then we apply Theorem 2 to prove the following bound on $\mathbb{E}_{\text{WP}}^g [\pi_{X_g}(t)]$ for large g and suitable $t = t(g)$. More precisely,

Theorem 3. *Let $t(g)$ be a function such that $t(g) > g^{12}$. Then for any $0 < \epsilon < 1$ and large enough g ,*

$$\mathbb{E}_{\text{WP}}^g \left[\left| 1 - \frac{\pi_{X_g}(t(g))}{\text{Li}(t(g))} \right| \right] = O_\epsilon(g^{-1+\epsilon})$$

where the implied constant only depends on ϵ .

For any $L > 0$, denote by $N(X_g, L)$ the number of unoriented primitive closed geodesics in X_g of length $\leq L$. So $N(X_g, L) = \frac{\pi_{X_g}(e^L)}{2}$. As introduced above, by the Prime Geodesic Theorem

$$N(X_g, L) \sim \frac{1}{2} \frac{e^L}{L} \text{ as } L \rightarrow \infty.$$

Denote by $N^s(X_g, L)$, $N_{nsep}^s(X_g, L)$, $N_{sep}^s(X_g, L)$ and $N^{ns}(X_g, L)$ the number of unoriented primitive *simple*, *simple and non-separating*, *simple and separating*, and *non-simple* closed geodesics in X_g of length $\leq L$ respectively. It is clear that $N(X_g, L) = N_{nsep}^s(X_g, L) + N_{sep}^s(X_g, L) + N^{ns}(X_g, L)$ and $N^s(X_g, L) = N_{nsep}^s(X_g, L) + N_{sep}^s(X_g, L)$. A celebrated result of Mirzakhani [Mir08] states that

$$N^s(X_g, L) \sim C(X_g) \cdot L^{6g-6} \text{ as } L \rightarrow \infty$$

where $C(X_g)$ is a constant depending on X_g , and as a function on \mathcal{M}_g , $C : \mathcal{M}_g \rightarrow \mathbb{R}$ is proper.

Based on Mirzakhani's Integration Formula [Mir07a], it is not hard to see that as $g \rightarrow \infty$, the expected values of $N^s(X_g, L)$ and $N_{nsep}^s(X_g, L)$ over \mathcal{M}_g satisfy that

$$\lim_{g \rightarrow \infty} \mathbb{E}_{\text{WP}}^g[N^s(X_g, L)] \sim \lim_{g \rightarrow \infty} \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)] \sim \frac{1}{2} \frac{e^L}{L} \text{ as } L \rightarrow \infty.$$

It is quite mysterious that both $N(X_g, L)$ and $\lim_{g \rightarrow \infty} \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]$ have the same asymptotic leading term $\frac{1}{2} \frac{e^L}{L}$ as $L \rightarrow \infty$. This suggests that for suitable $L = L(g)$ depending on g , as $g \rightarrow \infty$, in the set of closed geodesics of length $\leq L$ in a generic surface in \mathcal{M}_g , simple and non-separating ones are dominant. Meanwhile, as introduced above, $N^s(X_g, L) \sim C(X_g) \cdot L^{6g-6}$. This also seems to suggest that for suitable $L = L(g)$ depending on g , as $g \rightarrow \infty$, in the set of closed geodesics of length $\geq L$ in a generic surface in \mathcal{M}_g , non-simple ones are major part. Actually it was conjectured in [LW21, Conjecture 1.2] that

Conjecture (Lipnowski-Wright). As $g \rightarrow \infty$, on most closed hyperbolic surfaces in \mathcal{M}_g most geodesics of length significantly less than \sqrt{g} are simple and non-separating, and most geodesics of length significantly greater than \sqrt{g} are not simple.

It is known by [Mir13, Theorem 4.4] that for any $\epsilon > 0$,

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{M}_g; N_{\text{sep}}^s(X_g, (2 - \epsilon) \ln g) \geq 1) = 0.$$

And it is also known by [NWX20, Theorem 4] that for any $\epsilon > 0$,

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{M}_g; N^{ns}(X_g, (1 - \epsilon) \ln g) \geq 1) = 0.$$

The two limits above imply that if $L(g) \leq (1 - \epsilon_0) \ln g$ for some $\epsilon_0 > 0$, then

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{M}_g; N(X_g, L(g)) = N_{\text{nsep}}^s(X_g, L(g))) = 1.$$

This in particular tells that the first part of the conjecture of Lipnowski-Wright above is true when the length is less than $(1 - \epsilon_0) \ln g$. In this paper, we completely solve the conjecture above. More precisely, we prove

Theorem 4. *The following two probabilities hold:*

(1) *if $L(g)$ satisfies that for some fixed $\epsilon_0 > 0$,*

$$L(g) \geq (1 - \epsilon_0) \ln g \text{ and } \lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = 0,$$

then there exists a function $\delta(g) > 0$ satisfying $\lim_{g \rightarrow \infty} \delta(g) = 0$ such that

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N_{\text{nsep}}^s(X_g, L(g))}{N(X_g, L(g))} \right| < \delta(g) \right) = 1.$$

(2) *If $L(g)$ satisfies*

$$\lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = \infty,$$

then

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N^{ns}(X_g, L(g))}{N(X_g, L(g))} \right| < \frac{g}{L(g)^2} \right) = 1.$$

The proof of Part (1) of Theorem 4 above is divided into two parts: when $L(g) > 12 \ln g$, we use Theorem 1 and 3 which are obtained by spectral methods; when $L(g) \leq 12 \ln g$, we apply geometric methods developed in our previous works [NWX20, WX22].

Remark. It is known by [BS94] that $\sup_{X_{g_k} \in \mathcal{M}_{g_k}} \text{sys}(X_{g_k}) \geq \frac{4}{3} \ln g_k - C$ for a universal constant $C > 0$ and some sequence $\{g_k\}$ with $g_k \rightarrow \infty$ as $k \rightarrow \infty$. So for Part (1) of Theorem 4 above, $N(X_g, L(g))$ may vanish for certain $X_g \in \mathcal{M}_g$. However, it follows by [MP19, Theorem 5.1] that

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{M}_g; N(X_g, (1 - \epsilon) \ln g) \geq 1) = 1.$$

Thus, for Part (1) of Theorem 4 above, it suffices to consider $X_g \in \mathcal{M}_g$ with $N(X_g, (1 - \epsilon) \ln g) \geq 1$.

Recall that a closed geodesic $\gamma \subset X_g \in \mathcal{M}_g$ is called *filling* if each component of the complement $X_g \setminus \gamma$ of γ in X_g is homeomorphic to a disk. It is known by elementary isoperimetric inequality that the length of each filling closed geodesic in X_g grows at least like $2\pi(g - 1)$. We propose the following interesting question:

Question. As $g \rightarrow \infty$, on a generic closed hyperbolic surface in \mathcal{M}_g , are most closed geodesics of length significantly greater than g filling?

In the proof of Part (2) of Theorem 4, we establish a new bound on the Weil-Petersson volume $V_{g,n}(x_1, \dots, x_n)$ of moduli space of bordered hyperbolic surfaces with geodesic boundary components of length x_1, \dots, x_n , which relies on a novel bound on intersection numbers in this paper. Recall that on $\overline{\mathcal{M}}_{g,n}$ the compactified moduli space of surfaces of genus g with n punctures, there are n tautological line bundles $\{\mathcal{L}_i\}_{i=1}^n$. Let $\psi_i = c_1(\mathcal{L}_i) \in H_2(\overline{\mathcal{M}}_{g,n}, \mathbb{Q})$ be the first Chern class of \mathcal{L}_i and ω be the Weil-Petersson symplectic form on $\overline{\mathcal{M}}_{g,n}$. For any $d = (d_1, \dots, d_n)$ with each $d_i \in \mathbb{Z}_{\geq 0}$ and $|d| = d_1 + \dots + d_n \leq 3g - 3 + n$, the *intersection number* $[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}$ is defined as

$$[\tau_{d_1} \cdots \tau_{d_n}]_{g,n} = \frac{\prod_{i=1}^n 2^{2d_i} (2d_i + 1)!!}{(3g - 3 + n - |d|)!} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} \omega^{3g-3+n-|d|}.$$

In particular, $[\tau_0 \cdots \tau_0]_{g,n} = \text{Vol}(\mathcal{M}_{g,n})$, which is also denoted by $V_{g,n}$. There are deep connections between intersection numbers and Weil-Petersson volumes, especially for their asymptotics. Actually Mirzakhani [Mir07b] showed that Weil-Petersson volumes can be expressed in terms of intersection numbers. For asymptotic behaviors, one may see [Mir13, LX14, MZ15, DGZZ20, Agg21] and the references therein for related results. Based on her recursive formula, Mirzakhani in [Mir13, Page 286] showed that there exists a constant $c(n) > 0$ independent of g and d_i 's such that

$$1 - c(n) \frac{|d|^2}{g} \leq \frac{[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}}{V_{g,n}} \leq 1.$$

In particular, for fixed n and $|d| = o(\sqrt{g})$,

$$\frac{[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}}{V_{g,n}} \sim 1 \quad \text{as } g \rightarrow \infty.$$

In this paper, we study the case that $\max_{1 \leq i \leq n} \{d_i^2\}$ is large compared to $2g + n$, which seems to have not been presented generally before. We prove

Theorem 5. *There exists a universal constant $c' > 0$ such that*

$$\frac{[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}}{V_{g,n}} \leq c' \frac{2g - 2 + n}{\max_{1 \leq i \leq n} \{d_i^2\}}.$$

It is known by [MP19, Proposition 3.1] and [NWX20, Lemma 19] that there exists a constant $c(n) > 0$ only depending on n such that

$$\left(1 - c(n) \frac{\sum_{i=1}^n x_i^2}{g}\right) \prod_{i=1}^n \frac{\sinh(x_i/2)}{x_i/2} \leq \frac{V_{g,n}(x_1, \dots, x_n)}{V_{g,n}} \leq \prod_{i=1}^n \frac{\sinh(x_i/2)}{x_i/2}.$$

One may also see recent work [AM22] of Anantharaman-Monk for sharper results. In particular, for fixed n and $\sum x_i^2 = o(g)$,

$$\frac{V_{g,n}(x_1, \dots, x_n)}{V_{g,n}} \sim \prod_{i=1}^n \frac{\sinh(x_i/2)}{x_i/2} \quad \text{as } g \rightarrow \infty.$$

In this paper we apply Theorem 5 to study $V_{g,n}(x_1, \dots, x_n)$ when $\max_{1 \leq i \leq n} \{x_i^2\}$ is large compared to $2g + n$. The following bound is essential in the proof of Part (2) of Theorem 4.

Theorem 6. *For any $k \geq 1$, there exists a constant $c(k) > 0$ independent of g, n and x_i 's such that*

$$\frac{V_{g,n}(x_1, \dots, x_n)}{V_{g,n}} \leq \left(\prod_{i=1}^n \min \left\{ c(k) \frac{2g - 2 + n}{x_i^2}, 1 \right\} \right)^k \prod_{i=1}^n \frac{\sinh(x_i/2)}{x_i/2}.$$

We believe that both Theorem 5 and 6 will find further other applications on related problems.

Notations. We say two positive functions

$$f_1(g) \prec f_2(g) \quad \text{or} \quad f_2(g) \succ f_1(g)$$

if there exists a universal constant $C > 0$, independent of g , such that $f_1(g) \leq C \cdot f_2(g)$; and we say

$$f_1(g) \asymp f_2(g)$$

if $f_1(g) \prec f_2(g)$ and $f_2(g) \prec f_1(g)$. And $f_1(g) \prec_* f_2(g)$ means that the constant C depends on $*$.

We also say two functions

$$h_1 = O(h_2),$$

where h_1 may not be positive, if there exists a universal constant $C > 0$ such that $|h_1| \leq C \cdot h_2$. And $h_1 = O_*(h_2)$ means that C depends on $*$.

Plan of the paper. Section 2 will provide a review of relevant background materials. The proofs of Theorem 1 and 2 are independent. We will first prove Theorem 2 in Section 3, and then apply it to prove Theorem 3 in Section 4. In Section 5 we complete the proof of Theorem 1. Then in Section 6 we establish the two new bounds on intersection numbers and Weil-Petersson volumes, i.e., prove Theorem 5 and 6. In the last Section, we apply Theorem 1, 3, 5, 6 and our previous works [NW20, WX22] to prove Theorem 4. In Appendix A, we follow [WX22, Section 8] to prove a new counting result, i.e., Theorem 18, which will be used in the proof of Theorem 4 when $L(g) \asymp \ln g$.

Acknowledgement. We would like to thank all the participants in our seminar on Teichmüller theory for helpful discussions and comments on this project. We also would like to thank Zeév Rudnick for his useful comments. The first named author is partially supported by the NSFC grant No. 12171263.

CONTENTS

1. Introduction	1
Notations.	6
Plan of the paper.	7
Acknowledgement	7
2. Preliminaries	8
The Weil-Petersson metric	8
Mirzakhani's integration formula	9
The Weil-Petersson volume	9
Counting closed geodesics	10
Small eigenvalues	12
Selberg's trace formula	12
3. PGT for general hyperbolic surfaces	13
4. Expectation of the error term in PGT	21
5. PGT for random hyperbolic surfaces	24
5.1. First part of Theorem 31	25
5.2. Second part of Theorem 31	33
6. Intersection numbers and Weil-Petersson volumes	34
6.1. Intersection numbers	34
6.2. Weil-Petersson volumes	38
7. Proof of Theorem 4	40
7.1. Significantly greater than \sqrt{g}	40
7.2. Significantly less than \sqrt{g} and larger than $12 \ln g$	44
7.3. No more than $12 \ln g$	46
Appendix A. Counting filling k -tuples	57
References	60

2. PRELIMINARIES

In this section, we include certain necessary backgrounds and tools used in this paper.

The Weil-Petersson metric. We denote $S_{g,n}$ to be an oriented topological surface with g genus and n punctures or boundaries. Require $2g + n \geq 3$ so that $S_{g,n}$ admits hyperbolic metrics. Let $\mathcal{T}_{g,n}$ be the Teichmüller space of surfaces with g genus and n punctures. The moduli space of Riemann surfaces is $\mathcal{M}_{g,n} = \mathcal{T}_{g,n}/\text{Mod}_{g,n}$ where $\text{Mod}_{g,n}$ is the mapping class group of $S_{g,n}$ which fix the order of punctures. If $n = 0$, write $\mathcal{T}_g = \mathcal{T}_{g,0}$ and $\mathcal{M}_g = \mathcal{M}_{g,0}$ for simplicity. Given $L = (L_1, \dots, L_n) \in \mathbb{R}_{\geq 0}^n$, the Teichmüller space $\mathcal{T}_{g,n}(L)$ consists of bordered hyperbolic surfaces with geodesic boundary components of length L_1, \dots, L_n . Note that a geodesic boundary component of length 0 is a puncture, so $\mathcal{T}_{g,n}(0, \dots, 0) = \mathcal{T}_{g,n}$. The weighted moduli space is $\mathcal{M}_{g,n}(L) = \mathcal{T}_{g,n}(L)/\text{Mod}_{g,n}$.

Fix a pants decomposition $\{\alpha_i\}_{i=1}^{3g-3+n}$ of $S_{g,n}$, the Fenchel-Nielsen coordinates $X \mapsto (\ell_{\alpha_i}(X), \tau_{\alpha_i}(X))_{i=1}^{3g-3+n}$ is a global coordinates for the Teichmüller space $\mathcal{T}_{g,n}(L)$. Where $\ell_{\alpha_i}(X)$ is the length of α_i on X and $\tau_{\alpha_i}(X)$ is the twist along α_i (measured by length). Wolpert [Wol82] showed that the Weil-Petersson symplectic form of $\mathcal{T}_{g,n}(L)$ has a good expression under Fenchel-Nielsen coordinates:

Theorem 7 (Wolpert). *The Weil-Petersson symplectic form ω_{WP} on $\mathcal{T}_{g,n}(L)$ is given by*

$$\omega_{\text{WP}} = \sum_{i=1}^{3g-3+n} d\ell_{\alpha_i} \wedge d\tau_{\alpha_i}.$$

The Weil-Petersson volume form is

$$d\text{Vol}_{\text{WP}} = \frac{1}{(3g-3+n)!} \underbrace{\omega_{\text{WP}} \wedge \dots \wedge \omega_{\text{WP}}}_{3g-3+n \text{ copies}}.$$

This is a mapping class group invariant measure on $\mathcal{T}_{g,n}(L)$, and hence a measure on $\mathcal{M}_{g,n}(L)$, which still denoted by $d\text{Vol}_{\text{WP}}$. The total volume of $\mathcal{M}_{g,n}(L)$ under the Weil-Petersson metric is finite and denoted by $V_{g,n}(L)$. This also implies a probability measure $\text{Prob}_{\text{WP}}^g$ on \mathcal{M}_g :

$$\text{Prob}_{\text{WP}}^g(\mathcal{A}) := \frac{1}{V_g} \int_{\mathcal{M}_g} \mathbf{1}_{\mathcal{A}} dX$$

where $\mathcal{A} \subset \mathcal{M}_g$ is a Borel subset, $\mathbf{1}_{\mathcal{A}} : \mathcal{M}_g \rightarrow \{0, 1\}$ is its characteristic function, and dX is short for $d\text{Vol}_{\text{WP}}(X)$. One may see [Mir13, GPY11, MP19, NWX20, PWX21, SW22, HW22] and the references therein for recent studies of the geometry of Weil-Petersson random hyperbolic surfaces.

Mirzakhani's integration formula. Now we introduce an integration formula in [Mir07a, Mir13], which is an essential tool in the study of random hyperbolic surfaces of Weil-Petersson model.

Given any non-peripheral closed curve γ on topological surface $S_{g,n}$ and $X \in \mathcal{T}_{g,n}$, denote $\ell(\gamma) = \ell_\gamma(X)$ to be the hyperbolic length of the unique closed geodesic in the homotopy class γ on X . Let $\Gamma = (\gamma_1, \dots, \gamma_k)$ be an ordered k -tuple where the γ_i 's are distinct disjoint homotopy classes of nontrivial, non-peripheral, unoriented simple closed curves on $S_{g,n}$. Consider the orbit containing Γ under $\text{Mod}_{g,n}$ -action

$$\mathcal{O}_\Gamma = \{(h \cdot \gamma_1, \dots, h \cdot \gamma_k); h \in \text{Mod}_{g,n}\}.$$

Given a function $F : \mathbb{R}_{\geq 0}^k \rightarrow \mathbb{R}$, we define a function on $\mathcal{M}_{g,n}$

$$\begin{aligned} F^\Gamma : \mathcal{M}_{g,n} &\rightarrow \mathbb{R} \\ X &\mapsto \sum_{(\alpha_1, \dots, \alpha_k) \in \mathcal{O}_\Gamma} F(\ell_{\alpha_1}(X), \dots, \ell_{\alpha_k}(X)). \end{aligned}$$

Assume $S_{g,n} - \cup_{j=1}^k \gamma_j = \cup_{i=1}^s S_{g_i, n_i}$. For any $x = (x_1, \dots, x_k) \in \mathbb{R}_{\geq 0}^k$, we consider the moduli space $\mathcal{M}(S_{g,n}(\Gamma); \ell_\Gamma = x)$ of the hyperbolic surfaces (possibly disconnected) homeomorphic to $S_{g,n} - \cup_{j=1}^k \gamma_j$ with $\ell_{\gamma_i^1} = \ell_{\gamma_i^2} = x_i$ for every $i = 1, \dots, k$, where γ_i^1 and γ_i^2 are two boundary components of $S_{g,n} - \cup_{j=1}^k \gamma_j$ given by cutting along γ_i . Consider the Weil-Petersson volume

$$V_{g,n}(\Gamma, x) = \text{Vol}_{\text{WP}}(\mathcal{M}(S_{g,n}(\Gamma); \ell_\Gamma = x)) = \prod_{i=1}^s V_{g_i, n_i}(x^{(i)})$$

where $x^{(i)}$ is the list of those coordinates x_j of x such that γ_j is a boundary component of S_{g_i, n_i} . Using Theorem 7, Mirzakhani get the following integration formula. See [Mir07a, Theorem 7.1], [Mir13, Theorem 2.2], [MP19, Theorem 2.2], [Wri20, Theorem 4.1] for different versions.

Theorem 8 (Mirzakhani). *For any $\Gamma = (\gamma_1, \dots, \gamma_k)$, the integral of F^Γ over $\mathcal{M}_{g,n}$ with respect to Weil-Petersson metric is given by*

$$\int_{\mathcal{M}_{g,n}} F^\Gamma dX = C_\Gamma \int_{\mathbb{R}_{\geq 0}^k} F(x_1, \dots, x_k) V_{g,n}(\Gamma, x) x_1 \cdots x_k dx_1 \cdots dx_k$$

where the constant $C_\Gamma \in (0, 1]$ only depend on Γ . Moreover, $C_\Gamma = \frac{1}{2^k}$ if $g > 2$ and $S_{g,n} - \cup_{j=1}^k \gamma_j \cong S_{g-k, n+2k}$.

The Weil-Petersson volume. We denote $V_{g,n}(x_1, \dots, x_n)$ to be the Weil-Petersson volume of $\mathcal{M}_{g,n}(x_1, \dots, x_n)$ and $V_{g,n} = V_{g,n}(0, \dots, 0)$. Recall that the intersection numbers $[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}$ are integral of first Chern classes of tautological line bundles and Weil-Petersson symplectic form over $\overline{\mathcal{M}}_{g,n}$. Mirzakhani [Mir07b] showed that this volume can be expressed by intersection numbers:

Theorem 9.

$$V_{g,n}(2x_1, \dots, 2x_n) = \sum_{|d| \leq 3g-3+n} [\tau_{d_1} \cdots \tau_{d_n}]_{g,n} \prod_{i=1}^n \frac{x_i^{2d_i}}{(2d_i + 1)!}.$$

Mirzakhani gave some estimates about $V_{g,n}$:

Theorem 10. (1) [Mir13, Lemma 3.2] *For any $g, n \geq 0$,*

$$b_0 < \frac{(2g-2+n)V_{g,n}}{V_{g,n+1}} < b_1$$

where $b_0 = \frac{10-\pi^2}{120} \approx 0.001$ and $b_1 = \sum_{l=1}^{\infty} \frac{l\pi^{2l-2}}{(2l+1)!} \approx 0.401$.

(2) [Mir13, Theorem 3.5]

$$\frac{(2g-2+n)V_{g,n}}{V_{g,n+1}} = \frac{1}{4\pi^2} + O\left(\frac{1}{g}\right),$$

$$\frac{V_{g,n}}{V_{g-1,n+2}} = 1 + O\left(\frac{1}{g}\right).$$

Where the implied constants are related to n and independent of g .

Theorem 11 ([Mir13, Corollary 3.7]).

$$\sum_{\substack{g_1+g_2=g \\ 1 \leq g_1 \leq g_2}} \frac{V_{g_1,1}V_{g_2,1}}{V_g} \asymp \frac{1}{g} \quad \text{and} \quad \sum_{\substack{g_1+g_2=g-1 \\ 1 \leq g_1 \leq g_2}} \frac{V_{g_1,2}V_{g_2,2}}{V_g} \asymp \frac{1}{g^2}.$$

The implied constants are independent of g .

The asymptotic behavior of $V_{g,n}(x_1, \dots, x_n)$ was first studied in [MP19], also see [NWX20] and refined results in [AM22].

Theorem 12. *There exists a constant $c(n) > 0$ independent of g and x_i 's such that*

$$\left(1 - c(n) \frac{\sum_{i=1}^n x_i^2}{g}\right) \prod_{i=1}^n \frac{\sinh(x_i/2)}{x_i/2} \leq \frac{V_{g,n}(x_1, \dots, x_n)}{V_{g,n}} \leq \prod_{i=1}^n \frac{\sinh(x_i/2)}{x_i/2}.$$

Counting closed geodesics. Here we introduce some counting results. We want some upper bounds for the number of closed geodesics of length $\leq L$. First, the collar lemma (see e.g. [Bus10, Theorem 4.1.6]) states that:

Theorem 13. *In a hyperbolic surface $X_g \in \mathcal{M}_g$, there are at most $3g-3$ unoriented primitive closed geodesics of length $\leq 2 \operatorname{arcsinh} 1 \approx 1.7627$. Moreover, they are simple and disjoint with each other.*

Buser showed that:

Theorem 14 ([Bus10, Theorem 6.6.4]). *Let $X_g \in \mathcal{M}_g$ and $L > 0$. In X_g there are at most $(g-1)e^{L+6}$ oriented closed geodesics of length $\leq L$ which are not iterates of closed geodesics of length $\leq 2 \operatorname{arcsinh} 1$.*

For a hyperbolic surface $X_g \in \mathcal{M}_g$, denote $\mathcal{P}(X_g)$ to be the set of all oriented primitive closed geodesics in X_g . Then combine Theorem 13 and 14 we have:

Corollary 15. *For any $X_g \in \mathcal{M}_g$ and $L > 0$,*

$$\#\{\gamma \in \mathcal{P}(X_g); \ell(\gamma) \leq L\} \leq (g-1)e^{L+7}.$$

For compact hyperbolic surfaces with geodesic boundaries, the following counting result in our previous work [WX22] will be applied later.

Theorem 16 ([WX22, Theorem 4]). *For any $\epsilon_1 > 0$ and $m = 2g-2+n \geq 1$, there exists a constant $c(\epsilon_1, m) > 0$ only depending on m and ϵ_1 such that for all $L > 0$ and any compact hyperbolic surface X of genus g with n boundary simple closed geodesics, we have*

$$\#_f(X, L) \leq c(\epsilon_1, m) \cdot e^{L - \frac{1-\epsilon_1}{2}\ell(\partial X)}.$$

Where $\#_f(X, L)$ is the number of filling closed geodesics in X of length $\leq L$ and $\ell(\partial X)$ is the total length of the boundary closed geodesics of X .

Actually Theorem 16 also holds for filling multi-curves.

Definition 17. For $2g+n \geq 3$, let $\Gamma = (\gamma_1, \dots, \gamma_k)$ be an ordered k -tuple where γ_i 's are homotopy classes of nontrivial non-peripheral unoriented primitive closed curves on the topological surface $S_{g,n}$. We say Γ is *filling* in $S_{g,n}$ if each component of the complement $S_{g,n} \setminus \bigcup_{i=1}^k \gamma_i$ is homeomorphic to a disk or a punctured disk which is homotopic to a cusp of $S_{g,n}$.

In particular, a filling 1-tuple is a filling closed curve in $S_{g,n}$.

On a hyperbolic surface $X \in \mathcal{T}_{g,n}(L_1, \dots, L_n)$, we define the length of a k -tuple $\Gamma = (\gamma_1, \dots, \gamma_k)$ to be the total length of γ_i 's, that is,

$$\ell_\Gamma(X) := \sum_{i=1}^k \ell_{\gamma_i}(X).$$

Define the counting function $N_k^{\text{fill}}(X, L)$ for $L \geq 0$ and $X \in \mathcal{T}_{g,n}(L_1, \dots, L_n)$ to be

$$N_k^{\text{fill}}(X, L) := \#\left\{ \Gamma = (\gamma_1, \dots, \gamma_k); \begin{array}{l} \Gamma \text{ is a filling } k\text{-tuple in } X \\ \text{and } \ell_\Gamma(X) \leq L \end{array} \right\}.$$

Theorem 18. *For any $k \in \mathbb{Z}_{\geq 1}$, $0 < \epsilon < \frac{1}{2}$ and $m = 2g-2+n \geq 1$, there exists a constant $c(k, \epsilon, m) > 0$ only depending on k, ϵ and m such that for all $L > 0$ and any compact hyperbolic surface X of genus g with n boundary simple closed geodesics, the following holds:*

$$N_k^{\text{fill}}(X, L) \leq c(k, \epsilon, m) \cdot (1+L)^{k-1} e^{L - \frac{1-\epsilon}{2}\ell(\partial X)}.$$

Where $\ell(\partial X)$ is the total length of the boundary closed geodesics of X .

The proof of Theorem 18 is actually the same as the proof of Theorem 16 given in [WX22, Section 8]. We will give a sketch in Appendix A.

Small eigenvalues. For $X_g \in \mathcal{M}_g$, the spectrum of the Laplacian operator consists of discrete eigenvalues

$$0 = \lambda_0(X_g) < \lambda_1(X_g) \leq \lambda_2(X_g) \leq \cdots \rightarrow \infty.$$

Spectrum of hyperbolic surfaces are widely studied in the past century, see [Cha84, Bus10, Ber16] for example. Eigenvalues that are $\leq \frac{1}{4}$ are called small eigenvalues. Buser [Bus77] showed that $\lambda_{4g-2}(X_g) > \frac{1}{4}$ and constructed $\mathcal{X}_g \in \mathcal{M}_g$ which admits arbitrary small $\lambda_{2g-3}(\mathcal{X}_g)$. Later, Otal and Rosas [OR09] improved that $\lambda_{2g-2}(X_g) > \frac{1}{4}$. Here $2g-2$ is sharp. One may also see Ballmann-Matthiesen-Mondal [BMM16, BMM17] for more general statements on $\lambda_{2g-2}(X_g)$.

Theorem 19 ([OR09]). *There are at most $2g-2$ eigenvalues in $[0, \frac{1}{4}]$ for any $X_g \in \mathcal{M}_g$.*

Selberg's trace formula. We introduce the Selberg's trace formula [Sel56] of closed hyperbolic surfaces here, which is a useful tool when consider the spectrum of Laplacian and length spectrum. One may see [Bus10, Theorem 9.5.3] or [Ber16, Theorem 5.6] for the version we describe here.

Given a smooth function with compact support $\varphi \in C_c^\infty(\mathbb{R})$, its Fourier transform is defined as

$$\widehat{\varphi}(z) = \int_{\mathbb{R}} \varphi(t) e^{-itz} dt$$

for any $z \in \mathbb{C}$.

On a hyperbolic surface, a closed geodesic is called primitive if it is not an iterate of any other closed geodesic at least twice. For any hyperbolic surface X_g , we denote $\mathcal{P}(X_g)$ to be the set of all oriented primitive closed geodesics on X_g .

For $k \in \mathbb{Z}_{\geq 0}$, denote

$$(3) \quad r_k(X_g) = \begin{cases} \sqrt{\lambda_k(X_g) - \frac{1}{4}} & \text{if } \lambda_k(X_g) > \frac{1}{4} \\ \mathbf{i} \cdot \sqrt{\frac{1}{4} - \lambda_k(X_g)} & \text{if } \lambda_k(X_g) \leq \frac{1}{4} \end{cases}$$

where $\lambda_k(X_g)$ is the k -th eigenvalue of X_g . Then the remarkable trace formula of Selberg states:

Theorem 20 (Selberg's trace formula). *Let $X_g \in \mathcal{M}_g$ be a closed hyperbolic surface of genus g . For any even function $\varphi \in C_c^\infty(\mathbb{R})$, we have*

$$\begin{aligned} \sum_{k=0}^{\infty} \widehat{\varphi}(r_k(X_g)) &= (g-1) \int_{\mathbb{R}} r \tanh(\pi r) \widehat{\varphi}(r) dr \\ &+ \sum_{k=1}^{\infty} \sum_{\gamma \in \mathcal{P}(X_g)} \frac{\ell(\gamma)}{2 \sinh\left(\frac{k\ell(\gamma)}{2}\right)} \varphi(k\ell(\gamma)). \end{aligned}$$

And both sides of the formula are absolutely convergent.

3. PGT FOR GENERAL HYPERBOLIC SURFACES

In this section we finish the proof of Theorem 2. Actually we show

Theorem 21. *For any $X_g \in \mathcal{M}_g$ and $t > 2$,*

$$\pi_{X_g}(t) = \text{Li}(t) + \sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) + O \left(g \frac{t^{\frac{5}{6}}}{\ln t} + \frac{t^{\frac{2}{3}}}{\ln t} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < t^{-1/6}}} \ln \left(\frac{1}{\ell(\gamma)t^{\frac{1}{6}}} \right) \right)$$

where $s_j = s_j(X_g) = \frac{1}{2} + \sqrt{\frac{1}{4} - \lambda_j(X_g)}$, $\mathcal{P}(X_g)$ is the set of all oriented primitive closed geodesics in X_g , and the implied constant is universal.

Theorem 2 is a direct consequence of Theorem 21 above.

Proof of Theorem 2. By Theorem 13 we know that there exist at most $(3g-3)$ mutually disjoint primitive closed geodesics in X_g of length $\leq 2^{-1/6}$. So we have

$$\sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < t^{-\frac{1}{6}}}} \ln \left(\frac{1}{\ell(\gamma)t^{\frac{1}{6}}} \right) < g \cdot \max \left\{ 0, \ln \left(\frac{1}{\text{sys}(X_g)t^{\frac{1}{6}}} \right) \right\}$$

which together with Theorem 21 implies the conclusion. \square

The error term in Theorem 21 is explicitly given in terms of the geometry of X_g . In order to achieve it, we will choose suitable test functions for Selberg's trace formula such that we are able to distinguish all the geometric terms as shown in Theorem 21. Our method is motivated by the proof of [BP22, Corollary 1.4] by Bourque and Petri.

Before showing Theorem 21, we first make the following application.

Corollary 22. *For any $X_g \in \mathcal{M}_g$, if L satisfies $e^{\frac{1}{3}L} > \frac{1}{\text{sys}(X_g)}$, then*

$$\#\{\gamma \in \mathcal{P}(X_g); \ell(\gamma) \leq L\} < g \frac{e^L}{L}.$$

Proof. By Theorem 21 and 19,

$$\#\{\gamma \in \mathcal{P}(X_g); \ell(\gamma) \leq L\} < g \frac{e^L}{L} + \frac{e^{\frac{2}{3}L}}{L} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < e^{-L/6}}} \ln \left(\frac{1}{\ell(\gamma)e^{\frac{1}{6}L}} \right).$$

When $e^{\frac{1}{3}L} > \frac{1}{\text{sys}(X_g)}$, we have

$$\sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < e^{-L/6}}} \ln \left(\frac{1}{\ell(\gamma)e^{\frac{1}{6}L}} \right) < \max \left\{ 0, g \cdot \ln \left(\frac{1}{\text{sys}(X_g)} \right) \right\} \leq g e^{\frac{1}{3}L}.$$

Then the conclusion follows by the two inequalities above. \square

Now we return to prove Theorem 21.

For bounded t , the theorem holds naturally. So we only consider sufficiently large t . Let $\eta(x) \geq 0$ be a non-negative smooth even function with $\text{supp}(\eta) \subset (-\frac{1}{2}, \frac{1}{2})$, $\int_{\mathbb{R}} \eta(x) dx = 1$ and $\eta(0) = \max_{x \in \mathbb{R}} \eta(x) = 2$. Let $\eta_\epsilon(x) = \frac{1}{\epsilon} \eta(\frac{x}{\epsilon})$. For any $0 < \epsilon < 0.01$ and $L > 1$, let

$$(4) \quad f_\epsilon(x) = (\eta_\epsilon * \eta_\epsilon)(x),$$

$$(5) \quad \varphi_{L,\epsilon}^\pm(x) = \frac{1}{2}(f_\epsilon(x-L) + f_\epsilon(x+L)) \pm f_\epsilon(x).$$

We will use $\varphi_{L,\epsilon}^+$ and $\varphi_{L,\epsilon}^-$ as test functions to show the lower bound and upper bound of $\pi_{X_g}(t)$ respectively. By direct calculation, we have

$$0 \leq f_\epsilon(x) = \int_{\mathbb{R}} \eta_\epsilon(y) \eta_\epsilon(x-y) dy \leq \frac{2}{\epsilon} \int_{\mathbb{R}} \eta_\epsilon(y) dy = \frac{2}{\epsilon}.$$

So

$$(6) \quad 0 \leq \varphi_{L,\epsilon}^+(x) \leq \frac{4}{\epsilon} \quad \text{and} \quad -\frac{2}{\epsilon} \leq \varphi_{L,\epsilon}^-(x) \leq \frac{2}{\epsilon}.$$

It is clear that both f_ϵ and $\varphi_{L,\epsilon}^\pm$ are smooth even functions with compact supports. And their Fourier transforms satisfy

$$(7) \quad \widehat{f_\epsilon}(r) = (\widehat{\eta}(\epsilon r))^2 \geq 0,$$

$$(8) \quad \widehat{\varphi_{L,\epsilon}^+}(r) = (\cos(Lr) + 1)(\widehat{\eta}(\epsilon r))^2 \geq 0$$

and

$$(9) \quad \widehat{\varphi_{L,\epsilon}^-}(r) = (\cos(Lr) - 1)(\widehat{\eta}(\epsilon r))^2 \leq 0.$$

Lemma 23. *For any $L > 1$ and $\epsilon < 0.01$ we have*

$$\# \left\{ \begin{array}{l} \gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L \end{array} \right\} = 2 \int_{1-\epsilon}^{L+\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau,\epsilon}^\pm(\ell(\gamma)) d\tau.$$

Proof. For any $x > 1$ and $0 < \epsilon < 0.01$, since $\text{supp}(f_\epsilon) \subset (-\epsilon, \epsilon)$ we have

$$\begin{aligned} \int_0^\infty \varphi_{\tau,\epsilon}^\pm(x) \mathbf{1}_{[-\epsilon,\epsilon]}(x-\tau) d\tau &= \int_{x-\epsilon}^{x+\epsilon} \frac{1}{2} f_\epsilon(x-\tau) d\tau + \int_{x-\epsilon}^{x+\epsilon} \frac{1}{2} f_\epsilon(x+\tau) d\tau \\ &\quad \pm \int_{x-\epsilon}^{x+\epsilon} f_\epsilon(x) d\tau = \int_{-\epsilon}^\epsilon \frac{1}{2} f_\epsilon(\tau) d\tau = \frac{1}{2}. \end{aligned}$$

So

$$\begin{aligned} \# \left\{ \begin{array}{l} \gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L \end{array} \right\} &= \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} 1 \\ &= 2 \int_0^\infty \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau,\epsilon}^\pm(\ell(\gamma)) \mathbf{1}_{[-\epsilon,\epsilon]}(\ell(\gamma) - \tau) d\tau. \end{aligned}$$

Since $\varphi_{\tau,\epsilon}^{\pm}(\ell(\gamma)) = \frac{1}{2}f_{\epsilon}(\ell(\gamma) - \tau)$ when $1 < \ell(\gamma) \leq L$ and $\tau \geq 0$, we have

$$\begin{aligned}
\#\left\{ \begin{array}{l} \gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L \end{array} \right\} &= 2 \int_0^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \frac{1}{2} f_{\epsilon}(\ell(\gamma) - \tau) \mathbf{1}_{[-\epsilon, \epsilon]}(\ell(\gamma) - \tau) d\tau \\
&= 2 \int_0^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \frac{1}{2} f_{\epsilon}(\ell(\gamma) - \tau) d\tau \\
&= 2 \int_{1-\epsilon}^{L+\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \frac{1}{2} f_{\epsilon}(\ell(\gamma) - \tau) d\tau \\
&= 2 \int_{1-\epsilon}^{L+\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau,\epsilon}^{\pm}(\ell(\gamma)) d\tau.
\end{aligned}$$

Which finish the proof. \square

By Selberg's trace formula, i.e., Theorem 20,

$$\begin{aligned}
(10) \quad \sum_{k=1}^{\infty} \sum_{\gamma \in \mathcal{P}(X_g)} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} \varphi_{\tau,\epsilon}^{\pm}(k\ell(\gamma)) &= \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \widehat{\varphi_{\tau,\epsilon}^{\pm}}(r_j) + \sum_{\lambda_j > \frac{1}{4}} \widehat{\varphi_{\tau,\epsilon}^{\pm}}(r_j) \\
&\quad - (g-1) \int_{\mathbb{R}} r \tanh(\pi r) \widehat{\varphi_{\tau,\epsilon}^{\pm}}(r) dr.
\end{aligned}$$

We will estimate each term and give an upper bound for $\sum \varphi_{\tau,\epsilon}^{-}(\ell(\gamma))$ and a lower bound for $\sum \varphi_{\tau,\epsilon}^{+}(\ell(\gamma))$ which appear in Lemma 23.

Lemma 24. *For $\tau \in [1 - \epsilon, L + \epsilon]$ where $L > 1$ and $0 < \epsilon < 0.01$, we have*

$$\begin{aligned}
&\sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau,\epsilon}^{-}(\ell(\gamma)) \\
&\leq \frac{1}{2} \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \frac{e^{s_j \tau}}{\tau} + O_{\eta} \left(g\epsilon \frac{e^{\tau}}{\tau} + \frac{g}{\epsilon^2} \frac{e^{\frac{1}{2}\tau}}{\tau} + \frac{1}{\epsilon} \frac{e^{\frac{1}{2}\tau}}{\tau} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right).
\end{aligned}$$

Proof. Since $\varphi_{\tau,\epsilon}^{-}(x) \geq 0$ when $x > \epsilon$ and $\varphi_{\tau,\epsilon}^{-}(x) = 0$ when $x \geq \tau + \epsilon$, by applying $\varphi_{\tau,\epsilon}^{-}(x) \geq -\frac{2}{\epsilon}$ in (6) and the fact that the function $\frac{x}{2 \sinh \frac{x}{2}}$ is

decreasing on $(0, \infty)$, we have

$$\begin{aligned}
(11) \quad & \sum_{k=1}^{\infty} \sum_{\gamma \in \mathcal{P}(X_g)} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} \varphi_{\tau, \epsilon}^-(k\ell(\gamma)) \\
& \geq \frac{\tau + \epsilon}{2 \sinh \frac{\tau + \epsilon}{2}} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) > \epsilon}} \varphi_{\tau, \epsilon}^-(\ell(\gamma)) - \frac{2}{\epsilon} \sum_{k=1}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \leq \epsilon}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} \\
& \geq \frac{\tau + \epsilon}{2 \sinh \frac{\tau + \epsilon}{2}} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau, \epsilon}^-(\ell(\gamma)) - \frac{2}{\epsilon} \sum_{k=1}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \leq \epsilon}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}}.
\end{aligned}$$

Denote $\{\gamma_1, \dots, \gamma_m\}$ to be the set of all oriented primitive closed geodesics with length $\leq \epsilon$. By Theorem 13, $m \leq 6g - 6$. Then

$$\begin{aligned}
(12) \quad & \frac{2}{\epsilon} \sum_{k=1}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \leq \epsilon}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} = \frac{2}{\epsilon} \sum_{j=1}^m \sum_{k=1}^{[\epsilon/\ell(\gamma_j)]} \frac{\ell(\gamma_j)}{2 \sinh \frac{k\ell(\gamma_j)}{2}} \\
& \leq \frac{2}{\epsilon} \sum_{j=1}^m \sum_{k=1}^{[\epsilon/\ell(\gamma_j)]} \frac{1}{k} < \frac{1}{\epsilon} \sum_{j=1}^m \left(1 + \ln \frac{\epsilon}{\ell(\gamma_j)} \right) \\
& < \frac{g}{\epsilon} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)}.
\end{aligned}$$

Since $\widehat{\varphi_{\tau, \epsilon}^-}(r) \leq 0$ for $r \in \mathbb{R}$, we have

$$(13) \quad \sum_{\lambda_j > \frac{1}{4}} \widehat{\varphi_{\tau, \epsilon}^-}(r_j) \leq 0.$$

For each small eigenvalue $\lambda_j \in [0, \frac{1}{4}]$,

$$\begin{aligned}
\widehat{\varphi_{\tau, \epsilon}^-}(r_j) &= \left(\cosh(\tau \sqrt{\frac{1}{4} - \lambda_j}) - 1 \right) \left(\widehat{\eta}(\mathbf{i}\epsilon \sqrt{\frac{1}{4} - \lambda_j}) \right)^2 \\
&\leq \frac{1}{2} e^{(s_j - \frac{1}{2})\tau} (1 + O_{\eta}(\epsilon^2))
\end{aligned}$$

where $\widehat{\eta}(\mathbf{i}\epsilon \sqrt{\frac{1}{4} - \lambda_j}) = 1 + O_{\eta}(\epsilon^2)$ since $\widehat{\eta}(0) = 1$ and $\widehat{\eta}$ is even. And hence combine with Theorem 19 we have

$$(14) \quad \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \widehat{\varphi_{\tau, \epsilon}^-}(r_j) \leq \frac{1}{2} \sum_{0 \leq \lambda_j \leq \frac{1}{4}} e^{(s_j - \frac{1}{2})\tau} + O_{\eta}(g\epsilon^2 e^{\frac{1}{2}\tau}).$$

For the integral term,

$$\begin{aligned}
& \left| (g-1) \int_{\mathbb{R}} r \tanh(\pi r) \widehat{\varphi_{\tau,\epsilon}^-}(r) dr \right| \\
&= \left| 2(g-1) \int_0^\infty r \tanh(\pi r) (\cos(r\tau) - 1) (\widehat{\eta}(\epsilon r))^2 dr \right| \\
(15) \quad &\leq 4(g-1) \int_0^\infty r \cdot (\widehat{\eta}(\epsilon r))^2 dr \\
&= 4(g-1) \frac{1}{\epsilon^2} \int_0^\infty r \cdot (\widehat{\eta}(r))^2 dr \\
&\prec_\eta \frac{g}{\epsilon^2}.
\end{aligned}$$

Then plug (11), (12), (13), (14) and (15) into (10) we have

$$\begin{aligned}
& \frac{\tau + \epsilon}{2 \sinh \frac{\tau + \epsilon}{2}} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau,\epsilon}^-(\ell(\gamma)) \\
&\leq \frac{1}{2} \sum_{0 \leq \lambda_j \leq \frac{1}{4}} e^{(s_j - \frac{1}{2})\tau} + O_\eta \left(g\epsilon^2 e^{\frac{1}{2}\tau} + \frac{g}{\epsilon^2} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right).
\end{aligned}$$

Since $\frac{2 \sinh \frac{\tau + \epsilon}{2}}{\tau + \epsilon} \leq \frac{e^{\tau/2}}{\tau} e^{\epsilon/2} = \frac{e^{\tau/2}}{\tau} (1 + O(\epsilon))$, combining with Theorem 19 we get

$$\begin{aligned}
& \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau,\epsilon}^-(\ell(\gamma)) \\
&\leq \frac{1}{2} \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \frac{e^{s_j \tau}}{\tau} + O_\eta \left(g\epsilon \frac{e^\tau}{\tau} + \frac{g}{\epsilon^2} \frac{e^{\frac{1}{2}\tau}}{\tau} + \frac{1}{\epsilon} \frac{e^{\frac{1}{2}\tau}}{\tau} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right)
\end{aligned}$$

as desired. \square

Lemma 25. For $\tau \in [1 + \epsilon, L - \epsilon]$ where $L > 1$ and $0 < \epsilon < 0.01$, we have

$$\begin{aligned}
& \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau,\epsilon}^+(\ell(\gamma)) \\
&\geq \frac{1}{2} \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \frac{e^{s_j \tau}}{\tau} - O_\eta \left(g\epsilon \frac{e^\tau}{\tau} + g \frac{(1 + \epsilon\tau) e^{\frac{1}{2}\tau}}{\epsilon^2 \tau} + \frac{1}{\epsilon} \frac{e^{\frac{1}{2}\tau}}{\tau} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right).
\end{aligned}$$

Proof. Since $\varphi_{\tau,\epsilon}^+(x) = 0$ when $x \in [\epsilon, \tau - \epsilon] \cup [\tau + \epsilon, +\infty]$, by applying $\varphi_{\tau,\epsilon}^+(x) \leq \frac{4}{\epsilon}$ in (6) and the fact that the function $\frac{x}{2 \sinh \frac{x}{2}}$ is decreasing on

(0, ∞), we have

$$\begin{aligned}
(16) \quad & \sum_{k=1}^{\infty} \sum_{\gamma \in \mathcal{P}(X_g)} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} \varphi_{\tau, \epsilon}^+(k\ell(\gamma)) \\
& \leq \frac{\tau - \epsilon}{2 \sinh \frac{\tau - \epsilon}{2}} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) \in (\tau - \epsilon, \tau + \epsilon)}} \varphi_{\tau, \epsilon}^+(\ell(\gamma)) \\
& \quad + \frac{4}{\epsilon} \sum_{k=2}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \in (\tau - \epsilon, \tau + \epsilon)}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} + \frac{4}{\epsilon} \sum_{k=1}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \leq \epsilon}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} \\
& \leq \frac{\tau - \epsilon}{2 \sinh \frac{\tau - \epsilon}{2}} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau, \epsilon}^+(\ell(\gamma)) \\
& \quad + \frac{4}{\epsilon} \sum_{k=2}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \in (\tau - \epsilon, \tau + \epsilon)}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} + \frac{4}{\epsilon} \sum_{k=1}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \leq \epsilon}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}}.
\end{aligned}$$

With the same argument as (12),

$$(17) \quad \frac{4}{\epsilon} \sum_{k=1}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \leq \epsilon}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} \prec \frac{g}{\epsilon} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)}.$$

By Corollary 15,

$$\begin{aligned}
(18) \quad & \frac{4}{\epsilon} \sum_{k=2}^{\infty} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ k\ell(\gamma) \in (\tau - \epsilon, \tau + \epsilon)}} \frac{\ell(\gamma)}{2 \sinh \frac{k\ell(\gamma)}{2}} \leq \frac{4}{\epsilon} \frac{1}{2 \sinh \frac{\tau - \epsilon}{2}} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \frac{\tau + \epsilon}{2}}} \sum_{k=\lceil \frac{\tau - \epsilon}{\ell(\gamma)} \rceil}^{\lfloor \frac{\tau + \epsilon}{\ell(\gamma)} \rfloor} \ell(\gamma) \\
& \prec \frac{1}{\epsilon} e^{-\frac{\tau}{2}} \sum_{\gamma \in \mathcal{P}(X_g), \ell(\gamma) < \frac{\tau + \epsilon}{2}} \left(1 + \frac{2\epsilon}{\ell(\gamma)}\right) \ell(\gamma) \prec \frac{1}{\epsilon} e^{-\frac{\tau}{2}} \left((g-1)e^{\frac{\tau + \epsilon}{2} + \tau}\right) \tau \\
& \prec g \frac{\tau}{\epsilon}.
\end{aligned}$$

Since $\widehat{\varphi_{\tau, \epsilon}^+}(r) \geq 0$ for $r \in \mathbb{R}$, we have

$$(19) \quad \sum_{\lambda_j > \frac{1}{4}} \widehat{\varphi_{\tau, \epsilon}^+}(r_j) \geq 0.$$

For each small eigenvalue $\lambda_j \in [0, \frac{1}{4}]$,

$$\begin{aligned}
\widehat{\varphi_{\tau, \epsilon}^+}(r_j) &= \left(\cosh\left(\tau \sqrt{\frac{1}{4} - \lambda_j}\right) + 1 \right) \left(\widehat{\eta}(\mathbf{i}\epsilon \sqrt{\frac{1}{4} - \lambda_j}) \right)^2 \\
&\geq \frac{1}{2} e^{(s_j - \frac{1}{2})\tau} (1 + O_{\eta}(\epsilon^2))
\end{aligned}$$

where $\widehat{\eta}(i\epsilon\sqrt{\frac{1}{4}-\lambda_j}) = 1 + O_\eta(\epsilon^2)$ since $\widehat{\eta}(0) = 1$ and $\widehat{\eta}$ is even. And hence together with Theorem 19,

$$(20) \quad \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \widehat{\varphi}_{\tau, \epsilon}^+(r_j) \geq \frac{1}{2} \sum_{0 \leq \lambda_j \leq \frac{1}{4}} e^{(s_j - \frac{1}{2})\tau} - O_\eta \left(g\epsilon^2 e^{\frac{1}{2}\tau} \right).$$

For the integral term, similar as (15) we have

$$(21) \quad \begin{aligned} & \left| (g-1) \int_{\mathbb{R}} r \tanh(\pi r) \widehat{\varphi}_{\tau, \epsilon}^+(r) dr \right| \\ &= \left| 2(g-1) \int_0^\infty r \tanh(\pi r) (\cos(r\tau) + 1) (\widehat{\eta}(\epsilon r))^2 dr \right| \\ &\leq 4(g-1) \int_0^\infty r \cdot (\widehat{\eta}(\epsilon r))^2 dr \\ &= 4(g-1) \frac{1}{\epsilon^2} \int_0^\infty r \cdot (\widehat{\eta}(r))^2 dr \\ &\prec_\eta \frac{g}{\epsilon^2}. \end{aligned}$$

Then plug (16), (17), (18), (19), (20) and (21) into the trace formula (10), we have

$$\begin{aligned} & \frac{\tau - \epsilon}{2 \sinh \frac{\tau - \epsilon}{2}} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau, \epsilon}^+(\ell(\gamma)) \\ &\geq \frac{1}{2} \sum_{0 \leq \lambda_j \leq \frac{1}{4}} e^{(s_j - \frac{1}{2})\tau} - O_\eta \left(g\epsilon^2 e^{\frac{1}{2}\tau} + \frac{g}{\epsilon^2} + g\frac{\tau}{\epsilon} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right). \end{aligned}$$

Since $\frac{2 \sinh \frac{\tau - \epsilon}{2}}{\tau - \epsilon} \geq \frac{e^{\tau/2}}{\tau} (1 - O(\epsilon + e^{-\tau}))$, combining with Theorem 19 we get

$$\begin{aligned} & \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau, \epsilon}^+(\ell(\gamma)) \geq \frac{1}{2} \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \frac{e^{s_j \tau}}{\tau} \\ & - O_\eta \left(g\epsilon \frac{e^\tau}{\tau} + g \frac{(1 + \epsilon\tau) e^{\frac{1}{2}\tau}}{\epsilon^2 \tau} + \frac{1}{\epsilon} \frac{e^{\frac{1}{2}\tau}}{\tau} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right) \end{aligned}$$

as desired. \square

Now we are ready to prove Theorem 21.

Proof of Theorem 21. Lemma 23 and 24 give the upper bound:

$$\begin{aligned}
& \# \left\{ \begin{array}{l} \gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L \end{array} \right\} = 2 \int_{1-\epsilon}^{L+\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau, \epsilon}^-(\ell(\gamma)) d\tau \\
& \leq \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \int_{1-\epsilon}^{L+\epsilon} \frac{e^{s_j \tau}}{\tau} d\tau + O_\eta \left(g\epsilon e^{\frac{L}{2}} + \frac{g}{\epsilon^2} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right) \cdot \frac{e^{\frac{L}{2}}}{L} \\
& = \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \text{Li}(e^{s_j(L+\epsilon)}) + O_\eta \left(g\epsilon e^{\frac{L}{2}} + \frac{g}{\epsilon^2} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right) \cdot \frac{e^{\frac{L}{2}}}{L} \\
& = \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \text{Li}(e^{s_j L}) + O_\eta \left(g\epsilon e^{\frac{L}{2}} + \frac{g}{\epsilon^2} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right) \cdot \frac{e^{\frac{L}{2}}}{L}
\end{aligned}$$

where in the last equality we apply $\text{Li}(e^{s_j(L+\epsilon)}) = \text{Li}(e^{s_j L}) + O\left(\epsilon \frac{e^L}{L}\right)$.

For the other direction, Lemma 23 and 25 give the lower bound:

$$\begin{aligned}
& \# \left\{ \begin{array}{l} \gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L \end{array} \right\} = 2 \int_{1-\epsilon}^{L+\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau, \epsilon}^+(\ell(\gamma)) d\tau \\
& \geq 2 \int_{1+\epsilon}^{L-\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L}} \varphi_{\tau, \epsilon}^+(\ell(\gamma)) d\tau \geq \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \int_{1+\epsilon}^{L-\epsilon} \frac{e^{s_j \tau}}{\tau} d\tau \\
& \quad - O_\eta \left(g\epsilon e^{\frac{L}{2}} + g \frac{1+\epsilon L}{\epsilon^2} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right) \cdot \frac{e^{\frac{L}{2}}}{L} \\
& = \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \text{Li}(e^{s_j(L-\epsilon)}) \\
& \quad - O_\eta \left(g\epsilon e^{\frac{L}{2}} + g \frac{1+\epsilon L}{\epsilon^2} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right) \cdot \frac{e^{\frac{L}{2}}}{L} \\
& = \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \text{Li}(e^{s_j L}) - O_\eta \left(g\epsilon e^{\frac{L}{2}} + g \frac{1+\epsilon L}{\epsilon^2} + \frac{1}{\epsilon} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < \epsilon}} \ln \frac{\epsilon}{\ell(\gamma)} \right) \cdot \frac{e^{\frac{L}{2}}}{L}
\end{aligned}$$

where in the last equality we apply $\text{Li}(e^{s_j(L-\epsilon)}) = \text{Li}(e^{s_j L}) + O\left(\epsilon \frac{e^L}{L}\right)$. For sufficiently large L , take $\epsilon = e^{-\frac{1}{6}L} < 0.01$ and fix a function η , we have

$$\begin{aligned} \# \left\{ \begin{array}{l} \gamma \in \mathcal{P}(X_g), \\ 1 < \ell(\gamma) \leq L \end{array} \right\} &= \sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(e^{s_j L}) \\ &+ O \left(g \frac{e^{\frac{5}{6}L}}{L} + \frac{e^{\frac{2}{3}L}}{L} \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < e^{-L/6}}} \ln \frac{1}{\ell(\gamma) e^{\frac{1}{6}L}} \right). \end{aligned}$$

Then the proof is completed by taking $t = e^L$. \square

4. EXPECTATION OF THE ERROR TERM IN PGT

In this section we study the expected value of $\pi_{X_g}(t)$ over \mathcal{M}_g for large genus. We will apply Theorem 21 and our previous work [WX22] to complete the proof of Theorem 3.

We first see the following consequence of Theorem 21.

Corollary 26. *For any $t > 2$,*

$$\mathbb{E}_{\text{WP}}^g \left[\left| \pi_{X_g}(t) - \text{Li}(t) - \sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) \right| \right] \prec g \frac{t^{\frac{5}{6}}}{\ln t}.$$

Proof. By Theorem 13 we know that there are at most $6g - 6$ oriented primitive closed geodesics with length ≤ 1 . So for $t > 2$, we have

$$\sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < t^{-1/6}}} \ln \left(\frac{1}{\ell(\gamma) t^{\frac{1}{6}}} \right) \prec \max \left\{ 0, g \cdot \ln \left(\frac{1}{\text{sys}(X_g) t^{\frac{1}{6}}} \right) \right\} \leq \frac{g}{\text{sys}(X_g) t^{\frac{1}{6}}}.$$

It was shown by Mirzakhani in [Mir13, Corollary 4.3] that

$$\mathbb{E}_{\text{WP}}^g \left[\frac{1}{\text{sys}(X_g)} \right] \asymp 1.$$

Then the conclusion clearly follows by Theorem 21 and the two estimates above. \square

To prove Theorem 3, in light of Corollary 26 we need to estimate the term

$$\mathbb{E}_{\text{WP}}^g \left[\sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) \right].$$

We will apply our previous work [WX22] to control the term above. In [WX22] we proved

Theorem 27. *For any $\epsilon > 0$,*

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \lambda_1(X_g) > \frac{3}{16} - \epsilon \right) = 1.$$

One can also see an independent proof by Lipnowski-Wright in [LW21]. The first positive lower bound 0.0024 was established by Mirzakhani in [Mir13]. One may also see related results for random covers of a fixed closed hyperbolic surface in [MNP22] by Magee-Naud-Puder; and see a recent breakthrough by Hide-Magee [HM21] for optimal relative spectral gap of random covers of a fixed cusped hyperbolic surface.

Let ϕ_T be the test function on [WX22, Page 355]. Its Fourier transform satisfies

$$(22) \quad \widehat{\phi_T}(z) \geq 0 \text{ when } z \in \mathbb{R} \cup \mathbf{i} \cdot \mathbb{R}.$$

The proof of [WX22, Lemma 22] also gives that for any j with $\lambda_j(X_g) < \frac{1}{4}$ and any $0 < \epsilon < 1$ there exists a constant $C_\epsilon > 0$ only depending on ϵ such that

$$(23) \quad \widehat{\phi_{4 \ln g}}(r_j(X_g)) \geq C_\epsilon g^{4(1-\epsilon)\sqrt{\frac{1}{4}-\lambda_j(X_g)}} \ln g.$$

For the proof of Theorem 27, the essential part is the following estimation.

Theorem 28. *Let ϕ_T be the test function on [WX22, Page 355]. Then for any $\epsilon_1 > 0$, there exists a constant $C(\epsilon_1) > 0$ only depending on ϵ_1 such that for all $g > C(\epsilon_1)$,*

$$\mathbb{E}_{\text{WP}}^g \left[\sum_{0 < \lambda_j \leq \frac{1}{4}} \widehat{\phi_{4 \ln g}}(r_j(X_g)) \right] \leq g^{1+4\epsilon_1} (\ln g)^{68}.$$

Proof. This is exactly [WX22, Equation (56)]. \square

Now we apply Theorem 28 to prove the following result which is slightly stronger than Theorem 27 and essential in the proof of Theorem 3.

Theorem 29. *For any $\epsilon > 0$, $0 \leq r < \frac{1}{2}$ and $1 \leq j \leq 2g - 3$, there exists a constant $c(\epsilon) > 0$ independent of g , r and j such that for $g > 0$ large enough,*

$$\text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \lambda_j(X_g) \leq \frac{1}{4} - r^2 \right) \leq c(\epsilon) \frac{1}{j} g^{1-4r+\epsilon}.$$

Proof. By (22) and (23) we know that for any $0 < \epsilon < 1$ and any $0 \leq r < \frac{1}{2}$ there exists a constant $C_\epsilon > 0$ only depending on ϵ such that

$$\begin{aligned} & \mathbb{E}_{\text{WP}}^g \left[\sum_{0 < \lambda_j \leq \frac{1}{4}} \widehat{\phi_{4 \ln g}}(r_j(X_g)) \right] \\ & \geq \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \lambda_j(X_g) \leq \frac{1}{4} - r^2 \right) \cdot j \cdot C_\epsilon g^{4(1-\epsilon)r} \ln g. \end{aligned}$$

Which together with Theorem 28 implies that for any $\epsilon_1 > 0$ and $g > 0$ large enough we have

$$\text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \lambda_j(X_g) \leq \frac{1}{4} - r^2 \right) \leq \frac{(\ln g)^{67}}{j \cdot C_\epsilon} \cdot g^{1-4r+4(\epsilon_1+r\cdot\epsilon)}.$$

Then the conclusion follows by choosing suitable ϵ, ϵ_1 and $c(\epsilon)$. \square

Remark. If we choose $j = 1, r = \frac{1}{4} + \epsilon$, by letting $g \rightarrow \infty$ Theorem 29 implies Theorem 27.

Now we apply Theorem 29 to estimate $\mathbb{E}_{\text{WP}}^g \left[\sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) \right]$ for large $t = t(g)$. More precisely,

Theorem 30. *For any $\epsilon > 0$ and $t > 2g^4$, we have*

$$\mathbb{E}_{\text{WP}}^g \left[\sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) \right] \prec_\epsilon g^{-1+3\epsilon} \text{Li}(t).$$

Proof. For a non-negative random variable $\xi(X)$, the expectation is

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g[\xi] &= \frac{1}{V_g} \int_{\mathcal{M}_g} \xi(X) dX \\ &= \frac{1}{V_g} \int_{\mathcal{M}_g} \int_0^\infty \mathbf{1}_{\{\xi(X) \geq t\}} dt dX \\ &= \int_0^\infty \text{Prob}_{\text{WP}}^g(\xi(X) \geq t) dt. \end{aligned}$$

So

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g \left[\text{Li}(t^{s_j}) \mathbf{1}_{\{\frac{1}{2} \leq s_j < 1\}} \right] &= \int_0^\infty \text{Prob}_{\text{WP}}^g \left(\text{Li}(t^{s_j}) \mathbf{1}_{\{\frac{1}{2} \leq s_j < 1\}} \geq \rho \right) d\rho \\ &\leq \text{Li}(t^{\frac{1}{2}}) + \int_{\text{Li}(t^{\frac{1}{2}})}^{\text{Li}(t)} \text{Prob}_{\text{WP}}^g \left(\text{Li}(t^{s_j}) \mathbf{1}_{\{\frac{1}{2} \leq s_j < 1\}} \geq \rho \right) d\rho \\ &= \text{Li}(t^{\frac{1}{2}}) + \int_{\frac{1}{2}}^1 \text{Prob}_{\text{WP}}^g(s \leq s_j < 1) \left(\frac{d}{ds} \text{Li}(t^s) \right) ds \\ &= \text{Li}(t^{\frac{1}{2}}) + \int_{\frac{1}{2}}^1 \frac{t^s}{s} \text{Prob}_{\text{WP}}^g(s_j \geq s) ds \end{aligned}$$

where in the last equation we apply $\frac{d}{ds} \text{Li}(t^s) = \frac{d}{ds} \left(\int_2^{t^s} \frac{1}{\ln x} dx \right) = \frac{t^s}{s}$. By Theorem 29, for any $\epsilon > 0, \frac{1}{2} \leq s < 1$ and $1 \leq j \leq 2g - 3$, we have

$$\text{Prob}_{\text{WP}}^g(X_g \in \mathcal{M}_g; s_j(X_g) \geq s) \leq c(\epsilon) \frac{1}{j} g^{3-4s+\epsilon}.$$

Then it follows by Theorem 19 that for $t > 2g^4$,

$$\begin{aligned}
\mathbb{E}_{\text{WP}}^g \left[\sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) \right] &= \sum_{j=1}^{2g-3} \mathbb{E}_{\text{WP}}^g \left[\text{Li}(t^{s_j}) \mathbf{1}_{\{\frac{1}{2} \leq s_j < 1\}} \right] \\
&\leq (2g-3) \text{Li}(t^{\frac{1}{2}}) + \sum_{j=1}^{2g-3} \int_{\frac{1}{2}}^1 \frac{t^s}{s} c(\epsilon) \frac{1}{j} g^{3-4s+\epsilon} ds \\
&\asymp g \text{Li}(t^{\frac{1}{2}}) + c(\epsilon) \cdot g^{3-\epsilon} \cdot \left(\sum_{j=1}^{2g-3} \frac{1}{j} \right) \cdot \int_{\frac{1}{2}}^1 \left(\frac{t}{g^4} \right)^s ds \\
&\prec g \text{Li}(t^{\frac{1}{2}}) + c(\epsilon) g^{-1+\epsilon} \ln(g) \frac{t}{\ln t - 4 \ln g} \\
&\prec_{\epsilon} g \text{Li}(t^{\frac{1}{2}}) + g^{-1+2\epsilon} \text{Li}(t) \ln(g) \\
&\prec_{\epsilon} g^{-1+3\epsilon} \text{Li}(t).
\end{aligned}$$

The proof is complete. \square

Now we complete the proof of Theorem 3.

Proof of Theorem 3. By triangle inequality and Corollary 26 we know that for $t > g^{12}$,

$$\mathbb{E}_{\text{WP}}^g \left[|\pi_{X_g}(t) - \text{Li}(t)| \right] \prec \left(\mathbb{E}_{\text{WP}}^g \left[\sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) \right] + \frac{g}{t^{\frac{1}{6}}} \cdot \text{Li}(t) \right)$$

which together with Theorem 30 implies that for any $\epsilon > 0$,

$$\mathbb{E}_{\text{WP}}^g \left[\left| 1 - \frac{\pi_{X_g}(t)}{\text{Li}(t)} \right| \right] \prec_{\epsilon} \left(\frac{1}{g^{1-3\epsilon}} + \frac{g}{t^{\frac{1}{6}}} \right).$$

Then the conclusion clearly follows because $t > g^{12}$. \square

5. PGT FOR RANDOM HYPERBOLIC SURFACES

In this section, our main goal is to show that

Theorem 31. *There exists a subset $\mathcal{A}_g \subset \mathcal{M}_g$ with $\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g(\mathcal{A}_g) = 1$ such that for any $X_g \in \mathcal{A}_g$ and $t = t(g) > 2$ which may depend on g ,*

$$\pi_{X_g}(t) = \text{Li}(t) + \sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) + O\left(g \frac{t^{\frac{3}{4}}}{\ln t}\right)$$

where the implied constant is independent of t , g and X_g .

Theorem 1 is as a direct consequence.

Proof of Theorem 1. Since $\ln t > \ln 2$, it clearly follows by Theorem 19, 27 and 31. \square

For the proof of Theorem 31, we split into two parts. In the first subsection, with an assumption on Weyl's law which is based on the work of Monk [Mon22], we will show a Prime Geodesic Theorem with a complicated remainder. Then in the second subsection we will simplify the remainder to the form as shown in Theorem 31 for random hyperbolic surfaces.

5.1. First part of Theorem 31. We follow the strategy by Randol in [Cha84, Chapter XI].

For any $0 \leq a \leq b$, denote $N_{X_g}^\Delta(a, b)$ to be the number of eigenvalues of X_g in the interval $[a, b]$. The standard Weyl's law says that

$$\lim_{b \rightarrow \infty} \frac{N_{X_g}^\Delta(0, b)}{(g-1)b} = 1$$

for any fixed hyperbolic surface $X_g \in \mathcal{M}_g$. Monk [Mon22] gave a bound of the remainder of Weyl's law for random hyperbolic surfaces.

Theorem 32 ([Mon22, Theorem 5]). *There exists a universal constant $C > 0$ and a subset $\mathcal{B}_g \subset \mathcal{M}_g$ such that $1 - \text{Prob}_{\text{WP}}^g(\mathcal{B}_g) = O\left(g^{-\frac{1}{12}}(\ln g)^{\frac{9}{8}}\right)$. And for any large enough g , any $0 \leq a \leq b$ and any surface $X_g \in \mathcal{B}_g$, one can write*

$$\frac{N_{X_g}^\Delta(a, b)}{4\pi(g-1)} = \frac{1}{4\pi} \int_{\frac{1}{4}}^{\infty} \mathbf{1}_{[a, b]}(\lambda) \tanh\left(\pi\sqrt{\lambda - \frac{1}{4}}\right) d\lambda + R(X_g, a, b)$$

where

$$-C\sqrt{\frac{b+1}{\ln g}} \leq R(X_g, a, b) \leq C\sqrt{\frac{b+1}{\ln g}} \ln\left(2 + (b-a)\sqrt{\frac{\ln g}{b+1}}\right)^{\frac{1}{2}}.$$

Now we make the following definition.

Definition. For a constant $A \geq 1$, we say that a hyperbolic surface $X_g \in \mathcal{M}_g$ satisfies the condition \mathcal{P}_A if for any $T > 0$,

$$(24) \quad \#\left\{k; \frac{1}{4} < \lambda_k(X_g) \leq \frac{1}{4} + T\right\} \leq Ag(1+T).$$

If we choose $a = \frac{1}{4}$ and $b = \frac{1}{4} + T$ in Theorem 32, we know that there exists $A \geq 1$ such that as $g \rightarrow \infty$, the condition \mathcal{P}_A holds for a generic closed hyperbolic surface $X_g \in \mathcal{M}_g$.

In this subsection we aim to show that

Theorem 33. *If $X_g \in \mathcal{M}_g$ satisfies the condition \mathcal{P}_A for some $A \geq 1$, then for any $t > 2$,*

$$\pi_{X_g}(t) = \text{Li}(t) + \sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) + O\left(g\sqrt{A} \frac{t^{\frac{3}{4}}}{\ln t} + \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln\left(\frac{1}{\ell(\gamma)}\right)\right)$$

where the implied constant is independent of t , A and X_g .

For $X_g \in \mathcal{M}_g$ and $T > 0$, denote

$$(25) \quad \begin{aligned} H(T) = H(X_g, T) &:= \sum_{\ell(\gamma) \leq T} \frac{\Lambda(\gamma)}{2 \sinh(\frac{1}{2}\ell(\gamma))} 2 \cosh(\frac{1}{2}\ell(\gamma)) \\ &= \sum_{\ell(\gamma) \leq T} \Lambda(\gamma) \frac{1 + e^{-\ell(\gamma)}}{1 - e^{-\ell(\gamma)}} \end{aligned}$$

where the sum is taken over all oriented closed geodesics in X_g with length $\leq T$, the number $\Lambda(\gamma) = \ell(\gamma_0)$ where γ_0 is the unique oriented primitive closed geodesic such that γ is an iterate of γ_0 . Let

$$(26) \quad \varphi_T(x) := 2 \cosh\left(\frac{x}{2}\right) \cdot \mathbf{1}_{[-T, T]}(x)$$

where $\mathbf{1}_{[-T, T]}(x)$ is the characteristic function on $[-T, T]$. Since Selberg's trace formula only admits smooth functions, we do convolution to $\varphi_T(x)$. Let $\eta(x) \geq 0$ be a non-negative smooth even function satisfying that $\text{supp}(\eta) \subset (-1, 1)$ and $\int_{\mathbb{R}} \eta(x) dx = 1$. For any $0 < \epsilon < 1$, let

$$(27) \quad \eta_\epsilon(x) := \frac{1}{\epsilon} \eta\left(\frac{x}{\epsilon}\right)$$

and

$$(28) \quad \varphi_T^\epsilon(x) := \varphi_T * \eta_\epsilon(x) = \int_{\mathbb{R}} 2 \cosh\left(\frac{x-y}{2}\right) \mathbf{1}_{[-T, T]}(x-y) \eta_\epsilon(y) dy.$$

Then $\varphi_T^\epsilon(x)$ is a smooth even function with $\text{supp}(\varphi_T^\epsilon) \subset (-T - \epsilon, T + \epsilon)$. Their Fourier transforms satisfy

$$(29) \quad \begin{aligned} \widehat{\varphi_T}(r) &= \int_{-T}^T e^{-irx} 2 \cosh\left(\frac{x}{2}\right) dx \\ &= \frac{2 \sinh(T(\frac{1}{2} + ir))}{\frac{1}{2} + ir} + \frac{2 \sinh(T(\frac{1}{2} - ir))}{\frac{1}{2} - ir} \\ &= \frac{8}{1 + 4r^2} \left(2r \sin(rT) \cosh\left(\frac{T}{2}\right) + \cos(rT) \sinh\left(\frac{T}{2}\right) \right) \end{aligned}$$

and

$$(30) \quad \widehat{\varphi_T^\epsilon}(r) = \widehat{\varphi_T}(r) \widehat{\eta}_\epsilon(r) = \widehat{\varphi_T}(r) \widehat{\eta}(\epsilon r).$$

Denote

$$(31) \quad H_\epsilon(T) = H_\epsilon(X_g, T) := \sum_{\gamma} \frac{\Lambda(\gamma)}{2 \sinh(\frac{1}{2}\ell(\gamma))} \varphi_T^\epsilon(\ell(\gamma))$$

where the sum is taken over all oriented closed geodesics in X_g .

Proposition 34. *If $X_g \in \mathcal{M}_g$ satisfies the condition \mathcal{P}_A defined in (24) for some $A \geq 1$, then*

$$H_\epsilon(T) = e^T + \sum_{0 < \lambda_j \leq \frac{1}{4}} \frac{e^{s_j T}}{s_j} + O_\eta \left(Ag \frac{1}{\epsilon} e^{\frac{1}{2}T} + \epsilon^2 g e^T \right).$$

Proof. Plug φ_T^ϵ into Selberg's trace formula, i.e., Theorem 20, we have

$$(32) \quad H_\epsilon(T) = \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \widehat{\varphi}_T^\epsilon(r_j) + \sum_{\lambda_j > \frac{1}{4}} \widehat{\varphi}_T^\epsilon(r_j) - (g-1) \int_{\mathbb{R}} r \widehat{\varphi}_T^\epsilon(r) \tanh(\pi r) dr.$$

Now we estimate these three terms separately.

First we estimate the term $\sum_{0 \leq \lambda_j \leq \frac{1}{4}} \widehat{\varphi}_T^\epsilon(r_j)$. Note that $\widehat{\eta}(r) = 1 + O_\eta(r^2)$ since $\int_{\mathbb{R}} \eta(x) dx = 1$ and $\widehat{\eta}$ is even. For $0 \leq \lambda_j \leq \frac{1}{4}$ we have

$$(33) \quad \begin{aligned} \widehat{\varphi}_T^\epsilon(r_j) &= \widehat{\varphi}_T(r_j) \widehat{\eta}(\epsilon r_j) \\ &= \widehat{\eta}(\epsilon r_j) \left(\frac{2 \sinh(T(\frac{1}{2} + \sqrt{\frac{1}{4} - \lambda_j}))}{\frac{1}{2} + \sqrt{\frac{1}{4} - \lambda_j}} + \frac{2 \sinh(T(\frac{1}{2} - \sqrt{\frac{1}{4} - \lambda_j}))}{\frac{1}{2} - \sqrt{\frac{1}{4} - \lambda_j}} \right) \\ &= (1 + O_\eta(\epsilon^2)) \left(\frac{e^{s_j T}}{s_j} + O(e^{\frac{1}{2} T}) \right) \\ &= \frac{e^{s_j T}}{s_j} + O_\eta \left(e^{\frac{1}{2} T} + \epsilon^2 e^{s_j T} \right). \end{aligned}$$

By Theorem 19, there are at most $2g - 2$ eigenvalues in $[0, \frac{1}{4}]$. So

$$(34) \quad \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \widehat{\varphi}_T^\epsilon(r_j) = e^T + \sum_{0 < \lambda_j \leq \frac{1}{4}} \frac{e^{s_j T}}{s_j} + O_\eta \left(g e^{\frac{1}{2} T} + \epsilon^2 g e^T \right).$$

Next we estimate the integral term $(g-1) \int_{\mathbb{R}} r \widehat{\varphi}_T^\epsilon(r) \tanh(\pi r) dr$. For real r , since $|\widehat{\varphi}_T(r)| = O(\frac{1}{1+|r|} e^{\frac{1}{2} T})$ and $|\widehat{\eta}(\epsilon r)| = O_\eta(\frac{1}{1+|\epsilon r|^2})$, we have

$$(35) \quad \begin{aligned} \left| (g-1) \int_{\mathbb{R}} r \widehat{\varphi}_T^\epsilon(r) \tanh(\pi r) dr \right| &\prec_\eta g e^{\frac{1}{2} T} \int_{\mathbb{R}} \frac{1}{1+|\epsilon r|^2} dr \\ &\prec g \frac{1}{\epsilon} e^{\frac{1}{2} T}. \end{aligned}$$

For the term $\sum_{\lambda_j > \frac{1}{4}} \widehat{\varphi}_T^\epsilon(r_j)$ related to large eigenvalues, we first denote $N(r) = \#\{j; \frac{1}{4} < \lambda_j(X_g) \leq \frac{1}{4} + r^2\}$ and $dN(r) = \sum_{\lambda_j > \frac{1}{4}} \delta_{r_j}(r) dr$ to be a discrete measure where $r_j = \sqrt{\lambda_j - \frac{1}{4}}$ and $\delta(\cdot)$ is the Dirac function. Then

$$\sum_{\lambda_j > \frac{1}{4}} \widehat{\varphi}_T^\epsilon(r_j) = \int_0^\infty \widehat{\varphi}_T^\epsilon(r) dN(r).$$

Since X_g satisfies the condition \mathcal{P}_A ,

$$N(r) \leq A \cdot g \cdot (1 + r^2).$$

So we have $N(0) = 0$ and $\lim_{r \rightarrow \infty} \frac{N(r)}{(1+r)(1+\epsilon^2 r^2)} = 0$. Then using Integration by Parts we have

$$\begin{aligned}
\left| \sum_{\lambda_j > \frac{1}{4}} \widehat{\varphi}_T^\epsilon(r_j) \right| &\leq \int_0^\infty |\widehat{\varphi}_T^\epsilon(r)| dN(r) \\
&\prec_\eta \int_0^\infty \frac{e^{T/2}}{1+r} \frac{1}{1+|\epsilon r|^2} dN(r) \\
&= -e^{T/2} \int_0^\infty N(r) \frac{d}{dr} \left(\frac{1}{(1+r)(1+\epsilon^2 r^2)} \right) dr \\
&= e^{T/2} \int_0^\infty N(r) \left(\frac{1}{(1+r)^2(1+\epsilon^2 r^2)} + \frac{2\epsilon^2 r}{(1+r)(1+\epsilon^2 r^2)^2} \right) dr \\
&\prec e^{T/2} \int_0^\infty \frac{N(r)}{1+r^2} \frac{1}{1+\epsilon^2 r^2} dr.
\end{aligned}$$

So if $X_g \in \mathcal{M}_g$ satisfies the condition \mathcal{P}_A defined in (24), then

$$(36) \quad \left| \sum_{\lambda_j > \frac{1}{4}} \widehat{\varphi}_T^\epsilon(r_j) \right| \prec_\eta Ag \frac{1}{\epsilon} e^{\frac{1}{2}T}.$$

Then the conclusion follows by (32), (34), (35) and (36). \square

Lemma 35. *For any $0 < \epsilon < 1$, $T > \epsilon$, let φ_T and φ_T^ϵ be functions as above, then we have that for all $x > 0$,*

$$\frac{1}{\cosh \frac{\epsilon}{2}} \varphi_{T-\epsilon}^\epsilon(x) \leq \varphi_T(x) \leq \varphi_{T+\epsilon}^\epsilon(x).$$

Proof. If $x > T$, the conclusion clearly follows because

$$\varphi_{T-\epsilon}^\epsilon(x) = \varphi_T(x) = 0 \text{ and } \varphi_{T+\epsilon}^\epsilon(x) \geq 0.$$

Now we assume $x \in [0, T]$. By definition

$$\varphi_T(x) = 2 \cosh \frac{x}{2}.$$

Since η is even,

$$\begin{aligned}
&\int_{-\epsilon}^\epsilon 2 \cosh \frac{x-y}{2} \eta_\epsilon(y) dy \\
&= \int_0^\epsilon 2 \left(\cosh \frac{x-y}{2} + \cosh \frac{x+y}{2} \right) \eta_\epsilon(y) dy \\
&= 4 \int_0^\epsilon \cosh \frac{x}{2} \cosh \frac{y}{2} \eta_\epsilon(y) dy \\
&= 2 \cosh \frac{x}{2} \int_{-\epsilon}^\epsilon \cosh \frac{y}{2} \eta_\epsilon(y) dy.
\end{aligned}$$

Then we have

$$\begin{aligned}\varphi_{T+\epsilon}^\epsilon(x) &= \int_{-\epsilon}^\epsilon 2 \cosh \frac{x-y}{2} \eta_\epsilon(y) dy \\ &> 2 \cosh \frac{x}{2}\end{aligned}$$

and

$$\begin{aligned}\varphi_{T-\epsilon}^\epsilon(x) &\leq \int_{-\epsilon}^\epsilon 2 \cosh \frac{x-y}{2} \eta_\epsilon(y) dy \\ &< 2 \cosh \frac{x}{2} \cosh \frac{\epsilon}{2}.\end{aligned}$$

The proof is complete. \square

Proposition 36. *If $X_g \in \mathcal{M}_g$ satisfies the condition \mathcal{P}_A for some $A \geq 1$, then for $T > 0$ large enough,*

$$H(T) = e^T + \sum_{0 < \lambda_j \leq \frac{1}{4}} \frac{e^{s_j T}}{s_j} + O\left(g\sqrt{A}e^{\frac{3}{4}T}\right).$$

Proof. By the above Lemma 35,

$$\frac{1}{\cosh \frac{\epsilon}{2}} H_\epsilon(T - \epsilon) \leq H(T) \leq H_\epsilon(T + \epsilon).$$

Since $\frac{1}{\cosh(\epsilon/2)} = 1 + O(\epsilon^2)$ and $H_\epsilon(T)$ is increasing with respect to T ,

$$\begin{aligned}|H(T) - H_\epsilon(T)| &\leq |H_\epsilon(T + \epsilon) - H_\epsilon(T)| + \left| H_\epsilon(T + \epsilon) - \frac{1}{\cosh \frac{\epsilon}{2}} H_\epsilon(T - \epsilon) \right| \\ &\leq 2 |H_\epsilon(T + \epsilon) - H_\epsilon(T - \epsilon)| + O(\epsilon^2) H_\epsilon(T - \epsilon).\end{aligned}$$

Recall that $s_j \in [\frac{1}{2}, 1]$. Since $\frac{e^{s_j(T \pm \epsilon)}}{s_j} = \frac{e^{s_j T}}{s_j} (1 + O(\epsilon))$, it follows by Proposition 34 and Theorem 19 that for $X_g \in \mathcal{M}_g$ satisfying the condition \mathcal{P}_A defined in (24) we have

$$\epsilon^2 H_\epsilon(T - \epsilon) \prec Ag\epsilon e^{\frac{1}{2}T} + g\epsilon^2 e^T$$

and

$$\begin{aligned}& |H_\epsilon(T + \epsilon) - H_\epsilon(T - \epsilon)| \\ &= \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \frac{e^{s_j(T+\epsilon)}}{s_j} - \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \frac{e^{s_j(T-\epsilon)}}{s_j} + O_\eta \left(Ag \frac{1}{\epsilon} e^{\frac{1}{2}T} + g\epsilon^2 e^T \right) \\ &= O(\epsilon) \sum_{0 \leq \lambda_j \leq \frac{1}{4}} \frac{e^{s_j T}}{s_j} + O_\eta \left(Ag \frac{1}{\epsilon} e^{\frac{1}{2}T} + g\epsilon^2 e^T \right) \\ &= O_\eta \left(Ag \frac{1}{\epsilon} e^{\frac{1}{2}T} + g\epsilon e^T \right).\end{aligned}$$

So

$$|H(T) - H_\epsilon(T)| \prec_\eta \left(Ag \frac{1}{\epsilon} e^{\frac{1}{2}T} + g\epsilon e^T \right).$$

And hence

$$H(T) = e^T + \sum_{0 < \lambda_j \leq \frac{1}{4}} \frac{e^{s_j T}}{s_j} + O_\eta \left(Ag \frac{1}{\epsilon} e^{\frac{1}{2}T} + g\epsilon e^T \right).$$

Then the conclusion follows by taking $\epsilon = \sqrt{A}e^{-\frac{1}{4}T}$ and fixing η . \square

Now we transfer to prove for $\pi_{X_g}(t)$. Denote

$$(37) \quad \psi(T) = \psi(X_g, T) := \sum_{\ell(\gamma) \leq T} \Lambda(\gamma)$$

where the sum is taken over all oriented closed geodesics in X_g with length $\leq T$, and the number $\Lambda(\gamma) = \ell(\gamma_0)$ where γ_0 is the unique oriented primitive closed geodesic such that γ is an iterate of γ_0 . Then we have

Lemma 37.

$$0 \leq H(T) - \psi(T) \prec g \cdot (1 + T^2) + \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln \left(\frac{1}{\ell(\gamma)} \right)$$

where $\mathcal{P}(X_g)$ is the set of all oriented primitive closed geodesics in X_g .

Proof. Rewrite

$$\begin{aligned} H(T) &= \sum_{\ell(\gamma) \leq T} \Lambda(\gamma) \frac{1 + e^{-\ell(\gamma)}}{1 - e^{-\ell(\gamma)}} \\ &= \psi(T) + \sum_{\ell(\gamma) \leq T} \Lambda(\gamma) \frac{2}{e^{\ell(\gamma)} - 1}. \end{aligned}$$

So

$$0 \leq H(T) - \psi(T) = \sum_{\substack{\ell(\gamma) \leq T \\ \Lambda(\gamma) > 2 \operatorname{arcsinh} 1}} \frac{2\Lambda(\gamma)}{e^{\ell(\gamma)} - 1} + \sum_{\substack{\ell(\gamma) \leq T \\ \Lambda(\gamma) \leq 2 \operatorname{arcsinh} 1}} \frac{2\Lambda(\gamma)}{e^{\ell(\gamma)} - 1}.$$

For the first term, by Theorem 14,

$$\begin{aligned} \sum_{\substack{\ell(\gamma) \leq T \\ \Lambda(\gamma) > 2 \operatorname{arcsinh} 1}} \frac{2\Lambda(\gamma)}{e^{\ell(\gamma)} - 1} &\prec \sum_{k=1}^{[T]} \sum_{\substack{k < \ell(\gamma) \leq k+1 \\ \Lambda(\gamma) > 2 \operatorname{arcsinh} 1}} \frac{\ell(\gamma)}{e^{\ell(\gamma)}} \\ &\prec \sum_{k=1}^{[T]} g e^k \frac{k}{e^k} \\ &\prec g T^2. \end{aligned}$$

For the second term, let $\{\gamma_1, \dots, \gamma_m\}$ be the set of all oriented primitive closed geodesics in X_g with length $\leq 2 \operatorname{arcsinh} 1$. By Theorem 13, $m \leq 6g-6$. Then we have

$$\begin{aligned}
\sum_{\substack{\ell(\gamma) \leq T \\ \Lambda(\gamma) \leq 2 \operatorname{arcsinh} 1}} \frac{2\Lambda(\gamma)}{e^{\ell(\gamma)} - 1} &= \sum_{j=1}^m \sum_{k=1}^{\lfloor T/\ell(\gamma_j) \rfloor} \frac{2\ell(\gamma_j)}{e^{k\ell(\gamma_j)} - 1} \\
&\leq \sum_{j=1}^m \ell(\gamma_j) \left(\frac{2}{e^{\ell(\gamma_j)} - 1} + \int_1^\infty \frac{2}{e^{x\ell(\gamma_j)} - 1} dx \right) \\
&\leq \sum_{j=1}^m \ell(\gamma_j) \left(\frac{2}{\ell(\gamma_j)} + \frac{2}{\ell(\gamma_j)} \ln \left(\frac{1}{1 - e^{-\ell(\gamma_j)}} \right) \right) \\
&\prec \sum_{j=1}^m \left(1 + \max \left\{ 1, \ln \frac{1}{\ell(\gamma_j)} \right\} \right) \\
&\prec g + \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln \left(\frac{1}{\ell(\gamma)} \right).
\end{aligned}$$

Then combining the two parts above, we finish the proof. \square

Now denote

$$(38) \quad \nu(T) = \nu(X_g, T) := \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) \leq T}} \ell(\gamma)$$

where $\mathcal{P}(X_g)$ is the set of all oriented primitive closed geodesics in X_g . Then we have

Lemma 38.

$$0 \leq \psi(T) - \nu(T) \prec gTe^{\frac{1}{2}T}.$$

Proof. By definitions,

$$\begin{aligned}
0 \leq \psi(T) - \nu(T) &= \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) \leq T/2}} \sum_{k=2}^{\lfloor T/\ell(\gamma) \rfloor} \ell(\gamma) \\
&\leq T \cdot \# \left\{ \gamma \in \mathcal{P}(X_g); \ell(\gamma) \leq \frac{T}{2} \right\}.
\end{aligned}$$

By Corollary 15,

$$\# \left\{ \gamma \in \mathcal{P}(X_g); \ell(\gamma) \leq \frac{T}{2} \right\} \prec ge^{\frac{1}{2}T}.$$

Then the conclusion follows. \square

Now we are ready to prove Theorem 33.

Proof of Theorem 33. First combining Proposition 36, Lemma 37 and 38 we have

$$(39) \quad \nu(T) = e^T + \sum_{0 < \lambda_j \leq \frac{1}{4}} \frac{e^{s_j T}}{s_j} + O \left(g\sqrt{A}e^{\frac{3}{4}T} + \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln \left(\frac{1}{\ell(\gamma)} \right) \right).$$

We know that

$$(40) \quad \begin{aligned} \pi_{X_g}(t) &= \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) \leq \ln t}} \frac{1}{\ell(\gamma)} \ell(\gamma) \\ &= \#\{\gamma \in \mathcal{P}(X_g); \ell(\gamma) \leq \ln 2\} + \int_{T \in (\ln 2, \ln t]} \frac{1}{T} d\nu(T) \end{aligned}$$

where $d\nu(T) = \sum \ell(\gamma) \delta_{\ell(\gamma)}(T) dT$ is a discrete measure, and $\delta(\cdot)$ is the Dirac function. By using (39), Integration by Parts and Theorem 19, the second term satisfies

$$(41) \quad \begin{aligned} \int_{(\ln 2, \ln t]} \frac{d\nu(T)}{T} &= \frac{\nu(\ln t)}{\ln t} - \frac{\nu(\ln 2)}{\ln 2} + \int_{\ln 2}^{\ln t} \nu(T) \frac{1}{T^2} dT \\ &= \frac{t}{\ln t} + \sum_{0 < \lambda_j \leq \frac{1}{4}} \frac{t^{s_j}}{s_j \ln t} - \frac{2}{\ln 2} - \sum_{0 < \lambda_j \leq \frac{1}{4}} \frac{2^{s_j}}{s_j \ln 2} \\ &\quad + \int_{\ln 2}^{\ln t} \left(e^T + \sum_{0 < \lambda_j \leq \frac{1}{4}} \frac{e^{s_j T}}{s_j} \right) \frac{1}{T^2} dT \\ &\quad + O \left(g\sqrt{A} \frac{t^{\frac{3}{4}}}{\ln t} + \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln \frac{1}{\ell(\gamma)} \right) \\ &= \int_{\ln 2}^{\ln t} \frac{e^T}{T} dT + \sum_{0 < \lambda_j \leq \frac{1}{4}} \int_{\ln 2}^{\ln t} \frac{e^{s_j T}}{T} dT \\ &\quad + O \left(g\sqrt{A} \frac{t^{\frac{3}{4}}}{\ln t} + \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln \frac{1}{\ell(\gamma)} \right) \\ &= \text{Li}(t) + \sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) + O \left(g\sqrt{A} \frac{t^{\frac{3}{4}}}{\ln t} + \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln \frac{1}{\ell(\gamma)} \right). \end{aligned}$$

For the first term, it follows by Theorem 13 that for $t > 2$,

$$(42) \quad \#\{\gamma \in \mathcal{P}(X_g); \ell(\gamma) \leq \ln 2\} \leq 6g - 6 \prec g\sqrt{A} \frac{t^{\frac{3}{4}}}{\ln t}.$$

Then the conclusion follows by (39), (40), (41) and (42). \square

5.2. Second part of Theorem 31. In this subsection we aim to finish the proof of Theorem 31. For $X_g \in \mathcal{B}_g \subset \mathcal{M}_g$ given in Theorem 32, there exists a universal constant $A > 1$ such that the condition \mathcal{P}_A (24) holds. So using Theorem 33, it remain to estimate $\sum_{\ell(\gamma) < 1, \gamma \in \mathcal{P}(X_g)} \ln \frac{1}{\ell(\gamma)}$ for random hyperbolic surfaces.

Lemma 39.

$$\text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln \frac{1}{\ell(\gamma)} < g \right) = 1 - O\left(\frac{1}{g^{0.99}}\right).$$

Proof. For a hyperbolic surface $X_g \in \mathcal{M}_g$, denote $\{\gamma_1, \dots, \gamma_m\}$ to be the set of all unoriented primitive closed geodesics with length < 1 . By Theorem 13, they are all simple and $m \leq 3g - 3$. By [Mir13, Theorem 4.4],

$$\text{Prob}_{\text{WP}}^g (X_g \in \mathcal{M}_g; \ell_{\text{sys}}^s(X_g) < 0.01 \ln g) = O\left(\frac{(\ln g)^3}{g^{0.995}}\right) = O\left(\frac{1}{g^{0.99}}\right)$$

where $\ell_{\text{sys}}^s(X_g)$ is the length of the shortest simple separating geodesic in X_g . So for sufficiently large g ,

$$(43) \quad \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \begin{array}{l} \gamma_1, \dots, \gamma_m \text{ are all} \\ \text{simple non-separating} \end{array} \right) = 1 - O\left(\frac{1}{g^{0.99}}\right).$$

Then we consider unoriented primitive simple non-separating closed geodesics, which are all in the same orbit of Mod_g . By using Mirzakhani's Integration Formula, i.e., Theorem 8, and Theorem 12 and 10, we have

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g \left[\sum_{\ell(\gamma) < 1}^{n\text{sep}} \ln \frac{1}{\ell(\gamma)} \right] &= \frac{1}{V_g} \frac{1}{2} \int_{\mathbb{R}} \left(\ln \frac{1}{x} \right) \mathbf{1}_{(0,1)}(x) x V_{g-1,2}(x, x) dx \\ &\leq \frac{V_{g-1,2}}{V_g} \int_0^1 \left(\ln \frac{1}{x} \right) \frac{2}{x} (\sinh(x/2))^2 dx \\ &\prec 1 \end{aligned}$$

where the sum $\sum^{n\text{sep}}$ is taken over all unoriented primitive simple non-separating closed geodesics. Then by Markov's inequality we have

$$(44) \quad \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \sum_{\ell(\gamma) < 1}^{n\text{sep}} \ln \frac{1}{\ell(\gamma)} \geq \frac{1}{2}g \right) = O\left(\frac{1}{g}\right).$$

Together with (43) we finish the proof. \square

Now we are ready to prove Theorem 31.

Proof of Theorem 31. Take

$$\mathcal{A}_g = \mathcal{B}_g \cap \left\{ X_g \in \mathcal{M}_g; \sum_{\substack{\gamma \in \mathcal{P}(X_g), \\ \ell(\gamma) < 1}} \ln \frac{1}{\ell(\gamma)} < g \right\}$$

where \mathcal{B}_g is the subset given in Theorem 32. By Theorem 32 and Lemma 39,

$$\text{Prob}_{\text{WP}}^g(\mathcal{A}_g) = 1 - O\left(g^{-\frac{1}{12}}(\ln g)^{\frac{9}{8}}\right).$$

And by Theorem 33 it is clear that for any $X_g \in \mathcal{A}_g$ and $t > 2$,

$$\pi_{X_g}(t) = \text{Li}(t) + \sum_{0 < \lambda_j \leq \frac{1}{4}} \text{Li}(t^{s_j}) + O\left(g \frac{t^{\frac{3}{4}}}{\ln t}\right).$$

The proof is complete. \square

6. INTERSECTION NUMBERS AND WEIL-PETERSSON VOLUMES

In this section, based on Mirzakhani's recursive formula [Mir07a] we prove Theorem 5 and 6, which are essential in the proof of Theorem 4.

6.1. Intersection numbers. For any $d = (d_1, \dots, d_n)$ with each $d_i \in \mathbb{Z}_{\geq 0}$ and $|d| = d_1 + \dots + d_n \leq 3g - 3 + n$, Mirzakhani [Mir07a] gave a recursive formula as follow. Let

$$a_0 = \frac{1}{2}, \quad a_n = \zeta(2n)(1 - 2^{1-2n}) \text{ for } n \geq 1.$$

where $\zeta(\cdot)$ is the Riemann zeta function. Then

$$(45) \quad [\tau_{d_1} \cdots \tau_{d_n}]_{g,n} = \sum_{j=2}^n [\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^{A,j} + [\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^B + [\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^C.$$

where

$$(46) \quad \begin{aligned} [\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^{A,j} &= 8 \sum_{L=0}^{3g-3+n-|d|} (2d_j + 1) a_L \left[\tau_{d_1+d_j+L-1} \prod_{i \neq 1, j} \tau_{d_i} \right]_{g,n-1} \\ &= 8 \sum_{L=d_1+d_j-1}^{3g-4+n-|d|+d_1+d_j} (2d_j + 1) a_{L-d_1-d_2+1} \left[\tau_L \prod_{i \neq 1, j} \tau_{d_i} \right]_{g,n-1}, \end{aligned}$$

$$\begin{aligned}
[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^B &= 16 \sum_{L=0}^{3g-3+n-|d|} \sum_{k_1+k_2=L+d_1-2} a_L \left[\tau_{k_1} \tau_{k_2} \prod_{i \neq 1} \tau_{d_i} \right]_{g-1, n+1} \\
(47) \quad &= 16 \sum_{L=d_1-2}^{3g-5+n-|d|+d_1} \sum_{k_1+k_2=L} a_{L-d_1+2} \left[\tau_{k_1} \tau_{k_2} \prod_{i \neq 1} \tau_{d_i} \right]_{g-1, n+1},
\end{aligned}$$

$$\begin{aligned}
[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^C &= 16 \sum_{I \amalg J = \{2, \dots, n\}} \sum_{g_1+g_2=g} \sum_{L=0}^{3g-3+n-|d|} \sum_{k_1+k_2=L+d_1-2} \\
&\quad a_L \left[\tau_{k_1} \prod_{i \in I} \tau_{d_i} \right]_{g_1, |I|+1} \times \left[\tau_{k_2} \prod_{i \in J} \tau_{d_i} \right]_{g_2, |J|+1} \\
(48) \quad &= 16 \sum_{I \amalg J = \{2, \dots, n\}} \sum_{g_1+g_2=g} \sum_{L=d_1-2}^{3g-5+n-|d|+d_1} \sum_{k_1+k_2=L} \\
&\quad a_{L-d_1+2} \left[\tau_{k_1} \prod_{i \in I} \tau_{d_i} \right]_{g_1, |I|+1} \times \left[\tau_{k_2} \prod_{i \in J} \tau_{d_i} \right]_{g_2, |J|+1}.
\end{aligned}$$

For convenience, here we denote $a_i = 0$ if $i < 0$ and $[\tau_{d_1} \cdots \tau_{d_n}]_{g,n} = 0$ if some index $d_i < 0$.

Mirzakhani showed that

Lemma 40 ([Mir13, Lemma 3.1]). *The sequence $\{a_n\}_{n=0}^\infty$ is increasing. Moreover,*

$$\lim_{n \rightarrow \infty} a_n = 1 \quad \text{and} \quad a_{n+1} - a_n \asymp \frac{1}{2^{2n}}.$$

So clearly we have

$$\begin{aligned}
[\tau_{d_1+1} \cdots \tau_{d_n}]_{g,n}^{A,j} &< [\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^{A,j}, \\
[\tau_{d_1+1} \cdots \tau_{d_n}]_{g,n}^B &< [\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^B, \\
[\tau_{d_1+1} \cdots \tau_{d_n}]_{g,n}^C &< [\tau_{d_1} \cdots \tau_{d_n}]_{g,n}^C.
\end{aligned}$$

These directly yield to

$$[\tau_{d_1+1} \cdots \tau_{d_n}]_{g,n} < [\tau_{d_1} \cdots \tau_{d_n}]_{g,n} \leq V_{g,n}$$

which implies

$$(49) \quad [\tau_{d_1} \cdots \tau_{d_n}]_{g,n} \leq [\tau_0 \tau_{d_2} \cdots \tau_{d_n}]_{g,n} \leq V_{g,n}.$$

Our main result on intersection numbers in this section is as follows. Theorem 5 is as a direct consequence.

Theorem 41. *There exists a universal constant $c > 0$ such that for any $d = (d_1, \dots, d_n)$, assume that $I \neq \emptyset \subset \{1, \dots, n\}$ is the subset consisting of all i 's with $h_i = c \cdot \frac{d_i^2}{2g-2+n} > 1$, then we have*

$$\frac{[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}}{V_{g,n}} \leq \prod_{i \in I} \left(\frac{1}{h_i} \right)^{\frac{1}{2} \log_4(h_i)}.$$

First assuming Theorem 41, we prove Theorem 5.

Proof of Theorem 5. Let $I \subset \{1, \dots, n\}$ be the subset in Theorem 41. If $I = \emptyset$, we have $\frac{2g-2+n}{\max_{1 \leq i \leq n} \{d_i^2\}} \geq c$. Then the conclusion clearly holds by choosing $c' = \frac{1}{c}$ because $[\tau_{d_1} \cdots \tau_{d_n}]_{g,n} \leq V_{g,n}$. If $I \neq \emptyset$, let $i_0 \in [1, n]$ with $d_{i_0}^2 = \max_{1 \leq i \leq n} \{d_i^2\}$. It is clear that the function $f(x) = \left(\frac{1}{x}\right)^{\frac{1}{2} \log_4 x^{-1}}$ is uniformly bounded on $[1, \infty)$. Then it follows by Theorem 41 that

$$(50) \quad \frac{[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}}{V_{g,n}} \leq \left(\frac{1}{h_{i_0}} \right)^{\frac{1}{2} \log_4(h_{i_0})} \leq \frac{c''}{h_{i_0}}$$

for some universal constant $c'' > 0$. Then the conclusion follows by choosing $c' = \frac{c''}{c}$. \square

Now we return to prove Theorem 41.

Proof of Theorem 41. For convenience, we denote $a_i = 0$ if $i < 0$ and $[\tau_{d_1} \cdots \tau_{d_n}]_{g,n} = 0$ if some index $d_i < 0$. For any $d' = (d'_1, \dots, d'_{n+1})$, firstly by (46) we have

$$\begin{aligned} & \left[\tau_{d'_1} \tau_{d'_2} \cdots \tau_{d'_{n+1}} \right]_{g,n+1}^{A,j} - \left[\tau_{d'_{1+1}} \tau_{d'_2} \cdots \tau_{d'_{n+1}} \right]_{g,n+1}^{A,j} \\ &= 8 \sum_{L=d'_1+d'_j-1}^{3g-3+n-\sum_{i \neq 1,j} d'_i} (2d'_j + 1) (a_{L-d'_1-d'_2+1} - a_{L-d'_1-d'_2}) \left[\tau_L \prod_{i \neq 1,j} \tau_{d'_i} \right]_{g,n} \\ &\geq 8(2d'_j + 1) a_0 \left[\tau_{d'_1+d'_j-1} \prod_{i \neq 1,j} \tau_{d'_i} \right]_{g,n}. \end{aligned}$$

So

$$\begin{aligned}
& [\tau_0 \tau_{d'_2} \cdots \tau_{d'_{n+1}}]_{g,n+1}^{A,j} = [\tau_{d'_1} \cdots \tau_{d'_{n+1}}]_{g,n+1}^{A,j} \\
& + \sum_{i=0}^{d'_1-1} \left([\tau_i \tau_{d'_2} \cdots \tau_{d'_{n+1}}]_{g,n+1}^{A,j} - [\tau_{i+1} \tau_{d'_2} \cdots \tau_{d'_{n+1}}]_{g,n+1}^{A,j} \right) \\
(51) \quad & \geq \sum_{i=0}^{d'_1-1} 8(2d'_j + 1)a_0 \left[\tau_{i+d'_j-1} \prod_{i \neq 1,j} \tau_{d'_i} \right]_{g,n} \\
& \geq 8d'_1(2d'_j + 1)a_0 \left[\tau_{d'_1+d'_j-2} \prod_{i \neq 1,j} \tau_{d'_i} \right]_{g,n}.
\end{aligned}$$

And then for any $d'_1 \geq 0$ and $2 \leq j \leq n+1$,

$$(52) \quad [\tau_0 \tau_{d'_2} \cdots \tau_{d'_{n+1}}]_{g,n+1} \geq 8d'_1(2d'_j + 1)a_0 \left[\tau_{d'_1+d'_j-2} \prod_{i \neq 1,j} \tau_{d'_i} \right]_{g,n}.$$

For $d = (d_1, \dots, d_n)$, taking $d'_1 = \lfloor \frac{d_1+2}{2} \rfloor$, $d'_j = \lceil \frac{d_1+2}{2} \rceil$ and $j = 2$ we have

$$\begin{aligned}
(53) \quad [\tau_{d_1} \cdots \tau_{d_n}]_{g,n} & \leq \frac{[\tau_0 \tau_{\lceil \frac{d_1+2}{2} \rceil} \tau_{d_2} \cdots \tau_{d_n}]_{g,n+1}}{8a_0 \lfloor \frac{d_1+2}{2} \rfloor (2 \lceil \frac{d_1+2}{2} \rceil + 1)} \\
& \leq \frac{1}{2d_1^2} [\tau_0 \tau_{\lceil \frac{d_1+2}{2} \rceil} \tau_{d_2} \cdots \tau_{d_n}]_{g,n+1}
\end{aligned}$$

where the fact $a_0 = \frac{1}{2}$ is used in the last inequality. Now we apply this argument for $p_1 \geq 0$ times and each time d_1 is reduced by no more than $\frac{1}{2}$. So combine with (49), we get

$$[\tau_{d_1} \cdots \tau_{d_n}]_{g,n} \leq \prod_{k=1}^{p_1} \frac{1}{2(d_1/2^{k-1})^2} [\tau_0 \cdots \tau_0 \tau_{d_2} \cdots \tau_{d_n}]_{g,n+p_1}.$$

By induction, for any $p = (p_1, \dots, p_n)$,

$$(54) \quad [\tau_{d_1} \cdots \tau_{d_n}]_{g,n} \leq \prod_{i=1}^n \left(\prod_{k_i=1}^{p_i} \frac{1}{2(d_i/2^{k_i-1})^2} \right) V_{g,n+|p|}.$$

By Theorem 10 we know that

$$V_{g,n+|p|} \leq (b(2g - 2 + n + |p|))^{|p|} V_{g,n}$$

for a universal constant $b = \frac{120}{10-\pi^2} \approx 920.276$. So

$$(55) \quad [\tau_{d_1} \cdots \tau_{d_n}]_{g,n} \leq \prod_{i=1}^n \left(\prod_{k_i=1}^{p_i} \frac{b(2g - 2 + n + |p|)}{2(d_i/2^{k_i-1})^2} \right) V_{g,n}.$$

Let $h_i = \frac{2}{3b} \frac{d_i^2}{2g-2+n}$. Assume $I \subset \{1, \dots, n\}$ is the subset consisting of all i 's with $h_i > 1$. For any $i \in [1, n]$, take

$$p_i = \begin{cases} 0 & \text{if } h_i \leq 1 \\ 1 + \lceil \log_4(h_i) \rceil & \text{if } h_i > 1 \end{cases}.$$

Recall that the function $\log_4(x)$ is concave on $(0, \infty)$. In particular,

$$\frac{1}{|I|} \log_4 \left(\prod_{i \in I} d_i \right) \leq \log_4 \left(\frac{\sum_{i \in I} d_i}{|I|} \right).$$

Then we have

$$\begin{aligned} |p| &= \sum_{i \in I} p_i \leq |I| + \log_4 \left(\left(\frac{2}{3b(2g-2+n)} \right)^{|I|} \prod_{i \in I} d_i^2 \right) \\ &\leq |I| + |I| \log_4 \left(\frac{2}{3b(2g-2+n)} \left(\frac{\sum_{i \in I} d_i}{|I|} \right)^2 \right) \\ (56) \quad &\leq |I| + |I| \log_4 \left(\frac{2(3g-3+n)^2}{3b(2g-2+n)} \frac{1}{|I|^2} \right) \\ &\leq |I| + |I| \frac{3g-3+n}{b} \frac{1}{|I|^2} \\ &\leq 2(2g-2+n). \end{aligned}$$

And so for $i \in I$,

$$\begin{aligned} \prod_{k=1}^{p_i} \frac{b(2g-2+n+|p|)}{2(d_i/2^{k-1})^2} &\leq \prod_{k=1}^{p_i} \frac{3b(2g-2+n)}{2(d_i/2^{k-1})^2} \\ (57) \quad &= \prod_{k=1}^{p_i} \frac{4^{k-1}}{h_i} = \frac{4^{\frac{1}{2}(1+\lceil \log_4(h_i) \rceil) \lceil \log_4(h_i) \rceil}}{h_i^{1+\lceil \log_4(h_i) \rceil}} \\ &\leq (h_i)^{-\frac{1}{2} \log_4(h_i)}. \end{aligned}$$

Then the conclusion follows by (55), (56) and (57). \square

6.2. Weil-Petersson volumes. When $\sum x_i^2 = o(2g-2+n)$, for the asymptotic behavior of $V_{g,n}(x_1, \dots, x_n)$, the bounds in Theorem 12 are effective. In this subsection we apply Theorem 41 to prove Theorem 6, which is a more effective upper bound for $V_{g,n}(x_1, \dots, x_n)$ when some of x_i 's are larger than $\sqrt{2g-2+n}$. We first prove the following result.

Theorem 42. *For any $k \geq 1$, there exists a constant $c(k) > 0$ independent of g, n and x_i 's such that*

$$\frac{V_{g,n}(x_1, \dots, x_n)}{V_{g,n}} \leq \left(\prod_{i \in I} c(k) \frac{2g-2+n}{x_i^2} \right)^k \prod_{i=1}^n \frac{\sinh(x_i/2)}{x_i/2}$$

where we assume that $I \neq \emptyset \subset \{1, \dots, n\}$ is the subset consisting of all i 's satisfying that $x_i^2 > c(k)(2g - 2 + n)$.

Proof. As in Theorem 41 we denote $h_i = c \frac{d_i^2}{2g-2+n}$. Then similar as in the proof of Theorem 5 we have that for any fixed $k \geq 1$, $\left(\frac{1}{h_i}\right)^{\frac{1}{2} \log_4(h_i)} \prec_k \left(\frac{2g-2+n}{(2d_i+2k+1)^2}\right)^k$ whenever $h_i > 1$, and $1 \prec_k \left(\frac{2g-2+n}{(2d_i+2k+1)^2}\right)^k$ whenever $h_i \leq 1$. So combine with Theorem 41 we have that for any subset $I' \subset \{1, \dots, n\}$,

$$(58) \quad \frac{[\tau_{d_1} \cdots \tau_{d_n}]_{g,n}}{V_{g,n}} \leq \prod_{i \in I'} \left(c'(k) \frac{2g-2+n}{(2d_i+2k+1)^2} \right)^k$$

for some $c'(k) > 0$. Then it follows by Theorem 9 that

$$(59) \quad \begin{aligned} \frac{V_{g,n}(2x_1, \dots, 2x_n)}{V_{g,n}} &\leq \sum_{|d| \leq 3g-3+n} \prod_{i \in I'} \left(c'(k) \frac{2g-2+n}{(2d_i+2k+1)^2} \right)^k \prod_{i=1}^n \frac{x_i^{2d_i}}{(2d_i+1)!} \\ &\leq (c'(k)(2g-2+n))^{k|I'|} \cdot \left(\prod_{i \in I'} \sum_{d_i=0}^{\infty} \frac{x_i^{2d_i}}{(2d_i+2k+1)!} \right) \cdot \left(\prod_{i \notin I'} \sum_{d_i=0}^{\infty} \frac{x_i^{2d_i}}{(2d_i+1)!} \right). \end{aligned}$$

By Taylor's expansion,

$$\sum_{d=0}^{\infty} \frac{x^{2d}}{(2d+1)!} = \frac{\sinh(x)}{x}$$

and hence

$$\sum_{d=0}^{\infty} \frac{x^{2d}}{(2d+2k+1)!} \leq \frac{1}{x^{2k}} \sum_{d=0}^{\infty} \frac{x^{2d}}{(2d+1)!} = \frac{\sinh(x)}{x^{2k+1}}.$$

Then again by (59),

$$(60) \quad \frac{V_{g,n}(2x_1, \dots, 2x_n)}{V_{g,n}} \leq \left(\prod_{i \in I'} c'(k) \frac{2g-2+n}{x_i^2} \right)^k \prod_{i=1}^n \frac{\sinh(x_i)}{x_i}$$

for any subset $I' \subset \{1, \dots, n\}$. Now let $I' = I$, then the conclusion follows by choosing $c(k) = 4c'(k)$. The proof is complete. \square

Now we complete the proof of Theorem 6.

Proof of Theorem 6. Recall that the upper bound in Theorem 12 says that

$$(61) \quad \frac{V_{g,n}(x_1, \dots, x_n)}{V_{g,n}} \leq \prod_{i=1}^n \frac{\sinh(x_i/2)}{x_i/2}.$$

Then the conclusion directly follows by (61) and Theorem 42. \square

7. PROOF OF THEOREM 4

In this section we prove Theorem 4 confirming [LW21, Conjecture 1.2].

Similar as [NWX20, WX22], we first make the following notations. Let $X_g \in \mathcal{M}_g$ be a closed hyperbolic surface of genus g and $\mathcal{P}'(X_g)$ be the set of all unoriented primitive closed geodesics on X_g . For any $L > 0$, we define

- (1) $N(X_g, L) := \#\{\gamma \in \mathcal{P}'(X_g); \ell_\gamma(X_g) \leq L\}$,
- (2) $N^s(X_g, L) := \#\{\gamma \in \mathcal{P}'(X_g) \text{ is simple}; \ell_\gamma(X_g) \leq L\}$,
- (3) $N^{ns}(X_g, L) := \#\{\gamma \in \mathcal{P}'(X_g) \text{ is non-simple}; \ell_\gamma(X_g) \leq L\}$,
- (4) $N_{nsep}^s(X_g, L) := \#\{\gamma \in \mathcal{P}'(X_g) \text{ is simple and non-separating}; \ell_\gamma(X_g) \leq L\}$,
- (5) $N_{sep,k}^s(X_g, L) := \#\{\gamma \in \mathcal{P}'(X_g) \text{ and } X_g \setminus \gamma \cong S_{k,1} \cup S_{g-k,1}; \ell_\gamma(X_g) \leq L\}$ where $1 \leq k \leq \lfloor \frac{g}{2} \rfloor$.




FIGURE 1. Examples: α is simple and non-separating; β is simple and separating; γ is non-simple.

It is clear that

$$N(X_g, L) = \frac{\pi X_g(e^L)}{2}$$

and

$$\begin{aligned} N(X_g, L) &= N^s(X_g, L) + N^{ns}(X_g, L) \\ (62) \quad &= N_{nsep}^s(X_g, L) + \sum_{k=1}^{\lfloor \frac{g}{2} \rfloor} N_{sep,k}^s(X_g, L) + N^{ns}(X_g, L). \end{aligned}$$

We split the proof of Theorem 4 into three parts.

7.1. Significantly greater than \sqrt{g} . In this subsection we confirm the second part of [LW21, Conjecture 1.2]. That is, we show that as $g \rightarrow \infty$, on most surfaces in \mathcal{M}_g most geodesics of length significantly greater than \sqrt{g} are non-simple. More precisely,

Theorem 43. *Assume that $L = L(g)$ satisfies*

$$\lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = \infty.$$

Then for any $k > 0 \in \mathbb{Z}$, we have

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N^{ns}(X_g, L)}{N(X_g, L)} \right| < \left(\frac{g}{L^2} \right)^k \right) = 1.$$

We first begin with the following easy consequence of Theorem 6.

Lemma 44. *For any $x > 0$ and $k > 0 \in \mathbb{Z}$, we have that for all $g \geq 2$ and $1 \leq i \leq \lfloor \frac{g}{2} \rfloor$,*

$$\frac{V_{g-1,2}(x, x)}{V_{g-1,2}} \prec_k \left(\frac{g}{1+x^2} \right)^{k+1} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2$$

and

$$\frac{V_{g-i,1}(x) \cdot V_{i,1}(x)}{V_{g-i,1} \cdot V_{i,1}} \prec_k \left(\frac{g}{1+x^2} \right)^{k+1} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2.$$

Proof. By Theorem 6 we have

$$\begin{aligned} \frac{V_{g-1,2}(x, x)}{V_{g-1,2}} &\leq \left(\min \left\{ c(k+1) \frac{2g-2}{x^2}, 1 \right\} \right)^{2k+2} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2 \\ &\leq \left(\min \left\{ c(k+1) \frac{2g-2}{x^2}, 1 \right\} \right)^{k+1} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2 \\ &\prec_k \left(\frac{g}{1+x^2} \right)^{k+1} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2. \end{aligned}$$

Now we prove the second inequality. Recall that the upper bound in Theorem 12 implies that $\frac{V_{i,1}(x)}{V_{i,1}} \leq \frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}}$. Similarly, then it follows by Theorem 6 that for all $1 \leq i \leq \lfloor \frac{g}{2} \rfloor$,

$$\begin{aligned} \frac{V_{g-i,1}(x) \cdot V_{i,1}(x)}{V_{g-i,1} \cdot V_{i,1}} &\leq \frac{V_{g-i,1}(x)}{V_{g-i,1}} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right) \\ &\leq \left(\min \left\{ c(k+1) \frac{2g-2i-1}{x^2}, 1 \right\} \right)^{k+1} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2 \\ &\prec_k \left(\frac{g}{1+x^2} \right)^{k+1} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2. \end{aligned}$$

The proof is complete. \square

Lemma 45. *Assume that $L = L(g)$ satisfies*

$$\lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = \infty.$$

Then for any $k > 0 \in \mathbb{Z}$,

$$\mathbb{E}_{\text{WP}}^g [N_{nsep}^s(X_g, L)] \prec_k \text{Li}(e^L) \cdot \left(\frac{g}{L^2}\right)^{k+1}.$$

Proof. It follows by Mirzakhani's integration formula, i.e., Theorem 8, and Lemma 44 that

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g [N_{nsep}^s(X_g, L)] &= \frac{1}{V_g} \frac{1}{2} \int_0^L V_{g-1,2}(x, x) dx \\ &\prec_k \frac{V_{g-1,2}}{V_g} \cdot \int_0^L \left(\sinh\left(\frac{x}{2}\right)\right)^2 \cdot \frac{1}{x} \cdot \left(\frac{g}{1+x^2}\right)^{k+1} dx. \end{aligned}$$

By Theorem 10 we also know that

$$\frac{V_g}{V_{g-1,2}} = 1 + O\left(\frac{1}{g}\right).$$

Thus,

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g [N_{nsep}^s(X_g, L)] &\prec_k \left(1 + O\left(\frac{1}{g}\right)\right) \cdot \int_0^L \left(\sinh\left(\frac{x}{2}\right)\right)^2 \frac{1}{x} \left(\frac{g}{1+x^2}\right)^{k+1} dx \\ &\asymp \text{Li}(e^L) \cdot \left(\frac{g}{L^2}\right)^{k+1} \end{aligned}$$

where we apply $\lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = \infty$ in the last equation. The proof is complete. \square

Lemma 46. *Assume that $L = L(g)$ satisfies*

$$\lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = \infty.$$

Then for any $k > 0 \in \mathbb{Z}$,

$$\sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \mathbb{E}_{\text{WP}}^g [N_{sep,i}^s(X_g, L)] \prec_k \frac{\text{Li}(e^L)}{g} \cdot \left(\frac{g}{L^2}\right)^{k+1}.$$

Proof. It follows by Theorem 8 and Lemma 44 that

$$\begin{aligned} \sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \mathbb{E}_{\text{WP}}^g [N_{sep,i}^s(X_g, L)] &\leq \frac{1}{V_g} \sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \int_0^L V_{g-i,1}(x) V_{i,1}(x) dx \\ &\prec_k \left(\sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \frac{V_{g-i,1} V_{i,1}}{V_g} \right) \cdot \int_0^L \left(\sinh\left(\frac{x}{2}\right)\right)^2 \cdot \frac{1}{x} \cdot \left(\frac{g}{1+x^2}\right)^{k+1} dx. \end{aligned}$$

By Theorem 11 we also know that

$$\sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \frac{V_{g-i,1} V_{i,1}}{V_g} \asymp \frac{1}{g}.$$

Thus,

$$\begin{aligned} \sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \mathbb{E}_{\text{WP}}^g [N_{\text{sep},i}^s(X_g, L)] &\prec_k \frac{1}{g} \cdot \int_0^L \left(\sinh\left(\frac{x}{2}\right) \right)^2 \frac{1}{x} \left(\frac{g}{1+x^2} \right)^{k+1} dx \\ &\asymp \frac{\text{Li}(e^L)}{g} \cdot \left(\frac{g}{L^2} \right)^{k+1} \end{aligned}$$

where we apply $\lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = \infty$ in the last equation. The proof is complete. \square

Now we are ready to prove Theorem 43.

Proof of Theorem 43. First by Lemma 45 and Lemma 46 we know that for any $k > 0 \in \mathbb{Z}$,

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g [N^s(X_g, L)] &= \mathbb{E}_{\text{WP}}^g [N_{\text{nsep}}^s(X_g, L)] + \sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \mathbb{E}_{\text{WP}}^g [N_{\text{sep},i}^s(X_g, L)] \\ &\prec_k \text{Li}(e^L) \cdot \left(\frac{g}{L^2} \right)^{k+1}. \end{aligned}$$

Then by Markov's inequality we have

$$\begin{aligned} (63) \quad \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; N^s(X_g, L) \geq \frac{1}{4} \left(\frac{g}{L^2} \right)^k \cdot \text{Li}(e^L) \right) \\ \prec_k \frac{g}{L^2} \rightarrow 0 \text{ as } g \rightarrow \infty. \end{aligned}$$

Recall that $N(X_g, L) = \frac{\pi X_g(e^L)}{2}$. Let \mathcal{A}_g be in Theorem 31. Then we have

$$(64) \quad \lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{A}_g) = 1$$

and for any $X_g \in \mathcal{A}_g$ and large enough $g > 0$,

$$(65) \quad N(X_g, L) \geq \frac{1}{2} \cdot \text{Li}(e^L) \cdot \left(1 + O\left(\frac{g}{e^{\frac{L}{4}}}\right) \right) \geq \frac{\text{Li}(e^L)}{4}$$

where we apply $\lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = \infty$ in the last inequality. Since $N^s(X_g, L) = N(X_g, L) - N^{\text{nsep}}(X_g, L)$, it follows by (63) and (65) that for large enough

$g > 0$,

$$\begin{aligned} & \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N^{ns}(X_g, L)}{N(X_g, L)} \right| \geq \left(\frac{g}{L^2} \right)^k \right) \leq \text{Prob}_{\text{WP}}^g (X_g \notin \mathcal{A}_g) \\ & \quad + \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; N^s(X_g, L) \geq \frac{1}{4} \left(\frac{g}{L^2} \right)^k \cdot \text{Li}(e^L) \right) \\ & \prec_k \left((1 - \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{A}_g)) + \frac{g}{L^2} \right). \end{aligned}$$

By letting $g \rightarrow \infty$, then the conclusion follows by (64) and our assumption on $L = L(g)$. The proof is complete. \square

7.2. Significantly less than \sqrt{g} and larger than $12 \ln g$. In this and next subsection we confirm the first part of [LW21, Conjecture 1.2]. That is, we show that as $g \rightarrow \infty$, on most surfaces in \mathcal{M}_g most geodesics of length significantly less than \sqrt{g} are simple and non-separating. The proof is splitted into two subcases: when $L(g) > 12 \ln g$, we apply the spectral method in this paper; when $L(g) \leq 12 \ln g$, we apply our previous works [NWX20, WX22].

In this subsection we prove

Theorem 47. *Assume that $L = L(g)$ satisfies*

$$L(g) > 12 \ln g \text{ and } \lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = 0.$$

Then for any $\delta = \delta(g) > 0$ satisfying $\lim_{g \rightarrow \infty} \delta(g) = 0$ and $\lim_{g \rightarrow \infty} \frac{1}{\delta} \left(\frac{L^2}{g} + \frac{1}{g^{1-\epsilon_0}} \right) = 0$ for some $\epsilon_0 \in (0, 1)$, we have

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N_{nsep}^s(X_g, L)}{N(X_g, L)} \right| < \delta \right) = 1.$$

We first make the following lemma.

Lemma 48. *Assume that $L = L(g)$ satisfies*

$$\lim_{g \rightarrow \infty} L(g) = \infty \text{ and } \lim_{g \rightarrow \infty} \frac{L^2(g)}{g} = 0.$$

Then

$$\mathbb{E}_{\text{WP}}^g [N_{nsep}^s(X_g, L)] = \frac{\text{Li}(e^L)}{2} \cdot \left(1 + O \left(\frac{L^2}{g} + \frac{L \ln L}{e^L} \right) \right)$$

where the implied constant is universal.

Proof. First by Mirzakhani's integration formula, i.e., Theorem 8, we have

$$\mathbb{E}_{\text{WP}}^g [N_{nsep}^s(X_g, L)] = \frac{1}{V_g} \frac{1}{2} \int_0^L V_{g-1,2}(x, x) x dx.$$

Since $L = L(g) = o(\sqrt{g})$, by Theorem 12 we have that for all $x \in [0, L]$,

$$V_{g-1,2}(x, x) = V_{g-1,2} \cdot \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2 \cdot \left(1 + O\left(\frac{x^2}{g}\right) \right)$$

where the implied constant is universal. By Theorem 10 we also know that

$$\frac{V_g}{V_{g-1,2}} = 1 + O\left(\frac{1}{g}\right).$$

Thus,

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g [N_{nsep}^s(X_g, L)] &= \int_0^L \left(\sinh\left(\frac{x}{2}\right) \right)^2 \cdot \frac{2}{x} \cdot \left(1 + O\left(\frac{x^2}{g}\right) \right) \\ &\cdot \left(1 + O\left(\frac{1}{g}\right) \right) dx = \frac{\text{Li}(e^L)}{2} \cdot \left(1 + O\left(\frac{L^2}{g} + \frac{L \ln L}{e^L}\right) \right) \end{aligned}$$

where we apply $\lim_{g \rightarrow \infty} L = \lim_{g \rightarrow \infty} L(g) = \infty$ in the last equality.

The proof is complete. \square

Now we are ready to prove Theorem 47.

Proof of Theorem 47. Recall that $N(X_g, L) = \frac{\pi_{X_g}(e^L)}{2}$. Since $L > 12 \ln g$, it follows by Theorem 3 that for any $\epsilon \in (0, 1)$,

$$\mathbb{E}_{\text{WP}}^g [N(X_g, L)] = \frac{\mathbb{E}_{\text{WP}}^g [\pi_{X_g}(e^L)]}{2} = \frac{\text{Li}(e^L)}{2} \cdot \left(1 + O_\epsilon\left(\frac{1}{g^{1-\epsilon}}\right) \right)$$

which together with Lemma 48 imply that

$$\mathbb{E}_{\text{WP}}^g [|N(X_g, L) - N_{nsep}^s(X_g, L)|] = \text{Li}(e^L) \cdot O_\epsilon\left(\frac{L^2}{g} + \frac{1}{g^{1-\epsilon}}\right).$$

Now we choose $\epsilon = \epsilon_0$. Then by Markov's inequality we have

$$\begin{aligned} (66) \quad &\text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; |N(X_g, L) - N_{nsep}^s(X_g, L)| \geq \frac{\delta \cdot \text{Li}(e^L)}{4} \right) \\ &= O_{\epsilon_0} \left(\frac{1}{\delta} \left(\frac{L^2}{g} + \frac{1}{g^{1-\epsilon_0}} \right) \right) \rightarrow 0 \text{ as } g \rightarrow \infty. \end{aligned}$$

Let \mathcal{A}_g be in Theorem 31. Then we have

$$(67) \quad \lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{A}_g) = 1$$

and for any $X_g \in \mathcal{A}_g$ and large enough $g > 0$,

$$(68) \quad N(X_g, L) \geq \frac{1}{2} \cdot \text{Li}(e^L) \cdot \left(1 + O\left(\frac{g}{e^{\frac{L}{4}}}\right) \right) \geq \frac{\text{Li}(e^L)}{4}$$

where we apply $L > 12 \ln g$ in the last inequality. Thus, it follows by (66) and (68) that for large enough $g > 0$,

$$\begin{aligned} & \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N_{nsep}^s(X_g, L)}{N(X_g, L)} \right| \geq \delta \right) \leq \text{Prob}_{\text{WP}}^g (X_g \notin \mathcal{A}_g) \\ & + \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; |N(X_g, L) - N_{nsep}^s(X_g, L)| \geq \frac{\delta \cdot \text{Li}(e^L)}{4} \right) \\ & = (1 - \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{A}_g)) + O_{\epsilon_0} \left(\frac{1}{\delta} \left(\frac{L^2}{g} + \frac{1}{g^{1-\epsilon_0}} \right) \right). \end{aligned}$$

By letting $g \rightarrow \infty$, then the conclusion follows by (67) and our assumption on δ . The proof is complete. \square

7.3. No more than $12 \ln g$. As mentioned in the introduction, it follows by [Mir13, Theorem 4.4] and [NWX20, Theorem 4] that if $L = L(g) \leq (1 - \epsilon_0) \ln g$ for some constant $\epsilon_0 > 0$, then

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{M}_g; N(X_g, L) = N_{nsep}^s(X_g, L)) = 1.$$

In this subsection we prove

Theorem 49. *Assume $L = L(g)$ satisfies that for some $\epsilon_0 \in (0, 1)$,*

$$L(g) \geq (1 - \epsilon_0) \ln g \quad \text{and} \quad \limsup_{g \rightarrow \infty} \frac{L(g)}{\ln g} < \infty.$$

Then for any fixed $\epsilon > 0$ we have

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N_{nsep}^s(X_g, L)}{N(X_g, L)} \right| < \frac{1}{g^{1-\epsilon}} \right) = 1.$$

Recall that

$$N(X_g, L) = N_{nsep}^s(X_g, L) + \sum_{k=1}^{\lfloor \frac{g}{2} \rfloor} N_{sep,k}^s(X_g, L) + N^{ns}(X_g, L).$$

We will deal with the three terms on the RHS above separately.

Expected value on simple and separating closed geodesics. Similar as Lemma 46 we first show that

Lemma 50. *For any $L = L(g)$ with $\lim_{g \rightarrow \infty} L(g) = \infty$, we have*

$$\sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \mathbb{E}_{\text{WP}}^g [N_{sep,i}^s(X_g, L)] \prec \frac{\text{Li}(e^L)}{g}.$$

Proof. It follows by Theorem 8 and 12 that

$$\begin{aligned} \sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \mathbb{E}_{\text{WP}}^g [N_{\text{sep},i}^s(X_g, L)] &\leq \frac{1}{V_g} \sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \int_0^L V_{g-i,1}(x) V_{i,1}(x) dx \\ &\leq \left(\sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \frac{V_{g-i,1} V_{i,1}}{V_g} \right) \cdot \int_0^L \left(\sinh \left(\frac{x}{2} \right) \right)^2 \cdot \frac{4}{x} dx \\ &\asymp \left(\sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \frac{V_{g-i,1} V_{i,1}}{V_g} \right) \cdot \text{Li}(e^L) \end{aligned}$$

where we apply $\lim_{g \rightarrow \infty} L(g) = \infty$ in the last equation. By Theorem 11 we also know that

$$\sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \frac{V_{g-i,1} V_{i,1}}{V_g} \asymp \frac{1}{g}.$$

Thus, we get

$$\sum_{i=1}^{\lfloor \frac{g}{2} \rfloor} \mathbb{E}_{\text{WP}}^g [N_{\text{sep},i}^s(X_g, L)] \prec \frac{\text{Li}(e^L)}{g}$$

as desired. \square

Expected value on non-simple closed geodesics. For $N^{ns}(X_g, L)$ on non-simple closed geodesics, we use our previous work [WX22, Section 7]. Since the proofs here are almost identical to the ones in [WX22, Section 7], we only outline them. One may see [WX22, Section 7] for more details. More precisely, we show

Theorem 51. *Assume that $L = L(g) \leq 12 \ln g$ and $\lim_{g \rightarrow \infty} L(g) = \infty$. Then for any $\epsilon > 0$, as $g \rightarrow \infty$ we have*

$$\mathbb{E}_{\text{WP}}^g [N^{ns}(X_g, L)] \prec_{\epsilon} \frac{e^{(1+\epsilon)L}}{g}.$$

We use $X = X_g \in \mathcal{M}_g$ for simplicity. Consider Γ to be either a non-simple closed geodesic $\gamma \subset X$ or a pair of distinct closed geodesics (γ_1, γ_2) in X satisfying $\gamma_1 \cap \gamma_2 \neq \emptyset$. For the later case, denote the length of Γ to be the total length $\ell_{\Gamma}(X) := \ell_{\gamma_1}(X) + \ell_{\gamma_2}(X)$.

As in [WX22, Section 7], for any $T = T(g) > 0$ and Γ as above of length $\ell_{\Gamma}(X) \leq T$, let $X(\Gamma) \subset X$ be the connected subsurface of geodesic boundary (we warn here that two distinct simple closed geodesics on the boundary of $X(\Gamma)$ may correspond to a single simple closed geodesic in X) such that

- (1) $\Gamma \subset X(\Gamma)$ is filling;
- (2) $\ell(\partial X(\Gamma)) \leq 2\ell_{\Gamma}(X) \leq 2T$;
- (3) $\text{Area}(X(\Gamma)) \leq 4\ell_{\Gamma}(X) \leq 4T$.

Definition. For $X \in \mathcal{M}_g$ and $0 < T = T(g) \leq \frac{\text{Area}(X)}{8}$, we define

$$\text{Sub}_T(X) := \left\{ Y \subset X \text{ is a connected subsurface of geodesic} \right. \\ \left. \text{boundary such that } \ell(\partial Y) \leq 2T \text{ and } \text{Area}(Y) \leq 4T \right\}$$

where we allow two distinct simple closed geodesics on the boundary of Y to be a single simple closed geodesic in X .

As in [WX22, Section 7], let $\mathcal{P}^{ns}(X)$ be the set of oriented primitive non-simple closed geodesics of X . And let $\mathcal{P}^{ns,2}(X)$ be the set of pairs of oriented primitive closed geodesics (γ_1, γ_2) with $\gamma_1 \cap \gamma_2 \neq \emptyset$ on X . Consider the composition \mathcal{F} of the following two maps

$$\Gamma \mapsto X(\Gamma) \mapsto \partial X(\Gamma),$$

then we have that for $\gamma \in \mathcal{P}^{ns}(X)$ of length $\ell_\gamma(X) \leq T$,

$$(69) \quad \# \left\{ \gamma' \in \mathcal{P}^{ns}(X); \begin{array}{l} \ell_{\gamma'}(X) \leq T \\ \text{and } \mathcal{F}(\gamma') = \mathcal{F}(\gamma) \end{array} \right\} \leq 2\#_f(X(\gamma), T)$$

where $\#_f(X(\gamma), T)$ is the number of filling (unoriented) closed geodesics in $X(\gamma)$ of length $\leq T$. And for $\Gamma \in \mathcal{P}^{ns,2}(X)$ of length $\ell_\Gamma(X) \leq T$,

$$(70) \quad \# \left\{ \Gamma' \in \mathcal{P}^{ns,2}(X); \begin{array}{l} \ell_{\Gamma'}(X) \leq T \\ \text{and } \mathcal{F}(\Gamma') = \mathcal{F}(\Gamma) \end{array} \right\} \leq 4N_2^{\text{fill}}(X(\Gamma), T)$$

where $N_2^{\text{fill}}(X(\Gamma), T)$ is defined in Definition 17.

Denote

$$N^{ns,2}(X, T) := \frac{1}{4} \# \{ \Gamma \in \mathcal{P}^{ns,2}(X); \ell_\Gamma(X) \leq T \}$$

to be the number of pairs of intersected unoriented closed geodesics (γ_1, γ_2) of X of total length $\ell_{\gamma_1}(X) + \ell_{\gamma_2}(X) \leq T$. Similar to [WX22, Proposition 30], we first show that

Proposition 52. *For any $\epsilon_1 > 0$ and $M \in \mathbb{Z}_{\geq 1}$, there exists a constant $c(\epsilon_1, M) > 0$ only depending on ϵ_1 and M such that as $g \rightarrow \infty$,*

$$(71) \quad \max \left\{ \begin{array}{l} N^{ns}(X, T), \\ N^{ns,2}(X, T) \end{array} \right\} \prec T^3 e^T \left(\sum_{\substack{Y \in \text{Sub}_T(X); \\ |\chi(Y)| \geq M+1}} \mathbf{1}_{[0,2T]}(\ell(\partial Y)) \right) \\ + c(\epsilon_1, M) \left(\sum_{\substack{Y \in \text{Sub}_T(X); \\ 1 \leq |\chi(Y)| \leq M}} T e^{T - \frac{1-\epsilon_1}{2} \ell(\partial Y)} \mathbf{1}_{[0,2T]}(\ell(\partial Y)) \right).$$

Proof. By Collar Lemma, either a non-simple closed geodesic or a pair of intersecting closed geodesics has length $> \text{arcsinh } 1$. So it suffices to consider

for large T . Rewrite

$$\begin{aligned}
(72) \quad N^{ns}(X, T) &= \frac{1}{2} \cdot \sum_{\gamma \in \mathcal{P}^{ns}(X)} \mathbf{1}_{[0, T]}(\ell_\gamma(X)) \\
&= \frac{1}{2} \times \sum_{\substack{Y \in \text{Sub}_T(X); \\ |\chi(Y)| \geq M+1}} \sum_{\gamma \subset Y \text{ is filling}} \mathbf{1}_{[0, T]}(\ell_\gamma(X)) \\
&\quad + \frac{1}{2} \times \sum_{\substack{Y \in \text{Sub}_T(X); \\ 1 \leq |\chi(Y)| \leq M}} \sum_{\gamma \subset Y \text{ is filling}} \mathbf{1}_{[0, T]}(\ell_\gamma(X)).
\end{aligned}$$

Now we consider the first term in the RHS of (72). Since $\gamma \subset Y$ is filling,

$$(73) \quad \frac{\ell(\partial Y)}{2} \leq \ell_\gamma(Y) = \ell_\gamma(X).$$

For $Y \in \text{Sub}_T(X)$ we know that $\text{Area}(Y) \leq 4T$. So by [WX22, Lemma 10] we have

$$(74) \quad \#_f(Y, T) \prec T e^T.$$

By (73) and (74) we have

$$\begin{aligned}
(75) \quad &\sum_{\substack{Y \in \text{Sub}_T(X); \\ |\chi(Y)| \geq M+1}} \sum_{\gamma \subset Y \text{ is filling}} \mathbf{1}_{[0, T]}(\ell_\gamma(X)) \\
&\leq \sum_{\substack{Y \in \text{Sub}_T(X); \\ |\chi(Y)| \geq M+1}} \#_f(Y, T) \mathbf{1}_{[0, 2T]}(\ell(\partial Y)) \\
&\prec T e^T \left(\sum_{\substack{Y \in \text{Sub}_T(X); \\ |\chi(Y)| \geq M+1}} \mathbf{1}_{[0, 2T]}(\ell(\partial Y)) \right).
\end{aligned}$$

Now we consider the second term in the RHS of (72). Let $Y \in \text{Sub}_T(X)$ with $m = |\chi(Y)| \in [1, M]$. For any $\epsilon_1 > 0$, let $c'(\epsilon_1, M) = \max_{1 \leq m \leq M} c(\epsilon_1, m) > 0$ where $c(\epsilon_1, m) > 0$ is the constant in Theorem 16. Then we have that for large enough g ,

$$(76) \quad \sum_{\gamma \subset Y \text{ is filling}} \mathbf{1}_{[0, T]}(\ell_\gamma(X)) \prec c'(\epsilon_1, M) e^{T - \frac{1-\epsilon_1}{2} \ell(\partial Y)} \mathbf{1}_{[0, 2T]}(\ell(\partial Y))$$

which implies that

$$(77) \quad \sum_{\substack{Y \in \text{Sub}_T(X); \\ 1 \leq |\chi(Y)| \leq M}} \sum_{\gamma \subset Y \text{ is filling}} \mathbf{1}_{[0,T]}(\ell_\gamma(X)) \\ \prec c'(\epsilon_1, M) \left(\sum_{\substack{Y \in \text{Sub}_T(X); \\ 1 \leq |\chi(Y)| \leq M}} e^{T - \frac{1-\epsilon_1}{2} \ell(\partial Y)} \mathbf{1}_{[0,2T]}(\ell(\partial Y)) \right).$$

Then the conclusion for $N^{ns}(X, T)$ follows by (72), (75) and (77).

For $N^{ns,2}(X, T)$, the proof is identical as above through replacing (74) by

$$N_2^{\text{fill}}(Y, T) \prec T^3 e^T$$

obtained from Lemma 64 in Appendix A, and replacing (76) by

$$\sum_{\Gamma \subset Y \text{ is filling}} \mathbf{1}_{[0,T]}(\ell_\Gamma(X)) \prec c'(\epsilon_1, M) T e^{T - \frac{1-\epsilon_1}{2} \ell(\partial Y)} \mathbf{1}_{[0,2T]}(\ell(\partial Y))$$

obtained from Theorem 18. \square

Now we follow [WX22] to estimate the expected values $\mathbb{E}_{\text{WP}}^g [N^{ns}(X, T)]$ and $\mathbb{E}_{\text{WP}}^g [N^{ns,2}(X, T)]$ through taking an integral of (71) in Proposition 52 over \mathcal{M}_g . The proofs are identical to the ones in [WX22, Section 7]. So we only point out the necessary modifications. First we recall the Assumption (\star) in [WX22, Section 7.2]:

Assumption (\star) . let $Y_0 \in \text{Sub}_T(X)$ satisfying

- Y_0 is homeomorphic to $S_{g_0, k}$ for some $g_0 \geq 0$ and $k > 0$ with $m = |\chi(Y_0)| = 2g_0 - 2 + k \geq 1$;
- the boundary ∂Y_0 is a simple closed multi-geodesics in X consisting of k simple closed geodesics which has n_0 pairs of simple closed geodesics for some $n_0 \geq 0$ such that each pair corresponds to a single simple closed geodesic in X ;
- the interior of its complement $X \setminus Y_0$ consists of q components $S_{g_1, n_1}, \dots, S_{g_q, n_q}$ for some $q \geq 1$ where $\sum_{i=1}^q n_i = k - 2n_0$.

First we study a single orbit $\text{Mod}_g \cdot Y_0 \subset \text{Sub}_T(X)$.

Proposition 53. *Let $Y_0 \in \text{Sub}_T(X)$ satisfying Assumption (\star) . Then we have that as $g \rightarrow \infty$,*

$$\int_{\mathcal{M}_g} \sum_{Y \in \text{Mod}_g \cdot Y_0} \mathbf{1}_{[0,2T]}(\ell(\partial Y)) dX \prec e^{4T} V_{g_0, k} \frac{V_{g_1, n_1} \cdots V_{g_q, n_q}}{n_0! n_1! \cdots n_q!}.$$

Proof. The proof is identical to the proof of [WX22, Proposition 31] through replacing $e^{-\frac{\ell(\partial Y)}{4}}$ by 1. \square

As in [NWX20, WX22], for $r \geq 1$ we define

$$W_r := \begin{cases} V_{\frac{r}{2}+1} & \text{if } r \text{ is even,} \\ V_{\frac{r+1}{2},1} & \text{if } r \text{ is odd.} \end{cases}$$

Proposition 54. *Let $g_0 \geq 0$ and $k \geq 1$ be fixed with $m = 2g_0 - 2 + k \geq 1$. Then we have that as $g \rightarrow \infty$,*

$$\int_{\mathcal{M}_g} \sum_{\substack{Y \in \text{Sub}_T(X); \\ Y \cong S_{g_0, k}}} \mathbf{1}_{[0, 2T]}(\ell(\partial Y)) dX \prec e^{4T} W_{2g_0+k-2} W_{2g-2g_0-k}.$$

Proof. The proof is identical to the one of [WX22, Proposition 32] through replacing $e^{-\frac{\ell(\partial Y)}{4}}$ by 1 and applying Proposition 53. \square

For any $Y \in \text{Sub}_T(X)$ it is known that $|\chi(Y)| \leq \frac{4T}{2\pi}$. Now we bound the integral of $\sum_{Y \in \text{Sub}_T(X)} \mathbf{1}_{[0, 2T]}(\ell(\partial Y))$ over \mathcal{M}_g when $m \leq |\chi(Y)| \leq [\frac{4T}{2\pi}]$ where $m \geq 1$ is a fixed integer. That is, we allow g_0 and k in Assumption (\star) vary such that $m \leq 2g_0 + k - 2 \leq [\frac{4T}{2\pi}]$.

Proposition 55. *Let $m \geq 1$ be any fixed integer. Then we have that there exists a constant $c(m) > 0$ only depending on m such that as $g \rightarrow \infty$,*

$$\int_{\mathcal{M}_g} \sum_{\substack{Y \in \text{Sub}_T(X); \\ m \leq |\chi(Y)| \leq [\frac{4T}{2\pi}]}} \mathbf{1}_{[0, 2T]}(\ell(\partial Y)) dX \prec T e^{4T} c(m) \frac{V_g}{g^m}.$$

Proof. The proof is identical to the proof of [WX22, Proposition 33] through replacing $e^{-\frac{\ell(\partial Y)}{4}}$ by 1 and applying Proposition 54. \square

For the expected value of the RHS in (71), Proposition 55 implies that if $T \leq C \ln g$ for some constant $C > 0$ and m large enough, then we have

$$\frac{T^3 e^T}{V_g} \int_{\mathcal{M}_g} \sum_{\substack{Y \in \text{Sub}_T(X); \\ m \leq |\chi(Y)| \leq [\frac{4T}{2\pi}]}} \mathbf{1}_{[0, 2T]}(\ell(\partial Y)) dX \prec_{C, m} \frac{e^T}{g}.$$

Next we aim to show the expected value of the second term of the RHS of (71) in Proposition 52 also has the similar property for bounded m .

In the sequel, we always assume that $Y_0 \cong S_{g_0, k} \in \text{Sub}_T(X)$ satisfies Assumption (\star) with an additional assumption that

$$1 \leq 2g_0 + k - 2 \leq M$$

for a constant $M \in \mathbb{Z}_{\geq 1}$. Then $1 \leq k \leq M + 2$ and $0 \leq g_0 \leq \frac{1}{2}(M + 1)$.

Proposition 56. *Let $g_0 \geq 0$ and $k \geq 1$ be fixed with $m = 2g_0 - 2 + k$. For any $\epsilon_1 > 0$, then we have that as $g \rightarrow \infty$,*

$$\int_{\mathcal{M}_g} \sum_{\substack{Y \in \text{Sub}_T(X); \\ Y \cong S_{g_0, k}}} e^{T - \frac{1-\epsilon_1}{2} \ell(\partial Y)} \mathbf{1}_{[0, 2T]}(\ell(\partial Y)) dX \prec_m T^{4m+2} e^{(1+\epsilon_1)T} \frac{V_g}{g^m}.$$

Proof. The proof is identical to the one of [WX22, Proposition 34] in which the power 66 comes from the bound:

$$6g_0 - 6 + 4k = 3m + k \leq 66$$

which is above [WX22, Equation (46)]; in this current case, we have

$$6g_0 - 6 + 4k = 3m + k \leq 4m + 2$$

as desired. \square

Since we consider $|\chi(Y)| = m = 2g_0 + k - 2 \in [1, M]$, both g_0 and k are bounded. As a direct consequence of Proposition 56 we have

Proposition 57. *For any $\epsilon_1 > 0$ and fixed $M \in \mathbb{Z}_{\geq 1}$, then we have that as $g \rightarrow \infty$,*

$$\int_{\mathcal{M}_g} \sum_{\substack{Y \in \text{Sub}_T(X); \\ 1 \leq |\chi(Y)| \leq M}} T e^{T - \frac{1-\epsilon_1}{2} \ell(\partial Y)} \mathbf{1}_{[0, 2T]}(\ell(\partial Y)) dX \prec_M T^{4M+3} e^{(1+\epsilon_1)T} \frac{V_g}{g}.$$

Now we are ready to prove the following effective bound for the expected value $\mathbb{E}_{\text{WP}}^g [N^{ns}(X_g, L)]$ and $\mathbb{E}_{\text{WP}}^g [N^{ns,2}(X_g, L)]$.

Theorem 58. *Assume that $L = L(g) \leq C \ln g$ for some constant $C > 0$ and $\lim_{g \rightarrow \infty} L(g) = \infty$. Then for any $\epsilon > 0$, as $g \rightarrow \infty$ we have*

$$\mathbb{E}_{\text{WP}}^g [N^{ns}(X_g, L)] \prec_{C, \epsilon} \frac{e^{(1+\epsilon)L}}{g}$$

and

$$\mathbb{E}_{\text{WP}}^g [N^{ns,2}(X_g, L)] \prec_{C, \epsilon} \frac{e^{(1+\epsilon)L}}{g}.$$

Proof. The conclusion clearly follows by Proposition 52, Proposition 55 for any fixed $m \geq 100(1+C)$ and Proposition 57 for $M = m - 1$ and $\epsilon_1 = \frac{\epsilon}{2}$ where we assume g is large enough such that $L^{3+400(1+C)} \prec e^{\frac{\epsilon}{2}L}$. \square

Proof of Theorem 51. The conclusion clearly follows from Theorem 58 by letting $C = 12$. \square

Expected value on simple and non-separating closed geodesics. We have shown in Lemma 48 that

$$\mathbb{E}_{\text{WP}}^g [N_{nsep}^s(X_g, L)] = \frac{1}{2} \text{Li}(e^L) \cdot \left(1 + O\left(\frac{L^2}{g} + \frac{L \ln L}{e^L}\right) \right)$$

when $\lim_{g \rightarrow \infty} L(g) = \infty$ and $\lim_{g \rightarrow \infty} \frac{L(g)^2}{g} = 0$. Now we aim to show that

Theorem 59. *Assume $L = L(g)$ satisfies*

$$\lim_{g \rightarrow \infty} L(g) = \infty \quad \text{and} \quad \limsup_{g \rightarrow \infty} \frac{L(g)}{\ln g} < \infty.$$

Then for any fixed $\epsilon > 0$ we have

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; N_{nsep}^s(X_g, L) > \frac{1-\epsilon}{2} \text{Li}(e^L) \right) = 1.$$

Similar to the methods in [MP19, NWX20], we compute the variance of $N_{nsep}^s(X_g, L)$ by using the Chebyshev inequality. For any fixed $\epsilon > 0$ we have

$$\begin{aligned} & \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N_{nsep}^s(X_g, L)}{\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]} \right| \geq \frac{1}{2}\epsilon \right) \\ (78) \quad & \leq \frac{\text{Var}[N_{nsep}^s(X_g, L)]}{\frac{\epsilon^2}{4} \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2} \\ & = \frac{4 \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)^2] - \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2}{\epsilon^2 \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2} \end{aligned}$$

where $\text{Var}[\cdot] = \mathbb{E}_{\text{WP}}^g[\cdot^2] - \mathbb{E}_{\text{WP}}^g[\cdot]^2$ is the variance of a random variable.

We decompose $N_{nsep}^s(X_g, L)^2$ into four different parts as follows.

Definition 60. Let $\mathcal{P}'(X_g)$ to be the set of all unoriented primitive closed geodesics on X_g . For $1 \leq k \leq \lfloor \frac{g-1}{2} \rfloor$, define

$$\begin{aligned} Y_1(X_g, L) &:= \# \left\{ \begin{array}{l} \alpha, \beta \text{ are simple and non-separating,} \\ (\alpha, \beta) \in \mathcal{P}'(X_g)^2; \quad \ell_\alpha(X_g) \leq L, \ell_\beta(X_g) \leq L, \\ \alpha \cap \beta = \emptyset, \alpha \neq \beta, \\ X_g \setminus (\alpha \cup \beta) \cong S_{g-2,4} \end{array} \right\}, \\ Y_2^k(X_g, L) &:= \# \left\{ \begin{array}{l} \alpha, \beta \text{ are simple and non-separating,} \\ (\alpha, \beta) \in \mathcal{P}'(X_g)^2; \quad \ell_\alpha(X_g) \leq L, \ell_\beta(X_g) \leq L, \\ \alpha \cap \beta = \emptyset, \alpha \neq \beta, \\ X_g \setminus (\alpha \cup \beta) \cong S_{k,2} \cup S_{g-k-1,2} \end{array} \right\}, \\ Z(X_g, L) &:= \# \left\{ \begin{array}{l} \alpha, \beta \text{ are simple and non-separating,} \\ (\alpha, \beta) \in \mathcal{P}'(X_g)^2; \quad \ell_\alpha(X_g) \leq L, \ell_\beta(X_g) \leq L, \\ \alpha \cap \beta \neq \emptyset, \alpha \neq \beta \end{array} \right\}. \end{aligned}$$

Then

(79)

$$N_{nsep}^s(X_g, L)^2 = N_{nsep}^s(X_g, L) + Y_1(X_g, L) + \sum_{k=1}^{\lfloor \frac{g-1}{2} \rfloor} Y_2^k(X_g, L) + Z(X_g, L).$$

Next we estimate the expectation of each term of the RHS of (79) above.

Lemma 61. *Assume $L = L(g)$ satisfies $\lim_{g \rightarrow \infty} L(g) = \infty$ and $\lim_{g \rightarrow \infty} \frac{L(g)^2}{g} = 0$.*

Then

$$|\mathbb{E}_{\text{WP}}^g[Y_1(X_g, L)] - \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2| \prec \frac{1}{g}e^{2L}.$$

Proof. By Mirzakhani's integration formula, i.e., Theorem 8, we have

$$\mathbb{E}_{\text{WP}}^g[Y_1(X_g, L)] = \frac{1}{V_g} \frac{1}{4} \int_0^L \int_0^L V_{g-2,4}(x, x, y, y) xy dx dy.$$

Since $L = L(g) = o(\sqrt{g})$, by Theorem 12 we have that for all $x, y \in [0, L]$,

$$\frac{V_{g-2,4}(x, x, y, y)}{V_{g-2,4}} = \left(\frac{\sinh\left(\frac{x}{2}\right)}{\frac{x}{2}} \right)^2 \left(\frac{\sinh\left(\frac{y}{2}\right)}{\frac{y}{2}} \right)^2 \left(1 + O\left(\frac{x^2 + y^2}{g}\right) \right).$$

By Theorem 10,

$$\frac{V_{g-2,4}}{V_g} = \frac{V_{g-2,4}}{V_{g-1,2}} \frac{V_{g-1,2}}{V_g} = 1 + O\left(\frac{1}{g}\right).$$

So

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g[Y_1(X_g, L)] &= \frac{1}{4} \int_0^L \int_0^L \left(\sinh\left(\frac{x}{2}\right) \sinh\left(\frac{y}{2}\right) \right)^2 \frac{4}{xy} \\ &\quad \left(1 + O\left(\frac{1 + x^2 + y^2}{g}\right) \right) dx dy \\ &= \left(\int_0^L \left(\sinh\left(\frac{x}{2}\right) \right)^2 \frac{1}{x} dx \right)^2 \left(1 + O\left(\frac{L^2}{g}\right) \right). \end{aligned}$$

Similarly,

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)] &= \frac{1}{V_g} \frac{1}{2} \int_0^L V_{g-1,2}(x, x) x dx \\ &= \int_0^L \left(\sinh\left(\frac{x}{2}\right) \right)^2 \frac{1}{x} dx \cdot \left(1 + O\left(\frac{L^2}{g}\right) \right). \end{aligned}$$

Thus, we obtain

$$\begin{aligned} &|\mathbb{E}_{\text{WP}}^g[Y_1(X_g, L)] - \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2| \\ &= \left(\int_0^L \left(\sinh\left(\frac{x}{2}\right) \right)^2 \frac{1}{x} dx \right)^2 \cdot O\left(\frac{L^2}{g}\right) \\ &\prec \frac{1}{g}e^{2L} \end{aligned}$$

as desired. \square

Lemma 62. *Assume $L = L(g)$ satisfies $\lim_{g \rightarrow \infty} L(g) = \infty$. Then*

$$\sum_{k=1}^{\lfloor \frac{g-1}{2} \rfloor} \mathbb{E}_{\text{WP}}^g[Y_2^k(X_g, L)] \prec \frac{1}{g^2} \frac{e^{2L}}{L^2}.$$

Proof. By Mirzakhani's integration formula, i.e., Theorem 8, we have

$$\mathbb{E}_{\text{WP}}^g[Y_2^k(X_g, L)] \leq \frac{1}{V_g} \int_0^L \int_0^L V_{k,2}(x, y) V_{g-k-1,2}(x, y) xy dx dy.$$

Then by Theorem 12,

$$\begin{aligned} \mathbb{E}_{\text{WP}}^g[Y_2^k(X_g, L)] &\leq \frac{V_{k,2} V_{g-k-1,2}}{V_g} \int_0^L \int_0^L \left(\frac{\sinh(\frac{x}{2})}{\frac{x}{2}} \frac{\sinh(\frac{y}{2})}{\frac{y}{2}} \right)^2 xy dx dy \\ &\prec \frac{V_{k,2} V_{g-k-1,2}}{V_g} \cdot \frac{e^{2L}}{L^2}. \end{aligned}$$

Recall that Theorem 11 says that

$$\sum_{k=1}^{\lfloor \frac{g-1}{2} \rfloor} \frac{V_{k,2} V_{g-k-1,2}}{V_g} \asymp \frac{1}{g^2}.$$

Therefore, the proof is finished by taking a summation over k . \square

Lemma 63. *Assume $L = L(g)$ satisfies $\lim_{g \rightarrow \infty} L(g) = \infty$ and $L(g) \leq C \ln g$ for some constant $C > 0$. Then for any $\epsilon > 0$,*

$$\mathbb{E}_{\text{WP}}^g[Z(X_g, L)] \prec_{C, \epsilon} \frac{1}{g} e^{(2+\epsilon)L}.$$

Proof. The conclusion clearly follows by Theorem 58. Here we recall that $N^{ns,2}(X, T)$ is the number of pairs of intersected unoriented closed geodesics pair (γ_1, γ_2) of X of total length $\ell_{\gamma_1}(X) + \ell_{\gamma_2}(X) \leq T$, and hence $Z(X_g, L) \leq N^{ns,2}(X_g, 2L)$. \square

Now we prove Theorem 59.

Proof of Theorem 59. Now assume

$$\lim_{g \rightarrow \infty} L(g) = \infty \quad \text{and} \quad \limsup_{g \rightarrow \infty} \frac{L(g)}{\ln g} \leq C - 1 < \infty$$

for some constant $C > 1$. So $L \leq C \ln g$ for large enough g .

Then it follows by Lemma 48, 61, 62, 63 and Equation (79) that for any $\epsilon_1 > 0$,

$$\begin{aligned} &\frac{\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)^2] - \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2}{\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2} \\ &\prec_{\epsilon_1} \frac{\text{Li}(e^L) + \frac{1}{g} e^{2L} + \frac{1}{g^2} \frac{e^{2L}}{L^2} + \frac{1}{g} e^{(2+\epsilon_1)L}}{\text{Li}(e^L)^2} \\ &\prec L e^{-L} + \frac{1}{g} L^2 e^{\epsilon_1 L} \\ &\prec_C L e^{-L} + (\ln g)^2 g^{\epsilon_1 C - 1}. \end{aligned}$$

Now take $\epsilon_1 = \frac{1}{2C}$. For any fixed $\epsilon > 0$, by (78) as $g \rightarrow \infty$ we have

$$\begin{aligned} & \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{M}_g; N_{nsep}^s(X_g, L) \leq (1 - \frac{1}{2}\epsilon)\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]) \\ & \leq \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N_{nsep}^s(X_g, L)}{\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]} \right| \geq \frac{1}{2}\epsilon \right) \\ & \leq \frac{4}{\epsilon^2} \frac{\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)^2] - \mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2}{\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]^2} \\ & \prec_C \frac{1}{\epsilon^2} \left(Le^{-L} + (\ln g)^2 g^{-\frac{1}{2}} \right). \end{aligned}$$

By Lemma 48, we know that for g large enough,

$$\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)] \geq \frac{1 - \epsilon/2}{2} \text{Li}(e^L).$$

Thus, we have that for large enough g ,

$$\begin{aligned} & \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; N_{nsep}^s(X_g, L) > \frac{(1 - \epsilon/2)^2}{2} \text{Li}(e^L) \right) \\ & \geq 1 - \text{Prob}_{\text{WP}}^g (X_g \in \mathcal{M}_g; N_{nsep}^s(X_g, L) \leq (1 - \frac{1}{2}\epsilon)\mathbb{E}_{\text{WP}}^g[N_{nsep}^s(X_g, L)]) \\ & = 1 - O_C \left(\frac{1}{\epsilon^2} \left(Le^{-L} + (\ln g)^2 g^{-\frac{1}{2}} \right) \right). \end{aligned}$$

Then the conclusion follows by letting $g \rightarrow \infty$ because $\frac{(1 - \epsilon/2)^2}{2} \text{Li}(e^L) > \frac{1 - \epsilon}{2} \text{Li}(e^L)$. \square

Now we are ready to finish the proof of Theorem 49.

Proof of Theorem 49. Assume that

$$\lim_{g \rightarrow \infty} L(g) = \infty \quad \text{and} \quad \limsup_{g \rightarrow \infty} \frac{L(g)}{\ln g} \leq C - 1 < \infty$$

for some constant $C > 1$. So $L \leq C \ln g$ for large enough g .

By (62), Lemma 50 and Theorem 58, for any $\epsilon_1 > 0$, when g is large enough we have

$$\begin{aligned} & \mathbb{E}_{\text{WP}}^g [|N(X_g, L) - N_{nsep}^s(X_g, L)|] \\ & = \mathbb{E}_{\text{WP}}^g \left[\sum_{k=1}^{\lfloor g/2 \rfloor} N_{sep,k}^s(X_g, L) + N^{ns}(X_g, L) \right] \\ & \prec_{C, \epsilon_1} \frac{1}{g} \text{Li}(e^L) + \frac{1}{g} e^{(1 + \epsilon_1)L} \\ & \prec \frac{1}{g} e^{(1 + \epsilon_1)L}. \end{aligned}$$

Then by Markov's inequality we have

$$\text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; |N(X_g, L) - N_{nsep}^s(X_g, L)| \geq \frac{e^{(1 + 2\epsilon_1)L}}{g} \right) \prec_{C, \epsilon_1} e^{-\epsilon_1 L}$$

which converges to 0 as $g \rightarrow \infty$.

On the other hand, by Theorem 59 we have

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; N_{n\text{sep}}^s(X_g, L) > \frac{1}{4} \text{Li}(e^L) \right) = 1.$$

Thus,

$$\lim_{g \rightarrow \infty} \text{Prob}_{\text{WP}}^g \left(X_g \in \mathcal{M}_g; \left| 1 - \frac{N_{n\text{sep}}^s(X_g, L)}{N(X_g, L)} \right| < \frac{4e^{(1+2\epsilon_1)L}}{g \text{Li}(e^L)} \right) = 1.$$

Then the proof is finished by taking suitable $\epsilon_1 = \epsilon_1(C, \epsilon) > 0$. \square

Proof of Theorem 4. It clearly follows from Theorem 43, 47 and 49. \square

APPENDIX A. COUNTING FILLING k -TUPLES

In this Appendix, we give a brief sketch of the proof of Theorem 18 here. One may see [WX22, Section 8] for more details.

First, as a simple application of Corollary 15 we have

Lemma 64. *For fixed $k \in \mathbb{Z}_{\geq 1}$, there exists a constant $c_k > 0$ only depending on k such that for any hyperbolic surface $X \in \mathcal{T}_{g,n}(L_1, \dots, L_n)$ and $L > 0$, we have*

$$\# \left\{ (\gamma_1, \dots, \gamma_k) \in \mathcal{P}(X)^k; \sum_{i=1}^k \ell(\gamma_i) \leq L \right\} \leq c_k \cdot (2g - 2 + n)^k (1 + L)^{k-1} e^L$$

where $\mathcal{P}(X)$ is the set of all oriented non-peripheral primitive closed geodesics in X .

Proof. By doubling X along its boundaries, we get a closed hyperbolic surface $\bar{X} \in \mathcal{T}_{2g+n-1}$. Each non-peripheral primitive closed geodesic in X has two copies in \bar{X} . So by Corollary 15, for any $T > 0$ we have

$$\begin{aligned} \#\{\gamma \in \mathcal{P}(X); \ell(\gamma) \leq T\} &\leq \frac{1}{2} \#\{\gamma \in \mathcal{P}(\bar{X}); \ell(\gamma) \leq T\} \\ &\leq (2g - 2 + n) e^{T+7}. \end{aligned}$$

Now we prove by induction on k . First the lemma holds for $k = 1$ by the above inequality. Now assume that the lemma holds for $k - 1$. Then for

$k \geq 2$, we get

$$\begin{aligned}
& \# \left\{ (\gamma_1, \dots, \gamma_k) \in \mathcal{P}(X)^k; \sum_{i=1}^k \ell(\gamma_i) \leq L \right\} \\
& \leq \sum_{T=1}^{[L]+1} \# \{ \gamma_1 \in \mathcal{P}(X); T-1 < \ell(\gamma_1) \leq T \} \\
& \quad \times \# \left\{ (\gamma_2, \dots, \gamma_k) \in \mathcal{P}(X)^{k-1}; \sum_{i=1}^{k-1} \ell(\gamma_i) \leq L - T + 1 \right\} \\
& \prec_k \sum_{T=1}^{[L]+1} (2g-2+n)e^T \cdot (2g-2+n)^{k-1} (L-T+2)^{k-2} e^{L-T+1} \\
& \prec \sum_{T=1}^{[L]+1} (2g-2+n)^k (1+L)^{k-2} e^L \\
& \leq (2g-2+n)^k (1+L)^{k-1} e^L
\end{aligned}$$

as desired. \square

Similar to [WX22, Theorem 38], we study counting filling k -tuples instead of filling geodesics. We prove

Theorem 65. *There exists a universal constant $L_0 > 0$ such that for any $0 < \epsilon < 1$, $m = 2g - 2 + n \geq 1$, $\Delta \geq 0$ and $\sum_{i=1}^n x_i \geq \Delta + L_0 \cdot \frac{m \cdot n}{\epsilon}$, the following holds: for any hyperbolic surface $X \in \mathcal{T}_{g,n}(x_1, \dots, x_n)$, one can always find a new hyperbolic surface $Y \in \mathcal{T}_{g,n}(y_1, \dots, y_n)$ satisfying*

- (1) $y_i \leq x_i$ for all $1 \leq i \leq n$;
- (2) $\sum_{i=1}^n x_i - \sum_{i=1}^n y_i = \Delta$;
- (3) for any filling k -tuple $\Gamma = (\gamma_1, \dots, \gamma_k)$ in $S_{g,n}$, we have

$$\sum_{i=1}^k \ell_{\gamma_i}(X) - \sum_{i=1}^k \ell_{\gamma_i}(Y) \geq \frac{1}{2}(1-\epsilon)\Delta.$$

Proof. The proof is identical to the one of [WX22, Theorem 38] through replacing filling curves by filling k -tuples. We give a sketch about the proof here, and one may refer to [WX22, Section 8] for more details.

Consider a longest boundary component of X , denoted by α . Let w be the width of the maximal embedded half-collar of α in X , that is,

$$w = \sup \left\{ \begin{array}{l} \{p \in X; \text{dist}(p, \alpha) < d\} \\ d > 0; \text{ is homeomorphic to a cylinder and} \\ \text{does not intersect with } (\partial X \setminus \alpha) \end{array} \right\}.$$

Then the closed curve $\{p \in X; \text{dist}(p, \alpha) = w\}$ either touches another component of ∂X or has non-empty self-intersection. For both cases, it

induces a pair of pants, denoted by P , in which α is a boundary curve (see [WX22, Page 382 Construction] for details).

For any $k \in \mathbb{Z}_{\geq 1}$ and filling k -tuple $\Gamma = (\gamma_1, \dots, \gamma_k)$ of X , at least one of γ_i 's must intersect with the pair of pants P (actually this is the only place where we apply the filling property for k -tuples). Then as in [WX22, Page 384] we reduce the length of $\partial P \cap \partial X$ and keep the part $X \setminus P$ unchanged, the length of those γ_i 's which do not intersect with P are unchanged. For those γ_i 's which intersect with P , by [WX22, Remark 41], length of γ_i reduce at least the multiplication of $\frac{1-\epsilon}{2}$ and the reduction of $\partial P \cap \partial X$ if $\ell(\alpha)$ is large enough.

So if the initial total boundary length $\sum x_i$ is large enough, we can always find a long boundary component and apply the above process to reduce the total boundary length. The process is workable unless all boundary components are bounded by some constant related to ϵ and m . As argued above, similar as [WX22, Proposition 40], for each time the total length of any filling k -tuple must reduce at least the multiplication of $\frac{1-\epsilon}{2}$ and the reduction of the total boundary length. Then by using the same argument as in the proof of [WX22, Theorem 38], one can get a desired new hyperbolic surface Y satisfying all the three required properties. \square

Now applying Theorem 65, we finish the proof of Theorem 18.

Proof of Theorem 18. If $n = 0$, $\ell(\partial X) = 0$. Then the conclusion holds directly by Lemma 64. Now assume $n \geq 1$ and $X \in \mathcal{T}_{g,n}(x_1, \dots, x_n)$.

Case 1: the total boundary length $\sum_{i=1}^n x_i > L_0 \frac{mn}{\epsilon}$. By Theorem 65 we may take $\Delta = \sum x_i - L_0 \frac{mn}{\epsilon}$ and then get a hyperbolic surface $Y \in \mathcal{T}_{g,n}(y_1, \dots, y_n)$ such that $\sum y_i = L_0 \frac{mn}{\epsilon}$ and for any filling k -tuple $\Gamma = (\gamma_1, \dots, \gamma_k)$ in $S_{g,n}$ we have

$$\sum_{i=1}^k \ell_{\gamma_i}(X) - \sum_{i=1}^k \ell_{\gamma_i}(Y) \geq \frac{1-\epsilon}{2} \cdot \left(\sum_{i=1}^n x_i - L_0 \frac{mn}{\epsilon} \right).$$

Thus

$$N_k^{\text{fill}}(X, L) \leq N_k^{\text{fill}} \left(Y, L - \frac{1-\epsilon}{2} \cdot \left(\sum_{i=1}^n x_i - L_0 \frac{mn}{\epsilon} \right) \right)$$

which together with Lemma 64 and the fact that $n \leq m + 2$ implies that

$$\begin{aligned} N_k^{\text{fill}}(X, L) &\prec_k m^k \left(1 + L - \frac{1-\epsilon}{2} \left(\sum_{i=1}^n x_i - L_0 \frac{mn}{\epsilon} \right) \right)^{k-1} e^{L - \frac{1-\epsilon}{2} \left(\sum_{i=1}^n x_i - L_0 \frac{mn}{\epsilon} \right)} \\ &\prec_{k,\epsilon,m} (1+L)^{k-1} e^{L - \frac{1-\epsilon}{2} \sum x_i}. \end{aligned}$$

Case 2: the total boundary length $\sum_{i=1}^n x_i \leq L_0 \frac{mn}{\epsilon}$. Then by Lemma 64 directly we have

$$\begin{aligned} N_k^{\text{fill}}(X, L) &\prec_k m^k (1+L)^{k-1} e^L \\ &\prec_{k,\epsilon,m} (1+L)^{k-1} e^{L - \frac{1-\epsilon}{2} \sum x_i}. \end{aligned}$$

The proof is complete. \square

REFERENCES

- [Agg21] Amol Aggarwal, *Large genus asymptotics for intersection numbers and principal strata volumes of quadratic differentials*, Invent. Math. **226** (2021), no. 3, 897–1010.
- [AM22] Nalini Anantharaman and Laura Monk, *A high-genus asymptotic expansion of Weil-Petersson volume polynomials*, J. Math. Phys. **63** (2022), no. 4, Paper No. 043502, 26.
- [Ber16] Nicolas Bergeron, *The spectrum of hyperbolic surfaces*, Universitext, Springer, Cham; EDP Sciences, Les Ulis, 2016, Appendix C by Valentin Blomer and Farrell Brumley, Translated from the 2011 French original by Brumley [2857626].
- [BMM16] Werner Ballmann, Henrik Matthiesen, and Sugata Mondal, *Small eigenvalues of closed surfaces*, J. Differential Geom. **103** (2016), no. 1, 1–13.
- [BMM17] ———, *On the analytic systole of Riemannian surfaces of finite type*, Geom. Funct. Anal. **27** (2017), no. 5, 1070–1105.
- [BP22] Maxime Fortier Bourque and Bram Petri, *Kissing numbers of closed hyperbolic manifolds*, Amer. J. Math. **144** (2022), no. 4, 1067–1086.
- [BS94] P. Buser and P. Sarnak, *On the period matrix of a Riemann surface of large genus*, Invent. Math. **117** (1994), no. 1, 27–56, With an appendix by J. H. Conway and N. J. A. Sloane.
- [Bus77] Peter Buser, *Riemannsche Flächen mit Eigenwerten in $(0, 1/4)$* , Comment. Math. Helv. **52** (1977), no. 1, 25–34.
- [Bus10] ———, *Geometry and spectra of compact Riemann surfaces*, Modern Birkhäuser Classics, Birkhäuser Boston, Ltd., Boston, MA, 2010, Reprint of the 1992 edition.
- [Cai02] Yingchun Cai, *Prime geodesic theorem*, J. Théor. Nombres Bordeaux **14** (2002), no. 1, 59–72.
- [Cha84] Isaac Chavel, *Eigenvalues in Riemannian geometry*, Pure and Applied Mathematics, vol. 115, Academic Press, Inc., Orlando, FL, 1984, Including a chapter by Burton Randol, With an appendix by Jozef Dodziuk.
- [DGZZ20] Vincent Delecroix, Élise Goujard, Peter Zograf, and Anton Zorich, *Uniform lower bound for intersection numbers of ψ -classes*, SIGMA Symmetry Integrability Geom. Methods Appl. **16** (2020), 086, 13 pages.
- [GLMST21] Clifford Gilmore, Etienne Le Masson, Tuomas Sahlsten, and Joe Thomas, *Short geodesic loops and L^p norms of eigenfunctions on large genus random surfaces*, Geom. Funct. Anal. **31** (2021), no. 1, 62–110.
- [GPY11] Larry Guth, Hugo Parlier, and Robert Young, *Pants decompositions of random surfaces*, Geom. Funct. Anal. **21** (2011), no. 5, 1069–1090.
- [Hej76] Dennis A. Hejhal, *The Selberg trace formula for $\mathrm{PSL}(2, R)$. Vol. I*, Lecture Notes in Mathematics, Vol. 548, Springer-Verlag, Berlin-New York, 1976.
- [Hid21] Will Hide, *Spectral gap for Weil-Petersson random surfaces with cusps*, arXiv e-prints (2021), arXiv:2107.14555.
- [HM21] Will Hide and Michael Magee, *Near optimal spectral gaps for hyperbolic surfaces*, arXiv e-prints (2021), arXiv:2107.05292.
- [Hub61] Heinz Huber, *Zur analytischen Theorie hyperbolischer Raumformen und Bewegungsgruppen. II*, Math. Ann. **142** (1960/61), 385–398.
- [HW22] Yuxin He and Yunhui Wu, *On second eigenvalues of closed hyperbolic surfaces for large genus*, arXiv e-prints (2022), arXiv:2207.12919.
- [Iwa84] Henryk Iwaniec, *Prime geodesic theorem*, J. Reine Angew. Math. **349** (1984), 136–159.
- [LRS95] W. Luo, Z. Rudnick, and P. Sarnak, *On Selberg’s eigenvalue conjecture*, Geom. Funct. Anal. **5** (1995), no. 2, 387–401.

- [LS20] Etienne Le Masson and Tuomas Sahlsten, *Quantum ergodicity for Eisenstein series on hyperbolic surfaces of large genus*, arXiv e-prints (2020), arXiv:2006.14935.
- [LW21] Michael Lipnowski and Alex Wright, *Towards optimal spectral gaps in large genus*, arXiv e-prints (2021), arXiv:2103.07496.
- [LX14] Kefeng Liu and Hao Xu, *A remark on Mirzakhani's asymptotic formulae*, Asian J. Math. **18** (2014), no. 1, 29–52.
- [Mir07a] Maryam Mirzakhani, *Simple geodesics and Weil-Petersson volumes of moduli spaces of bordered Riemann surfaces*, Invent. Math. **167** (2007), no. 1, 179–222.
- [Mir07b] ———, *Weil-Petersson volumes and intersection theory on the moduli space of curves*, J. Amer. Math. Soc. **20** (2007), no. 1, 1–23.
- [Mir08] ———, *Growth of the number of simple closed geodesics on hyperbolic surfaces*, Ann. of Math. (2) **168** (2008), no. 1, 97–125.
- [Mir13] ———, *Growth of Weil-Petersson volumes and random hyperbolic surfaces of large genus*, J. Differential Geom. **94** (2013), no. 2, 267–300.
- [MNP22] Michael Magee, Frédéric Naud, and Doron Puder, *A random cover of a compact hyperbolic surface has relative spectral gap $\frac{3}{16} - \epsilon$* , Geom. Funct. Anal. **32** (2022), no. 3, 595–661.
- [Mon22] Laura Monk, *Benjamini-Schramm convergence and spectra of random hyperbolic surfaces of high genus*, Anal. PDE **15** (2022), no. 3, 727–752.
- [MP19] Maryam Mirzakhani and Bram Petri, *Lengths of closed geodesics on random surfaces of large genus*, Comment. Math. Helv. **94** (2019), no. 4, 869–889.
- [MZ15] Maryam Mirzakhani and Peter Zograf, *Towards large genus asymptotics of intersection numbers on moduli spaces of curves*, Geom. Funct. Anal. **25** (2015), no. 4, 1258–1289.
- [NWX20] Xin Nie, Yunhui Wu, and Yuhao Xue, *Large genus asymptotics for lengths of separating closed geodesics on random surfaces*, arXiv e-prints (2020), arXiv:2009.07538.
- [OR09] Jean-Pierre Otal and Eulalio Rosas, *Pour toute surface hyperbolique de genre g , $\lambda_{2g-2} > 1/4$* , Duke Math. J. **150** (2009), no. 1, 101–115.
- [PWX21] Hugo Parlier, Yunhui Wu, and Yuhao Xue, *The simple separating systole for hyperbolic surfaces of large genus*, Journal of the Institute of Mathematics of Jussieu (2021), to appear.
- [Ran77] Burton Randol, *On the asymptotic distribution of closed geodesics on compact Riemann surfaces*, Trans. Amer. Math. Soc. **233** (1977), 241–247.
- [Rud22] Zeév Rudnick, *GOE statistics on the moduli space of surfaces of large genus*, arXiv e-prints (2022), arXiv:2202.06379.
- [Sar80] Peter Clive Sarnak, *PRIME GEODESIC THEOREMS*, ProQuest LLC, Ann Arbor, MI, 1980, Thesis (Ph.D.)–Stanford University. MR 2630950
- [Sel56] A. Selberg, *Harmonic analysis and discontinuous groups in weakly symmetric Riemannian spaces with applications to Dirichlet series*, J. Indian Math. Soc. (N.S.) **20** (1956), 47–87.
- [SW22] Yang Shen and Yunhui Wu, *Arbitrarily small spectral gaps for random hyperbolic surfaces with many cusps*, arXiv e-prints (2022), arXiv:2203.15681.
- [SY13] K. Soundararajan and Matthew P. Young, *The prime geodesic theorem*, J. Reine Angew. Math. **676** (2013), 105–120.
- [Tho20] Joe Thomas, *Delocalisation of eigenfunctions on large genus random surfaces*, Israel Journal of Mathematics, to appear (2020).
- [Wol82] Scott Wolpert, *The Fenchel-Nielsen deformation*, Ann. of Math. (2) **115** (1982), no. 3, 501–528.
- [Wri20] Alex Wright, *A tour through Mirzakhani's work on moduli spaces of Riemann surfaces*, Bull. Amer. Math. Soc. (N.S.) **57** (2020), no. 3, 359–408.

- [WX22] Yunhui Wu and Yuhao Xue, *Random hyperbolic surfaces of large genus have first eigenvalues greater than $\frac{3}{16} - \epsilon$* , *Geom. Funct. Anal.* **32** (2022), no. 2, 340–410.

YAU MATHEMATICAL SCIENCES CENTER AND DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING, CHINA

Email address, (Y. W.): yunhui_wu@tsinghua.edu.cn

Email address, (Y. X.): xueyh18@mails.tsinghua.edu.cn