

STOCHASTIC PROCESSES UNDER PARAMETER UNCERTAINTY

DAVID CRIENS

ABSTRACT. In this paper we study a family of nonlinear (conditional) expectations that can be understood as a stochastic process with uncertain parameters. We develop a general framework which can be seen as a version of the martingale problem method of Stroock and Varadhan with parameter uncertainty. In particular, we show that our methodology includes the important class of nonlinear Lévy processes as introduced by Neufeld and Nutz, and we introduce the new classes of stochastic partial differential equations and spin systems with parameter uncertainty. Moreover, we study properties of the nonlinear expectations. We prove the dynamic programming principle, i.e., the tower property, and we establish conditions for the (strong) USC_b -Feller property and a strong Markov selection principle. Finally, we investigate ergodicity for spin systems with uncertain parameters such as the nonlinear contact process.

1. INTRODUCTION

We study conditional nonlinear expectations on the path space Ω of continuous or càdlàg functions from \mathbb{R}_+ into a Polish space F . More specifically, we are interested in nonlinear expectations of the form

$$(1.1) \quad \mathcal{E}_t(\psi)(\omega) = \sup_{P \in \mathcal{C}(t, \omega)} E^P[\psi],$$

where ψ is a suitable real-valued function on Ω and $\mathcal{C}(t, \omega)$ is a set of laws of stochastic processes whose paths coincide with $\omega \in \Omega$ till time $t \in \mathbb{R}_+$. We think of the nonlinear conditional expectation \mathcal{E} as a *nonlinear stochastic process*.

The systematic study of nonlinear stochastic processes started with the seminal work of Peng [43, 44] on the G -Brownian motion. More recently, larger classes of nonlinear semimartingales have been investigated in [8, 9, 10, 19, 22, 23, 29, 38]. At this point we also highlight the articles [16, 37, 40] where abstract tools for the study were developed. Next to the measure theoretic approach based on the nonlinear expectation (1.1), nonlinear Markov processes have also been constructed by analytic methods via nonlinear semigroups, see [14, 35, 36]. This construction is based on fundamental ideas of Nisio [39]. In the recent paper [30], it was shown that the two approaches are equivalent for the class of nonlinear Lévy processes from [14, 38]. A key property of the construction via (1.1) is its relation to a class of stochastic processes given through the set \mathcal{C} . This connection opens the door to analyze properties of \mathcal{E} with powerful tools from probability theory such as stochastic calculus and stochastic comparison.

Date: September 22, 2022.

2020 Mathematics Subject Classification. 60G65, 60G44, 60G07, 60H15, 60J25, 60K35.

Key words and phrases. nonlinear stochastic process; martingale problem; stochastic partial differential equations; spin system; interacting particle system; sublinear expectation; nonlinear expectation; parameter uncertainty; Knightian uncertainty; nonlinear contact process; nonlinear Ising model.

The author acknowledges financial support from the DFG project SCHM 2160/15-1.

In this paper, we propose a new framework for the set \mathcal{C} which extends the nonlinear expectation approach to a much larger class of nonlinear stochastic processes than nonlinear semimartingales. The framework is inspired by the martingale problem method of Stroock and Varadhan [50]. Consider the family

$$(1.2) \quad f(X) - \int_0^\cdot g(X_s)ds, \quad (f, g) \in A \subset C_b(F; \mathbb{R}) \times C_b(F; \mathbb{R}),$$

of test processes, where X is the coordinate process on Ω . The set A of test functions is often called a pregenerator. A probability measure P is said to be a solution to the martingale problem associated to A if all test processes in (1.2) are local P -martingales. Our idea is to incorporate uncertainty to the pregenerator A .

Let U be a countable index set and let $\{Y^u: u \in U\}$ be a family of càdlàg processes on Ω , which serve as test processes for the martingale problem under uncertainty. If P is a probability measure such that Y^u is a *special* P -semimartingale, then there exists a P -unique predictable càdlàg process $A^P(Y^u)$ of finite variation, starting in zero, such that $Y^u - Y_0^u - A^P(Y^u)$ is a local P -martingale. In case the process in (1.2) is a local P -martingale, the process $f(X)$ is a special semimartingale and $A^P(f(X)) = \int_0^\cdot g(X_s)ds$. Motivated by this observation, we define \mathcal{C} to be the set of probability measures P under that each test process Y^u is a special P -semimartingale with absolutely continuous compensator $A^P(Y^u)$ such that $(\lambda \otimes P)$ -a.e.

$$(dA^P(Y^u)/d\lambda)_{u \in U} \in \Theta.$$

Here, Θ is a time and path-dependent set-valued mapping on $\mathbb{R}_+ \times \Omega$, which captures the uncertainty.

Our framework is tailor made for adding uncertainty to any class of stochastic processes which can be characterized by a martingale problem. For instance, this is the case for semimartingales, solutions to stochastic partial differential equations and many Markov processes. In particular, it is possible to recover any nonlinear semimartingale setting which is built from uncertain absolutely continuous semimartingale characteristics. To illustrate this, we show that the important class of nonlinear Lévy processes from [38] is included in our framework. Further, we introduce two novel classes of nonlinear Markov processes which are not necessarily semimartingales: the class of *nonlinear (in the sense of uncertainty) stochastic partial differential equations (NSPDE)* and the class of *nonlinear spin systems*, which are interacting particle systems with uncertain parameters. We think that both of these classes are of specific interest and therefore deserve further investigations.

Our first main result is the *dynamic programming principle (DPP)* for the nonlinear expectation (1.1), i.e., the tower property

$$\mathcal{E}_s(\mathcal{E}_t(\psi))(\omega) = \mathcal{E}_s(\psi)(\omega), \quad \omega \in \Omega, \quad s \leq t.$$

In case the uncertainty set Θ has a Markovian structure in the sense that $\Theta(t, \omega)$ depends on (t, ω) only through the value $\omega(t-)$, the DPP induces a *nonlinear Markov property* given by

$$\mathcal{E}^x(\mathcal{E}^{X_t}(\psi(X_s))) = \mathcal{E}^x(\psi(X_{t+s})), \quad \mathcal{E}^x(\psi) \triangleq \mathcal{E}_0(\psi)(\omega)|_{\omega(0)=x}.$$

As in the theory of (linear) Markov processes, there is a strong link to semigroups. Indeed, in this Markovian case, the nonlinear Markov property ensures the semigroup property $T_t T_s = T_{s+t}$ of the family $(T_t)_{t \geq 0}$ defined by

$$(1.3) \quad T_t(\psi)(x) \triangleq \mathcal{E}^x(\psi(X_t)),$$

where ψ runs through the class of bounded upper semianalytic functions. It is an important question when the semigroup $(T_t)_{t \geq 0}$ preserves some regularity. For a continuous path setting, we provide general conditions for the *USC_b-Feller property* of the semigroup $(T_t)_{t \geq 0}$, which means that it is a nonlinear semigroup on the space of bounded upper semicontinuous functions. Further, we establish a *strong Markov selection principle*, i.e., we prove that, for every bounded upper semicontinuous function $\psi: F \rightarrow \mathbb{R}$ and any $t > 0$, there exists a strong Markov family $\{P_{(s,x)}: (s,x) \in \mathbb{R}_+ \times F\}$ such that $P_{0,x} \in \mathcal{C}(0,x)$ and

$$T_t(\psi)(x) = E^{P_{(0,x)}}[\psi(X_t)].$$

We stress that the nonlinear structure of our setting is reflected by the fact that the strong Markov selection $\{P_{(s,x)}: (s,x) \in \mathbb{R}_+ \times F\}$ depends on the input elements ψ and t .

Thereafter, we investigate the Markovian classes of NSPDEs and nonlinear spin systems in more detail. For the former, we establish continuity and linear growth conditions for the *USC_b-Feller property* and the strong Markov selection principle. Moreover, for the important subclass of infinite-dimensional nonlinear Cauchy problems with drift uncertainty, we derive conditions for the *strong USC_b-Feller property*, which means that $T_t(\text{USC}_b) \subset C_b$ for all $t > 0$. The strong *USC_b-Feller property* can be seen as a *smoothing property* of the nonlinear semigroup $(T_t)_{t \geq 0}$. To the best of our knowledge, such an effect was first discovered in [8] for nonlinear one-dimensional diffusions. We also emphasize that the strong *USC_b-Feller property* entails the *C_b-Feller property*, i.e., $T_t(C_b) \subset C_b$ for all $t \in \mathbb{R}_+$. This property is considered to be of fundamental importance for the study of nonlinear semigroups via pointwise generators, see [9, Section 2.4] for some comments.

Finally, we investigate some ergodic theory of the class of nonlinear spin systems, which is motivated by the statistical mechanics origin of these processes. Ergodic theory deals with the set of invariant distributions and its domain of attraction. More specifically, we say that a nonlinear semigroup $(T_t)_{t \geq 0}$ is ergodic if it has a unique invariant distribution μ such that $\lim_{t \rightarrow \infty} \int T_t f d\nu = \int f d\mu$ for all probability measures ν and all functions f taken from a suitably rich class. In this paper, we establish ergodicity for nonlinear spin systems with finitely many sites that can be dominated by attractive and ergodic linear spin systems. For illustration, we show that our condition provides a simple criterion for the ergodicity of the contact process with uncertain infection rates.

Let us now comment on some technical aspects of our proofs. To establish the DPP, we invoke a general theorem from [16] which requires that we check a measurable graph condition and stability under conditioning and concatenation. The Markov property of the semigroup (1.3) can be deduced from the DPP, which, to the best of our knowledge, was first discovered in [22]. For the *USC_b-Feller property*, we rely on the theory of correspondences and, for the strong Markov selection principle, we invoke a technique from [15, 21] for controlled diffusions, which is based on ideas of Krylov [28] and Stroock and Varadhan [50] about Markovian selection. The strong *USC_b-Feller property* for nonlinear Cauchy problems is proved by means of a strong Feller selection principle, which adapts ideas from [9, 10] for finite-dimensional diffusions to an infinite-dimensional non-semimartingale setting. The concept of ergodicity of nonlinear spin systems seems to be new. Our criterion is based on a comparison with linear spin systems.

We end this section with a summary of the structure of the paper. In Section 2, we introduce our framework and formulate the DPP. Furthermore, in Section 2.3, we show that the class of nonlinear Lévy processes from [38] is included in our setting and we introduce the new classes of NSPDEs and nonlinear spin systems. Thereafter, in Section 3, we discuss nonlinear Markov processes. In Section 3.2, we establish the *USC_b-Feller property* and,

in Section 3.3, we provide the strong Markov selection principle. In Section 3.4, these results are tailored to the special case of NSPDEs and the strong USC_b -Feller property for nonlinear Cauchy problems with drift uncertainty is established. The ergodicity of nonlinear spin systems is discussed in Section 4. The remaining sections of this paper are devoted to the proofs of our results. The location of each proof is indicated before the statement of the respective result.

2. NONLINEAR PROCESSES AND THE DYNAMIC PROGRAMMING PRINCIPLE

2.1. The Ingredients. Let F be a Polish space and define Ω to be either the space of càdlàg functions $\mathbb{R}_+ \rightarrow F$ or the space of all continuous functions $\mathbb{R}_+ \rightarrow F$ endowed with the Skorokhod J_1 topology.¹ The canonical process on Ω is denoted by X , i.e., $X_t(\omega) = \omega(t)$ for $\omega \in \Omega$ and $t \in \mathbb{R}_+$. It is well-known that $\mathcal{F} \triangleq \mathcal{B}(\Omega) = \sigma(X_t, t \geq 0)$. We define $\mathbf{F} \triangleq (\mathcal{F}_t)_{t \geq 0}$ to be the canonical filtration generated by X , i.e., $\mathcal{F}_t \triangleq \sigma(X_s, s \leq t)$ for $t \in \mathbb{R}_+$. The set of all probability measures on (Ω, \mathcal{F}) is denoted by $\mathfrak{P}(\Omega)$ and endowed with the usual topology of convergence in distribution, i.e., the weak topology. The shift operator on Ω is denoted by $\theta = (\theta_t)_{t \geq 0}$, i.e., $\theta_t(\omega) = \omega(\cdot + t)$ for all $\omega \in \Omega$ and $t \in \mathbb{R}_+$. Let \mathbb{K} be either the real line \mathbb{R} or the complex plane \mathbb{C} . Further, let U be a countable index set and, for every $u \in U$, let Y^u be a \mathbb{K} -valued càdlàg \mathbf{F} -adapted process on Ω such that $Y_{+t}^u = Y^u \circ \theta_t$ for all $t \in \mathbb{R}_+$. We endow \mathbb{K}^U with the product Euclidean topology. Let $\Theta: \mathbb{R}_+ \times \Omega \rightarrow \mathbb{K}^U$ be a correspondence, i.e., a set-valued mapping.

Standing Assumption 2.1. The correspondence Θ has a measurable graph, i.e., the graph

$$\text{gr } \Theta = \{(t, \omega, g) \in \mathbb{R}_+ \times \Omega \times \mathbb{K}^U : g \in \Theta(t, \omega)\}$$

is Borel.

Recall that a \mathbb{K} -valued càdlàg adapted process $Y = (Y_t)_{t \geq 0}$ is said to be a *special semimartingale* if $Y = Y_0 + M + A$, where M is a local martingale and A is a predictable process of (locally) finite variation, both starting in zero. The decomposition is unique up to indistinguishability. A \mathbb{K} -valued càdlàg process Y is called a *special semimartingale after a time $t^* \in \mathbb{R}_+$* if the process $Y_{+t^*} = (Y_{t+t^*})_{t \geq 0}$ is a special semimartingale for the right-continuous natural filtration of $X_{+t^*} = (X_{t+t^*})_{t \geq 0}$. Notice that it comes without loss of generality that we consider the right-continuous version of the filtration ([37, Proposition 2.2]). For a given probability measure P on (Ω, \mathcal{F}) , the set of special semimartingales after time t^* on the setup $(\Omega, \mathcal{F}, \mathbf{F}, P)$ is denoted by $\mathcal{S}^{\text{sp}}(t^*, P)$. We write $\mathcal{S}^{\text{sp}}(P) \triangleq \mathcal{S}^{\text{sp}}(0, P)$. For given P and $Y \in \mathcal{S}^{\text{sp}}(t^*, P)$, denote by $A^P(Y_{+t^*})$ the predictable part of (locally) finite variation in the semimartingale decomposition of Y_{+t^*} , and denote by $\mathcal{S}_{\text{ac}}^{\text{sp}}(t^*, P)$ the set of all $Y \in \mathcal{S}^{\text{sp}}(t^*, P)$ such that $A^P(Y_{+t^*})$ is absolutely continuous w.r.t. to the Lebesgue measure λ . Again, we write $\mathcal{S}_{\text{ac}}^{\text{sp}}(P) \triangleq \mathcal{S}_{\text{ac}}^{\text{sp}}(0, P)$. For $\omega, \omega' \in \Omega$ and $t \in \mathbb{R}_+$, we define the concatenation

$$\omega \otimes_t \omega' \triangleq \omega \mathbb{1}_{[0, t)} + (\omega(t) + \omega' - \omega'(t)) \mathbb{1}_{[t, \infty)},$$

and, finally, for $(t, \omega) \in \llbracket 0, \infty \llbracket$, we define $\mathcal{C}(t, \omega) \subset \mathfrak{P}(\Omega)$ by

$$\begin{aligned} \mathcal{C}(t, \omega) \triangleq \{P \in \mathfrak{P}(\Omega) : P(X = \omega \text{ on } [0, t]) = 1, \forall u \in U \ Y^u \in \mathcal{S}_{\text{ac}}^{\text{sp}}(t, P), \\ (\lambda \otimes P)\text{-a.e. } (dA^P(Y_{+t}^u)/d\lambda)_{u \in U} \in \Theta(\cdot + t, \omega \otimes_t X)\}. \end{aligned}$$

¹When restricted to the space of continuous functions $\mathbb{R}_+ \rightarrow F$, the Skorokhod J_1 topology coincides with the local uniform topology.

Standing Assumption 2.2. $\mathcal{C}(t, \omega) \neq \emptyset$ for all $(t, \omega) \in \llbracket 0, \infty \llbracket$.

Remark 2.3.

- (i) For every $(t, \omega) \in \llbracket 0, \infty \llbracket$, the set $\mathcal{C}(t, \omega)$ depends only on the path ω up to time t .
- (ii) Notice that

$$\mathcal{C}(t, \omega) = \left\{ P \in \mathfrak{P}(\Omega) : P(X = \omega \text{ on } [0, t]) = 1, \forall u \in U \ Y^u \in \mathcal{S}_{ac}^{sp}(t, P), \right. \\ \left. (\lambda \otimes P)\text{-a.e. } (dA^P(Y_{\cdot+t}^u)/d\lambda)_{u \in U} \in \Theta(\cdot + t, X) \right\}.$$

We define $\mathcal{C}(t, \omega)$ with $\Theta(\cdot + t, \omega \otimes_t X)$, instead of $\Theta(\cdot + t, X)$, to prevent confusion about measurability. Namely, both $(dA^P(Y_{\cdot+t}^u)/d\lambda)_{u \in U}$ and $\Theta(\cdot + t, \omega \otimes_t X)$ are measurable w.r.t. $\sigma(X_s, s \geq t)$, while $\Theta(\cdot + t, X)$ might also depend on $X_s, s < t$.

Discussion (Relation to Martingale Problems). The idea behind the set \mathcal{C} is inspired by the fact that many classes of stochastic processes can be modeled via martingale problems. Take a set $A \subset C_b(F; \mathbb{R}) \times B(F; \mathbb{R})$, which is often called *pregenerator*. Here, $B(F; \mathbb{R})$ denotes the set of all bounded Borel functions $F \rightarrow \mathbb{R}$. Then, following the monograph [18], a probability measure P on (Ω, \mathcal{F}) is said to solve the martingale problem associated to A if all processes of the form

$$(2.1) \quad f(X) - \int_0^\cdot g(X_s) ds, \quad (f, g) \in A,$$

are local P -martingales. To see the connection to the set \mathcal{C} , suppose that P solves the martingale problem associated to A . Then, for every $(f, g) \in A$, the process $f(X)$ is a special P -semimartingale, as it decomposes into the local P -martingale given in (2.1) and the continuous (hence, predictable) finite variation process $\int_0^\cdot g(X_s) ds$, which, in particular, means that $dA^P(f(X))/d\lambda = g(X)$. Roughly speaking, the set \mathcal{C} mimics the idea of the martingale problem but it allows some part of the pregenerator, namely the density of the compensator, to take values in the set Θ , which incorporates uncertainty. For the important special case where Θ is generated by a controlled coefficient, this interpretation is made explicit by the following proposition.

Proposition 2.4. *Suppose that G is a compact metrizable space and assume that there exists a predictable Carathéodory function $\mathfrak{z} = (\mathfrak{z}^u)_{u \in U} : G \times \llbracket 0, \infty \llbracket \rightarrow \mathbb{K}^U$, i.e., \mathfrak{z} is continuous in the first and predictable in the second variable, such that*

$$\Theta(t, \omega) = \{ \mathfrak{z}(g, t, \omega) : g \in G \}, \quad (t, \omega) \in \llbracket 0, \infty \llbracket.$$

Then, for every $\omega^ \in \Omega$ and $P \in \mathcal{C}(0, \omega^*)$, there exists a predictable function $\mathfrak{g} : \llbracket 0, \infty \llbracket \rightarrow G$ such that, for all $u \in U$, the process*

$$Y^u - Y_0^u - \int_0^\cdot \mathfrak{z}^u(\mathfrak{g}(s, X), s, X) ds$$

is a local P -martingale.

Proof. Let \mathcal{P} be the predictable σ -field. Thanks to [8, Lemma 2.12], the graph $\text{gr } \Theta$ is $\mathcal{P} \otimes \mathcal{B}(\mathbb{K}^U)$ -measurable. Thus,

$$H \triangleq \{ (t, \omega) \in \llbracket 0, \infty \llbracket : (t, \omega, (dA_t^P(Y^u)(\omega)/d\lambda)_{u \in U}) \notin \text{gr } \Theta \} \in \mathcal{P}.$$

For some arbitrary, but fixed, $g_0 \in G$, we set

$$\pi(t, \omega) \triangleq \begin{cases} \mathfrak{z}(g_0, t, \omega), & \text{if } (t, \omega) \in H, \\ (dA_t^P(Y^u)(\omega)/d\lambda)_{u \in U}, & \text{if } (t, \omega) \notin H. \end{cases}$$

As G is compact and \mathfrak{z} is a predictable Carathéodory function, the measurable implicit function theorem [1, Theorem 18.17] shows that the correspondence $\gamma: \llbracket 0, \infty \llbracket \rightarrow G$ defined by $\gamma(t, \omega) \triangleq \{g \in G: \mathfrak{z}(g, t, \omega) = \pi(t, \omega)\}$ is \mathcal{P} -measurable and that it admits a measurable selector. In other words, there exists a \mathcal{P} -measurable function $\mathfrak{g}: \llbracket 0, \infty \llbracket \rightarrow G$ such that $\pi(t, \omega) = \mathfrak{z}(\mathfrak{g}(t, \omega), t, \omega)$ for all $(t, \omega) \in \llbracket 0, \infty \llbracket$. Since $P \in \mathcal{C}(0, \omega^*)$, we have $(\lambda \otimes P)$ -a.e. $\pi = (dA^P(Y^u)/d\lambda)_{u \in U}$ and the claim follows. \square

Discussion (Relation to Control Rules). In the following, we relate our framework to controlled martingale problems as, e.g., discussed in [15, 16]. Take a compact Polish space Q and denote by $\mathbb{M}(\mathbb{R}_+ \times Q)$ the collection of all Radon measures on $\mathbb{R}_+ \times Q$. We define \mathbb{M} as the set of all measures in $\mathbb{M}(\mathbb{R}_+ \times Q)$ whose projections on \mathbb{R}_+ coincide with the Lebesgue measure. The identity map on \mathbb{M} is denoted by M . We define the σ -field $\mathcal{M} \triangleq \sigma(M_t(\phi); t \in \mathbb{R}_+, \phi \in C_c(\mathbb{R}_+ \times Q; \mathbb{R}))$, where

$$M_t(\phi) \triangleq \int_0^t \int_Q \phi(s, q) M(ds, dq).$$

Let H be a Polish space, which serves as state space, and take a countable set $\mathbb{G} \subset C_b(H; \mathbb{R}) \times B(\mathbb{R}_+ \times C(\mathbb{R}_+; H) \times Q; \mathbb{R})$. We denote the coordinate process on $C(\mathbb{R}_+; H)$ by X and set $\mathcal{B}(C(\mathbb{R}_+; H)) \triangleq \sigma(X_t, t \geq 0)$. For $(f, g) \in \mathbb{G}$, define

$$C(f, g) \triangleq f(X) - \int_0^\cdot \int_Q g(s, X_{\cdot \wedge s}, q) M(ds, dq).$$

A *control rule* with initial value $(t, \omega) \in \mathbb{R}_+ \times C(\mathbb{R}_+; H)$ is a probability measure P on the product space $(C(\mathbb{R}_+; H) \times \mathbb{M}, \mathcal{B}(C(\mathbb{R}_+; H)) \otimes \mathcal{M})$ such that $P(X = \omega \text{ on } [0, t]) = 1$ and under which the processes $(C_s(f, g))_{s \geq t}$, with $(f, g) \in \mathbb{G}$, are martingales for the filtration

$$\mathcal{F}_s^{(X, M)} \triangleq \sigma(X_r, M_r(\phi); r \leq s, \phi \in C_c(\mathbb{R}_+ \times Q; \mathbb{R})), \quad s \in \mathbb{R}_+.$$

Finally, we define

$$\mathcal{P}(t, \omega) \triangleq \{\text{all control rules with initial value } (t, \omega)\}.$$

By virtue of Proposition 2.4, both sets \mathcal{P} and \mathcal{C} are closely connected to martingale problems. In contrast to those for \mathcal{C} , the martingale problem related to \mathcal{P} is controlled in the sense that it depends on X and the second variable M . In other words, while the randomness in \mathcal{C} arises solely through the paths of X , exogenous randomness coming from the controls is added in \mathcal{P} .

Finally, let us explain how some kind of control problems can be modeled with our framework. Let \mathbb{V} be the set of all Radon measures on $(Q, \mathcal{B}(Q))$ endowed with the vague (or, equivalently, weak) topology, which turns it into a Polish space ([2, Theorem 31.5]). We use the product space $F \triangleq H \times \mathbb{V}$ to incorporate a control via the second coordinate. More specifically, this can be done with $U \triangleq \mathbb{G}$, $Y^u(X, Y) \triangleq f(X)$ for $u = (f, g) \in \mathbb{G}$ and $(X, Y) \in C(\mathbb{R}_+; H \times \mathbb{V})$, and

$$\Theta(t, \omega, \alpha) \triangleq \left\{ \left(\int_Q g(t, \omega(\cdot \wedge t), q) \alpha(t)(dq) \right)_{(f, g) \in U} \right\}$$

for $(t, \omega, \alpha) \in \mathbb{R}_+ \times C(\mathbb{R}_+; H) \times C(\mathbb{R}_+; \mathbb{V})$. We emphasize that although the correspondence Θ is singleton-valued, this is not necessarily the case for \mathcal{C} as the associated martingale problem might not be well-posed.

2.2. Dynamic Programming Principle. We now define a nonlinear conditional expectation and we provide the corresponding dynamic programming principle (DPP), i.e., the tower property of the nonlinear conditional expectation. Suppose that $\psi: \Omega \rightarrow \mathbb{R}$ is an upper semianalytic function, i.e., the set $\{\omega \in \Omega: \psi(\omega) > c\}$ is analytic for every $c \in \mathbb{R}$, and define the *value function* v by

$$v(t, \omega) \triangleq \sup_{P \in \mathcal{C}(t, \omega)} E^P[\psi], \quad (t, \omega) \in \llbracket 0, \infty \llbracket.$$

The following theorem is proved in Section 5.

Theorem 2.5 (Dynamic Programming Principle). *The value function v is upper semianalytic. Moreover, for every $(t, \omega) \in \llbracket 0, \infty \llbracket$ and every stopping time τ with $t \leq \tau < \infty$, we have*

$$(2.2) \quad v(t, \omega) = \sup_{P \in \mathcal{C}(t, \omega)} E^P[v(\tau, X)].$$

The value function v induces a nonlinear expectation \mathcal{E} via the formula

$$\mathcal{E}_t(\psi)(\omega) \triangleq v(t, \omega), \quad (t, \omega) \in \llbracket 0, \infty \llbracket,$$

and the DPP provides its tower property, i.e., (2.2) means that

$$\mathcal{E}_t(\psi)(\omega) = \mathcal{E}_t(\mathcal{E}_\tau(\psi))(\omega), \quad t \leq \tau \leq \infty.$$

For $x \in F$, we define

$$\begin{aligned} \mathcal{R}(x) \triangleq \{ & P \in \mathfrak{P}(\Omega): P(X_0 = x) = 1, \forall_{u \in U} Y^u \in \mathcal{S}_{\text{ac}}^{\text{sp}}(P), \\ & (\lambda \otimes P)\text{-a.e. } (dA^P(Y^u)/d\lambda)_{u \in U} \in \Theta \}, \end{aligned}$$

and

$$\mathcal{E}^x(\psi) \triangleq \sup_{P \in \mathcal{R}(x)} E^P[\psi].$$

Thinking of classical martingale problems, we interpret $\{\mathcal{E}^x: x \in F\}$ as a *nonlinear stochastic process*. In Section 3, we specify our setting to a Markovian situation and we establish more properties of $\{\mathcal{E}^x: x \in F\}$ to justify our interpretation. Before we start this program, let us discuss some important examples

2.3. Examples. In this section we illustrate that our framework covers the class of *nonlinear Lévy processes* from [38] and we introduce some new classes of nonlinear processes, namely the class of *nonlinear Markov chains*, *nonlinear spin systems* and the class of *nonlinear (in the sense of uncertainty) stochastic partial differential equations (NSPDEs)*.

2.3.1. Nonlinear Lévy processes. The class of nonlinear Lévy processes has been introduced in [38] via an uncertain set of Lévy–Khinchine triplets. We now explain how this class can be constructed with our framework. We emphasize that our discussion can be extended readily to more general classes of semimartingales under parametric uncertainty.

To fix the basic setting, we consider $F \triangleq \mathbb{R}^d$, $\mathbb{K} \triangleq \mathbb{C}$, $U \triangleq \mathbb{Q}^d$ and we take Ω to be space of all càdlàg functions $\mathbb{R}_+ \rightarrow F$. Similar as in [38], let $\tilde{\Theta}$ be a given set of Lévy–Khinchine triplets. Now, for every $u \in \mathbb{Q}^d$, we set $Y^u \triangleq e^{i\langle u, X \rangle}$ and we define

$$A^u(b, c, F) \triangleq -i\langle u, b \rangle + \frac{1}{2}\langle u, cu \rangle - \int (e^{i\langle u, x \rangle} - 1 - i\langle u, h(x) \rangle) F(dx),$$

and

$$\Theta(t, \omega) \triangleq \{(e^{i\langle u, \omega(t^-) \rangle} A^u(b, c, F))_{u \in U}: (b, c, F) \in \tilde{\Theta}\}, \quad (t, \omega) \in \llbracket 0, \infty \llbracket,$$

where $h: \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a fixed truncation function and i denotes the imaginary number.

We now relate this setting to those from [38]. First, for a given $P \in \mathcal{C}(0, x)$, we show that the coordinate process X is an \mathbb{R}^d -valued P -semimartingale whose characteristics (B^P, C^P, ν^P) are absolutely continuous with densities (b^P, c^P, F^P) such that $(\lambda \otimes P)$ -a.e. $(b^P, c^P, F^P) \in \tilde{\Theta}$. For every $u \in \mathbb{Q}^d$, we define

$$A^P(u) \triangleq -i\langle u, b^P \rangle + \frac{1}{2}\langle u, c^P u \rangle - \int (e^{i\langle u, x \rangle} - 1 - i\langle u, h(x) \rangle) F^P(dx).$$

Thanks to [26, Theorem II.2.42], since $P \in \mathcal{C}(0, x)$, $(\lambda \otimes P)$ -a.e. $(e^{i\langle u, X_{-} \rangle} A^P(u))_{u \in U} \in \Theta$. Take $(t, \omega) \in \llbracket 0, \infty \llbracket$ such that $(e^{i\langle u, \omega(t-) \rangle} A_t^P(u)(\omega))_{u \in U} \in \Theta(t, \omega)$. By definition of Θ , there exists a triplet $(b(t, \omega), c(t, \omega), F(t, \omega)) \in \tilde{\Theta}$ such that, for all $u \in \mathbb{Q}^d$,

$$A_t^P(u)(\omega) = A^u(b(t, \omega), c(t, \omega), F(t, \omega)).$$

The uniqueness lemma of Gnedenko and Kolmogorov ([26, Lemma II.2.44]) yields that

$$(b_t^P(\omega), c_t^P(\omega), F_t^P(\omega)) = (b(t, \omega), c(t, \omega), F(t, \omega)) \in \tilde{\Theta}.$$

Consequently, $(\lambda \otimes P)$ -a.e. $(b^P, c^P, F^P) \in \tilde{\Theta}$.

Conversely, if P is the law of a semimartingale (for its canonical filtration), starting at $x \in \mathbb{R}^d$, with absolutely continuous characteristics whose densities (b^P, c^P, F^P) are such that $(\lambda \otimes P)$ -a.e. $(b^P, c^P, F^P) \in \tilde{\Theta}$, then [26, Theorem II.2.42] yields that $P \in \mathcal{C}(0, x)$. We conclude that our setting includes the Lévy framework from [38].

2.3.2. Nonlinear Markov Chains. A (continuous-time) Markov chain is a strong Markov process on some countable discrete state space F . Typically, Markov chains are modeled via Q -matrices, which encode their infinitesimal dynamics. In particular, provided the Markov chain is a Feller–Dynkin process, the Q -matrix corresponds to the generator of the chain (see [47]). In the following we adapt the underlying martingale problem (see, e.g., [48, Theorem IV.20.6] or [51, Lemma 2.4]) to a nonlinear setting. For simplicity, we will restrict our attention to nonlinear Markov chains with a *finite* state space F .

Let $U \triangleq F \triangleq \{1, \dots, N\}$ with $N \in \mathbb{N}$, let Ω be the space of all càdlàg functions $\mathbb{R}_+ \rightarrow F$, and set $Y^u = \mathbf{1}\{X = u\}$ for $u \in U$. Denote by \mathbb{M}_q the set of all Q -matrices on F , i.e., the set of all matrices $Q = (q(k, n))_{k, n=1}^N \subset \mathbb{R}^{N \times N}$ such that $q(k, n) \geq 0$ for all $k \neq n$ and $Q\mathbf{1} = 0$. Let G be some topological space, let $Q = (q(k, n))_{k, n=1}^N: G \rightarrow \mathbb{M}_q$ be a Borel map and set, for $(t, \omega) \in \llbracket 0, \infty \llbracket$,

$$\Theta(t, \omega) \triangleq \{(q(\omega(t-), u)(g))_{u \in U}: g \in G\}.$$

In case G is the singleton $\{g_0\}$, then $\mathcal{C}(0, \omega)$ is also a singleton whose only element is the law of the continuous-time Markov chain with starting value $\omega(0)$ and Q -matrix $Q(g_0)$, see, e.g., [51, Lemma 2.4]. By virtue of Proposition 2.4, this observation confirms our interpretation as a nonlinear Markov chain, i.e., a Markov chain with uncertain Q -matrix.

2.3.3. Nonlinear Spin Systems. A spin system is an interacting particle system with totally disconnected state space $F \triangleq \{0, 1\}^S$ whose individual transitions only involve one site. Here, the site space S is some countable set and the state space F is endowed with the product discrete topology, which turns it into a compact Polish space. For profound introductions to the theory of spin systems we refer to the classical monographs [32, 33]. In the traditional setting without uncertainty, a spin system is constructed either as a

Feller–Dynkin process via a generator construction ([32, 33]), or via a martingale problem ([32]). In both cases, the second coordinate of the pregenerator has the form

$$\mathcal{L}(f)(\eta) = \sum_{x \in S} c(x, \eta) [f(\eta_x) - f(\eta)], \quad \eta \in \{0, 1\}^S = F,$$

where f is taken from a suitable class of functions. Here, $(c(x, \eta))_{x \in S, \eta \in F}$ is the so-called *speed function*. The value $c(x, \eta)$ is the rate the configuration $\eta \in F$ changes to $\eta_x \in F$, where latter equals η with the state x flipped, i.e.,

$$\eta_x(y) \triangleq \begin{cases} \eta(y), & y \neq x, \\ 1 - \eta(y), & y = x. \end{cases}$$

In the following we use the above description of the pregenerator to introduce *spin systems with uncertain speed functions*, which we call *nonlinear spin systems*. Let G be a topological space and let $c: G \times S \times F \rightarrow \mathbb{R}_+$ be a bounded Borel function, let Ω be the space of all càdlàg functions $\mathbb{R}_+ \rightarrow F$, set

$$D \triangleq \left\{ C(F; \mathbb{R}) : \sum_{x \in S} \sup_{\eta \in F} |f(\eta_x) - f(\eta)| < \infty \right\},$$

and let U be a countable subset of D . Finally, we set $Y^f \triangleq f(X)$ for $f \in U$, and

$$\Theta(t, \omega) \triangleq \left\{ \left(\sum_{x \in S} c(g, x, \omega(t-)) [f((\omega(t-))_x) - f(\omega(t-))] \right)_{f \in U} : g \in G \right\}.$$

Remark 2.6. By virtue of [8, Lemma 2.12], Standing Assumption 2.1 holds once G is a compact metrizable space and

$$G \times \{0, 1\}^S \ni (g, \eta) \mapsto \left(\sum_{x \in S} c(g, x, \eta) [f(\eta_x) - f(\eta)] \right)_{f \in U} \in \mathbb{R}^U$$

is a Carathéodory function, i.e., continuous in the first and Borel in the second variable. A condition for Standing Assumption 2.2 can be deduced from [32, Theorem I.6.7] or [33, Theorem 4.3].

Let us also provide two more explicit examples, namely a *nonlinear Ising model* and a *nonlinear contact process*.

Example 2.7 (Nonlinear Ising model). *Take $S \triangleq \mathbb{Z}^d$, let $\beta: G \rightarrow (0, \infty)$ be a Borel function, and let*

$$c(g, x, \eta) \triangleq \exp \left(-\beta(g) \sum_{\substack{y \in S \\ y \sim x}} [2\eta(x) - 1] [2\eta(y) - 1] \right),$$

where $(g, x, \eta) \in G \times S \times F$. The quantity $2\eta(x) - 1 \in \{-1, 1\}$ is the spin of an atom at site x , and $\beta(g)$ is the reciprocal temperature of the system. In contrast to the classical case, we allow the reciprocal temperature to take uncertain values in the set $\{\beta(g): g \in G\}$.

Example 2.8 (Nonlinear contact process). *Suppose that S is a graph of bounded degree, let $\lambda: G \rightarrow (0, \infty)$ be a Borel function and suppose that*

$$c(g, x, \eta) \triangleq \begin{cases} 1, & \eta(x) = 1, \\ \lambda(g) [\# \text{ neighbors } y \text{ of } x \text{ with } \eta(y) = 1], & \eta(x) = 0, \end{cases}$$

with $(g, x, \eta) \in G \times S \times F$. Typically, $\eta(x) = 1$ means that the site x is infected, while $\eta(x) = 0$ means that x is healthy. Healthy sites get affected at a rate that is proportional to the number of infected neighbors. The proportionality is given by the parameter $\lambda(g)$, which is now taken to be uncertain in our framework.

2.3.4. Nonlinear SPDEs. Let $(H_1, \|\cdot\|_{H_1}, \langle \cdot, \cdot \rangle_{H_1})$ and $(H_2, \|\cdot\|_{H_2}, \langle \cdot, \cdot \rangle_{H_2})$ be two separable Hilbert spaces over the real numbers, and let $(A, D(A))$ be the generator of a C_0 -semigroup on H_2 . In the following we introduce drift and volatility uncertainty to the class of (semi-linear) stochastic partial differential equations (SPDEs) of the type

$$dY_t = AY_t dt + \mu(Y_t) dt + \sigma(Y_t) dW_t,$$

where W is a cylindrical Brownian motion over H_1 and μ and σ are suitable measurable coefficients. To define our setting, we rely on the cylindrical martingale problem associated to semilinear SPDEs, see, e.g., [6, 11] for more details.

We denote the set of linear bounded operators from H_1 into H_2 by $L(H_1, H_2)$. Let $F \triangleq H_2$, let Ω be the space of continuous functions $\mathbb{R}_+ \rightarrow F$, and let $\mu: G \times H_2 \rightarrow H_2$ and $\sigma: G \times H_2 \rightarrow L(H_1, H_2)$ be Borel functions, where, for the latter, we mean that σh_1 is a Borel function for every $h_1 \in H_1$. The adjoint of $(A, D(A))$ is denoted by $(A^*, D(A^*))$. By virtue of [7, Lemma 7.3], there exists a countable set $D \subset D(A^*) \subset H_2$ such that, for every $y \in D(A^*)$, there exists a sequence $(y_n)_{n=1}^\infty \subset D$ such that $y_n \rightarrow y$ and $A^* y_n \rightarrow A^* y$. In other words, D is a countable subset of $D(A^*)$ which is dense in $D(A^*)$ for the graph norm. We set $f_i(x) \triangleq x^i$ for $x \in \mathbb{R}$ and $i = 1, 2$,

$$U \triangleq \{(H_2 \ni x \mapsto f_i(\langle y, x \rangle_{H_2})) : y \in D, i = 1, 2\}, \quad Y^u \triangleq u(X), \quad u \in U,$$

and

$$\mathcal{L}^{u,g}(x) \triangleq (\langle A^* y, x \rangle_{H_2} + \langle y, \mu(g, x) \rangle_{H_2}) f'_i(\langle y, x \rangle_{H_2}) + \frac{1}{2} \|\sigma^*(g, x) y\|_{H_1}^2 f''_i(\langle y, x \rangle_{H_2}),$$

for $(x, g) \in H_2 \times G$ and $u = f_i(\langle y, \cdot \rangle_{H_2}) \in U$. Finally, for $(t, \omega) \in \llbracket 0, \infty \llbracket$, we define

$$\Theta(t, \omega) \triangleq \{(\mathcal{L}^{u,g}(\omega(t)))_{u \in U} : g \in G\}.$$

In Lemma 10.2 below, we show that, under suitable assumptions on the coefficients, any $P \in \mathcal{R}(x)$ is the law of a mild solution processes (see [13, Section 6.1]) to an SPDE of the form

$$dY_t = AY_t dt + \mu(\mathbf{g}(t, Y), Y_t) dt + \sigma(\mathbf{g}(t, Y), Y_t) dW_t, \quad Y_0 = x,$$

where $\mathbf{g} = \mathbf{g}(P): \llbracket 0, \infty \llbracket \rightarrow G$ is a predictable function. This observation confirms our interpretation as an SPDE with uncertain coefficients.

3. NONLINEAR MARKOV PROCESSES

In this section we investigate a class of nonlinear Markov processes. That is, we study the family $\{\mathcal{E}^x : x \in F\}$ under the following standing assumption.

Standing Assumption 3.1. For every $(t, \omega) \in \llbracket 0, \infty \llbracket$, the set $\Theta(t, \omega)$ depends on (t, ω) only through $\omega(t-)$, i.e., $\Theta(t, \omega) \equiv \Theta(\omega(t-))$, where $\Theta: F \rightarrow \mathbb{K}^U$.

Our program in this section is the following. First, we explain the Markov property of $\{\mathcal{E}^x : x \in F\}$, which further leads to a nonlinear semigroup $(T_t)_{t \geq 0}$ on the cone of bounded upper semianalytic functions. Afterwards, we establish conditions for $(T_t)_{t \geq 0}$ to be a nonlinear semigroup on the cone of bounded upper semicontinuous functions and we provide a strong Markov selection principle.

3.1. The nonlinear Markov property. The following proposition provides the Markov property of $\{\mathcal{E}^x : x \in F\}$. The proof is given in Section 7.

Proposition 3.2. *For every upper semianalytic function $\psi : \Omega \rightarrow \mathbb{R}$, the equality*

$$\mathcal{E}^x(\psi \circ \theta_t) = \mathcal{E}^x(\mathcal{E}^{X_t}(\psi))$$

holds for every $(t, x) \in \mathbb{R}_+ \times F$.

To the best of our knowledge, the nonlinear Markov property was first observed in [22, Lemma 4.32] for nonlinear semimartingales, see also [9, Proposition 2.8] for a setting with continuous paths.

Similar to the fact that linear Markov processes have a canonical relation to linear semigroups, nonlinear Markov processes can be related to nonlinear semigroups.

Definition 3.3. *Let \mathcal{H} be a convex cone of functions $f : F \rightarrow \mathbb{R}$ containing all constant functions. A family of sublinear operators $T_t : \mathcal{H} \rightarrow \mathcal{H}$, $t \in \mathbb{R}_+$, is called a sublinear Markovian semigroup on \mathcal{H} if it satisfies the following properties:*

- (i) $(T_t)_{t \geq 0}$ has the semigroup property, i.e., $T_s T_t = T_{s+t}$ for all $s, t \in \mathbb{R}_+$ and $T_0 = \text{id}$;
- (ii) T_t is monotone for each $t \in \mathbb{R}_+$, i.e., $f, g \in \mathcal{H}$ with $f \leq g$ implies $T_t f \leq T_t g$;
- (iii) T_t preserves constants for each $t \in \mathbb{R}_+$, i.e., $T_t(c) = c$ for each $c \in \mathbb{R}$.

For a bounded upper semianalytic function $\psi : F \rightarrow \mathbb{R}$, $x \in F$ and $t \in \mathbb{R}_+$, we set

$$(3.1) \quad T_t(\psi)(x) \triangleq \mathcal{E}^x(\psi(X_t)) = \sup_{P \in \mathcal{R}(x)} E^P[\psi(X_t)].$$

The following proposition should be compared to [22, Remark 4.33] and [9, Proposition 2.9], where it has been established for nonlinear semimartingale frameworks. Its proof can be found in Section 8.

Proposition 3.4. *The family $(T_t)_{t \geq 0}$ defines a sublinear Markovian semigroup on the set of bounded upper semianalytic functions from F into \mathbb{R} .*

In the following section we extend Proposition 3.4 via conditions for $(T_t)_{t \geq 0}$ to be a nonlinear semigroup on the space of upper semicontinuous functions from F into \mathbb{R} .

3.2. The USC_b -Feller Property. It is natural to ask whether the nonlinear semigroup $(T_t)_{t \geq 0}$ preserves some regularity. In case F is endowed with the discrete topology, as it is the case for the class of nonlinear Markov chains from Section 2.3.2 and the class of nonlinear spin systems from Section 2.3.3 when the site space S is finite, it is trivial that T_t , for every $t \in \mathbb{R}_+$, is a selfmap on the space of bounded continuous functions. In general, however, such a preservation property is non-trivial to establish. In the following we provide general conditions for a type of regularity preservation property of nonlinear stochastic processes with *continuous* paths, namely the so-called *USC_b -Feller property*. Before we formulate our conditions, let us introduce a last bit of notation. We fix an \mathbb{R}_+ -valued continuous adapted process $L = (L_t)_{t \geq 0}$ on $(\Omega, \mathcal{F}, \mathbf{F})$ and, for $M > 0$, we set

$$\rho_M(\omega) \triangleq \inf\{t \geq 0 : L_t \geq M\} \wedge M, \quad \omega \in \Omega.$$

When Ω is the Wiener space of continuous paths, a typical choice for L could be $d_F(X, y_0)$, where d_F is a metric on F and $y_0 \in F$ is an arbitrary reference point.

Condition 3.5.

- (i) The underlying path space Ω is the space of continuous functions $\mathbb{R}_+ \rightarrow F$, for every $u \in U$, the process Y^u has continuous paths and $\Omega \ni \omega \mapsto Y^u(\omega) \in C(\mathbb{R}_+; \mathbb{K})$ and $\Omega \ni \omega \mapsto L(\omega) \in C(\mathbb{R}_+; \mathbb{R})$ are continuous (where the image and the inverse image spaces are endowed with the local uniform topology). Furthermore, for every $u \in U, M > 0$ and any compact set $K \subset F$, the process $Y^u \wedge \rho_M \mathbb{1}_{\{X_0 \in K\}}$ is bounded.
- (ii) The correspondence $\Theta: F \rightarrow \mathbb{K}^U$ is convex-valued.
- (iii) For every $M > 0$, there exists a family $\{K^{u,M}: u \in U\} \subset \mathbb{K}$ of bounded sets such that $\Theta(\omega(t)) \subset \prod_{u \in U} K^{u,M}$ for all $(t, \omega) \in \llbracket 0, \rho_M \llbracket$. Moreover, for every compact set $K \subset F$ and every $T > 0$, there exists a number $q = q(K, T) > 0$ such that

$$\sup_{x \in K} \sup_{P \in \mathcal{R}(x)} E^P \left[\sup_{s \in [0, T]} |L_s|^q \right] < \infty.$$

- (iv) For every $\omega \in \Omega$, the correspondence $\mathbb{R}_+ \ni t \mapsto \Theta(\omega(t))$ is upper hemicontinuous and compact-valued, and, for every $(t, \omega) \in \llbracket 0, \infty \llbracket$, the closed convex hull of

$$\Theta(\omega(\llbracket t, t+1 \llbracket)) \triangleq \bigcup_{s \in \llbracket t, t+1 \llbracket} \Theta(\omega(s))$$

is compact.

- (v) For every compact set $K \subset F$, the set $\bigcup_{x \in K} \mathcal{R}(x) \subset \mathfrak{P}(\Omega)$ is relatively compact.

For nonlinear one-dimensional diffusions, a version of the following theorem was established in [9]. Moreover, for nonlinear semimartingales *with jumps* some conditions were proved in [22, Theorem 4.41, Lemma 4.42]. We provide a result which reaches beyond semimartingale settings. Its proof can be found in Section 9.

Theorem 3.6. *Suppose that Condition 3.5 holds. Then, $(T_t)_{t \geq 0}$ is a nonlinear semigroup on the space $USC_b(F; \mathbb{R})$ of bounded upper semicontinuous functions $F \rightarrow \mathbb{R}$.*

Remark 3.7. It is interesting to note that Condition 3.5 is not sufficient for the C_b -Feller property of $(T_t)_{t \geq 0}$, i.e., it does not imply $T_t(C_b(F; \mathbb{R})) \subset C_b(F; \mathbb{R})$ for all $t > 0$. We refer to [50, Exercise 12.4.2] for a counterexample.

In Section 3.4 below, we return to the class of NSPDEs from Section 2.3.4 and we provide some general parametric conditions which imply Condition 3.5. From a practical point of view, (i) – (iv) from Condition 3.5 are structural assumptions, while checking part (v) requires some argument.

3.3. The strong Markov selection principle. For a probability measure P on (Ω, \mathcal{F}) , a kernel $\Omega \ni \omega \mapsto Q_\omega \in \mathfrak{P}(\Omega)$, and a finite stopping time τ , we define the pasting measure

$$(P \otimes_\tau Q)(A) \triangleq \iint \mathbb{1}_A(\omega \otimes_{\tau(\omega)} \omega') Q_\omega(d\omega') P(d\omega)$$

for all $A \in \mathcal{F}$.

Definition 3.8 (Time inhomogeneous Markov Family). *A family $\{P_{(s,x)}: (s,x) \in \mathbb{R}_+ \times F\} \subset \mathfrak{P}(\Omega)$ is said to be a strong Markov family if $(t,x) \mapsto P_{(t,x)}$ is Borel and the strong Markov property holds, i.e., for every $(s,x) \in \mathbb{R}_+ \times F$ and every finite stopping time $\tau \geq s$,*

$$P_{(s,x)}(\cdot | \mathcal{F}_\tau)(\omega) = \omega \otimes_{\tau(\omega)} P_{(\tau(\omega), \omega(\tau(\omega)))}$$

for $P_{(s,x)}$ -a.a. $\omega \in \Omega$.

Further, we introduce a correspondence $\mathcal{K}: \mathbb{R}_+ \times F \rightarrow \mathfrak{P}(\Omega)$ by

$$\mathcal{K}(t, x) \triangleq \{P \in \mathfrak{P}(\Omega): P(X_s = x \text{ for all } s \in [0, t]) = 1, \forall u \in U \ Y^u \in \mathcal{S}_{\text{ac}}^{\text{SP}}(t, P), \\ (\lambda \otimes P)\text{-a.e. } (dA^P(Y_{\cdot+t}^u)/d\lambda)_{u \in U} \in \Theta(\cdot + t, X)\},$$

where $(t, x) \in \mathbb{R}_+ \times F$.

Theorem 3.9 (Strong Markov Selection Principle). *Suppose that Condition 3.5 holds. For every $\psi \in USC_b(\mathbb{R}; F)$ and every $t > 0$, there exists a strong Markov family $\{P_{(s,x)}: (s, x) \in \mathbb{R}_+ \times F\}$ such that, for all $(s, x) \in \mathbb{R}_+ \times F$, $P_{(s,x)} \in \mathcal{K}(s, x)$ and*

$$E^{P_{(s,x)}}[\phi(X_t)] = \sup_{P \in \mathcal{K}(s,x)} E^P[\phi(X_t)].$$

In particular, for every $x \in F$,

$$T_t(\psi)(x) = E^{P_{(0,x)}}[\psi(X_t)].$$

To the best of our knowledge, the first strong Markov selection principle for nonlinear processes appeared in [9] for nonlinear one-dimensional diffusions.

In the following section we tailor the Theorems 3.6 and 3.9 to the NSPDE setting from Section 2.3.4. Furthermore, we are going to discuss the strong USC_b -Feller property, which provides a smoothing effect.

3.4. NSPDEs: USC_b -Feller Properties and Strong Markov Selections. We return to the class of nonlinear SPDEs from Section 2.3.4 and provide explicit conditions (in terms of the drift and diffusion coefficients) for the USC_b -Feller property and the strong Markov selection principle.

We recall the precise setting. Let $(H_1, \|\cdot\|_{H_1}, \langle \cdot, \cdot \rangle_{H_1})$ and $(H_2, \|\cdot\|_{H_2}, \langle \cdot, \cdot \rangle_{H_2})$ be two separable Hilbert spaces over the real numbers, and let $(A, D(A))$ be the generator of a C_0 -semigroup $(S_t)_{t \geq 0}$ on the Hilbert space H_2 . Moreover, let Ω be the space of all continuous functions $\mathbb{R}_+ \rightarrow F$, and let $\mu: G \times H_2 \rightarrow H_2$ and $\sigma: G \times H_2 \rightarrow L(H_1, H_2)$ be Borel functions. Further, let $D \subset D(A^*)$ be a countable set such that for every $y \in D(A^*)$ there exists a sequence $(y_n)_{n=1}^\infty \subset D$ such that $y_n \rightarrow y$ and $A^*y_n \rightarrow A^*y$. Recall that such a set exists by [7, Lemma 7.3]. We set $f_i(x) \triangleq x^i$ for $x \in \mathbb{R}$ and $i = 1, 2$,

$$U \triangleq \{(H_2 \ni x \mapsto f_i(\langle y, x \rangle_{H_2})): y \in D, i = 1, 2\}, \quad Y^u \triangleq u(X), \quad u \in U,$$

and

$$\mathcal{L}^{u,g}(x) \triangleq (\langle A^*y, x \rangle_{H_2} + \langle y, \mu(g, x) \rangle_{H_2}) f'_i(\langle y, x \rangle_{H_2}) + \frac{1}{2} \|\sigma^*(g, x)y\|_{H_1}^2 f''_i(\langle y, x \rangle_{H_2}),$$

for $(x, g) \in H_2 \times G$ and $u = f_i(\langle y, \cdot \rangle_{H_2}) \in U$. Finally, for $x \in H_2$ and $(t, \omega) \in \llbracket 0, \infty \llbracket$, we define

$$\Theta(x) \triangleq \{(\mathcal{L}^{u,g}(x))_{u \in U}: g \in G\}, \quad \Theta(t, \omega) \triangleq \Theta(\omega(t)).$$

We denote the operator norm on $L(H_1, H_2)$ by $\|\cdot\|_{L(H_1, H_2)}$ and the Hilbert-Schmidt norm by $\|\cdot\|_{L_2(H_1, H_2)}$.

Condition 3.10.

- (i) G is a compact metrizable space.
- (ii) For every $x \in H_2$, the set $\{(\mu(g, x), (\sigma\sigma^*)(g, x)): g \in G\}$ is convex.
- (iii) For every $y \in D$, the functions $\langle y, \mu \rangle_{H_2}$ and $\|\sigma^*y\|_{H_1}$ are continuous.

(iv) *There exists a constant $C > 0$ such that*

$$\|\mu(g, x)\|_{H_2} + \|\sigma(g, x)\|_{L(H_1, H_2)} \leq C(1 + \|x\|_{H_2})$$

for all $(g, x) \in G \times H_2$.

(v) *The semigroup S is compact, i.e., S_t is compact for every $t > 0$, and there exists an $\alpha \in (0, 1/2)$ and a Borel function $f: (0, \infty) \rightarrow [0, \infty]$ such that*

$$\int_0^T \left[\frac{f(s)}{s^\alpha} \right]^2 ds < \infty, \quad \forall T > 0,$$

and, for all $t > 0, x \in H_2$ and $g \in G$,

$$\|S_t \sigma(g, x)\|_{L_2(H_1, H_2)} \leq f(t)(1 + \|x\|_{H_2}).$$

Lemma 3.11. *If Condition 3.10 is in force, both Standing Assumptions 2.1 and 2.2 are satisfied.*

Proof. Thanks to (i) and (iii) of Condition 3.10, Standing Assumption 2.1 is implied by [8, Lemma 2.12]. Moreover, thanks to (iii) – (v) from Condition 3.10, Standing Assumption 2.2 follows from [7, Theorem 2.5] and Lemmata 6.3 and 6.9 below. \square

The following proposition is proved in Section 10.

Proposition 3.12. *Condition 3.10 implies Condition 3.5 with $L \equiv \|X\|_{H_2}$.*

Thanks to Proposition 3.12, the Theorems 3.6 and 3.9 yield the following result.

Corollary 3.13. *Suppose that Condition 3.10 holds. Then, $(T_t)_{t \geq 0}$ is a nonlinear semigroup on $USC_b(H_2; \mathbb{R})$ and, for every $\psi \in USC_b(H_2; \mathbb{R})$ and every $t > 0$, there exists a strong Markov family $\{P_{(s,x)}: (s,x) \in \mathbb{R}_+ \times H_2\}$ such that, for every $(s,x) \in \mathbb{R}_+ \times H_2$, $P_{(s,x)} \in \mathcal{K}(s,x)$ and*

$$E^{P_{(s,x)}}[\phi(X_t)] = \sup_{P \in \mathcal{K}(s,x)} E^P[\phi(X_t)].$$

In particular, $T_t(\psi)(x) = E^{P_{(0,x)}}[\psi(X_t)]$ for every $x \in H_2$.

Finally, we establish a smoothing effect for stochastic nonlinear Cauchy problems with drift uncertainty, which are infinite-dimensional processes of the form

$$dY_t = AY_t dt + \mu(Y_t) dt + dW_t,$$

with drift uncertainty.

Definition 3.14. *We say that $(T_t)_{t \geq 0}$ has the strong USC_b -Feller property if*

$$T_t(USC_b(F; \mathbb{R})) \subset C_b(F; \mathbb{R})$$

for every $t > 0$.

To the best of our knowledge, the strong USC_b -Feller property of nonlinear semigroups was first discovered in [9] for nonlinear one-dimensional elliptic diffusions. In [10], it was established for nonlinear multidimensional diffusions with fixed (strongly) elliptic diffusion coefficients. Adapting some ideas from [9, 10], we establish the first result for infinite-dimensional nonlinear stochastic processes.

Condition 3.15.

- (i) *G is a compact metrizable space and $H_1 \equiv H_2 \equiv H$.*
- (ii) *For every $x \in H$, the set $\{\mu(g, x): g \in G\}$ is convex.*

- (iii) For every $y \in D$, $\langle y, \mu \rangle_H$ is continuous and σ equals constantly the identity operator on H , i.e., $\sigma(g, x) \equiv \text{id}$ for all $(g, x) \in G \times H$.
- (iv) There exists a constant $C > 0$ such that

$$\|\mu(g, x)\|_H \leq C(1 + \|x\|_H)$$

for all $(g, x) \in G \times H$.

- (v) There exists an $\alpha \in (0, 1/2)$ such that

$$\int_0^T \frac{\|S_s\|_{L_2(H,H)}^2 ds}{s^{2\alpha}} < \infty, \quad \forall T > 0.$$

Remark 3.16 ([6]). Part (v) of Condition 3.15 implies that S is a compact semigroup. Moreover, the condition

$$\int_0^T \frac{\|S_s\|_{L_2(H,H)}^2 ds}{s^{2\alpha}} < \infty$$

holds for all $T > 0$ once it holds for some $T > 0$.

The proof for the next theorem is given in Section 11.

Theorem 3.17. *If Condition 3.15 holds, then $(T_t)_{t \geq 0}$ has the strong USC_b-Feller property.*

Under suitable additional conditions (see [13, Hypothesis 9.1]), Theorem 3.17 can be extended to NSPDEs with *certain* diffusion coefficient, i.e., to the case $\sigma(g, x) \equiv \sigma(x)$.

Example 3.18 ([20]). *Let $\mathcal{O} \subset \mathbb{R}^d$ be a bounded domain with smooth boundary. Part (v) of Condition 3.15 holds when $H = L^2(\mathcal{O})$ and A is a strongly elliptic operator of order $2m > d$, see [20, Remark 3, Example 3] for some details. In particular, this is the case when $d = 1$ and A is the Laplacian, i.e., Condition 3.15 holds for one-dimensional stochastic heat equations with drift uncertainty.*

4. ERGODICITY OF NONLINEAR SPIN SYSTEMS

In this section we return to the setting from Section 2.3.3 and we investigate ergodicity of nonlinear spin systems with finitely many sites. First of all, we recall our precise setting.

Take a topological space G , a finite set S , which we endow with the discrete topology, and a Borel function $c: G \times S \times F \rightarrow \mathbb{R}_+$, where $F \triangleq \{0, 1\}^S$ is also endowed with the discrete topology. Let Ω be the space of all càdlàg functions from \mathbb{R}_+ into F .

Standing Assumption 4.1. G is a compact metrizable space and $g \mapsto c(g, x, \eta)$ is continuous for every $(x, \eta) \in S \times F$.

Let $U \subset C(F; \mathbb{R}) = \mathbb{R}^F$ be a countable set which is dense in the uniform (equivalently, product) topology. Define $Y^f \triangleq f(X)$ for $f \in U$ and set

$$\Theta(\eta) \triangleq \left\{ \left(\sum_{x \in S} c(g, x, \eta) [f(\eta_x) - f(\eta)] \right)_{f \in U} : g \in G \right\}$$

for $\eta \in F$. Finally, in the spirit of Standing Assumption 3.1, we define $\Theta(t, \omega) \triangleq \Theta(\omega(t-))$ for $(t, \omega) \in \llbracket 0, \infty \llbracket$.

Lemma 4.2. *The correspondence $(t, \omega) \mapsto \Theta(t, \omega)$ has a measurable graph, i.e., Standing Assumption 2.1 holds. Moreover, $\mathcal{C}(t, \omega) \neq \emptyset$ for all $(t, \omega) \in \llbracket 0, \infty \llbracket$, i.e., Standing Assumption 2.2 holds.*

Proof. The measurable graph property follows from [8, Lemma 2.12]. The second part follows from [33, Theorem 4.3] (or [33, Corollary 2.34, Theorem 2.37]) and Lemmata 6.3 and 6.9 below. \square

By virtue of Lemma 4.2 and Propositions 3.2 and 3.4, the nonlinear spin system $(T_t)_{t \geq 0}$ given by (3.1) is well-defined as a nonlinear Markov process.

For $\eta, \xi \in F$, we write $\xi \leq \eta$ if $\xi(x) \leq \eta(x)$ for all $x \in S$. Furthermore, a function $f \in C(F; \mathbb{R})$ is called *increasing* in case $\xi \leq \eta$ implies that $f(\xi) \leq f(\eta)$. The set of increasing functions $F \rightarrow \mathbb{R}$ is denoted by $C_\uparrow(F; \mathbb{R})$. The set $C_\uparrow(F; \mathbb{R})$ is measure determining ([33, Exercise 4.16]).

Definition 4.3. Let $(S_t)_{t \geq 0}$ be a nonlinear semigroup on $C(F; \mathbb{R})$.

- (i) A probability measure μ on $(F, \mathcal{B}(F))$ is called an invariant distribution for $(S_t)_{t \geq 0}$ if

$$\int S_t f d\mu = \int f d\mu$$

for all $t \in \mathbb{R}_+$ and $f \in C_\uparrow(F; \mathbb{R})$. The set of invariant distributions is denoted by \mathcal{I} .

- (ii) The semigroup $(S_t)_{t \geq 0}$ is said to be ergodic if $\mathcal{I} = \{\mu\}$ and

$$(4.1) \quad \lim_{t \rightarrow \infty} \int S_t f d\nu = \int f d\mu$$

for every $f \in C_\uparrow(F; \mathbb{R})$ and all probability measures ν on $(F, \mathcal{B}(F))$.

Remark 4.4.

- (i) We emphasize that invariant distributions are *measures*, i.e., they correspond to *linear* expectations. Moreover, as the set $C_\uparrow(F; \mathbb{R})$ is measure determining, a nonlinear semigroup $(S_t)_{t \geq 0}$ is ergodic if and only if there exists an invariant distribution μ such that (4.1) holds for all $(f, \nu) \in C_\uparrow(F; \mathbb{R}) \times \mathfrak{P}(F)$. Indeed, whenever μ^* is another invariant distribution, (4.1) yields that $\int f d\mu^* = \int f d\mu$ for all $f \in C_\uparrow(F; \mathbb{R})$ and hence, $\mu^* = \mu$.
- (ii) For *linear* C_0 -semigroups on $C(F; \mathbb{R})$ it is well-known that an invariant distribution exists due to the compactness of F ([33, Theorem 3.38]). The classical proof for this fact is based on a Cesàro mean approximation. This argument cannot be used in our nonlinear setting, as, due to the nonlinearity of the semigroup, the Cesàro means are not measures.

In the following we investigate ergodicity of the nonlinear spin system $(T_t)_{t \geq 0}$.

Definition 4.5 (Linear Spin System). A linear C_0 -semigroup $(S_t)_{t \geq 0}$ on $C(F; \mathbb{R})$ is said to be a linear spin system with speed function $(\tilde{c}(x, \eta))_{x \in S, \eta \in F}$ if its generator is given by

$$\mathcal{L}(f)(\eta) = \sum_{x \in S} \tilde{c}(x, \eta) [f(\eta_x) - f(\eta)], \quad (f, \eta) \in C(F; \mathbb{R}) \times F.$$

We define two auxiliary speed functions by

$$\bar{c}(x, \eta) \triangleq \begin{cases} \sup_{g \in G} c(g, x, \eta), & \eta(x) = 0, \\ \inf_{g \in G} c(g, x, \eta), & \eta(x) = 1, \end{cases}$$

$$\underline{c}(x, \eta) \triangleq \begin{cases} \inf_{g \in G} c(g, x, \eta), & \eta(x) = 0, \\ \sup_{g \in G} c(g, x, \eta), & \eta(x) = 1. \end{cases}$$

Lemma 4.6. *There exist two linear spin systems $(\overline{T}_t)_{t \geq 0}$ and $(\underline{T}_t)_{t \geq 0}$ with speed functions \overline{c} and \underline{c} , respectively.*

Proof. This follows from [33, Theorem 4.3] (or [33, Corollary 2.34, Theorem 2.37]). \square

Let us further recall a classical terminology for spin systems ([32, 33]).

Definition 4.7. *A linear spin system with speed function $(\tilde{c}(x, \eta))_{x \in S, \eta \in F}$ is called attractive if*

$$\begin{aligned} \tilde{c}(x, \eta) &\leq \tilde{c}(x, \xi) \text{ for } \eta(x) = \xi(x) = 0, \\ \tilde{c}(x, \eta) &\geq \tilde{c}(x, \xi) \text{ for } \eta(x) = \xi(x) = 1, \end{aligned}$$

whenever $\xi \leq \eta$.

We are in the position to state the main result of this section, whose proof is given in Section 12.

Theorem 4.8. *The nonlinear spin system $(T_t)_{t \geq 0}$ is ergodic if the linear spin systems $(\overline{T}_t)_{t \geq 0}$ and $(\underline{T}_t)_{t \geq 0}$ are attractive and ergodic with the same invariant distribution, which then is also the (unique) invariant distribution of the nonlinear spin system $(T_t)_{t \geq 0}$.*

The idea behind Theorem 4.8 is to wedge the nonlinear spin system $(T_t)_{t \geq 0}$ between the linear spin systems $(\underline{T}_t)_{t \geq 0}$ and $(\overline{T}_t)_{t \geq 0}$ via stochastic domination. Stochastic order properties are often used for the study of attractive linear spin systems (see, e.g., [31, Section 2.3] for an application in the spirit of Theorem 4.8). In contrast to the classical approach for the linear case, where coupling techniques for Markov processes are the key, to obtain the desired stochastic domination, we use a stochastic calculus approach, which we learned from [3]. To illustrate an application of the previous theorem, we provide a condition for ergodicity of the nonlinear contact process.

Corollary 4.9. *Provided $\lambda: G \rightarrow (0, \infty)$ is continuous, the nonlinear contact process from Example 2.8 is ergodic if*

$$\sup_{g \in G} \lambda(g) < 1 / \max_{x \in S} \deg(x),$$

and the unique invariant distribution is given by δ_0 , i.e., the Dirac measure on the configuration without infected sites.

Proof. Notice that \overline{c} and \underline{c} are speed functions of contact processes with parameters $\sup \lambda$ and $\inf \lambda$, respectively. Consequently, in case $\sup \lambda < 1 / \max \deg$, it is well-known that $(\overline{T}_t)_{t \geq 0}$ and $(\underline{T}_t)_{t \geq 0}$ are attractive and ergodic with (unique) invariant distribution δ_0 (see [33, Example 4.7]). Thus, the claim follows from Theorem 4.8. \square

5. PROOF OF THE DYNAMIC PROGRAMMING PRINCIPLE: THEOREM 2.5

Recall that the Standing Assumptions 2.1 and 2.2 are in force. The proof of Theorem 2.5 is based on an application of the general [16, Theorem 2.1], which provides three abstract conditions on the set \mathcal{C} which imply the DPP. In the following we verify these conditions. For reader's convenience, let us restate them.

- (i) *Measurable graph condition:* The set $\{(t, \omega, P) \in [0, \infty[\times \mathfrak{P}(\Omega) : P \in \mathcal{C}(t, \omega)\}$ is Borel.²

²In fact, to use the result from [16] it suffices that this set is analytic.

- (ii) *Stability under conditioning:* For any $t \in \mathbb{R}_+$, any stopping time τ with $t \leq \tau < \infty$, and any $P \in \mathcal{C}(t, \alpha)$ there exists a family $\{P(\cdot | \mathcal{F}_\tau)(\omega) : \omega \in \Omega\}$ of regular P -conditional probabilities given \mathcal{F}_τ such that P -a.s. $P(\cdot | \mathcal{F}_\tau) \in \mathcal{C}(\tau, X)$.
- (iii) *Stability under pasting:* For any $t \in \mathbb{R}_+$, any stopping time τ with $t \leq \tau < \infty$, any $P \in \mathcal{C}(t, \alpha)$ and any \mathcal{F}_τ -measurable map $\Omega \ni \omega \mapsto Q_\omega \in \mathfrak{P}(\Omega)$ the following implication holds:

$$P\text{-a.s. } Q \in \mathcal{C}(\tau, X) \implies P \otimes_\tau Q \in \mathcal{C}(t, \alpha).$$

In the following three sections we check these properties. In the fourth (and last) section, we finalize the proof of Theorem 2.5.

5.1. Measurable graph condition. The proof of the measurable graph condition is split into several parts.

Lemma 5.1. *For every \mathbb{K} -valued càdlàg \mathbf{F} -adapted process Y^* , the set*

$$\mathfrak{P}_{\text{sp sem}}^{\text{ac}}(Y^*) \triangleq \{P \in \mathfrak{P}(\Omega) : Y^* \in \mathcal{S}_{\text{ac}}^{\text{sp}}(P)\}$$

is Borel. Moreover, there exists a Borel map

$$[0, \infty[\times \mathfrak{P}_{\text{sp sem}}^{\text{ac}}(Y^*) \ni (t, \omega, P) \mapsto a_t^P(Y^*)(\omega) \in \mathbb{K}$$

such that, for every $P \in \mathfrak{P}_{\text{sp sem}}^{\text{ac}}(Y^)$, a^P is predictable and $(\lambda \otimes P)$ -a.e. $a^P(Y^*) = d\mathbf{A}^P(Y^*)/d\lambda$.*

Proof. For $P \in \mathfrak{P}(\Omega)$, denote the set of \mathbb{K} -valued P - \mathbf{F}_+ -semimartingales by $\mathcal{S}(P)$. The set $\mathfrak{P}_{\text{sem}} \triangleq \{P \in \mathfrak{P}(\Omega) : Y^* \in \mathcal{S}(P)\}$ is Borel by [37, Theorem 2.5]. Furthermore, by the very same theorem, there are Borel maps (with values in a suitable Polish space)

$$\mathfrak{P}_{\text{sem}} \times \Omega \times \mathbb{R}_+ \ni (P, \omega, t) \mapsto (B_t^P(\omega), C_t(\omega), \nu^P(\omega))$$

such that (B^P, C, ν^P) are the P - \mathbf{F}_+ -semimartingale characteristics of Y^* (corresponding to a given truncation function h). By virtue of [26, Proposition II.2.29], we have

$$\{P \in \mathfrak{P}(\Omega) : Y^* \in \mathcal{S}^{\text{sp}}(P)\} = \{P \in \mathfrak{P}(\Omega) : Y^* \in \mathcal{S}(P), \forall t \in \mathbb{N} (|x|^2 \wedge |x|) * \nu_t^P < \infty\},$$

where, as usual,

$$(|x|^2 \wedge |x|) * \nu_t^P \triangleq \int (|x|^2 \wedge |x|) \nu^P([0, t] \times dx).$$

Consequently, $\{P \in \mathfrak{P}(\Omega) : Y^* \in \mathcal{S}^{\text{sp}}(P)\}$ is Borel. Thanks again to [26, Proposition II.2.29], in case $Y^* \in \mathcal{S}^{\text{sp}}(P)$, the predictable part of (locally) finite variation in the semimartingale decomposition is given by

$$(5.1) \quad \mathbf{A}^P(Y^*) = B^P + (x - h(x)) * \nu^P.$$

Hence,

$$\{P \in \mathfrak{P}(\Omega) : Y^* \in \mathcal{S}_{\text{ac}}^{\text{sp}}(P)\} = \{P \in \mathfrak{P}(\Omega) : Y^* \in \mathcal{S}^{\text{sp}}(P), B^P + (x - h(x)) * \nu^P \ll \lambda\},$$

and it follows as in the proof of [37, Lemma 7.1] that the latter set is Borel.

Finally, the existence of the Borel map $(t, \omega, P) \mapsto a_t^P(Y^*)(\omega)$ follows from (5.1) and [37, Theorem 2.6]. The proof is complete. \square

Lemma 5.2. *The map $(t, P) \mapsto P \circ \theta_t^{-1} \triangleq P_t$ is Borel.*

Proof. As the map $(t, \omega) \mapsto \theta_t(\omega)$ is Borel, the claim follows from [5, Theorem 8.10.61]. \square

The following lemma is a version of [8, Lemma 3.3] for càdlàg F -valued, instead of continuous real-valued, processes. The proof is verbatim the same and omitted here.

Lemma 5.3. *Let $Y = (Y_t)_{t \geq 0}$ be an F -valued càdlàg process and set $\mathcal{F}_t^Y \triangleq \sigma(Y_s, s \leq t)$ for $t \in \mathbb{R}_+$. Then, $Y^{-1}(\mathcal{F}_{s+}) = \mathcal{F}_{s+}^Y$ for all $s \in \mathbb{R}_+$.*

The following lemma is a restatement of [25, Lemma 2.9 a)].

Lemma 5.4. *Take two filtered probability spaces $\mathbb{B}^* = (\Omega^*, \mathcal{F}^*, \mathbf{F}^* = (\mathcal{F}_t^*)_{t \geq 0}, P^*)$ and $\mathbb{B}' = (\Omega', \mathcal{F}', \mathbf{F}' = (\mathcal{F}_t')_{t \geq 0}, P')$ with right-continuous filtrations and the property that there is a map $\phi: \Omega' \rightarrow \Omega^*$ such that $\phi^{-1}(\mathcal{F}^*) \subset \mathcal{F}'$, $P^* = P' \circ \phi^{-1}$ and $\phi^{-1}(\mathcal{F}_t^*) = \mathcal{F}_t'$ for all $t \in \mathbb{R}_+$. Then, the following hold: X^* is a d -dimensional semimartingale on \mathbb{B}^* if and only if $X' = X^* \circ \phi$ is a d -dimensional semimartingale on \mathbb{B}' . Moreover, (B^*, C^*, ν^*) are the characteristics of X^* if and only if $(B^* \circ \phi, C^* \circ \phi, \nu^* \circ \phi)$ are the characteristics of $X' = X^* \circ \phi$.*

Lemma 5.5. *The set*

$$\left\{ (t, \omega, P) \in \llbracket 0, \infty \llbracket \times \mathfrak{P}(\Omega) : \forall_{u \in U} Y^u \in \mathcal{S}_{\text{ac}}^{\text{SP}}(t, P), \right. \\ \left. (\lambda \otimes P)\text{-a.e. } (dA^P(Y_{\cdot+t}^u)/d\lambda)_{u \in U} \in \Theta(\cdot + t, \omega \otimes_t X) \right\}$$

is Borel.

Proof. For $\omega, \omega' \in \Omega$ and $t \in \mathbb{R}_+$, we define the concatenation

$$\omega \tilde{\otimes}_t \omega' \triangleq \omega \mathbf{1}_{[0, t)} + (\omega(t) + \omega'(\cdot - t) - \omega'(0)) \mathbf{1}_{[t, \infty)},$$

and we notice that $(\omega \tilde{\otimes}_t X) \circ \theta_t = \omega \otimes_t X$ for all $(t, \omega) \in \llbracket 0, \infty \llbracket$. Further, recall that, for every $u \in U$, we assume that $Y_{\cdot+t}^u = Y^u \circ \theta_t$ for all $t \in \mathbb{R}_+$. Thanks to Lemmata 5.3 and 5.4, we get that

$$\left\{ (t, \omega, P) : \forall_{u \in U} Y^u \in \mathcal{S}_{\text{ac}}^{\text{SP}}(t, P), (\lambda \otimes P)\text{-a.e. } (dA^P(Y_{\cdot+t}^u)/d\lambda)_{u \in U} \in \Theta(\cdot + t, \omega \otimes_t X) \right\} \\ = \left\{ (t, \omega, P) : \forall_{u \in U} P_t \in \mathfrak{P}_{\text{spsem}}^{\text{ac}}(Y^u), (\lambda \otimes P_t)\text{-a.e. } (a^{P_t}(Y^u))_{u \in U} \in \Theta(\cdot + t, \omega \tilde{\otimes}_t X) \right\}.$$

The latter set is Borel by Lemmata 5.1 and 5.2, and [5, Theorem 8.10.61]. \square

Corollary 5.6. *The measurable graph condition holds.*

5.2. Stability under Conditioning. Next, we check stability under conditioning. In this section, we fix $t^* \in \mathbb{R}_+$, a stopping time τ with $t^* \leq \tau < \infty$, and a probability measure P on (Ω, \mathcal{F}) . We denote by $P(\cdot | \mathcal{F}_\tau)$ a version of the regular conditional P -probability given \mathcal{F}_τ . We recall some simple observations.

Lemma 5.7.

- (i) *There exists a P -null set $N \in \mathcal{F}_\tau$ such that $P(A | \mathcal{F}_\tau)(\omega) = \mathbf{1}_A(\omega)$ for all $A \in \mathcal{F}_\tau$ and $\omega \notin N$.*
- (ii) *If $\omega \mapsto Q_\omega$ is a kernel from \mathcal{F} into \mathcal{F} such that $Q_\omega(\omega = X \text{ on } [0, \tau(\omega)]) = 1$ for P -a.a. $\omega \in \Omega$, then $Q_\omega(\omega \otimes_{\tau(\omega)} X = X) = 1$ for P -a.a. $\omega \in \Omega$. In particular, there exists a P -null set $N \in \mathcal{F}_\tau$ such that $P(\omega \otimes_{\tau(\omega)} X = X | \mathcal{F}_\tau)(\omega) = 1$ for all $\omega \notin N$.*
- (iii) *Take a predictable process $H = (H_s)_{s \geq 0}$, $t \in \mathbb{R}_+$ and $\omega \in \Omega$. Then, $H_{\cdot+t}(\omega \otimes_t X)$ is predictable for the (right-continuous) filtration generated by $X_{\cdot+t}$.*

Proof. Part (i) is classical ([50, Theorem 1.1.8]), part (ii) is obvious and part (iii) follows from [24, Proposition 10.35]. \square

The next lemma is a partial restatement of [50, Theorem 1.2.10].

Lemma 5.8. *If $M - M^{t^*}$ is a P -martingale, then there exists a P -null set N such that, for all $\omega \notin N$, $M - M^{\tau(\omega)}$ is a $P(\cdot|\mathcal{F}_\tau)(\omega)$ -martingale.*

Corollary 5.9. *If $M - M^{t^*}$ is a local P -martingale, then there exists a P -null set N such that, for all $\omega \notin N$, $M - M^{\tau(\omega)}$ is a local $P(\cdot|\mathcal{F}_\tau)(\omega)$ -martingale.*

Proof. Let $(\tau_n)_{n=1}^\infty$ be a localizing sequence for $M - M^{t^*}$. By Lemma 5.8, there exists a P -null set N such that, for all $\omega \notin N$, $M^{\tau_n} - M^{\tau_n \wedge \tau(\omega)}$ is a $P(\cdot|\mathcal{F}_\tau)(\omega)$ -martingale for every $n \in \mathbb{N}$. Since

$$E^P \left[P \left(\liminf_{n \rightarrow \infty} \tau_n < \infty \mid \mathcal{F}_\tau \right) \right] = P \left(\liminf_{n \rightarrow \infty} \tau_n < \infty \right) = 0,$$

we can enlarge the P -null set N such that, for all $\omega \notin N$, $P(\tau_n \rightarrow \infty | \mathcal{F}_\tau)(\omega) = 1$. The claim is proved. \square

Lemma 5.10. *Suppose that $Y \in \mathcal{S}^{\text{sp}}(t^*, P)$. Then, there exists a P -null set N such that, for every $\omega \notin N$, $Y \in \mathcal{S}^{\text{sp}}(\tau(\omega), P(\cdot|\mathcal{F}_\tau)(\omega))$ and $P(\cdot|\mathcal{F}_\tau)(\omega)$ -a.s.*

$$A^{P(\cdot|\mathcal{F}_\tau)(\omega)}(Y_{\cdot+\tau(\omega)}) = (A_{\cdot+\tau(\omega)-t^*}^P(Y_{\cdot+t^*}) - A_{\tau(\omega)-t^*}^P(Y_{\cdot+t^*}))(\omega \otimes_{\tau(\omega)} X).$$

Moreover, when ν^P denotes the third P -characteristic of $Y_{\cdot+t^*}$, then, for every $\omega \notin N$, the third $P(\cdot|\mathcal{F}_\tau)(\omega)$ -characteristic of $Y_{\cdot+\tau(\omega)}$ (for the right-continuous natural filtration of $X_{\cdot+\tau(\omega)}$) is given by

$$\nu^P(\omega \otimes_{\tau(\omega)} X; \tau(\omega) - t^* + dt, dx).$$

Proof. Thanks to Corollary 5.9, there exists a P -null set N such that, for all $\omega \notin N$, the process

$$Y_{\cdot+t^*} - Y_{\cdot \wedge (\tau(\omega) - t^*) + t^*} - (A^P(Y_{\cdot+t^*}) - A_{\cdot \wedge (\tau(\omega) - t^*)}^P(Y_{\cdot+t^*}))$$

is a local $P(\cdot|\mathcal{F}_\tau)(\omega)$ -martingale. Inserting $s \mapsto s + \tau(\omega) - t^*$ yields that the process

$$Y_{\cdot+\tau(\omega)} - Y_{\tau(\omega)} - (A_{\cdot+\tau(\omega)-t^*}^P(Y_{\cdot+t^*}) - A_{\tau(\omega)-t^*}^P(Y_{\cdot+t^*}))$$

is a local $P(\cdot|\mathcal{F}_\tau)(\omega)$ -martingale, too. Thanks to part (ii) of Lemma 5.7, we can enlarge the P -null set N such that, for all $\omega \notin N$, $P(\cdot|\mathcal{F}_\tau)(\omega)$ -a.s.

$$A_{\cdot+\tau(\omega)-t^*}^P(Y_{\cdot+t^*}) - A_{\tau(\omega)-t^*}^P(Y_{\cdot+t^*}) = (A_{\cdot+\tau(\omega)-t^*}^P(Y_{\cdot+t^*}) - A_{\tau(\omega)-t^*}^P(Y_{\cdot+t^*}))(\omega \otimes_{\tau(\omega)} X).$$

By part (iii) of Lemma 5.7, the process on the r.h.s. is predictable for the right-continuous natural filtration of $X_{\cdot+\tau(\omega)}$. Hence, for every $\omega \notin N$, we have $Y \in \mathcal{S}^{\text{sp}}(\tau(\omega), P(\cdot|\mathcal{F}_\tau)(\omega))$ and also the formula for $A^{P(\cdot|\mathcal{F}_\tau)(\omega)}(Y_{\cdot+\tau(\omega)})$ follows.

To prove the final claim, let $g: \mathbb{R} \rightarrow \mathbb{R}$ be a bounded Borel function which vanishes around the origin. Then, by virtue of Corollary 5.9, possibly making N a bit larger, for every $\omega \notin N$, the process

$$\sum_{s \leq \cdot} g(\Delta(Y_{\cdot+\tau(\omega)})_s) - \int_0^\cdot \int g(x) \nu^P(\omega \otimes_{\tau(\omega)} X; \tau(\omega) - t^* + dt, dx)$$

is a local $P(\cdot|\mathcal{F}_\tau)(\omega)$ -martingale. The third characteristic can be characterized by this local martingale property for countably many test functions g , see [26, Theorem II.2.21]. Thus, the formula for the third $P(\cdot|\mathcal{F}_\tau)(\omega)$ -characteristic follows. The proof is complete. \square

We are in the position to deduce stability under conditioning.

Lemma 5.11. *If $P \in \mathcal{C}(t^*, \omega^*)$, then P -a.s. $P(\cdot|\mathcal{F}_\tau) \in \mathcal{C}(\tau, X)$.*

Proof. Let N be as in part (i) of Lemma 5.7 and take $\omega \notin N$. Then, $P(\tau = \tau(\omega)|\mathcal{F}_\tau)(\omega) = 1$ and $P(X = \omega \text{ on } [0, \tau(\omega)]|\mathcal{F}_\tau)(\omega) = 1$. Thanks to Lemma 5.10, as the set U is assumed to be countable, for P -a.a. $\omega \in \Omega$ and all $u \in U$, we have $Y^u \in \mathcal{S}_{\text{ac}}^{\text{sp}}(\tau(\omega), P(\cdot|\mathcal{F}_\tau)(\omega))$ and $P(\cdot|\mathcal{F}_\tau)(\omega)$ -a.s.

$$d\mathbf{A}^{P(\cdot|\mathcal{F}_\tau)(\omega)}(Y_{+\tau(\omega)}^u)/d\lambda = (d\mathbf{A}_{+\tau(\omega)-t^*}^P(Y_{+t^*}^u)/d\lambda)(\omega \otimes_{\tau(\omega)} X).$$

We get from part (ii) of Lemma 5.7, Fubini's theorem, part (i) of Lemma 5.7, the tower rule, and $P \in \mathcal{C}(t^*, \omega^*)$ that

$$\begin{aligned} & \iint_0^\infty P((d\mathbf{A}^{P(\cdot|\mathcal{F}_\tau)(\omega)}(Y_{+\tau(\omega)}^u)/d\lambda)_{u \in U}(t, X) \notin \Theta(t + \tau(\omega), X)|\mathcal{F}_\tau)(\omega) dt P(d\omega) \\ &= \iint_0^\infty P((d\mathbf{A}^P(Y_{+t^*}^u)/d\lambda)_{u \in U}(t + \tau(\omega) - t^*, X) \notin \Theta(t + \tau(\omega), X)|\mathcal{F}_\tau)(\omega) dt P(d\omega) \\ &= \int E^P \left[\int_{\tau(\omega)-t^*}^\infty \mathbf{1}\{(d\mathbf{A}^P(Y_{+t^*}^u)/d\lambda)_{u \in U}(t, X) \notin \Theta(t + t^*, X)\} dt | \mathcal{F}_\tau \right](\omega) P(d\omega) \\ &= \int E^P \left[\int_{\tau-t^*}^\infty \mathbf{1}\{(d\mathbf{A}^P(Y_{+t^*}^u)/d\lambda)_{u \in U}(t, X) \notin \Theta(t + t^*, X)\} dt | \mathcal{F}_\tau \right](\omega) P(d\omega) \\ &= E^P \left[\int_{\tau-t^*}^\infty \mathbf{1}\{(d\mathbf{A}^P(Y_{+t^*}^u)/d\lambda)_{u \in U}(t, X) \notin \Theta(t + t^*, X)\} dt \right] = 0. \end{aligned}$$

This observation completes the proof of P -a.s. $P(\cdot|\mathcal{F}_\tau) \in \mathcal{C}(\tau, X)$. \square

5.3. Stability under Pasting. In this section we check stability under pasting. The proof is split into several steps. Throughout this section, we fix a probability measure P on (Ω, \mathcal{F}) , $t^* \in \mathbb{R}_+$, a stopping time τ with $t^* \leq \tau < \infty$, and an \mathcal{F}_τ -measurable map $\Omega \ni \omega \mapsto Q_\omega \in \mathfrak{P}(\Omega)$ such that $Q_\omega(X = \omega \text{ on } [0, \tau(\omega)]) = 1$ for P -a.a. $\omega \in \Omega$. To simplify our notation, we set $\bar{P} \triangleq P \otimes_\tau Q$. The following corollary is an immediate consequence of part (ii) of Lemma 5.7.

Corollary 5.12. $\bar{P} = E^P [Q(\cdot)]$.

The next lemma follows identically to [8, Lemma 3.14]. We omit the details.

Lemma 5.13. $P = \bar{P}$ on \mathcal{F}_τ and \bar{P} -a.s. $\bar{P}(\cdot|\mathcal{F}_\tau) = Q$.

Lemma 5.14. Suppose that $Y \in \mathcal{S}_{\text{ac}}^{\text{sp}}(t^*, P)$. If $Y \in \mathcal{S}_{\text{ac}}^{\text{sp}}(\tau(\omega), Q_\omega)$ for P -a.a. $\omega \in \Omega$, then $Y \in \mathcal{S}_{\text{ac}}^{\text{sp}}(t^*, \bar{P})$.

Proof. Step 1: Semimartingale Property. Let $T > t^*$, and take a sequence $(H^n)_{n=0}^\infty$ of simple predictable processes on $[[t^*, T]]$ such that $H^n \rightarrow H^0$ uniformly in time and ω . Then, by Lemma 5.13, we get

$$\begin{aligned} & E^{\bar{P}} \left[\left| \int_{t^*}^T (H_s^n - H_s) dY_s \right| \wedge 1 \right] \\ & \leq E^P \left[\left| \int_{t^*}^{\tau \wedge T} (H_s^n - H_s) dY_s \right| \wedge 1 \right] \\ & \quad + \int E^{Q_\omega} \left[\left| \int_{\tau(\omega) \wedge T}^T (H_s^n - H_s)(\omega \otimes_{\tau(\omega)} X) dY_s \right| \wedge 1 \right] P(d\omega). \end{aligned}$$

The first term converges to zero as $Y \in \mathcal{S}^{\text{sp}}(t^*, P)$ and thanks to the Bichteler–Dellacherie (BD) Theorem ([46, Theorem III.43]). The second term converges to zero by dominated

convergence, the assumption that P -a.s. $Y \in \mathcal{S}^{\text{sp}}(\tau, Q)$ and, by virtue of part (iii) of Lemma 5.7, again the BD Theorem. Consequently, invoking the BD Theorem a third time, but this time the converse direction, yields that $Y \in \mathcal{S}(t^*, \bar{P})$.

Step 2: Special Semimartingale Property. Denote by $\nu^{\bar{P}}$ (resp. ν^P) the third characteristic of $Y_{\cdot+t^*}$ under \bar{P} (resp. under P). By Lemma 5.13, we obtain

$$(5.2) \quad \bar{P}(|x|^2 \wedge |x|) * \nu_{\tau-t^*}^{\bar{P}} < \infty = P(|x|^2 \wedge |x|) * \nu_{\tau-t^*}^P < \infty = 1.$$

Furthermore, as $Y \in \mathcal{S}(t^*, \bar{P})$ and \bar{P} -a.s. $\bar{P}(\cdot | \mathcal{F}_\tau) = Q$ by Lemma 5.13, we deduce from Lemma 5.10 that, for \bar{P} -a.a. $\omega \in \Omega$, the third Q_ω -characteristic of $Y_{\cdot+\tau(\omega)}$ is given by

$$\nu^{\bar{P}}(\omega \otimes_{\tau(\omega)} X; \tau(\omega) - t^* + dt, dx).$$

As P -a.s. $Y \in \mathcal{S}^{\text{sp}}(\tau, Q)$, we have, for \bar{P} -a.a. $\omega \in \Omega$, that, for all $T > \tau(\omega) - t^*$,

$$Q_\omega \left(\int_{\tau(\omega)-t^*}^T \int (|x|^2 \wedge |x|) \nu^{\bar{P}}(\omega \otimes_{\tau(\omega)} X; dt, dx) < \infty \right) = 1.$$

By Lemmata 5.11, 5.7 [(i), (ii)] and 5.13, this yields, for all $T > 0$, that

$$(5.3) \quad \begin{aligned} & \bar{P} \left(\int_{\tau-t^*}^T \int (|x|^2 \wedge |x|) \nu^{\bar{P}}(dt, dx) < \infty, T > \tau - t^* \right) \\ &= \int_{\{T > \tau - t^*\}} \bar{P} \left(\int_{\tau(\omega)-t^*}^T \int (|x|^2 \wedge |x|) \nu^{\bar{P}}(\omega \otimes_{\tau(\omega)} X; dt, dx) < \infty | \mathcal{F}_\tau \right) (\omega) \bar{P}(d\omega) \\ &= \int_{\{T > \tau - t^*\}} Q_\omega \left(\int_{\tau(\omega)-t^*}^T \int (|x|^2 \wedge |x|) \nu^{\bar{P}}(\omega \otimes_{\tau(\omega)} X; dt, dx) < \infty \right) \bar{P}(d\omega) \\ &= \bar{P}(T > \tau - t^*). \end{aligned}$$

Finally, (5.2) and (5.3) yield that

$$\bar{P}(|x|^2 \wedge |x|) * \nu_T^{\bar{P}} < \infty = 1, \quad T > 0.$$

By virtue of [26, Proposition II.2.29], this implies that $Y \in \mathcal{S}^{\text{sp}}(t^*, \bar{P})$.

Step 3: Absolutely Continuous Compensator. Let

$$\mathbf{A}^{\bar{P}}(Y_{\cdot+t^*}) = \int_0^\cdot \phi_s ds + \psi$$

be the Lebesgue decomposition of (the paths) of $\mathbf{A}^{\bar{P}}(Y_{\cdot+t^*})$. Since $Y \in \mathcal{S}_{\text{ac}}^{\text{sp}}(t^*, P)$ and $\bar{P} = P$ on \mathcal{F}_τ by Lemma 5.13, we get that $\mathbf{A}^{\bar{P}}(Y_{\cdot+t^*}) \ll \lambda$ on $\llbracket 0, \tau - t^* \rrbracket$. Hence, by Corollary 5.12, it suffices to show that

$$D \triangleq \left\{ \mathbf{A}_{\cdot+\tau-t^*}^{\bar{P}}(Y_{\cdot+t^*}) - \mathbf{A}_{\tau-t^*}^{\bar{P}}(Y_{\cdot+t^*}) \neq \int_{\tau-t^*}^{\cdot+\tau-t^*} \phi_s ds \right\}$$

is a Q_ω -null set for \bar{P} -a.a. $\omega \in \Omega$. Due to Lemmata 5.7 (ii), 5.10 and 5.13, for \bar{P} -a.a. $\omega \in \Omega$, we have Q_ω -a.s.

$$\mathbf{A}_{\cdot+\tau(\omega)-t^*}^{\bar{P}}(Y_{\cdot+t^*}) - \mathbf{A}_{\tau(\omega)-t^*}^{\bar{P}}(Y_{\cdot+t^*}) = \mathbf{A}^{Q_\omega}(Y_{\cdot+\tau(\omega)}).$$

Since P -a.s. $Y \in \mathcal{S}_{\text{ac}}^{\text{sp}}(\tau, Q)$, the uniqueness of the Lebesgue decomposition yields that, for \bar{P} -a.a. $\omega \in \Omega$,

$$\begin{aligned} Q_\omega \left(\mathbf{A}_{\cdot, +\tau(\omega)-t^*}^{\bar{P}}(Y_{\cdot, +t^*}) - \mathbf{A}_{\tau(\omega)-t^*}^{\bar{P}}(Y_{\cdot, +t^*}) \neq \int_{\tau(\omega)-t^*}^{\cdot + \tau(\omega) - t^*} \phi_s ds \right) \\ = Q_\omega \left(\mathbf{A}^{Q_\omega}(Y_{\cdot, +\tau(\omega)}) \neq \int_{\tau(\omega)-t^*}^{\cdot + \tau(\omega) - t^*} \phi_s ds \right) = 0. \end{aligned}$$

Since $Q_\omega(\tau = \tau(\omega)) = 1$ for \bar{P} -a.a. $\omega \in \Omega$ by Lemma 5.13, we have \bar{P} -a.s. $Q(D) = 0$. This shows that $\mathbf{A}^{\bar{P}}(Y_{\cdot, +t^*}) \ll \lambda$ and hence, completes the proof of $Y \in \mathcal{S}_{\text{ac}}^{\text{sp}}(t^*, \bar{P})$. \square

Lemma 5.15. *Let $P \in \mathcal{C}(t^*, \omega^*)$. If $Q_\omega \in \mathcal{C}(\tau(\omega), \omega)$ for P -a.e. $\omega \in \Omega$, then $\bar{P} \in \mathcal{C}(t^*, \omega^*)$.*

Proof. Lemma 5.14 implies that $Y^u \in \mathcal{S}_{\text{ac}}^{\text{sp}}(t^*, \bar{P})$ for all $u \in U$. Since $\{X = \omega^* \text{ on } [0, t^*]\} \in \mathcal{F}_\tau$, as $\tau \geq t^*$, Lemma 5.13 yields $\bar{P}(X = \omega^* \text{ on } [0, t^*]) = P(X = \omega^* \text{ on } [0, t^*]) = 1$. Consequently, as $\bar{P} = P$ on \mathcal{F}_τ , it suffices to show that

$$R \triangleq \{(t, \omega) \in \llbracket \tau - t^*, \infty \llbracket : (d\mathbf{A}^{\bar{P}}(Y_{\cdot, +t^*}^u)/d\lambda)_{u \in U}(t, \omega) \notin \Theta(t + t^*, \omega)\}$$

is a $(\lambda \otimes \bar{P})$ -null set. By virtue of Lemmata 5.10 and 5.13, the assumption that P -a.s. $Q \in \mathcal{C}(\tau, X)$ yields, for P -a.a. $\omega \in \Omega$, that

$$\begin{aligned} (\lambda \otimes Q_\omega)((t, \omega') \in \llbracket \tau(\omega) - t^*, \infty \llbracket : (d\mathbf{A}^{\bar{P}}(Y_{\cdot, +t^*}^u)/d\lambda)_{u \in U}(t, \omega') \notin \Theta(t + t^*, \omega')) \\ = (\lambda \otimes Q_\omega)((t, \omega') \in \llbracket 0, \infty \llbracket : (d\mathbf{A}_{+\tau(\omega)-t^*}^{\bar{P}}(Y_{\cdot, +t^*}^u)/d\lambda)_{u \in U}(t, \omega') \notin \Theta(t + \tau(\omega), \omega')) \\ = (\lambda \otimes Q_\omega)((t, \omega') \in \llbracket 0, \infty \llbracket : (d\mathbf{A}^{Q_\omega}(Y_{\cdot, +\tau(\omega)}^u)/d\lambda)_{u \in U}(t, \omega') \notin \Theta(t + \tau(\omega), \omega')) = 0. \end{aligned}$$

Finally, as $Q_\omega(\tau = \tau(\omega)) = 1$ for P -a.a. $\omega \in \Omega$, Corollary 5.12 and Fubini's theorem yield that

$$\begin{aligned} (\lambda \otimes \bar{P})(R) &= E^{\bar{P}} \left[\int_{\tau-t^*}^{\infty} \mathbb{1}\{(d\mathbf{A}^{\bar{P}}(Y_{\cdot, +t^*}^u)/d\lambda)_{u \in U}(t, X) \notin \Theta(t + t^*, X)\} dt \right] \\ &= E^P \left[E^Q \left[\int_{\tau-t^*}^{\infty} \mathbb{1}\{(d\mathbf{A}^{\bar{P}}(Y_{\cdot, +t^*}^u)/d\lambda)_{u \in U}(t, X) \notin \Theta(t + t^*, X)\} dt \right] \right] \\ &= E^P \left[\int_{\tau-t^*}^{\infty} Q((d\mathbf{A}^{\bar{P}}(Y_{\cdot, +t^*}^u)/d\lambda)_{u \in U}(t, X) \notin \Theta(t + t^*, X)) dt \right] = 0. \end{aligned}$$

This completes the proof. \square

5.4. Proof of Theorem 2.5. Corollary 5.6 and Lemmata 5.11 and 5.15 yield that the prerequisites of [16, Theorem 2.1] are fulfilled and this theorem implies the DPP. \square

6. PROOF OF THE STRONG MARKOV SELECTION PRINCIPLE: THEOREM 3.9

The proof of the strong Markov selection principle is based on some fundamental ideas of Krylov [28] for Markovian selection as worked out in the monograph [50] of Stroock and Varadhan. The main technical steps in the argument are to establish stability under conditioning and pasting of a suitable sequence of correspondences. Hereby, we adapt the proof of [9, Theorem 2.19] to our more general and more abstract framework.

This section is split into two parts. In the first, we study some properties of the correspondence \mathcal{K} and thereafter, we finalize the proof.

6.1. Preparations. Let us stress that in this section we do not assume Condition 3.5. In each of the following results, we indicate precisely which prerequisites are used (except for our Standing Assumptions 2.1, 2.2 and 3.1, which we assume).

For $t \in \mathbb{R}_+$, we define $\gamma_t: \Omega \rightarrow \Omega$ by $\gamma_t(\omega) \triangleq \omega((\cdot - t)^+)$ for $\omega \in \Omega$. Moreover, for $P \in \mathfrak{P}(\Omega)$ and $t \in \mathbb{R}_+$, we set

$$P^t \triangleq P \circ \gamma_t^{-1}.$$

Recall also the notation $P_t = P \circ \theta_t^{-1}$. In case the path space Ω is the space of continuous functions from \mathbb{R}_+ into F , we have the following extension of Lemma 5.2.

Lemma 6.1. *Suppose that $\Omega = C(\mathbb{R}_+; F)$. The maps $(t, P) \mapsto P_t$ and $(t, P) \mapsto P^t$ are continuous.*

Proof. Notice that $(t, \omega) \mapsto \theta_t(\omega)$ and $(t, \omega) \mapsto \gamma_t(\omega)$ are continuous by the Arzelà–Ascoli theorem. Now, the claim follows from [5, Theorem 8.10.61]. \square

Lemma 6.2. *For every $(t, \omega) \in \llbracket 0, \infty \llbracket$, $P \in \mathcal{C}(t, \omega)$ implies $P_t \in \mathcal{K}(0, \omega(t))$.*

Proof. Let $(t, \omega) \in \llbracket 0, \infty \llbracket$, take $P \in \mathcal{C}(t, \omega)$ and fix $u \in U$. It is clear that $P_t \circ X_0^{-1} = \delta_{\omega(t)}$. Since, by hypothesis, $Y_{\cdot+t}^u = Y^u \circ \theta_t$, we deduce from Lemma 5.4 and [26, Proposition II.2.29] that $Y^u \in \mathcal{S}_{\text{ac}}^{\text{sp}}(P_t)$ and P -a.s. $A^P(Y_{\cdot+t}^u) = A^{P_t}(Y^u) \circ \theta_t$. Hence, using also Standing Assumption 3.1, we deduce that

$$\begin{aligned} (\lambda \otimes P_t)((A^{P_t}(Y^u))_{u \in U} \notin \Theta) &= (\lambda \otimes P)((A^{P_t}(Y^u) \circ \theta_t)_{u \in U} \notin \Theta(\cdot, X \circ \theta_t)) \\ &= (\lambda \otimes P)((A^P(Y_{\cdot+t}^u))_{u \in U} \notin \Theta(\cdot + t, X)) = 0. \end{aligned}$$

We conclude that $P_t \in \mathcal{K}(0, \omega(t))$. \square

Lemma 6.3. *For every $(t, x) \in \mathbb{R}_+ \times F$, we have $P \in \mathcal{K}(0, x)$ if and only if $P^t \in \mathcal{K}(t, x)$.*

Proof. Let $x \in F$ and $P \in \mathcal{K}(0, x)$. As $\gamma_t^{-1}(\{X = x \text{ on } [0, t]\}) = \{X_0 = x\}$, we have $P^t(X = x \text{ on } [0, t]) = 1$. Take $u \in U$ and notice that $Y^u = Y^u \circ \theta_t \circ \gamma_t = Y_{\cdot+t}^u \circ \gamma_t$, as $\theta_t \circ \gamma_t = \text{id}$. Hence, by Lemma 5.4 and [26, Proposition II.2.29], $Y^u \in \mathcal{S}_{\text{ac}}^{\text{sp}}(t, P^t)$ and P -a.s. $A^P(Y^u) = A^{P^t}(Y_{\cdot+t}^u) \circ \gamma_t$. From the last equality, and Standing Assumption 3.1, we deduce that

$$\begin{aligned} (\lambda \otimes P^t)((A^{P^t}(Y_{\cdot+t}^u))_{u \in U} \notin \Theta(\cdot + t, X)) &= (\lambda \otimes P)((A^{P^t}(Y_{\cdot+t}^u))_{u \in U} \circ \gamma_t \notin \Theta((X \circ \gamma_t)(\cdot + t))) \\ &= (\lambda \otimes P)((A^P(Y^u))_{u \in U} \notin \Theta) \\ &= 0. \end{aligned}$$

This proves that $P^t \in \mathcal{K}(t, x)$. Conversely, take $P^t \in \mathcal{K}(t, x)$. Due to the identity $\theta_t \circ \gamma_t = \text{id}$, we have $P = (P^t)_t$ and Lemma 6.2 implies that $P \in \mathcal{K}(0, x)$. The proof is complete. \square

Lemma 6.4. *For every $(t, x) \in \mathbb{R}_+ \times F$, we have $\mathcal{K}(t, x) = \{P^t: P \in \mathcal{K}(0, x)\}$.*

Proof. Take $(t, x) \in \mathbb{R}_+ \times F$. Lemma 6.3 yields the inclusion $\{P^t: P \in \mathcal{K}(0, x)\} \subset \mathcal{K}(t, x)$. Conversely, take $P \in \mathcal{K}(t, x)$. As $P(X = x \text{ on } [0, t]) = 1$, we have $P = (P_t)^t$. Now, since $P_t \in \mathcal{K}(0, x)$ by Lemma 6.2, we get $\mathcal{K}(t, x) \subset \{P^t: P \in \mathcal{K}(0, x)\}$. \square

Lemma 6.5. *Suppose that $\Omega = C(\mathbb{R}_+; F)$ and that $\Omega \ni \omega \mapsto L(\omega) \in C(\mathbb{R}_+; \mathbb{R}_+)$ is continuous. Let P be a Borel probability measure on $\Omega \times C(\mathbb{R}_+; \mathbb{R})$ and set*

$$\zeta_M(\omega, \alpha) \triangleq \sup \left(\frac{|\alpha(t \wedge \rho_M(\omega)) - \alpha(s \wedge \rho_M(\omega))|}{t - s} : 0 \leq s < t \right),$$

for $(\omega, \alpha) \in \Omega \times C(\mathbb{R}_+; \mathbb{R})$, where

$$\rho_M = \inf(t \geq 0: L_t \geq M), \quad M > 0.$$

Then, there exists a dense set $D \subset \mathbb{R}_+$ such that, for every $M \in D$, there exists a P -null set $N = N(M)$ such that ζ_M is lower semicontinuous at every $\omega \notin N$.

Proof. In the first part of this proof, we adapt an argument from [50, Lemma 11.1.2]. Notice that $\omega \mapsto \rho_M(\omega)$ is lower semicontinuous. To see this, recall that

$$\{\rho_M \leq t\} = \left\{ \inf_{s \in \mathbb{Q} \cap [0, t]} \inf_{z \geq M} |L_s - z| = 0 \right\}, \quad t \in \mathbb{R}_+,$$

and notice that the set on the right hand side is closed, as $\omega \mapsto \inf_{s \in \mathbb{Q} \cap [0, t]} \inf_{z \geq M} |L_s(\omega) - z|$ is continuous. Next, set

$$\rho_M^+ \triangleq \inf(t \geq 0: L_t > M).$$

Since, for every $t > 0$,

$$\{\rho_M^+ < t\} = \bigcup_{\substack{s < t \\ s \in \mathbb{Q}_+}} \{L_s > M\},$$

where the set on the right hand side is open by the continuity of $\omega \mapsto L(\omega)$, we conclude that ρ_M^+ is upper semicontinuous. Notice that $\rho_M^+ = \rho_{M+} \triangleq \lim_{K \searrow M} \rho_K$. Let $\Psi: \Omega \times C(\mathbb{R}_+; \mathbb{R}) \rightarrow \Omega$ be defined by $\Psi(\omega, \alpha) = \omega$, let $\phi: \mathbb{R}_+ \rightarrow [0, 1]$ be defined by $\phi(M) \triangleq E^P[e^{-\rho_M \circ \Psi}]$, and set

$$D \triangleq \left\{ M \in \mathbb{R}_+ : E^P[e^{-\rho_M \circ \Psi}] = E^P[e^{-\rho_{M+} \circ \Psi}] \right\}.$$

The dominated convergence theorem yields that

$$D = \left\{ M \in \mathbb{R}_+ : \phi(M) = \phi(M+) \right\}, \quad \phi(M+) \triangleq \lim_{K \searrow M} \phi(K).$$

The set $\mathbb{R}_+ \setminus D$ is countable, as monotone functions, such as ϕ , have at most countably many discontinuities. Since $\rho_M \leq \rho_M^+$, for every $M \in D$ we have P -a.s. $\rho_M \circ \Psi = \rho_M^+ \circ \Psi$. To see this, notice that

$$E^P[(e^{-\rho_M^+ \circ \Psi} - e^{-\rho_M \circ \Psi}) \mathbb{1}_{\{\rho_M \circ \Psi < \rho_M^+ \circ \Psi\}}] = E^P[e^{-\rho_M^+ \circ \Psi} - e^{-\rho_M \circ \Psi}] = 0,$$

which implies that $P(\rho_M \circ \Psi < \rho_M^+ \circ \Psi) = 0$.

Now, take $\omega \in \{\rho_M = \rho_M^+\}$. Since $\rho_M \leq \rho_M^+$, for every sequence $(\omega^n)_{n=1}^\infty \subset \Omega$ with $\omega^n \rightarrow \omega$, we obtain

$$\rho_M(\omega) \leq \liminf_{n \rightarrow \infty} \rho_M(\omega^n) \leq \limsup_{n \rightarrow \infty} \rho_M(\omega^n) \leq \limsup_{n \rightarrow \infty} \rho_M^+(\omega^n) \leq \rho_M^+(\omega) = \rho_M(\omega),$$

which implies that ρ_M is continuous at ω .

Finally, let $\Omega \times C(\mathbb{R}_+; \mathbb{R}) \ni (\omega^n, \alpha^n) \rightarrow (\omega, \alpha) \in \{\rho_M \circ \Psi = \rho_M^+ \circ \Psi\}$. Then,

$$\begin{aligned} \zeta_M(\omega, \alpha) &= \sup \left(\liminf_{n \rightarrow \infty} \frac{|\alpha^n(t \wedge \rho_M(\omega^n)) - \alpha^n(s \wedge \rho_M(\omega^n))|}{t - s} : 0 \leq s < t \right) \\ &\leq \liminf_{n \rightarrow \infty} \zeta_M(\omega^n, \alpha^n). \end{aligned}$$

The proof is complete. \square

Proposition 6.6. *Assume that (i) – (iv) from Condition 3.5 hold. Then, for every closed set $K \subset F$, the set $\mathcal{R}(K) \triangleq \bigcup_{x \in K} \mathcal{R}(x)$ is closed (in $\mathfrak{P}(\Omega)$ endowed with the weak topology).*

Proof. Let $(P^n)_{n=1}^\infty \subset \mathcal{R}(K)$ be such that $P^n \rightarrow P$ weakly. By definition of $\mathcal{R}(K)$, for every $n \in \mathbb{N}$, there exists a point $x^n \in K$ such that $P^n \in \mathcal{R}(x^n)$. Since $P^n \rightarrow P$ and $\{\delta_x: x \in K\}$ is closed ([1, Theorem 15.8]), there exists a point $x^0 \in K$ such that $P \circ X_0^{-1} = \delta_{x^0}$. In particular, $x^n \rightarrow x^0$. We now prove that $P \in \mathcal{R}(x^0)$. Before we start our program, let us fix some auxiliary notation. We set $\Omega^* \triangleq \Omega \times C(\mathbb{R}_+; \mathbb{R})^U$ and endow this space with the product local uniform topology. In this case, the Borel σ -field $\mathcal{B}(\Omega^*) \triangleq \mathcal{F}^*$ coincides with $\sigma(Z_t, t \geq 0)$, where $Z = (Z^{(1)}, (Z^{(2,u)})_{u \in U})$ denotes the coordinate process on Ω^* . Furthermore, we define $\mathbf{F}^* \triangleq (\mathcal{F}_t^*)_{t \geq 0}$ to be the right-continuous filtration generated by the coordinate process Z .

Step 1. In this first step, we show that the family $\{P^n \circ (X, (A^{P^n}(Y^u))_{u \in U})^{-1}: n \in \mathbb{N}\}$ is tight when seen as a family of probability measures on $(\Omega^*, \mathcal{F}^*)$. As Ω^* is endowed with a product topology and since $P^n \rightarrow P$, it suffices to prove that, for every fixed $u \in U$, the family $\{P^n \circ A^{P^n}(Y^u)^{-1}: n \in \mathbb{N}\}$ is tight when seen as a family of Borel probability measures on $C(\mathbb{R}_+; \mathbb{R})$, endowed with the local uniform topology. First, for every $M > 0$, we deduce tightness of $\{P^n \circ A_{s \wedge \rho_M}^{P^n}(Y^u)^{-1}: n \in \mathbb{N}\}$ from Kolmogorov's tightness criterion ([27, Theorem 23.7]). By the first part from Condition 3.5 (iii), there exists a constant $C > 0$ such that, for all $s < t$,

$$\begin{aligned} \sup_{n \in \mathbb{N}} E^{P^n} [|A_{s \wedge \rho_M}^{P^n}(Y^u) - A_{t \wedge \rho_M}^{P^n}(Y^u)|^2] &\leq \sup_{n \in \mathbb{N}} E^{P^n} \left[\left(\int_{s \wedge \rho_M}^{t \wedge \rho_M} \left| \frac{dA^{P^n}(Y^u)}{d\lambda} \right| d\lambda \right)^2 \right] \\ &\leq C(t - s)^2. \end{aligned}$$

We conclude the tightness of $\{P^n \circ A_{s \wedge \rho_M}^{P^n}(Y^u)^{-1}: n \in \mathbb{N}\}$, for every $M > 0$. Using the Arzelà–Ascoli tightness criterion given by [27, Theorem 23.4], we transfer this observation to the global family $\{P^n \circ A^{P^n}(Y^u)^{-1}: n \in \mathbb{N}\}$. For a moment, we fix $\varepsilon, N > 0$. By virtue of Condition 3.5 (iii), there exists an $M_o > N$ such that

$$\sup_{n \in \mathbb{N}} P^n(\rho_{M_o} \leq N) = \sup_{n \in \mathbb{N}} P^n \left(\sup_{s \in [0, N]} L_s \geq M_o \right) \leq \frac{1}{M_o^q} \sup_{n \in \mathbb{N}} E^{P^n} \left[\sup_{s \in [0, N]} |L_s|^q \right] \leq \varepsilon/2,$$

where $q = q(\{x^n: n \in \mathbb{Z}_+\}, N) > 0$ is the exponent from Condition 3.5 (iii). Thanks to the tightness of $\{P^n \circ A_{s \wedge \rho_{M_o}}^{P^n}(Y^u)^{-1}: n \in \mathbb{N}\}$, there exists a compact set $K_N \subset \mathbb{R}$ such that

$$\sup_{n \in \mathbb{N}} P^n(A_{N \wedge \rho_{M_o}}^{P^n}(Y^u) \notin K_N) \leq \varepsilon/2.$$

Consequently, we obtain that

$$\begin{aligned} \sup_{n \in \mathbb{N}} P^n(A_N^{P^n}(Y^u) \notin K_N) &\leq \sup_{n \in \mathbb{N}} P^n(A_{N \wedge \rho_{M_o}}^{P^n}(Y^u) \notin K_N) + \sup_{n \in \mathbb{N}} P^n(\rho_{M_o} \leq N) \\ &\leq \varepsilon/2 + \varepsilon/2 = \varepsilon. \end{aligned}$$

This observation shows that the family $\{P^n \circ A_N^{P^n}(Y^u)^{-1}: n \in \mathbb{N}\}$ is tight. Let d_F be a metric on F which induces its topology. For a set $K \subset \mathbb{R}_+$, $\omega \in \Omega$ and $h > 0$, we define

$$w_K(\omega, h) \triangleq \sup \{ d_F(\omega(s), \omega(t)): s, t \in K, |s - t| \leq h \}.$$

As $\{P^n \circ A_{s \wedge \rho_{M_o}}^{P^n}(Y^u)^{-1}: n \in \mathbb{N}\}$ is tight, [27, Theorem 23.4] yields that

$$\begin{aligned} \sup_{n \in \mathbb{N}} E^{P^n} \left[w_{[0, N]}(A^{P^n}(Y^u), h) \wedge 1 \right] \\ \leq \sup_{n \in \mathbb{N}} E^{P^n} \left[w_{[0, N]}(A_{s \wedge \rho_{M_o}}^{P^n}(Y^u), h) \wedge 1 \right] + \sup_{n \in \mathbb{N}} P^n(\rho_{M_o} \leq N) \end{aligned}$$

$$\leq \sup_{n \in \mathbb{N}} E^{P^n} \left[w_{[0, N]}(\mathbf{A}_{\cdot \wedge \rho_{M_0}}^{P^n}(Y^u), h) \wedge 1 \right] + \varepsilon/2 \rightarrow \varepsilon/2$$

as $h \rightarrow 0$. Using [27, Theorem 23.4] once again, but this time the converse direction, we conclude that $\{P^n \circ \mathbf{A}^{P^n}(Y^u)^{-1} : n \in \mathbb{N}\}$, and hence also $\{P^n \circ (X, (\mathbf{A}^{P^n}(Y^u))_{u \in U})^{-1} : n \in \mathbb{N}\}$, is tight (on the respective probability space). Up to passing to a subsequence, we can assume that $(P^n \circ (X, (\mathbf{A}^{P^n}(Y^u))_{u \in U})^{-1})_{n=1}^\infty$ converges weakly (on the space $(\Omega^*, \mathcal{F}^*)$) to a probability measure Q .

Step 2. Recall that $Z = (Z^{(1)}, (Z^{(2,u)})_{u \in U})$ denotes the coordinate process on Ω^* and fix some $u \in U$. Next, we show that $Z^{(2,u)}$ is Q -a.s. locally Lipschitz continuous. Thanks to Lemma 6.5, there exists a dense set $D \subset \mathbb{R}_+$ such that, for every $M \in D$, the map ζ_M is $Q \circ (Z^{(1)}, Z^{(2,u)})^{-1}$ -a.s. lower semicontinuous. By part (iii) of Condition 3.5, for every $M > 0$, there exists a constant $C = C(M) > 0$ such that $P^n(\zeta_M(X, \mathbf{A}^{P^n}(Y^u)) \leq C) = 1$ for all $n \in \mathbb{N}$. Hence, for every $M \in D$, using the $Q \circ (Z^{(1)}, Z^{(2,u)})^{-1}$ -a.s. lower semicontinuity of ζ_M , we deduce from [45, Example 17, p. 73] that

$$0 = \liminf_{n \rightarrow \infty} P^n(\zeta_M(X, \mathbf{A}^{P^n}(Y^u)) > C) \geq Q(\zeta_M(Z^{(1)}, Z^{(2,u)}) > C).$$

Further, since D is dense in \mathbb{R}_+ , we can conclude that $Z^{(2,u)}$ is Q -a.s. locally Lipschitz continuous and hence, in particular, locally absolutely continuous and of finite variation.

Step 3. Define the map $\Phi: \Omega^* \rightarrow \Omega$ by $\Phi(\omega^{(1)}, \omega^{(2)}) \triangleq \omega^{(1)}$ for $\omega = (\omega^{(1)}, \omega^{(2)}) \in \Omega^*$. In this step, we prove that $(\lambda \otimes Q)$ -a.e. $(dZ^{(2,u)}/d\lambda)_{u \in U} \in \Theta \circ \Phi$. By virtue of [49, Theorems II.4.3 and II.6.2], P^n -a.s. for λ -a.a. $t \in \mathbb{R}_+$, we have

$$(6.1) \quad \begin{aligned} (m(\mathbf{A}_{t+1/m}^{P^n}(Y^u) - \mathbf{A}_t^{P^n}(Y^u)))_{u \in U} &\in \overline{\text{co}}((d\mathbf{A}^{P^n}(Y^u)/d\lambda)_{u \in U}([t, t+1/m])) \\ &\subset \overline{\text{co}} \Theta([t, t+1/m], X). \end{aligned}$$

Here, $\overline{\text{co}}$ denotes the closure of the convex hull. By Skorokhod's coupling theorem ([27, Theorem 5.31]), there exist random variables

$$(X^0, (B^{0,u})_{u \in U}), (X^1, (B^{1,u})_{u \in U}), (X^2, (B^{2,u})_{u \in U}), \dots$$

defined on some probability space $(\Sigma, \mathcal{G}, \mathbb{P})$ such that, for every $n \in \mathbb{N}$, $(X^n, (B^{n,u})_{u \in U})$ has distribution $P^n \circ (X, (\mathbf{A}^{P^n}(Y^u))_{u \in U})^{-1}$, $(X^0, (B^{0,u})_{u \in U})$ has distribution Q , and \mathbb{P} -a.s. $(X^n, (B^{n,u})_{u \in U}) \rightarrow (X^0, (B^{0,u})_{u \in U})$. For every $n \in \mathbb{N}$, we have \mathbb{P} -a.s.

$$(dB^{n,u}/d\lambda)_{u \in U} = \lim_{m \rightarrow \infty} (m(B_{\cdot+1/m}^n - B^n))_{u \in U}.$$

Now, recalling (6.1), we deduce from (ii) and (iv) of Condition 3.5 and [9, Lemma 3.5] that Q -a.s. for λ -a.a. $t \in \mathbb{R}_+$

$$(dB^{0,u}/d\lambda)_{u \in U}(t) \in \bigcap_{m \in \mathbb{N}} \overline{\text{co}} \Theta([t, t+1/m], X^0) \subset \Theta(t, X^0).$$

This proves that $(\lambda \otimes Q)$ -a.e. $(dZ^{(2,u)}/d\lambda)_{u \in U} \in \Theta \circ \Phi$.

Step 4. In the final step of the proof, we show that $Y^u \in \mathcal{S}_{\text{ac}}^{\text{sp}}(P)$ and we relate $B^{0,u}$ to $\mathbf{A}^P(Y^u)$. Notice that $Q \circ \Phi^{-1} = P$. For a moment, we fix $u \in U$. As in the proof of Lemma 6.5, we obtain the existence of a dense set $D = D^u \subset \mathbb{R}_+$ such that $\rho_M \circ \Phi$ is Q -a.s. continuous for all $M \in D$. Take some $M \in D$. Since $Y^u \in \mathcal{S}_{\text{ac}}^{\text{sp}}(P^n)$, the process $Y_{\cdot \wedge \rho_M}^u - \mathbf{A}_{\cdot \wedge \rho_M}^{P^n}(Y^u)$ is a local P^n - \mathbf{F}_+ -martingale. Furthermore, by (i) and (iii) from Condition 3.5, we see that $Y_{\cdot \wedge \rho_M}^u - \mathbf{A}_{\cdot \wedge \rho_M}^{P^n}(Y^u)$ is P^n -a.s. bounded by a constant independent of n , which, in particular, implies that it is a true P^n - \mathbf{F}_+ -martingale. Recall

from part (i) of Condition 3.5, that $\omega \mapsto Y^u(\omega)$ is continuous. Now, it follows³ from [26, Proposition IX.1.4] that

$$Y^u_{\cdot \wedge \rho_M} \circ \Phi - Z^{\cdot \wedge \rho_M \circ \Phi}_{(2,u)}$$

is a Q - \mathbf{F}^* -martingale. Since $Z^{(2,u)}$ is Q -a.s. locally absolutely continuous by Step 2, this means that $Y^u \circ \Phi$ is a Q - \mathbf{F}^* -semimartingale whose first characteristic is given by $Z^{(2,u)}$. Next, we relate this observation to the probability measure P and the filtration \mathbf{F}_+ . For a process A on $(\Omega^*, \mathcal{F}^*)$, we denote by $A^{p, \Phi^{-1}(\mathbf{F}_+)}$ its dual predictable projection to the filtration $\Phi^{-1}(\mathbf{F}_+)$. Recall from [24, Lemma 10.42] that, for every $t \in \mathbb{R}_+$, a random variable V on $(\Omega^*, \mathcal{F}^*)$ is $\Phi^{-1}(\mathcal{F}_{t+})$ -measurable if and only if it is \mathcal{F}_t^* -measurable and $V(\omega^{(1)}, \omega^{(2)})$ does not depend on $\omega^{(2)}$. Thanks to Stricker's theorem ([25, Lemma 2.7]), the process $Y^u \circ \Phi$ is a Q - $\Phi^{-1}(\mathbf{F}_+)$ -semimartingale. Recall from Step 3 that $(\lambda \otimes Q)$ -a.e. $(dZ^{(2,u)}/d\lambda)_{u \in U} \in \Theta \circ \Phi$. By virtue of (iii) from Condition 3.5, for every $M \in D$, we have

$$E^Q[\text{Var}(Z^{(2,u)})_{\rho_M \circ \Phi}] = E^Q\left[\int_0^{\rho_M \circ \Phi} \left| \frac{dZ^{(2,u)}}{d\lambda} \right| d\lambda\right] < \infty,$$

where $\text{Var}(\cdot)$ denotes the total variation process. Hence, we get from [24, Proposition 9.24] that the first Q - $\Phi^{-1}(\mathbf{F}_+)$ -characteristic of $Y^u \circ \Phi$ is given by $(Z^{(2,u)})^{p, \Phi^{-1}(\mathbf{F}_+)}$. Lemma 5.4 yields that Y^u is a P - \mathbf{F}_+ -semimartingale whose first characteristic $\mathbf{A}^P(Y^u)$ satisfies

$$\mathbf{A}^P(Y^u) \circ \Phi = (Z^{(2,u)})^{p, \Phi^{-1}(\mathbf{F}_+)}.$$

Consequently, we deduce from the Steps 2 and 3 that P -a.s. $\mathbf{A}^P(Y^u) \ll \lambda$ and

$$\begin{aligned} (\lambda \otimes P)((\mathbf{A}^P(Y^u))_{u \in U} \notin \Theta) &= (\lambda \otimes Q \circ \Phi^{-1})((d\mathbf{A}^P(Y^u)/d\lambda)_{u \in U} \notin \Theta) \\ &= (\lambda \otimes Q)(E^Q[(dZ^{(2,u)}/d\lambda)_{u \in U} | \Phi^{-1}(\mathbf{F}_+)_-] \notin \Theta \circ \Phi) = 0, \end{aligned}$$

where we use (ii) and (iii) from Condition 3.5 and [49, Theorems II.4.3 and II.6.2] for the final equality. This observation completes the proof. \square

Proposition 6.7. *Suppose that Condition 3.5 holds. The correspondence $(t, x) \mapsto \mathcal{K}(t, x)$ is upper hemicontinuous with non-empty and compact values.*

Proof. The correspondence $x \mapsto \mathcal{K}(0, x)$ has non-empty values by Standing Assumption 2.2. Further, it has compact values by Proposition 6.6 and part (v) of Condition 3.5. Thus, Lemmata 6.1 and 6.4 yield that the same is true for $(t, x) \mapsto \mathcal{K}(t, x)$.

It remains to show that \mathcal{K} is upper hemicontinuous. Let $G \subset \mathfrak{P}(\Omega)$ be closed. We need to show that $\mathcal{K}^l(G) = \{(t, x) \in \mathbb{R}_+ \times F : \mathcal{K}(t, x) \cap G \neq \emptyset\}$ is closed. Suppose that the sequence $(t^n, x^n)_{n=1}^\infty \subset \mathcal{K}^l(G)$ converges to $(t^0, x^0) \in \mathbb{R}_+ \times F$. For each $n \in \mathbb{N}$, there exists a probability measure $P^n \in \mathcal{K}(t^n, x^n) \cap G$. By part (v) of Condition 3.5 and Proposition 6.6, the set $\mathcal{R}^\circ \triangleq \bigcup_{n=0}^\infty \mathcal{R}(x^n)$ is compact. Hence, by Lemma 6.1, so is the set

$$\mathcal{K}^\circ \triangleq \{P^t : (t, P) \in \{t^n : n \in \mathbb{Z}_+\} \times \mathcal{R}^\circ\}.$$

By virtue of Lemma 6.4, we conclude that $\{P^n : n \in \mathbb{N}\} \subset \mathcal{K}^\circ$ is relatively compact. Hence, passing to a subsequence if necessary, we can assume that $P^n \rightarrow P$ weakly for some $P \in \mathcal{K}^\circ \cap G$. Let d_F be a metric on F which induces its topology. For every $\varepsilon \in (0, t^0)$, the

³[26, Proposition IX.1.4] is stated for $F = \mathbb{R}^d$ but the argument needs no change to work for more general state spaces.

set $\{d_F(X_s, x^0) \leq \varepsilon \text{ for all } s \in [0, t^0 - \varepsilon]\} \subset \Omega$ is closed. Consequently, by the Portmanteau theorem, for every $\varepsilon \in (0, t^0)$, we get

$$\begin{aligned} 1 &= \limsup_{n \rightarrow \infty} P^n(d_F(X_s, x^0) \leq \varepsilon \text{ for all } s \in [0, t^0 - \varepsilon]) \\ &\leq P(d_F(X_s, x^0) \leq \varepsilon \text{ for all } s \in [0, t^0 - \varepsilon]). \end{aligned}$$

Consequently, $P(X = x^0 \text{ on } [0, t^0]) = 1$, which implies that $P = (P_{t^0})^{t^0}$. By Lemmata 6.1 and 6.2, we have $(P^n)_{t^n} \in \mathcal{K}(0, x^n)$ and $(P^n)_{t^n} \rightarrow P_{t^0}$ weakly. Further, since $P_{t^0} \circ X_0^{-1} = \lim_{n \rightarrow \infty} \delta_{x^n} = \delta_{x^0}$, Proposition 6.6 yields that $P_{t^0} \in \mathcal{K}(0, x^0)$. Thus, by virtue of Lemma 6.3, $P \in \mathcal{K}^\circ \cap G \cap \mathcal{K}(t^0, x^0) = \mathcal{K}(t^0, x^0) \cap G$, which implies $(t^0, x^0) \in \mathcal{K}^l(G)$. We conclude that \mathcal{K} is upper hemicontinuous. \square

Lemma 6.8. *Suppose that part (ii) of Condition 3.5 holds. The correspondence $(t, x) \mapsto \mathcal{K}(t, x)$ has convex values.*

Proof. By virtue of Lemma 6.4, it suffices to prove that $\mathcal{K}(0, x)$ is convex for every $x \in F$. Indeed, for every $P, Q \in \mathcal{K}(t, x)$ and $\alpha \in (0, 1)$, there are probability measures $\mathbb{P}, \mathbb{Q} \in \mathcal{K}(0, x)$ such that $\mathbb{P}^t = P$ and $\mathbb{Q}^t = Q$. Then, $\alpha P + (1 - \alpha)Q = (\alpha \mathbb{P} + (1 - \alpha)\mathbb{Q})^t$ and consequently, from Lemma 6.3, we get $\alpha P + (1 - \alpha)Q \in \mathcal{K}(t, x)$ once $\alpha \mathbb{P} + (1 - \alpha)\mathbb{Q} \in \mathcal{K}(0, x)$. As Θ is assumed to be convex-valued by Condition 3.5 (ii), the convexity of $\mathcal{K}(0, x)$ follows from [26, Lemma III.3.38, Theorem III.3.40]. To make the argument precise, take $\mathbb{P}, \mathbb{Q} \in \mathcal{K}(0, x)$ and $\alpha \in (0, 1)$, and set $\mathbb{R} \triangleq \alpha \mathbb{P} + (1 - \alpha)\mathbb{Q}$. Clearly, $\mathbb{R} \circ X_0^{-1} = \delta_x$. Moreover, we have $\mathbb{P} \ll \mathbb{R}$ and $\mathbb{Q} \ll \mathbb{R}$ and, by [26, Lemma III.3.38], there are versions $Z^{\mathbb{P}}$ and $Z^{\mathbb{Q}}$ of the Radon–Nikodym densities $d\mathbb{P}/d\mathbb{R}$ and $d\mathbb{Q}/d\mathbb{R}$, respectively, such that

$$\alpha Z^{\mathbb{P}} + (1 - \alpha)Z^{\mathbb{Q}} = 1, \quad 0 \leq Z^{\mathbb{P}} \leq 1/\alpha, \quad 0 \leq Z^{\mathbb{Q}} \leq 1/(1 - \alpha).$$

For every $u \in U$, [26, Proposition II.2.29, Theorem III.3.40] yields that $Y^u \in \mathcal{S}_{\text{ac}}^{\text{SP}}(\mathbb{R})$ and that $(\lambda \otimes \mathbb{R})$ -a.e.

$$d\mathbb{A}^{\mathbb{R}}(Y^u)/d\lambda = \alpha Z^{\mathbb{P}} d\mathbb{A}^{\mathbb{P}}(Y^u)/d\lambda + (1 - \alpha)Z^{\mathbb{Q}} d\mathbb{A}^{\mathbb{Q}}(Y^u)/d\lambda.$$

Notice that

$$\begin{aligned} &\iint Z^{\mathbb{P}} \mathbb{1}\{(d\mathbb{A}^{\mathbb{P}}(Y^u)/d\lambda)_{u \in U} \notin \Theta, Z^{\mathbb{P}} > 0\} d(\lambda \otimes \mathbb{R}) \\ &= (\lambda \otimes \mathbb{P})((d\mathbb{A}^{\mathbb{P}}(Y^u)/d\lambda)_{u \in U} \notin \Theta, Z^{\mathbb{P}} > 0) = 0. \end{aligned}$$

Consequently, $(\lambda \otimes \mathbb{R})$ -a.e. $\mathbb{1}\{(d\mathbb{A}^{\mathbb{P}}(Y^u)/d\lambda)_{u \in U} \notin \Theta, Z^{\mathbb{P}} > 0\} = 0$. In the same manner, we obtain that $(\lambda \otimes \mathbb{R})$ -a.e. $\mathbb{1}\{(d\mathbb{A}^{\mathbb{Q}}(Y^u)/d\lambda)_{u \in U} \notin \Theta, Z^{\mathbb{Q}} > 0\} = 0$. Finally, as $\alpha Z^{\mathbb{P}} + (1 - \alpha)Z^{\mathbb{Q}} = 1$, using the convexity of Θ , we get that

$$\begin{aligned} &(\lambda \otimes \mathbb{R})((d\mathbb{A}^{\mathbb{R}}(Y^u)/d\lambda)_{u \in U} \notin \Theta) \\ &= (\lambda \otimes \mathbb{R})((d\mathbb{A}^{\mathbb{P}}(Y^u)/d\lambda)_{u \in U} \notin \Theta, Z^{\mathbb{P}} > 0, Z^{\mathbb{Q}} = 0) \\ &\quad + (\lambda \otimes \mathbb{R})((d\mathbb{A}^{\mathbb{Q}}(Y^u)/d\lambda)_{u \in U} \notin \Theta, Z^{\mathbb{P}} = 0, Z^{\mathbb{Q}} > 0) \\ &\quad + (\lambda \otimes \mathbb{R})((d\mathbb{A}^{\mathbb{R}}(Y^u)/d\lambda)_{u \in U} \notin \Theta, Z^{\mathbb{P}} > 0, Z^{\mathbb{Q}} > 0) \\ &= (\lambda \otimes \mathbb{R})((\alpha Z^{\mathbb{P}}(d\mathbb{A}^{\mathbb{P}}(Y^u)/d\lambda)_{u \in U} + (1 - \alpha)Z^{\mathbb{Q}}(d\mathbb{A}^{\mathbb{Q}}(Y^u)/d\lambda)_{u \in U}) \notin \Theta, \\ &\quad (d\mathbb{A}^{\mathbb{P}}(Y^u)/d\lambda)_{u \in U} \in \Theta, (d\mathbb{A}^{\mathbb{Q}}(Y^u)/d\lambda)_{u \in U} \in \Theta, Z^{\mathbb{P}} > 0, Z^{\mathbb{Q}} > 0) \\ &= 0. \end{aligned}$$

We conclude that $\mathbb{R} \in \mathcal{K}(0, x)$. The proof is complete. \square

Lemma 6.9. *Let $Q \in \mathfrak{P}(\Omega)$ and take $t \in \mathbb{R}_+$, $\omega, \alpha \in \Omega$ such that $\omega(t) = \alpha(t)$. Then,*

$$\delta_\alpha \otimes_t Q \in \mathcal{C}(t, \alpha) \iff \delta_\omega \otimes_t Q \in \mathcal{C}(t, \omega).$$

Proof. Set $\mathbf{Q} \triangleq \delta_\alpha \otimes_t Q$ and $\mathbf{P} \triangleq \delta_\omega \otimes_t Q$. Suppose that $\mathbf{Q} \in \mathcal{C}(t, \alpha)$. Thanks to Lemma 6.2, we have $\mathbf{Q}_t \in \mathcal{C}(0, \alpha(t)) = \mathcal{C}(0, \omega(t))$. Since $\mathbf{Q}_t = \mathbf{P}_t$, we also have $\mathbf{P}_t \in \mathcal{C}(0, \omega(t))$. We deduce from Lemma 5.4 that $Y^u \in \mathcal{S}_{ac}^{sp}(t, \mathbf{P})$ and that P-a.s. $\mathbf{A}^{\mathbf{P}}(Y^u_{\cdot+t}) = \mathbf{A}^{\mathbf{P}_t}(Y^u) \circ \theta_t$. Hence, using Standing Assumption 3.1, we get that

$$\begin{aligned} (\lambda \otimes \mathbf{P})((\mathbf{A}^{\mathbf{P}}(Y^u_{\cdot+t}))_{u \in U} \notin \Theta(\cdot + t, X)) &= (\lambda \otimes \mathbf{P})((\mathbf{A}^{\mathbf{P}_t}(Y^u) \circ \theta_t)_{u \in U} \notin \Theta(\cdot, X \circ \theta_t)) \\ &= (\lambda \otimes \mathbf{P}_t)((\mathbf{A}^{\mathbf{P}_t}(Y^u))_{u \in U} \notin \Theta) = 0. \end{aligned}$$

We conclude that $\mathbf{P} \in \mathcal{C}(t, \omega)$. The converse implication follows by symmetry. \square

Definition 6.10. *A correspondence $\mathcal{U}: \mathbb{R}_+ \times F \rightarrow \mathfrak{P}(\Omega)$ is said to be*

- (i) *stable under conditioning if for any $(t, x) \in \mathbb{R}_+ \times F$, any stopping time τ with $t \leq \tau < \infty$, and any $P \in \mathcal{U}(t, x)$, there exists a P -null set $N \in \mathcal{F}_\tau$ such that $\delta_{\omega(\tau(\omega))} \otimes_{\tau(\omega)} P(\cdot | \mathcal{F}_\tau)(\omega) \in \mathcal{U}(\tau(\omega), \omega(\tau(\omega)))$ for all $\omega \notin N$;*
- (ii) *stable under pasting if for any $(t, x) \in \mathbb{R}_+ \times F$, any stopping time τ with $t \leq \tau < \infty$, any $P \in \mathcal{U}(t, x)$ and any \mathcal{F}_τ -measurable map $\Omega \ni \omega \mapsto Q_\omega \in \mathfrak{P}(\Omega)$ the following implication holds:*

$$P\text{-a.a. } \omega \in \Omega \quad \delta_{\omega(\tau(\omega))} \otimes_{\tau(\omega)} Q_\omega \in \mathcal{U}(\tau(\omega), \omega(\tau(\omega))) \implies P \otimes_\tau Q \in \mathcal{U}(t, x).$$

Lemma 6.11. *The correspondence \mathcal{K} is stable under conditioning and pasting.*

Proof. Stability under conditioning follows from Lemmata 5.11 and 6.9, and stability under pasting follows from Lemmata 5.15 and 6.9. \square

Recall from [1, Definition 18.1] that a correspondence $\mathcal{U}: \mathbb{R}_+ \times F \rightarrow \mathfrak{P}(\Omega)$ is called *measurable* if the lower inverse $\{(t, x) \in \mathbb{R}_+ \times F: \mathcal{U}(t, x) \cap G \neq \emptyset\}$ is Borel measurable for every closed set $G \subset \mathfrak{P}(\Omega)$. The proof of the following result is similar to those of [9, Lemma 5.12]. We added it for reader's convenience and because it explains why both stability properties, i.e., stability under conditioning and pasting, are needed, see also Remark 6.13 below.

Lemma 6.12. *Suppose that $\mathcal{U}: \mathbb{R}_+ \times F \rightarrow \mathfrak{P}(\Omega)$ is a measurable correspondence with non-empty and compact values such that, for all $(t, x) \in \mathbb{R}_+ \times F$ and $P \in \mathcal{U}(t, x)$, $P(X_s = x \text{ for all } s \in [0, t]) = 1$. Suppose that \mathcal{U} is stable under conditioning and pasting. Then, for every $\phi \in USC_b(F; \mathbb{R})$, the correspondence*

$$\mathcal{U}^*(t, x) \triangleq \left\{ P \in \mathcal{U}(t, x): E^P[\phi(X_T)] = \sup_{Q \in \mathcal{U}(t, x)} E^Q[\phi(X_T)] \right\}, \quad (t, x) \in \mathbb{R}_+ \times F,$$

is also measurable with non-empty and compact values and it is stable under conditioning and pasting. Further, if \mathcal{U} has convex values, then so does \mathcal{U}^ .*

Proof. We adapt the proof of [50, Lemma 12.2.2]. As ψ is assumed to be upper semicontinuous, [1, Theorem 2.43] implies that \mathcal{U}^* has non-empty and compact values. Further, [1, Theorem 18.10] and [50, Lemma 12.1.7] imply that \mathcal{U}^* is measurable. The final claim about convexity is obvious. It is left to show that \mathcal{U}^* is stable under conditioning and pasting. Take $(t, x) \in \mathbb{R}_+ \times F$, $P \in \mathcal{U}^*(t, x)$ and let τ be a stopping time such that $t \leq \tau < \infty$. We define

$$N \triangleq \left\{ \omega \in \Omega: P_\omega \triangleq \delta_{\omega(\tau(\omega))} \otimes_{\tau(\omega)} P(\cdot | \mathcal{F}_\tau)(\omega) \notin \mathcal{U}(\tau(\omega), \omega(\tau(\omega))) \right\},$$

$$A \triangleq \{\omega \in \Omega \setminus N : P_\omega \notin \mathcal{U}^*(\tau(\omega), \omega(\tau(\omega)))\}.$$

As \mathcal{U} is stable under conditioning, we have $P(N) = 0$. By [50, Lemma 12.1.9], $N, A \in \mathcal{F}_\tau$. As we already know that \mathcal{U}^* is measurable, by virtue of [50, Theorem 12.1.10], there exists a measurable map $(s, y) \mapsto R(s, y)$ such that $R(s, y) \in \mathcal{U}^*(s, y)$. We set $R_\omega \triangleq R(\tau(\omega), \omega(\tau(\omega)))$, for $\omega \in \Omega$, and note that $\omega \mapsto R_\omega$ is \mathcal{F}_τ -measurable. Further, we set

$$Q_\omega \triangleq \begin{cases} R_\omega, & \omega \in N \cup A, \\ P_\omega, & \omega \notin N \cup A. \end{cases}$$

By definition of R and N , $Q_\omega \in \mathcal{U}(\tau(\omega), \omega(\tau(\omega)))$ for all $\omega \in \Omega$. As \mathcal{U} is stable under pasting, $P \otimes_\tau Q \in \mathcal{U}(t, x)$. We obtain

$$\begin{aligned} & \sup_{Q^* \in \mathcal{U}(t, x)} E^{Q^*} [\phi(X_T)] \\ & \geq E^{P \otimes_\tau Q} [\phi(X_T)] \\ & = \int_{N \cup A} E^{\delta_\omega \otimes_{\tau(\omega)} R_\omega} [\phi(X_T)] P(d\omega) + E^P [\mathbf{1}_{N^c \cap A^c} E^P [\phi(X_T) | \mathcal{F}_\tau]] \\ & = \int_A [E^{\delta_\omega \otimes_{\tau(\omega)} R_\omega} [\phi(X_T)] - E^{\delta_\omega \otimes_{\tau(\omega)} P_\omega} [\phi(X_T)]] P(d\omega) + \sup_{Q^* \in \mathcal{U}(t, x)} E^{Q^*} [\phi(X_T)] \\ & = \int_A [E^{R_\omega} [\phi(X_T)] - E^{P_\omega} [\phi(X_T)]] P(d\omega) + \sup_{Q^* \in \mathcal{U}(t, x)} E^{Q^*} [\phi(X_T)]. \end{aligned}$$

As $E^{R_\omega} [\phi(X_T)] > E^{P_\omega} [\phi(X_T)]$ for all $\omega \in A$, we conclude that $P(A) = 0$. This proves that \mathcal{U}^* is stable under conditioning.

Next, we prove stability under pasting. Let $(t, x) \in \mathbb{R}_+ \times F$, take a stopping time τ with $t \leq \tau < \infty$, a probability measure $P \in \mathcal{U}^*(t, x)$ and an \mathcal{F}_τ -measurable map $\Omega \ni \omega \mapsto Q_\omega \in \mathfrak{P}(\Omega)$ such that, for P -a.a. $\omega \in \Omega$, $\delta_{\omega(\tau(\omega))} \otimes_{\tau(\omega)} Q_\omega \in \mathcal{U}^*(\tau(\omega), \omega(\tau(\omega)))$. As \mathcal{U} is stable under pasting, we have $P \otimes_\tau Q \in \mathcal{U}(t, x)$. Further, recall that $\delta_{\omega(\tau(\omega))} \otimes_{\tau(\omega)} P(\cdot | \mathcal{F}_\tau)(\omega) \in \mathcal{U}(\tau(\omega), \omega(\tau(\omega)))$ for P -a.a. $\omega \in \Omega$, as \mathcal{U} is stable under conditioning. Thus, we get

$$\begin{aligned} & \sup_{Q^* \in \mathcal{U}(t, x)} E^{Q^*} [\phi(X_T)] \\ & \geq E^{P \otimes_\tau Q} [\phi(X_T)] \\ & = \int E^{\delta_\omega \otimes_{\tau(\omega)} Q_\omega} [\phi(X_T)] P(d\omega) \\ & = \int E^{\delta_\omega \otimes_{\tau(\omega)} \otimes_{\tau(\omega)} Q_\omega} [\phi(X_T)] \mathbf{1}_{\{\tau(\omega) < T\}} P(d\omega) + E^P [\phi(X_T) \mathbf{1}_{\{T \leq \tau\}}] \\ & = \int \sup_{Q^* \in \mathcal{U}(\tau(\omega), \omega(\tau(\omega)))} E^{Q^*} [\phi(X_T)] \mathbf{1}_{\{\tau(\omega) < T\}} P(d\omega) + E^P [\phi(X_T) \mathbf{1}_{\{T \leq \tau\}}] \\ & \geq \int E^{\delta_\omega \otimes_{\tau(\omega)} \otimes_{\tau(\omega)} P(\cdot | \mathcal{F}_\tau)(\omega)} [\phi(X_T)] \mathbf{1}_{\{\tau(\omega) < T\}} P(d\omega) + E^P [\phi(X_T) \mathbf{1}_{\{T \leq \tau\}}] \\ & = E^P [E^P [\phi(X_T) | \mathcal{F}_\tau] \mathbf{1}_{\{\tau < T\}}] + E^P [\phi(X_T) \mathbf{1}_{\{T \leq \tau\}}] \\ & = E^P [\phi(X_T)] \\ & = \sup_{Q^* \in \mathcal{U}(t, x)} E^{Q^*} [\phi(X_T)]. \end{aligned}$$

This implies that $P \otimes_\tau Q \in \mathcal{U}^*(t, x)$. The proof is complete. \square

6.2. Proof of Theorem 3.9. We follow the proof of [9, Theorem 2.19], which adapts the proofs of [50, Theorems 6.2.3 and 12.2.3], cf. also the proofs of [15, Proposition 6.6] and [21, Proposition 3.2]. In the following, Condition 3.5 is in force.

We fix a finite time horizon $T > 0$ and a function $\psi \in USC_b(F; \mathbb{R})$. Since F is a Polish space, there exists an equivalent metric d_F with respect to which F is totally bounded (see [42, p. 43] or [50, p. 9]). Let $U_{d_F}(F)$ be the space of all bounded uniformly continuous functions on (F, d_F) endowed with the uniform topology. Then, by [42, Lemma 6.3], $U_{d_F}(F)$ is separable. Furthermore, by [42, Theorem 5.9], $U_{d_F}(F)$ is measure determining. Let $\{\sigma_n : n \in \mathbb{N}\}$ be a dense subset of $(0, \infty)$, let $\{\phi_n : n \in \mathbb{N}\}$ be a dense subset of $U_{d_F}(F)$ and let $(\lambda_N, f_N)_{N=1}^\infty$ be an enumeration of $\{(\sigma_m, \phi_n) : n, m \in \mathbb{N}\}$. For $(t, x) \in \mathbb{R}_+ \times F$, define inductively

$$\mathcal{U}_0(t, x) \triangleq \left\{ P \in \mathcal{K}(t, x) : E^P[\psi(X_T)] = \sup_{P \in \mathcal{K}(t, x)} E^P[\psi(X_T)] \right\}$$

and

$$\mathcal{U}_{N+1}(t, x) \triangleq \left\{ P \in \mathcal{U}_N(t, x) : E^P[f_N(X_{\lambda_N})] = \sup_{Q \in \mathcal{U}_N(t, x)} E^Q[f_N(X_{\lambda_N})] \right\}, \quad N \in \mathbb{Z}_+.$$

Furthermore, we set

$$\mathcal{U}_\infty(t, x) \triangleq \bigcap_{N=0}^\infty \mathcal{U}_N(t, x).$$

Thanks to Proposition 6.7 and Lemmata 6.8 and 6.11, the correspondence \mathcal{K} is measurable with non-empty convex and compact values and it is further stable under conditioning and pasting. Thus, by Lemma 6.12, the same is true for \mathcal{U}_0 and, by induction, also for every \mathcal{U}_N , $N \in \mathbb{N}$. As (arbitrary) intersections of convex and compact sets are itself convex and compact, \mathcal{U}_∞ has convex and compact values. By Cantor's intersection theorem, \mathcal{U}_∞ has non-empty values, and, by [1, Lemma 18.4], \mathcal{U}_∞ is measurable. Moreover, it is clear that \mathcal{U}_∞ is stable under conditioning, as this is the case for every \mathcal{U}_N , $N \in \mathbb{Z}_+$.

We now show that \mathcal{U}_∞ is singleton-valued. Take $P, Q \in \mathcal{U}_\infty(t, x)$ for some $(t, x) \in \mathbb{R}_+ \times F$. By definition of \mathcal{U}_∞ , we have

$$E^P[f_N(X_{\lambda_N})] = E^Q[f_N(X_{\lambda_N})], \quad N \in \mathbb{N}.$$

This implies that $P \circ X_s^{-1} = Q \circ X_s^{-1}$ for all $s \in \mathbb{R}_+$. Next, we prove that

$$E^P \left[\prod_{k=1}^n g_k(X_{t_k}) \right] = E^Q \left[\prod_{k=1}^n g_k(X_{t_k}) \right]$$

for all $g_1, \dots, g_n \in C_b(F; \mathbb{R})$, $t \leq t_1 < t_2 < \dots < t_n < \infty$ and $n \in \mathbb{N}$. We use induction over n . For $n = 1$ the claim is implied by the equality $P \circ X_s^{-1} = Q \circ X_s^{-1}$ for all $s \in \mathbb{R}_+$. Suppose the claim holds for $n \in \mathbb{N}$ and take test functions $g_1, \dots, g_{n+1} \in C_b(\mathbb{R}; F)$ and times $t \leq t_1 < \dots < t_{n+1} < \infty$. We define

$$\mathcal{G}_n \triangleq \sigma(X_{t_k}, k = 1, \dots, n).$$

Since

$$E^P \left[\prod_{k=1}^{n+1} g_k(X_{t_k}) \right] = E^P \left[E^P[g_{n+1}(X_{t_{n+1}}) | \mathcal{G}_n] \prod_{k=1}^n g_k(X_{t_k}) \right],$$

it suffices to show that P -a.s.

$$E^P[g_{n+1}(X_{t_{n+1}}) | \mathcal{G}_n] = E^Q[g_{n+1}(X_{t_{n+1}}) | \mathcal{G}_n].$$

As \mathcal{U}_∞ is stable under conditioning, there exists a null set $N_1 \in \mathcal{F}_{t_n}$ such that $\delta_{\omega(t_n)} \otimes_{t_n} P(\cdot | \mathcal{F}_{t_n})(\omega) \in \mathcal{U}_\infty(t_n, \omega(t_n))$ for all $\omega \notin N_1$. Notice that, by the tower rule, there exists a P -null set $N_2 \in \mathcal{G}_n$ such that, for all $\omega \notin N_2$ and all $A \in \mathcal{F}$,

$$\begin{aligned}
(6.2) \quad & \int \delta_{\omega'(t_n)} \otimes_{t_n} P(A | \mathcal{F}_{t_n})(\omega') P(d\omega' | \mathcal{G}_n)(\omega) \\
&= \iint \mathbf{1}_A(\omega'(t_n) \otimes_{t_n} \alpha) P(d\alpha | \mathcal{F}_{t_n})(\omega') P(d\omega' | \mathcal{G}_n)(\omega) \\
&= \iint \mathbf{1}_A(\omega(t_n) \otimes_{t_n} \alpha) P(d\alpha | \mathcal{F}_{t_n})(\omega') P(d\omega' | \mathcal{G}_n)(\omega) \\
&= \int \mathbf{1}_A(\omega(t_n) \otimes_{t_n} \omega') P(d\omega' | \mathcal{G}_n)(\omega) \\
&= (\delta_{\omega(t_n)} \otimes_{t_n} P(\cdot | \mathcal{G}_n)(\omega))(A).
\end{aligned}$$

Let $N_3 \triangleq \{P(N_1 | \mathcal{G}_n) > 0\} \in \mathcal{G}_n$. Clearly, $E^P[P(N_1 | \mathcal{G}_n)] = P(N_1) = 0$, which implies that $P(N_3) = 0$. For a moment, take $\omega \notin N_2 \cup N_3$. Using that \mathcal{U}_∞ has convex and compact values and that $\delta_{\omega'(t_n)} \otimes_{t_n} P(\cdot | \mathcal{F}_{t_n})(\omega') \in \mathcal{U}_\infty(t_n, \omega'(t_n))$ for all $\omega' \notin N_1$, we obtain that

$$\int \delta_{\omega'(t_n)} \otimes_{t_n} P(A | \mathcal{F}_{t_n})(\omega') P(d\omega' | \mathcal{G}_n)(\omega) \in \mathcal{U}(t_n, \omega(t_n)).$$

Consequently, by virtue of (6.2), we also have $\delta_{\omega(t_n)} \otimes_{t_n} P(\cdot | \mathcal{G}_n)(\omega) \in \mathcal{U}_\infty(t_n, \omega(t_n))$. Similarly, there exists a Q -null set $N_4 \in \mathcal{G}_n$ such that $\delta_{\omega(t_n)} \otimes_{t_n} Q(\cdot | \mathcal{G}_n)(\omega) \in \mathcal{U}_\infty(t_n, \omega(t_n))$ for all $\omega \notin N_4$. Set $N \triangleq N_2 \cup N_3 \cup N_4$. As $P = Q$ on \mathcal{G}_n , we get that $P(N) = 0$. For all $\omega \notin N$, the induction base implies that

$$\begin{aligned}
E^P[g_{n+1}(X_{t_{n+1}}) | \mathcal{G}_n](\omega) &= E^{\delta_{\omega(t_n)} \otimes_{t_n} P(\cdot | \mathcal{G}_n)(\omega)}[g_{n+1}(X_{t_{n+1}})] \\
&= E^{\delta_{\omega(t_n)} \otimes_{t_n} Q(\cdot | \mathcal{G}_n)(\omega)}[g_{n+1}(X_{t_{n+1}})] \\
&= E^Q[g_{n+1}(X_{t_{n+1}}) | \mathcal{G}_n](\omega).
\end{aligned}$$

The induction step is complete and hence, $P = Q$.

We proved that \mathcal{U}_∞ is singleton-valued and we write $\mathcal{U}_\infty(s, y) = \{P_{(s, y)}\}$. By the measurability of \mathcal{U}_∞ , the map $(s, y) \mapsto P_{(s, y)}$ is measurable. It remains to show the strong Markov property of the family $\{P_{(s, y)} : (s, y) \in \mathbb{R}_+ \times F\}$. Take $(s, y) \in \mathbb{R}_+ \times F$. As \mathcal{U}_∞ is stable under conditioning, for every finite stopping time $\tau \geq s$, there exists a $P_{(s, x)}$ -null set N such that, for all $\omega \notin N$,

$$\delta_{\omega(\tau(\omega))} \otimes_{\tau(\omega)} P_{(s, y)}(\cdot | \mathcal{F}_\tau)(\omega) \in \mathcal{U}_\infty(\tau(\omega), \omega(\tau(\omega))) = \{P_{(\tau(\omega), \omega(\tau(\omega)))}\}.$$

This yields, for all $\omega \notin N$, that

$$P_{(s, y)}(\cdot | \mathcal{F}_\tau)(\omega) = \delta_\omega \otimes_{\tau(\omega)} [\delta_{\omega(\tau(\omega))} \otimes_{\tau(\omega)} P_{(s, y)}(\cdot | \mathcal{F}_\tau)(\omega)] = \delta_\omega \otimes_{\tau(\omega)} P_{(\tau(\omega), \omega(\tau(\omega)))}.$$

This is the strong Markov property and consequently, the proof is complete. \square

Remark 6.13. Notice that the strong Markov property of the selection $\{P_{(s, y)} : (s, y) \in \mathbb{R}_+ \times F\}$ follows solely from the stability under conditioning property of \mathcal{U}_∞ . We emphasize that the stability under pasting property of each $\mathcal{U}_N, N \in \mathbb{Z}_+$, is crucial for its proof. Indeed, in Lemma 6.12, the fact that \mathcal{U} is stable under pasting has been used to establish that \mathcal{U}^* is stable under conditioning.

7. PROOF OF THE NONLINEAR MARKOV PROPERTY: PROPOSITION 3.2

In this section, the Standing Assumptions 2.1, 2.2 and 3.1 are in force. We need a refinement of Lemma 6.2.

Lemma 7.1. *For every $(t, \omega) \in \llbracket 0, \infty \llbracket$, $\mathcal{K}(0, \omega(t)) = \{P_t : P \in \mathcal{C}(t, \omega)\}$.*

Proof. The inclusion $\mathcal{K}(0, \omega(t)) \supset \{P_t : P \in \mathcal{C}(t, \omega)\}$ follows from Lemma 6.2. For the converse inclusion, take $P \in \mathcal{K}(0, \omega(t))$. Then, by Lemmata 6.3 and 6.9, we have $Q \triangleq \delta_\omega \otimes_t P^t \in \mathcal{C}(t, \omega)$. Since, for every $A \in \mathcal{F}$, it holds that

$$Q_t(A) = \int \mathbb{1}_{\theta_t^{-1}A}(\omega \otimes_t \omega') P^t(d\omega') = (P^t)_t(A) = P(A),$$

the proof is complete. \square

We can now follow the proof of [9, Proposition 2.8] to deduce Proposition 3.2 from the DPP (Theorem 2.5).

Proof of Proposition 3.2. For every upper semianalytic function $\psi : \Omega \rightarrow \mathbb{R}$ and any $(t, \omega) \in \llbracket 0, \infty \llbracket$, we deduce from Lemma 7.1 that

$$\mathcal{E}_t(\psi \circ \theta_t)(\omega) = \sup_{P \in \mathcal{C}(t, \omega)} E^{P^t}[\psi] = \sup_{P \in \mathcal{K}(0, \omega(t))} E^P[\psi],$$

which means nothing else than

$$\mathcal{E}_t(\psi \circ \theta_t)(\omega) = \mathcal{E}^{\omega(t)}(\psi).$$

Now, the DPP (Theorem 2.5) yields that

$$\mathcal{E}^x(\psi \circ \theta_t) = \mathcal{E}^x(\mathcal{E}_t(\psi \circ \theta_t)) = \mathcal{E}^x(\mathcal{E}^{X_t}(\psi)).$$

The proof is complete. \square

8. PROOF OF THE NONLINEAR SEMIGROUP PROPERTY: PROPOSITION 3.4

First, let us show that T_t , for every $t \in \mathbb{R}_+$, is a selfmap on the space of bounded upper semianalytic functions. Take a bounded upper semianalytic function $\psi : F \rightarrow \mathbb{R}$ and $t \in \mathbb{R}_+$. Clearly, $T_t(\psi)$ is bounded. It remains to show that $T_t(\psi)$ is also upper semianalytic. Define a function $\phi : F \times \Omega \rightarrow [-\infty, \infty)$ by

$$\phi(x, \omega) \triangleq \psi(\omega) \mathbb{1}_{\{\omega(0)=x\}} + (-\infty) \mathbb{1}_{\{\omega(0) \neq x\}}.$$

Evidently, $F \times \Omega \ni (x, \omega) \mapsto \mathbb{1}_{\{\omega(0)=x\}}$ is Borel. Hence, by [4, Lemma 7.30], the function ϕ is upper semianalytic. We set

$$\mathcal{K} \triangleq \{P \in \mathfrak{P}(\Omega) : P \in \mathcal{C}(0, \omega) \text{ for some } \omega \in \Omega\}.$$

By Corollary 5.6, the correspondence \mathcal{C} has a Borel measurable graph. Thus, the set

$$K \triangleq \{(\omega, P) \in \Omega \times \mathfrak{P}(\Omega) : P \in \mathcal{C}(0, \omega)\}$$

is Borel measurable, and consequently, the set \mathcal{K} is analytic as it is the image of K under the projection to the second coordinate ([4, Proposition 7.39]). Notice that

$$T_t(\phi)(x) = \sup_{P \in \mathcal{K}} E^P[\phi(x, X)], \quad x \in F.$$

We conclude from [4, Propositions 7.47 and 7.48] that $x \mapsto T_t(\psi)(x)$ is upper semianalytic.

Finally, we discuss the properties (i) – (iii) from Definition 3.3. The properties (ii) and (iii) are trivially satisfied and the first property (i) is implied by Proposition 3.2. The proof is complete. \square

9. PROOF OF THE USC_b -FELLER PROPERTY: THEOREM 3.6

Fix $\psi \in USC_b(F; \mathbb{R})$ and $t \in \mathbb{R}_+$, and notice that $\omega \mapsto \psi(\omega(t))$ is upper semicontinuous and bounded. Thus, by [5, Theorem 8.10.61], the map $\mathfrak{P}(\Omega) \ni P \mapsto E^P[\psi(X_t)]$ is upper semicontinuous, too. Thanks to Proposition 6.7, the correspondence $x \mapsto \mathcal{R}(x)$ is upper hemicontinuous and compact-valued. Thus, upper semicontinuity of $x \mapsto T_t(\psi)(x)$ follows from [1, Lemma 17.30]. The proof is complete. \square

10. PROOF OF PROPOSITION 3.12

The following section is devoted to the proof of Proposition 3.12, which is split into several parts. In this section, we presume that Condition 3.10 holds.

10.1. Proof of parts (i) and (ii) from Condition 3.5. Take $u = \langle y, \cdot \rangle_{H_2}^i \in U$. By virtue of [18, Problem 13, p. 151], the maps $\omega \mapsto Y^u(\omega) = \langle y, \omega \rangle_{H_2}^i$ and $\omega \mapsto L(\omega) = \|\omega\|_{H_2}$ are continuous. Furthermore, for every $R, M > 0$, we have

$$|Y_{\cdot \wedge \rho_M}^u| \mathbb{1}_{\{\|X_0\|_{H_2} \leq R\}} \leq \|y\|_{H_2}^i (R + M)^i.$$

Thus, part (i) of Condition 3.5 holds.

The correspondence Θ is convex-valued by part (ii) of Condition 3.10. Hence, also part (ii) of Condition 3.5 holds. \square

10.2. Proof of part (iii) from Condition 3.5. The linear growth assumption (iv) from Condition 3.10 implies the first part of Condition 3.5 (iii). The second part, i.e., the moment bound for $L = \|X\|_{H_2}$, follows directly from the following lemma. \square

Lemma 10.1. *For every bounded set $K \subset H_2$, every $T > 0$ and every $p > 1/\alpha$, where $\alpha \in (0, 1/2)$ is as in part (v) of Condition 3.10,*

$$\sup_{x \in K} \sup_{P \in \mathcal{R}(x)} E^P \left[\sup_{s \in [0, T]} \|X_s\|_{H_2}^p \right] < \infty.$$

Before we prove this lemma, we record another useful observation which is needed in the proof.

Lemma 10.2. *For every $x \in H_2$ and $P \in \mathcal{R}(x)$, there exists a predictable map $\mathfrak{g}: \llbracket 0, \infty \rrbracket \rightarrow G$ such that, possibly on a standard extension of the filtered probability space $(\Omega, \mathcal{F}, \mathbf{F}, P)$, the coordinate process X admits the dynamics*

$$X_t = x + \int_0^t S_{t-s} \mu(\mathfrak{g}(s, X), X_s) ds + \int_0^t S_{t-s} \sigma(\mathfrak{g}(s, X), X_s) dW_s, \quad t \in \mathbb{R}_+,$$

where W is a cylindrical standard Brownian motion.

Proof. Take $x \in H_2$ and $P \in \mathcal{R}(x)$. Thanks to Proposition 2.4, there exists a predictable function $\mathfrak{g}: \llbracket 0, \infty \rrbracket \rightarrow G$ such that the processes

$$u(X) - \int_0^\cdot \mathcal{L}^{u, \mathfrak{g}(s, X)}(X_s) ds, \quad u = \langle y, \cdot \rangle_{H_2}^i, y \in D, i = 1, 2,$$

are local P -martingales. Recall that $D \subset D(A^*)$ has the property that for every $y \in D(A^*)$ there exists a sequence $(y_n)_{n=1}^\infty \subset D$ such that $y_n \rightarrow y$ and $A^* y_n \rightarrow A^* y$. Hence, as ucp

(uniformly on compacts in probability) limits of continuous local martingales are again continuous local martingales, we conclude that all processes of the form

$$(10.1) \quad u(X) - \int_0^\cdot \mathcal{L}^{u, \mathfrak{g}(s, X)}(X_s) ds, \quad u = \langle y, \cdot \rangle_{H_2}^i, y \in D(A^*), i = 1, 2,$$

are local P -martingales. Thanks to [41, Theorem 13], in our setting the concepts of analytically weak and mild solutions are equivalent for the SPDE

$$dY_t = AY_t dt + \mu(\mathfrak{g}(t, Y), Y_t) dt + \sigma(\mathfrak{g}(t, Y), Y_t) dW_t,$$

where W is a cylindrical standard Brownian motion. Using the local P -martingale properties of the processes from (10.1), the claim of the lemma now follows as in the proof of [11, Lemma 3.6]. We omit the details. \square

Proof of Lemma 10.1. Step 1: Recap on the Factorization Method. In this step, let P be a probability measure on some filtered probability space which supports a cylindrical standard P -Brownian motion W over the Hilbert space H_2 . Take $1/p < \lambda \leq 1$ and set, for $h \in L^p([0, T]; H_2)$,

$$(10.2) \quad R_\lambda(h)(t) \triangleq \int_0^t (t-s)^{\lambda-1} S_{t-s} h(s) ds, \quad t \in [0, T].$$

Notice that R_λ is well-defined, as, by Hölder's inequality,

$$(10.3) \quad \begin{aligned} & \int_0^t (t-s)^{\lambda-1} \|S_{t-s} h(s)\|_{H_2} ds \\ & \leq \left(\int_0^T s^{p(\lambda-1)/(p-1)} \|S_s\|_{L(H_2, H_2)}^{p/(p-1)} ds \right)^{(p-1)/p} \left(\int_0^t \|h(s)\|_{H_2}^p ds \right)^{1/p}. \end{aligned}$$

This inequality shows that

$$\|R_\lambda h(t)\|_{H_2} \leq \left(\int_0^T s^{p(\lambda-1)/(p-1)} \|S_s\|_{L(H_2, H_2)}^{p/(p-1)} ds \right)^{(p-1)/p} \left(\int_0^T \|h(s)\|_{H_2}^p ds \right)^{1/p},$$

which means that R_λ is a bounded linear operator on $L^p([0, T]; H_2)$. The following lemma provides some more precise properties of R_λ , which turn out to be useful in Section 10.4 below.

Lemma 10.3 ([20, Proposition 1]). *The operator R_λ maps $L^p([0, T]; H_2)$ into $C([0, T]; H_2)$. Moreover, if the semigroup S is compact, then R_λ is compact.*

We now recall the factorization formula from [12]. For a predictable $L(H_1, H_2)$ -valued process ϕ such that

$$E^P \left[\int_0^T \|\phi_s\|_{L(H_1, H_2)}^p ds \right] < \infty,$$

the factorization formula given in [13, Theorem 5.10] yields that

$$(10.4) \quad \int_0^t S_{t-s} \phi_s dW_s = \frac{\sin(\pi\alpha)}{\pi} R_\alpha(Y)(t), \quad t \in [0, T],$$

with

$$Y_t \triangleq \int_0^t (t-s)^{-\alpha} S_{t-s} \phi_s dW_s.$$

Let $\mathfrak{f}: (0, T] \rightarrow [0, \infty]$ be a Borel function such that, for some $\alpha \in (0, 1/2)$,

$$\int_0^T \left[\frac{\mathfrak{f}(s)}{s^\alpha} \right]^2 ds < \infty,$$

and such that

$$\|S_t \phi_s\|_{L_2(H_1, H_2)} \leq \mathfrak{f}(t) \|\phi_s\|_{L(H_1, H_2)}$$

for all $t, s \in (0, T]$. Then, we obtain

$$\begin{aligned} E^P \left[\int_0^T \|Y_t\|_E^p dt \right] &\leq c_p \int_0^T E^P \left[\left(\int_0^t (t-s)^{-2\alpha} \|S_{t-s} \phi_s\|_{L_2(H_1, H_2)}^2 ds \right)^{p/2} \right] dt \\ (10.5) \quad &\leq c_p E^P \left[\int_0^T \left(\int_0^t \left[\frac{\mathfrak{f}(t-s)}{(t-s)^\alpha} \right]^2 \|\phi_s\|_{L(H_1, H_2)}^2 ds \right)^{p/2} dt \right] \\ &\leq c_p \left(\int_0^T \left[\frac{\mathfrak{f}(s)}{s^\alpha} \right]^2 ds \right)^{p/2} E^P \left[\int_0^T \|\phi_s\|_{L(H_1, H_2)}^p ds \right], \end{aligned}$$

where we use Burkholder's inequality (with constant c_p) in the first and Young's inequality in the last line. In particular, (10.3) and (10.5) yield, for all $r \in [0, T]$, that

$$(10.6) \quad E^P \left[\sup_{t \in [0, r]} \left\| \int_0^t S_{t-s} \phi_s dW_s \right\|_{H_2}^p \right] \leq C E^P \left[\int_0^r \|\phi_s\|_{L(H_1, H_2)}^p ds \right],$$

where $C > 0$ is a constant which only depends on $T, S, \mathfrak{f}, \alpha$ and p .

Step 2: A Gronwall Argument. We are in the position to prove Lemma 10.1. Let $x \in K$ and $P \in \mathcal{R}(x)$ and let \mathfrak{g} be as in Lemma 10.2. Then, using the inequalities (10.3) and (10.6), for every $M > 0$ and all $r \in [0, T]$, we obtain

$$\begin{aligned} E^P \left[\sup_{t \in [0, r \wedge \rho_M]} \|X_t\|_{H_2}^p \right] &\leq C \left(\|x\|_{H_2}^p + E^P \left[\sup_{t \in [0, r \wedge \rho_M]} \left\| \int_0^t S_{t-s} \mu(\mathfrak{g}(s, X), X_s) ds \right\|_{H_2}^p \right] \right. \\ &\quad \left. + E^P \left[\sup_{t \in [0, r \wedge \rho_M]} \left\| \int_0^t S_{t-s} \sigma(\mathfrak{g}(s, X), X_s) dW_s \right\|_{H_2}^p \right] \right) \\ &\leq C \left(\|x\|_{H_2}^p + E^P \left[\int_0^r \mu(\mathfrak{g}(s, X), X_{s \wedge \rho_M}) ds \right] \right. \\ &\quad \left. + E^P \left[\sup_{t \in [0, r]} \left\| \int_0^t S_{t-s} \sigma(\mathfrak{g}(s, X), X_{s \wedge \rho_M}) dW_s \right\|_{H_2}^p \right] \right) \\ &\leq C \left(\|x\|_{H_2}^p + \int_0^r E^P \left[\sup_{s \leq t \wedge \rho_M} \|X_s\|_{H_2}^p \right] dt \right), \end{aligned}$$

where the constant $C > 0$ depends only on the constant from Condition 3.10 (iv), $T, S, \mathfrak{f}, \alpha$ and p . Finally, we conclude the claim of the lemma from Gronwall's and Fatou's lemma. The proof is complete. \square

10.3. Proof of part (iv) from Condition 3.5. By part (iii) of Condition 3.10, the map $(g, x) \mapsto (\mathcal{L}^{u, g}(x))_{u \in U}$ is continuous. Thus, the correspondence Θ is continuous by [9, Lemma 3.2]. Furthermore, for every $(t, \omega) \in [0, \infty[$, the set $\omega([t, t+1])$ is compact and hence, again by the continuity of $(g, x) \mapsto (\mathcal{L}^{u, g}(x))_{u \in U}$, the set $\Theta(\omega([t, t+1]))$ is compact. As \mathbb{R}^U is a Fréchet space ([1, p. 206]), it follows that also the closed convex hull of $\Theta(\omega([t, t+1]))$ is compact ([1, Theorem 5.35]). \square

10.4. Proof of part (v) from Condition 3.5. We adapt the compactness method from [20]. Fix a finite time horizon $T > 0$ and take $p > 1/\alpha$, where α is as in part (v) of Condition 3.10. Let $K \subset H_2$ be a compact set and, for $R > 0$, define

$$G_R \triangleq \left\{ \omega \in C([0, T]; H_2) : \omega = S \cdot x_0 + R_1(\psi) + \frac{\sin(\pi\alpha)}{\pi} R_\alpha(\phi), x_0 \in K, \right. \\ \left. \phi, \psi \in L^p([0, T]; H_2) \text{ with } \int_0^T \|\psi(s)\|_{H_2}^p ds \leq R, \int_0^T \|\phi(s)\|_{H_2}^p ds \leq R \right\},$$

where the operators R_1 and R_α are defined as in (10.2). For every $t \in [0, T]$, the set $\{S_t x_0 : x_0 \in K\}$ is compact, since part (v) from Condition 3.10 implies that S is a compact semigroup. By [17, Lemma I.5.2], the map $[0, T] \times K \ni (t, x) \mapsto S_t x \in E$ is uniformly continuous. Thus, the Arzelà–Ascoli theorem ([27, Theorem A.5.2]) yields that the set $\{([0, T] \ni t \mapsto S_t x_0) : x_0 \in K\}$ is relatively compact in $C([0, T]; H_2)$. Recall from Lemma 10.3 that R_1 and R_α are compact operators from $L^p([0, T]; H_2)$ into $C([0, T]; H_2)$. Hence, for every $R > 0$, the set G_R is relatively compact in $C([0, T]; H_2)$.

Finally, thanks to Chebyshev’s inequality, we deduce from (10.3) and (10.5), the linear growth assumptions on μ and σ , and Lemma 10.1, that, for every $\varepsilon > 0$, there exists an $R > 0$ such that

$$\sup_{x \in K} \sup_{P \in \mathcal{R}(x)} P(\{[0, T] \ni t \mapsto X_t \in G_R\}) \geq 1 - \varepsilon.$$

By virtue of [27, Theorem 23.4], since $T > 0$ was arbitrary, we conclude that the set $\bigcup_{x \in K} \mathcal{R}(x)$ is tight and hence, relatively compact by Prohorov’s theorem. The proof is complete. \square

11. PROOF OF STRONG USC_b -FELLER PROPERTY: THEOREM 3.17

Throughout this proof, we fix a function $\psi \in USC_b(F; \mathbb{R})$ and a time $T > 0$. W.l.o.g., we assume that $|\psi| \leq 1$. We start with some auxiliary observations. Notice that Condition 3.15 implies Condition 3.10. Hence, by Corollary 3.13, there exists a strong Markov family $\{P_{(s,x)} : (s, x) \in \mathbb{R}_+ \times H\}$ such that $P_{(s,x)} \in \mathcal{K}(s, x)$ and $T_T(\psi)(x) = E^{P_{(0,x)}}[\psi(X_T)]$. To simplify our notation, we set $P_x \triangleq P_{(0,x)}$. By Lemma 10.2, with some abuse of notation, for every $x \in H$, there exists a predictable map $\mathfrak{g}^x : \llbracket 0, \infty \llbracket \rightarrow G$ and a cylindrical standard P_x -Brownian motion W such that P_x -a.s.

$$X_t = x + \int_0^t S_{t-s} \mu(\mathfrak{g}^x(s, X), X_s) ds + \int_0^t S_{t-s} dW_s, \quad t \in \mathbb{R}_+.$$

We define

$$Z^{P_x} \triangleq \exp \left(- \int_0^\cdot \langle \mu(\mathfrak{g}^x(s, X), X_s), dW_s \rangle_H - \frac{1}{2} \int_0^\cdot \|\mu(\mathfrak{g}^x(s, X), X_s)\|_H^2 ds \right),$$

and, for $M > 0$, we set

$$\rho_M \triangleq \inf(t \geq 0 : \|X_t\|_H \geq M) \wedge M.$$

For every $M > 0$, [24, Theorem 8.25] (or Novikov’s condition [27, Theorem 19.24]) yields that $Z^{\cdot \wedge \rho_M}$ is a uniformly integrable P_x -martingale. We define a probability measure Q_x^M on (Ω, \mathcal{F}) via the Radon–Nikodym derivative $dQ_x^M = Z_{\rho_M}^x dP_x$. Thanks to [34, Lemma I.0.6], the process

$$B^M \triangleq W + \int_0^{\cdot \wedge \rho_M} \mu(\mathfrak{g}^x(s, X), X_s) ds$$

is a cylindrical standard Q_x^M -Brownian motion. Thus, under Q_x^M , we have

$$X_t = x + \int_0^t S_{t-s} dB_s^M, \quad t \leq \rho_M.$$

Now, for every $t \in [0, M)$, we obtain

$$\begin{aligned} (11.1) \quad Q_x^M(\rho_M \leq t) &\leq \frac{1}{M} E^{Q_x^M} \left[\sup_{s \in [0, t \wedge \rho_M]} \|X_s\|_H \right] \\ &= \frac{1}{M} E^{Q_x^M} \left[\sup_{s \in [0, t \wedge \rho_M]} \left\| x + \int_0^s S_{s-r} B_r^M \right\|_H \right] \\ &\leq \frac{1}{M} E^{Q_x^M} \left[\sup_{s \in [0, t]} \left\| x + \int_0^s S_{s-r} B_r^M \right\|_H \right] \\ &= \frac{1}{M} E^{P_x} \left[\sup_{s \in [0, t]} \left\| x + \int_0^s S_{s-r} W_r \right\|_H \right] \rightarrow 0 \text{ as } M \rightarrow \infty. \end{aligned}$$

Hence,

$$E^{P_x} [Z_t^{P_x}] = \lim_{M \rightarrow \infty} E^{P_x} [Z_t^{P_x} \mathbf{1}_{\{\rho_M > t\}}] = \lim_{M \rightarrow \infty} Q_x^M(\rho_M > t) = 1,$$

where we use the monotone convergence theorem for the first equality. This shows that Z^{P_x} is a true P_x -martingale. Consequently, using a standard extension theorem ([27, Lemma 19.19]), there exists a probability measure Q_x on (Ω, \mathcal{F}) such that $Q_x(G) = E^{P_x} [Z_T^{P_x} \mathbf{1}_G]$ for all $G \in \mathcal{F}_T$ and $T > 0$. Using again [34, Lemma I.0.6], we obtain that the process

$$B \triangleq W + \int_0^\cdot \mu(\mathfrak{g}^x(s, X), X_s) ds$$

is a cylindrical standard Q_x -Brownian motion and, under Q_x ,

$$X_t = x + \int_0^t S_{t-s} dB_s, \quad t \in \mathbb{R}_+.$$

Now, we are in the position to prove Theorem 3.17. Let $(x^n)_{n=0}^\infty \subset H$ be a sequence such that $x^n \rightarrow x^0$ and fix $\varepsilon > 0$. By virtue of (11.1), there exists an $M' > 0$ such that

$$(11.2) \quad \sup_{n \in \mathbb{Z}_+} Q_{x^n}(\rho_{M'} \leq T) \leq \varepsilon.$$

Furthermore, by Lemma 10.1, there exists an $M^\circ > 0$ such that

$$(11.3) \quad \sup_{n \in \mathbb{Z}_+} P_{x^n}(\rho_{M^\circ} \leq T) \leq \varepsilon.$$

We set $M \triangleq M' \vee M^\circ$. By the linear growth part of Condition 3.15, there exists a constant $C = C_M > 0$ such that

$$\|\mu(g, x)\|_H \leq C$$

for all $g \in G$ and $x \in H$: $\|x\|_H \leq M$. Take $\alpha \in (0, T)$ and notice that, for all $n \in \mathbb{Z}_+$,

$$(E^{P_{x^n}} [1 - Z_{\alpha \wedge \rho_M}^{P_{x^n}}])^2 \leq E^{P_{x^n}} [1 - Z_{\alpha \wedge \rho_M}^{P_{x^n}}]^2 = E^{P_{x^n}} [(Z_{\alpha \wedge \rho_M}^{P_{x^n}})^2] - 1 \leq e^{\alpha C^2} - 1.$$

Now, take $\alpha \in (0, T)$ small enough such that, for all $n \in \mathbb{Z}_+$,

$$(11.4) \quad E^{P_{x^n}} [1 - Z_{\alpha \wedge \rho_M}^{P_{x^n}}] \leq \varepsilon.$$

Define

$$\Psi_\alpha(x) \triangleq E^{P_{\alpha, x}} [\psi(X_T)], \quad x \in H.$$

Then, for every $n \in \mathbb{N}$, using the (strong) Markov property of $\{P_{(s,x)} : (s,x) \in \mathbb{R}_+ \times H\}$ and (11.2), (11.3) and (11.4), we obtain

$$\begin{aligned}
& |E^{P_{x^n}}[\psi(X_T)] - E^{P_{x^0}}[\psi(X_T)]| \\
&= |E^{P_{x^n}}[\Psi_\alpha(X_\alpha)] - E^{P_{x^0}}[\Psi_\alpha(X_\alpha)]| \\
&\leq |E^{P_{x^n}}[\Psi_\alpha(X_\alpha)\mathbb{1}_{\{\rho_M > \alpha\}}] - E^{P_{x^0}}[\Psi_\alpha(X_\alpha)\mathbb{1}_{\{\rho_M > \alpha\}}]| + 2\varepsilon \\
&\leq |E^{P_{x^n}}[Z_\alpha^{P_{x^n}}\Psi_\alpha(X_\alpha)\mathbb{1}_{\{\rho_M > \alpha\}}] - E^{P_{x^0}}[Z_\alpha^{P_{x^0}}\Psi_\alpha(X_\alpha)\mathbb{1}_{\{\rho_M > \alpha\}}]| \\
&\quad + E^{P_{x^n}}[|1 - Z_\alpha^{P_{x^n}}|] + E^{P_{x^0}}[|1 - Z_\alpha^{P_{x^0}}|] + 2\varepsilon \\
&\leq |E^{Q_{x^n}}[\Psi_\alpha(X_\alpha)\mathbb{1}_{\{\rho_M > \alpha\}}] - E^{Q_{x^0}}[\Psi_\alpha(X_\alpha)\mathbb{1}_{\{\rho_M > \alpha\}}]| + 4\varepsilon \\
&\leq |E^{Q_{x^n}}[\Psi_\alpha(X_\alpha)] - E^{Q_{x^0}}[\Psi_\alpha(X_\alpha)]| + 6\varepsilon.
\end{aligned}$$

Thanks to [13, Theorem 9.32], we have

$$|E^{Q_{x^n}}[\Psi_\alpha(X_\alpha)] - E^{Q_{x^0}}[\Psi_\alpha(X_\alpha)]| \rightarrow 0$$

as $n \rightarrow \infty$. Hence, there exists an $N \in \mathbb{N}$ such that, for all $n \geq N$,

$$|E^{P_{x^n}}[\psi(X_T)] - E^{P_{x^0}}[\psi(X_T)]| \leq 7\varepsilon.$$

We conclude that the map $x \mapsto E^{P_x}[\psi(X_T)]$ is continuous. This finishes the proof of Theorem 3.17. \square

12. PROOF OF ERGODICITY OF NONLINEAR SPIN SYSTEMS: THEOREM 4.8

The following lemma provides the main observation needed for the proof of Theorem 4.8. Its proof is given latter in this section.

Lemma 12.1. *For every $f \in C_\uparrow(F; \mathbb{R})$ and every $t \in \mathbb{R}_+$,*

$$(12.1) \quad \underline{T}_t(f) \leq T_t(f) \leq \overline{T}_t(f).$$

Take $f \in C_\uparrow(F; \mathbb{R})$ and let μ be the invariant distribution of $(\overline{T}_t)_{t \geq 0}$ and $(\underline{T}_t)_{t \geq 0}$. Thanks to Lemma 12.1, for every $f \in C_\uparrow(F; \mathbb{R})$, $t \in \mathbb{R}_+$ and any probability measure ν on $(F, \mathcal{B}(F))$, we have

$$(12.2) \quad \int \underline{T}_t(f) d\nu \leq \int T_t(f) d\nu \leq \int \overline{T}_t(f) d\nu.$$

Since μ is the invariant distribution of $(\overline{T}_t)_{t \geq 0}$ and $(\underline{T}_t)_{t \geq 0}$, we have, for all $f \in C_\uparrow(F; \mathbb{R})$ and $t \in \mathbb{R}_+$,

$$\int \underline{T}_t(f) d\mu = \int \overline{T}_t(f) d\mu = \int f d\mu.$$

Hence, taking $\nu = \mu$ in (12.2) yields that μ is an invariant distribution of $(T_t)_{t \geq 0}$.

It remains to prove that

$$\lim_{t \rightarrow \infty} \int T_t(f) d\nu = \int f d\mu$$

for any probability measure ν on $(F, \mathcal{B}(F))$ and any $f \in C_\uparrow(F; \mathbb{R})$, see Remark 4.4 (i).

As $(\overline{T}_t)_{t \geq 0}$ and $(\underline{T}_t)_{t \geq 0}$ are ergodic with unique invariant distribution μ , we have

$$\lim_{t \rightarrow \infty} \int \underline{T}_t(f) d\nu = \lim_{t \rightarrow \infty} \int \overline{T}_t(f) d\nu = \int f d\mu,$$

for any $f \in C_\uparrow(F; \mathbb{R})$ and any probability measure ν on $(F, \mathcal{B}(F))$. By virtue of (12.2), this completes the proof of Theorem 4.8. \square

Proof of Lemma 12.1. We use a strategy we learned from the proof of [3, Theorem 2.2]. Fix $f \in C_\uparrow(F; \mathbb{R})$ and fix a finite time horizon $T > 0$. Set

$$g(t, \eta) \triangleq \underline{T}_{T-t}(f)(\eta), \quad (t, \eta) \in [0, T] \times F.$$

Notice that $t \mapsto g(t, \eta)$ is continuously differentiable for every $\eta \in F$ ([33, Theorem 3.16]).

Lemma 12.2. *For every $\eta_0 \in F$ and $P \in \mathcal{R}(\eta_0)$, there exists a predictable function $\mathbf{g}: [0, \infty[\rightarrow G$ such that*

$$(12.3) \quad g(\cdot, X) - \int_0^\cdot \left(\frac{dg}{dt}(s, X_s) + \sum_{x \in S} c(\mathbf{g}(s, X), x, X_s) [g(s, (X_s)_x) - g(s, X_s)] \right) ds$$

is a P -martingale on the time interval $[0, T]$.

Proof. By Proposition 2.4, there exists a predictable function $\mathbf{g}: [0, \infty[\rightarrow G$ such that

$$f(X) - f(\eta_0) - \int_0^\cdot \sum_{x \in S} c(\mathbf{g}(s, X), x, X_s) [f((X_s)_x) - f(X_s)] ds, \quad f \in U,$$

are P -martingales. As U is dense in $C(F; \mathbb{R})$, all processes of the form

$$(12.4) \quad f(X) - f(\eta_0) - \int_0^\cdot \sum_{x \in S} c(\mathbf{g}(s, X), x, X_s) [f((X_s)_x) - f(X_s)] ds, \quad f \in C(F; \mathbb{R}),$$

are P -martingales. For $(t, x) \in \mathbb{R}_+ \times F$, we have

$$g(t, x) = \sum_{k \in F} g(t, k) \mathbf{1}\{x = k\} \triangleq \sum_{k \in F} h_k(t) f_k(x).$$

Using integration by parts and the fact that all processes in (12.4) are P -martingales, we obtain that

$$\begin{aligned} dg(s, X_s) &= d[P\text{-martingale}]_s \\ &\quad + \sum_{k \in F} \left(h'_k(s) f_k(X_s) + h_k(s) \sum_{x \in S} c(\mathbf{g}(s, X), x, X_s) [f_k((X_s)_x) - f_k(X_s)] \right) ds \\ &= d[P\text{-martingale}]_s \\ &\quad + \left(\frac{dg}{dt}(s, X_s) + \sum_{x \in S} c(\mathbf{g}(s, X), x, X_s) [g(s, (X_s)_x) - g(s, X_s)] \right) ds. \end{aligned}$$

This implies our claim. \square

Fix $\eta_0 \in F$ and take an arbitrary $P \in \mathcal{R}(\eta_0)$. From [33, Theorem 3.16], we get that

$$-\frac{dg}{dt}(s, \eta) = \sum_{x \in S} \underline{c}(x, \eta) [g(s, \eta_x) - g(s, \eta)]$$

for all $(s, \eta) \in [0, T] \times F$. Hence, we obtain

$$\begin{aligned} g(\cdot, X) &= P\text{-martingale} \\ &\quad + \int_0^\cdot \sum_{x \in S} (c(\mathbf{g}(s, X), x, X_s) - \underline{c}(s, X_s)) [g(s, (X_s)_x) - g(s, X_s)] ds, \end{aligned}$$

where \mathfrak{g} is as in Lemma 12.2. By our assumption that $(\underline{T}_t)_{t \geq 0}$ is attractive, [32, Theorem II.2.2] yields that $(x \mapsto g(t, x)) \in C_{\uparrow}(F; \mathbb{R})$ for all $t \in [0, T]$. Thus, for every $t \in [0, T]$,

$$g(t, \eta_x) - g(t, \eta) \begin{cases} \geq 0, & \eta(x) = 0, \\ \leq 0, & \eta(x) = 1, \end{cases}$$

which, by definition of \underline{c} , implies that

$$g(\cdot, X) = P\text{-martingale} + \text{increasing process} = P\text{-submartingale.}$$

Consequently, we get

$$\underline{T}_T(f)(\eta_0) = g(0, \eta_0) \leq E^P[g(T, X_T)] = E^P[f(X_T)] \leq T_T(f)(\eta_0).$$

We conclude that $\underline{T}_t(f)(\eta) \leq T_t(f)(\eta)$ for all $(t, f, \eta) \in \mathbb{R}_+ \times C_{\uparrow}(F; \mathbb{R}) \times F$, which is the first inequality from (12.1). The second inequality follows the same way. We omit the details. \square

ACKNOWLEDGMENT

The author thanks Lars Niemann for many interesting discussions.

REFERENCES

- [1] C. D. Aliprantis and K. B. Border. *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer Berlin Heidelberg, 3rd ed., 2006.
- [2] H. Bauer. *Measure and Integration Theory*. Walter de Gruyter Berlin New York, 2001.
- [3] J. Bergenthum and L. Rüschemdorf. Comparison of semimartingales and Lévy processes. *The Annals of Probability*, 35(1):228–254, 2007.
- [4] D. P. Bertsekas and S. E. Shreve. *Stochastic Optimal Control: The Discrete Time Case*. Academic Press, New York San Francisco London, 1978.
- [5] V. I. Bogachev. *Measure Theory*. Springer Berlin Heidelberg, 2007.
- [6] D. Criens. Limit theorems for cylindrical martingale problems associated with Lévy generators. *Journal of Theoretical Probability*, 33:866–905, 2020.
- [7] D. Criens. Propagation of chaos for weakly interacting mild solutions to stochastic partial differential equations. arXiv:2107.13397v4, 2021.
- [8] D. Criens and L. Niemann. Nonlinear continuous semimartingales. arXiv:2204.07823v2, 2022.
- [9] D. Criens and L. Niemann. Markov selections and Feller properties of nonlinear diffusions. arXiv:2205.15200v2, 2022.
- [10] D. Criens and L. Niemann. A class of multidimensional nonlinear diffusions with the Feller property. arXiv:2205.15692v1, 2022.
- [11] D. Criens and M. Ritter. On a theorem by A.S. Cherny for semilinear stochastic partial differential equations. *Journal of Theoretical Probability*, online first, 2021.
- [12] G. Da Prato, S. Kwapien, and J. Zabczyk. Regularity of solutions of linear stochastic equations in Hilbert spaces. *Stochastics*, 23(1):1–23, 1988.
- [13] G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions*. Cambridge University Press, 2nd edition, 2014.
- [14] R. Denk, M. Kupper, and M. Nendel. A semigroup approach to nonlinear Lévy processes. *Stochastic Processes and their Applications*, 130:1616–1642, 2020.

- [15] N. El Karoui, D. Nguyen and M. Jeanblanc-Picqué. Compactification methods in the control of degenerate diffusions: existence of an optimal control. *Stochastics*, 20(3):169–219, 1987.
- [16] N. El Karoui and X. Tan. Capacities, measurable selection and dynamic programming part II: application in stochastic control problems. arXiv:1310.3364v2, 2015.
- [17] K.-J. Engel and R. Nagel. *One-Parameter Semigroups for Linear Evolution Equations*. Springer New York, 2000.
- [18] S. N. Ethier and T. G. Kurtz. *Markov Processes: Characterization and Convergence*. Wiley, 2005.
- [19] T. Fadina, A. Neufeld, and T. Schmidt. Affine processes under parameter uncertainty. *Probability, Uncertainty and Quantitative Risk*, 4(5), 2019.
- [20] D. Gatarek and B. Goldys. On weak solutions of stochastic equations in Hilbert spaces. *Stochastics and Stochastic Reports*, 46(1-2):41–51, 1994.
- [21] U. G. Haussmann. Existence of optimal markovian controls for degenerate diffusions. In: Christopheit, N., Helmes, K., Kohlmann, M. (eds.) *Stochastic Differential Systems*, Lecture Notes in Control and Information Sciences, vol 78, Springer Berlin Heidelberg, 1986.
- [22] J. Hollender. *Lévy-Type Processes under Uncertainty and Related Nonlocal Equations*. PhD thesis, TU Dresden, 2016.
- [23] M. Hu and S. Peng. G-Lévy processes under sublinear expectations. *Probability, Uncertainty and Quantitative Risk*, 6(1), 2021.
- [24] J. Jacod. *Calcul stochastique et problèmes de martingales*. Springer Berlin Heidelberg New York, 1979.
- [25] J. Jacod. Weak and strong solutions of stochastic differential equations. *Stochastics*, 3:171–191, 1980.
- [26] J. Jacod and A. N. Shiryaev. *Limit Theorems for Stochastic Processes*. Springer Berlin Heidelberg, 2nd ed., 2003.
- [27] O. Kallenberg. *Foundations of Modern Probability*. Springer International Publishing, 3rd edition, 2021.
- [28] N. Krylov. On the selection of a Markov process from a system of processes and the construction of quasi-diffusion processes. *Mathematics of the USSR-Izvestiya*, 7(3):691–708, 1973.
- [29] F. Kühn. Viscosity solutions to Hamilton–Jacobi–Bellman equations associated with sublinear Lévy(-type) processes. *ALEA*, 16:531–559, 2019.
- [30] F. Kühn. On infinitesimal generators of sublinear Markov semigroups. *Osaka Journal of Mathematics*, 58(3):487–508, 2021.
- [31] T. M. Liggett. The stochastic evolution of infinite systems of interacting particles. In: *Ecole d’Été de Probabilités de Saint-Flour VI 1976*, pp. 188 – 249, Springer Berlin Heidelberg New York, 1977.
- [32] T. M. Liggett. *Interacting Particle Systems*. Springer Berlin Heidelberg, reprint of 1985 ed., 2005.
- [33] T. M. Liggett. *Continuous Time Markov Processes*. American Mathematical Society, 2010.
- [34] W. Liu and M. Röckner. *Stochastic Partial Differential Equations: An Introduction*. Springer International Publishing Switzerland, 2015.
- [35] M. Nendel. Markov chains under nonlinear expectation. *Mathematical Finance*, 31(1):474–507, 2021.

- [36] M. Nendel and M. Röckner. Upper envelopes of families of Feller semigroups and viscosity solutions to a class of nonlinear Cauchy problems. *SIAM Journal on Control and Optimization*, 59(6):4400–4428, 2021.
- [37] A. Neufeld and M. Nutz. Measurability of semimartingale characteristics with respect to the probability law. *Stochastic Processes and their Applications*, 124:3819–3845, 2014.
- [38] A. Neufeld and M. Nutz. Nonlinear Lévy processes and their characteristics. *Transactions of the American Mathematical Society*, 369:69–95, 2017.
- [39] M. Nisio. On a Non-linear semi-group attached to stochastic optimal control. *Publications of the Research Institute for Mathematical Sciences, Kyoto University*, 13:513–537, 1976.
- [40] M. Nutz and R. van Handel. Constructing sublinear expectations on path space. *Stochastic Processes and their Applications*, 123(8):3100–3121, 2013.
- [41] M. Ondreját. Uniqueness for stochastic evolution equations in Banach spaces. *Dissertationes Mathematicae (Rozprawy Matematyczne)*, 426:63, 2004.
- [42] K. R. Parthasarathy. *Probability Measures on Metric Space*. Academic Press, New York London, 1976.
- [43] S. Peng. G-expectation, G-Brownian motion and related stochastic calculus of Itô type. In F. E. Benth et. al., editors, *Stochastic Analysis and Applications: The Abel Symposium 2005*, pages 541–567, Springer Berlin Heidelberg, 2007.
- [44] S. Peng. Multi-dimensional G-Brownian motion and related stochastic calculus under G-expectation. *Stochastic Processes and their Applications*, 118:2223–2253, 2008.
- [45] D. Pollard. *Convergence of Stochastic Processes*. Springer New York, 1984.
- [46] P. E. Protter. *Stochastic Integration and Differential Equations*. Springer Berlin Heidelberg, 2nd ed., 2004.
- [47] G. Reuter and P. Riley. The Feller property for Markov semigroups on a countable state space. *Journal of the London Mathematical Society*, s2-5(2), 267–275, 1972.
- [48] L. C. G. Rogers and D. Williams. *Diffusions, Markov Processes, and Martingales. Volume 2: Itô Calculus*. Cambridge University Press, 2nd ed., 2000.
- [49] M. Sion. *A Theory of Semigroup Valued Measures*. Springer Berlin Heidelberg, 1973.
- [50] D. W. Stroock and S.R.S. Varadhan. *Multidimensional Diffusion Processes*. Springer Berlin Heidelberg, reprint of 1997 ed., 2006.
- [51] G. G. Yin and Q. Zhang. *Continuous-Time Markov Chains and Applications*. Springer Science+Business Media, 2nd edition, 2013.