

On the Turán number of $K_m \vee C_{2k-1}$

Jingru Yan *

Department of Mathematics, East China Normal University, Shanghai 200241, China

Abstract

Given a graph H and a positive integer n , the Turán number of H for the order n , denoted $ex(n, H)$, is the maximum size of a simple graph of order n not containing H as a subgraph. Given graphs G and H , the notation $G \vee H$ means the joint of G and H . $\chi(G)$ denotes the chromatic number of a graph G . Since $\chi(K_m \vee C_{2k-1}) = m + 3$ and there is an edge $e \in E(K_m \vee C_{2k-1})$ such that $\chi(K_m \vee C_{2k-1} - e) = m + 2$, by the Simonovits theorem, $ex(n, K_m \vee C_{2k-1}) = \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor$ for sufficiently large n . In this paper, we prove that $2(m+2)k - 3(m+2) - 1$ is large enough for n .

Keywords. Turán number, even wheels, generalized wheels

Mathematics Subject Classification. 05C35, 05C38

1 Introduction

We will deal with only finite nontrivial simple graphs. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. The order n and size $e(G)$ of a graph G are its number of vertices and edges, respectively. The degree of $v \in V(G)$ is $d_G(v)$. $\Delta(G)$ and $\chi(G)$ denote the maximum degree and chromatic number of a graph G , respectively. For terminology and notations not explicitly described in this paper, the reader is referred the books [3, 11].

Given a graph H and a positive integer n , the *Turán number* of H for the order n , denoted $ex(n, H)$, is the maximum size of a simple graph of order n not containing H as a subgraph. A graph is *H-free* if it contains no H as a subgraph. A graph G of order n is an *extremal graph* for H if G is H -free with $ex(n, H)$ edges. Determining the Turán number of various graphs H is one of the main topics in extremal graph theory [2]. A classic result of Mantel [6] from 1907 states that $ex(n, K_3) = \lfloor n^2/4 \rfloor$ and the balanced complete bipartite graph $K_{\lfloor n/2 \rfloor, \lfloor n/2 \rfloor}$ is the unique extremal graph. The famous Turán theorem [8] states that $ex(n, K_{r+1}) = \lfloor (r-1)n^2/(2r) \rfloor$ and the balanced complete r -partite graph is the unique extremal graph.

*E-mail address: mathyjr@163.com

Given graphs G and H , the notation $G \vee H$ means the *joint* of G and H . K_n and C_n stand for the complete graph and cycle of order n , respectively. A *wheel* W_n is a graph obtained by connecting a single vertex to all vertices of C_{n-1} , that is $W_n = K_1 \vee C_{n-1}$. A *generalized wheel* $W_{s,t} = K_s \vee C_t$. In recent years, several authors reported results on wheels [1, 4, 5, 13], generalized wheels [9, 10] and disjoint union of wheels [12]. In this paper, we consider the Turán number of the generalized wheel $W_{m,2k-1} = K_m \vee C_{2k-1}$.

Theorem 1. [7] *Let H be a graph with $\chi(H) \geq r + 1$. If there is an edge e such that $\chi(H - e) = r$, then there exists an n_0 such that if $n > n_0$ then the balanced complete r -partite graph is the unique extremal graph for H .*

Since $\chi(W_{m,2k-1}) = m+3$ and there is an edge $e \in E(W_{m,2k-1})$ such that $\chi(W_{m,2k-1} - e) = m + 2$, by Theorem 1 and the Turán theorem,

$$ex(n, W_{m,2k-1}) = ex(n, K_{m+3}) = \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor$$

for sufficiently large n . In this paper, we prove that $2(m+2)k - 3(m+2) - 1$ is large enough for n .

Theorem 2. [4] *For all $k \geq 3$ and $n \geq 6k - 10$, $ex(n, W_{2k}) = \lfloor n^2/3 \rfloor$.*

Note that when $k = 2$, $K_m \vee C_{2k-1} = K_{m+3}$. Thus it suffices to consider the case $k \geq 3$.

Theorem 3. *If $k \geq 3$ and $n \geq 2(m+2)k - 3(m+2) - 1$, then*

$$ex(n, W_{m,2k-1}) = \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor.$$

2 Proof of Theorem 3

Proof. First note that the case $m = 1$ of Theorem 3 is exactly Theorem 2. Thus it suffices to consider the case $m \geq 2$.

It is easy to check that the balanced complete $(m+2)$ -partite graph of order $n \geq 2(m+2)k - 3(m+2) - 1$ does not contain $W_{m,2k-1}$ as a subgraph, where $m \geq 2$ and $k \geq 3$. Then we have

$$ex(n, W_{m,2k-1}) \geq \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor$$

for $n \geq 2(m+2)k - 3(m+2) - 1$.

Let G be a $W_{m,2k-1}$ -free graph with order $n \geq 2(m+2)k - 3(m+2) - 1$ and size $\lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor + 1$. We use induction on m .

Next we show that the theorem holds for $m = 2$. Let G_1 be a $W_{2,2k-1}$ -free graph with order $n \geq 8k - 13$ and size $\lfloor 3n^2/8 \rfloor + 1$. Since

$$\frac{(n - \lfloor \frac{n}{4} \rfloor - 1)n}{2} < \lfloor \frac{3n^2}{8} \rfloor + 1,$$

we have $\Delta(G_1) \geq n - \lfloor n/4 \rfloor$.

Suppose that $\Delta(G_1) = n - 1$. Let $v \in V(G_1)$ with degree $d_{G_1}(v) = n - 1$. Then $e(G_1 - v) = \lfloor 3n^2/8 \rfloor + 1 - (n - 1)$. Note that $n - 1 \geq 8k - 14 > 6k - 10$ for $k \geq 3$ and

$$e(G_1 - v) = \lfloor \frac{3n^2}{8} \rfloor + 1 - (n - 1) > \lfloor \frac{(n - 1)^2}{3} \rfloor = ex(n - 1, W_{2k}).$$

By Theorem 2, $G_1 - v$ contains W_{2k} and then G_1 contains $W_{2,2k-1}$, a contradiction.

Next suppose $\Delta(G_1) = n - p$, where $2 \leq p \leq \lfloor n/4 \rfloor$. Let $v \in V(G_1)$ with degree $d_{G_1}(v) = n - p$. Let G'_1 be a graph of order $n - p$, obtained from G_1 by deleting v and $p - 1$ vertices not adjacent to v . Then

$$e(G'_1) \geq e(G_1) - p(n - p) = \lfloor \frac{3n^2}{8} \rfloor + 1 - p(n - p).$$

A simple calculation shows that the maximum value of the expression $p(n - p) + \lfloor (n - p)^2/3 \rfloor$ when $2 \leq p \leq \lfloor n/4 \rfloor$ is equal to $\lfloor 3n^2/8 \rfloor$. Note that $n - p \geq n - \lfloor n/4 \rfloor \geq 6k - 10$ for $n \geq 8k - 13$ and

$$e(G'_1) \geq \lfloor \frac{3n^2}{8} \rfloor + 1 - p(n - p) > \lfloor \frac{(n - p)^2}{3} \rfloor = ex(n - p, W_{2k}).$$

By Theorem 2, G'_1 contains W_{2k} and then G_1 contains $W_{2,2k-1}$, a contradiction. This completes the proof of the case $m = 2$ of Theorem 3.

Therefore, by the induction hypothesis, if $n \geq 2(m+1)k - 3(m+1) - 1$, then

$$ex(n, W_{m-1,2k-1}) = \lfloor \frac{mn^2}{2(m+1)} \rfloor.$$

Now we show that the theorem holds for m . Recall that G is a $W_{m,2k-1}$ -free graph with order $n \geq 2(m+2)k - 3(m+2) - 1$ and size $\lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor + 1$. Since

$$\frac{(n - \lfloor \frac{n}{m+2} \rfloor - 1)n}{2} < \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor + 1,$$

we have $\Delta(G) \geq n - \lfloor n/(m+2) \rfloor$.

Suppose that $\Delta(G) = n - 1$. Let $v \in V(G)$ with degree $d_G(v) = n - 1$. Then $e(G - v) = \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor + 1 - (n - 1)$. Note that $n - 1 \geq 2(m+2)k - 3(m+2) - 1 - 1 > 2(m+1)k - 3(m+1) - 1$ for $k \geq 3$ and

$$e(G - v) = \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor + 1 - (n - 1) > \lfloor \frac{m(n-1)^2}{2(m+1)} \rfloor = ex(n-1, W_{m-1, 2k-1}).$$

By Theorem 2, $G - v$ contains $W_{m-1, 2k-1}$ and then G contains $W_{m, 2k-1}$, a contradiction.

Next suppose $\Delta(G) = n - p$, where $2 \leq p \leq \lfloor n/(m+2) \rfloor$. Let $v \in V(G)$ with degree $d_G(v) = n - p$. Let G' be a graph of order $n - p$, obtained from G by deleting v and $p - 1$ vertices not adjacent to v . Then

$$e(G') \geq e(G) - p(n - p) = \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor + 1 - p(n - p).$$

A simple calculation shows that the maximum value of the expression $p(n - p) + \lfloor \frac{m(n-p)^2}{2(m+1)} \rfloor$ when $2 \leq p \leq \lfloor n/(m+2) \rfloor$ is equal to $\lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor$. Note that $n - p \geq n - \lfloor n/(m+2) \rfloor \geq 2(m+1)k - 3(m+1) - 1$ for $n \geq 2(m+2)k - 3(m+2) - 1$ and

$$e(G') \geq \lfloor \frac{(m+1)n^2}{2(m+2)} \rfloor + 1 - p(n - p) > \lfloor \frac{m(n-p)^2}{2(m+1)} \rfloor = ex(n-p, W_{m-1, 2k-1}).$$

By Theorem 2, G' contains $W_{m-1, 2k-1}$ and then G contains $W_{m, 2k-1}$, a contradiction. This completes the proof of Theorem 3. \square

Acknowledgement This research was supported by the NSFC grant 12271170 and Science and Technology Commission of Shanghai Municipality (STCSM) grant 22DZ2229014.

References

- [1] M. S. A. Bataineh, M. M. M. Jaradat, A. M. M. Jaradat, Edge maximal graphs containing no specific wheels, *Jordan J. Math. Stat.* 8 (2015), no. 2, 107–120.
- [2] B. Bollobás, *Extremal Graph Theory*, Academic Press, London-New York, 1978.
- [3] J. A. Bondy, U.S.R. Murty, *Graph Theory*, in: GTM, vol. 244, Springer, 2008.
- [4] T. Dzido, A note on Turán numbers for even wheels, *Graphs Combin.* 29 (2013), no. 5, 1305–1309.

- [5] T. Dzido, A. Jastrzębski, Turán numbers for odd wheels, *Discrete Math.* 341 (4) (2018), pp. 1150-1154
- [6] W. Mantel, Problem 28, *Wiskundige Opgaven*, 10(1907), 60-61.
- [7] M. Simonovits, A method for solving extremal problems in graph theory, stability problems. In: *Theory of Graphs*. Academic Press, New York (1968)
- [8] P. Turán, On an extremal problem in graph theory, *Mat. Fiz. Lapok* 48, 436–452 (1941)
- [9] L. Wang, The Ramsey Numbers of Trees Versus Generalized 6-Wheels or Generalized 7-Wheels, *Graphs Combin.* 38 (2022), no. 5, Paper No. 153.
- [10] L. Wang, Y. Chen, The Ramsey numbers of trees versus generalized wheels, *Graphs Combin.* 35 (2019), no. 1, 189–193.
- [11] D. B. West, *Introduction to Graph Theory*, Prentice Hall, Inc., Upper Saddle River, NJ, (1996).
- [12] C. Xiao, O. Zamora, A note on the Turán number of disjoint union of wheels, *Discrete Math.* 344 (2021), no. 11, Paper No. 112570, 7 pp.
- [13] L. Yuan, Extremal graphs for odd wheels, *J. Graph Theory* 98 (2021), no. 4, 691–707.