

ON INDUCED MATCHING NUMBERS OF STACKED-BOOK GRAPHS

TAYO CHARLES ADEFOKUN¹

ABSTRACT. Suppose that G is a simple, undirected graph. An induced matching in G is a set of edges M in the edge set $E(G)$ of G such that if $e_1, e_2 \in M$, then no endpoint v_1, v_2 of e_1 and e_2 respectively is incident to any edge $e_k \in E(G)$ such that e_k is incident to any edge in M . Denoted by $im(G)$, the maximum cardinal number of M is known as the induced matching number of G . In this work, we probe $im(G)$ where $G = G_{m,n}$, which is the stacked-book graph obtained by the Cartesian product of the star graph S_m and path P_n .

1. INTRODUCTION

Suppose that G is a graph with $E(G)$ as the edge set of G while $V(G)$ denotes the vertex set of G . Let M be subset of $E(G)$ such that for every $e_1, e_2 \in M$ there is no such edge in $E(G)$ to which any of the end points of e_1 and e_2 are commonly adjacent. Maximum Induced matching (MIM) problem is the generalization of the older graph matching problem, and it was introduced in [8].

Suppose that M is the largest induced matching in G then the cardinal number of M , denoted by $im(G)$ is called the maximum induced matching number of G . Many work has has been on this subject. It has attracted interest mostly because of it is theoretically interested and it has a number of direct applications. In [8], the authors described MIM problem as "risk free" marriage where married couples who are perfectly matched are identified. Its usefulness in cryptography is also obvious. Cameron in her earlier work [3] showed that even though the MIM problem is NP-complete for bipartite graphs, it is easier to resolve for chordal graphs. This was also confirmed for circular graph in [5]. Golubic and Lewenstein [6] established that the a relationship between MIM number and redundancy number in graphs and also showed that the MIM problem is polynomial-time solvable for tree graphs, while [4] investigated the MIM problem in intersection graphs. Recent works on MIM problem include [7] where

Key words and phrases. Stacked-book graphs, Maximum Induced Matching Number, Cartesian product Star graph paths

2010 *Mathematics Subject Classification.* Primary: 05C70, 05C15.

the MIM number was extensively probed for grids $G_{n,m} = P_n \square P_m$, the Cartesian product of paths P_n and P_m . For odd nm , a bound $im(G_{n,m}) \leq \lfloor \frac{nm+1}{4} \rfloor$ was obtained. The bound was tightened in [2] and further in [1]. In [9] investigation was made into obtaining exact algorithm for MIM problem of graphs on n -vertices.

In this work, we probe the maximum induced matching problem for stacked-book graph $G_{m,n}$ class which are graphs obtained from the Cartesian product of star graphs S_m and paths P_n . The MIM numbers are obtained for the initial range of these graphs while lower bounds of MIM number are derived for the general class.

2. DEFINITIONS

To make this works self-contained, we give the following definitions, which we shall adopt in the course of the paper. Definitions that are not considered as general will be given at the point of application.

The vertex set of graph G is $V(G)$ and M is a subset of $E(G)$, the edge set of G , and M is the induced matching of G . A vertex $v \in V(G)$ is called saturated if $v \in V(G)$ and unsaturated if otherwise. A star graph S_m contains a central vertex v_1 (except if specifically indicated otherwise) with $m-1$ leaves, which are all incident to v_1 as pendants. A path P_n contains n edges and $n-1$ paths, while a cycle C_m contains m vertices and n edges. Supposed that u and v are members of $V(G)$, then $d(u, v)$ is a positive integer, which is the distance between u and v in G . A vertex $v \in V(G)$ is called unstaurable if by the virtue of its position, can not be saturated either because of its distance from a saturated vertex or it is a the the right distance but not adjacent to a vertex that can be saturated in other to form an edge in the induced matching. A saturable vertex therefore, is the opposite of an un saturable vertex. The diameter of a graph is the largest distance between any two vertices on a graph u and v , demoted by $diam(G)$. The set $[a, b]$ denoted set of integers from a to b while $[a]$ is a shortened for for $[1, a]$.

2.1. Structure of a Stacked-book graph. The stacked-book graph is the Cartesian product $S_m \square P_n$ of a star graph S_m and path P_n . Structurally, a $S_m \square P_n$ contains n number of S_m stars such that there exist the $E(G') \in E(S_m \square P_n)$, where $E(G') = \{v_i u_i : v_i \in V(S_m(i)); u_i \in V(S_m(i+1)), i \in [n]\}$. Clearly, $E(S_m \square P_n) = E(G') \cup E(\cup_{i=1}^n S_m(i))$, where $S_m(i)$ is designated as the the i th S_m star graph for all $1 \leq i \leq n$

2.2. Initial Results. The following results are obvious

Theorem 2.1. *Let P_n be a path graph on n vertices. Then, $im(P_n) = \lfloor \frac{n-1}{3} \rfloor$.*

Theorem 2.2. *Let C_n be a circle graph on n vertices. Then $im(C_n) = \lfloor \frac{n}{3} \rfloor$.*

Theorem 2.3. [7] *Suppose that $G_{3,n}$ is a grid graph obtained by the Cartesian product $P_3 \square P_n$, where n is even or odd. Then for a positive integer k ,*

$$im(P_3 \square P_n) = \begin{cases} \lceil \frac{3n}{4} \rceil & \text{if } n \text{ is even;} \\ \frac{3(n-1)}{4} & \text{if } n = 4k + 1 \\ \frac{3(n-1)+2}{4} & \text{if } n = 4k + 3 \end{cases}$$

3. RESULT

Now we present the results we have obtained in this work. First we show a result on induced matching on star graph S_m .

Theorem 3.1. *Let S_m be a given a star graph such that the central vertex is v_1 and it is adjacent to $m - 1$ leaves. Then $im(S_m) = 1$*

(The implication of this result is that every star contains at most one element in its induced matching set.)

Proof. Let S_m be a start with v_1 being the central vertex. Then v_1 is saturated. Suppose that $v_k \in V(S_m)$, $k \leq m$, such that v_k is saturated. Then $v_1 v_k \in M$. Now for all i , $i \neq k$, $v_i \in V(G)$ is unsaturated since the $diam(S_m) = 2$. Thus $im(S_m) = |m| = 1$. \square

Now we present our first results on the induced matching of stacked-book graph $G_{m,n}$.

Lemma 3.1. *Suppose that $G_{m,n}$ is a stacked-book graph. Then if the induced matching number of $G_{m,n}$ is obtained, then the central vertices $v_1(1)$, $v_1(2)$ of factor stars $S_m(1)$ and $S_m(2)$ of $G_{m,n}$ are not unsaturated.*

Proof. Suppose that v_1 and u_1 are the central vertices of $S_m(1)$ and $S_m(2)$ stars. Now, suppose that v_1 is saturated. Then either $v_1 v_i \in M$, $v_i \in V(S_m(1))$ for some $2 \leq i < m$ or $v_1 u_1 \in M$. Suppose that $v_1 v_i \in M$. By Theorem 3.1, if v_1 is saturated, then at least $m - 2$ vertices on S_m will be unsaturated. Thus, for all $v_i \in V(S_m(1))$, $v_i u_i \notin M$. Same argument holds if $u_1 \in S_m(2)$ is saturated. Thus, $im(G_{m,2}) = 1$. Now, suppose that $v_1 u_1 \in M$. Since $d(v_1, v_i) = 1 = d(u_1, u_i)$ for all $i \in [2, m]$, then v_i, u_i are unsaturated for all $i \in [2, m]$. Clearly there exists a path P_5 in $G_{m,2}$. From Theorem 2.1, P_5 contains two edges in M of $G_{m,2}$. Thus a contradiction. \square

Now we present the first theorem,

Theorem 3.2. *Let $G_{m,2}$ be a stacked-book graph. Then $im(G_{m,2}) = m - 1$.*

Proof. Let $G_{m,2}$ be a stacked-book graph. Then there exist $S_m(1), S_m(2) \subseteq G_{m,2}$ with vertices $v_1, v_2 \cdots v_m$ and $u_1, u_2, \cdots u_m$ and a path $P_5(i) = v_i \rightarrow u_i \rightarrow u_1 \rightarrow u_{i+1} \rightarrow v_{i+1}$, for all $i \in [2, m]$. Thus, there exists, the set $\bar{P} = P_5(2), P_5(3), \cdots, P_5(\frac{m-1}{2})$, if m is odd. Thus, there are $\frac{m-1}{2}$ number of P_5 -paths.

Now, by Theorem 2.1, $im(P_5) = 2$. Clearly, \bar{P} consists of all the edges in $E(G_{m,2})$ that can be in M . Therefore, $im(G_{m,2}) \leq 2 \binom{m-1}{2} = m - 1$. Suppose that m is even. Then, set $P* = \{P_5(2), P_5(3), \dots, P_5(\frac{m-2}{2}), P_3(t)\}$, where $P_3(t) = v_k \rightarrow u_k \rightarrow u_1$. So, $im(P* \setminus P_3(t)) = 2 \binom{m-2}{2} = m - 2$. By an earlier result, $im(P_3(t)) = 1$. Therefore, $im(P*) = m - 1$. Hence, for any integer m , $im(G_{m,2}) \leq m - 1$. Conversely, by definition of induced matching and stacked-book graph, $v_2u_2, v_3u_3, \dots, v_mu_m$, satisfying the distance conditions to belong to M . Thus, $im(G_{m,2}) \geq m - 1$ and hence the claim. \square

Next we consider the induced matching in $G_{m,3}$, where m is either even or odd and show that the graph contains the same induced matching as $G_{m,2}$.

Theorem 3.3. *Let $G_{m,3}$ be a stacked-book graph. Then $im(G_{m,3}) = m - 1$.*

To proof Theorem 3.3, we need two results, the first one, which is about about the nature of induced matching and distances between vertices of graphs, is more like a folklore because it follows from the definitions of induced matching of graphs.

Lemma 3.2. *Let e_1 be a member of M of a graph G . Then some edge $e_2 \in E(G)$ also belongs to M if there exists $v_1 \in e_1$ and $u_1 \in e_2$ such that $d(v_1, u_1) \geq 3$ and $v_2 \in e_1$ and $u_2 \in e_2$, such that $d(v_2, u_2) \geq 2$.*

Proof. The proof follows from the definition of induced matching M of graph G \square

Lemma 3.3. *Let $G_{m,3}$ be a stacked-book graph with factor star graphs $S_m(1)$, $S_m(2)$ and $S_m(3)$ such that $v_1 \rightarrow u_1 \rightarrow w_1$ is a P_3 path in $G_{m,3}$, where v_1, u_1 and w_1 are the central vertices of the respective factor star graphs. Then if u_1 is saturated, and $u_1v_k \in M$ for some $v_k \in V(G_{m,3})$, then $|M| = 1$ and thus, not the maximum induced matching of $G_{m,3}$.*

Proof. For $v_k \in V(G_{m,3})$, $v_k \neq u_1$, for which $v_i \in G_{m,3}$ such that $d(v_k, v_i) = 3$ since the $diam(G_{m,3}) = 3$. However, suppose that $v_iv_j \in E(G_{m,3})$, for which $d(v_k, v_i) = 3$. It is clear that v_i is a leaf if some $S_m(t)$, $t \in \{1, 3\}$. Thus, $d(u_1, v_j) = 1$, hence a contradiction to Lemma 3.2 and hence the result. \square

3.1. Proof of Theorem 3.3. Now we proceed to proof Theorem 3.3.

Proof. Suppose that $|M| > m - 1$. Let v_1, u_1 and w_1 be the central vertices of $S_m(1)$, $S_m(2)$ and $S_m(3)$ respectively. Clearly, $v_1u_1, u_1w_1 \notin M$ from Lemma 3.3. Now, first we show that v_1 is not saturable. Suppose that v_1 is saturable, then $v_1v_q \in M$, where v_q is a leaf on $S_m(1)$. By an earlier result, subgraph induced $S_m(1)$ and $S_m(2)$ does not contain another member of M . Also, let $v_qu_q \in E(G_{m,3})$, with $u_q \in S_m(2)$ and $u_qw_q \in E(G)$, with $w_q \in S_m(3)$. By earlier result, $u_qw_q \notin M$. In like manner, if w_1 is saturated, and $w_1w_q \in M$

no other edge in subgraph of $G_{m,3}$ induced by $S_m(2)$ and $S_m(3)$ is a member of M , and $v_q u_q \notin M$. Without loss of generality, suppose that $v_1, v_q \in M$, then only $\bar{M} = \{u_i w_i : i \in [2, m]; i \neq q\} \subset E(G_{m,3})$ will be member of M . Thus $|\bar{M}| = m - 2$ and so $|M| = m - 1$, which is a contradiction. Now it has been established that none of the pendants of $S_m(1), S_m(2)$ and $S_m(3)$ can be in M . Thus, the possible members of M are $\{v_i u_i : i \in [2, m]\} \cup \{u_i w_i : i \in [2, m]\} = M'$. Clearly, $|\bar{M}| = 2(m - 1)$. By Lemma 3.2, only half of the members of \bar{M} can be in M . Thus, $im(G_{m,3}) \leq m - 1$. Reasonably, $im(G_{m,2}) \leq im(G_{m,3})$. By earlier result, therefore, $iu(G_{m,3}) \geq m - 1$ and thus $im(G_{m,3}) = m - 1$. \square

Next we investigate the induced matching number of $G_{m,4}$. We start with a lemma that will be employed in the main result.

Lemma 3.4. *Let $G_{m,4}$ be a stacked-book graph such that $S_m(1), S_m(2), S_m(3)$ and $S_m(4)$ are the factor stars of $G_{m,4}$. Suppose that $im(G_{m,4}) \geq m$. Then if $M' = \{u_i w_i : i \in [2, m]; u_i \in S_m(2), w_i \in S_m(3)\}$, then M' is not a subset of M .*

Proof. It is easy to see that $|M'| = m - 1$. Now, suppose that $M' \subset M$, then u_i, w_i are saturated for all $i \in [2, m]$. Thus, no vertex $v_i \in S_m(1)$ and $r_i \in S_m(4)$ is saturable, for $i \in [2, m]$, which implies that $im(G_{m,4}) = m - 1$ and thus, a contradiction. \square

Next we consider the main theorem.

Theorem 3.4. *Let $S_m(1), S_m(2), S_m(3)$ and $S_m(4)$ be the factor star graphs of the stacked-book graph $G_{m,4}$. Then, $im(G_{m,4}) = m$.*

Proof. By Lemma 3.4, suppose that at least some edge in $M' = \{u_i w_i : i \in [2, m]; u_i \in S_m(2), w_i \in S_m(3)\}$ is not in M . Suppose therefore that $u_k w_k \in M$. Then for $v_k \in S_m(1)$, and $r_k \in S_m(4)$, $v_1 v_k, r_1 r_k \in M$, where v_1 and r_1 are the central vertices of $S_m(1)$ and $S_m(4)$ respectively. Thus, $im(G_{m,4}) \geq m$. Conversely, suppose that $im(G_{m,4}) = m + 1$. Now, let u_1, w_1 be the central vertices of $S_m(2)$ and $S_m(3)$ respectively. Suppose that one of u_i, w_i , say u_i is saturated such that $u_1 u_i \in M$. Then, from earlier result, no edge in the subgraph of $G_{m,4}$ induced by $S_m(1), S_m(2)$ and $S_m(3)$ is contained in M . Likewise, if $w_1 w_i \in M$, then all other vertices on the subgraph of $G_{m,4}$ induced by $S_m(2), S_m(3)$ and $S_m(4)$ are unsaturable. If any of the pendant of $S_m(2)$ and $S_m(3)$ is in M , then $M = 2$. Now, note as well that if $u_1 w_1 \in M$, then by the distances of u_1 and w_1 to the rest of vertices on $S_m(1), S_m(2), S_m(3)$ and $S_m(4)$, only $u_1 w_1$ will be in M . Thus for optimal M , some members of $M'' = \{v_i u_i; i \in [2, m]\}$ or $M''' = \{w_i r_i : i \in [2, m]\}$ will have to be in M .

Now clearly, it can be seen that $|M' \cup M''| = 2(m - 1)$ and only $m - 1$ members of $M' \cup M''$ can be in M . Based on this observable fact, at least there will exist a $w_i \in S_m(3)$ that is not saturable. Thus, there exist a saturable vertex

$r_i \in S_m(4)$, such that $r_1 r_i \in M$. By earlier result, there is no other pendant of $S_m(4)$ that is in M . Thus, $im(G_{m,4}) < m + 1$ and hence a contradiction. Therefore, $im(G_{m,4}) \leq m$ and the claim follows. \square

Now we consider the case of $G_{m,5}$. We shall need some new results to aid the proof.

Lemma 3.5. *Suppose that $w_1 \in S_m(3)$ is the central vertex of $S_m(3)$, where $\{S_m(i) : i \in [1, 5]\}$ is the set of factor stars of $G_{m,5}$. If w_1 is saturated, then for M of $G_{m,5}$, $|M| \leq 2m - 3$.*

Suppose that w_1 is the central vertex of $S_m(3)$ and it is saturated. Then one of the $w_1 w_k, u_1 w_1$ and $w_1 r_1$ belongs to M where u_1, r_1 are central vertices of $S_m(2)$ and $S_m(4)$ respectively. Suppose that $w_1 w_k \in M$, where $k \leq m$. Now for all $i \in [2, m], i \neq k$, $w_i \in S_m(3)$ is unsaturable by earlier results. Thus members of $\{u_i w_i : i \in [2, m]\}$ and $\{w_i r_i : r_i \in S_m(4), i \in [2, m]\}$ do not belong in M . Also it clear to see that both edges $v_k u_k, r_k t_k \notin M$, where $t_k \in S_m(5)$. Now, from earlier technique, it can be deduced that $v_1 v_i, t_1 t_i \notin M$ for all $i \in [2, m]$. Thus, only $E' = \{v_i u_i : i \in [2, m], i \neq k\}$ and $E'' = \{r_i t_i : i \in [2, m], i \neq k\}$ can be in M . Clearly, $|E' \cup E''| = 2(m - 2)$. Thus $|M| = 2m - 3$. Also, if $u_1 w_i \in M$, it can be seen by following the definitions of induced matching that no other edges in the subgraph of $G_{m,5}$ induced by $S_m(1), S_m(2)$ and $S_m(3)$ is a member of M and from earlier results, only $m - 1$ edges of the subgraph of $G_{m,5}$ induced by $S_m(3), S_m(4)$ and $S_m(5)$ can be in M . Thus, M consists of at most m edges, which is not more than $2m - 3$, since $m \geq 3$.

Lemma 3.6. *Suppose that $im(G_{m,5}) \geq 2(m - 1)$. Then u_1, w_1 and r_1 , the central vertices of $S_m(2), S_m(3)$ and $S_m(4)$ respectively are unsaturated.*

Proof. Proof follows from last theorem and an earlier result. \square

Now we proceed to the probe the induced matching of $G_{m,5}$.

Theorem 3.5. *Let $G_{m,5}$ be a stacked-book graph. Then, $im(G_{m,5}) = 2(m - 1)$.*

Proof. From the last results, we see that if u_1, w_1, r_1 are unsaturated, then $|M| \geq 2m - 3$. Now we show that $im(G_{m,5}) \geq 2(m - 1)$. Note that there exists a path $P_5(i) = v_i \rightarrow u_i \rightarrow w_i \rightarrow r_i \rightarrow t_i$, for all $i \in [2, m]$. Therefore, there are $m - 1$ such paths in $G_{m,5}$. From earlier results, $im(P_5) = 2$. Thus, $im(G_{m,5}) \geq 2(m - 1)$. Conversely, u_1, w_1, r_1 are established not to be saturated for the claim to hold. The edges in $E(G_{m,5})$ left to be members of M the pendants of $S_m(1)$ and $S_m(5)$ and the paths $P_5(i)$ defined earlier. Suppose that a pendant each from $S_m(1)$ and $S_m(5)$ belong to M , then by definition of induced matching, at most one edge on each of the paths $P_5(i)$ can be a member of M . Thus $|M| = m + 1$. The only alternative is if no pendant of $S_m(1)$ and

$S_m(5)$ is a member of M . Thus, at most two edges on each member of $P_5(i)$ will be in M . Thus, $|M| \leq 2(m - 1)$ and so, $im(G_{m,5}) = 2(m - 1)$. \square

Now we generalize the results.

Theorem 3.6. *Let $G_{m,n}$ be a stacked-book graph such that n is even. Then*

$$im(G_{m,n}) \geq \begin{cases} m\lfloor \frac{n}{4} \rfloor - 1 & \text{if } n \equiv 2 \pmod{4}; \\ \frac{mn}{4} & \text{if } n \equiv 0 \pmod{4}. \end{cases}$$

Proof. the claims follow from combining the results earlier proved where n are even numbers. \square

Theorem 3.7. *Let $G_{m,n}$ be stacked-book graph with n odd. Then*

$$im(G_{m,n}) \geq \begin{cases} m\lfloor \frac{n}{4} \rfloor + 2 & \text{if } n \equiv 3 \pmod{4}; \\ \frac{mn+3m-8}{4} & \text{if } n \equiv 1 \pmod{4}. \end{cases}$$

We have established the lower bound for the the MIM numbers for the stacked-book graphs. From our preliminary work into establishing the tighter bounds, we have reasons to suggest that the results in the last two theorems may coincide with the upper bounds, and thus we come up with the conjectures below.

Conjecture 3.1. *Let $G_{m,n}$ be a stacked-book graph such that n is even. Then*

$$im(G_{m,n}) = \begin{cases} m\lfloor \frac{n}{4} \rfloor - 1 & \text{if } n \equiv 2 \pmod{4}; \\ \frac{mn}{4} & \text{if } n \equiv 0 \pmod{4}. \end{cases}$$

Conjecture 3.2. *Let $G_{m,n}$ be stacked-book graph with n odd. Then*

$$im(G_{m,n}) = \begin{cases} m\lfloor \frac{n}{4} \rfloor + 2 & \text{if } n \equiv 3 \pmod{4}; \\ \frac{mn+3m-8}{4} & \text{if } n \equiv 1 \pmod{4}. \end{cases}$$

4. CONCLUSION

We have obtained the MIM number of stacked-book graphs $G_{m,n}$ for all m and for $n \in [1, 5]$. These results are building blocks for obtaining the lower bounds for the cases where $n \geq 6$. The conjecture at the end of the work suggests that the lower bounds obtained in this work will in fact be equal to the upper bounds if those can be found. It must be noted that finding the lower bounds or the MIM numbers for the complete stacked-book graphs class will take rigorous effort and therefore may worth considering as a new task.

REFERENCES

- [1] Adefokun, T.C., Ajayi, D.O *On maximum induced matching number of special grids*, Journal of Mathematics and Applications. 41 (2018) 5-18.
- [2] Ajayi, D.O., Adefokun, T.C. *Some bounds on the maximum induced matching numbers of certain grids*, Acta Universitatis Matthiae Belii, Series Mathematics. 25 (2017) 63-71.

- [3] Cameron, K. *Induced matching in intersection graphs*, Discrete Mathematics. 278 (1)(3) (2004) 1-9.
- [4] Cameron, K. *Induced matching*, Discrete Applied Mathematics. 24 (1989) 97-102.
- [5] Golumbic, M.C., Laskar, R. *New results on induced matching*, Discrete Applied Mathematics. 44 (1993) 79-89.
- [6] Golumbic, M.C., Lewenstein, M. *New results on induced matching*, Discrete Applied Mathematics. 101 (2000) 157-165.
- [7] Marinescu-Ghemaci, R. *Maximum induced matching on grids*, Optimization Theory, Decision Making and Operation Research Applications, Springer New York. (2013) 177-187.
- [8] Stockmeyer, L.J., Vazirani, V.V. *NP-Completeness of some generalization of the maximum matching problem*, Inf. Process. Lett. 15(1) (1982) 14-19.
- [9] Xia, M., Tan, H. *Exact algorithm for maximum induced matching*, Information and Computation. 256 (2017) 196-211.

¹DEPARTMENT OF COMPUTER AND MATHEMATICAL SCIENCES,
CRAWFORD UNIVERSITY,
NIGERIA
Email address: tayoadefokun@crawforduniversity.edu.ng