

On the Equivalence, Stabilisers, and Feet of Buekenhout-Tits Unitals

Jake Faulkner

Geertrui Van de Voorde*

Abstract

This paper addresses a number of problems concerning Buekenhout-Tits unitals in $\text{PG}(2, q^2)$, where $q = 2^{2e+1}$ and $e \geq 1$. We show that all Buekenhout-Tits unitals are PGL -equivalent (addressing an open problem in [4]), explicitly describe their $\text{P}\Gamma\text{L}$ -stabiliser (expanding Ebert's work in [8]), and show that lines meet the feet of points not on ℓ_∞ in at most four points. Finally, we show that feet of points not on ℓ_∞ are not always a $\{0, 1, 2, 4\}$ -set, in contrast to what happens for Buekenhout-Metz unitals [1].

MSC2020: 51E20

Keywords: Unital, Tits ovoid, Buekenhout-Tits unital, feet

1 Introduction

1.1 Background

Let $\text{PG}(2, q^2)$ denote the Desarguesian projective plane over the finite field with q^2 elements, \mathbb{F}_{q^2} , where q is a prime power. A *unital* U in $\text{PG}(2, q^2)$ is a set of $q^3 + 1$ points such that every line of $\text{PG}(2, q^2)$ meets U in 1 or $q + 1$ points. The *classical* or *Hermitian* unital, usually denoted by $\mathcal{H}(2, q^2)$, arises by taking the absolute points of a non-degenerate Hermitian polarity. Each point P not lying on a unital U , lies on $q + 1$ tangent lines to U ; the $q + 1$ points of U whose tangent lines contain P are called the *feet* of P , and are denoted by $\tau_P(U)$.

It is well-known that $\text{PG}(2, q^2)$ can be modelled using a Desarguesian line spread of $\text{PG}(3, q)$ embedded in $\text{PG}(4, q)$ via the *André/Bruck-Bose (ABB)* construction. A wide class of unitals in $\text{PG}(2, q^2)$, called *Buekenhout unitals*, arise as follows from the ABB construction; starting in $\text{PG}(4, q)$ fixing a hyperplane Σ , and a Desarguesian spread of Σ , we take any ovoidal cone \mathcal{C} such that $\mathcal{C} \cap \Sigma$ is a spread line of Σ . Then in $\text{PG}(2, q^2)$, \mathcal{C} gives rise to a unital U . If the base of \mathcal{C} is an elliptic quadric, the unital is called a *Buekenhout-Metz unital*. The family of Buekenhout-Metz unitals contains the Hermitian unitals, but there are many non-equivalent Buekenhout-Metz unitals (see [3], [9]). If

*This author is supported by the Marsden Fund Council administered by the Royal Society of New Zealand.

$q = 2^{2e+1}$, $e \geq 1$, and the base of \mathcal{C} is a Tits ovoid, the unital is called a *Buekenhout-Tits unital*. For more information on unitals and their constructions, see [4].

Unitals can be characterised by the combinatorial properties of their feet. It is easy to see that for the classical unital $\mathcal{H}(2, q^2)$, the feet of a point not on the unital are always collinear. Thas [12] showed the converse, namely, that a unital U is classical if and only if for all points, not on U , the feet are collinear. This was improved by Aguglia and Ebert [2] who showed that a unital U is classical if and only if there exist two tangent lines ℓ_1, ℓ_2 such that for all points $P \in (\ell_1 \cup \ell_2) \setminus U$ the feet of P are collinear. It is known (see e.g. [4]) that if U is a non-classical Buekenhout-Metz unital, the feet of a point $P \notin U$ are collinear if and only if $P \in \ell_\infty$. Furthermore, it is shown in [1] that if U is Buekenhout-Metz unital, a line meets the feet of a point $P \notin \ell_\infty$ in either 0, 1, 2, or 4 points. Ebert [8] showed for a Buekenhout-Tits unital, the feet of $P \notin U$ are collinear if and only if $P \in \ell_\infty$. It is then natural to ask how a line may meet the feet of a point $P \notin \ell_\infty$ for Buekenhout-Tits unitals. We will answer this question in Theorem 3.

Many characterisations of unitals make use of their stabilisers in PGL, resp. PTL. In [7] it is shown that a unital is classical if its stabiliser contains a cyclic group of order $q^2 - q + 1$. Several other characterisations of unitals by their stabiliser group are listed in [4]. In [8], Ebert determined the stabiliser of a Buekenhout-Tits unital in $\text{PGL}(3, q^2)$ (see Result 1). We will extend this work in this paper.

1.2 Summary of this paper

In this paper we present three main results:

1. We show that all Buekenhout-Tits unitals are projectively equivalent (see Theorem 1). This addresses an open problem in [4], and is alluded to in [10] (see Remark 1).
2. A description of the full stabiliser group of a Buekenhout-Tits unital in $\text{PFL}(3, q^2)$ (see Theorem 2). Ebert [8] only provides a description of stabiliser of the Buekenhout-Tits unital in PGL (Result 1). The stabiliser of the classical unital in $\text{PFL}(3, q^2)$ is $\text{PFU}(3, q^2)$, and the stabiliser of the Buekenhout-Metz unital in $\text{PFL}(3, q^2)$ is described in [9] for q even and [3] for q odd.
3. If U is a Buekenhout-Tits unital, then a line ℓ meets the feet of a point $P \notin (\ell_\infty \cup U)$ in at most 4 points. Moreover, there exists a point P and line ℓ such that the feet of P meet ℓ in exactly three points (see Theorem 3). This highlights a difference between Buekenhout-Metz unitals and Buekenhout-Tits unitals. It also solves an open problem posed by Aguglia and Ebert [2] and later listed in [4].

1.3 Coordinates for a Buekenhout-Tits unital

In [8], Ebert derives coordinates for a Buekenhout-Tits unital \mathcal{U}_{BT} in $\text{PG}(2, q^2)$, $q = 2^{2e+1}$. Pick $\epsilon \in \mathbb{F}_{q^2}$ such that $\epsilon^q = \epsilon + 1$, and $\epsilon^2 = \epsilon + \delta$ for some $1 \neq \delta \in \mathbb{F}_q$ with absolute trace equal to one. Then the

following set of points in $\text{PG}(2, q^2)$ is a Buekenhout-Tits unital,

$$\mathcal{U}_{\text{BT}} = \{(0, 0, 1)\} \cup \{P_{r,s,t} = (1, s + t\epsilon, r + (s^{\sigma+2} + t^\sigma + st)\epsilon) \mid r, s, t \in \mathbb{F}_q\}, \quad (1)$$

where $\sigma = 2^{e+1}$ has the property that σ^2 induces the automorphism $x \mapsto x^2$ of \mathbb{F}_{q^2} . In addition, it can be verified that $\sigma + 1$, $\sigma + 2$, $\sigma - 1$, and $\sigma - 2$ all induce permutations of \mathbb{F}_q with inverses induced by $\sigma - 1$, $1 - \sigma/2$, $\sigma + 1$ and $-(\sigma/2 + 1)$ respectively.

The following theorem describes the group of projectivities (that is, elements of $\text{PGL}(3, q^2)$) stabilising \mathcal{U}_{BT} .

Result 1. [8, Theorem 4] *Let $G = \text{PGL}(3, q^2)_{\mathcal{U}_{\text{BT}}}$, $q = 2^{2e+1}$, be the group of projectivities stabilising the Buekenhout-Tits unital \mathcal{U}_{BT} . Then G is an abelian group of order q^2 , consisting of the projectivities induced by the matrices*

$$M_{u,v} = \left\{ \left[\begin{array}{ccc} 1 & u\epsilon & v + u^\sigma\epsilon \\ 0 & 1 & u + u\epsilon \\ 0 & 0 & 1 \end{array} \right] \mid u, v \in \mathbb{F}_q \right\}, \quad (2)$$

where $\sigma = 2^{e+1}$ and matrices act on the homogeneous coordinates of points by multiplication from the right.

2 On the Projective Equivalence of Buekenhout-Tits Unitals

In this section, we show that all Buekenhout-Tits unitals are projectively equivalent to the unital \mathcal{U}_{BT} given in equation (1).

Remark 1. The authors of [10] give this result without proof and state it can be derived using the same techniques employed by Ebert in [8]. Ebert however, lists the equivalence of Buekenhout-Tits unital as an open problem in [4] which appeared about ten years after his original paper [8].

It is easy to see that the Buekenhout-Tits unital \mathcal{U}_{BT} is tangent to the line $\ell_\infty : x = 0$ in the point $P_\infty = (0, 0, 1)$. From the ABB construction it follows that P_∞ has the following property with respect to \mathcal{U}_{BT} .

Property 1. Given any unital U , a point $P \in U$ has Property 1 if all secant lines through P meet U in Baer sublines.

It is shown in [5] that if two different points of U have Property 1, then U is classical. Hence, the point P_∞ is the unique point of \mathcal{U}_{BT} admitting this property. We will count all Buekenhout-Tits unitals tangent to ℓ_∞ at a point P_∞ having Property 1.

Lemma 1. *There are $q^4(q^2 - 1)^2$ unitals projectively equivalent to \mathcal{U}_{BT} in $\text{PG}(2, q^2)$ tangent to $\ell_\infty : x = 0$, and containing the point $P_\infty = (0, 0, 1)$ with Property 1.*

Proof. First note that any projectivity mapping \mathcal{U}_{BT} to a unital tangent to ℓ_∞ in P_∞ necessarily is contained in the group H of projectivities fixing ℓ_∞ line-wise and P_∞ point-wise. The elements of H are induced by all matrices of the following form,

$$\begin{bmatrix} 1 & x_{12} & x_{13} \\ 0 & x_{22} & x_{23} \\ 0 & 0 & x_{33} \end{bmatrix},$$

where $x_{22}x_{33} \neq 0$ and matrices act on homogeneous coordinates by multiplication on the right. It follows that $|H| = (q^2 - 1)^2 q^6$. Furthermore, from the description of $G = \text{PGL}(3, q^2)_{\mathcal{U}_{\text{BT}}}$ in Result 1, we know that $H_{\mathcal{U}_{\text{BT}}} = G$, and hence, $H_{\mathcal{U}_{\text{BT}}}$ has order q^2 . By the orbit-stabiliser theorem, we find that there are $(q^2 - 1)^2 q^4$ unitals in the orbit of \mathcal{U}_{BT} under H . \square

Consider $\text{PG}(2, q^2)$ modelled using the ABB construction with fixed hyperplane H_∞ . Let p_∞ be the spread line corresponding to P_∞ . Then any Buekenhout-Tits unital U tangent to ℓ_∞ at P_∞ with Property 1 corresponds uniquely to an ovoidal cone \mathcal{C} meeting H_∞ at p_∞ .

Lemma 2. *There are $q^4(q^2 - 1)^2$ ovoidal cones \mathcal{C} in $\text{PG}(4, q)$ with base a Tits ovoid, such that \mathcal{C} meets H_∞ in the spread element p_∞ .*

Proof. Let V be a point on the line p_∞ , and $H \neq H_\infty$ a hyperplane not containing V . Then, H meets H_∞ in a plane containing a point $R \in p_\infty \setminus \{V\}$. Any ovoidal cone \mathcal{C} with vertex V and base a Tits ovoid, such that \mathcal{C} meets H_∞ precisely in p_∞ , meets H in a Tits ovoid tangent to $H \cap H_\infty$ at the point R . We will count all cones of this form, for all $V \in p_\infty$.

Consider the pairs of planes Π and Tits ovoids \mathcal{O} , (Π, \mathcal{O}) , where $\Pi, \mathcal{O} \subset H$ and Π is tangent to \mathcal{O} . On the one hand, there are $|\text{PGL}(4, q)|/|\mathcal{O}_{\text{PGL}(4, q)}| = (q + 1)^2 q^4 (q - 1)^2 (q^2 + q + 1)$ Tits ovoids in $\text{PG}(3, q)$, and each has $q^2 + 1$ tangent planes. On the other hand, $\text{PGL}(4, q)$ is transitive on hyperplanes of $\text{PG}(3, q)$, so each plane is tangent to the same number of Tits ovoids. It thus follows, that there are

$$\frac{(q + 1)^2 q^4 (q - 1)^2 (q^2 + q + 1)(q^2 + 1)}{q^4 + q^3 + q^2 + q + 1} = (q - 1)^2 q^4 (q + 1)(q^2 + q + 1)$$

Tits ovoids tangent to $H \cap H_\infty$ contained in H .

Furthermore, since $\text{PGL}(4, q)_{H \cap H_\infty}$ is transitive on points of $H \cap H_\infty$, each point of $H \cap H_\infty$ is contained in the same number of Tits ovoids \mathcal{O} , so it follows that the number of Tits ovoids tangent to $H \cap H_\infty$ at $R = p_\infty \cap H$ is $(q - 1)^2 q^4 (q + 1)$. Hence, there is an equal number of ovoidal cones with base a Tits ovoid, vertex V , and meeting H_∞ at p_∞ . As the choice of V was arbitrary, and there are $q + 1$ points on p_∞ , there are $(q^2 - 1)^2 q^4$ ovoidal cones with base a Tits ovoid, and meeting H_∞ at p_∞ . \square

Theorem 1. *All Buekenhout-Tits unitals in $\text{PG}(2, q^2)$ are PGL-equivalent.*

Proof. From Lemmas 1 and 2, we see that the number of ovoidal cones with vertex a Tits ovoid, tangent to H_∞ at p_∞ is equal to the number of Buekenhout-Tits unitals that are PGL equivalent to \mathcal{U}_{BT} and tangent to l_∞ at P_∞ with Property 1. The result follows. \square

Corollary 1. *Let U be a Buekenhout-Tits unital, then the projectivity group stabilising U is isomorphic to the group G in Theorem 1.*

In showing that all Buekenhout-Tits unitals are projectively equivalent, we may use \mathcal{U}_{BT} to verify statements about general Buekenhout-Tits unitals.

3 On the Stabiliser of the Buekenhout-Tits Unital

We now describe the stabiliser of the Buekenhout-Tits unital \mathcal{U}_{BT} in $\text{PFL}(3, q^2)$.

Lemma 3. *Let $M_{u,v}, M_{s,t}$ be matrices inducing collineations of G as defined in Result 1, then $M_{u,v}M_{s,t} = M_{u+s, t+v+su\delta}$.*

Proof. Using equation (2), we find

$$M_{u,v}M_{s,t} = \begin{bmatrix} 1 & (s+u)\epsilon & (t+v+su\delta) + (s+u)\sigma \\ 0 & 1 & (u+s) + (u+s)\epsilon \\ 0 & 0 & 1 \end{bmatrix}. \quad (3)$$

Thus, we have $M_{u,v}M_{s,t} = M_{u+s, t+v+su\delta}$. \square

Corollary 2. *The order of any collineation of G induced by a matrix $M_{u,v}$ as defined in Result 1 is four if and only if $u \neq 0$, and two if and only if $u = 0$ and $v \neq 0$.*

Proof. Firstly note that $M_{0,0} = I$. Direct calculation shows that $M_{u,v}^2 = M_{0, u^2\delta}$, $M_{u,v}^3 = M_{u, v+u^2\delta}$ and $M_{u,v}^4 = M_{0,0}$. \square

Corollary 3. *The stabiliser group G as defined in Result 1 is isomorphic to $(C_4)^{2e+1}$.*

Proof. Recall from Result 1 that $|G| = q^2 = 2^{4e+2}$. From Corollary 2, we have that $G \cong (C_4)^k (C_2)^l$ for some integers k, l such that $2^{2k+l} = |G| = 2^{4e+2}$, and hence,

$$l = 2(e+1-k). \quad (4)$$

Furthermore, we see that the number of elements of order four in G is $q^2 - q$ as they correspond to all matrices $M_{u,v}$ with $u, v \in \mathbb{F}_q$ and $u \neq 0$. The number of elements of order four in a group isomorphic to $(C_4)^k (C_2)^l$ is $(4^k - 2^k)2^l$. Thus,

$$(4^k - 2^k)2^l = 4^{2e+1} - 2^{2e+1}. \quad (5)$$

Using (4), we find that $k = 2e + 1$, and therefore $G \cong (C_4)^{2e+1}$. \square

Theorem 2. *Let U be a Buekenhout-Tits unital in $\text{PG}(2, q^2)$, $q = 2^{2e+1}$, then the stabiliser group of U in $\text{P}\Gamma\text{L}(3, q^2)$ is the order $q^2(4e + 2)$ group GK , where K is a cyclic subgroup of order $16e + 8$ generated by*

$$\psi : \mathbf{x} \mapsto \mathbf{x}^2 \begin{bmatrix} 1 & 1 & \epsilon \\ 0 & \delta^{\sigma/2}(1 + \epsilon) & \delta^{\sigma/2}(1 + \epsilon) \\ 0 & 0 & \delta^{\sigma+1} \end{bmatrix}. \quad (6)$$

(Here, \mathbf{x} denotes the row vector containing the three homogeneous coordinates of a point, and \mathbf{x}^2 denotes its elementwise power.)

Proof. From Lemma 2, we have that the number of Buekenhout-Tits unitals is $q^4(q^2 - 1)^2$. Since all of those unitals are PGL -equivalent by Theorem 1, and $\text{PGL}(3, q^2) \triangleleft \text{P}\Gamma\text{L}(3, q^2)$, we have that

$$q^4(q^2 - 1)^2 = \frac{|\text{PGL}(3, q^2)|}{|\text{PGL}(3, q^2)_U|} = \frac{|\text{P}\Gamma\text{L}(3, q^2)|}{|\text{P}\Gamma\text{L}(3, q^2)_U|}. \quad (7)$$

So $\text{P}\Gamma\text{L}(3, q^2)_U$ must have order $q^2(4e + 2)$. Direct calculation shows that ψ stabilises \mathcal{U}_{BT} . We have $\psi^{4e+2} \in G$ as $\mathbf{x}^{2^{4e+2}} = \mathbf{x}^{q^2} = \mathbf{x}$. Hence, $|\psi| = (4e + 2)|\psi^{4e+2}|$. From Corollary 2, it follows that $|\psi^{4e+2}| \in \{1, 2, 4\}$, with $|\psi^{4e+2}| = 4$ if and only if ψ^{4e+2} is induced by $M_{u,v}$ for some $u \neq 0$. Hence, $|\psi^{4e+2}| = 4$ if and only if $\psi^{4e+2}(0, 1, 0) \neq (0, 1, 0)$ as $(0, 1, 0)M_{u,v} = (0, 1, u + u\epsilon)$. Consider the point $(0, 1, z)$ for some arbitrary $z \in \mathbb{F}_q$. Direct calculation shows that $\psi(0, 1, z) = (0, 1, 1 + \mu z^2)$, where $\mu = \frac{\delta^{\sigma+1}}{\delta^{\sigma/2}(1+\epsilon)} = \delta^{\sigma/2}\epsilon$. Thus,

$$\psi^k(0, 1, z) = (0, 1, \sum_{i=0}^k \mu^{2^i-1} + zg(z)) \quad (8)$$

for some polynomial $g(z)$ depending on k . If $z = 0$ and $k = 4e + 2$ we thus find:

$$\psi^{4e+2}(0, 1, 0) = (0, 1, \sum_{i=0}^{4e+2} \mu^{2^i-1}) \quad (9)$$

$$= (0, 1, \frac{\text{Tr}(\mu)}{\mu}). \quad (10)$$

Recall that $\epsilon^q = \epsilon + 1$, so $\text{Tr}_{\mathbb{F}_{q^2}/\mathbb{F}_q}(\epsilon) = 1$. We have that, $\text{Tr}_{\mathbb{F}_{q^2}/\mathbb{F}_2}(\delta^{\sigma/2}\epsilon) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(\text{Tr}_{\mathbb{F}_{q^2}/\mathbb{F}_q}(\delta^{\sigma/2}\epsilon)) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(\delta^{\sigma/2} \text{Tr}_{\mathbb{F}_{q^2}/\mathbb{F}_q}(\epsilon)) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(\delta^{\sigma/2}) = 1$. Hence, $\psi((0, 1, 0)) \neq (0, 1, 0)$, so $|\psi^{4e+2}| = 4$ and $|\psi| = 16e + 8$. Let $K = \langle \psi \rangle$, because $|K \cap G| = 4$, it follows that $|GK| = q^2(4e + 2)$ and thus $GK = \text{P}\Gamma\text{L}(3, q^2)_U$. \square

4 On the Feet of the Buekenhout-Tits Unital

The feet of the Buekenhout-Tits unital \mathcal{U}_{BT} are first described by Ebert in [8]. He shows that the feet of a point $P = (1, y_1 + y_2\epsilon, z_1 + z_2\epsilon)$ is the following set of points:

$$\begin{aligned} \tau_P(\mathcal{U}_{\text{BT}}) = \{ & (1, s + t\epsilon, s^2 + t^2\delta + st + y_1s + y_1t + y_2\delta t + z_1 + (s^{\sigma+2} + t^\sigma + st)\epsilon) \\ & | s, t \in \mathbb{F}_q, s^{\sigma+2} + t^\sigma + st = y_2s + y_1t + z_2\}. \end{aligned} \quad (11)$$

If the line ℓ has equation $\alpha x + y = 0$, where $\alpha \in \mathbb{F}_{q^2}$, Ebert shows that $|\ell \cap \tau_P(\mathcal{U}_{\text{BT}})| \leq 1$. Otherwise, ℓ has equation $(a_1 + a_2\epsilon)x + (b_1 + b_2\epsilon)y + z = 0$, with $a_1, a_2, b_1, b_2 \in \mathbb{F}_q$, and Ebert shows that ℓ meets $\tau_P(\mathcal{U}_{\text{BT}})$ in the points $P_{r,s,t} \in \mathcal{U}_{\text{BT}}$, where $r, s, t \in \mathbb{F}_q$ satisfy

$$s^2 + \delta t^2 + st + (y_1 + b_1)s + (y_1 + y_2\delta + b_2\delta)t + z_1 + a_1 = 0, \quad (12)$$

$$s^{\sigma+2} + t^\sigma + st = b_2s + (b_1 + b_2)t + a_2, \quad (13)$$

$$y_2s + y_1t + z_2 = b_2s + (b_1 + b_2)t + a_2. \quad (14)$$

We will show that for all choices of points $P \notin \ell_\infty$ and lines ℓ , $|\tau_P(\mathcal{U}_{\text{BT}}) \cap \ell| \leq 4$.

Lemma 4. *Let G be the group of projectivities stabilising the Buekenhout-Tits unital as described in Result 1. Then, the set of $q^2 - q$ points $\{P_{a,b} = (1, a, b\epsilon) \mid a, b \in \mathbb{F}_q, b \neq a^{\sigma+2}\}$ are points from $q^2 - q$ distinct point orbits of order q^2 under G .*

Proof. Suppose there exists a collineation of G induced by a matrix $M_{u,v}$ such that $P_{a,b}M_{u,v} = P_{c,d}$. Then,

$$(1, a, b\epsilon) \begin{bmatrix} 1 & u\epsilon & v + u^\sigma\epsilon \\ 0 & 1 & u + u\epsilon \\ 0 & 0 & 1 \end{bmatrix} = (1, c, d\epsilon).$$

However, it is clear that $P_{a,b}M_{u,v} = (1, a + u\epsilon, v + u^\sigma\epsilon + a(u + u\epsilon) + b\epsilon)$, so $a + u\epsilon = c$. Therefore, $a = c$ and $u = 0$. If $u = 0$, then $v + b\epsilon = d\epsilon$, and we have $b = d$. Hence, $P_{a,b} = P_{c,d}$ and the lemma follows. \square

There are $q^4 - q^3 = q^2(q^2 - q)$ points of $\text{PG}(2, q)$ not on ℓ_∞ or \mathcal{U}_{BT} . By Lemma 4, each of these points lies in the orbit of a point of the form $(1, a, b\epsilon)$. Therefore, in order to study the feet of a point P , we may assume that the point $P = (1, y_1 + y_2\epsilon, z_1 + z_2\epsilon)$ has $y_2 = z_1 = 0$.

The following lemma shows that the feet of a point $P = (1, y_1, z_2\epsilon)$ meets almost all lines in at most 2 points.

Lemma 5. *Let $\ell : \alpha x + \beta y + z$ be a line in $\text{PG}(2, q^2)$, where $\alpha = a_1 + a_2\epsilon$, $\beta = b_1 + b_2\epsilon$ and $a_1, a_2, b_1, b_2 \in \mathbb{F}_q$. Let $P = (1, y_1, z_2\epsilon)$, with $y_1, z_2 \in \mathbb{F}_q$ such that $z_2 \neq y_1$. Unless $b_2 = 0$, $y_1 = b_1$ and $a_2 = z_2$, we have $|\tau_P(\mathcal{U}_{\text{BT}}) \cap \ell| \leq 2$.*

Proof. From the description given in (11), we see that the points $P_{r,s,t} \in \tau_P(\mathcal{U}_{\text{BT}})$ satisfy $s^{\sigma+2} + t^\sigma + st = y_1t + z_2$, and this equation has $q + 1$ solutions. Substituting this into equation (13), the points $P_{r,s,t} \in \tau_P(\mathcal{U}_{\text{BT}}) \cap \ell$ have s, t satisfying

$$s^2 + \delta t^2 + st + (y_1 + b_1)s + (y_1 + b_2\delta)t + a_1 = 0 \quad (15)$$

$$b_2s + (y_1 + b_1 + b_2)t + a_2 + z_2 = 0 \quad (16)$$

$$s^{\sigma+2} + t^\sigma + st + y_1t + z_2 = 0. \quad (17)$$

Recall that the points $(1, s, t, s^{\sigma+2} + t^\sigma + st)$, where $s, t \in \mathbb{F}_q$ are the q^2 affine points of a Tits ovoid. Hence, (17) represents an affine section of a Tits ovoid. Since it has $q+1$ points, it is an oval projectively equivalent to the translation oval $\mathcal{D}_\sigma = \{(1, t, t^\sigma) \mid t \in \mathbb{F}_q\}$. Unless $b_2 = 0$ and $y_1 = b_1$, equation (16) represents a line in $\text{AG}(2, q)$ which meets the oval (17) in at most two points, so we have at most two solutions to the system. If $b_2 = 0$, $y_1 = b_1$, and $a_2 \neq z_2$, then equation (16) has no solutions. \square

Remark 2. Lemma 5 is a refinement of [4, Theorem 4.33], where Barwick and Ebert rework Ebert's earlier proof in [8] that the feet of a point $P \notin (\ell_\infty \cup \mathcal{U}_{\text{BT}})$ are not collinear. This reworked proof asserts that the feet cannot be collinear because the line given by equation (16) and the conic from equation (15) cannot have $q+1$ common solutions. However, we can see that this logic is not complete, and leaves an interesting case to examine when equation (16) vanishes. Ebert's original proof in [8] does not contain this error, instead arguing that equations (15) and (17) cannot have $q+1$ common solutions.

It follows from Lemma 5 that the feet of a point $P \notin (\ell_\infty \cup \mathcal{U}_{\text{BT}})$ is a set of $q+1$ points such that every line meets $\tau_P(\mathcal{U}_{\text{BT}})$ in at most two points except for a set of q concurrent lines.

To this end, assume that $b_2 = 0$, $y_1 = b_1$ and $a_2 = z_2$. In this case, equation (16) vanishes. The system describing $\ell \cap \tau_P(\mathcal{U}_{\text{BT}})$ is thus

$$s^2 + \delta t^2 + st = y_1 t + a_1 \tag{18}$$

$$s^{\sigma+2} + t^\sigma + st = y_1 t + z_2. \tag{19}$$

The lines that produce these cases are the lines with dual coordinates $[a_1 + z_2 \epsilon, y_1, 1]$. These lines are concurrent at the point $(0, 1, y_1)$ which lies on ℓ_∞ . We will show in Corollary 4 that these latter lines meet $\tau_P(\mathcal{U}_{\text{BT}})$ in at most four points.

We require the following lemma, which adapts arguments found in [6, Lemma 2.1].

Lemma 6. *Let \mathcal{O} be a translation oval in $\text{PG}(2, q)$ projectively equivalent to \mathcal{D}_σ , and let \mathcal{C} be a non-degenerate conic. If the nucleus of \mathcal{O} is also the nucleus of \mathcal{C} , then $|\mathcal{O} \cap \mathcal{C}| \leq 4$.*

Proof. Without loss of generality we may take $\mathcal{O} = \mathcal{D}_\sigma$, so that the nucleus of \mathcal{O} is $N = (0, 1, 0)$. If N is also the nucleus of \mathcal{C} , then \mathcal{C} is a conic of the following form,

$$a_1 x^2 + a_2 y^2 + a_3 z^2 + xz = 0, \tag{20}$$

for some $a_1, a_2, a_3 \in \mathbb{F}_q$ with $a_2 \neq 0$. Suppose that $(0, 0, 1) \notin \mathcal{C}$. Then $a_3 \neq 0$, and the point $(1, t, t^\sigma) \in \mathcal{O}$ if and only if t satisfies

$$a_1 + a_2 t^2 + a_3 t^{2\sigma} + t^\sigma = 0, \tag{21}$$

that is

$$0 = (a_1 + a_2 t^2 + a_3 t^{2\sigma} + t^\sigma)^{\sigma/2} = a_1^{\sigma/2} + a_2^{\sigma/2} t^\sigma + t^2 + t. \tag{22}$$

Therefore,

$$t^\sigma = \left(\frac{a_3}{a_2}\right)^{2^e} t^2 + \frac{1}{a_2^{2^e}} t + \left(\frac{a_1}{a_2}\right)^{2^e} \quad (23)$$

and substituting into equation (21), we find that this equation has at most four solutions. If instead $(0, 0, 1) \in \mathcal{C}$, then $a_3 = 0$ and arguing as above we find that equation (21) has at most two solutions, so $|\mathcal{O} \cap \mathcal{C}| \leq 3$. \square

Corollary 4. *The feet of a point $P \notin (\ell_\infty \cup \mathcal{U}_{\text{BT}})$ meet a line ℓ in at most four points.*

Proof. From Lemma 5, we know we can restrict ourselves to the case $b_2 = 0, y_1 = b_1, a_2 = z_2$ which means we are looking at the points $P_{r,s,t} \in \tau_P(\mathcal{U}_{\text{BT}}) \cap \ell$ have s, t satisfying

$$s^2 + \delta t^2 + st = y_1 t + a_1 \quad (24)$$

$$s^{\sigma+2} + t^\sigma + st = y_1 t + z_2, \quad (25)$$

where equation (24) represents a conic \mathcal{C} , and equation (25) represents an oval \mathcal{O} in $\text{AG}(2, q)$. If the conic is degenerate, it's easy to see that the oval and conic have at most four points in common. So we may assume that the conic is non-degenerate. The nucleus of \mathcal{C} is $N = (y_1, 0, 1)$. We now show that N is the nucleus of the oval \mathcal{O} . The line $t = 0$ goes through N and meets the oval (25) when $s^{\sigma+2} = z_2$, which has one solution as $\sigma + 2$ is a permutation of \mathbb{F}_q . The line $s + y_1 = 0$ through N meets the oval (25) when $t^\sigma = y_1^{\sigma+2} + z_2$ which has one solution for t . Therefore, N is the nucleus, as it is the intersection of two tangent lines to the oval. It now follows from Lemma 6 that equations (24) and (25) have at most four common solutions. \square

We now show the existence of a point $P \notin (\mathcal{U}_{\text{BT}} \cup \ell_\infty)$ and a line ℓ such that $|\ell \cap \tau_P(\mathcal{U}_{\text{BT}})| = 3$, and demonstrate that our bound is sharp.

Lemma 7. *Let $y_1 = 0$, then the points of the oval given by equation (25) are*

$$\left\{ P_u = \left(\frac{z_2^{1-\sigma/2} u^\sigma}{1+u+u^\sigma}, \frac{z_2^{\sigma/2} (1+u^\sigma)}{1+u+u^\sigma} \right) \mid u \in \mathbb{F}_q \right\} \cup \left\{ (z_2^{1-\sigma/2}, z_2^{\sigma/2}) \right\}. \quad (26)$$

Proof. If $y_1 = 0$, then equation (25) reduces to

$$s^{\sigma+2} + t^\sigma + st + z_2 = 0 \quad (27)$$

Using the properties of σ described in Section 1.3, one can show the point $(z_2^{1-\sigma/2}, z_2^{\sigma/2})$ satisfies equation (27). Furthermore, the points $\overline{P}_u = (z_2^{1-\sigma/2} u^\sigma, z_2^{\sigma/2} (1+u^\sigma), 1+u+u^\sigma)$, where $u \in \mathbb{F}_q$, are projective points satisfying the following homogeneous equation

$$x^{\sigma+2} + y^\sigma z^2 + xyz^\sigma + z_2 z^{\sigma+2} = 0. \quad (28)$$

Because $\text{Tr}(u + u^\sigma) = 0$, and $\text{Tr}(1) = 1$ when $q = 2^{2e+1}$, we have $u^\sigma + u + 1 \neq 0$ for all $u \in \mathbb{F}_q$. Thus, normalising so $z = 1$, the points \overline{P}_u have the form $(s, t, 1)$ where s and t satisfy equation (27). \square

Corollary 5. *Let $y_1 = 0$ and consider the points P_u as described in Lemma 7. A point P_u lies on the conic given by equation (24), if and only if u is a root of the following polynomial*

$$a_1^{\sigma/2}u^\sigma + (z_2^{\sigma-1} + \delta^{\sigma/2}z_2 + z_2^{\sigma/2} + a_1^{\sigma/2})u^2 + z_2^{\sigma/2}u + \delta^{\sigma/2}z_2 + a_1^{\sigma/2} \quad (29)$$

Proof. By directly substituting P_u into equation (24) we have

$$(z_2^{2-\sigma} + \delta z_2^\sigma + z_2 + a_1)u^{2\sigma} + z_2u^\sigma + a_1u^2 + (\delta z_2^\sigma + a_1) = 0 \quad (30)$$

Raising both sides of equation (30) to the power of $\sigma/2$ yields our result. \square

Theorem 3. *Let U be a Buekenhout-Tits unital in $\text{PG}(2, q^2)$. The feet of a point $P \notin (\ell_\infty \cup U)$ meet a line ℓ in at most four points. Moreover, there exists a line ℓ and point P such that $|\ell \cap \tau_P(U)| = k$ for each $k \in \{0, 1, 2, 3, 4\}$.*

Proof. By Theorem 1 we may assume that $U = \mathcal{U}_{\text{BT}}$. The first part of the proof comes from Corollary 4. Let $P = (1, y_1, z_2\epsilon)$. All lines through P meet $\tau_P(U)$ in at most one point by definition, so it is clear that there exists lines ℓ such that $|\ell \cap \tau_P(U)|$ is zero or one. Because the points of $\tau_P(U)$ are not collinear, there exists a pair of points $Q, R \in \tau_P(U)$ such that the line QR does not contain $(0, 1, y_1)$. Hence, the line QR meets in precisely two points by Lemma 5.

Now consider a line ℓ with equation $(\delta + \epsilon)x + z = 0$ and let P be the point $(1, 0, \epsilon)$ (that is, $a_1 = \delta, a_2 = 1, b_1 = b_2 = y_1 = 0, z_2 = 1$). The number of points of $\ell \cap \tau_P(U)$ is the same as the number of solutions to equations (18) and (19). By Lemma 7 the points P_u satisfying equation (19) lie on the conic (18) when

$$\delta^{\sigma/2}u^\sigma + u = u(\delta^{\sigma/2}u^{\sigma-1} + 1) = 0, \quad (31)$$

which has two roots as $\sigma - 1$ is a permutation of \mathbb{F}_q . It can also be shown that $(z_2^{1-\sigma/2}, z_2^{\sigma/2}) = (1, 1)$ satisfies both equations. Hence, the intersection of the feet of the point $(1, 0, \epsilon)$ and ℓ has exactly three points.

Finally, consider the point $P(1, 0, \frac{1}{\delta^\sigma}\epsilon)$ and the line ℓ with dual coordinates $[\frac{1}{\delta} + \frac{1}{\delta^2}\epsilon, 0, 1]$. By Corollary 5, the number of feet of P on the line ℓ is the number of solutions to the equation (29), where $a_1 = \frac{1}{\delta}$ and $z_2 = \frac{1}{\delta^\sigma}$ which is

$$\frac{1}{\delta^{\sigma/2}}u^\sigma + \left(\frac{1}{\delta^{2-\sigma}} + \frac{1}{\delta}\right)u^2 + \frac{1}{\delta}u = 0. \quad (32)$$

Since equation (32) is a \mathbb{F}_2 -linearised polynomial, and there are at most 4 roots, we have that equation (29) has 1, 2, or 4 roots. We will show that, under the condition $\text{Tr}(\delta) = 1$, it has four roots. Multiplying equation (32) by δ yields $\delta^{1-\sigma/2}u^\sigma + (\delta^{\sigma-1} + 1)u^2 + u = 0$ and now substituting $a = \delta^{\sigma-1} + 1$ gives

$$(a^{\sigma/2} + 1)u^\sigma + au^2 + u = 0. \quad (33)$$

We find that $u = 0$ and $u = \frac{1}{a^{1+\sigma/2}}$ are solutions to equation (33). Now consider

$$u^\sigma + au^2 + 1 = 0. \quad (34)$$

Any solution to equation (34) also satisfies $(u^\sigma + au^2 + 1)^{\sigma/2} + u^\sigma + au^2 + 1 = 0$ which is precisely equation (33). Multiply equation (34) with $a^{\sigma+1}$, then we find $(a^{\sigma/2+1}u)^\sigma + (a^{\sigma/2+1}u)^2 + a^{\sigma+1} = 0$, and letting $z = (a^{\sigma/2+1}u)^2$,

$$z^{\sigma/2} + z + a^{\sigma+1} = 0, \quad (35)$$

which is known (see [11]) to have solutions if and only if $\text{Tr}(a^{\sigma+1}) = 0$. As $z = 0$ and $z = 1$ are not solutions of equation (35), no solutions of equation (35) correspond to the solutions $u = 0$ or $u = \frac{1}{a^{1+\sigma/2}}$ of equation (32). Furthermore, recall that equation (32) has 1, 2 or 4 solutions and that we have assumed that $\text{Tr}(\delta) = 1$. Since $\delta^{\sigma-1} = a + 1$, it follows that $\delta = (a + 1)^{\sigma+1}$ and $\text{Tr}(\delta) = \text{Tr}(a^{\sigma+1} + a^\sigma + a + 1) = \text{Tr}(a^{\sigma+1}) + \text{Tr}(1) = \text{Tr}(a^{\sigma+1}) + 1$. Hence, the conditions $\text{Tr}(\delta) = 1$ and $\text{Tr}(a^{\sigma+1}) = 0$ are equivalent, and we find exactly four roots to equation (32). \square

References

- [1] N. Abarzúa, R. Pomareda, and O. Vega. “Feet in orthogonal-Buekenhout-Metz unitals”. In: *Adv. Geom.* 18.2 (2018), pp. 229–236. DOI: 10.1515/advgeom-2017-0050.
- [2] A. Aguglia and G. L. Ebert. “A combinatorial characterization of classical unitals”. In: *Arch. Math. (Basel)* 78.2 (2002), pp. 166–172. DOI: 10.1007/s00013-002-8231-3.
- [3] R. D. Baker and G. L. Ebert. “On Buekenhout-Metz unitals of odd order”. In: *J. Combin. Theory Ser. A* 60.1 (1992), pp. 67–84. DOI: 10.1016/0097-3165(92)90038-V.
- [4] S. Barwick and G. L. Ebert. *Unitals in projective planes*. Springer Monographs in Mathematics. Springer, New York, 2008.
- [5] S. G. Barwick and Catherine T. Quinn. “Generalising a characterisation of Hermitian curves”. In: *J. Geom.* 70.1-2 (2001), pp. 1–7. DOI: 10.1007/PL00000978.
- [6] M. Ceria et al. *On near-MDS codes and caps*. 2021. DOI: 10.48550/ARXIV.2106.03402. URL: <https://arxiv.org/abs/2106.03402>.
- [7] A. Cossidente, G. L. Ebert, and G. Korchmáros. “A group-theoretic characterization of classical unitals”. In: *Arch. Math. (Basel)* 74.1 (2000), pp. 1–5. DOI: 10.1007/PL00000403.
- [8] G. L. Ebert. “Buekenhout-Tits unitals”. In: *J. Algebraic Combin.* 6.2 (1997), pp. 133–140. DOI: 10.1023/A:1008691020874.
- [9] G. L. Ebert. “On Buekenhout-Metz unitals of even order”. In: *European J. Combin.* 13.2 (1992), pp. 109–117. DOI: 10.1016/0195-6698(92)90042-X.
- [10] T. Feng and W. Li. “On the existence of O’Nan configurations in ovoidal Buekenhout-Metz unitals in $\text{PG}(2, q^2)$ ”. In: *Discrete Math.* 342.8 (2019), pp. 2324–2332. DOI: 10.1016/j.disc.2019.05.013.
- [11] G. Menichetti. “Roots of affine polynomials”. In: *Combinatorics ’84 (Bari, 1984)*. Vol. 123. North-Holland Math. Stud. North-Holland, Amsterdam, 1986, pp. 303–310.

- [12] J. A. Thas. “A combinatorial characterization of Hermitian curves”. In: *J. Algebraic Combin.* 1.1 (1992), pp. 97–102. DOI: 10.1023/A:1022437415099.

Address of the authors:

Jake Faulkner `jake.faulkner@pg.canterbury.ac.nz`

Geertrui Van de Voorde `geertrui.vandevoorde@canterbury.ac.nz`

School of Mathematics and Statistics

University of Canterbury

Private bag 4800

8140 Christchurch

New Zealand