

# On PFH and HF spectral invariants

Guanheng Chen

## Abstract

In this note, we define the link spectral invariants by using the cylindrical formulation of the quantitative Heegaard Floer homology. We call them HF spectral invariants. We deduce a relation between the HF spectral invariants and the PFH spectral invariants by using closed-open morphisms and open-closed morphisms. For the sphere, we prove that the homogenized HF spectral invariants at the unit are equal to the homogenized PFH spectral invariants. Moreover, we show that the homogenized PFH spectral invariants are quasimorphisms.

## Contents

<b>1</b>	<b>Introduction and main results</b>	<b>2</b>
<b>2</b>	<b>Preliminaries</b>	<b>8</b>
2.1	Periodic Floer homology . . . . .	8
2.2	Quantitative Heegaard Floer homology . . . . .	13
2.2.1	Cylindrical formulation of QHF . . . . .	13
2.2.2	Filtered QFH and spectral invariants . . . . .	15
<b>3</b>	<b>Product structure on HF</b>	<b>16</b>
3.1	Moduli space of HF curves . . . . .	16
3.2	Quantum product . . . . .	20
3.2.1	Reference relative homology classes . . . . .	22
3.2.2	Continuous morphisms . . . . .	23
3.2.3	Quantum product on HF . . . . .	25
3.2.4	Unit . . . . .	25
<b>4</b>	<b>Proof of Theorem 1</b>	<b>27</b>
4.1	The HF action spectrum . . . . .	27
4.2	Proof of Theorem 1 . . . . .	28

<b>5</b>	<b>Open-closed morphisms</b>	<b>32</b>
5.1	Construction and invariance of $\mathcal{OC}$	35
5.2	Computations of $\mathcal{OC}$	36
<b>6</b>	<b>Spectral invariants</b>	<b>42</b>
6.1	Comparing the PFH and HF spectral invariants	42
6.2	Homogenized spectral invariants	44
6.3	Quasimorphisms	45
	Bibliography	48

# 1 Introduction and main results

Let  $\Sigma$  be a closed surface with genus  $g$  and  $\omega$  a volume form of volume 1 (of course, the number 1 can be replaced by any positive number). Given a volume-preserving diffeomorphism  $\varphi : \Sigma \rightarrow \Sigma$ , M. Hutchings defines a version of Floer homology for  $(\Sigma, \omega, \varphi)$  which he calls **periodic Floer homology** [18, 20], abbreviated as PFH. Recently, D. Cristofaro-Gardiner, V. Humilière, and S. Seyfaddini use a twisted version of PFH to define a family of numerical invariants

$$c_d^{pfh} : C^\infty(S^1 \times \Sigma) \times \widetilde{PFH}(\Sigma, \varphi_H, \gamma_0) \rightarrow \{-\infty\} \cup \mathbb{R}.$$

called **PFH spectral invariants** [4] (also see [6, 16] for the non-Hamiltonian case).

A **link**  $\underline{L}$  on  $\Sigma$  is a disjoint union of simple closed curves. Under certain monotone assumptions, D. Cristofaro-Gardiner, V. Humilière, C. Mak, S. Seyfaddini and I. Smith show that the Lagrangian Floer homology of Lagrangian pair  $(Sym^d \varphi_H(\underline{L}), Sym^d \underline{L})$  in  $Sym^d \Sigma$ , denoted by  $HF(Sym^d \Sigma, Sym^d \underline{L}, Sym^d \varphi_H)$ , is well-defined and non-vanishing [7]. Here  $\varphi_H$  is a Hamiltonian symplectomorphism. They call the Floer homology  $HF(Sym^d \Sigma, Sym^d \underline{L}, Sym^d \varphi_H)$  **quantitative Heegaard Floer homology**, abbreviated as QHF. For any two different Hamiltonian symplectomorphisms, the corresponding QHF are canonically isomorphic to each other. Let  $HF(Sym^d \underline{L})$  denote an abstract group that is a union of all the QHF defined by Hamiltonian symplectomorphisms, modulo the canonical isomorphisms. Using R. Leclercq and F. Zapolsky's general results [30], they define a set of numerical invariants parameterized by QFH

$$c_{\underline{L}}^{hf} : C^\infty([0, 1] \times \Sigma) \times HF(Sym^d \underline{L}) \rightarrow \{-\infty\} \cup \mathbb{R}.$$

These numerical invariants are called **link spectral invariants**.

Even though these two spectral invariants come from different Floer theories, they satisfy many parallel properties. So it is natural to study whether they have any relation. To this end, our strategy is to construct morphisms between these two Floer

homologies. Because these two Floer theories are defined by counting holomorphic curves in manifolds of different dimensions, it is hard to define the morphisms directly. To overcome this issue, the author follows R. Lipshitz's idea [28] to define a homology by counting holomorphic curves in a 4-manifold, denoted by  $HF(\Sigma, \varphi_H(\underline{L}), \underline{L})$  [13]. Moreover, the author proves that there is an isomorphism

$$\Phi_H : HF(\Sigma, \varphi_H(\underline{L}), \underline{L}) \rightarrow HF(\text{Sym}^d \Sigma, \text{Sym}^d \underline{L}, \text{Sym}^d \varphi_H). \quad (1.1)$$

Therefore, this can be viewed as an alternative formulation of the quantitative Heegaard Floer homology. When the context is clear, we also call it QHF. It serves as a bridge between the QHF and PFH. The author establishes a homomorphism from PFH to QHF

$$(\mathcal{CO}_{\mathcal{Z}_0}(W, \Omega_H, \mathcal{L}_H)_J)_* : \widetilde{PFH}(\Sigma, \varphi_H, \gamma_0)_J \rightarrow HF(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})_J \quad (1.2)$$

which is called the **closed-open morphism**. The map (1.2) is an analogy of the usual closed-open morphism from the symplectic Floer homology to Lagrangian Floer homology defined by P. Albers [1]. A version of (1.2) also has been constructed by V. Colin, P. Ghiggini, and K. Honda [15] for a different setting. Using these morphisms, we obtain a partial result on the relation between PFH spectral invariants and HF spectral invariants [13].

In this note, we define the quantum product structures and spectral invariants for  $HF(\Sigma, \varphi_H(\underline{L}), \underline{L})$  as the Lagrangian Floer homology. Similar to the QHF, for any  $\varphi_H$ , the group  $HF(\Sigma, \varphi_H(\underline{L}), \underline{L})$  is isomorphic to an abstract group  $HF(\Sigma, \underline{L})$  canonically. The spectral invariants defined by  $HF(\Sigma, \varphi_H(\underline{L}), \underline{L})$  are denoted by  $c_{\underline{L}}$ . To distinguish with the link spectral invariants  $c_{\underline{L}}^{hf}$ , we call  $c_{\underline{L}}$  the **HF spectral invariants** instead. Via the isomorphism (1.1), we know that  $c_{\underline{L}}$  is equivalent to  $c_{\underline{L}}^{hf}$  in certain sense (see (2.16)).

The purpose of this paper is to study the properties of  $c_{\underline{L}}$  and try to understand the relations between  $c_{\underline{L}}$  and  $c_d^{pfh}$ . Before we state the main results, let us recall the assumptions on a link.

**Definition 1.1.** *Let  $\underline{L} = \sqcup_{i=1}^d L_i$  be a  $d$ -disjoint union of simple closed curves on  $\Sigma$ . We call  $\underline{L}$  a **link** on  $\Sigma$ . We say a link  $\underline{L}$  is **admissible** if it satisfies the following properties:*

- A.1** *The integer satisfies  $d = k + g$ , where  $g$  is the genus of  $\Sigma$  and  $k > 1$ .  $\sqcup_{i=1}^k L_i$  is a disjoint contractible simple curves. For  $k + 1 \leq i \leq d$ ,  $L_i$  is the cocore of the 1-handle. For each 1-handle, we have exactly one corresponding  $L_i$ .*
- A.2** *We require that  $\Sigma - \underline{L} = \cup_{i=1}^{k+1} \mathring{B}_i$ . Let  $B_i$  be the closure of  $\mathring{B}_i$ . Then  $B_i$  is a disk for  $1 \leq i \leq k$  and  $B_{k+1}$  is a planar domain with  $2g + k$  boundary components. For  $1 \leq i \leq k$ , the circle  $L_i$  is the boundary of  $B_i$ .*

**A.3**  $\mathring{B}_i \cap \mathring{B}_j = \emptyset$ .

**A.4** For  $1 \leq i < j \leq k+1$ , we require  $\int_{B_i} \omega = \int_{B_j} \omega = \frac{1}{k+1}$ .

A picture of an admissible link is shown in Figure 1. Note that if  $\underline{L}$  is admissible,




Figure 1: The red circles are the admissible link.

so is  $\varphi(\underline{L})$ , where  $\varphi$  is any Hamiltonian symplectomorphism. We assume that the link is admissible throughout.

**Remark 1.1.** To define  $HF(\Sigma, \varphi_H(\underline{L}), \underline{L})$ , we need much stronger assumptions on  $\underline{L}$  than [7] for technical reasons.

In the first part of this note, we study the properties of the spectral invariants  $c_{\underline{L}}$ . The results are summarized in the following theorem. These properties are parallel to the one in [7, 30].

**Theorem 1.** The spectral invariant  $c_{\underline{L}} : C^\infty([0, 1] \times \Sigma) \times HF(\Sigma, \underline{L}) \rightarrow \{-\infty\} \cup \mathbb{R}$  satisfies the following properties:

1. (Spectrality) For any  $H$  and  $a \neq 0 \in HF(\Sigma, \underline{L})$ , we have  $c_{\underline{L}}(H, a) \in \text{Spec}(H : \underline{L})$ .
2. (Hofer-Lipschitz) For  $a \neq 0 \in HF(\Sigma, \underline{L})$ , we have

$$d \int_0^1 \min_{\Sigma} (H_t - K_t) dt \leq c_{\underline{L}}(H, a) - c_{\underline{L}}(K, a) \leq d \int_0^1 \max_{\Sigma} (H_t - K_t) dt.$$

3. (Shift) Fix  $a \neq 0 \in HF(\Sigma, \underline{L})$ . Let  $c : [0, 1]_t \rightarrow \mathbb{R}$  be a function only dependent on  $t$ . Then

$$c_{\underline{L}}(H + c, a) = c_{\underline{L}}(H, a) + \int_0^1 c(t) dt.$$

4. (Homotopy invariance) Let  $H, K$  are two mean-normalized Hamiltonian functions. Suppose that they are homotopic in the sense of Definition 4.1. Then

$$c_{\underline{L}}(H, a) = c_{\underline{L}}(K, a).$$

5. (Lagrangian control) If  $H_t|_{L_i} = c_i(t)$ , then

$$c_{\underline{L}}(H, a) = c_{\underline{L}}(0, a) + \sum_{i=1}^d \int_0^1 c_i(t) dt.$$

Moreover, for any Hamiltonian function  $H$ , we have

$$\sum_{i=1}^d \int_0^1 \min_{L_i} H_t dt + c_{\underline{L}}(0, a) \leq c_{\underline{L}}(H, a) \leq c_{\underline{L}}(0, a) + \sum_{i=1}^d \int_0^1 \max_{L_i} H_t dt.$$

6. (Triangle inequality) For any Hamiltonian functions  $H, K$  and  $a, b \in HF(\Sigma, \underline{L})$ , we have

$$c_{\underline{L}}(H \# K, \mu_2(a \otimes b)) \leq c_{\underline{L}}(H, a) + c_{\underline{L}}(K, b),$$

where  $\mu_2$  is the quantum product defined in Section 3.

7. (Normalization) For the unit  $e$ , we have  $c_{\underline{L}}(0, e) = 0$ .

8. (Calabi property) Let  $\{\underline{L}_m\}_{m=1}^{\infty}$  be a sequence of admissible links. Suppose that  $\{\underline{L}_m\}_{m=1}^{\infty}$  is equidistributed in the sense of [7]. Let  $d_m$  denote the number of components of  $\underline{L}_m$ . Then for  $a \neq 0 \in HF(\Sigma, \underline{L})$ , we have

$$\lim_{m \rightarrow \infty} \frac{1}{d_m} (c_{\underline{L}_m}(H, a) - c_{\underline{L}_m}(0, a)) = \int_0^1 \int_{\Sigma} H_t dt \wedge \omega.$$

**Remark 1.2.** We don't know whether the isomorphism (1.1) is canonical (but we expect this is true). Also, we don't know whether the product  $\mu_2$  agrees with the usual quantum product in monotone Lagrangian Floer homology. So we cannot deduce Theorem 1 from (1.1) and the results in [7, 30] directly. The methods in the proof of Theorem 1 basically the same as [7, 30].

In [13], we define the closed-open morphisms (1.2). We use the same techniques to construct a “reverse” of the closed-open morphisms, called **open-closed morphisms**.

**Theorem 2.** Let  $\underline{L}$  be an admissible link and  $\varphi_H$  a  $d$ -nondegenerate Hamiltonian symplectomorphism. Fix a reference 1-cycle  $\gamma_0$  with degree  $d$  and a base point  $\mathbf{x} \in \underline{L}$ . Let  $\mathcal{Z}_{ref} \in H_2(W, \mathbf{x}_H, \gamma_0)$  be a reference relative homology class. Let  $(W, \Omega_H, \mathcal{L}_H)$  be the open-closed symplectic cobordism defined in Section 5. Then for a generic admissible

almost complex structure  $J \in \mathcal{J}(W, \Omega_H)$ , the triple  $(W, \Omega_H, \mathcal{L}_H)$  induces a homomorphism

$$(\mathcal{OC}_{Z_{ref}}(W, \Omega_H, \mathcal{L}_H)_J)_* : HF(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})_J \rightarrow \widetilde{PFH}(\Sigma, \varphi_H, \gamma_0)_J$$

satisfying the following properties:

- (**Partial invariance**) Suppose that  $\varphi_H, \varphi_G$  satisfy the following conditions: (see Definition 2.1)

♠.1 Each periodic orbit of  $\varphi_H$  with degree less than or equal  $d$  is either  $d$ -negative elliptic or hyperbolic.

♠.2 Each periodic orbit of  $\varphi_G$  with degree less than or equal  $d$  is either  $d$ -positive elliptic or hyperbolic.

Fix reference relative homology classes  $Z_{ref} \in H_2(X, \gamma_1, \gamma_0)$ ,  $Z_0 \in H_2(W, \mathbf{x}_G, \gamma_0)$  and  $Z_1 \in H_2(W, \mathbf{x}_H, \gamma_1)$  satisfying  $A_{ref} \# Z_0 = Z_{ref} \# Z_1$ , where  $A_{ref}$  is the class defining the continuous morphism  $\mathcal{I}_{0,0}^{H,G}$ . Then for any generic admissible almost complex structures  $J_H \in \mathcal{J}(W, \Omega_H)$  and  $J_G \in \mathcal{J}(W, \Omega_G)$ , we have the following commutative diagram:

$$\begin{array}{ccc} HF_*(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})_{J_H} & \xrightarrow{(\mathcal{OC}_{Z_1}(W, \Omega_H, \mathcal{L}_H)_{J_H})_*} & \widetilde{PFH}_*(\Sigma, \varphi_H, \gamma_1)_{J_H} \\ \downarrow \mathcal{I}_{0,0}^{H,G} & & \downarrow PFH_{Z_{ref}}^{sw}(X, \Omega_X) \\ HF_*(\Sigma, \varphi_G(\underline{L}), \underline{L}, \mathbf{x})_{J_G} & \xrightarrow{(\mathcal{OC}_{Z_0}(W, \Omega_G, \mathcal{L}_G)_{J_G})_*} & \widetilde{PFH}_*(\Sigma, \varphi_G, \gamma_0)_{J_G} \end{array}$$

Here  $PFH_{Z_{ref}}^{sw}(X, \Omega_X)$  is the PFH cobordism map induced by symplectic cobordism (2.10) and  $\mathcal{I}_{0,0}^{H,G}$  is the continuous morphism on QHF defined in Section 3.

- (**Non-vanishing**) There are nonzero classes  $e_\diamond \in HF(\Sigma, \underline{L})$  and  $\sigma_{\diamond H}^{\mathbf{x}} \in \widetilde{PFH}(\Sigma, \varphi_H, \gamma_H^{\mathbf{x}})$  such that if  $\varphi_H$  satisfies the condition (♠.2), then we have

$$\mathcal{OC}_{Z_{ref}}(W, \varphi_H, \mathcal{L}_H)_*((j_H^{\mathbf{x}})^{-1}(e_\diamond)) = \sigma_{\diamond H}^{\mathbf{x}},$$

where  $j_H^{\mathbf{x}}$  is the canonical isomorphism (2.14). In particular, the open-closed morphism is non-vanishing.

There is a spectral class  $e \in HF(\Sigma, \underline{L})$  called the **unit** (see Definition 3.6). Define spectral invariants

$$c_{\underline{L}}^-(H) := c_{\underline{L}}(H, e_\diamond) \quad \text{and} \quad c_{\underline{L}}^+(H) := c_{\underline{L}}(H, e).$$

**Corollary 1.2.** For any Hamiltonian function  $H$ , let  $\sigma_{\diamond H}^{\mathbf{x}} \in \widetilde{PFH}(\Sigma, \varphi_H, \gamma_H^{\mathbf{x}})$  be the class defined in Section 7 of [13]. We have

$$c_d^{pjh}(H, \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x})dt \leq c_{\underline{L}}^-(H) \leq c_{\underline{L}}^+(H) \leq c_d^{pjh}(H, \sigma_{\heartsuit H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x})dt,$$

where  $\int_0^1 H_t(\mathbf{x})dt$  is short for  $\sum_{i=1}^d \int_0^1 H_t(x_i)dt$ .

**Corollary 1.3.** Suppose that  $\Sigma = \mathbb{S}^2$ . Then for any Hamiltonian function  $H$ , we have

$$c_d^{pjh}(H, \sigma_{\heartsuit H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x})dt - 1 \leq c_{\underline{L}}^-(H) \leq c_{\underline{L}}^+(H) \leq c_d^{pjh}(H, \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x})dt.$$

Moreover, for any  $\varphi \in \text{Ham}(\mathbb{S}^2, \omega)$ , we have

$$\mu_{\underline{L}}(\varphi, e) = \mu_d^{pjh}(\varphi), \quad (1.3)$$

where  $\mu_{\underline{L}}, \mu_d^{pjh}$  are the homogenized spectral invariants (see Section 6 for details). In particular, for any two admissible links  $\underline{L}, \underline{L}'$  with same number of components, then we have  $\mu_{\underline{L}}(\varphi, e) = \mu_{\underline{L}'}(\varphi, e)$

The terms  $c_d^{pjh}(H, \sigma_{\heartsuit H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x})dt$ ,  $c_d^{pjh}(H, \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x})dt$  and  $c_{\underline{L}}(H, a)$  are independent of the choice of the base point. See discussion in Section 7 of [13].

**Remark 1.3.** For technical reasons, the cobordism maps on PFH are defined by using the Seiberg-Witten theory [27] and the isomorphism “PFH=SWF” [29]. Nevertheless, the proof of the Theorem 2 needs a holomorphic curves definition. The assumptions  $\spadesuit.1, \spadesuit.2$  are used to guarantee that the PFH cobordism maps can be defined by counting holomorphic curves. According to the results in [11], the Seiberg-Witten definition agrees with the holomorphic curves definition in these special cases. We believe that the assumptions  $\spadesuit.1, \spadesuit.2$  can be removed if one could define the PFH cobordism maps by pure holomorphic curve methods.

By Proposition 3.7 of [11], the conditions  $\spadesuit.1, \spadesuit.2$  can be achieved by a  $C^1$ -perturbation of the Hamiltonian functions. More precisely, fix a metric  $g_Y$  on  $S^1 \times \Sigma$ . For any  $\delta > 0$  and Hamiltonian function  $H$ , there is a Hamiltonian function  $H'$  such that

- $\varphi_{H'}$  satisfies  $\spadesuit.1$  ( $\spadesuit.2$ ).
- $|H' - H| + |dH - dH'|_{g_Y} \leq \delta$ .

Even Theorem 2 relies on the conditions  $\spadesuit.1, \spadesuit.2$ , the above estimates and the Hofer-Lipschitz property imply that Corollaries 1.2, 1.3 work for a general Hamiltonian function.

From [7], we know that the homogenized link spectral invariants are homogeneous quasimorphisms. We show that this is also true for  $\mu_d^{pfh}(\varphi)$ . Recall that a **homogeneous quasimorphism** on a group  $G$  is a map  $\mu : G \rightarrow \mathbb{R}$  such that

1.  $\mu(g^n) = n\mu(g)$ ;
2. there exists a constant  $D = D(\mu) \geq 0$ , called the **defect** of  $\mu$ , satisfying

$$|\mu(gh) - \mu(g) - \mu(h)| \leq D.$$

**Theorem 3.** *The homogenized spectral invariants  $\mu_d^{pfh} : Ham(\mathbb{S}^2, \omega) \rightarrow \mathbb{R}$  are homogeneous quasimorphisms with defect 1.*

**Relevant results** The Calabi property in Theorem 1 in fact is an analogy of the “ECH volume property” for embedded contact homology, it was first discovered by D. Cristofaro-Gardiner, M. Hutchings, and V. Ramos [3]. **Embedded contact homology** (short for “ECH”) is a sister version of the periodic Floer homology. The construction of ECH and PFH are the same. The only difference is that they are defined for different geometric structures. If a result holds for one of them, then one could expect that there should be a parallel result for another one. The Calabi property also holds for PFH. This is proved by O. Edtmair and Hutchings [16], also by D. Cristofaro-Gardiner, R. Prasad and B. Zhang independently [6]. The Calabi property for QHF is discovered in [7].

Recently, D. Cristofaro-Gardiner, V. Humilière, C. Mak, S. Seyfaddini and I. Smith show that the homogenized link spectral invariants satisfy the “two-terms Weyl law” for a class of automatous Hamiltonian functions [8] on the sphere. We believe that the HF spectral invariants agree with the link spectral invariants. If one could show that this is true, Corollary 1.3 implies that homogenized PFH spectral invariants agree with the homogenized link spectral invariants. This suggests that homogenized PFH spectral invariants should also satisfy the “two-terms Weyl law”.

## 2 Preliminaries

### 2.1 Periodic Floer homology

In this section, we review the definition of twisted periodic Floer homology and PFH spectral invariants. For more details, please refer to [20, 21, 4, 16].

Suppose that  $\Sigma$  is a closed surface and  $\omega$  is a volume form of volume 1. Given a **Hamiltonian function**  $H : S_t^1 \times \Sigma \rightarrow \mathbb{R}$ , then we have a unique vector field  $X_{H_t}$ , called the **Hamiltonian vector field**, satisfying the relation  $\omega(X_{H_t}, \cdot) = d_\Sigma H_t$ . Let



$\varphi_H^t$  be the flow generated by  $X_{H_t}$ , i.e.,  $\partial_t \varphi_H^t = X_{H_t} \circ \varphi_H^t$  and  $\varphi_H^0 = id$ . For each  $t$ ,  $\varphi_H^t$  is a symplectomorphism. The time-1 flow is denoted by  $\varphi_H := \varphi_H^1$ . A Hamiltonian function is called **automatous** if it is  $t$ -independent.

Fix a symplectomorphism  $\varphi$ . Define the **mapping torus** by

$$Y_\varphi := [0, 1]_t \times \Sigma / (0, \varphi(x)) \sim (1, x).$$

There is a natural vector field  $R := \partial_t$  and a closed 2-form  $\omega_\varphi$  on  $Y_\varphi$  induced from the above quotient. The pair  $(dt, \omega_\varphi)$  forms a stable Hamiltonian structure and  $R$  is the Reeb vector field. Let  $\xi := \ker \pi_*$  denote the vertical bundle of  $\pi : Y_\varphi \rightarrow S^1$ .

A **periodic orbit** is a map  $\gamma : \mathbb{R}/q\mathbb{Z} \rightarrow Y_\varphi$  satisfying the ODE  $\partial_t \gamma(t) = R \circ \gamma(t)$ . The number  $q \geq 0$  is called the **period** or **degree** of  $\gamma$ . Note that  $q$  is equal to the intersection number  $[\gamma] \cdot [\Sigma]$ .

A periodic orbit is called **nondegenerate** if the linearized return map does not have 1 as an eigenvalue. The nondegenerate periodic orbits are classified as either **elliptic** or **hyperbolic** according to the eigenvalues of linearized return maps. The symplectomorphism  $\varphi$  is called  **$d$ -nondegenerate** if every closed orbit with degree at most  $d$  is nondegenerate.

Let  $\gamma$  be an elliptic periodic orbit with period  $q$ . We can find a trivialization of  $\xi$  such that the linearized return map is a rotation  $e^{i2\pi\theta}$ , where  $\{\theta_t\}_{t \in [0, q]}$  is a continuous function with  $\theta_0 = 0$ . The number  $\theta = \theta_t|_{t=q} \in \mathbb{R}/\mathbb{Z}$  is called the **rotation number** of  $\gamma$  (see [21] for details). The following definition explains the terminologies in the assumptions **♠.1**, **♠.2**.

**Definition 2.1.** (see [22] Definition 4.1) Fix  $d > 0$ . Let  $\gamma$  be an embedded elliptic orbit with degree  $q \leq d$ .

- $\gamma$  is called  *$d$ -positive elliptic* if the rotation number  $\theta$  is in  $(0, \frac{q}{d}) \pmod{1}$ .
- $\gamma$  is called  *$d$ -negative elliptic* if the rotation number  $\theta$  is in  $(-\frac{q}{d}, 0) \pmod{1}$ .

**For our purpose, we assume that  $\varphi$  is Hamiltonian throughout** (but the construction of PFH works for a general symplectomorphism). Under the Hamiltonian assumption, we have the following diffeomorphism

$$\begin{aligned} \Psi_H : S_t^1 \times \Sigma &\rightarrow Y_{\varphi_H} \\ (s, t, x) &\rightarrow (s, t, (\varphi_H^t)^{-1}(x)). \end{aligned} \tag{2.4}$$

It is easy to check that  $\Psi_H^*(\omega_{\varphi_H}) = \omega + d(H_t dt)$  and  $(\Psi_H)_*(\partial_t + X_H) = R$ .

**PFH complex** An **orbit set** is a finite set of pairs  $\gamma = \{(\gamma_i, m_i)\}$ , where  $\{\gamma_i\}$  are distinct embedded periodic orbits and  $\{m_i\}$  are positive integers. An orbit set is called a **PFH generator** if it satisfies a further condition: If  $\gamma_i$  is hyperbolic, then  $m_i = 1$ .

Let  $H_2(Y_\varphi, \gamma_+, \gamma_-)$  denote the set of 2-chains  $Z$  in  $Y_\varphi$  with  $\partial Z = \gamma_+ - \gamma_-$ , modulo the boundary of 3-chains. We call the element  $Z \in H_2(Y_\varphi, \gamma_+, \gamma_-)$  a **relative homology classes**. This is an affine space of  $H_2(Y_\varphi, \mathbb{Z}) \cong \mathbb{Z}[\Sigma] \oplus (H_1(S^1) \otimes H_1(\Sigma))$ . Let  $[Z]$  denote the equivalent class in  $H_2(Y_\varphi, \gamma_+, \gamma_-)/\ker \omega_\varphi$ . Note that  $H_2(Y_\varphi, \gamma_+, \gamma_-)/\ker \omega_\varphi$  is an affine space of  $\mathbb{Z}[\Sigma]$ .

Fix a reference 1-cycle  $\gamma_0$  transversed to  $\xi$  positively. Assume that  $[\gamma_0] \cdot [\Sigma] > g(\Sigma)$  throughout. An **anchored orbit set** is a pair  $(\gamma, [Z])$ , where  $\gamma$  is an orbit set and  $[Z] \in H_2(Y_\varphi, \gamma, \gamma_0)/\ker \omega_\varphi$ . We call it an **anchored PFH generator** if  $\gamma$  is a PFH generator.

The chain complex  $\widetilde{PFC}(\Sigma, \varphi, \gamma_0)$  is the set of the formal sums (possibly infinity)

$$\sum a_{(\gamma, [Z])}(\gamma, [Z]), \quad (2.5)$$

where  $a_{(\gamma, [Z])} \in \mathbb{Z}/2\mathbb{Z}$  and each  $(\gamma, [Z])$  is an anchored PFH generator. Also, for any  $C \in \mathbb{R}$ , we require that there is only finitely many  $(\gamma, [Z])$  such that  $\int_Z \omega_{\varphi_H} > C$  and  $a_{(\gamma, [Z])} \neq 0$ .

Let  $\Lambda = \{\sum_i a_i q^{b_i} | a_i \in \mathbb{Z}/2\mathbb{Z}, b_0 < b_1 \dots\}$  be the Novikov ring. Then the  $\widetilde{PFC}(\Sigma, \varphi, \gamma_0)$  is  $\Lambda$ -module because we define an action

$$\sum_i a_i q^{b_i}(\gamma, [Z]) := \sum_i a_i(\gamma, [Z - b_i \Sigma]). \quad (2.6)$$

In most of the time, it is convenient to take

$$\gamma_0 = \Psi_H(S^1 \times \mathbf{x}), \quad (2.7)$$

denoted by  $\gamma_H^{\mathbf{x}}$ , where  $\mathbf{x} = (x_1, \dots, x_d)$  is  $d$ -points on  $\Sigma$  (not necessarily to be distinct).

**Differential on PFH** To define the differential, consider the symplectization

$$X := \mathbb{R}_s \times Y_\varphi, \quad \Omega := \omega_\varphi + ds \wedge dt.$$

An almost complex structure on  $X$  is called **admissible** if it preserves  $\xi$ , is  $\mathbb{R}$ -invariant, sends  $\partial_s$  to  $R$ , and its restriction to  $\xi$  is compatible with  $\omega_\varphi$ . The set of admissible almost complex structures is denoted by  $\mathcal{J}(Y_\varphi, \omega_\varphi)$ .

Given  $J \in \mathcal{J}(Y_\varphi, \omega_\varphi)$  and orbit sets  $\gamma_+ = \{(\gamma_{+,i}, m_i)\}$ ,  $\gamma_- = \{(\gamma_{-,j}, n_j)\}$ , let  $\mathcal{M}^J(\gamma_+, \gamma_-, Z)$  be the moduli space of punctured holomorphic curves  $u : \dot{F} \rightarrow X$  with the following properties:  $u$  has positive ends at covers of  $\gamma_{+,i}$  with total multiplicity  $m_i$ ,

negative ends at covers of  $\gamma_{-,j}$  with total multiplicity  $n_j$ , and no other ends. Also, the relative homology class of  $u$  is  $Z$ . Note that  $\mathcal{M}^J(\gamma_+, \gamma_-, Z)$  admits a natural  $\mathbb{R}$ -action.

For a relative homology class  $Z \in H_2(Y_\varphi, \gamma_+, \gamma_-)$ , Hutchings defines a topology index called **ECH index** [18, 19]. Fix a trivialization  $\tau$  of  $\xi$ . The ECH index is given by the following formula:

$$I(Z) := c_\tau(\xi|_Z) + Q_\tau(Z) + \sum_i \sum_{p=1}^{m_i} CZ_\tau(\gamma_{+,i}^p) - \sum_j \sum_{q=1}^{n_j} CZ_\tau(\gamma_{-,j}^q),$$

where  $c_\tau(\xi|_Z)$  is the relative Chern number,  $Q_\tau(Z)$  is the relative self-intersection number and  $CZ_\tau$  is the Conley-Zehnder index. We refer readers to [18, 19] for the precise definition of these quantities.

The differential  $\partial_J$  on  $\widetilde{PFC}(\Sigma, \varphi, \gamma_0)$  is defined by

$$\partial_J(\gamma_+, [Z_+]) := \sum_{\gamma_-} \sum_{Z, I(Z)=1} \#(\mathcal{M}^J(\gamma_+, \gamma_-, Z)/\mathbb{R})(\gamma_-, [Z_+ - Z]).$$

The homology of  $(\widetilde{PFC}(\Sigma, \varphi, \gamma_0), \partial_J)$  is called the **twisted periodic Floer homology**, denoted by  $\widetilde{PFH}(\Sigma, \varphi, \gamma_0)_J$ . By Corollary 1.1 of [29], PFH is independent of the choice of almost complex structures and Hamiltonian isotopic of  $\varphi$ . Note that  $\widetilde{PFH}(\Sigma, \varphi, \gamma_0)$  is a  $\Lambda$ -module because the action (2.6) descends to the homology.

**The U-map** There is a well-defined map

$$U : \widetilde{PFH}(\Sigma, \varphi_H, \gamma_0) \rightarrow \widetilde{PFH}(\Sigma, \varphi_H, \gamma_0).$$

Fix  $z \in \mathbb{R} \times Y_{\varphi_H}$ . The definition of the U-map is similar to the differential. Instead of counting  $I = 1$  holomorphic curves modulo  $\mathbb{R}$  translation, the U-map is defined by counting  $I = 2$  holomorphic curves that pass through the fixed point  $z$  and modulo  $\mathbb{R}$  translation. The homotopy argument can show that the U-map is independent of the choice of  $z$ . For more details, please see Section 2.5 of [26].

**Filtered PFH** We define a functional  $\mathbb{A}_H$  on the anchored orbit sets by

$$\mathbb{A}_H(\gamma, [Z]) := \int_Z \omega_\varphi.$$

Let  $\widetilde{PFC}^L(\Sigma, \varphi, \gamma_0)$  be the set of formal sum (2.5) satisfying  $\mathbb{A}_H(\gamma, [Z]) < L$ . This is a subcomplex of  $(\widetilde{PFC}(\Sigma, \varphi, \gamma_0), \partial_J)$ . The homology is denoted by  $\widetilde{PFH}^L(\Sigma, \varphi, \gamma_0)$ . Let  $\mathbf{i}_L : \widetilde{PFH}^L(\Sigma, \varphi, \gamma_0) \rightarrow \widetilde{PFH}(\Sigma, \varphi, \gamma_0)$  be the map induced by the inclusion. Fix  $\sigma \in \widetilde{PFH}(\Sigma, \varphi, \gamma_0)$ . The **PFH spectral invariant** is defined by

$$c_d^{pfh}(H, \sigma, \gamma_0) := \inf\{L \in \mathbb{R} | \sigma \text{ belongs to the image of } \mathbf{i}_L\}.$$

We extend the PFH spectral invariant for a degenerate  $\varphi_H$  by setting

$$c_d^{pfh}(H, \sigma, \gamma_0) = \lim_{n \rightarrow \infty} c_d^{pfh}(H_n, \sigma_n, \gamma_0), \quad (2.8)$$

where  $\varphi_{H_n}$  are  $d$ -nondegenerate,  $\{H_n\}_{n=1}^\infty$   $C^\infty$ -converges to  $H$ , and  $\sigma_n \in \widetilde{PFH}(\Sigma, \varphi_{H_n}, \gamma_0)$  is the class corresponding to  $\sigma$ .

**Cobordism maps on PFH** Let  $(X, \Omega_X)$  be a symplectic 4-manifold. Suppose that there exists a compact subset  $K$  such that

$$(X - K, \Omega_X) \cong ([0, \infty) \times Y_{\varphi_+}, \omega_{\varphi_+} + ds \wedge dt) \cup ((-\infty, 0] \times Y_{\varphi_-}, \omega_{\varphi_-} + ds \wedge dt) \quad (2.9)$$

We allow  $Y_{\varphi_+} = \emptyset$  or  $Y_{\varphi_-} = \emptyset$ . We call  $(X, \Omega_X)$  a **symplectic cobordism** from  $(Y_{\varphi_+}, \omega_{\varphi_+})$  to  $(Y_{\varphi_-}, \omega_{\varphi_-})$ . Fix a reference homology class  $Z_{ref} \in H_2(X, \gamma_0, \gamma_1)$ . The symplectic manifold  $(X, \Omega_X)$  induces a homomorphism

$$PFH_{Z_{ref}}^{sw}(X, \Omega_X) : \widetilde{PFH}(\Sigma, \varphi_+, \gamma_0) \rightarrow \widetilde{PFH}(\Sigma, \varphi_-, \gamma_1).$$

This homomorphism is called a **PFH cobordism map**.

Following Hutchings-Taubes's idea [23], the cobordism map  $PFH_{Z_{ref}}^{sw}(X, \Omega_X)$  is defined by using the Seiberg-Witten theory [27] and Lee-Taubes's isomorphism [29]. Even the cobordism maps are defined by Seiberg-Witten theory, they satisfy some nice properties called **holomorphic curves axioms**. It means that the PFH cobordism maps count holomorphic curves in certain sense. For the precise statement, we refer readers to [11] and Appendix of [13].

In this paper, we will focus on the following special cases of  $(X, \Omega_X)$ .

1. Given two Hamiltonian functions  $H_+, H_-$ , define a homotopy  $H_s := \chi(s)H_+ + (1 - \chi(s))H_-$ , where  $\chi$  is a cut off function such that  $\chi = 1$  for  $s \geq R_0 > 0$  and  $\chi = 0$  for  $\chi \leq 0$ . Define

$$\begin{aligned} X &:= \mathbb{R}_s \times S_t^1 \times \Sigma, \\ \omega_X &:= \omega + dH_s \wedge dt, \quad \Omega_X := \omega_X + ds \wedge dt. \end{aligned} \quad (2.10)$$

This is a symplectic cobordism if  $R_0$  is large enough. Note that we identify  $Y_{\varphi_{H_\pm}}$  with  $S^1 \times \Sigma$  implicitly by using (2.4). Fix a reference relative homology class  $Z_{ref} \in H_2(X, \gamma_0, \gamma_1)$ . Then we have a cobordism map

$$PFH_{Z_{ref}}^{sw}(X, \Omega_X) : \widetilde{PFH}(\Sigma, \varphi_{H_+}, \gamma_0) \rightarrow \widetilde{PFH}(\Sigma, \varphi_{H_-}, \gamma_1).$$

This map only depends on  $H_+, H_-$  and the relative homology class  $Z_{ref}$ .

2. Let  $(B_-, \omega_{B_-}, j_{B_-})$  be a sphere with a puncture  $p$ . Suppose that we have neighbourhood  $U$  of  $p$  so that we have the following identification

$$(B_-, \omega_{B_-}, j_{B_-})|_U \cong ([0, \infty)_s \times S_t^1, ds \wedge dt, j),$$

where  $j$  is a complex structure that maps  $\partial_s$  to  $\partial_t$ . Let  $\chi : \mathbb{R} \rightarrow \mathbb{R}$  be cut off function such that  $\chi = 1$  when  $s \leq -R_0$  and  $\chi(s) = 0$  when  $s \geq 0$ . Take

$$\begin{aligned} X_- &:= B_- \times \Sigma \\ \omega_{X_-} &:= \omega + d(\chi(s)Hdt) \quad \Omega_{X_-} := \omega_{X_-} + \omega_{B_-}. \end{aligned} \tag{2.11}$$

For sufficiently large  $R_0 > 0$ ,  $(X_-, \Omega_{X_-})$  is a symplectic manifold satisfying (2.9).

In the case (2.10), if  $H_+$  satisfies  $\spadesuit.1$  and  $H_-$  satisfies  $\spadesuit.2$ , the author shows that the cobordism map  $PFH_{Z_{ref}}^{sw}(X, \Omega_X)$  can be defined alternatively by using the pure holomorphic curve methods [11]. The holomorphic curves definition will be used to prove Theorem 2. That is why we need the assumptions  $\spadesuit.1$ ,  $\spadesuit.2$  in the statement.

## 2.2 Quantitative Heegaard Floer homology

In this section, we review the cylindrical formulation of QHF defined in [13].

### 2.2.1 Cylindrical formulation of QHF

Fix an admissible link  $\underline{L} = \cup_{i=1}^d L_i$  and a Hamiltonian symplectomorphism  $\varphi_H$ . We always assume that  $\varphi_H$  is **nondegenerate** in the sense that  $\varphi_H(\underline{L})$  intersects  $\underline{L}$  transversely.

A **Reeb chord** of  $\varphi_H$  is a  $d$ -union of paths

$$\mathbf{y} = [0, 1] \times (y_1, \dots, y_d) \subset [0, 1] \times \Sigma,$$

where  $y_i \in L_i \cap \varphi_H(L_{\sigma(i)})$  and  $\sigma : \{1, \dots, d\} \rightarrow \{1, \dots, d\}$  is a permutation. Obviously, a Reeb chord is determined by  $d$ -distinct intersection points  $(y_1, \dots, y_d)$ . Thus, we don't distinguish the Reeb chords and  $d$ -intersection points.

Fix a base point  $\mathbf{x} = (x_1, \dots, x_d)$ , where  $x_i \in L_i$ . It determines a reference chord

$$\mathbf{x}_H(t) = \varphi_H \circ (\varphi_H^t)^{-1}(\mathbf{x}) \subset [0, 1]_t \times \Sigma$$

from  $\{0\} \times \varphi_H(\underline{L})$  to  $\{1\} \times \underline{L}$ .

Let  $(E := \mathbb{R}_s \times [0, 1]_t \times \Sigma, \Omega := \omega + ds \wedge dt)$  be a symplectic manifold. Let  $\mathcal{L} = \mathbb{R} \times (\{0\} \times \varphi_H(\underline{L}) \cup \{1\} \times \underline{L})$  be a union of Lagrangian submanifolds in  $(E, \Omega)$ . Let  $\mathbf{y}_\pm$  be two Reeb chords. Then we have a concept called  $d$ -**multisection** in  $E$ . Roughly

speaking, this is a map  $u : \dot{F} \rightarrow E$  which is asymptotic to  $\mathbf{y}_\pm$  as  $s \rightarrow \pm\infty$  and satisfies the Lagrangian boundary conditions, where  $\dot{F}$  is a Riemann surface with boundary punctures. If a  $d$ -multisection is holomorphic, we call it an **HF curve**. The set of equivalence classes of the  $d$ -multisections is denoted by  $H_2(E, \mathbf{y}_+, \mathbf{y}_-)$ . An element in  $H_2(E, \mathbf{y}_+, \mathbf{y}_-)$  is also called a **relative homology class**. Fix  $A \in H_2(E, \mathbf{y}_+, \mathbf{y}_-)$ . The ECH index also can be generalized to the current setting, denoted by  $I(A)$ . The definition of relative homology class, HF curves and ECH index will be postponed to Section 3. We will define these concepts for a slightly general setting.

Given a Reeb chord  $\mathbf{y}$ , a **capping** of  $\mathbf{y}$  is an equivalence class  $[A]$  in  $H_2(E, \mathbf{x}_H, \mathbf{y}) / \ker \omega$ . Define the complex  $CF(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})$  be the set of formal sums of capping

$$\sum_{(\mathbf{y}, [A])} a_{(\mathbf{y}, [A])}(\mathbf{y}, [A]) \quad (2.12)$$

satisfying that  $a_{(\mathbf{y}, [A])} \in \mathbb{Z}/2\mathbb{Z}$  and for any  $C \in \mathbb{R}$ , there are only finitely  $(\mathbf{y}, [A])$  such that  $-\int_A \omega > C$  and  $a_{(\mathbf{y}, [A])} \neq 0$ .

For  $1 \leq i \leq k$ , let  $v_i : [0, 1]_s \times [0, 1]_t \rightarrow \Sigma$  be a map such that  $v_i(0, t) = v_i(1, t) = v_i(s, 0) = x_i$  and  $v_i(s, 1) \in L_i$  and represents the class  $[B_i] \in H_2(\Sigma, L_i, \mathbb{Z})$ . Define

$$\begin{aligned} u_{x_i} : [0, 1]_s \times [0, 1]_t &\rightarrow [0, 1]_s \times [0, 1]_t \times \Sigma \\ (s, t) &\rightarrow (s, t, \varphi_H \circ (\varphi_H^t)^{-1} \circ v_i(s, t)). \end{aligned}$$

Together with the trivial strip at  $x_j$  ( $j \neq i$ ),  $u_{x_i}$  represents a class in  $H_2(E, \mathbf{x}_H, \mathbf{x}_H)$ , still denoted by  $[B_i]$ . We also replace the map  $v_i$  by  $v'_i$ , where  $v'_i$  satisfies  $v'_i(0, t) = v'_i(1, t) = v'_i(s, 1) = x_i$  and  $v'_i(s, 0) \in L_i$  and represents the class  $[B_i] \in H_2(\Sigma, L_i)$ . Using the same construction, we have another map  $u'_{x_i}$ . The slightly different between  $u_{x_i}$  and  $u'_{x_i}$  is that  $u_{x_i}|_{t=1}$  wraps  $\partial B_i$  one time while  $u'_{x_i}|_{t=0}$  wraps  $\partial \varphi_H(B_i)$  one time. So we denote the equivalence class of  $u'_{x_i}$  in  $H_2(E, \mathbf{x}_H, \mathbf{x}_H)$  by  $[\varphi_H(B_i)]$ . By the monotone assumption **(A.4)**, all the classes  $[B_i]$  and  $[\varphi_H(B_i)]$  are equivalent in  $H_2(E, \mathbf{x}_H, \mathbf{x}_H) / \ker \omega$ , written as  $\mathcal{B}$ .

Let  $R = \{\sum_i a_i T^{b_i} | a_i \in \mathbb{Z}/2\mathbb{Z}, b_0 < b_1 \dots\}$  be the Novikov ring. To distinguish the one for PFH, we use different notations to denote the ring and the formal variable. Then  $CF(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})$  is a  $R$ -module because we have the following action

$$\sum_i a_i T^{b_i} \cdot (\mathbf{y}, [A]) := \sum_i a_i (\mathbf{y}, [A] + b_i \mathcal{B}). \quad (2.13)$$

Let  $\mathcal{J}_E$  denote the set of  $\Omega$ -compatible almost complex structures satisfying that  $J$  is  $\mathbb{R}_s$ -invariant,  $J(\partial_s) = \partial_t$ ,  $J$  sends  $T\Sigma$  to itself and  $J|_{T\Sigma}$  is  $\omega$ -compatible. Fix  $J \in \mathcal{J}_E$ . Let  $\mathcal{M}^J(\mathbf{y}_+, \mathbf{y}_-, A)$  denote the moduli space of HF curves that are asymptotic to  $\mathbf{y}_\pm$  as  $s \rightarrow \pm\infty$  and have relative homology class  $A$ . Because  $J$  is  $\mathbb{R}_s$ -invariant, this induces a natural  $\mathbb{R}$ -action on  $\mathcal{M}^J(\mathbf{y}_+, \mathbf{y}_-, A)$ .

Fix a generic  $J \in \mathcal{J}_E$ . The differential is defined by

$$d_J(\mathbf{y}_+, [A_+]) = \sum_{A \in H_2(E, \mathbf{y}_+, \mathbf{y}_-), I(A)=1} \# (\mathcal{M}^J(\mathbf{y}_+, \mathbf{y}_-, A)/\mathbb{R}) (\mathbf{y}_-, [A_+ \# A]).$$

The homology of  $(CF_*(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x}), d_J)$  is well defined [13], denoted by  $HF_*(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})_J$ . Again, the Floer homology is a  $R$ -module.

By Proposition 3.9 of [13], the homology is independent of the choices of  $J$  and  $H$ . For different choices of  $(J, H)$ , there is an isomorphism between the corresponding QHF called a **continuous morphism**. More details about this point are given in Section 3 later. For two different choices of base points, the corresponding homologies are also isomorphic. Let  $HF(\Sigma, \underline{L})$  be the direct limit of the continuous morphisms. For any  $H$ , we have an isomorphism

$$j_H^{\mathbf{x}} : HF(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x}) \rightarrow HF(\Sigma, \underline{L}). \quad (2.14)$$

Combining the isomorphism (1.1) with Lemma 6.10 of [7], we know that  $HF_*(\Sigma, \underline{L})$  is isomorphic to  $H^*(\mathbb{T}^d, R)$  as an  $R$ -vector space, where  $\mathbb{T}^d$  is the  $d$ -torus.

**Remark 2.1.** *Even though we only define the QHF for a Hamiltonian symplectic morphism  $\varphi_H$ , the above construction also works for a pair of Hamiltonian symplectic morphisms  $(\varphi_H, \varphi_K)$ . Because  $\varphi_K(\underline{L})$  is also an admissible link, we just need to replace  $\underline{L}$  by  $\varphi_K(\underline{L})$ . The result is denoted by  $HF(\Sigma, \varphi_H(\underline{L}), \varphi_K(\underline{L}), \mathbf{x})$ .*

## 2.2.2 Filtered QFH and spectral invariants

Similar as [30, 7], we define an **action functional** on the generators by

$$\mathcal{A}_H(\mathbf{y}, [A]) := - \int_A \omega + \int_0^1 H_t(\mathbf{x}) dt.$$

Let  $CF^L(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})$  be the set of formal sums (2.13) satisfying  $\mathcal{A}_H(\mathbf{y}, [A]) < L$ . It is easy to check that it is a subcomplex. The **filtered** QHF, denoted by  $HF^L(\Sigma, \varphi_H(\underline{L}), \underline{L})$ , is the homology  $(CF^L(\Sigma, \varphi_H(\underline{L}), \underline{L}), d_J)$ . Let

$$i_L : HF^L(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x}) \rightarrow HF(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})$$

be the homomorphism induced by the inclusion.

**Definition 2.2.** *Fix  $a \in HF(\Sigma, \underline{L})$ . The **HF spectral invariant** is*

$$c_{\underline{L}}(H, a) := \inf\{L \in \mathbb{R} \mid (j_H^{\mathbf{x}})^{-1}(a) \text{ belongs to the image of } i_L\}.$$

Let  $\mathbf{c} = \sum a_{(\mathbf{y}, [A])}(\mathbf{y}, [A])$  be a cycle in  $CF(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x})$ . The action of this cycle is defined by

$$\mathcal{A}_H(\mathbf{c}) = \max\{\mathcal{A}_H(\mathbf{y}, [A]) \mid a_{(\mathbf{y}, [A])} \neq 0\}.$$

Then the spectral invariant can be expressed as

$$c_{\underline{L}}(H, a) = \inf\{\mathcal{A}_H(\mathbf{c}) \mid [\mathbf{c}] = i_H^{-1}(a)\}. \quad (2.15)$$

Let  $HF(Sym^d \Sigma, Sym^d \underline{L}, Sym^d \varphi_H)$  denote the QHF defined in [7]. Because QHF is independent of the choices of  $\varphi_H$  and  $\mathbf{x}$ , we have an abstract group  $HF(Sym^d \underline{L})$  and a canonical isomorphism

$$\mathbf{j}_H^{\mathbf{x}} : HF(Sym^d \Sigma, Sym^d \underline{L}, Sym^d \varphi_H) \rightarrow HF(Sym^d \underline{L}).$$

Since the isomorphism (1.1) also preserves the action filtrations, we have

$$\frac{1}{d} c_{\underline{L}}(H, a) = c_{\underline{L}}^{hf}(H, \mathbf{j}_H^{\mathbf{x}} \circ \Phi_H \circ (j_H^{\mathbf{x}})^{-1}(a)). \quad (2.16)$$

We want to emphasize that the isomorphism  $\mathbf{j}_H^{\mathbf{x}} \circ \Phi_H \circ (j_H^{\mathbf{x}})^{-1} : HF(\Sigma, \underline{L}) \rightarrow HF(Sym^d \underline{L})$  may depend on  $H$  and  $\mathbf{x}$ . Therefore, the relation (2.16) is not strong enough to transfer all the properties of  $c_{\underline{L}}^{hf}$  to  $c_{\underline{L}}$ .

### 3 Product structure on HF

In this section, we define the continuous morphisms, quantum product and unit on  $HF(\Sigma, \underline{L})$ .

#### 3.1 Moduli space of HF curves

In this subsection, we give the definition of HF curves, relative homology class, and the ECH index.

Let  $D_m$  be a disk with boundary punctures  $(p_0, p_1, \dots, p_m)$ , the order of the punctures is counter-clockwise. See Figure 2. Let  $\partial_i D_m$  denote the boundary of  $D_m$  connecting  $p_{i-1}$  and  $p_i$  for  $1 \leq i \leq m$ . Let  $\partial_{m+1} D_m$  be the boundary connecting  $p_m$  and  $p_0$ .

Fix a complex structure  $j_m$  and a Kähler form  $\omega_{D_m}$  over  $D_m$  throughout. We say that  $D_m$  is a **disk with strip-like ends** if for each  $p_i$  we have a neighborhood  $U_i$  of  $p_i$  such that

$$(U_i, \omega_{D_m}, j_m) \cong (\mathbb{R}_{\epsilon_i} \times [0, 1], ds \wedge dt, j), \quad (3.17)$$

where  $j$  is the standard complex structure on  $\mathbb{R} \times [0, 1]$  that  $j(\partial_s) = \partial_t$ , where  $\epsilon_i = +$  for  $1 \leq i \leq m$  and  $\epsilon_0 = -$ . Here  $\mathbb{R}_+ = [0, \infty)$  and  $\mathbb{R}_- = (-\infty, 0]$ .






Figure 2: A picture of the case  $m = 5$ .

Let  $\pi : E_m = D_m \times \Sigma \rightarrow D_m$  be the trivial fibration. A closed 2-form  $\omega_{E_m}$  is called **admissible** if  $\omega_{E_m}|_{\Sigma} = \omega$  and  $\omega_{E_m} = \omega$  over the strip-like ends. Note that  $\Omega_{E_m} = \omega_{E_m} + \omega_{D_m}$  is a symplectic form on  $E_m$  if  $\omega_{D_m}$  is large enough. As a result,  $(\pi : E_m \rightarrow D_m, \Omega_m)$  over  $U_i$  can be identified with

$$(\pi : U_i \times \Sigma \rightarrow U_i, \Omega_{D_m}) \cong (\pi_{\mathbb{R} \times [0,1]} : \mathbb{R}_{\epsilon_i} \times [0, 1] \times \Sigma \rightarrow \mathbb{R}_{\epsilon_i} \times [0, 1], \omega + ds \wedge dt). \quad (3.18)$$

We call it a **(strip-like) end** of  $(E_m, \Omega_{E_m})$  at  $p_i$ .

Let  $\mathcal{L} = (\mathcal{L}_1, \dots, \mathcal{L}_{m+1})$  be a chain of  $d$ -disjoint union Lagrangian submanifolds in  $\partial E_m$  satisfying the following conditions:

**C.1** Let  $\mathcal{L}_i = \mathcal{L}|_{\partial_i D_m} \subset \pi^{-1}(\partial_i D_m)$ .  $\mathcal{L}_i$  consists of  $d$ -disjoint union of Lagrangian submanifolds.

**C.2** For  $1 \leq i \leq m$ , over the end at  $p_i$  (under the identification 3.18), we have

$$\mathcal{L} = (\mathbb{R}_+ \times \{0\} \times \underline{L}_{p_{i-1}}) \cup (\mathbb{R}_+ \times \{1\} \times \underline{L}_{p_i}).$$

**C.3** Over the end at  $p_0$  (under the identification 3.18), we have

$$\mathcal{L} = (\mathbb{R}_- \times \{0\} \times \underline{L}_{p_0}) \cup (\mathbb{R}_- \times \{1\} \times \underline{L}_{p_m}).$$

**C.4** The links  $\{\underline{L}_{p_i}\}_{i=0}^m$  are admissible and they are Hamiltonian isotropic to each other.

**C.5** For  $z \in \partial D_m$ ,  $\mathcal{L} \cap \pi^{-1}(z)$  is an admissible link.

Let  $(E_m, \Omega_m, \mathcal{L}_m)$  and  $(E_n, \Omega_n, \mathcal{L}_n)$  be two symplectic fibrations. Suppose that the negative end of  $(E_m, \Omega_m, \mathcal{L}_m)$  agrees with the  $i$ -th positive end of  $(E_n, \Omega_n, \mathcal{L}_n)$ , i.e.,

$$\begin{aligned} (E_m, \Omega_m, \mathcal{L}_m)|_{U_0} &\cong (\mathbb{R}_{s_- \leq 0} \times [0, 1]_t \times \Sigma, \omega + ds_- \wedge dt, \mathbb{R}_{s_- \leq 0} \times (\{0\} \times \underline{L}_0 \cup \{1\} \times \underline{L}_1)) \\ (E_n, \Omega_n, \mathcal{L}_n)|_{U_i} &\cong (\mathbb{R}_{s_+ \geq 0} \times [0, 1]_t \times \Sigma, \omega + ds_+ \wedge dt, \mathbb{R}_{s_+ \geq 0} \times (\{0\} \times \underline{L}_0 \cup \{1\} \times \underline{L}_1)). \end{aligned}$$

Fix  $R \geq 0$ . Define the  $R$ -**stretched composition**  $(E, \Omega, \mathcal{L}) := (E_n, \Omega_n, \mathcal{L}_n) \circ_R (E_m, \Omega_m, \mathcal{L}_m)$  by

$$(E, \Omega, \mathcal{L}) = (E_n, \Omega_n, \mathcal{L}_n)|_{s_+ \leq R} \cup_{s_+ - R = s_- + R} (E_m, \Omega_m, \mathcal{L}_m)|_{s_- \geq -R}. \quad (3.19)$$

In most of the time, the number  $R$  is not important, we suppress it from the notation.

**Definition 3.1.** Fix Reeb chords  $\mathbf{y}_i \in \underline{L}_{p_{i-1}} \cap \underline{L}_{p_i}$  and  $\mathbf{y}_0 \in \underline{L}_{p_0} \cap \underline{L}_{p_m}$ . Let  $(\dot{F}, j)$  be a Riemann surface (possibly disconnected) with boundary punctures. A  **$d$ -multisection** is a smooth map  $u : (\dot{F}, \partial\dot{F}) \rightarrow E_m$  such that

1.  $u(\partial\dot{F}) \subset \mathcal{L}$ . Let  $\{L_j^i\}_{i=1}^d$  be the connected components of  $\mathcal{L}|_{\partial_j D}$ . For each  $1 \leq i \leq d$ ,  $u^{-1}(L_j^i)$  consists of exactly one component of  $\partial\dot{F}$ .
2. For  $1 \leq i \leq m$ ,  $u$  is asymptotic to  $\mathbf{y}_i$  as  $s \rightarrow \infty$ .
3.  $u$  is asymptotic to  $\mathbf{y}_0$  as  $s \rightarrow -\infty$ .
4.  $\int_F u^* \omega_{E_m} < \infty$ .

Let  $H_2(E_m, \mathbf{y}_1, \dots, \mathbf{y}_m, \mathbf{y}_0)$  be the set of continuous maps

$$u : (\dot{F}, \partial\dot{F}) \rightarrow (E, \mathcal{L} \cup_{i=1}^m \{\infty\} \times \mathbf{y}_i \cup \{-\infty\} \times \mathbf{y}_0)$$

satisfying the conditions 1), 2), 3) and modulo a relation  $\sim$ . Here  $u_1 \sim u_2$  if and only if their compactifications are equivalent in  $H_2(E_m, \mathcal{L} \cup_{i=1}^m \{\infty\} \times \mathbf{y}_i \cup \{-\infty\} \times \mathbf{y}_0; \mathbb{Z})$ . An element in  $H_2(E_m, \mathbf{y}_1, \dots, \mathbf{y}_m, \mathbf{y}_0)$  is called a **relative homology class**. An easy generalization is that one could replace the Reeb chords by the reference chords  $\mathbf{x}_H$  in the above definition.

**Definition 3.2.** An almost complex structure is called adapted to fibration if

1.  $J$  is  $\Omega_{E_m}$ -tame.
2. Over the strip-like ends,  $J$  is  $\mathbb{R}_s$ -invariant,  $J(\partial_s) = \partial_t$ ,  $J$  preserves  $T\Sigma$  and  $J|_{T\Sigma}$  is compatible with  $\omega$ .
3.  $\pi$  is complex linear with respect to  $(J, j_m)$ , i.e.,  $j_m \circ \pi_* = \pi_* \circ J$ .

Let  $\mathcal{J}_{tame}(E_m)$  denote the set of the almost complex structures adapted to fibration. Fix an almost complex structure  $J$ . If  $u$  is a  $J$ -holomorphic  $d$ -multisection, then  $u$  is called an **HF curve**.

Using the admissible 2-form  $\omega_{E_m}$ , we have a splitting  $TE_m = TE_m^h \oplus TE_m^v$ , where  $TE_m^v = \ker \pi_*$  and  $TE_m^h = \{v \in TE_m \mid \omega_{E_m}(v, w) = 0, w \in T^v E_m\}$ . With respect to this splitting, an almost complex structure  $J \in \mathcal{J}_{tame}(E_m)$  can be written as  $J = \begin{pmatrix} J^{hh} & 0 \\ J^{hv} & J^{vv} \end{pmatrix}$ . Therefore,  $J$  is  $\Omega_{E_m}$ -compatible if and only if  $J^{hv} = 0$ .

Let  $\mathcal{J}_{comp}(E_m) \subset \mathcal{J}_{tame}(E_m)$  denote the set of almost complex structures which are adapted to fibration and  $\Omega_{E_m}$ -compatible. We will use the almost complex structures in  $\mathcal{J}_{comp}(E_m)$  for later computations.

**Fredholm index and ECH index** We begin to define the index of an HF curve. There are two types of index defined for an HF curve, called Fredholm index and ECH index.

To begin with, fix a trivialization of  $u^*T\Sigma$  as follows. Fix a non-singular vector  $v$  on  $\underline{L}$ . Then  $(v, j_\Sigma(v))$  gives a trivialization of  $T\Sigma|_{\mathcal{L}}$ , where  $j_\Sigma$  is a complex structure on  $\Sigma$ . We extend the trivialization arbitrarily along  $\mathbf{y}_i$ . Such a trivialization is denoted by  $\tau$ .

Define a real line bundle  $\mathfrak{L}$  over  $\partial F$  as follows. Take  $\mathfrak{L}|_{\partial \hat{F}} := u^*(T\mathcal{L} \cap T\Sigma)$ . Extend  $\mathfrak{L}$  to  $\partial F - \partial \hat{F}$  by rotating in the counter-clockwise direction from  $u^*TL_{p_{j-1}}^i$  and  $u^*TL_{p_j}^i$  by the minimum amount. Then  $(u^*T\Sigma, \mathfrak{L})$  forms a bundle pair over  $\partial F$ . With respect to the trivialization  $\tau$ , we have a well-defined Maslov index  $\mu_\tau(u) := \mu(u^*T\Sigma, \mathfrak{L}, \tau)$  and relative Chern number  $c_1(u^*T\Sigma, \tau)$ . The number  $2c_1(u^*T\Sigma, \tau) + \mu_\tau(u)$  is independent of the trivialization  $\tau$ .

The **Fredholm index** of an HF curve is defined by

$$indu := -\chi(F) + 2c_1(u^*T\Sigma, \tau) + \mu_\tau(u) + d(2 - m).$$

The above index formula can be obtained by the doubling argument in Proposition 5.5.2 of [15].

Given  $A \in H_2(E_m, \mathbf{y}_+, \mathbf{y}_-)$ , an oriented immersed surface  $C \subset E_m$  is a  $\tau$ -representative of  $A$  if

1.  $C$  intersects the fibers positively along  $\partial C$ ;
2.  $\pi_{[0,1] \times \Sigma}|_C$  is an embedding near infinity;
3.  $C$  satisfies the  $\tau$ -trivial conditions in the sense of Definition 4.5.2 in [15].

Let  $C$  be a  $\tau$ -trivial representative of  $A$ . Let  $\psi$  be a section of the normal bundle  $N_C$  such that  $\psi|_{\partial C} = J\tau$ . Let  $C'$  be a push-off of  $C$  in the direction of  $\psi$ . Then the

relative self-intersection number is defined by

$$Q_\tau(A) := \#(C \cap C').$$

Suppose that  $A \in H_2(E_m, \mathbf{y}_+, \mathbf{y}_-)$  admits a  $\tau$ -representative. The **ECH index** of a relative homology class is

$$I(A) := c_1(T\Sigma|_A, \tau) + Q_\tau(A) + \mu_\tau(A) + d(1 - m).$$

Using the relative adjunction formula, we have the following result.

**Theorem 3.3.** *(Theorem 4.5.13 of [15]) Let  $u$  be a  $J$ -holomorphic HF curve. Then the ECH index and the Fredholm index satisfy the following relation:*

$$I(u) = \text{ind}u + 2\delta(u),$$

where  $\delta(u) \geq 0$  is a count of the singularities of  $u$  with positive weight. Moreover,  $I(u) = \text{ind}(u)$  if and only if  $u$  is embedded.

*Proof.* By the same argument in Lemma 4.5.9 [15], we have

$$\begin{aligned} c_1(u^*TE_m, (\tau, \partial_t)) &= c_1(du(TF), \partial_t) + c_1(N_u, J\tau) \\ &= \chi(F) - d + Q_\tau(u) - 2\delta(u), \end{aligned}$$

where  $N_u$  is the normal bundle of  $u$  and  $\partial_t$  is a trivialization of  $TD_m$  such that it agrees with  $\partial_t$  over the ends. On the other hand, we have

$$c_1(u^*TE_m, (\tau, \partial_t)) = c_1(u^*T\Sigma, \tau) + c_1(u^*TD_m, \partial_t) = c_1(u^*T\Sigma, \tau).$$

Combine the above two equations; then we obtain the ECH equality.  $\square$

## 3.2 Quantum product

With the above preliminaries, we now define the quantum product. To begin with, let us define the cobordism maps on QHF induced by  $(E_m, \Omega_m, \mathcal{L}_m)$ . Assume that  $\underline{L}_{p_i} = \varphi_{H_i}(\underline{L})$ . Define reference chords by  $\delta_i(t) := \varphi_{H_i}(\mathbf{x}_{\bar{H}_i \# H_{i-1}}(t))$  for  $1 \leq i \leq m$  and  $\delta_0(t) = \varphi_{H_m}(\mathbf{x}_{\bar{H}_m \# H_0}(t))$ , where  $\bar{H}_i(x) = -H_t(\varphi_H^t(x))$ . Here

$$H \# K(t, x) := H_t(x) + K_t((\varphi_H^t)^{-1}(x))$$

is the **composition** of two Hamiltonian functions. By the chain rule, we have  $\varphi_{H \# K}^t = \varphi_H^t \circ \varphi_K^t$ . There is another operation on Hamiltonian functions called the **join**. The join of  $H$  and  $K$  is defined by

$$H_t \diamond K_t(x) = \begin{cases} 2\rho'(2t)K_{\rho(2t)}(x) & \text{if } 0 \leq t \leq \frac{1}{2} \\ 2\rho'(2t-1)H_{\rho(2t-1)}(x) & \text{if } \frac{1}{2} \leq t \leq 1, \end{cases}$$

where  $\rho : [0, 1] \rightarrow [0, 1]$  is a fixed non-decreasing smooth function that is equal to 0 near 0 and equal to 1 near 1. As with the composition, the time 1-map of  $H_t \diamond K_t$  is  $\varphi_H \circ \varphi_K$ .

The following proposition is similar to the result in Section 4 of [14].

**Proposition 3.4.** *Let  $(\pi : E_m = D_m \times \Sigma \rightarrow D_m, \Omega_m)$  be the symplectic fiber bundle with strip-like ends. Let  $\mathcal{L}_m \subset \pi^{-1}(\partial D_m)$  be Lagrangian submanifolds of  $(E_m, \Omega_m)$  satisfying **C.1**, **C.2**, **C.3**, **C.4**, **C.5**. Fix a reference homology class  $A_{ref} \in H_2(E_m, \delta_1, \dots, \delta_m, \delta_0)$  and a generic almost complex structure  $J \in \mathcal{J}_{tam\epsilon}(E_m)$ . Then  $(\pi : E_m \rightarrow D_m, \Omega_m, \mathcal{L}_m)$  induces a homomorphism*

$$HF_{A_{ref}}(E_m, \Omega_m, \mathcal{L}_m)_J : \bigotimes_{i=1}^m HF(\Sigma, \underline{L}_{p_{i-1}}, \underline{L}_{p_i}) \rightarrow HF(\Sigma, \underline{L}_{p_0}, \underline{L}_{p_m}).$$

satisfying the following properties:

1. (Invariance) *If there exists a family of symplectic form  $\{\Omega_\tau\}_{\tau \in [0,1]}$  and a family of  $\Omega_\tau$ -Lagrangians  $\{\mathcal{L}\}_{\tau \in [0,1]} \subset \partial E_m$  satisfying **C.1**, **C.2**, **C.3**, **C.4**, **C.5** and  $\{(\Omega_\tau, \mathcal{L}_\tau)\}_{\tau \in [0,1]}$  is  $\tau$ -independent over the strip-like ends. Assume  $\{J_\tau\}_{\tau \in [0,1]}$  is a general family of almost complex structures. Then*

$$HF_{A_{ref}}(E_m, \Omega_0, \mathcal{L}_0)_{J_0} = HF_{A_{ref}}(E_m, \Omega_1, \mathcal{L}_1)_{J_1}.$$

*In particular, the cobordism maps are independent of the choice of almost complex structures.*

2. (Composition rule) *Suppose that the negative end of  $(E_m, \Omega_m, \mathcal{L}_m)$  agrees with the  $j$ -th positive end of  $(E_n, \Omega_n, \mathcal{L}_n)$ . Then we have*

$$HF_{A_2}(E_n, \Omega_n, \mathcal{L}_n) \circ HF_{A_1}(E_m, \Omega_m, \mathcal{L}_m) = HF_{A_1 \# A_2}(E_{m+n-1}, \Omega_{m+n-1}, \mathcal{L}_{m+n-1}).$$

*Proof.* In chain level, define

$$CF_{A_{ref}}(E_m, \Omega_m, \mathcal{L}_m)_J((\mathbf{y}_1, [A_1]) \otimes \dots \otimes (\mathbf{y}_m, [A_m])) = \sum_{I(A)=0} \# \mathcal{M}^J(\mathbf{y}_1, \dots, \mathbf{y}_m; \mathbf{y}_0, A)(\mathbf{y}_0, [A_0]).$$

Here  $A_0$  is determined by the relation  $A_1 \# \dots \# A_m \# A \# (-A_0) = A_{ref}$ .

To see the above map is well defined, first note that the HF curves are simple because they are asymptotic to the Reeb chords. Therefore, the transversality of moduli space can be obtained by a generic choice of an almost complex structure. By Theorem 3.3, the ECH indices of HF curves are nonnegative.

Secondly, consider a sequence of HF curves  $\{u_n : \dot{F} \rightarrow E_m\}_{n=1}^\infty$  in  $\mathcal{M}^J(\mathbf{y}_1, \dots, \mathbf{y}_m; \mathbf{y}_0, A)$ . Apply the Gromov compactness [2] to  $\{u_n\}_{n=1}^\infty$ . To rule out the bubbles, our assumptions on the links play a key role here. Note that the bubbles arise from pinching an

arc or an interior simple curve in  $F_n$ . Since the Lagrangians  $\mathcal{L}_m$  are pairwise disjoint, if an irreducible component  $v$  of the bubbles comes from pinching an arc  $a$ , then the endpoints of  $a$  must lie inside the same component of  $\mathcal{L}_m$ . By the open mapping theorem,  $v$  lies in a fiber  $\pi^{-1}(z)$ , where  $z \in \partial_j D_m$ . Let  $\mathcal{L}_m \cap \pi^{-1}(z) = \cup_{i=1}^d L_z^i$ . By assumptions,  $L_z^i$  bounded a disk  $B_z^i$  for  $1 \leq i \leq k$ . Then the image of  $v$  is either  $B_z^i$  or  $\Sigma - B_z^i$  for some  $1 \leq i \leq k$ , or  $\Sigma - L_z^j$  for some  $k+1 \leq j \leq d$ . Similarly, if  $v$  comes from pinching an interior simple curve in  $F_n$ , then the image of  $v$  must be a fiber  $\Sigma$ . The index formula in Lemma 3.3 [13] can be generalized to the current setting easily. As a result, the bubble  $v$  contributes at least 2 to the ECH index. This is because the Maslov index of a disk is 2 and adding a  $\Sigma$  will increase the ECH index by  $2(k+1)$ . This violates the condition that  $I = 0$ . Hence, the bubbles can be ruled out. Therefore,  $\mathcal{M}^J(\mathbf{y}_1, \dots, \mathbf{y}_m; \mathbf{y}_0, A)$  is compact. Similarly, the bubbles cannot appear in the module space of HF curves with  $I = 1$ . The standard neck-stretching and gluing argument [28] shows that  $CF_{A_{ref}}(E_m, \Omega_m, \mathcal{L}_m)$  is a chain map.

The invariance and the composition rule follow from the standard homotopy and neck-stretching argument. Again, the bubbles can be ruled by the index reason as the above.

□

### 3.2.1 Reference relative homology classes

Obviously, the cobordism maps depend on the choice of the reference relative homology class  $A_{ref}$ . For any two different reference homology classes, the cobordism maps defined by them are differed by a shifting (2.13). Here we fix a reference relative homology class in the following way:

Let  $\chi_+(s) : \mathbb{R}_s \rightarrow \mathbb{R}$  be a function such that  $\chi_+ = 1$  when  $s \leq -R_0$  and  $\chi_+ = 0$  when  $s \geq -1$ . Define a diffeomorphism

$$F_+ : \mathbb{R}_- \times [0, 1] \times \Sigma \rightarrow \mathbb{R}_- \times [0, 1] \times \Sigma$$

$$(s, t, x) \rightarrow (s, t, \varphi_K \circ \varphi_{\chi(s)H} \circ (\varphi_{\chi(s)H}^t)^{-1}(x)).$$

We view  $F_+$  as a map on the end of  $E_0$ , then extend  $F_+$  to be  $(z, x) \rightarrow (z, \varphi_K(x))$  over the rest of  $E_0$ . Let  $\mathcal{L}_+ := F_+(\partial D_0 \times \underline{L}) \subset \partial E_0$  be a submanifold. Note that  $\mathcal{L}_+|_{s \leq -R_0} = \mathbb{R}_{s \leq -R_0} \times (\{0\} \times \varphi_K \circ \varphi_H(\underline{L}) \cup \{1\} \times \varphi_K(\underline{L}))$ . The surface  $F_+(D_0 \times \{\mathbf{x}\})$  represent a relative homology class  $A^+ \in H_2(E_0, \emptyset, \varphi_K(\mathbf{x}_{\bar{K}\#(K\#H)}))$ .

For any Hamiltonian functions  $H_1, H_2$ , we find a suitable  $H$  such that  $H_1 = H\#K$  and  $H_2 = K$ . So the above construction gives us a class  $A_{H_1, H_2}^+ \in H_2(E_0, \emptyset, \varphi_{H_2}(\mathbf{x}_{\bar{H}_2\#H_1}))$ .

Let  $\bar{D}_0$  be a disk with a strip-like positive end. Define  $\bar{E}_0 := \bar{D}_0 \times \Sigma$ . By a similar construction, we have a fiber-preserving diffeomorphism  $F_- : \bar{E}_0 \rightarrow \bar{E}_0$ . Let

$\mathcal{L}_- := F_-(\partial\bar{D}_0 \times \underline{L})$ . Then  $A_{H_1, H_2}^- := [F_-(\bar{D}_0 \times \{\mathbf{x}\})]$  gives a relative homology class in  $H_2(\bar{E}_0, \varphi_{H_2}(\mathbf{x}_{\bar{H}_2 \# H_1}), \emptyset)$ .

Using  $A_{H_1, H_2}^\pm$ , we determine a unique reference homology class  $A_{ref} \in H_2(E_m, \delta_1, \dots, \delta_m, \delta_0)$  as follows: For  $i$ -th positive end of  $(E_m, \mathcal{L}_m)$ , we glue it with  $(E_0, \mathcal{L}_+)$  as in (3.19), where  $\mathcal{L}_+$  is determined by  $H_{i-1}, H_i$ . Similarly, we glue the negative end of  $(E_m, \mathcal{L}_m)$  with  $(\bar{E}_0, \mathcal{L}_-)$ . Then this gives us a pair  $(\bar{E} = \bar{D} \times \Sigma, \bar{\mathcal{L}})$ , where  $\bar{D}$  is a closed disk without puncture. Note that  $H_2(\bar{E}, \bar{\mathcal{L}}; \mathbb{Z}) \cong H_2(\bar{E}, \partial\bar{D} \times \underline{L}; \mathbb{Z})$ . Under this identification, we have a canonical class  $A_{can} = [\bar{D} \times \{\mathbf{x}\}] \in H_2(\bar{E}, \bar{\mathcal{L}}; \mathbb{Z})$ . We pick  $A_{ref} \in H_2(E_m, \delta_1, \dots, \delta_m, \delta_0)$  to be a unique class such that

$$A_{H_0, H_m}^- \# A_{ref} \#_{i=1}^m A_{H_{i-1}, H_i}^+ = A_{can}.$$

### 3.2.2 Continuous morphisms

In the case that  $m = 1$ , we identify  $\pi : E_1 \rightarrow D_1$  with  $\pi : \mathbb{R}_s \times [0, 1]_t \times \Sigma \rightarrow \mathbb{R}_s \times [0, 1]_t$ . Given two pairs of symplectomorphisms  $(\varphi_{H_1}, \varphi_{K_1})$  and  $(\varphi_{H_2}, \varphi_{K_2})$ , we can use the same argument in Lemma 6.1.1 of [14] to construct a pair  $(\Omega_1, \mathcal{L}_1)$  such that

1.  $\Omega_1$  is a symplectic form such that  $\Omega_1|_{|s| \geq R_0} = \omega + ds \wedge dt$ .
2.  $\mathcal{L}_1 \subset \mathbb{R} \times \{0, 1\} \times \Sigma$  are two  $d$ -disjoint union of  $\Omega_1$ -Lagrangian submanifolds,
3.  $\mathcal{L}_1|_{s \geq R_0} = (\mathbb{R}_{s \geq R_0} \times \{0\} \times \varphi_{H_1}(\underline{L})) \cup (\mathbb{R}_{s \geq R_0} \times \{1\} \times \varphi_{K_1}(\underline{L}))$ ,
4.  $\mathcal{L}_1|_{s \leq -R_0} = (\mathbb{R}_{s \leq -R_0} \times \{0\} \times \varphi_{H_2}(\underline{L})) \cup (\mathbb{R}_{s \leq -R_0} \times \{1\} \times \varphi_{K_2}(\underline{L}))$ ,

We call the above triple  $(E_1, \Omega_1, \mathcal{L}_1)$  a **Lagrangian cobordism** from  $(\varphi_{H_1}(\underline{L}), \varphi_{K_1}(\underline{L}))$  to  $(\varphi_{H_2}(\underline{L}), \varphi_{K_2}(\underline{L}))$ .

Recall that the reference class  $A_{ref}$  is the unique class defined in Section 3.2.1. By the invariance property in Proposition 3.4, the cobordism map  $HF_{A_{ref}}(E_1, \Omega_1, \mathcal{L}_1)$  only depends on  $\{(H_i, K_i)\}_{i=1,2}$ . We call it a **continuous morphism**, denoted by  $\mathcal{I}_{K_1, K_2}^{H_1, H_2}$ . The continuous morphisms satisfy  $\mathcal{I}_{K_2, K_3}^{H_2, H_3} \circ \mathcal{I}_{K_1, K_2}^{H_1, H_2} = \mathcal{I}_{K_1, K_3}^{H_1, H_3}$ . Thus,  $\mathcal{I}_{K_1, K_2}^{H_1, H_2}$  is an isomorphism.

The direct limit of  $HF(\Sigma, \varphi_H(\underline{L}), \varphi_K(\underline{L}), \mathbf{x})$  is denoted by  $HF(\Sigma, \underline{L})$ . Because  $HF(\Sigma, \varphi_H(\underline{L}), \varphi_K(\underline{L}), \mathbf{x})$  is independent of  $\mathbf{x}$ , so is  $HF(\Sigma, \underline{L})$ . We have a canonical isomorphism

$$j_{H, K}^\mathbf{x} : HF(\Sigma, \varphi_H(\underline{L}), \varphi_K(\underline{L}), \mathbf{x}) \rightarrow HF(\Sigma, \underline{L}). \quad (3.20)$$

that is induced by the direct limit.

Let  $H$  be a Hamiltonian function. We consider another homomorphism

$$I_H : CF(\Sigma, \varphi_K(\underline{L}), \underline{L}) \rightarrow CF(\Sigma, \varphi_{H \# K}(\underline{L}), \varphi_H(\underline{L})) \quad (3.21)$$

defined by mapping  $(\mathbf{y}, [A])$  to  $(\varphi_H(\mathbf{y}), [\varphi_H(A)])$ . Obviously, it induces an isomorphism  $(I_H)_*$  in the homology level. We call it the **naturality isomorphism**. In the following lemma, we show that it agrees with the continuous morphism.

**Lemma 3.5.** *The naturality isomorphisms satisfy the following diagram.*

$$\begin{array}{ccc} HF(\Sigma, \varphi_{K_1}(\underline{L}), \underline{L}) & \xrightarrow{(I_{H_1})_*} & HF(\Sigma, \varphi_{H_1 \# K_1}(\underline{L}), \varphi_{H_1}(\underline{L})) \\ \downarrow \mathcal{I}_{0,0}^{K_1, K_2} & & \downarrow \mathcal{I}_{H_1, H_2}^{H_1 \# K_1, H_2 \# K_2} \\ HF(\Sigma, \varphi_{K_2}(\underline{L}), \underline{L}) & \xrightarrow{(I_{H_2})_*} & HF(\Sigma, \varphi_{H_2 \# K_2}(\underline{L}), \varphi_{H_2}(\underline{L})) \end{array}$$

In particular, we have  $(I_{H_1})_* = \mathcal{I}_{H_1, 0}^{H_1 \# K_1, K_1}$ .

*Proof.* To prove the statement, we first split the diagram into two:

$$\begin{array}{ccc} HF(\Sigma, \varphi_{K_1}(\underline{L}), \underline{L}) & \xrightarrow{(I_{H_1})_*} & HF(\Sigma, \varphi_{H_1 \# K_1}(\underline{L}), \varphi_{H_1}(\underline{L})) \\ \downarrow \mathcal{I}_{0,0}^{K_1, K_2} & & \downarrow \mathcal{I}_{H_1, H_1}^{H_1 \# K_1, H_1 \# K_2} \\ HF(\Sigma, \varphi_{K_2}(\underline{L}), \underline{L}) & \xrightarrow{(I_{H_1})_*} & HF(\Sigma, \varphi_{H_1 \# K_2}(\underline{L}), \varphi_{H_1}(\underline{L})) \\ \downarrow Id & & \downarrow \mathcal{I}_{H_1, H_2}^{H_1 \# K_2, H_2 \# K_2} \\ HF(\Sigma, \varphi_{K_2}(\underline{L}), \underline{L}) & \xrightarrow{(I_{H_2})_*} & HF(\Sigma, \varphi_{H_2 \# K_2}(\underline{L}), \varphi_{H_2}(\underline{L})) \end{array}$$

To prove the first diagram, we define a diffeomorphism

$$\begin{aligned} F_{H_1} : \mathbb{R} \times [0, 1] \times \Sigma &\rightarrow \mathbb{R} \times [0, 1] \times \Sigma \\ (s, t, x) &\rightarrow (s, t, \varphi_{H_1}(x)) \end{aligned}$$

Let  $(\mathbb{R} \times [0, 1] \times \Sigma, \Omega_1, \mathcal{L})$  be a Lagrangian cobordism from  $(\varphi_{K_1}(\underline{L}), \underline{L})$  to  $(\varphi_{K_2}(\underline{L}), \underline{L})$ . Note that if  $u \in \mathcal{M}^J(\mathbf{y}_+, \mathbf{y}_-)$  is an HF curve in  $(\mathbb{R} \times [0, 1] \times \Sigma, \Omega_1)$  with Lagrangian boundaries  $\mathcal{L}$ , then  $F_{H_1}(u)$  is a  $F_{H_1}^* J$ -holomorphic HF curve in  $(\mathbb{R} \times [0, 1] \times \Sigma, (F_{H_1}^{-1})^* \Omega_1)$  with Lagrangian boundaries  $F_{H_1}(\mathcal{L})$ . This gives a 1-1 correspondence between the curves in  $(E_1, \Omega_1, \mathcal{L})$  and curves in  $(E_1, (F_{H_1}^{-1})^* \Omega_1, F_{H_1}(\mathcal{L}))$ . Note that  $F_{H_1}(u)$  is a holomorphic curve contributed to the cobordism map  $CF_{A_{ref}}(E_1, (F_{H_1}^{-1})^* \Omega_1, F_{H_1}(\mathcal{L}))$ . Also, the cobordism map induces  $\mathcal{I}_{H_1, H_1}^{H_1 \# K_1, H_1 \# K_2}$ . Hence, the first diagram is true.

To prove the second diagram, the idea is the same. Let  $H_s : [0, 1] \times \Sigma \rightarrow \mathbb{R}$  be a family of Hamiltonian functions such that  $H_s = H_1$  for  $s \geq R_0$  and  $H_s = H_2$  for  $s \leq -R_0$ . Define a diffeomorphism

$$\begin{aligned} F_{\{H_s\}} : \mathbb{R} \times [0, 1] \times \Sigma &\rightarrow \mathbb{R} \times [0, 1] \times \Sigma \\ (s, t, x) &\rightarrow (s, t, \varphi_{H_s}(x)) \end{aligned}$$



Let  $\mathcal{L} = \mathbb{R} \times ((\{0\} \times \varphi_K(\underline{L})) \cup (\{1\} \times \underline{L}))$  be Lagrangians in  $(\mathbb{R} \times [0, 1] \times \Sigma, \Omega = \omega + ds \wedge dt)$ . Then  $F_{\{H_s\}}(\mathcal{L})$  is a disjoint union  $(F_{\{H_s\}}^{-1})^* \Omega$ -Lagrangian submanifolds such that

$$F_{\{H_s\}}(\mathcal{L}) = \begin{cases} \mathbb{R}_{\geq R_0} \times ((\{0\} \times \varphi_{H_1 \# K}(\underline{L})) \cup \{1\} \times \varphi_{H_1}(\underline{L})) & \text{when } s \geq R_0 \\ \mathbb{R}_{\leq -R_0} \times ((\{0\} \times \varphi_{H_2 \# K}(\underline{L})) \cup \{1\} \times \varphi_{H_2}(\underline{L})) & \text{when } s \leq -R_0 \end{cases}$$

Therefore, we define the continuous morphism  $\mathcal{I}_{H_1, H_2}^{H_1 \# K_2, H_2 \# K_2}$  by counting the holomorphic curves in  $(\mathbb{R} \times [0, 1] \times \Sigma, (F_{\{H_s\}}^{-1})^* \Omega, \mathcal{I}_{\{H_s\}}(\mathcal{L}))$ . Similar as the previous case, the map  $F_{\{H_s\}}$  establishes a 1-1 correspondence between the curves in  $(\mathbb{R} \times [0, 1] \times \Sigma, \Omega, \mathcal{L})$  and curves in  $(\mathbb{R} \times [0, 1] \times \Sigma, (F_{\{H_s\}}^{-1})^* \Omega, \mathcal{I}_{\{H_s\}}(\mathcal{L}))$ . This gives us the second diagram.

To see  $(I_{H_1})_* = \mathcal{I}_{H_1, 0}^{H_1 \# K_1, K_1}$ , we just need to take  $K_2 = K_1$  and  $H_2 = 0$  in the diagram.  $\square$

### 3.2.3 Quantum product on HF

Consider  $E_2 = D_2 \times \Sigma$  with a symplectic form  $\Omega_{E_2} = \omega + \omega_{D_2}$ . Take

$$\mathcal{L}_2 = (\partial_1 D_2 \times \varphi_{H_1}(\underline{L})) \cup (\partial_2 D_2 \times \varphi_{H_2}(\underline{L})) \cup (\partial_3 D_2 \times \varphi_{H_3}(\underline{L})).$$

Define  $\mu_2^{H_1, H_2, H_3} := HF_{A_{ref}}(E_2, \Omega_2, \mathcal{L}_2)$ , where  $A_{ref}$  is the reference class in Section 3.2.1. Then  $\mu_2^{H_1, H_2, H_3}$  is a map

$$\mu_2^{H_1, H_2, H_3} : HF(\Sigma, \varphi_{H_1}(\underline{L}), \varphi_{H_2}(\underline{L})) \otimes HF(\Sigma, \varphi_{H_2}(\underline{L}), \varphi_{H_3}(\underline{L})) \rightarrow HF(\Sigma, \varphi_{H_1}(\underline{L}), \varphi_{H_3}(\underline{L})).$$

By Proposition 3.4, we have the following diagram:

$$\begin{array}{ccc} HF(\Sigma, \varphi_{H_1}(\underline{L}), \varphi_{H_2}(\underline{L})) \otimes HF(\Sigma, \varphi_{H_2}(\underline{L}), \varphi_{H_3}(\underline{L})) & \xrightarrow{\mu_2^{H_1, H_2, H_3}} & HF(\Sigma, \varphi_{H_1}(\underline{L}), \varphi_{H_3}(\underline{L})) \\ \downarrow \mathcal{I}_{H_2, K_2}^{H_1, K_1} \otimes \mathcal{I}_{H_3, K_3}^{H_2, K_2} & & \downarrow \mathcal{I}_{H_3, K_3}^{H_1, K_1} \\ HF(\Sigma, \varphi_{K_1}(\underline{L}), \varphi_{K_2}(\underline{L})) \otimes HF(\Sigma, \varphi_{K_2}(\underline{L}), \varphi_{K_3}(\underline{L})) & \xrightarrow{\mu_2^{K_1, K_2, K_3}} & HF(\Sigma, \varphi_{K_1}(\underline{L}), \varphi_{K_3}(\underline{L})) \end{array}$$

Therefore,  $\mu_2^{H_1, H_2, H_3}$  descends to a bilinear map  $\mu_2 : HF(\Sigma, \underline{L}) \otimes HF(\Sigma, \underline{L}) \rightarrow HF(\Sigma, \underline{L})$ .

We call  $\mu_2$  the **quantum product** on QHF.

### 3.2.4 Unit

In this subsection, we define the unit of the quantum product  $\mu_2$ .

Consider the case that  $m = 0$ . Let  $\mathcal{L}_0 \subset \partial E_0 = \partial D_0 \times \Sigma$  be  $d$ -disjoint union of submanifolds such that

$$\mathcal{L}_0|_{s \leq -R_0} = \mathbb{R}|_{s \leq -R_0} \times (\{0\} \times \varphi_H(\underline{L}) \cup \{1\} \times \varphi_K(\underline{L})).$$

Take a symplectic form  $\Omega_0$  such that  $\Omega_0|_{s \leq -R_0} = \omega + ds \wedge dt$  and  $\mathcal{L}_0$  is a union of  $\Omega_0$ -Lagrangians. The tuple  $(E_0, \Omega_0, \mathcal{L}_0)$  can be constructed as follows: First, we take a Lagrangian cobordism  $(E_1, \Omega_1, \mathcal{L}_1)$  from  $(\underline{L}, \underline{L})$  to  $(\varphi_H(\underline{L}), \varphi_K(\underline{L}))$ . Then take  $(E_0, \Omega_0, \mathcal{L}_0)$  to be the composition of  $(E_1, \Omega_1, \mathcal{L}_1)$  and  $(E_0, \omega + \omega_{D_0}, \partial D_0 \times \underline{L})$ .

These data induce a cobordism map

$$HF_{A_{ref}}(E_0, \Omega_0, \mathcal{L}_0) : R \rightarrow HF(\Sigma, \varphi_H(\underline{L}), \varphi_K(\underline{L})).$$

Again,  $A_{ref}$  is the reference class in Section 3.2.1. Define

$$e_{H,K} := HF_{A_{ref}}(E_0, \Omega_0, \mathcal{L}_0)(1).$$

By Proposition 3.4, we have

$$\mathcal{I}_{K_1, K_2}^{H_1, H_2}(e_{H_1, K_1}) = e_{H_2, K_2} \text{ and } \mu_2^{H_1, H_2, H_3}(a \otimes e_{H_2, H_3}) = \mathcal{I}_{H_2, H_3}^{H_1, H_1}(a),$$

where  $a \in HF(\Sigma, \varphi_{H_1}(\underline{L}), \varphi_{H_2}(\underline{L}))$ . These identities imply that the following definition makes sense.

**Definition 3.6.** *The class  $e_{H,K}$  descends to a class  $e \in HF(\Sigma, \underline{L})$ . We call it the **unit**.*

Note that the unit satisfies  $\mu_2(a \otimes e) = a$ .

We now describe the unit when  $H$  is a small Morse function. Fix perfect Morse functions  $f_{L_i} : L_i \rightarrow \mathbb{R}$  with a maximum point  $y_i^+$  and a minimum point  $y_i^-$ . Extend  $\cup_i f_{L_i}$  to be a Morse function  $f : \Sigma \rightarrow \mathbb{R}$  satisfying the following conditions:

**M.1**  $(f, g_\Sigma)$  satisfies the Morse–Smale condition, where  $g_\Sigma$  is a fixed metric on  $\Sigma$ .

**M.2**  $f|_{L_i}$  has a unique maximum  $y_i^+$  and a unique minimum  $y_i^-$ .

**M.3**  $\{y_i^+\}$  are the only maximum points of  $f$ . Also,  $f \leq 0$  and  $f(y_i^+) = 0$  for  $1 \leq i \leq d$ .

**M.4**  $f = f_{L_i} - \frac{1}{2}y^2$  in a neighborhood of  $L_i$ , where  $y$  is the coordinate of the normal direction.

Take  $H = \epsilon f$ , where  $0 < \epsilon \ll 1$ . By Lemma 6.1 in [13], the set of Reeb chords of  $\varphi_H$  is

$$\{\mathbf{y} = [0, 1] \times (y_1, \dots, y_d) \mid y_i \in \text{Crit}(f|_{L_i})\} \quad (3.22)$$

For each  $\mathbf{y} = [0, 1] \times (y_1, \dots, y_d)$ , we construct a relative homology class  $A_{\mathbf{y}}$  as follows: Let  $\eta = \cup_{i=1}^d \eta_i : \oplus_i [0, 1]_s \rightarrow \underline{L}$  be a  $d$ -union of paths in  $\underline{L}$ , where  $\eta_i \subset L_i$  satisfies  $\eta_i(0) = y_i$  and  $\eta_i(1) = x_i$ . Let  $u_i(s, t) = (s, t, \varphi_H \circ (\varphi_H^t)^{-1}(\eta_i(s)))$ . Then  $u = \cup_{i=1}^d u_i$  is a  $d$ -multisection and it gives rise a relative homology class  $A_{\mathbf{y}} \in H_2(E, \mathbf{x}_H, \mathbf{y})$ . It is easy to show that (see Equation (3.18) [13])

$$\mathcal{A}_H(\mathbf{y}, [A_{\mathbf{y}}] + \sum_{i=1}^d c_i [B_i] + m[\Sigma]) = H(\mathbf{y}) - \frac{1}{k+1} \sum_{i=1}^d c_i - m, \quad (3.23)$$

**Lemma 3.7.** *Take  $H = \epsilon f$ . Let  $\mathbf{y}_\heartsuit = [0, 1] \times (y_1^+, \dots, y_d^+)$ . Let  $A_{ref}$  be the reference homology class defined in Section 3.2.1. Then we have a suitable pair  $(\Omega_{E_0}, \mathcal{L}_0)$  such that for a generic  $J \in \mathcal{J}_{comp}(E_0)$ , we have*

$$CF_{A_{ref}}(E_0, \Omega_{E_0}, \mathcal{L}_0)_J(1) = (\mathbf{y}_\heartsuit, [A_{\mathbf{y}_\heartsuit}]).$$

*In particular,  $(\mathbf{y}_\heartsuit, [A_{\mathbf{y}_\heartsuit}])$  is a cycle that represents the unit.*

The idea of the proof is to use index and energy constraint to show that the union of horizontal sections is the only  $I = 0$  holomorphic curve contributed to the cobordism map  $CF_{A_{ref}}(E_0, \Omega_{E_0}, \mathcal{L}_0)_J(1)$ . Since the proof Lemma 3.7 is the same as Lemma 6.6 in [13], we omit the details here.

## 4 Proof of Theorem 1

In this section, we study the properties of the spectral invariants  $c_{\underline{L}}$ . These properties and their proof are parallel to the one in [7, 30].

### 4.1 The HF action spectrum

Fix a base point  $\mathbf{x}$ . Define the **action spectrum** to be

$$Spec(H : \underline{L}, \mathbf{x}) := \{\mathcal{A}_H(\mathbf{y}, [A]) \mid A \in H_2(E, \mathbf{x}_H, \mathbf{y})\}.$$

For different base points  $\mathbf{x}, \mathbf{x}'$ , we have an isomorphism

$$\Psi_{A_{\mathbf{x}, \mathbf{x}'}} : H_2(E, \mathbf{x}_H, \mathbf{y}) \rightarrow H_2(E, \mathbf{x}'_H, \mathbf{y})$$

preserving the action functional (see Equation 3.17 of [13]). In particular, the action spectrum is independent of the base point. So we omit  $\mathbf{x}$  from the notation.

A Hamiltonian function  $H$  is called **mean-normalized** if  $\int_{\Sigma} H_t \omega = 0$  for any  $t$ .

**Definition 4.1.** *Two mean-normalized Hamiltonians  $H^0, H^1$  are said to be homotopic if there exists a smooth path of Hamiltonians  $\{H^s\}_{s \in [0, 1]}$  connecting  $H^0$  to  $H^1$  such that  $H^s$  is normalized and  $\varphi_{H^s} = \varphi_{H^0} = \varphi_{H^1}$  for all  $s$ .*

**Lemma 4.2.** *If two mean-normalized Hamiltonian functions  $H, K$  are homotopic, then we have*

$$Spec(H : \underline{L}) = Spec(K : \underline{L}).$$

*Proof.* Fix a base point  $\mathbf{x} = (x_1, \dots, x_d)$ . Let  $\{\varphi_{s,t} := \varphi_{H^s}^t\}_{s \in [0, 1], t \in [0, 1]}$  be the homotopic such that  $\varphi_{0,t} = \varphi_H^t$ ,  $\varphi_{1,t} = \varphi_K^t$  and  $\varphi_{H^s}^1 = \varphi_H = \varphi_K$  for all  $s \in [0, 1]$ . For fixed

$t$ ,  $\{\varphi_{s,t}\}_{s \in [0,1]}$  is also a family of Hamiltonian symplectomorphisms. Let  $F_t^s$  be the Hamiltonian function in  $s$ -direction, i.e.,

$$X_{F_t^s} = \partial_s \varphi_{s,t} \circ \varphi_{s,t}^{-1}.$$

$F_t^s$  is unique if we require that  $F_t^s$  is mean-normalized. Note that  $X_{F_t^s} = 0$  along  $t = 0, 1$  because  $\varphi_{s,0} = id$  and  $\varphi_{s,1} = \varphi_H = \varphi_K = \varphi$ . By the mean-normalized condition, we have  $F_0^s = F_1^s = 0$ .

Let  $u_i(s, t) = (s, t, \varphi \circ \varphi_{s,t}^{-1}(x_i))$ . Then  $u := \cup_{i=1}^d u_i$  represents a class  $A_0 \in H_2(E, \mathbf{x}_K, \mathbf{x}_H)$ . This induces an isomorphism

$$\Psi_{A_0} : CF(\Sigma, \varphi_H(\underline{L}), \underline{L}, \mathbf{x}) \rightarrow CF(\Sigma, \varphi_K(\underline{L}), \underline{L}, \mathbf{x})$$

by mapping  $(\mathbf{y}, [A])$  to  $(\mathbf{y}, [A_0 \# A])$ .

By a direct computation, we have

$$\begin{aligned} \int u_i^* \omega &= \int_0^1 \int_0^1 \omega(\partial_s \varphi_{s,t}^{-1}(x_i), \partial_t \varphi_{s,t}^{-1}(x_i)) ds \wedge dt \\ &= \int_0^1 \int_0^1 \omega(X_{F_t^s}(x_i), X_{H_t^s}(x_i)) ds \wedge dt = \int_0^1 \int_0^1 \{F_t^s, H_t^s\}(x_i) ds \wedge dt \end{aligned}$$

Because  $H, K$  are mean-normalized, we have  $\partial_s H_t^s - \partial_t F_t^s - \{F_t^s, H_t^s\} = 0$  (see (18.3.17) of [31]). Therefore,

$$\begin{aligned} \int u_i^* \omega &= \int_0^1 \int_0^1 (\partial_s H_t^s(x_i) - \partial_t F_t^s(x_i)) ds \wedge dt \\ &= \int_0^1 H_t^1(x_i) dt - \int_0^1 H_t^0(x_i) dt = \int_0^1 K_t(x_i) dt - \int_0^1 H_t(x_i) dt. \end{aligned}$$

This implies that  $\mathcal{A}_K(\mathbf{y}, [\Psi_{A_0}(A)]) = \mathcal{A}_H(\mathbf{y}, [A])$ . In particular,  $Spec(H : \underline{L}) = Spec(K : \underline{L})$ . □

## 4.2 Proof of Theorem 1

*Proof.* 1. Suppose that  $\varphi_H$  is nondegenerate. Then  $Spec(H : \underline{L})$  is a discrete set over  $\mathbb{R}$ . The spectrality follows directly from the expression (2.15). For the case that  $\varphi_H$  is degenerate, the statement can be deduced from the limit argument in [30].

2. To prove the Hofer-Lipschitz, we first need to construct a Lagrangian cobordism so that we could estimate the energy of holomorphic curves.

Let  $\chi(s) : \mathbb{R}_s \rightarrow \mathbb{R}$  be a non-decreasing cut-off function such that

$$\chi(s) = \begin{cases} 0 & \text{if } s \leq -R_0 \\ 1. & \text{if } s \geq R_0. \end{cases} \quad (4.24)$$

Let  $H^s := \chi(s)H_+ + (1 - \chi(s))H_-$ . Define a diffeomorphism

$$\begin{aligned} F : \mathbb{R} \times [0, 1] \times \Sigma &\rightarrow \mathbb{R} \times [0, 1] \times \Sigma \\ (s, t, x) &\rightarrow (s, t, \varphi_{H^s} \circ (\varphi_{H^s}^t)^{-1}(x)). \end{aligned} \quad (4.25)$$

Let

$$\begin{aligned} \mathcal{L} &:= F(\mathbb{R} \times \{0, 1\} \times \underline{L}) \\ \omega_E &:= (F^{-1})^*(\omega + d(H_t^s dt)) \text{ and } \Omega_E = \omega_E + ds \wedge dt. \end{aligned}$$

Then  $\mathcal{L} \subset \mathbb{R} \times \{0, 1\} \times \Sigma$  is a disjoint union of  $\Omega_E$ -Lagrangians such that

$$\begin{aligned} \mathcal{L}|_{s \geq R_0} &= \mathbb{R}_{s \geq R_0} \times ((\{0\} \times \varphi_{H_+}(\underline{L})) \cup (\{1\} \times \underline{L})) \\ \mathcal{L}|_{s \leq -R_0} &= \mathbb{R}_{s \leq -R_0} \times ((\{0\} \times \varphi_{H_-}(\underline{L})) \cup (\{1\} \times \underline{L})). \end{aligned}$$

Let  $A_{ref} = F(\mathbb{R} \times [0, 1] \times \{\mathbf{x}\}) \in H_2(E_1, \mathbf{x}_{H_+}, \mathbf{x}_{H_-})$ . Take a generic  $J \in \mathcal{J}_{comp}(E_1)$ . Then we have a cobordism  $HF_{A_{ref}}(E_1, \Omega_E, \mathcal{L})_J$  and it is the continuous morphism  $\mathcal{I}_{0,0}^{H_+, H_-}$ . Let  $u \in \mathcal{M}^J(\mathbf{y}_+, \mathbf{y}_-)$  be an HF curve in  $(E_1, \Omega, \mathcal{L})$ .

The energy of  $u$  satisfies

$$\begin{aligned} \int u^* \omega_E &= \int_{F^{-1}(u)} \omega + d_\Sigma H^s \wedge dt + \dot{\chi}(s)(H_+ - H_-) ds \wedge dt \\ &\geq \int_{F^{-1}(u)} \dot{\chi}(s)(H_+ - H_-) ds \wedge dt \\ &\geq d \int_0^1 \min_\Sigma (H_+ - H_-) dt. \end{aligned}$$

The inequality in the second step follows the same argument in Lemma 3.8 of [4].

On the other hand, we have

$$\int_{A_{ref}} \omega_E = \int_{A_+} \omega + \int u^* \omega_E - \int_{A_-} \omega$$

because of the relation  $A_+ \# u \# (-A_-) = A_{ref}$ . Note that

$$\int_{A_{ref}} \omega_E = \int_0^1 H_+(t, \mathbf{x}) dt - \int_0^1 H_-(t, \mathbf{x}) dt.$$

Then we have

$$\int u^* \omega_E = \mathcal{A}_{H_+}(\mathbf{y}_+, A_+) - \mathcal{A}_{H_-}(\mathbf{y}_-, A_-).$$

Therefore,  $\mathcal{A}_{H_+}(\mathbf{y}_+, A_+) - \mathcal{A}_{H_-}(\mathbf{y}_-, A_-) \geq d \int_0^1 \min_{\Sigma}(H_+ - H_-) dt..$

Fix  $a \neq 0 \in HF(\Sigma, \underline{L})$ . Take a cycle  $\mathbf{c}_+ \in CF(\Sigma, \varphi_{H_+}(\underline{L}), (\underline{L}))$  represented  $(j_{H_+}^{\mathbf{x}})^{-1}(a)$  and satisfying

$$\mathcal{A}_H(\mathbf{c}_+) \leq c_{\underline{L}}(H_+, a) + \delta.$$

Let  $\mathbf{c}_- = \mathcal{I}_{0,0}^{H_+, H_-}(\mathbf{c}_+)$ . Then it is a cycle represented  $(j_{H_-}^{\mathbf{x}})^{-1}(a)$ . Take a factor  $(\mathbf{y}_-, [A_-])$  in  $\mathbf{c}_-$  such that  $\mathcal{A}_H(\mathbf{y}_-, [A_-]) = \mathcal{A}_H(\mathbf{c}_-)$ . Find a factor  $(\mathbf{y}_+, [A_+])$  in  $\mathbf{c}_+$  such that  $\langle \mathcal{I}_{0,0}^{H_+, H_-}(\mathbf{y}_+, [A_+]), (\mathbf{y}_-, [A_-]) \rangle = 1$ . Then the above discussion implies that

$$\delta + c_{\underline{L}}(H_+, a) - c_{\underline{L}}(H_-, a) \geq d \int_0^1 \min_{\Sigma}(H_+ - H_-) dt.$$

Take  $\delta \rightarrow 0$ . Interchange the positions of  $H_+$  and  $H_-$ ; then we obtain the Hofer-Lipschitz property.

3. Since  $H$  and  $K$  are homotopic, we have a family of Hamiltonian functions  $\{H_t^s\}_{s \in [0,1]}$  with  $H_t^0 = H_t$  and  $H_t^1 = K_t$ . By Lemma 4.2, we have

$$Spec(H : \underline{L}) = Spec(H^s : \underline{L}) = Spec(K : \underline{L}).$$

On the other hand,  $c_{\underline{L}}(H^s, a)$  is continuous with respect to  $s$ . Therefore, it must be constant.

4. Define a family of functions  $H^s = H + sc$ . Note that  $\mathcal{A}_{H+sc}(\mathbf{y}, A) = \mathcal{A}_H(\mathbf{y}, A) + s \int_0^1 c(t) dt$ . Therefore the spectral  $Spec(H^s : \underline{L})$  is  $Spec(H : \underline{L})$  translated by  $s \int_0^1 c(t) dt$ . As a result,  $c_{\underline{L}}(H^s, a) - s \int_0^1 c(t) dt \in Spec(H : \underline{L})$ . Then the shift property follows directly from Hofer-Lipschitz property.
5. If  $H_t|_{L_i} = c_i(t)$ , then  $\varphi_H(L_i) = L_i$ . The Reeb chords are corresponding to  $\mathbf{y} \in \underline{L}$ . Hence,  $Spec(H : \underline{L}) = \{\frac{m}{k+1} + \sum_{i=1}^d \int_0^1 c_i(t) dt | m \in \mathbb{Z}\}$ . Define a family of Hamiltonians functions  $\{H^s := sH\}_{s \in [0,1]}$ . By the spectrality, we have  $c_{\underline{L}}(H^s, a) = \frac{m_0}{k+1} + \sum_{i=1}^d \int_0^1 s c_i(t) dt$ . Here  $m_0$  is a constant due to the Hofer-Lipschitz continuity. We know that  $\frac{m_0}{k+1} = c_{\underline{L}}(0, a)$  by taking  $s = 0$ . Then the Lagrangian control property follows from taking  $s = 1$ .
6. Let  $a, b \in HF(\Sigma, \underline{L})$ . Take

$$\Omega_2 = \omega + \omega_{D_2}$$

$$\mathcal{L}_2 = (\partial_1 D_2 \times \varphi_H \circ \varphi_K(\underline{L})) \cup (\partial_2 D_2 \times \varphi_H(\underline{L})) \cup (\partial_3 D_2 \times \underline{L}).$$

Then  $\mu_2 : HF(\Sigma, \varphi_H \circ \varphi_K(\underline{L}), \varphi_H(\underline{L})) \otimes HF(\Sigma, \varphi_H(\underline{L}), \underline{L}) \rightarrow HF(\Sigma, \varphi_H \circ \varphi_K(\underline{L}), \underline{L})$ .

Let us first consider the following special case: Suppose that there is a base point  $\mathbf{x} = (x_1, \dots, x_d) \in \underline{L}$  such that

$$\begin{aligned} d_\Sigma H_t(x_i) &= d_\Sigma K_t(x_i) = 0 \\ \nabla^2 H_t(x_i), \nabla^2 K_t(x_i) &\text{ are non-degenerate.} \end{aligned} \quad (4.26)$$

for  $1 \leq i \leq d$ . This assumption implies that  $\varphi_H^t(x_i) = x_i$ ,  $\varphi_K^t(x_i) = x_i$  and  $d_\Sigma(H_t \diamond K_t)(x_i) = 0$ . In particular,  $\mathbf{x}$  is a non-degenerate Reeb chord of  $\varphi_H$ ,  $\varphi_K$  and  $\varphi_H \circ \varphi_K$ . Also, the reference chords become  $\mathbf{x}_H = \varphi_H(\mathbf{x}_K) = \mathbf{x}_{H \diamond K} = \mathbf{x}$ .

Take  $A_{ref} = [D_2 \times \{\mathbf{x}\}] \in H_2(\varphi_H(\mathbf{x}_K), \mathbf{x}_H, \mathbf{x}_{H \diamond K})$  be the reference homology class. Obviously, we have

$$\int_{A_{ref}} \omega = \int_{D_2 \times \{\mathbf{x}\}} \omega = 0. \quad (4.27)$$

Let  $u \in \mathcal{M}^J((\mathbf{y}_1, [A_1]) \otimes (\mathbf{y}_2, [A_2]), (\mathbf{y}_0, [A_0]))$  be an HF curve with  $I = 0$ . Here the relative homology classes satisfy  $A_1 \# A_2 \# u \# (-A_0) = A_{ref}$ . Therefore, the energy of  $u$  is

$$\int u^* \omega = - \int_{A_1} \omega - \int_{A_2} \omega + \int_{A_0} \omega + \int_{A_{ref}} \omega. \quad (4.28)$$

Take  $J \in \mathcal{J}_{comp}(E_2)$ . Then  $\int u^* \omega \geq 0$ . Combine this with (4.27), (4.28); then we have

$$\mathcal{A}_{H \diamond K}(\mathbf{y}_0, [A_0]) \leq \mathcal{A}_H(\mathbf{y}_1, [A_1]) + \mathcal{A}_K(\varphi_H^{-1}(\mathbf{y}_2), [\varphi_H^{-1}(A_2)]). \quad (4.29)$$

Assume that  $\mu_2(a \otimes b) \neq 0$ . Let  $\mathbf{c}_0 \in CF(\Sigma, \varphi_H \circ \varphi_K(\underline{L}), \underline{L})$ ,  $\mathbf{c}_1 \in CF(\Sigma, \varphi_H(\underline{L}), \underline{L})$  and  $\mathbf{c}_2 \in CF(\Sigma, \varphi_H \circ \varphi_K(\underline{L}), \varphi_H(\underline{L}))$  be cycles represented  $j_{H \diamond K}^{-1}(\mu_2(a \otimes b))$ ,  $j_H^{-1}(a)$  and  $j_{H \diamond K, H}^{-1}(b)$  respectively. By Lemma 3.5,  $\varphi_H^{-1}(\mathbf{c}_2)$  is a cycle represented  $j_K^{-1}(b)$ . We choose  $\mathbf{c}_1, \mathbf{c}_2$  such that

$$\begin{aligned} \mathcal{A}_H(\mathbf{c}_1) &\leq c_{\underline{L}}(H, a) + \delta \\ \mathcal{A}_K(\varphi_H^{-1}(\mathbf{c}_2)) &\leq c_{\underline{L}}(K, b) + \delta. \end{aligned}$$

Therefore, (4.29) implies that  $\mathcal{A}_{H \diamond K}(\mathbf{c}_0) \leq \mathcal{A}_H(\mathbf{c}_1) + \mathcal{A}_K(\varphi_H^{-1}(\mathbf{c}_2)) + 2\delta$ . Take  $\delta \rightarrow 0$ . We have

$$c_{\underline{L}}(H \diamond K, \mu_2(a \otimes b)) \leq c_{\underline{L}}(H, a) + c_{\underline{L}}(K, b).$$

For general Hamiltonians  $H_t, K_t$ , we first construct approximations  $H_t^\delta, K_t^\delta$  satisfying the assumptions (4.26).

Fix local coordinates  $(x, y)$  around  $x_i$ . Then we can write

$$H_t(x, y) = H_t(0) + \partial_x H_t(0)x + \partial_y H_t(0)y + R_t(x, y).$$

We may assume that  $\nabla^2 H_t(0)$  is non-degenerate; otherwise, we can achieve this by perturbing  $H_t$  using a small Morse function with a critical point at  $x_i$ .

Pick a cut-off function  $\chi_\delta(r) : \mathbb{R}_+ \rightarrow \mathbb{R}$  such that  $\chi_\delta(0) = 1$ ,  $\chi'_\delta(0) = 0$  and  $\chi_\delta = 0$  for  $r \geq \delta$ , where  $r = \sqrt{x^2 + y^2}$ . Define  $H_t^\delta$  by

$$H_t^\delta(x, y) = H_t(0) + (1 - \chi_\delta(r))(\partial_x H_t(0)x + \partial_y H_t(0)y) + R_t(x, y).$$

We perform the same construction for  $K_t$ . Apparently, we have

$$\begin{aligned} d_\Sigma H_t^\delta(x_i) &= d_\Sigma K_t^\delta(x_i) = 0, \\ \nabla^2 H_t^\delta(x_i) &= \nabla^2 H_t(x_i), \quad \nabla^2 K_t^\delta(x_i) = \nabla^2 K_t(x_i), \\ |H_t^\delta - H_t| &\leq c_0 \delta, \quad |K_t^\delta - K_t| \leq c_0 \delta, \\ |H_t \diamond K_t - H_t^\delta \diamond K_t^\delta| &\leq c_0 \delta, \end{aligned}$$

By the Hofer-Lipschitz continuity, we have

$$c_{\underline{L}}(H \diamond K, \mu_2(a \otimes b)) \leq c_{\underline{L}}(H, a) + c_{\underline{L}}(K, b) + O(\delta).$$

Note that the above construction works for any  $\delta$ , we can take  $\delta \rightarrow 0$ .

Since the normalization of  $H \diamond K$  and  $H \# K$  are homotopic, we can replace  $H \diamond K$  in the triangle equality by  $H \# K$ .

7. By the triangle inequality, we have

$$c_{\underline{L}}(0, e) = c_{\underline{L}}(0, \mu_2(e \otimes e)) \leq c_{\underline{L}}(0, e) + c_{\underline{L}}(0, e).$$

Hence, we get  $c_{\underline{L}}(0, e) \geq 0$ . On the other hand, Lemma 3.7 and (2.15) imply that  $c_{\underline{L}}(0, e) \leq 0$ .

8. The proof of the Calabi property relies on the Hofer-Lipschitz and the Lagrangian control properties. We have obtained these properties. One can follow the same argument in [7] to prove the Calabi property. We skip the details here.  $\square$

## 5 Open-closed morphisms

In this section, we prove Theorem 2. Most of the arguments here are parallel to [15] and the counterparts of the closed-open morphisms [13]. Therefore, we will just outline



the construction of the open-closed morphisms and the proof of partial invariance. We will focus on proving the non-vanishing of the open-closed morphisms.

To begin with, let us introduce the open-closed symplectic manifold and the Lagrangian submanifolds. The construction follows [15]. Define a base surface  $B \subset \mathbb{R}_s \times (\mathbb{R}_t/(2\mathbb{Z}))$  by  $B := \mathbb{R}_s \times (\mathbb{R}_t/(2\mathbb{Z})) - B^c$ , where  $B^c$  is  $(2, \infty)_s \times [1, 2]_t$  with the corners rounded. See Figure 3.




Figure 3: The open-closed surface

Let  $Y_{\varphi_H} := [0, 2] \times \Sigma/(0, \varphi_H(x)) \sim (2, x)$  be the mapping torus of  $\varphi_H$ . Then  $\pi : \mathbb{R}_s \times Y_{\varphi_H} \rightarrow \mathbb{R}_s \times (\mathbb{R}_t/(2\mathbb{Z}))$  is a surface bundle over the cylinder. Define a surface bundle  $W_H$  by

$$\pi_W = \pi|_W : W_H := \pi^{-1}(B) \rightarrow B.$$

The symplectic form  $\Omega_H$  on  $W_H$  is defined to be the restriction of  $\omega_{\varphi_H} + ds \wedge dt$ . Note that  $W_H$  is diffeomorphic (preserving the fibration structure) to the  $B \times \Sigma$ . So we denote  $W_H$  by  $W$  instead when the context is clear.

We place a copy of  $\underline{L}$  on the fiber  $\pi_W^{-1}(3, 1)$  and take its parallel transport along  $\partial B$  using the symplectic connection. The parallel transport sweeps out an  $\Omega_H$ -Lagrangian submanifold  $\mathcal{L}_H$  in  $W$ . Then  $\mathcal{L}_H$  consists of  $d$ -disjoint connected components. Moreover, we have

$$\begin{aligned} \mathcal{L}_H|_{s \geq 3 \times \{0\}} &= \mathbb{R}_{s \geq 3} \times \{0\} \times \varphi_H(\underline{L}) \\ \mathcal{L}_H|_{s \geq 3 \times \{1\}} &= \mathbb{R}_{s \geq 3} \times \{1\} \times \underline{L}. \end{aligned}$$

We call the triple  $(W_H, \Omega_H, \mathcal{L}_H)$  an **open-closed cobordism**.

**Definition 5.1** (Definition 5.4.3 of [15]). *Fix a Reeb chord  $\mathbf{y}$  and an orbit set  $\gamma$  with degree  $d$ . Let  $(\dot{F}, j)$  be a Riemann surface (possibly disconnected) with punctures. A  $d$ -multisection in  $W$  is a smooth map  $u : (\dot{F}, \partial\dot{F}) \rightarrow W$  such that*

1.  $u(\partial\dot{F}) \subset \mathcal{L}_H$ . Write  $\mathcal{L}_H = \cup_{i=1}^d L_H^i$ , where  $L_H^i$  is a connected component of  $\mathcal{L}_H$ . For each  $1 \leq i \leq d$ ,  $u^{-1}(L_H^i)$  consists of exactly one component of  $\partial\dot{F}$ .

2.  $u$  is asymptotic to  $\mathbf{y}$  as  $s \rightarrow \infty$ .
3.  $u$  is asymptotic to  $\gamma$  as  $s \rightarrow -\infty$ .
4.  $\int_F u^* \omega_{\varphi_H} < \infty$ .

A  $J$ -holomorphic  $d$ -multisection is called an **HF-PFH curve**. We remark that the HF-PFH curves are simple because they are asymptotic to Reeb chords. This observation is crucial in the proof of Theorem 2.

Let

$$Z_{\mathbf{y}, \gamma} := \mathcal{L}_H \cup (\{\infty\} \times \mathbf{y}) \cup (\{-\infty\} \times \gamma) \subset W.$$

We denote  $H_2(W, \mathbf{y}, \gamma)$  the equivalence classes of continuous maps  $u : (\dot{F}, \partial \dot{F}) \rightarrow (W, Z_{\mathbf{y}, \gamma})$  satisfying 1), 2), 3) in the above definition. Two maps are equivalent if they represent the same element in  $H_2(W, Z_{\mathbf{y}, \gamma}; \mathbb{Z})$ . Note that  $H_2(W, \mathbf{y}, \gamma)$  is an affine space of  $H_2(W, \mathcal{L}_H; \mathbb{Z})$ . The difference of any two classes can be written as

$$\mathcal{Z}' - \mathcal{Z} = \sum_{i=1}^k c_i [B_i] + m[\Sigma] + [S],$$

where  $[B_i]$  is the class represented by the parallel translation of  $B_i$  and  $[S]$  is a class in the  $H_1(S^1, \mathbb{Z}) \otimes H_1(\Sigma, \mathbb{Z})$ -component of  $H_2(Y_{\varphi_H}, \mathbb{Z})$ .

Fix a nonvanishing vector field on  $\underline{L}$ . This gives a trivialization  $\tau$  of  $T\Sigma|_{\underline{L}}$ . We extend it to  $T\Sigma|_{\mathcal{L}_H}$  by using the symplectic parallel transport. We then extend the trivialization of  $T\Sigma|_{\mathcal{L}_H}$  in an arbitrary manner along  $\{\infty\} \times \mathbf{y}$  and along  $\{-\infty\} \times \gamma$ . Then we can define the relative Chern number  $c_1(u^*T\Sigma, \tau)$ . This is the obstruction of extending  $\tau$  to  $u$ .

Define a real line bundle  $\mathfrak{L}$  of  $T\Sigma$  along  $\mathcal{L}_H \cup \{\infty\} \times \mathbf{y}$  as follows. We set  $\mathfrak{L}|_{\mathcal{L}_H} := T\mathcal{L}_H \cap T\Sigma$ . Then extend  $\mathfrak{L}$  across  $\{\infty\} \times \mathbf{y}$  by rotating in the counterclockwise direction from  $T\varphi_H(\underline{L})$  to  $T\underline{L}$  in  $T\Sigma$  by the minimum amount. With respect to the trivialization  $\tau$ , we have Maslov index for the bundle pair  $(u^*\mathfrak{L}, u^*T\Sigma)$ , denoted by  $\mu_\tau(u)$ .

The **Fredholm index** for an HF-PFH curve is

$$\text{indu} := -\chi(\dot{F}) - d + 2c_1(u^*T\Sigma, \tau) + \mu_\tau(u) - \mu_\tau^{\text{ind}}(\gamma).$$

The notation  $\mu_\tau^{\text{ind}}(\gamma)$  is explained as follows. Let  $\gamma = \{(\gamma_i, m_i)\}$ . Suppose that for each  $i$ ,  $u$  has  $k_i$ -negative ends and each end is asymptotic to  $\gamma_i^{q_j}$ . Then the total multiplicity is  $m_i = \sum_{j=1}^{k_i} q_j$ . Define

$$\mu_\tau^{\text{ind}}(\gamma) := \sum_i \sum_{j=1}^{k_i} CZ_\tau(\gamma_i^{q_j}),$$

where  $CZ_\tau$  is the Conley-Zehnder index.

Given  $\mathcal{Z} \in H_2(W, \mathbf{y}, \gamma)$ , the **ECH index** is (see Definition 5.6.5 of [15])

$$I(\mathcal{Z}) := c_1(TW|_{\mathcal{Z}}, \tau) + Q_\tau(\mathcal{Z}) + \mu_\tau(\mathcal{Z}) - \mu_\tau^{ech}(\gamma) - d,$$

where  $\mu_\tau^{ech}(\gamma) := \sum_i \sum_{p=1}^{m_i} CZ_\tau(\gamma_i^p)$ .

**Theorem 5.2.** (Theorem 5.6.9 of [15]) *Let  $u$  be an HF-PFH curve in  $(W_H, \Omega_H, \mathcal{L}_H)$ . Then we have*

$$indu + 2\delta(u) \leq I(u).$$

Moreover, the equality holds if and only if  $u$  satisfies the ECH partition condition.

In this paper, we don't need the details on "ECH partition condition". For the readers who are interested in it, please refer to [18, 19].

## 5.1 Construction and invariance of $\mathcal{OC}$

In this subsection, we outline the construction of the open-closed morphisms. Also, we will explain why it satisfies the partial invariance.

To begin with, we need the following lemma to rule out the bubbles.

**Lemma 5.3.** *Let  $\mathcal{Z}', \mathcal{Z} \in H_2(W, \mathbf{y}, \gamma)$  be relative homology classes such that  $\mathcal{Z}' - \mathcal{Z} = m[\Sigma] + \sum_{i=1}^k c_i[B_i] + [S]$ , where  $[S] \in H_1(S^1, \mathbb{Z}) \otimes H_1(\Sigma, \mathbb{Z})$ . Then we have*

$$I(\mathcal{Z}') = I(\mathcal{Z}) + \sum_i^k 2c_i + 2m(k+1). \quad (5.30)$$

*Proof.* Using the same argument in Lemma 3.3 of [13], we know that adding  $B_i$  to a relative homology class  $\mathcal{Z}$  will increase the ECH index by 2 because the Maslov index of  $B_i$  is 2.

If we add  $[\Sigma]$  to  $\mathcal{Z}$ , then the ECH index will increase by  $2(k+1)$ . See the index ambiguity formula in Proposition 1.6 of [18]. Similarly, adding  $[S]$  doesn't change the ECH index.  $\square$

Fix a reference relative homology class  $\mathcal{Z}_{ref} \in H_2(W, \mathbf{x}_H, \gamma_{ref})$ . The open-closed morphism in chain level is defined by

$$\mathcal{OC}_{\mathcal{Z}_{ref}}(W_H, \Omega_H, \mathcal{L}_H)_J(\mathbf{y}, [A]) = \sum_{(\gamma, [Z])} \sum_{\mathcal{Z}, I(\mathcal{Z})=0} \#\mathcal{M}^J(\mathbf{y}, \gamma, \mathcal{Z})(\gamma, [Z]),$$

The class  $Z$  is characterized by  $A\#\mathcal{Z}\#Z = \mathcal{Z}_{ref}$ . The arguments in [15] (also see the relevant argument for closed-open morphisms [13]) show that this is well defined and it is a chain map. The main difference here with [15] is that the bubbles would appear, but these can be ruled out by Lemma 5.3 and the argument in Proposition 3.4.

To prove the partial invariance, the arguments consist of the following key steps:

1. If we deform the open-closed morphism smoothly over a compact set of  $W$  (the deformation needs to be generic), then the standard homotopy arguments show that the open-closed morphism is invariant.
2. Assume that  $\varphi_H$  satisfies  $\spadesuit.1$  and  $\varphi_G$  satisfies  $\spadesuit.2$ . Let  $(E_1, \Omega_1, \mathcal{L}_1)$  be a Lagrangian cobordism from  $(\varphi_G(\underline{L}), \underline{L})$  to  $(\varphi_H(\underline{L}), \underline{L})$ . Let  $(X, \Omega_X)$  be a symplectic cobordism from  $(Y_{\varphi_H}, \omega_{\varphi_H})$  to  $(Y_{\varphi_G}, \omega_{\varphi_G})$  defined by (2.10). Consider the  $R$ -stretched composition of  $(E_1, \Omega_1, \mathcal{L}_1)$ ,  $(W_H, \Omega_H, \mathcal{L}_H)$  and  $(X, \Omega_X)$ , denoted by  $(W_R, \Omega_R, \mathcal{L}_R)$ . As  $R \rightarrow \infty$ , the  $I = 0$  HF-PFH curves in  $(W_R, \Omega_R, \mathcal{L}_R)$  converges to a holomorphic building. Under assumptions  $\spadesuit.1$ ,  $\spadesuit.2$ , the holomorphic curves in  $(X, \Omega_X)$  have nonnegative ECH index (see Section 7.1 of [11]). Combining this with Theorems 3.3, 5.2, the holomorphic curves in each level have nonnegative ECH index. As a result, these holomorphic curves have zero ECH index. They are either embedded or branched covers of trivial cylinders. By Huctings-Taubes's gluing argument [24, 25], the open-closed morphism defined by  $(W_R, \Omega_R, \mathcal{L}_R)$  is equal to  $I_{0,0}^{G,H} \circ (\mathcal{OC}_{Z_{ref}}(W_H, \Omega_H, \mathcal{L}_H))_* \circ PFH_{Z_{ref}}^{hol}(X, \Omega_X)$  for  $R \gg 1$ . Here  $PFH_{Z_{ref}}^{hol}(X, \Omega_X)$  is the PFH cobordism map defined by counting embedded holomorphic curves in  $X$ . By Theorem 3 in [11], we can replace it by  $PFH_{Z_{ref}}^{sw}(X, \Omega_X)$ . Finally, by the homotopy invariance in step 1, we get the partial invariance.

For more details, we refer the readers to [13].

## 5.2 Computations of $\mathcal{OC}$

In this subsection, we compute the open-closed morphism for a special Hamiltonian function  $H$  satisfying  $\spadesuit.1$ . Using partial invariance, we deduce the non-vanishing result under the assumption  $\spadesuit.2$ . The main idea here is also the same as [13].

Suppose that  $f$  is a Morse function satisfying **M.1**, **M.2**, **M.3**, and **M.4**. Define  $H_\epsilon = -\epsilon f$ , where  $0 < \epsilon \ll 1$ .  $H_\epsilon$  is a slight perturbation of the height function in Figure 1. This is a nice candidate for computation because we can describe the periodic orbits and Reeb chords in terms of the critical points and the index of holomorphic curves are computable. However, the  $H_\epsilon$  doesn't satisfy  $\spadesuit.1$  or  $\spadesuit.2$ . We need to follow the discussion in Section 6.1 of [13] to modify  $H_\epsilon$ .

Fix numbers  $0 < \epsilon_0 \ll 1$  and  $\delta, \delta_0 > 0$ . By [13], we have a function  $\varepsilon : \Sigma \rightarrow \mathbb{R}$  such that  $0 < \epsilon \leq \varepsilon \leq \epsilon_0$  and the new autonomous Hamiltonian function  $H_\varepsilon = -\varepsilon f$  satisfies the following conditions:

- F.1** There is an open set  $\mathcal{U}^{\delta+\delta_0} = \cup_p U_p^{\delta+\delta_0}$  such that  $H_\varepsilon|_{\Sigma-\mathcal{U}^{\delta+\delta_0}} = H_\epsilon|_{\Sigma-\mathcal{U}^{\delta+\delta_0}}$ , where  $p$  runs over all the local maximums of  $-f$  and  $U_p^{\delta+\delta_0}$  is a  $(\delta + \delta_0)$ -neighbourhood of  $p$ .

**F.2**  $H_\varepsilon$  is still a Morse function satisfying the Morse-Smale conditions. Moreover,  $Crit(H_\varepsilon) = Crit(-f)$ .

**F.3**  $\varphi_{H_\varepsilon}$  is  $d$ -nondegenerate. The periodic orbits of  $\varphi_{H_\varepsilon}$  with period at most  $d$  are covers of the constant orbits at critical points of  $H_\varepsilon$ .

**F.4** For each local maximum  $p$ ,  $\varphi_{H_\varepsilon}$  has a family periodic orbits  $\gamma_{r_0, \theta}(t)$  that foliates  $S_t^1 \times \partial \mathcal{U}_p^{r_0}$ , where  $\delta + \delta_0 \leq r_0 \leq \delta + 2\delta_0$ . Moreover, the period of  $\gamma_{r_0, \theta}(t)$  is strictly greater than  $d$ .

**F.5** The Reeb chords of  $\varphi_{H_\varepsilon}$  are still corresponding to the critical points of  $\cup_{i=1}^d f_{L_i}$ . See (3.22).

By Proposition 3.7 of [11], we perturb  $H_\varepsilon$  to a new Hamiltonian function  $H'_\varepsilon$  (may depend on  $t$ ) such that it satisfies the following properties:

1.  $H'_\varepsilon|_{\Sigma - \mathcal{U}^\delta} = H_\varepsilon|_{\Sigma - \mathcal{U}^\delta}$ .
2.  $H'_\varepsilon$  still satisfies **F.4** and **F.5**.
3.  $|H'_\varepsilon - H_\varepsilon| \leq c_0\delta$  and  $|dH'_\varepsilon - dH_\varepsilon| \leq c_0\delta$ .
4. The periodic orbits of  $\varphi_{H'_\varepsilon}$  with period less than or equal to  $d$  are either hyperbolic or  $d$ -negative elliptic. In other words,  $\varphi_{H'_\varepsilon}$  is  $d$ -nondegenerate and satisfies  $\spadesuit.1$ .

**Remark 5.1.** *Because we take  $H_\varepsilon = -\varepsilon f$ , the maximum points  $\{y_i^+\}$  of  $f$  are the minimum points of  $H_\varepsilon$ . We use  $\{y_i^-\}$  to denote the minimum points of  $H_\varepsilon$  from now on.*

Let  $y$  be a critical point of  $H_\varepsilon$ . Let  $\gamma_y$  denote the constant simple periodic orbit at the critical point  $y$ . We define PFH generators and a Reeb chord as follows:

1. Let  $I = (i_1, \dots, i_d)$ . Here we allow  $i_j = i_k$  for  $j \neq k$ . Let  $\alpha_I = \gamma_{y_{i_1}^-} \dots \gamma_{y_{i_d}^-}$ . When  $I = (1, 2, \dots, d)$ , we denote  $\alpha_I$  by  $\alpha_\diamond$ . Here we use multiplicative notation to denote an orbit set instead.
2.  $\mathbf{y}_\diamond = [0, 1] \times (y_-^1, \dots, y_-^d)$ .

Let  $\alpha = \gamma_{p_1} \dots \gamma_{p_d}$  and  $\beta = \gamma_{q_1} \dots \gamma_{q_d}$  be two orbit sets, where  $p_i, q_j \in Crit(H_\varepsilon)$ . Following [13], we define a relative homology class  $Z_{\alpha, \beta}$  as follows: Let  $\eta = \sqcup_{i=1}^d \eta_i : \sqcup_{i=1}^d [0, 1] \rightarrow \Sigma$  be a union of paths with  $d$  components such that  $\eta_i(1) = p_i$  and  $\eta_i(0) = q_i$ . Define a relative homology class

$$Z_{\alpha, \beta} := [S^1 \times \eta] \in H_2(Y_{\varphi_{H_\varepsilon}}, \alpha, \beta). \quad (5.31)$$

We also use this way to define  $Z_\alpha \in H_2(Y_{\varphi_{H_\varepsilon}}, \alpha, \gamma_H^\mathbf{x})$ .

Let  $\mathcal{J}(W, \Omega_{H'_\varepsilon}) \subset \mathcal{J}_{tame}(W, \Omega_{H'_\varepsilon})$  be a set of almost complex structures which are the restriction of admissible almost complex structures in  $\mathcal{J}(Y_{\varphi_{H'_\varepsilon}}, \omega_{\varphi_{H'_\varepsilon}})$ .

Take a  $J \in \mathcal{J}(W, \Omega_{H'_\varepsilon})$ . Let  $u_{y_i}$  be the restriction of  $\mathbb{R} \times \gamma_{y_i}$  to  $W$ . Obviously, it is a  $J$ -holomorphic curve in  $\mathcal{M}^J(y_i, \gamma_{y_i})$ . It is called a **horizontal section** of  $(W, \Omega_{H'_\varepsilon}, \mathcal{L}_{H'_\varepsilon}, J)$ . Moreover, it is easy to check that  $ind u_{y_i} = 0$  from the definition.

**Lemma 5.4.** *Let  $u : F \rightarrow W$  be a  $J$ -holomorphic HF-PFH curve in  $(W, \Omega_H, \mathcal{L}_H)$  and  $J \in \mathcal{J}(W, \Omega_H)$ . Then*

$$E_{\omega_{\varphi_H}}(u) = \int_F u^* \omega_{\varphi_H} \geq 0.$$

Moreover, when  $H = H'_\varepsilon$ ,  $E_{\omega_{\varphi_H}}(u) = 0$  if and only if  $u$  is a union of the horizontal sections.

*Proof.* The proof is the same as Lemma 6.6 in [13]. □

The horizontal sections  $\cup_{i=1}^d u_{y_i}$  represent a relative homology class  $\mathcal{Z}_{hor}$ . We take the reference relative homology class to be  $\mathcal{Z}_{ref} = A_{\mathbf{y}_\diamond} \# \mathcal{Z}_{hor} \# Z_{\alpha_\diamond} \in H_2(W, \mathbf{x}_H, \gamma_H^\mathbf{x})$ .

**Lemma 5.5.** *For a generic  $J \in \mathcal{J}(W, \Omega_{H'_\varepsilon})$ , we have*

$$\mathcal{OC}_{\mathcal{Z}_{ref}}(W, \Omega_{H'_\varepsilon}, \mathcal{L}_{H'_\varepsilon})_J(\mathbf{y}_\diamond, A_{\mathbf{y}_\diamond}) = (\alpha_\diamond, Z_{\alpha_\diamond}) + \sum (\beta, Z),$$

Here  $(\beta, Z)$  satisfies  $\beta \neq \alpha_\diamond$  and  $\mathbb{A}_{H'_\varepsilon}(\alpha_\diamond, Z_{\alpha_\diamond}) > \mathbb{A}_{H'_\varepsilon}(\beta, Z)$ .

*Proof.* Consider the moduli space of HF-PFH curves  $\mathcal{M}_0^J(\mathbf{y}_\diamond, \alpha_\diamond)$  with  $I = 0$ . Let  $u \in \mathcal{M}_0^J(\mathbf{y}_\diamond, \alpha_\diamond)$ . By Lemma 5.3,  $I(u) = 0$  implies that  $\sum_i c_i + m(k+1) = 0$ .

On the other hand,  $E_{\omega_{H'_\varepsilon}}(u) = \int |d^{vert} u|^2 = \sum \frac{c_i}{k+1} + m = 0$ . Lemma 5.4 implies that  $u = \cup_i u_{y_i}$  is a union of horizontal sections. In other words, the union of horizontal sections is the unique element in  $\mathcal{M}_0^J(\mathbf{y}_\diamond, \alpha_\diamond)$ .

If  $u$  is an HF-PFH curve in  $\mathcal{M}_0^J(\mathbf{y}_\diamond, \beta)$  and  $\beta \neq \alpha_\diamond$ , then  $E_{\omega_{\varphi_{H'_\varepsilon}}}(u) > 0$ ; otherwise,  $u$  is horizontal and  $u$  must be asymptotic to  $\alpha_\diamond$ . Note that

$$\begin{aligned} 0 < E_{\omega_{H'_\varepsilon}}(u) &= \int_{\mathcal{Z}_{ref}} \omega_{\varphi_{H'_\varepsilon}} - \int_{A_{\mathbf{y}_\diamond}} \omega - \int_Z \omega_{\varphi_{H'_\varepsilon}} \\ &= \int_{Z_{\alpha_\diamond}} \omega_{\varphi_{H'_\varepsilon}} - \int_Z \omega_{\varphi_{H'_\varepsilon}} = \mathbb{A}_{H'_\varepsilon}(\alpha_\diamond, Z_{\alpha_\diamond}) - \mathbb{A}_{H'_\varepsilon}(\beta, Z). \end{aligned}$$

□

The following lemma tells us that  $(\mathbf{y}_\diamond, A_{\mathbf{y}_\diamond})$  is a cycle.

**Lemma 5.6.** *Let  $d_J$  be the differential of  $CF(\Sigma, \varphi_{H'_\varepsilon}(\underline{L}), \underline{L})$ . Then  $d_J = 0$ . In particular,  $(\mathbf{y}_\diamond, A_{\mathbf{y}_\diamond})$  is a cycle.*

*Proof.* By **F.5**, we know that  $CF(\Sigma, \varphi_{H'_\varepsilon}(\underline{L}), \underline{L}) \cong \oplus^{2^d} R$ . According to Lemma 6.8 in [7], we know that

$$CF(\Sigma, \varphi_{H'_\varepsilon}(\underline{L}), \underline{L}) \cong H^*(\mathbb{T}^d, R) \cong HF(\Sigma, \varphi_{H'_\varepsilon}(\underline{L}), \underline{L})$$

as vector spaces. Since  $\dim_R HF(\Sigma, \varphi_{H'_\varepsilon}(\underline{L}), \underline{L}) \leq \dim_R \ker d_J \leq \dim_R CF(\Sigma, \varphi_{H'_\varepsilon}(\underline{L}), \underline{L})$ , we must have  $d_J = 0$ .  $\square$

Let  $\mathbf{c} := \mathcal{OC}_{Z_{ref}}(W, \Omega_{\varphi_{H'_\varepsilon}}, \mathcal{L}_{\omega_{\varphi_{H'_\varepsilon}}})_{J(\mathbf{y}_\diamond, A_{\mathbf{y}_\diamond})}$ . By Lemma 5.6,  $\mathbf{c}$  is a cycle. To show that  $\mathbf{c}$  is non-exact, we first need to find the corresponding cycle  $\mathbf{c}' \in \widetilde{PFC}(\Sigma, \varphi_{H_\varepsilon}, \gamma_0)$  because the elements in  $\widetilde{PFC}(\Sigma, \varphi_{H_\varepsilon}, \gamma_0)$  can be figured out easily.

Let  $(X, \Omega_X)$  be the symplectic cobordism in (2.10). We take  $H_+ = H'_\varepsilon$  and  $H_- = H_\varepsilon$ . Note that  $\Omega_X = \omega + dH_\varepsilon \wedge dt + ds \wedge dt$  is  $\mathbb{R}$ -invariant over  $\mathbb{R} \times S^1_t \times (\Sigma - \mathcal{U}^{\delta+\delta_0})$ . This region is called a **product region**. Take  $Z_{ref} = [\mathbb{R} \times S^1 \times \mathbf{x}] \in H_2(X, \gamma_{H'_\varepsilon}^{\mathbf{x}}, \gamma_{H_\varepsilon}^{\mathbf{x}})$  be the reference homology class.

Let  $\mathcal{J}_{comp}(X, \Omega_X)$  be the set of  $\Omega_X$ -compatible almost complex structures such that

1.  $J_X|_{s \geq R_0} \in \mathcal{J}(Y_{\varphi_{H'_\varepsilon}}, \omega_{\varphi_{H'_\varepsilon}})$  and  $J_X|_{s \leq 0} \in \mathcal{J}(Y_{\varphi_{H_\varepsilon}}, \omega_{\varphi_{H_\varepsilon}})$ .
2.  $j \circ \pi_* = \pi_* \circ J_X$ , where  $j$  is the complex structure on  $\mathbb{R}_s \times S^1_t$  that  $j(\partial_s) = \partial_t$ .

**Lemma 5.7.** *Let  $J_X \in \mathcal{J}_{comp}(X, \Omega_X)$  be an almost complex structure such that it is  $\mathbb{R}$ -invariant in the product region  $\mathbb{R} \times S^1_t \times (\Sigma - \mathcal{U}^{\delta+\delta_0})$ . Let  $\overline{\mathcal{M}_0^{J_X}(\alpha_\diamond, \alpha_I)}$  be the moduli space of broken holomorphic curves with  $I = 0$ . Then  $\mathcal{M}_0^{J_X}(\alpha_\diamond, \alpha_I) = \emptyset$  unless  $\alpha_I = \alpha_\diamond$ . In the case that  $\alpha_I = \alpha_\diamond$ , the trivial cylinder is the unique element in  $\overline{\mathcal{M}_0^{J_X}(\alpha_\diamond, \alpha_\diamond)}$ .*

*Proof.* Let  $\mathcal{C} \in \overline{\mathcal{M}_0^{J_X}(\alpha_\diamond, \alpha_I)}$  be a (broken) holomorphic curve. Let  $Z \in H_2(X, \alpha_\diamond, \alpha_I)$  denote the relative homology class of  $\mathcal{C}$ . Then  $Z$  can be written as  $Z_{\alpha_\diamond, \alpha_I} + m(Z)[\Sigma] + [S]$ , where  $[S] \in H_1(S^1, \mathbb{Z}) \otimes H_1(\Sigma, \mathbb{Z})$ . It is easy to show that

$$I(\alpha_\diamond, \alpha_I, Z) = 2m(Z)(k+1) \text{ and } \int_Z \omega_X = m(Z).$$

Then  $m(Z) = 0$  because  $I = 0$ . Also, we have

$$\#(\mathcal{C} \cap \mathbb{R} \times \gamma_{r_0, \theta}) = \#(Z_{\alpha_\diamond, \alpha_I} \cap \mathbb{R} \times \gamma_{r_0, \theta}) = 0.$$

Note that the above intersection numbers are well defined because  $\gamma_{r_0, \theta}$  and  $\alpha_I$  are disjoint. Because  $\mathbb{R} \times \gamma_{r_0, \theta}$  is holomorphic by the choice of  $J_X$ , the above equality implies that  $\mathcal{C}$  doesn't intersect  $\mathbb{R} \times \gamma_{r_0, \theta}$ . In particular,  $\mathcal{C}$  is contained in the product region  $\mathbb{R} \times S^1_t \times (\Sigma - \mathcal{U}^{\delta+\delta_0})$ . Then  $\int_{\mathcal{C}} \omega_X = 0$  implies that  $\mathcal{C}$  is a union of trivial cylinders (Proposition 9.1 of [18]). Thus we must have  $\alpha_I = \alpha_\diamond$ .  $\square$

**Lemma 5.8.** *Let  $J_X$  be a generic almost complex structure in  $\mathcal{J}_{comp}(X, \Omega_X)$  and  $J_X$  is  $\mathbb{R}$ -invariant in the product region  $\mathbb{R} \times S_t^1 \times (\Sigma - \mathcal{U}^{\delta+\delta_0})$ . Then we have*

$$PFC_{Z_{ref}}^{sw}(X, \Omega_X)_{J_X}(\alpha_\diamond, Z_{\alpha_\diamond}) = (\alpha_\diamond, Z_{\alpha_\diamond}) + \sum(\beta', Z'),$$

where  $(\beta', Z')$  satisfies  $\mathbb{A}_{H'_\varepsilon}(\alpha_\diamond, Z_{\alpha_\diamond}) - \mathbb{A}_{H_\varepsilon}(\beta', Z') \geq \frac{1}{4(k+1)}$  and  $\beta' \neq \alpha_I$ .

*Proof.* By the holomorphic axioms (Theorem 1 of [11] and Appendix of [13]) and Lemma 5.7, we know that

$$\langle PFC_{Z_{ref}}^{sw}(X, \Omega_X)_{J_X}(\alpha_\diamond, Z_{\alpha_\diamond}), (\alpha_I, Z_I) \rangle = 0$$

when  $(\alpha_I, Z_I) \neq (\alpha_\diamond, Z_{\alpha_\diamond})$ , and

$$\langle PFC_{Z_{ref}}^{sw}(X, \Omega_X)_{J_X}(\alpha_\diamond, Z_{\alpha_\diamond}), (\alpha_\diamond, Z_{\alpha_\diamond}) \rangle = 1.$$

Assume that  $\langle PFC_{Z_{ref}}^{sw}(X, \Omega_X)_{J_X}(\alpha_\diamond, Z_{\alpha_\diamond}), (\beta', Z') \rangle = 1$  for some  $(\beta', Z')$  and  $\beta' \neq \alpha_I$ . Again by the holomorphic axioms, we have a holomorphic curve  $\mathcal{C} \in \overline{\mathcal{M}}_0^J(\alpha_\diamond, \beta')$ .

It is easy to check that

$$\begin{aligned} I(\mathcal{C}) &= -h(\beta') - 2e_+(\beta') + 2m(k+1) = 0 \\ \mathbb{A}_{H'_\varepsilon}(\alpha_\diamond, Z_{\alpha_\diamond}) - \mathbb{A}_{H_\varepsilon}(\beta', Z) &= -H_\varepsilon(\beta') + m, \end{aligned} \tag{5.32}$$

where  $h(\beta')$  is the total multiplicities of the simple hyperbolic orbits in  $\beta'$  and  $e_+(\beta')$  is the total multiplicities of the simple elliptic orbits at local maximum of  $H_\varepsilon$ .

Because  $\beta' \neq \alpha_I$ , we have  $h(\beta') + 2e_+(\beta') \geq 1$ . Therefore, we have

$$\mathbb{A}_{H_\varepsilon}(\alpha_\diamond, Z_{\alpha_\diamond}) - \mathbb{A}_{H_\varepsilon}(\beta', Z') = -H_\varepsilon(\beta') + \frac{h(\beta') + 2e_+(\beta')}{2(k+1)} \geq \frac{1}{2(k+1)} + O(d\varepsilon_0) \geq \frac{1}{4(k+1)}.$$

□

**Lemma 5.9.** *Let  $J_X$  be the almost complex structure in Lemma 5.8. Then we have*

$$PFC_{Z_{ref}}^{sw}(X, \Omega_X)_{J_X}(\beta, Z) = \sum(\beta', Z'),$$

where  $(\beta', Z')$  satisfies  $\mathbb{A}_{H_\varepsilon}(\alpha_\diamond, Z_{\alpha_\diamond}) - \mathbb{A}_{H_\varepsilon}(\beta', Z') \geq \frac{1}{4(k+1)}$  and  $\beta' \neq \alpha_I$ .

*Proof.* First, we show that  $\beta'$  cannot be  $\alpha_I$ . Assume that

$$\langle PFC_{Z_{ref}}^{sw}(X, \Omega_X)_{J_X}(\beta, Z), (\alpha_I, Z') \rangle = 1.$$

Then we have a broken holomorphic curve  $\mathcal{C} = (C, \mathcal{C}_0)$ , where  $C \in \overline{\mathcal{M}}_0^J(\mathbf{y}_\diamond, \beta)$  is an HF-PFH curve and  $\mathcal{C}_0 \in \overline{\mathcal{M}}_0^{J_X}(\beta, \alpha_I)$ . The holomorphic curve gives us a relative homology class  $\mathcal{Z} \in H_2(W, \mathbf{y}_\diamond, \alpha_I)$ .



Reintroduce the periodic orbits  $\gamma_{r_0, \theta_0}^i$  near the local maximums of  $H_\varepsilon$ . The superscript “ $i$ ” indicates that the local maximum lies in the domain  $\mathring{B}_i$ , where  $1 \leq i \leq k+1$ . In particular,  $\gamma_{r_0, \theta_0}^i$  lies in  $S^1 \times \mathring{B}_i$ . Define a curve  $v_i := (\mathbb{R} \times \gamma_{r_0, \theta_0}^i) \cap W$ . Note that it is  $J$ -holomorphic and  $\partial v_i$  is disjoint from the Lagrangian  $\mathcal{L}_{H_\varepsilon}$ . Then for any relative homology class  $\mathcal{Z}' \in H_2(W, \mathbf{y}_\diamond, \alpha_I)$ , we have a well-defined intersection number

$$n_i(\mathcal{Z}') := \#(\mathcal{Z}' \cap v_i).$$

The relative homology class  $\mathcal{Z} \in H_2(W, \mathbf{y}_\diamond, \alpha_I)$  can be written as  $\mathcal{Z} = \mathcal{Z}_{hor} \# Z_{\alpha_\diamond, \alpha_I} + \sum_{i=1}^k c_i [B_i] + m[\Sigma] + [S]$ , where  $[S] \in H_1(S^1, \mathbb{Z}) \otimes H_1(\Sigma, \mathbb{Z})$  and  $\mathcal{Z}_{hor}$  is the class represented by the union of horizontal sections. By (5.30), the ECH index of  $\mathcal{Z}$  is

$$I(\mathcal{Z}) = \sum_i 2c_i + 2m(k+1) = I(C) + I(\mathcal{C}_0) = 0. \quad (5.33)$$

Let  $q_i$  denote the period of  $\gamma_{r_0, \theta_0}^i$ . From the construction in [13], the period of  $\gamma_{r_0, \theta_0}^i$  is determined by the function  $\varepsilon$ . For a suitable choice of  $\varepsilon$ , we can choose  $q_i = q$  for  $1 \leq i \leq k+1$ . By definition, we have

$$n_i(\mathcal{Z}_{hor} \# Z_{\alpha_\diamond, \alpha_I}) = 0, \quad n_i([B_j]) = \delta_{ij} q \quad n_i([S]) = 0 \quad \text{and} \quad n_i([\Sigma]) = q. \quad (5.34)$$

for  $1 \leq i, j \leq k+1$ . From (5.33) and (5.34), we know that

$$\#(\mathcal{C} \cap (\sqcup_{i=1}^{k+1} v_i)) = \sum_{i=1}^{k+1} n_i(\mathcal{Z}) = \sum_{i=1}^k c_i q + (k+1)m q = 0.$$

By the intersection positivity of holomorphic curves,  $\mathcal{C}$  doesn't intersect  $\mathbb{R} \times \gamma_{r_0, \theta_0}^i$ . In particular,  $\mathcal{C}_0$  lies inside the product region of  $X$  and  $E_{\omega_{\varphi_{H'_\varepsilon}}}(\mathcal{C}_0) \geq 0$ . By Lemma 5.4, we have

$$\int_{\mathcal{Z}} \omega_{\varphi_{H'_\varepsilon}} = \int_{\mathcal{Z}_{hor} \# Z_{\alpha_\diamond, \alpha_I}} \omega_{\varphi_{H'_\varepsilon}} = 0 = E_{\omega_{\varphi_{H'_\varepsilon}}}(C) + E_{\omega_{\varphi_{H'_\varepsilon}}}(\mathcal{C}_0) > 0.$$

We obtain a contradiction.

Now we consider the case that  $\langle PFC_{Z_{ref}}^{sw}(X, \Omega_X)_{J_X}(\beta, Z), (\beta', Z') \rangle = 1$  and  $\beta' \neq \alpha_I$ . As before, we have a broken holomorphic curve  $\mathcal{C} = (C, \mathcal{C}_0)$ , where  $C \in \mathcal{M}_0^J(\mathbf{y}_\diamond, \beta)$  is an HF-PFH curve and  $\mathcal{C}_0 \in \overline{\mathcal{M}^{J_X}(\beta, \beta')}$ . Similar as (5.32), we have

$$0 = I(\mathcal{C}) = I(C) + I(\mathcal{C}_0) = -h(\beta') - 2e_+(\beta') + \sum_{i=1}^k 2c_i + 2m(k+1)$$

$$\mathcal{A}_{H'_\varepsilon}(\mathbf{y}_\diamond, A_{\mathbf{y}_\diamond}) - \mathbb{A}_{H_\varepsilon}(\beta', Z') = \int_C \omega_{\varphi_{H'_\varepsilon}} + \int_{\mathcal{C}_0} \omega_X = -H_\varepsilon(\beta') + m + \sum_{i=1}^k \frac{c_i}{k+1}.$$

Note that  $\mathcal{A}_{H'_\varepsilon}(\mathbf{y}_\diamond, [A_{\mathbf{y}_\diamond}]) = \mathbb{A}_{H'_\varepsilon}(\alpha_\diamond, Z_{\alpha_\diamond})$ . Therefore, we have

$$\mathbb{A}_{H'_\varepsilon}(\alpha_\diamond, Z_{\alpha_\diamond}) - \mathbb{A}_{H_\varepsilon}(\beta', Z') \geq \frac{1}{4(k+1)}.$$

□

**Lemma 5.10.** *Let  $\mathfrak{c}' = PFC_{Z_{ref}}^{sw}(X, \Omega_X)(\mathfrak{c})$ . Then the cycle  $\mathfrak{c}'$  is non-exact, i.e., it represents a non-zero class in  $\widetilde{PFH}(\Sigma, \varphi_{H_\varepsilon}, \gamma_{H_\varepsilon}^x)$ .*

*Proof.* Let  $(X_-, \Omega_{X_-})$  be the symplectic cobordism from  $(Y_{\varphi_{H_\varepsilon}}, \omega_{\varphi_{H_\varepsilon}})$  to  $\emptyset$  in (2.11). Fix  $J_{X_-} \in \mathcal{J}_{comp}(X_-, \Omega_{X_-})$ . Using the same argument as in [12], we define a homomorphism

$$PFC_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}} : \widetilde{PFC}(\Sigma, \omega_{\varphi_H}, \gamma_{H_\varepsilon}^x) \rightarrow \Lambda,$$

by counting  $I = 0$  (unbroken) holomorphic curves in  $(X_-, \Omega_{X_-})$ . Moreover, this is a chain map. Therefore,  $PFC_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}}$  induces a homomorphism in homology level:

$$PFH_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}} : \widetilde{PFH}(\Sigma, \omega_{\varphi_{H_\varepsilon}}, \gamma_{H_\varepsilon}^x) \rightarrow \Lambda,$$

Using Taubes's techniques [32, 33] and C. Gerig's generalization [10],  $PFH_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}}$  should agree with the PFH cobordism map  $PFH_{Z_{ref}}^{sw}(X_-, \Omega_{X_-})_{J_{X_-}}$  (see Remark 1.3 of [12]). But we don't need this to prove the lemma.

To show that  $\mathfrak{c}'$  is non-exact, it suffices to prove  $PFC_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}}(\mathfrak{c}') \neq 0$ . In [12], the author computes the map  $PFC_{Z_{ref}}^{hol}(X, \Omega_X)_{J_X}$  for the elementary Lefschetz fibration (a symplectic fibration over a disk with a single singularity). The current situation is an easier version of [12]. By the argument in [12], we have

$$\begin{aligned} PFC_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}}(\alpha_I, Z_I) &= 1, \\ PFC_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}}(\beta', Z') &= 0 \quad \text{for } (\beta', Z') \neq (\alpha_I, Z_I). \end{aligned} \tag{5.35}$$

Therefore,  $PFC_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}}(\mathfrak{c}') = 1$ .

Here let us explain a little more about how to get (5.35). Basically, the idea is the same as Lemma 3.7. From the computation of the ECH index, we know that  $I = 0$  implies that the holomorphic curves must be asymptotic to  $\alpha_I$ . Also, the energy is zero. Therefore, the unbranched covers of the horizontal sections are the only curves that contribute to  $PFC_{Z_{ref}}^{hol}(X_-, \Omega_{X_-})_{J_{X_-}}$ , and this leads to (5.35). Even these holomorphic curves may not be simple, they are still regular (see [9]). □

## 6 Spectral invariants

### 6.1 Comparing the PFH and HF spectral invariants

Let  $H_\varepsilon$  be the function satisfying **F.1**, **F.2**, **F.3**, **F.4**, **F.5**. Let  $H'_\varepsilon$  be the perturbation. Let  $\mathbf{y}_\diamond = [0, 1] \times (y_-^1, \dots, y_-^d)$  be the Reeb chord of  $\varphi_{H'_\varepsilon}$ , where  $\{y_-^i\}_{i=1}^d$  are minimums

of  $H_\epsilon$ . By Lemma 5.6,  $[(\mathbf{y}_\diamond, [A_{\mathbf{y}_\diamond}])]$  represents a class  $e_\diamond \in HF(\Sigma, \underline{L})$ , i.e.,  $e_\diamond = j_{H'_\epsilon}^{\mathbf{x}}([(y_\diamond, [A_{y_\diamond}])])$ . Let  $e$  be the unit of  $HF(\Sigma, \underline{L})$ . Reintroduce

$$c_{\underline{L}}^+(H) := c_{\underline{L}}(H, e) \text{ and } c_{\underline{L}}^-(H) := c_{\underline{L}}(H, e_\diamond).$$

Fix a base point  $\mathbf{x} = (x_1, \dots, x_d)$ . Define a reference 1-cycle  $\gamma_H^{\mathbf{x}} := \Psi_H(S^1 \times \mathbf{x})$ . Let  $(X, \Omega_X)$  symplectic cobordism (2.9) from  $(Y_{\varphi_H}, \omega_{\varphi_H})$  to  $(Y_{\varphi_G}, \omega_{\varphi_G})$ . Let  $Z_{\mathbf{x}}^{H,G} = [\mathbb{R} \times S^1 \times \mathbf{x}] \in H_2(X, \gamma_H^{\mathbf{x}}, \gamma_G^{\mathbf{x}})$  be a reference class. Given another base point  $\mathbf{x}'$ , let  $\eta$  be  $d$  union of paths starting from  $\mathbf{x}$  and ending at  $\mathbf{x}'$ . Let  $Z_{\mathbf{x}', \mathbf{x}} = [S^1 \times \eta] \in H_2(X, \gamma_H^{\mathbf{x}'}, \gamma_H^{\mathbf{x}})$ . The cobordism map  $PfH_{Z_{\mathbf{x}}^{H,G}}^{sw}(X, \Omega_X)(\sigma_\diamond)$  only depends on  $\mathbf{x}, H, G$ . Thus we denote it by  $\mathfrak{J}_{H,G}^{\mathbf{x}}$ .

The cycle  $\mathbf{c} = (\alpha_\diamond, Z_{\alpha_\diamond}) + \sum(\beta, Z)$  represents a class  $\sigma_\diamond \neq 0 \in PfH(\Sigma, H'_\epsilon, \gamma_{H'_\epsilon}^{\mathbf{y}_\diamond})$ . Define

$$\sigma_{\diamond H}^{\mathbf{x}} := \Psi_{Z_{\mathbf{y}_\diamond, \mathbf{x}}} \circ \mathfrak{J}_{H'_\epsilon, H}^{\mathbf{y}_\diamond}(\sigma_\diamond).$$

Let  $H_-$  and  $H_+$  be Hamiltonian functions satisfying conditions  $\spadesuit.1$  and  $\spadesuit.2$  respectively. Let  $W_+$  be the open-closed cobordism and  $W_-$  the closed-open cobordism. Then by [13] and Theorem 2, we have

$$\begin{aligned} \mathcal{CO}_{Z_{ref}^-}(W_-, \varphi_{H_-}, \mathcal{L}_{H_-}) * (\sigma_{\diamond H_-}^{\mathbf{x}})^{-1}(e) \\ \mathcal{OC}_{Z_{ref}^+}(W_+, \Omega_{H_+}, \mathcal{L}_{H_+}) * ((j_{H_+}^{\mathbf{x}})^{-1}(e_\diamond)) = \sigma_{\diamond H_+}^{\mathbf{x}}. \end{aligned}$$

*Proof of Corollary 1.2.* The inequality  $c_{\underline{L}}^+(H) \leq c_d^{pjh}(H, \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x}) dt$  is the Corollary 1.1 of [13].

To prove  $c_d^{pjh,-}(H, \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x}) dt \leq c_{\underline{L}}^-(H)$ , it suffices the inequality for a Hamiltonian function  $H$  satisfying  $\spadesuit.2$  because of the Hofer-Lipschitz property and Remark 1.3. By using Theorem 2 and the argument in Section 7 of [13], we obtain the inequality.

By definition,  $(\mathbf{y}_\diamond, A_{\mathbf{y}_\diamond})$  represents  $(j_{H'_\epsilon}^{\mathbf{x}})^{-1}(e_\diamond) = (j_{-\epsilon f}^{\mathbf{x}})^{-1}(e_\diamond)$ . Therefore,

$$c_{\underline{L}}(-\epsilon f, e_\diamond) \leq \mathcal{A}_{-\epsilon f}(\mathbf{y}_\diamond, A_{\mathbf{y}_\diamond}) = O(\epsilon).$$

Let  $\epsilon \rightarrow 0$ . We have  $c_{\underline{L}}(0, e_\diamond) \leq 0$ . By the triangle inequality, we have

$$c_{\underline{L}}^-(H) \leq c_{\underline{L}}^+(H) + c_{\underline{L}}(0, e_\diamond) \leq c_{\underline{L}}^+(H).$$

□

## 6.2 Homogenized spectral invariants

Let  $\widetilde{Ham}(\Sigma, \omega)$  be the universal cover of  $Ham(\Sigma, \omega)$ . A element in  $\widetilde{Ham}(\Sigma, \omega)$  is a homotopy class of paths  $\{\varphi_t\}_{t \in [0,1]} \subset Ham(\Sigma, \omega)$  with fixed endpoints  $\varphi_0 = id$  and  $\varphi_1 = \varphi$ . Let  $\tilde{\varphi} \in \widetilde{Ham}(\Sigma, \omega)$  be a class represented by a path generated by a mean-normalized Hamiltonian  $H$ . Define

$$c_{\underline{L}}(\tilde{\varphi}, a) := c_{\underline{L}}(H, a).$$

By Theorem 1, this is well defined. Thus, the HF spectral invariants descend to invariants on  $\tilde{\varphi} \in \widetilde{Ham}(\Sigma, \omega)$ . But in general, the spectral invariants cannot descend to  $Ham(\Sigma, \omega)$ . This is also true for PFH spectral invariants. To obtain numerical invariants for the elements in  $Ham(\Sigma, \omega)$  rather than its universal cover, we need the homogenized spectral invariants. It is well known that  $\widetilde{Ham}(\Sigma, \omega) = Ham(\Sigma, \omega)$  when  $g(\Sigma) \geq 1$ . Therefore, we only consider the case that  $\Sigma = \mathbb{S}^2$ . Fix  $\varphi \in Ham(\Sigma, \omega)$ . We define the **homogenized HF spectral invariant** by

$$\mu_{\underline{L}}(\varphi, a) := \limsup_{n \rightarrow \infty} \frac{c_{\underline{L}}(\tilde{\varphi}^n, a)}{n},$$

where  $\tilde{\varphi}$  is a lift of  $\varphi$ . One can define the PFH spectral invariant for  $\tilde{\varphi}$  in the same manner, denoted by  $c_d^{pfh}(\tilde{\varphi}, \sigma, \gamma_0)$ . Similarly, the **homogenized PFH-spectral invariant** is

$$\mu_d^{pfh}(\varphi) = \limsup_{n \rightarrow \infty} \frac{1}{n} \left( c_d^{pfh}(\#^n H, \sigma_{\heartsuit \#^n H}^{\mathbf{x}}, \gamma_{\#^n H}^{\mathbf{x}}) + \int_0^1 \#^n H_t(\mathbf{x}) dt \right),$$

where  $H$  is a mean-normalized Hamiltonian function generated  $\tilde{\varphi}$ . This is well defined by Proposition 3.6 of [5].

**Remark 6.1.** Let  $\mathbf{x} = (x_1, \dots, x_d)$ . Suppose that each  $x_i$  is the south pole of the sphere and  $H_t(x_i) = 0$ . Then  $c_d^{pfh}(H, \sigma_{\heartsuit H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}})$  agrees with spectral invariant  $c_{d,k}(H)$  in [5], where  $k$  is the grading of  $\sigma_{\heartsuit H}^{\mathbf{x}}$ .

*Proof of Corollary 1.3.* There is a natural trivialization  $\tau_H$  of  $\xi|_{\gamma_H^{\mathbf{x}}}$  defined by pushing forward the  $S^1$ -invariant trivialization over  $S^1 \times \{\mathbf{x}\}$ . Then we have a well-defined grading  $gr(\alpha, [Z])$  for each anchored orbit set (see (11) of [5]). We claim that

$$gr(\sigma_{\heartsuit H}^{\mathbf{x}}) - gr(\sigma_{\diamond H}^{\mathbf{x}}) = 2d.$$

Because the cobordism maps  $\mathfrak{J}_{H,G}^{\mathbf{x}}$  preserve the grading, one only needs to check this for a special case that  $H$  is a small Morse function. Take  $H = H_\varepsilon$ . Then  $\sigma_{\diamond H}^{\mathbf{x}}$  is represented by  $(\alpha_\diamond, Z_{\alpha_\diamond})$ . The class  $\sigma_{\heartsuit H}^{\mathbf{x}} = PFH_{Z_{ref}}^{sw}(X_+, \Omega_{X_+})(1)$  (see Remark 6.1 of [13]), where  $X_+ = B_+ \times \Sigma$  and  $B_+$  is a punctured sphere with a negative end. The construction of  $(X_+, \Omega_{X_+})$  is similar to (2.11). By index reason, the class  $\sigma_{\heartsuit H}^{\mathbf{x}}$  can be

represented by a cycle that is a certain combination of constant orbits at the maximum points. It is not difficult to show that the claim is true.

According to Example 2.16 of [16], we know that

$$U^d \sigma_{\diamond H}^{\mathbf{x}} = \sigma_{\diamond H}^{\mathbf{x}} \text{ and } U^{d+1} \sigma_{\diamond H}^{\mathbf{x}} = q \sigma_{\diamond H}^{\mathbf{x}}.$$

According to Proposition 4.2 of [16], we have

$$c_d^{pjh}(H, \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) - 1 + \int_0^1 H_t(\mathbf{x}) dt = c_d^{pjh}(H, q \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x}) dt \leq c_{\underline{L}}^-(H).$$

Here the second inequality follows from that the U-map decreases the spectral numbers. Therefore, we have

$$c_d^{pjh}(H, \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x}) dt - 1 \leq c_{\underline{L}}^-(H) \leq c_{\underline{L}}^+(H) \leq c_d^{pjh}(H, \sigma_{\diamond H}^{\mathbf{x}}, \gamma_H^{\mathbf{x}}) + \int_0^1 H_t(\mathbf{x}) dt.$$

This implies that (1.3).  $\square$

### 6.3 Quasimorphisms

In this section, we show that  $\mu_d^{pjh}$  is a quasimorphism on  $Ham(\mathbb{S}^2, \omega)$ . The argument is similar to M. Entov and L. Polterovich [17]. Before we prove the result, let us recall some facts about the duality in Floer homology.

Let  $\mathfrak{c}$  be a **graded filtered Floer-Novikov complex** over a field  $\mathbb{F}$  in the sense of [34]. We can associate  $\mathfrak{c}$  with a graded chain complex  $(C_*(\mathfrak{c}), \partial)$ . One can define the homology and spectral numbers for  $(C_*(\mathfrak{c}), \partial)$ . It is not hard to see that the PFH chain complex is an example of graded filtered Floer-Novikov complexes. Roughly speaking,  $\mathfrak{c}$  is an abstract complex that is characterized by the common properties of Floer homology.

For  $\mathfrak{c}$ , M. Usher defines another graded filtered Floer-Novikov complex  $\mathfrak{c}^{op}$  called the opposite complex. Roughly speaking, the homology of  $(C_*(\mathfrak{c}^{op}), \delta)$  is the Poincare duality of  $H_*(C_*(\mathfrak{c}))$  in the following sense: There is a non-degenerate pairing  $\Delta : H_{-k}(C_*(\mathfrak{c}^{op})) \times H_k(C_*(\mathfrak{c})) \rightarrow \mathbb{F}$ . We refer the readers to [34] for the details of the graded filtered Floer-Novikov complex and opposite complex.

Let  $\mathfrak{c}_1, \mathfrak{c}_2$  be graded filtered Floer-Novikov complexes. Let  $I : C_*(\mathfrak{c}_1) \rightarrow C_*(\mathfrak{c}_2)$  be a 0-degree chain map given by

$$Ip_1 = \sum_{p_2} n(p_1, p_2)p_2,$$

where  $p_i$  are generators of  $C_*(\mathfrak{c}_i)$  and  $n(p_1, p_2) \in \mathbb{F}$ . Define  $I^{op} : C_*(\mathfrak{c}_2^{op}) \rightarrow C_*(\mathfrak{c}_1^{op})$  by

$$I^{op}p_2 = \sum_{p_1} n(p_1, p_2)p_1.$$

**Lemma 6.1.** *The map  $I^{op} : C_*(\mathfrak{c}_2^{op}) \rightarrow C_*(\mathfrak{c}_1^{op})$  satisfies the following properties:*

- *$I^{op}$  is a chain map. It descends to a map  $I_*^{op} : H_*(C_*(\mathfrak{c}_2^{op})) \rightarrow H_*(C_*(\mathfrak{c}_1^{op}))$ .*
- *Let  $I_1 : C_*(\mathfrak{c}_1) \rightarrow C_*(\mathfrak{c}_2)$  and  $I_2 : C_*(\mathfrak{c}_2) \rightarrow C_*(\mathfrak{c}_3)$  be two 0-degree chain maps. Then  $(I_1 \circ I_2)^{op} = I_2^{op} \circ I_1^{op}$ . In particular, if  $I_*$  is an isomorphism, so is  $I_*^{op}$ .*
- *Let  $a \in H_{-k}(C_*(\mathfrak{c}_2^{op}))$  and  $b \in H_k(C_*(\mathfrak{c}_1))$ . Then we have*

$$\Delta(a, I_*(b)) = \Delta(I_*^{op}(a), b).$$

The proof of this lemma is straightforward (see Proposition 2.4 in [34] for the case  $\mathfrak{c}_1 = \mathfrak{c}_2$ ), we left the details to the readers.

Now we construct the opposite complex of  $(PFC_*(\mathbb{S}^2, \varphi_H, \gamma_H^x), \partial_J)$ . Let  $\bar{H}_t = -H_{1-t}$ . This is another function generated  $\varphi_H^{-1}$ . Define a diffeomorphism

$$\begin{aligned} \iota : S_t^1 \times \Sigma &\rightarrow S_\tau^1 \times \Sigma \\ (t, x) &\rightarrow (1-t, x). \end{aligned}$$

Note that  $(\iota^{-1})^*(\omega + dH_t \wedge dt) = \omega + d\bar{H}_\tau \wedge d\tau$ . If  $\gamma$  is a  $\varphi_H$  periodic orbit, then  $\bar{\gamma} := \iota \circ \gamma$  is a  $\varphi_H^{-1}$  periodic orbit. Here we orient  $\bar{\gamma}$  such that it transverse  $\Sigma$  positively.

Recall that the symplectic cobordism  $(X = \mathbb{R} \times S^1 \times \Sigma, \Omega_X = \omega + d(H_t^s dt) + ds \wedge dt)$ . We extend the map  $\iota$  to be

$$\begin{aligned} \iota : \mathbb{R}_s \times S_t^1 \times \Sigma &\rightarrow \mathbb{R}_r \times S_\tau^1 \times \Sigma \\ (s, t, x) &\rightarrow (-s, 1-t, x). \end{aligned}$$

Note that  $(\iota^{-1})^*\Omega_X = \omega - d(H_{1-\tau}^{-r} \wedge d\tau) + dr \wedge d\tau$ . Therefore,  $(X, (\iota^{-1})^*\Omega_X)$  is a symplectic cobordism from  $(Y_{\varphi_{\bar{H}^-}}, \omega_{\varphi_{\bar{H}^-}})$  to  $(Y_{\varphi_{\bar{H}^+}}, \omega_{\varphi_{\bar{H}^+}})$

Consider the case that  $H_t^s = H_t$ . Let  $PFC_*(\mathbb{S}^2, \varphi_H^{-1}, \gamma_H^x)$  be the complex generated by  $(\bar{\gamma}, -\iota_*Z)$ . Note that  $\iota_*Z \in H_2(Y_{\varphi_H^{-1}}, \gamma_H^x, \bar{\gamma})$ . Here  $-\iota_*Z$  denote the unique class in  $H_2(Y_{\varphi_H^{-1}}, \bar{\gamma}, \gamma_H^x)$  such that  $(-\iota_*Z) \# \iota_*Z = [\mathbb{R} \times \bar{\gamma}]$ . Note that we have

**O.1**  $\mathcal{A}_{\bar{H}}(\bar{\gamma}, -\iota_*Z) = -\mathcal{A}(\gamma, [Z]).$

**O.2**  $gr(\bar{\gamma}, -\iota_*Z) = -gr(\gamma, Z).$

**O.3** Let  $u \in \mathcal{M}^J(\gamma_+, \gamma_-, Z)$  be a holomorphic curve in  $(X, \Omega_X)$ . Then  $\bar{u} = \iota \circ u \in \mathcal{M}^{\bar{J}}(\bar{\gamma}_-, \bar{\gamma}_+, \iota_*Z)$  be a holomorphic curve in  $(X, (\iota^{-1})^*\Omega_X)$ , where  $\bar{J} = \iota_* \circ J \circ \iota_*^{-1}$ . This establishes a 1-1 correspondence between  $\mathcal{M}^J(\gamma_+, \gamma_-, Z)$  and  $\mathcal{M}^{\bar{J}}(\bar{\gamma}_-, \bar{\gamma}_+, \iota_*Z)$ .

These three points implies that  $PFC_*(\mathbb{S}^2, \varphi_H^{-1}, \gamma_H^{\mathbf{x}})$  is the opposite complex of  $PFC_*(\mathbb{S}^2, \varphi_H^{-1}, \gamma_{\bar{H}}^{\mathbf{x}})$ .

The pairing  $\Delta : PFC_{-k}(\mathbb{S}^2, \varphi_H^{-1}, \gamma_{\bar{H}}^{\mathbf{x}}) \otimes PFC_k(\mathbb{S}^2, \varphi_H, \gamma_H^{\mathbf{x}}) \rightarrow \mathbb{Z}/2\mathbb{Z}$  is defined by

$$\Delta \left( \sum a_{(\bar{\gamma}, -\iota_*[Z])}(\bar{\gamma}, -\iota_*[Z]), \sum b_{(\gamma, [Z])}(\gamma, [Z]) \right) = \sum_{(\gamma, [Z])} a_{(\bar{\gamma}, -\iota_*[Z])} b_{(\gamma, [Z])}.$$

This pairing descends to the homologies. By Usher's result [34], we have

$$c_d(H, \sigma_{\heartsuit H}^{\mathbf{x}}) = -\inf\{c_d(\bar{H}, \sigma) \mid \sigma \in PFH_{-k}(\mathbb{S}^2, \varphi_H^{-1}, \gamma_H^{\mathbf{x}}), \Delta(\sigma, \sigma_{\heartsuit H}^{\mathbf{x}}) \neq 0\}.$$

**Lemma 6.2.** *For any Hamiltonian function  $H$ , we have*

$$\begin{aligned} c_d^{pfh}(H, \sigma_{\heartsuit H}^{\mathbf{x}}) + c_d^{pfh}(\bar{H}, \sigma_{\heartsuit \bar{H}}^{\mathbf{x}}) &\leq 1 \\ c_L^+(H) + c_L^+(\bar{H}) &\leq 1. \end{aligned}$$

*Proof.* Let  $g : \mathbb{S}^2 \rightarrow \mathbb{R}$  be a Morse function with two critical points  $x_+, x_-$ , where  $x_+$  is the maximum point and  $x_-$  is the minimum point. Let  $\bar{G}_\epsilon := \epsilon g$ . Take  $\mathbf{x} = (x_-, \dots, x_-)$  be the base point. Then

$$gr((\gamma_{x_+}^{d_+} \gamma_{x_-}^{d_-}, Z_{\gamma_{x_+}^{d_+} \gamma_{x_-}^{d_-}} + m[\mathbb{S}^2])) = 2d_+ + 2m(d+1) - d,$$

where  $d_\pm \geq 0$  such that  $d_+ + d_- = d$ . The grading formula implies that  $\partial = 0$ . Also, we have  $\sigma_{\heartsuit \bar{G}_\epsilon}^{\mathbf{x}} = (\gamma_{x_+}^d, dZ_{\gamma_{x_+}^d})$  and  $\sigma_{\heartsuit \bar{G}_\epsilon}^{\mathbf{x}} = (\gamma_{x_-}^d, dZ_{\gamma_{x_-}^d})$ . Then for any  $H$ ,  $\sigma_{\heartsuit \bar{H}}^{\mathbf{x}} = \mathfrak{J}_{\bar{G}_\epsilon, \bar{H}}^{\mathbf{x}}((\gamma_{x_+}^d, dZ_{\gamma_{x_+}^d}))$ .

By the observation **O.3**, we have  $(\mathfrak{J}_{\bar{G}_\epsilon, \bar{H}}^{\mathbf{x}})^{op} = \mathfrak{J}_{H, G_\epsilon}^{\mathbf{x}}$ . Therefore,

$$\Delta(\sigma, \sigma_{\heartsuit \bar{H}}^{\mathbf{x}}) = \Delta(\sigma, \mathfrak{J}_{\bar{G}_\epsilon, \bar{H}}^{\mathbf{x}}((\gamma_{x_+}^d, dZ_{\gamma_{x_+}^d}))) = \Delta(\mathfrak{J}_{H, G_\epsilon}^{\mathbf{x}}(\sigma), (\gamma_{x_+}^d, dZ_{\gamma_{x_+}^d})).$$

Note that  $(\bar{\gamma}_{x_+}^d, -\iota_* Z_{\gamma_{x_+}^d})$  is the only class with  $gr = -d$ . Hence,  $\Delta(\sigma, \sigma_{\heartsuit \bar{H}}^{\mathbf{x}}) \neq 0$  if and only if  $\mathfrak{J}_{H, G_\epsilon}^{\mathbf{x}}(\sigma) = (\bar{\gamma}_{x_+}^d, -\iota_* Z_{\gamma_{x_+}^d})$ . Therefore,  $\sigma = \sigma_{\heartsuit H}^{\mathbf{x}}$ . We have

$$-c_d^{pfh}(\bar{H}, \sigma_{\heartsuit \bar{H}}^{\mathbf{x}}) = c_d^{pfh}(H, \sigma_{\heartsuit H}^{\mathbf{x}}) \geq c_d^{pfh}(H, \sigma_{\heartsuit H}^{\mathbf{x}}) - 1.$$

By Corollary 1.3, we get the second inequality for  $c_L^+$ .  $\square$

*Proof of Theorem 3.* By Theorem 1, we have

$$\begin{aligned} &c_L^+(H) + c_L^+(K) \\ &= c_L^+(H) + c_L^+(\bar{H} \diamond H \diamond K) \\ &\leq c_L^+(H) + c_L^+(\bar{H}) + c_L^+(H \diamond K) \leq c_L^+(H \diamond K) + 1. \end{aligned}$$

These two inequalities implies that  $\mu_L = \mu_d^{pfh}$  is a quasimorphism with defect 1.  $\square$

E-mail address: 103200077@seu.edu.cn

## References

- [1] P. Albers, A Lagrangian Piunikhin-Salamon-Schwarz morphism and two comparison homomorphisms in Floer homology, *Int. Math. Res. Not. IMRN* 2008, no. 4, 2008.
- [2] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki, and E. Zehnder, *Compactness results in symplectic field theory*, *Geom. Topol.* **7** (2003), 799–888.
- [3] D. Cristofaro-Gardiner, M. Hutchings, and V. Ramos, *The asymptotics of ECH capacities*, *Invent. Math.* **199**(2015), 187–214.
- [4] D. Cristofaro-Gardiner, V. Humilière, and S. Seyfaddini, *Proof of the simplicity conjecture*. *arXiv:2001.01792*, 2020.
- [5] D. Cristofaro-Gardiner, V. Humilière, and S. Seyfaddini *PFH spectral invariants on the two-sphere and the large scale geometry of Hofer’s metric*. *arXiv:2102.04404v1*, 2021.
- [6] D. Cristofaro-Gardiner, R. Prasad, B. Zhang *The smooth closing lemma for area-preserving surface diffeomorphisms*. *arXiv:2110.02925*, 2021.
- [7] D. Cristofaro-Gardiner, V. Humilière, C. Mak, S. Seyfaddini, I. Smith *Quantitative Heegaard Floer cohomology and the Calabi invariant*. *arXiv:2105.11026*, 2022.
- [8] D. Cristofaro-Gardiner, V. Humilière, C. Mak, S. Seyfaddini, I. Smith *Sub-leading asymptotics of link spectral invariants and homeomorphism groups of surfaces*. *arXiv:2206.10749*
- [9] C. Gerig, *Taming the pseudoholomorphic beasts in  $\mathbb{R} \times S^1 \times S^2$* , *Geom. Topol.* **24**(2020) 1791–1839.
- [10] C. Gerig, *Seiberg–Witten and Gromov invariants for self-dual harmonic 2-forms*, *arXiv:1809.03405*, (2018).
- [11] G. Chen, *On cobordism maps on periodic Floer homology*, *Algebr. Geom. Topol.* **21** (1) 1–103, 2021.
- [12] G. Chen, *Cobordism maps on periodic Floer homology induced by elementary Lefschetz fibrations*. *Topology Appl.* **302** (2021), Paper No. 107818, 23 pp.
- [13] G. Chen, *Closed-open morphisms on periodic Floer homology*, *arXiv:2111.11891*, 2021.



- [14] V. Colin, K. Honda, and Y. Tian, *Applications of higher-dimensional Heegaard Floer homology to contact topology*. *arXiv:2006.05701*, 2020.
- [15] V. Colin, P. Ghiggini, and K. Honda, *The equivalence of Heegaard Floer homology and embedded contact homology via open book decompositions I*, *arXiv:1208.1074*, 2012.
- [16] O. Edtmair, M. Hutchings, *PFH spectral invariants and  $C^\infty$ -closing lemmas*. *arXiv:2110.02463*, 2021
- [17] M. Entov and L. Polterovich. *Calabi quasimorphism and quantum homology*. *Int. Math. Res. Not.* 2003, no. 30, 1635–1676. MR1979584.
- [18] M. Hutchings, *An index inequality for embedded pseudoholomorphic curves in symplectizations*, *J. Eur. Math. Soc.* **4** (2002) 313–361.
- [19] M. Hutchings, *The embedded contact homology index revisited*, *New perspectives and challenges in symplectic field theory*, 263–297, *CRM Proc. Lecture Notes* 49, Amer. Math. Soc., 2009.
- [20] M. Hutchings, M. Sullivan, *The periodic Floer homology of Dehn twist*, *Algebr. Geom. Topol.* **5** (2005), pp. 301 – 354. issn: 1472 – 2747.
- [21] M. Hutchings, *Lecture notes on embedded contact homology*, *Contact and Symplectic Topology*, *Bolyai Society Mathematical Studies*, vol. 26, Springer, 2014, 389–484.
- [22] M. Hutchings, *Beyond ECH capacities*, *Geometry and Topology* **20** (2016) 1085–1126.
- [23] M. Hutchings and C. H. Taubes, *Proof of the Arnold chord conjecture in three dimensions II*, *Geom, Topol.* **17** (2003), 2601–2688.
- [24] M. Hutchings and C. H. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders I*, *J. Symplectic Geom.* 5 (2007) 43–137.
- [25] M. Hutchings and C. H. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders II*, *J. Symplectic Geom.* 7 (2009) 29–133.
- [26] M. Hutchings and C. H. Taubes, *The Weinstein conjecture for stable Hamiltonian structures*, *Geom. Topol.* **13** (2009), 901–941.
- [27] P. Kronheimer, T. Mrowka, *Monopoles and three-manifolds*, *New Math. Monogr.* 10, Cambridge Univ. Press (2007).

- [28] R. Lipshitz, *A cylindrical reformulation of Heegaard Floer homology*, *Geom. Topol.* **10** (2006).
- [29] Y.-J. Lee and C. H. Taubes, *Periodic Floer homology and Seiberg–Witten–Floer cohomology*, *J. Symplectic Geom.* **10** (2012), no. 1, 81–164.
- [30] R. Leclercq and F. Zapolsky, *Spectral invariants for monotone Lagrangians*. *J. Topol. Anal.*, 10(3):627–700, 2018.
- [31] Y.-G. Oh *Symplectic topology and Floer homology. Vol. 2, New Mathematical Mono- graphs, vol. 28, Cambridge University Press, Cambridge, 2015, Symplectic geometry and pseudoholomorphic curves.*
- [32] C. H. Taubes, *Seiberg–Witten and Gromov invariants for symplectic 4-manifolds*, *First International Press Lecture Series 2, International Press, Somerville, MA (2000) MR1798809*
- [33] C. H. Taubes, *Embedded contact homology and Seiberg–Witten Floer cohomology I–V*, *Geometry and Topology* 14 (2010).
- [34] M. Usher, *Duality in filtered floer-novikov complexes*. *Journal of Topology and Analysis* (2011).