

# Work extraction from unknown quantum sources

Dominik Šafránek,<sup>1,\*</sup> Dario Rosa,<sup>1,2,†</sup> and Felix C. Binder<sup>3,‡</sup>

<sup>1</sup>*Center for Theoretical Physics of Complex Systems,  
Institute for Basic Science (IBS), Daejeon - 34126, Korea*

<sup>2</sup>*Basic Science Program, Korea University of Science and Technology (UST), Daejeon - 34113, Korea*

<sup>3</sup>*School of Physics, Trinity College Dublin, Dublin 2, Ireland*

(Dated: October 12, 2022)

Energy extraction is a central task in thermodynamics. In quantum physics, ergotropy measures the amount of work extractable under cyclic Hamiltonian control. As its full extraction requires perfect knowledge of the initial state, however, it does not characterize the work value of unknown or untrusted quantum sources. Fully characterizing such sources would require quantum tomography, which is prohibitively costly in experiments due to the exponential growth of required measurements and operational limitations. Here, we therefore derive a new notion of ergotropy applicable when nothing is known about the quantum states produced by the source, apart from what can be learned by performing only a single type of coarse-grained measurement. We find that in this case the extracted work is defined by the Boltzmann and observational entropy, in cases where the measurement outcomes are, or are not, used in the work extraction, respectively. This notion of ergotropy represents a realistic measure of extractable work, which can be used as the relevant figure of merit to characterize a quantum battery.

Efficient energy extraction is a key quest for living beings and modern technology alike. In recent years the advent of quantum technology has spurred the study of energy sources beyond the classical realm and the emerging field of quantum thermodynamics [1–6] has investigated the role of quantum features in this task. At the same time while modern quantum technology already finds applications in secure communication [7–9], sensing [10–12], and computing [13–17] these devices need to be powered, conceivably with non-equilibrium, quantum sources of energy. An example are recently experimentally-demonstrated [18, 19] quantum batteries [20–23], which offer a significant quantum advantage in charging power [24–28]. Generally, if one wants to make use of energy from an energy source, the first step is to characterize it. In the quantum regime, the energetic potential of the source is given by the quantum state it produces and measured by the Hamiltonian. Energy can be extracted by performing operations that transform this state into a state of lower energy, and collecting the surplus in the process.

Here, we consider a quantifier of work potential in the quantum regime called ergotropy  $W$  which equals the amount of energy extractable from a known quantum state  $\rho$  under the application of cyclic Hamiltonian control  $H(t)$  (where  $H(t) = 0$  for  $t < 0$  and  $t > T$  for a protocol duration  $T$ ) [29]. Since the resulting overall unitary evolution  $U$  is reversible, no entropy or heat is produced and the energy change  $\text{tr}[H(0)(\rho - U\rho U^\dagger)]$  exclusively manifests as work.

Ergotropy has been widely studied and measured in experiments [30, 31] where it quantifies the energy de-

posited onto a quantum load. However, a conceptual hurdle remains in this framework: the assumption of perfect knowledge of the state from which energy is extracted. In practice, the energy source may be unknown or uncharacterized and prohibit such idealized energy extraction. To fully characterize the state that is produced, one would require complete state tomography on a large number of identically-prepared states before the actual work extraction procedure, involving measurements in a number of non-commuting bases [32–34]. In many-body systems, which constitute quantum batteries, this number is enormous and thus these measurements are practically unrealistic. This is also the reason why entanglement entropy is difficult to measure, with a few exceptions in small-dimensional systems [35, 36]. In many-body systems, only the second order Rényi entropy has been measured instead [37–40]. In many experiments only limited types of measurements can be performed [37, 41–45]. Further, the required measurements can be prohibitively costly, may require long [46] or even infinite time [47], and may in fact be fundamentally incompatible with the laws of thermodynamics [48]. Thus, we here ask how much work may be extracted when only a single type of coarse-grained measurement is available to characterize the energy source.

We derive the corresponding quantities that describe the maximally extracted work under this operational constraint, and dub them Boltzmann and observational ergotropy, because they are implicitly defined by the average Boltzmann and observational entropy [49–53], respectively. The first applies to a situation when the measurement outcomes are employed in the work extraction process, while the second to a situation when the partially characterized source is no longer measured. Finally, we illustrate the effects of the operational constraints on work extraction from an evolving quantum state.

**Work extraction.** The amount of work that may be

\* [dsafrane@kibs.ac.kr](mailto:dsafrane@kibs.ac.kr)

† [dario\\_rosa@ibs.re.kr](mailto:dario_rosa@ibs.re.kr)

‡ [quantum@felix-binder.net](mailto:quantum@felix-binder.net)

extracted from a quantum system is contingent on what type of system manipulation is experimentally possible [54]. Here, we will be concerned with a quantum system's ergotropy, which measures the amount of energy that can be extracted by a unitary transformation. It is defined as [29]

$$W(\rho, H) = \text{tr}[H\rho] - \min_U \text{tr}[HU\rho U^\dagger]. \quad (1)$$

This can be written as a closed expression:

$$W(\rho, H) = \text{tr}[H(\rho - \pi)], \quad (2)$$

where, in the case of non-degenerate energy levels  $\pi$  is the unique passive state for the tuple  $(\rho, H)$ , meaning that it has the same eigenvalues as  $\rho$ , multiplying energy eigenvectors in decreasing order (in the degenerate case,  $\pi$  is a member of a family of passive states minimizing Eq. (1)). Interestingly, it is more efficient to extract the energy simultaneously from  $N$  copies, in which case we obtain an asymptotic expression [20, 55, 56],

$$W^\infty(\rho) := \lim_{N \rightarrow \infty} \frac{W(\rho^{\otimes N}, H_N)}{N} = \text{tr}[H(\rho - \rho_\beta)]. \quad (3)$$

Here,  $\rho_\beta = e^{-\beta H}/Z$  is the thermal state with inverse temperature  $\beta$  defined by the von Neumann entropy,  $S_{vN}(\rho_\beta) = S_{vN}(\rho)$ , and  $H_N := \sum_{i=1}^N h_i$  with  $h_i \cong H$  for all  $i$  (That is,  $H_N$  is a sum of isomorphic local terms; we notationally omit trivial terms on all other Hilbert spaces).  $\rho_\beta$  always lower-bounds  $\pi$  energetically:  $\text{tr}[H(\pi - \rho_\beta)] \geq 0$ . For the remainder of the paper we simplify the notation to  $W(\rho) \equiv W(\rho, H)$ .

Note that Eq. (3) is equivalent to the expression

$$\lim_{N \rightarrow \infty} \min_U \frac{\text{tr}[H_N U \rho^{\otimes N} U^\dagger]}{N} = \text{tr}[H\rho_\beta], \quad (4)$$

where  $S_{vN}(\rho_\beta) = S_{vN}(\rho)$ .

In the Supplemental Material, we derive its generalization,

$$\lim_{N \rightarrow \infty} \min_U \frac{\text{tr}[H_N U \otimes_i \rho_i^{\otimes p_i N} U^\dagger]}{N} = \text{tr}[H\rho_\beta], \quad (5)$$

where  $S_{vN}(\rho_\beta) = \sum_i p_i S_{vN}(\rho_i)$ , assuming that the limit exists [57].  $\{p_i\}$  is a set of probabilities,  $\sum_i p_i = 1$ , and  $\{\rho_i\}$  is any set of density matrices. The original formula is recovered by taking  $p_1 = 1$ .

**Extraction scheme.** Consider a scenario where only a single type of measurement is available to characterize the source of quantum states, potentially comprising significantly fewer outcomes than the system's microscopic degrees of freedom. In particular, consider a set of orthogonal projectors  $\mathcal{C} = \{P_i\}$  that sum to unity ( $\sum_i P_i = I$ ). The probability of outcome  $i$  when measured on an unknown state  $\rho$  is given by  $p_i = \text{tr}[P_i \rho]$ . After obtaining this outcome, the corresponding state is projected onto  $\rho_i = P_i \rho P_i / p_i$ . However, if one does not keep




FIG. 1. Scheme of work extraction from unknown sources. In the limit of the large number of iterations  $N$ , the ergotropy per state is larger in Stage 1 at the cost of having to store the measurement outcomes. It is given by  $W = \text{tr}[H(\rho - \rho_\beta)]$ , where  $\rho_\beta$  is a thermal state with temperature  $\beta$  implicitly defined by the mean Boltzmann entropy through  $S_{vN}(\rho_\beta) = S_C^B$  in Stage 1, and by observational entropy through  $S_{vN}(\rho_\beta) = S_C$  in Stage 2, respectively. See Eqs. (16) and (21). The energy  $\text{tr}[H\rho]$  is unknown, but it can be estimated from Eq. (24).

track of the outcome for each specific copy, the state after the measurement is best described by an averaged density matrix  $\rho_{av} = \sum_i p_i \rho_i = \sum_i P_i \rho P_i$ . The coarse-graining also naturally defines the decomposition of the Hilbert space into subspaces-macrostates,  $\mathcal{H} = \bigoplus_i \mathcal{H}_i$ , where each macrostate is given by

$$\mathcal{H}_i = P_i \mathcal{H} P_i = \text{span}_{|\psi\rangle} \{P_i |\psi\rangle\}. \quad (6)$$

We will study two work-extraction stages, depicted in Fig. 1:

**Stage 1.** The first stage combines characterization the source with work extraction. We characterize the source by measuring a sufficiently large number  $N$  of states. We record the outcomes and identify the corresponding probabilities  $p_i$ . Then, we extract work from these  $N$  states, by a protocol that makes use of the records.

**Stage 2.** In the second stage, we use the already characterized source to extract energy from the quantum states it produces, without further measurements. Given that we already know the probabilities  $p_i$ , we extract energy from any number of states produced by the source.

**Coarse-grained extraction operations.** The original notion of ergotropy assumes perfect knowledge of the density matrix from which energy is extracted. This is to design the best extraction unitary for that particular state. Here, however, the state produced by the source is unknown; only the outcomes  $i$ , or probabilities  $p_i$ , are known. Thus, we are unable to find the unitary that extracts the energy perfectly. We solve this by designing a protocol that creates a density matrix known to the ex-

perimeter, from which a certain amount of energy can be extracted.

The work extraction protocol consists of two parts: first, a random unitary that, on average, randomizes all states within each subspace, and then a global extraction unitary that extracts the remaining available energy,

$$U \bigoplus_i \tilde{U}_i. \quad (7)$$

$U$  is a unitary acting on the entire Hilbert space and  $\tilde{U}_i$  are random unitaries, each acting on the corresponding macrostate-subspace  $\mathcal{H}_i$ . We choose operations  $\tilde{U}_i$  to be completely random, i.e., we assume they are picked with the Haar measure. This choice ensures that averaging over this measure (computing the average over different realizations) leads to a non-unitary operation

$$\mathcal{U}(\rho) = U \left( \int \left( \bigoplus_i \tilde{U}_i \right) \rho \left( \bigoplus_i \tilde{U}_i^\dagger \right) d\mu \left( \bigoplus_i \tilde{U}_i \right) \right) U^\dagger = U \rho_{\text{cg}} U^\dagger. \quad (8)$$

$d\mu(\bigoplus_i \tilde{U}_i)$  is the Haar measure,

$$\rho_{\text{cg}} = \sum_i \frac{p_i}{V_i} P_i \quad (9)$$

is the coarse-grained state that resulted from the averaging, and  $V_i = \text{tr}[P_i] = \dim \mathcal{H}_i$  the volume of the macrostate (the number of its constituent distinct microstates). Note that the  $p_i$  are experimentally available, unlike the full original state  $\rho$ .

In the case of simultaneous extraction from multiple copies, we choose the random unitaries to act on each of the individual copies, mixing up every subspace so that the global extraction operation (before averaging) amounts to  $U(\bigoplus_i \tilde{U}_i)^{\otimes N}$ .  $U$  is a global unitary acting on an  $N$ -partite state. After averaging this corresponds to

$$\mathcal{U}(\rho^{(1)} \otimes \dots \otimes \rho^{(N)}) = U \rho_{\text{cg}}^{(1)} \otimes \dots \otimes \rho_{\text{cg}}^{(N)} U^\dagger. \quad (10)$$

**Extracted work in Stage 1: with measurement.** Consider the energetics of Stage 1. The total extracted work is given by the difference between the initial and the final energy of the state, which for measurement outcome  $i$  is

$$W_{\text{single}}^i = \text{tr}[H\rho] - \text{tr} \left[ HU \left( \bigoplus_j \tilde{U}_j \right) \rho_i \left( \bigoplus_j \tilde{U}_j^\dagger \right) U^\dagger \right]. \quad (11)$$

On average, the experimenter extracts

$$W^i = \int W_{\text{single}}^i d\mu \left( \bigoplus_j \tilde{U}_j \right) = \text{tr}[H\rho] - \text{tr} \left[ HU(P_i/V_i)U^\dagger \right]. \quad (12)$$

Here we used  $\mathcal{U}(\rho_i) = P_i/V_i$ . Maximizing over the global unitary and using Eq. (2), we define *Boltzmann ergotropy corresponding to outcome  $i$* ,

$$W_{\mathcal{C}}^{B,i}(\rho) = \max_U \text{tr} \left[ H(\rho - U(P_i/V_i)U^\dagger) \right] = \text{tr} \left[ H(\rho - \pi_i) \right]. \quad (13)$$

$\pi_i$  is a passive state for the tuple  $(P_i/V_i, H)$ . Averaging over all possible outcomes defines *Boltzmann ergotropy*,

$$W_{\mathcal{C}}^B(\rho) = \text{tr} \left[ H \left( \rho - \sum_i p_i \pi_i \right) \right]. \quad (14)$$

This quantity describes the maximal amount of work extractable from a state produced by an unknown source, when measuring the state and using the outcome for the coarse-grained extraction protocol [58].

In the case of simultaneous extraction in the limit of large  $N$ , the state after a series of measurements is (*almost surely*) described by

$$\rho_N = \bigotimes_i \rho_i^{\otimes p_i N} \quad (15)$$

This is up reordering the outcomes: while the order affects the specific extraction unitary, all of such permutations lead to the same ergotropy (since these states are energetically equivalent). This is derived by using the law of large numbers; see Supplemental Material.

Using the same logic as in Eq. (12) when applying Eq. (10) on Eq. (15), and then using Eq. (5) we derive the Boltzmann ergotropy in the large- $N$  limit,

$$W_{\mathcal{C}}^{B\infty}(\rho) := \lim_{N \rightarrow \infty} \frac{W_{\mathcal{C}}^B(\rho^{\otimes N})}{N} = \text{tr} \left[ H(\rho - \rho_\beta) \right]. \quad (16)$$

Temperature of the thermal state  $\rho_\beta = e^{-\beta H}/Z$  is implicitly defined by requiring that its von Neumann entropy equals the mean *Boltzmann entropy* with coarse-graining  $\mathcal{C}$ ,

$$S_{\text{vN}}(\rho_\beta) = \sum_i p_i \ln V_i =: S_{\mathcal{C}}^B. \quad (17)$$

See Supplemental Material for details. The extractable work depends largely on the amount of coarse-graining. For a fine-grained measurement projecting onto a pure state, we have  $V_i = 1$  and thus  $S_{\mathcal{C}}^B = 0$ . This means that all the mean energy can be extracted. On the other hand, for a very coarse measurement in which  $S_{\mathcal{C}}^B$  is large, very little can be gained. Energy can even be lost, if  $\text{tr}[H\rho_\beta] > \text{tr}[H\rho]$ .

**Extracted work in Stage 2: no measurement.** Assuming that the source has been characterized, with probabilities  $p_i$  known, how much energy can be extracted without further measurement? As in Eq. (12), we derive this energy as the energy difference between the initial and the final state,

$$W_{\text{single}}^{\text{no-meas.}} = \text{tr}[H\rho] - \text{tr} \left[ HU \left( \bigoplus_i \tilde{U}_i \right) \rho \left( \bigoplus_i \tilde{U}_i^\dagger \right) U^\dagger \right]. \quad (18)$$

Averaging over the random unitaries using Eq. (8), and maximizing over the extraction unitary defines *observational ergotropy*,

$$W_{\mathcal{C}}(\rho) = \text{tr}[H\rho] - \min_U \text{tr} \left[ HU \rho_{\text{cg}} U^\dagger \right] = \text{tr} \left[ H(\rho - \pi_{\text{cg}}) \right]. \quad (19)$$

$\pi_{\text{cg}}$  is a passive state for the tuple  $(\rho_{\text{cg}}, H)$ . This quantity describes the maximal extractable work from an unknown source, characterized by a set of probabilities  $\{p_i\}$ , without further measurements.

In the case of simultaneous extraction, we have

$$\begin{aligned} W_{\mathcal{C}}(\rho^{\otimes N}) &= \text{tr}[H_N \rho^{\otimes N}] - \min_U \text{tr}[H_N \mathcal{U}(\rho^{\otimes N})] \\ &= N \text{tr}[H \rho] - \min_U \text{tr}[H_N U \rho_{\text{cg}}^{\otimes N} U^\dagger]. \end{aligned} \quad (20)$$

From this, we obtain observational ergotropy in the large- $N$  limit,

$$W_{\mathcal{C}}^\infty(\rho) := \lim_{N \rightarrow \infty} \frac{W_{\mathcal{C}}(\rho^{\otimes N})}{N} = \text{tr}[H(\rho - \rho_{\beta'})]. \quad (21)$$

As per Eq. (4), temperature  $\beta'$  of the thermal state is implicitly defined by requiring that its von Neumann entropy equals the observational entropy  $S_{\mathcal{C}}$  [52],

$$\begin{aligned} S_{\text{vN}}(\rho_{\beta'}) &= S_{\text{vN}}(\rho_{\text{cg}}) = - \sum_i p_i \ln p_i + \sum_i p_i \ln V_i \\ &= S_{\mathcal{C}}^{\text{Sh}} + S_{\mathcal{C}}^{\text{B}} = S_{\mathcal{C}}, \end{aligned} \quad (22)$$

which is the sum of Shannon entropy and the average Boltzmann entropy. Due to the conditional extraction, Stage 1 leads to a larger extractable work than Stage 2, where the difference in Eqs. (17) and (22) is given by the Shannon entropy of measurement. As a result of identity  $\mathcal{U}(\rho) = \mathcal{U}(\rho_{\text{av}})$ , if in Stage 1 the measurement outcomes are not used in the extraction, the same protocol can be employed in both stages, leading to the same extracted energy. An example of observational ergotropy and the corresponding entropy is depicted in Fig. 2.

**Bounds on observational ergotropy.** To compute the Boltzmann or observational ergotropy, one needs to know the mean initial energy, which depends on the unknown density matrix  $\rho$ . While full knowledge of the state is not available, it is possible to estimate it from the partial knowledge given by distribution  $\{p_i\}$ .

Consider local energy coarse-grainings,

$$\mathcal{C} = \{P_{E_1} \otimes P_{E_2}\}, \quad (23)$$

studied in observational entropy literature [50, 52].  $P_{E_1} = \sum_{E'_1 \in [E_1, E_1 + \Delta E]} |E'_1\rangle\langle E'_1|$  and  $P_{E_2}$  (analogous) are the coarse-grained projectors on local energies with resolution  $\Delta E$ . The mean energy can be estimated from

$$|\text{tr}[H \rho] - \text{tr}[H \rho_{\text{cg}}]| \leq 2 \|H_{\text{int}}\| + 2\Delta E, \quad (24)$$

where the full Hamiltonian is  $H = H_1 + H_2 + H_{\text{int}}$ , and  $\|\cdot\|$  denotes the operator norm. When increasing the number of partitions to  $k$  instead of 2, we obtain  $|\text{tr}[H \rho] - \text{tr}[H \rho_{\text{cg}}]| \leq 2 \|H_{\text{int}}\| + k\Delta E$ . The first term  $\|H_{\text{int}}\|$  also scales linearly with  $k$ . Thus, this correction represents a finite-size effect. See Supplemental Material for details.

Energy can be estimated also in the case of a completely general coarse-graining [59].




FIG. 2. Observational ergotropy and observational entropy as a function of time, for a thermalizing system of four particles. The coarse-graining  $\mathcal{C}$  is chosen to be a local energy coarse-graining, Eq. (23), and the initial state  $|\psi_1\rangle \otimes |\psi_2\rangle$ ,  $|\psi_1\rangle = |000000\rangle$  and  $|\psi_2\rangle = |111100\rangle$ , evolves with the Hamiltonian given by Eq. (25), with  $T = V = 1$  and  $T' = V' = 0.96$ . We choose the resolution in energy  $\Delta E = (E_1 - E_0)/2$ , where  $E_0$  and  $E_1$  are the ground state and the first excited state energy. True ergotropy  $W_{\mathcal{C}}^\infty(\rho)$  (solid purple) is unknown to the experimenter. However, they can estimate the value of  $W_{\mathcal{C}}^\infty(\rho_{\text{cg}})$  (red dotted),  $\rho_{\text{cg}} = \sum_{E_1, E_2} \frac{P_{E_1} E_2(t)}{V_{E_1} V_{E_2}} P_{E_1} \otimes P_{E_2}$ , and be sure that the true value lies within the light-red shaded region, in between lines  $W_{\mathcal{C}}^\infty(\rho_{\text{cg}}) \pm 2(\|H_{\text{int}}\| + \Delta E)$  (red dashed). We compare this with observational entropy  $S_{\mathcal{C}}(\rho)$  (blue long-dashed). For this particular coarse-graining, it is called non-equilibrium thermodynamic entropy [52], and it is bounded by the sum of initial thermodynamic entropies (dark green dot-dashed) and by the final thermodynamic entropy (green dot-dashed),  $S_{\text{th}} \equiv S_{\mathcal{C}_E}$ , which is defined as observational entropy with global energy coarse-graining with the same resolution  $\Delta E$ .

**Example: ergotropy of local energy coarse-graining.** We illustrate the essence of our result by showing observational ergotropy as a function of time for a thermalizing system. We assume that we can measure local energies, described by coarse-graining (23). A suitable model is given by a widely-used and generic one-dimensional fermionic Hamiltonian [60], with the nearest neighbor and next-nearest neighbor interaction, describing interacting particles hopping between  $k$ 'th and  $l$ 'th site as

$$H^{(k:l)} = \sum_{i=k}^l -T f_i^\dagger f_{i+1} - T' f_i^\dagger f_{i+2} + h.c. + V n_i n_{i+1} + V' n_i n_{i+2}. \quad (25)$$

(Terms with  $f_{l+1}$ ,  $f_{l+2}$ ,  $f_{k-1}$ , and  $f_{k-2}$  are not included in the sum.)  $f_i$  and  $f_i^\dagger$  are the fermionic annihilation and creation operators for site  $i$ .  $n_i = f_i^\dagger f_i$  is the local density operator. We take the full Hamiltonian  $H \equiv H^{(1:L)}$ , where  $L$  is the length of the chain, with a division between two equal sized subsystems  $H_1 \equiv H^{(1:L/2)}$ ,  $H_2 \equiv H^{(L/2+1:L)}$ , and  $H_{\text{int}} = H - H_1 - H_2$ . We also employ

hard wall boundary conditions.

In Fig. 2 we plot observational ergotropy  $W_C^\infty(|\psi_t\rangle)$  for extracting energy from state  $|\psi_t\rangle = \exp(-iHt)|\psi_0\rangle$ , together with its estimates. We compare this with observational entropy, which is naturally connected to observational ergotropy through Eq. (21), to which it is reciprocally related. We can spot the point where the lower bound on ergotropy crosses zero, in which the experimenter, given the available information, will stop characterizing the state as useful, i.e., they cannot be certain that it provides energy. As the state evolves, the system thermalizes, and the opportunity to extract work diminishes.

The full Hamiltonian preserves the total number  $n$  of particles, thus the relevant Hilbert space explored by the system during time evolution is  $\mathcal{H} = \bigoplus_{k=0}^n \mathcal{H}_k \otimes \mathcal{H}_{n-k}$ . This is important for correctly computing the accessible macrostate volumes  $V_{E_1 E_2}$ . This condition can be relaxed to apply to situations with an unknown particle number, by employing additional coarse-graining in local particle numbers (as suggested in [52]).

**Discussion and Conclusions.** To relate ergotropy to realistic scenarios of work extraction, previous works have considered constraints such as a restriction to local [20, 24, 25, 61–63], Gaussian [64], or incoherent operations [65]. Here, we have addressed the remaining open problem which is the assumption of perfect knowledge of the initial state.

Assuming that a source of unknown states can be characterized only by a single type of coarse-grained measurement, we designed the extraction protocol as follows. A random unitary is applied on each measurement subspace. In effect, this creates a uniform state in each subspace, turning the unknown state into a known coarse-grained state. Then we apply a global unitary operation that optimizes the energy extraction from the coarse-grained state.

The protocol results in two notions of ergotropy: Boltzmann ergotropy, which measures the extracted energy when the measurement result is conditionally taken into account (Eq. 16), and observational ergotropy for unconditional extraction (Eq. 21). The energy difference between the two cases results from the difference in entropy of corresponding thermal states (Eq. 17 and Eq. 22). The two deviate exactly by the (Shannon) entropy of

measurement  $S^{Sh} = -\sum_i p_i \ln p_i$  that lower-bounds the work required for erasing the measurement record [66–68]:  $W_{\text{erasure}} \geq \beta^{-1} S^{Sh}$  with  $\beta$  the inverse temperature of the heat bath used for the erasure process [69].

This work applies in two cases. First is the case of a high-dimensional external source, i.e., not prepared by an experimenter, on which full quantum tomography is not viable. The second case is that of imperfectly controlled systems, such as quantum batteries. Having perfect control over the charging procedure, one knows exactly the state of the charged battery and therefore ergotropy is the relevant figure of merit [70–73]. However, certain lack of control is inevitable, for example in the form of an unknown disorder [27, 74–80], uncertain time of charging [81, 82], or batteries charged from an initially unknown state [83]. In all these cases, the final battery state is not fully determined. Observational ergotropy then gives an experimentally verifiable lower bound on the amount of energy that can be extracted. As such, it is a realistic figure of merit for characterizing quantum batteries.

One can also find applications from the theoretical perspective. These appear whenever there is a limit on which measurement the experimenter can or is willing to do, much alongside E.T. Jaynes’ statement regarding the Gibbs (mixing) paradox [84]: “The amount of useful work that we can extract from any system depends - obviously and necessarily - on how much ”subjective” information we have about its microstate because that tells us which interactions will extract energy and which will not.” Here as well, depending on the measurement, the outcome will determine the best unitary to perform and in turn the amount of work the experimenter can extract.

**Acknowledgments.** DR and DŠ acknowledge the support from the Institute for Basic Science in Korea (IBS-R024-D1). FCB acknowledges support from grant number FQXi-RFP-IPW-1910 from the Foundational Questions Institute and Fetzer Franklin Fund, a donor-advised fund of the Silicon Valley Community Foundation. DŠ thanks Anthony Aguirre, Joshua M. Deutsch, Joseph Schindler, and Susanne Still for discussing theory years in the making. We acknowledge Philipp Strasberg for the excellent feedback on the first version of this manuscript and for discussing with us the examples mentioned in the Supplemental Material.

- 
- [1] J. Gemmer, M. Michel, and G. Mahler, *Quantum thermodynamics: Emergence of thermodynamic behavior within composite quantum systems*, Vol. 784 (Springer, 2009).
  - [2] R. Kosloff, *Entropy* **15**, 2100 (2013).
  - [3] J. Goold, M. Huber, A. Riera, L. del Rio, and P. Skrzypczyk, *Journal of Physics A: Mathematical and Theoretical* **49**, 143001 (2016).
  - [4] S. Vinjanampathy and J. Anders, *Contemporary Physics* **57**, 545 (2016).
  - [5] F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso, *Fundamental Theories of Physics* **195**, 1 (2018).
  - [6] S. Deffner and S. Campbell, *Quantum Thermodynamics*, 2053-2571 (Morgan & Claypool Publishers, 2019).
  - [7] M. Minder, M. Pittaluga, G. Roberts, M. Lucamarini, J. Dynes, Z. Yuan, and A. Shields, *Nature Photonics* **13**, 334 (2019).

- [8] S. Pirandola, U. L. Andersen, L. Banchi, M. Berta, D. Bunandar, R. Colbeck, D. Englund, T. Gehring, C. Lupo, C. Ottaviani, J. L. Pereira, M. Razavi, J. S. Shaari, M. Tomamichel, V. C. Usenko, G. Vallone, P. Villoresi, and P. Wallden, *Adv. Opt. Photon.* **12**, 1012 (2020).
- [9] Y.-A. Chen, Q. Zhang, T.-Y. Chen, W.-Q. Cai, S.-K. Liao, J. Zhang, K. Chen, J. Yin, J.-G. Ren, Z. Chen, *et al.*, *Nature* **589**, 214 (2021).
- [10] P. Cheiney, L. Fouché, S. Templier, F. Napolitano, B. Battelier, P. Bouyer, and B. Barrett, *Phys. Rev. Applied* **10**, 034030 (2018).
- [11] V. Ménoret, P. Vermeulen, N. Le Moigne, S. Bonvalot, P. Bouyer, A. Landragin, and B. Desruelle, *Scientific reports* **8**, 1 (2018).
- [12] M. Tse *et al.*, *Phys. Rev. Lett.* **123**, 231107 (2019).
- [13] F. Arute *et al.*, *Nature (London)* **574**, 505 (2019), [arXiv:1910.11333 \[quant-ph\]](https://arxiv.org/abs/1910.11333).
- [14] H.-S. Zhong, H. Wang, Y.-H. Deng, M.-C. Chen, L.-C. Peng, Y.-H. Luo, J. Qin, D. Wu, X. Ding, Y. Hu, P. Hu, X.-Y. Yang, W.-J. Zhang, H. Li, Y. Li, X. Jiang, L. Gan, G. Yang, L. You, Z. Wang, L. Li, N.-L. Liu, C.-Y. Lu, and J.-W. Pan, *Science* **370**, 1460 (2020).
- [15] D. Maslov, J.-S. Kim, S. Bravyi, T. J. Yoder, and S. Sheldon, *Nature Physics* **17**, 894 (2021), [arXiv:2008.06478 \[quant-ph\]](https://arxiv.org/abs/2008.06478).
- [16] J. Randall, C. E. Bradley, F. V. van der Gronden, A. Galicia, M. H. Abobeih, M. Markham, D. J. Twitchen, F. Machado, N. Y. Yao, and T. H. Taminiou, *Science* **374**, 1474 (2021).
- [17] X. Mi *et al.*, *Nature (London)* **601**, 531 (2022), [arXiv:2107.13571 \[quant-ph\]](https://arxiv.org/abs/2107.13571).
- [18] J. Q. Quach, K. E. McGhee, L. Ganzer, D. M. Rouse, B. W. Lovett, E. M. Gauger, J. Keeling, G. Cerullo, D. G. Lidzey, and T. Virgili, *Science Advances* **8**, eabk3160 (2022).
- [19] C.-K. Hu, J. Qiu, P. J. P. Souza, J. Yuan, Y. Zhou, L. Zhang, J. Chu, X. Pan, L. Hu, J. Li, Y. Xu, Y. Zhong, S. Liu, F. Yan, D. Tan, R. Bachelard, C. J. Villas-Boas, A. C. Santos, and D. Yu, *arXiv e-prints* (2021), [arXiv:2108.04298 \[quant-ph\]](https://arxiv.org/abs/2108.04298).
- [20] R. Alicki and M. Fannes, *Phys. Rev. E* **87**, 042123 (2013).
- [21] F. Campaioli, F. A. Pollock, and S. Vinjanampathy, in *Fundamental Theories of Physics*, Vol. 195, edited by F. Binder, L. A. Correa, C. Gogolin, J. Anders, and G. Adesso (Springer, 2018) pp. 207–225.
- [22] S. Bhattacharjee and A. Dutta, *The European Physical Journal B* **94**, 1 (2021).
- [23] H.-L. Shi, S. Ding, Q.-K. Wan, X.-H. Wang, and W.-L. Yang, *Phys. Rev. Lett.* **129**, 130602 (2022), [arXiv:2205.11080 \[quant-ph\]](https://arxiv.org/abs/2205.11080).
- [24] F. C. Binder, S. Vinjanampathy, K. Modi, and J. Goold, *New Journal of Physics* **17**, 75015 (2015).
- [25] F. Campaioli, F. A. Pollock, F. C. Binder, L. Céleri, J. Goold, S. Vinjanampathy, and K. Modi, *Phys. Rev. Lett.* **118**, 150601 (2017).
- [26] D. Ferraro, M. Campisi, G. M. Andolina, V. Pellegrini, and M. Polini, *Phys. Rev. Lett.* **120**, 117702 (2018).
- [27] D. Rossini, G. M. Andolina, D. Rosa, M. Carrega, and M. Polini, *Phys. Rev. Lett.* **125**, 236402 (2020).
- [28] J.-Y. Gyhm, D. Šafránek, and D. Rosa, *Phys. Rev. Lett.* **128**, 140501 (2022).
- [29] A. E. Allahverdyan, R. Balian, and T. M. Nieuwenhuizen, *Europhysics Letters (EPL)* **67**, 565 (2004).
- [30] D. Von Lindenfels, O. Gräß, C. T. Schmiegelow, V. Kaushal, J. Schulz, M. T. Mitchison, J. Goold, F. Schmidt-Kaler, and U. G. Poschinger, *Physical Review Letters* **123**, 80602 (2019).
- [31] N. Van Horne, D. Yum, T. Dutta, P. Hänggi, J. Gong, D. Poletti, and M. Mukherjee, *npj Quantum Information* **6**, 37 (2020), [arXiv:1812.01303](https://arxiv.org/abs/1812.01303).
- [32] G. M. D’Ariano, M. G. Paris, and M. F. Sacchi, *Advances in Imaging and Electron Physics* **128**, 206 (2003).
- [33] R. B. A. Adamson and A. M. Steinberg, *Phys. Rev. Lett.* **105**, 030406 (2010).
- [34] E. Toninelli, B. Ndagano, A. Vallés, B. Sephton, I. Nape, A. Ambrosio, F. Capasso, M. J. Padgett, and A. Forbes, *Adv. Opt. Photon.* **11**, 67 (2019).
- [35] B. Lanyon, C. Maier, M. Holzäpfel, T. Baumgratz, C. Hempel, P. Jurcevic, I. Dhand, A. Buyskikh, A. Daley, M. Cramer, *et al.*, *Nature Physics* **13**, 1158 (2017).
- [36] C. A. Sackett, D. Kielpinski, B. E. King, C. Langer, V. Meyer, C. J. Myatt, M. Rowe, Q. Turchette, W. M. Itano, D. J. Wineland, *et al.*, *Nature* **404**, 256 (2000).
- [37] A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, *Science* **353**, 794 (2016).
- [38] T. Brydges, A. Elben, P. Jurcevic, B. Vermersch, C. Maier, B. P. Lanyon, P. Zoller, R. Blatt, and C. F. Roos, *Science* **364**, 260 (2019).
- [39] R. Islam, R. Ma, P. M. Preiss, M. Eric Tai, A. Lukin, M. Rispoli, and M. Greiner, *Nature* **528**, 77 (2015).
- [40] G.-X. Su, H. Sun, A. Hudomal, J.-Y. Desaulles, Z.-Y. Zhou, B. Yang, J. C. Halimeh, Z.-S. Yuan, Z. Papić, and J.-W. Pan, *arXiv e-prints*, [arXiv:2201.00821 \(2022\)](https://arxiv.org/abs/2201.00821), [arXiv:2201.00821 \[cond-mat.quant-gas\]](https://arxiv.org/abs/2201.00821).
- [41] S. Hofferberth, I. Lesanovsky, B. Fischer, T. Schumm, and J. Schmiedmayer, *Nature* **449**, 324 (2007).
- [42] A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, *Rev. Mod. Phys.* **83**, 863 (2011).
- [43] S. Trotzky, Y.-A. Chen, A. Fleisch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, *Nature physics* **8**, 325 (2012).
- [44] M. Schreiber, S. S. Hodgman, P. Bordia, H. P. Lüschen, M. H. Fischer, R. Vosk, E. Altman, U. Schneider, and I. Bloch, *Science* **349**, 842 (2015).
- [45] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, *Nature (London)* **551**, 579 (2017), [arXiv:1707.04344 \[quant-ph\]](https://arxiv.org/abs/1707.04344).
- [46] P. Strasberg, K. Modi, and M. Skotiniotis, *European Journal of Physics* **43**, 035404 (2022).
- [47] P. Busch, *Foundations of physics* **20**, 1 (1990).
- [48] Y. Guryanova, N. Friis, and M. Huber, *Quantum* **4**, 222 (2020), [arXiv:1805.11899](https://arxiv.org/abs/1805.11899).
- [49] D. Šafránek, J. M. Deutsch, and A. Aguirre, *Physical Review A* **99**, 010101(R) (2019), [arXiv:1707.09722](https://arxiv.org/abs/1707.09722).
- [50] D. Šafránek, J. M. Deutsch, and A. Aguirre, *Phys. Rev. A* **99**, 012103 (2019), [arXiv:1803.00665 \[quant-ph\]](https://arxiv.org/abs/1803.00665).
- [51] P. Strasberg and A. Winter, *PRX Quantum* **2**, 030202 (2021), [arXiv:2002.08817 \[quant-ph\]](https://arxiv.org/abs/2002.08817).
- [52] D. Šafránek, A. Aguirre, J. Schindler, and J. M. Deutsch, *Foundations of Physics* **51**, 101 (2021), [arXiv:2008.04409](https://arxiv.org/abs/2008.04409).
- [53] F. Buscemi, J. Schindler, and D. Šafránek, [arXiv:2209.03803 \(2022\)](https://arxiv.org/abs/2209.03803), [10.48550/arXiv.2209.03803](https://arxiv.org/abs/2209.03803).
- [54] W. Niedenzu, M. Huber, and E. Boukobza, *Quantum* **3**, 195 (2019), [arXiv:1907.01353](https://arxiv.org/abs/1907.01353).
- [55] K. V. Hovhannisyanyan, M. Perarnau-Llobet, M. Huber,

- and A. Acín, *Phys. Rev. Lett.* **111**, 240401 (2013), [arXiv:1303.4686 \[quant-ph\]](#).
- [56] F. Campaioli, F. A. Pollock, and S. Vinjanampathy, in *Thermodynamics in the Quantum Regime* (Springer, 2018) pp. 207–225.
- [57] Here and in the following,  $\rho_i^{\otimes p_i N}$  is shorthand for  $\rho_i^{\otimes [p_i N]}$  and  $\rho_i^{\otimes 0} = 1$  inside the limit. See Supplemental Material for details.
- [58] Note here, that this quantity does not include the (Landauer) work that would be required to reset the measurement record [85].
- [59] D. Šafránek et al., “Energy measurement from any other measurement (in preparation),” (2022).
- [60] L. F. Santos and M. Rigol, *Phys. Rev. E* **81**, 036206 (2010), [arXiv:0910.2985 \[cond-mat.stat-mech\]](#).
- [61] K. V. Hovhannisyán, M. Perarnau-Llobet, M. Huber, and A. Acín, *Physical Review Letters* **111**, 240401 (2013), [arXiv:arXiv:1303.4686](#).
- [62] M. Alimuddin, T. Guha, and P. Parashar, *Phys. Rev. A* **99**, 052320 (2019).
- [63] S. Puliylil, M. Banik, and M. Alimuddin, *Phys. Rev. Lett.* **129**, 070601 (2022).
- [64] E. G. Brown, N. Friis, and M. Huber, *New Journal of Physics* **18**, 113028 (2016), [arXiv:1608.04977](#).
- [65] G. Francica, F. C. Binder, G. Guarnieri, M. T. Mitchison, J. Goold, and F. Plastina, *Physical Review Letters* **125**, 180603 (2020), [arXiv:2006.05424](#).
- [66] T. Sagawa and M. Ueda, *Physical Review Letters* **102**, 250602 (2009).
- [67] D. Reeb and M. M. Wolf, *New Journal of Physics* **16**, 103011 (2014), [arXiv:1306.4352](#).
- [68] J. M. R. Parrondo, J. M. Horowitz, and T. Sagawa, *Nature Physics* **11**, 131 (2015).
- [69] Note that we have here *not* attempted to quantify the energy associated with the measurement itself (which indeed diverges for perfect projective measurement [48]).
- [70] G. M. Andolina, M. Keck, A. Mari, M. Campisi, V. Giovannetti, and M. Polini, *Phys. Rev. Lett.* **122**, 047702 (2019).
- [71] F. Barra, *Phys. Rev. Lett.* **122**, 210601 (2019).
- [72] K. V. Hovhannisyán, F. Barra, and A. Imparato, *Phys. Rev. Research* **2**, 033413 (2020).
- [73] A. Delmonte, A. Crescente, M. Carrega, D. Ferraro, and M. Sassetti, *Entropy* **23** (2021).
- [74] D. Rossini, G. M. Andolina, and M. Polini, *Phys. Rev. B* **100**, 115142 (2019).
- [75] S. Ghosh, T. Chanda, and A. Sen(De), *Phys. Rev. A* **101**, 032115 (2020).
- [76] D. Rosa, D. Rossini, G. M. Andolina, M. Polini, and M. Carrega, *Journal of High Energy Physics* **2020**, 67 (2020).
- [77] F. Caravelli, G. Coulter-De Wit, L. P. García-Pintos, and A. Hamma, *Phys. Rev. Research* **2**, 023095 (2020).
- [78] F. Zhao, F.-Q. Dou, and Q. Zhao, *Phys. Rev. A* **103**, 033715 (2021).
- [79] J. Kim, J. Murugan, J. Olle, and D. Rosa, *Phys. Rev. A* **105**, L010201 (2022).
- [80] M. B. Arjmandi, H. Mohammadi, and A. C. Santos, *Phys. Rev. E* **105**, 054115 (2022).
- [81] M. T. Mitchison, J. Goold, and J. Prior, *Quantum* **5**, 500 (2021).
- [82] Y. Yao and X. Q. Shao, *Phys. Rev. E* **106**, 014138 (2022).
- [83] G. T. Landi, *Entropy* **23**, 1627 (2021), [arXiv:2110.10512 \[quant-ph\]](#).
- [84] E. T. Jaynes, “The Gibbs Paradox,” in *Maximum Entropy and Bayesian Methods: Seattle, 1991*, edited by C. R. Smith, G. J. Erickson, and P. O. Neudorfer (Springer Netherlands, Dordrecht, 1992) pp. 1–21.
- [85] R. Landauer, *IBM Journal* **July**, 183 (1961).
- [86] T. M. Cover, *Elements of information theory* (John Wiley & Sons, 1999).
- [87] M. Lundberg and L. Svensson, in *Processing Workshop Proceedings, 2004 Sensor Array and Multichannel Signal* (IEEE, 2004) pp. 114–118.

## SUPPLEMENTAL MATERIAL

In this Supplemental Material, we provide several proofs and derivations, as well as examples. It contains: Sec. I: generalized formula for the maximal extracted work, Sec. II: limit state after  $N$  measurements in stage 1, Sec. III: Boltzmann ergotropy in the large  $N$  limit, Sec. IV: examples on a three-level system, with a special focus on understanding the coarse-grained unitary, Sec. V: bound on energy for local energy coarse-graining.

### I. GENERALIZATION OF THE MAXIMALLY EXTRACTED WORK

Here, we will derive a generalization of the formula for the maximally extracted work,

$$\lim_{N \rightarrow \infty} \min_U \frac{\text{tr}[H_N U \otimes_i \rho_i^{\otimes p_i N} U^\dagger]}{N} = \text{tr}[H \rho_\beta], \quad (26)$$

where  $S_{vN}(\rho_\beta) = \sum_i p_i S_{vN}(\rho_i)$ , assuming that the limit exists.  $\{p_i\}$  is a set of probabilities,  $\sum_i p_i = 1$ , and  $\{\rho_i\}$  is any set of density matrices. Eq. (26) corresponds to Eq. (5) in the main text.

As  $p_i N$  is not necessarily an integer (which is irrelevant for large  $N$ ), to be mathematically exact, we define

$$\rho^N \equiv \bigotimes_i \rho_i^{\otimes p_i N} := \rho_1^{\otimes (N - \sum_{i \geq 2} \lfloor p_i N \rfloor)} \bigotimes_{i \geq 2} \rho_i^{\otimes \lfloor p_i N \rfloor}. \quad (27)$$

This ensures that there are exactly  $N$  states and each state is exponentiated to an integer.

The Hamiltonian is defined as

$$H_N := \sum_{i=1}^N h_i = H \otimes I \otimes \dots \otimes I + I \otimes H \otimes I \otimes \dots \otimes I + \dots \quad (28)$$

with  $h_i \cong H$  for all  $i$  (that is,  $H_N$  is a sum of isomorphic local terms).

We have to assume that the limit in Eq. (26) exists because, while it seems quite natural, it is difficult to prove. Unlike in the original statement, the sequence (26) is not monotonously decreasing. This is because of implementation of  $\rho^N$  using integers, Eq. (27). Elements of the sequence may temporarily spike due to the discrete nature  $\rho^N$ , even though the sequence is expected to go down most of the time. It is clear though that the right hand side of Eq. (26) is a lower bound, which follows from the variational principle of statistical mechanics, which asserts that the Gibbs canonical density matrix minimizes the free energy,

$$\text{tr}[\rho H] - \beta^{-1} S_{vN}(\rho) \geq \text{tr}[\rho_\beta H] - \beta^{-1} S_{vN}(\rho_\beta). \quad (29)$$

To prove Eq. (26), we are going to show that for any  $\epsilon > 0$  we find an element of the sequence closer to  $\text{tr}[H \rho_\beta]$  than  $\epsilon$ . Because we assume that the limit exists, it must be equal to this number. We assume that  $\rho^N$  is given by

Eq. (27), although the exact way how the  $\rho^N$  is represented using floor functions in order to be properly defined will not matter for the argument.

We pick a subsequence of the sequence obtained by substitution  $N \rightarrow MN$  (product of  $M$  and  $N$ ). This gives

$$\rho^{MN} = \rho_1^{\otimes (MN - \sum_{i \geq 2} \lfloor p_i MN \rfloor)} \bigotimes_{i \geq 2} \rho_i^{\otimes \lfloor p_i MN \rfloor}. \quad (30)$$

For now, we assume that  $M$  is some large integer. Consider a similar state

$$(\rho^M)^{\otimes N} = \left( \rho_1^{\otimes (M - \sum_{i \geq 2} \lfloor p_i M \rfloor)} \bigotimes_{i \geq 2} \rho_i^{\otimes \lfloor p_i M \rfloor} \right)^{\otimes N}. \quad (31)$$

Energies of  $\rho^{MN}$  and  $(\rho^M)^{\otimes N}$  are very similar. In fact, we have

$$\text{tr}[H_{MN} \rho^{MN}] = \text{tr}[H_{MN} (\rho^M)^{\otimes N}] + k(NMg_M + Nf_N). \quad (32)$$

where  $\lim_{N \rightarrow \infty} f_N = \lim_{M \rightarrow \infty} g_M = 0$  and  $k$  is the number of different  $i$ 's (number of elements in sets  $\{p_i\}$  or  $\{\rho_i\}$ ; this is by construction independent of  $N$  and  $M$ ). This is because the exponents associated with a given  $i$  differ only slightly for large enough  $N$  and  $M$  (Recall that permutation of density matrices does not change the energy.) For example, for an  $i \geq 2$  we have

$$\begin{aligned} \lfloor p_i MN \rfloor - \lfloor p_i M \rfloor N &= N \left( \frac{\lfloor p_i MN \rfloor}{N} - \lfloor p_i M \rfloor \right) \\ &= N(p_i M - \lfloor p_i M \rfloor + f_N) \\ &= NM \left( p_i - \frac{\lfloor p_i M \rfloor}{M} \right) + Nf_N \\ &= NMg_M + Nf_N \end{aligned} \quad (33)$$

where  $\lim_{N \rightarrow \infty} f_N = \lim_{M \rightarrow \infty} g_M = 0$ , because  $\lim_{N \rightarrow \infty} \frac{\lfloor xN \rfloor}{N} = x$ . Since there are  $k$  indices, the total difference is of order  $k$  times the above.

According to  $\lim_{N \rightarrow \infty} \min_U \text{tr}[H_N U \rho^N U^\dagger]/N = \text{tr}[H \rho_\beta]$ , where  $S_{vN}(\rho_\beta) = S_{vN}(\rho)$ , (Eq. (4) in the main text), we have

$$\lim_{N \rightarrow \infty} \min_U \frac{\text{tr}[H_{MN} U (\rho^M)^{\otimes N} U^\dagger]}{MN} = \frac{\text{tr}[H_M \rho_{\beta'}^{\otimes M}]}{M} = \text{tr}[H \rho_{\beta'}]. \quad (34)$$

where  $S_{vN}(\rho_{\beta'}^{\otimes M}) = S_{vN}(\rho^M)$ , which written explicitly gives

$$S_{vN}(\rho_{\beta'}) = \left( 1 - \sum_{i \geq 2} \frac{\lfloor p_i M \rfloor}{M} \right) S(\rho_1) + \sum_{i \geq 2} \frac{\lfloor p_i M \rfloor}{M} S(\rho_i). \quad (35)$$

Eq. (32) also holds for the transformed states, since the same argument can be made. Thus we have,

$$\begin{aligned} &\lim_{N \rightarrow \infty} \min_U \frac{\text{tr}[H_{MN} U \rho^{MN} U^\dagger]}{MN} \\ &= \lim_{N \rightarrow \infty} \min_U \frac{\text{tr}[H_{MN} U (\rho^M)^{\otimes N} U^\dagger]}{MN} + kg_M + k \frac{f_N}{M} \\ &= \text{tr}[H \rho_{\beta'}] + kg_M, \end{aligned} \quad (36)$$

where  $\rho_{\beta'}$  is given by Eq. (35). Taking the limit  $M \rightarrow \infty$ , the thermal state energy  $\text{tr}[H\rho_{\beta'}]$  converges to  $\text{tr}[H\rho_{\beta}]$  and  $g_M$  converges to zero. Therefore, for any  $\epsilon > 0$  we can find  $M$  and  $N$  such that an element of sequence Eq. (26), defined by  $\rho^{MN}$ , is closer to  $\text{tr}[H\rho_{\beta}]$  than  $\epsilon$ . Because we assume that the sequence converges, it must converge to this value.

## II. DERIVATION OF THE LIMIT STATE IN STAGE 1.

In this section, we show that the limiting state after  $N$  measurements in Stage 1. is given by

$$\rho_N = \bigotimes_i \rho_i^{\otimes p_i N} \equiv \rho^N \quad (37)$$

up to a permutation in the order of tensor products, as  $N$  to infinity. This is Eq. (15) in the main text. The left hand side is a random variable (random state)  $\rho_N = \rho_{i_1} \otimes \dots \otimes \rho_{i_N}$  which is produced with probability  $p_{i_1 \dots i_N} = p_{i_1} \dots p_{i_N}$  after measuring each copy produced by the source (see the proof below for details). The right hand side is defined by Eq. (27). In mathematical terms, we show that  $\rho_N$  converges to  $\bigotimes_i \rho_i^{\otimes p_i N}$  almost surely. To show that, we first need to introduce relevant terminology and results.

For independent and identical distributed (i.i.d.) source  $X$  (meaning that all  $X_i = X$  are the same random variables), the Asymptotic Equipartition Property states that for any  $\epsilon > 0$ ,

$$\lim_{N \rightarrow \infty} \Pr \left[ \left| -\frac{1}{N} \ln p(X_1, \dots, X_N) - H(X) \right| > \epsilon \right] = 0. \quad (38)$$

where  $H(X) = -\sum_X p(X) \ln p(X)$  is the Shannon entropy of the outcomes, and  $p_i$  probability of event  $X_i$ . (See, e.g. [86].) In other words, for any fixed  $\epsilon$ , probability that we find sequence  $(X_1, \dots, X_N)$  to have probability  $p(X_1, \dots, X_N)$  outside these bounds:

$$e^{-N(H(X)+\epsilon)} \leq p(X_1, \dots, X_N) \leq e^{-N(H(X)-\epsilon)} \quad (39)$$

is zero as  $N$  goes to infinity, i.e., vast majority of probabilities will fall within these bounds for large  $N$ . Eq. (39) is the defining property of the typical set: typical set is defined as  $B_N^\epsilon = \{(X_1, \dots, X_N) | \text{Eq. (39) holds}\}$ .

For i.i.d. sources we actually have a stronger property: almost sure convergence,

$$\Pr \left[ \lim_{N \rightarrow \infty} -\frac{1}{N} \ln p(X_1, \dots, X_N) = H(X) \right] = 1, \quad (40)$$

which is implied by the strong law of large numbers. This states that any sequence of events  $(X_1, \dots, X_N)$  for which

$$p(X_1, \dots, X_N) \neq e^{-NH(X)} \quad (41)$$

have probability zero, as  $N$  goes to infinity.

We can now proceed with the proof.

*Proof.* We will show that in the limit of large  $N$ , the resulting state is almost surely of form (27) up to a different order of tensor products (i.e., up to a permutation).

For a single measurement, the resulting state is  $\rho_i$  with probability  $p_i \equiv p(\rho_i)$ , for two measurements, the resulting state is  $\rho_{i_1} \otimes \rho_{i_2}$  with probability  $p_{i_1 i_2} = p_{i_1} p_{i_2}$ , for  $N$  measurements, it is  $\rho_{i_1} \otimes \dots \otimes \rho_{i_N}$  with probability  $p_{i_1 \dots i_N} = p_{i_1} \dots p_{i_N}$ . The sequence  $\rho_N = \rho_{i_1} \otimes \dots \otimes \rho_{i_N}$  is therefore generated by an i.i.d. process in which the next element is given by  $\rho_i$  with probability  $p_i$ . We define a set

$$A_N = \left\{ \pi \left( \rho_1^{\otimes (N - \sum_{i \geq 2} \lfloor p_i N \rfloor)} \bigotimes_{i \geq 2} \rho_i^{\otimes \lfloor p_i N \rfloor} \right) \right\}_\pi, \quad (42)$$

where  $\pi$  defines all permutations on the tensor product. The probability of any state from this set is

$$p_{i_1 \dots i_N} = p_1^{N - \sum_{i \geq 2} \lfloor p_i N \rfloor} \prod_{i \geq 2} p_i^{\lfloor p_i N \rfloor}. \quad (43)$$

Taking the logarithm we have

$$\begin{aligned} -\frac{1}{N} \ln p_{i_1 \dots i_N} &= -\frac{1}{N} \left( \left( N - \sum_{i \geq 2} \lfloor p_i N \rfloor \right) \ln p_1 + \sum_{i \geq 2} \lfloor p_i N \rfloor \ln p_i \right) \\ &= -\sum_i p_i \ln p_i + \mathcal{O}\left(\frac{1}{N}\right) = H(\rho) + \mathcal{O}\left(\frac{1}{N}\right). \end{aligned} \quad (44)$$

Since  $\mathcal{O}\left(\frac{1}{N}\right)$  will be smaller than any  $\epsilon$ , Eq. (38) is satisfied for every element of set  $A_N$ . Further, due to the number of permutations  $\pi$  that generate a different sequence, the number of elements in the set is given by a multinomial,

$$|A_N| = \frac{N!}{(N - \sum_{i \geq 2} \lfloor p_i N \rfloor)! \lfloor p_2 N \rfloor! \dots \lfloor p_k N \rfloor!} \quad (45)$$

Taking the logarithm and using Stirling's approximation  $\ln N! = n \ln N + N + \frac{1}{2} \ln(2\pi N)$  we derive

$$\ln |A_N| = N \left( H(\rho) + \mathcal{O}\left(\frac{\ln(N)}{N}\right) \right). \quad (46)$$

Again,  $\mathcal{O}\left(\frac{\ln(N)}{N}\right)$  will be smaller than any  $\epsilon$  for large enough  $N$ . This gives,

$$e^{N(H(\rho)-\epsilon)} \leq |A_N| \leq e^{N(H(\rho)+\epsilon)}, \quad (47)$$

where  $H(\rho) = -\sum_i p_i \ln p_i$ . This means that set  $A_N$  contains elements with the same probability as the typical set

$$B_N^\epsilon = \left\{ \rho_{i_1} \otimes \dots \otimes \rho_{i_N} \mid e^{-N(H(\rho)+\epsilon)} \leq p_{i_1 \dots i_N} \leq e^{-N(H(\rho)-\epsilon)} \right\} \quad (48)$$

does, and its size (cardinality) is also the same as the typical set. This means that there will be a large overlap between the two and that they will share the same properties. Namely, for any arbitrarily small  $\delta > 0$ , the probability of finding the randomly generated sequence  $\rho_{i_1} \otimes \dots \otimes \rho_{i_N}$  in  $A_N$  is  $\Pr\{A_N\} \geq 1 - \delta$ , if  $N$  is taken to

be large enough [86]. Even the stronger condition of almost sure convergence, Eq. (40), holds, meaning that for a large enough  $N$ , sequences that do not fall into set  $A_N$  have probability zero. This completes the proof that for a large  $N$ , all states have the form (27) (or Eq. (37) if we gloss over the non-integer exponents), up to a permutation in the order of tensor products.  $\square$

### III. BOLTZMANN ERGOTROPY IN THE LARGE $N$ LIMIT

In this section, we combine Eqs. (26) and (37) (corresponding to Eqs. (5) and (15) in the main text) to derive expression for the Boltzmann ergotropy in the limit of large  $N$ ,

$$W_{\mathcal{C}}^{B\infty}(\rho) := \lim_{N \rightarrow \infty} \frac{W_{\mathcal{C}}^B(\rho^{\otimes N})}{N} = \text{tr}[H(\rho - \rho_\beta)], \quad (49)$$

where temperature  $\beta$  is implicitly defined by  $S_{\text{vN}}(\rho_\beta) = S_{\mathcal{C}}^B := \sum_i p_i \ln V_i$ .

To clarify, note that  $\rho^{\otimes N}$  above denotes the initial state produced by the source, i.e., the source produced  $N$  identical copies of the same unknown state  $\rho$ . Compare it to a random variable (random density matrix)  $\rho_N = \rho_{i_1} \otimes \dots \otimes \rho_{i_N}$  which is produced with probability  $p_{i_1 \dots i_N} = p_{i_1} \dots p_{i_N}$  after measuring each copy produced by the source, and the limiting state  $\rho^N = \otimes_i \rho_i^{p_i N}$ , defined by Eq. (27), to which  $\rho_N$  almost surely converges.

Recall the definition of the multipartite unitary extraction, Eq. (10) in the main text:

$$\mathcal{U}(\rho^{(1)} \otimes \dots \otimes \rho^{(N)}) = U \rho_{\text{cg}}^{(1)} \otimes \dots \otimes \rho_{\text{cg}}^{(N)} U^\dagger, \quad (50)$$

Using the same logic as in Eq. (12) in the main text, we apply  $\mathcal{U}$  to  $\rho_N$  to obtain

$$\begin{aligned} W_{\mathcal{C}}^B(\rho^{\otimes N}) &= \text{tr}[H_N \rho^{\otimes N}] - \min_U \text{tr}[H_N \mathcal{U}(\rho_N)] \\ &= \text{tr}[H_N \rho^{\otimes N}] - \min_U \text{tr}[H_N \mathcal{U}(\rho^N)] \\ &= N \text{tr}[H \rho] - \min_U \text{tr} \left[ H_N U \left( \otimes_i (P_i/V_i)^{\otimes p_i N} \right) U^\dagger \right]. \end{aligned} \quad (51)$$

where we used Eq. (37) in the second equality, assuming that  $N$  is large. Dividing this equation by  $N$  and using Eq. (26), we have

$$\lim_{N \rightarrow \infty} \frac{W_{\mathcal{C}}^B(\rho^{\otimes N})}{N} = \text{tr}[H \rho] - \text{tr}[H \rho_\beta], \quad (52)$$

where

$$S_{\text{vN}}(\rho_\beta) = \sum_i p_i S_{\text{vN}}(P_i/V_i) = \sum_i p_i \ln V_i =: S_{\mathcal{C}}^B. \quad (53)$$

This concludes the proof.




FIG. 3. Example of a three-level system with two-outcome coarse-graining.  $P_0$  and  $P_1$  represents two macrostates,  $|0\rangle$ ,  $|1\rangle$ ,  $|2\rangle$  energy eigenstates,  $\tilde{U}_0$  and  $\tilde{U}_1$  random unitaries acting on the first and the second subspace-macrostate, respectively, and  $U$  is the global extraction unitary.

### IV. UNDERSTANDING THE COARSE-GRAINED EXTRACTION OPERATION

Here we illustrate the coarse-grained extraction unitary,

$$U \oplus \tilde{U}_i, \quad (54)$$

where  $U_i$  are random unitary operators picked with the Haar measure acting on macrostates, and  $U$  is the global unitary, on an example depicted in Fig 3. This is Eq. (7) in the main text.

Consider a three level system described by Hamiltonian

$$H = E_0|0\rangle\langle 0| + E_1|1\rangle\langle 1| + E_2|2\rangle\langle 2|. \quad (55)$$

We will form a two-outcome coarse-graining by grouping together the two excited states, while the ground state will form its own macrostate. Mathematically, we have

$$\mathcal{C} = \{P_0, P_1\} \quad (56)$$

as our coarse-graining, where

$$P_0 = |0\rangle\langle 0|, \quad P_1 = |1\rangle\langle 1| + |2\rangle\langle 2| \quad (57)$$

are the corresponding projectors. The corresponding subspaces-macrostates are

$$\mathcal{H}_0 = \text{span}\{|0\rangle\}, \quad \mathcal{H}_1 = \text{span}\{|1\rangle, |2\rangle\}. \quad (58)$$

Clearly, we have  $\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_1$ .

The random unitaries are of the form

$$\tilde{U}_0 \oplus \tilde{U}_1 = \begin{pmatrix} \tilde{U}_0 & 0 & 0 \\ 0 & & \\ 0 & \tilde{U}_1 & \end{pmatrix}, \quad (59)$$

where  $\tilde{U}_0$  is a  $1 \times 1$  random unitary matrix and  $\tilde{U}_1$  is a  $2 \times 2$  random unitary matrix. (For an efficient sampling method of unitary operators from the Haar measure see, e.g., Ref. [87].) The global extraction unitary  $U$  is a general,  $3 \times 3$  unitary matrix.

### A. Three pure states that are indistinguishable by the measurement

We illustrate the difference the effects of the coarse-grained extraction unitary and a generic extraction unitary on several different states that could be produced by an unknown source.

Consider three states:

$$\rho_A = |1\rangle\langle 1|, \quad \rho_B = |2\rangle\langle 2|, \quad \rho_C = |+\rangle\langle +|, \quad (60)$$

where  $|+\rangle = (|1\rangle + |2\rangle)/\sqrt{2}$ . All of these have the same probabilities of outcomes when performing a measurement given by the coarse-graining (57),

$$p_0 = 0, \quad p_1 = 1. \quad (61)$$

Because the measurement outcome is  $i = 1$  with certainty, in all cases the post-measurement state is the same as the initial state,  $\rho_1^X = \rho_X$  ( $X = A, B, C$ ). It also means that the corresponding coarse-grained density matrix obtained by averaging over the random unitaries picked with the Haar measure,

$$\begin{aligned} \rho_{\text{cg}} &= \int (\tilde{U}_0 \oplus \tilde{U}_1) \rho_X (\tilde{U}_0^\dagger \oplus \tilde{U}_1^\dagger) d\mu(\tilde{U}_0 \oplus \tilde{U}_1) \\ &= \frac{p_0}{V_0} P_0 + \frac{p_1}{V_1} P_1 \\ &= 0 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{2} \end{pmatrix}, \end{aligned} \quad (62)$$

is the same of all of them (where  $X = A, B, C$ ,  $V_0 = 1$ ,  $V_1 = 2$ ). The optimal work extraction unitary from this state is given by

$$U = |0\rangle\langle 2| + |2\rangle\langle 0| + |1\rangle\langle 1| = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}. \quad (63)$$

This unitary turns the coarse-grained state into a passive state,

$$\pi_{\text{cg}} = U \rho_{\text{cg}} U^\dagger = \mathcal{U}(\rho_X) = \begin{pmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (64)$$

The initial energies of these three states are different (and unknown to the experimenter), and given by

$$\begin{aligned} \text{tr}[H\rho_A] &= E_1, \\ \text{tr}[H\rho_B] &= E_2, \\ \text{tr}[H\rho_C] &= (E_1 + E_2)/2. \end{aligned} \quad (65)$$

However, in all of these cases the energy of the respective state is reduced to the same value through the optimal work extraction unitary  $U$ , i.e.,

$$\text{tr}[H\pi_{\text{cg}}] = (E_0 + E_1)/2, \quad (66)$$

where  $X = A, B, C$ .

The corresponding Boltzmann ergotropy is given by

$$W_C^B(\rho_X) = \text{tr}[H(\rho_X - \pi_{\text{cg}})]. \quad (67)$$

This yields

$$\begin{aligned} W_C^B(\rho_A) &= (E_1 - E_0)/2, \\ W_C^B(\rho_B) &= E_2 - (E_0 + E_1)/2, \\ W_C^B(\rho_C) &= (E_2 - E_0)/2. \end{aligned} \quad (68)$$

Note that the observational ergotropy is the same as the Boltzmann ergotropy in this case. This is because  $p_0 = 0$  and  $p_1 = 1$ .

### B. Comparison with blind direct extraction

Let us compare the above strategy to the blind strategy of extraction, without the averaging using random unitaries picked with the Haar measure. The experimenter does not know the actual state of the system; their only knowledge is of the probabilities  $p_0 = 0$  and  $p_1 = 1$ . Let us say that they decide to extract energy directly by performing a unitary operation  $U$  which transfers the energy from the second excited state to the ground state, given by Eq. (63). In this case the final state energy is

$$\text{tr}[H\rho_X^{\text{final}}] = \text{tr}[HU\rho_X U^\dagger]. \quad (69)$$

This gives

$$\begin{aligned} \text{tr}[H\rho_A^{\text{final}}] &= E_1 \\ \text{tr}[H\rho_B^{\text{final}}] &= E_0, \\ \text{tr}[H\rho_C^{\text{final}}] &= (E_0 + E_1)/2. \end{aligned} \quad (70)$$

The total extracted work is then given by

$$W(\rho_X) = \text{tr}[H(\rho_X - \rho_X^{\text{final}})]. \quad (71)$$

This gives

$$\begin{aligned} W(\rho_A) &= 0 \\ W(\rho_B) &= E_2 - E_0, \\ W(\rho_C) &= (E_2 - E_0)/2. \end{aligned} \quad (72)$$

Clearly, this makes a very unreliable source, strongly depending on the unknown state of the system and on how lucky is the experimenter in guessing which extraction unitary  $U$  to apply. While in some cases, the extracted energy is quite high due to being lucky, in some other cases the strategy fails.

### C. What if the initial state was known?

Since all the states  $A, B, C$  are pure, if the experimenter knew what the states are, they would be able to

extract all the mean energy by transferring the state into the ground state. The extracted work would be given by the usual notion of ergotropy, which leads to

$$W_X = \text{tr}[H\rho_X] - E_0. \quad (73)$$

This is, however, not the case in our setup.

#### D. Another example

##### 1. Boltzmann ergotropy vs blind extraction

Finally, we show an example that can be very unfavorable to the blind direct extraction. Consider a state

$$\rho_D = \frac{1}{8}|0\rangle\langle 0| + \frac{7}{8}|1\rangle\langle 1| = \begin{pmatrix} \frac{1}{8} & 0 & 0 \\ 0 & \frac{7}{8} & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (74)$$

which has the mean initial energy

$$\text{tr}[H\rho_D] = \frac{1}{8}E_0 + \frac{7}{8}E_1. \quad (75)$$

We have  $p_0 = \frac{1}{8}$  and  $p_1 = \frac{7}{8}$ .

If the outcome of the measurement is  $i = 1$  we obtain the projected state  $\rho_1 = |1\rangle\langle 1|$  after the measurement. The corresponding coarse-grained state is given by

$$\rho_{\text{cg}}^1 = P_1/V_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & \frac{1}{2} \end{pmatrix}. \quad (76)$$

The optimal extraction unitary is again Eq. (63), which turns this coarse-grained state into a passive state

$$\pi_1 = \begin{pmatrix} \frac{1}{2} & 0 & 0 \\ 0 & \frac{1}{2} & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (77)$$

If the outcome is  $i = 0$  we obtain the projected state  $\rho_0 = |0\rangle\langle 0|$  after the measurement. The corresponding coarse-grained state is already passive and equal to the projected state,

$$\rho_{\text{cg}}^0 = \pi_0 = \rho_0. \quad (78)$$

As a result, no energy can be extracted in this case.

The total extractable work is given by Boltzmann ergotropy,

$$\begin{aligned} W_C^B(\rho_D) &= \text{tr}[H(\rho_D - p_0\pi_0 - p_1\pi_1)] \\ &= \frac{1}{8}E_0 + \frac{7}{8}E_1 - \frac{1}{8}E_0 - \frac{7}{16}(E_0 + E_1) \\ &= \frac{7}{16}(E_1 - E_0). \end{aligned} \quad (79)$$

Compare this to the blind extraction protocol, assuming that the experimenter chooses unitary operator Eq. (63) to extract the energy directly, in the case when they receive outcome  $i = 1$  from the measurement. In the

case they receive outcome  $i = 0$ , they decide to do nothing, because they already know from the outcome that the state is passive. Thus, the final states are

$$\rho_1^{\text{final}} = U|1\rangle\langle 1|U^\dagger = |1\rangle\langle 1|, \quad \rho_0^{\text{final}} = |0\rangle\langle 0|. \quad (80)$$

The average extracted work is

$$W(\rho_D) = \text{tr}[H\rho_D] - p_0 \text{tr}[H\rho_0^{\text{final}}] - p_1 \text{tr}[H\rho_1^{\text{final}}] = 0. \quad (81)$$

Thus, blind extraction does not extract anything in this case, while the coarse-grained extraction does.

##### 2. Observational ergotropy vs blind extraction

Now consider the experimenter moved to stage 2., in which the probabilities  $p_i$  are already identified, and no further measurements will be performed.

The coarse-grained state corresponding to the initial state  $\rho_D$  is given by

$$\rho_{\text{cg}} = \frac{1}{8}P_0 + \frac{7}{16}P_1 = \begin{pmatrix} \frac{1}{8} & 0 & 0 \\ 0 & \frac{7}{16} & 0 \\ 0 & 0 & \frac{7}{16} \end{pmatrix}. \quad (82)$$

The optimal extraction unitary, which is again of form (63), will turn this into a passive state

$$\pi_{\text{cg}} = \begin{pmatrix} \frac{7}{16} & 0 & 0 \\ 0 & \frac{7}{16} & 0 \\ 0 & 0 & \frac{1}{8} \end{pmatrix}. \quad (83)$$

The extracted work is given by observational ergotropy,

$$\begin{aligned} W_C(\rho_D) &= \text{tr}[H(\rho_D - \pi_{\text{cg}})] \\ &= \frac{1}{8}E_0 + \frac{7}{8}E_1 - \frac{7}{16}E_0 - \frac{7}{16}E_1 - \frac{1}{8}E_2 \\ &= (7E_1 - 5E_0 - 2E_2)/16. \end{aligned} \quad (84)$$

The extracted work is positive if  $7E_1 > 5E_0 + 2E_2$ . It is negative in the opposite case, signifying that the experimenter would lose energy. Of course, the computation crucially depends on the unknown initial energy, but as we mentioned in the main text, this can be estimated, either from Eq. (24) in the main text in the case of local energy coarse-grainings, or using methods in [59] in the case of general coarse-grainings.

We compare this to the blind extraction, in which the unitary (63) is applied directly on the state produced by the source. In this case, the final state is

$$\rho_D^{\text{final}} = U\rho_D U^\dagger = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{7}{8} & 0 \\ 0 & 0 & \frac{1}{8} \end{pmatrix}, \quad (85)$$

and the extracted work is given by

$$W(\rho_D) = \text{tr}[H(\rho_D - \rho_D^{\text{final}})] = (E_0 - E_2)/8. \quad (86)$$

This is always negative, so the experimenter is bound to lose energy.

## V. BOUND ON ENERGY FOR LOCAL ENERGY COARSE-GRAININGS

Here we prove the bound on energy that can be obtained when knowing the probabilities of outcomes of local energy measurements,

$$|\text{tr}[H\rho] - \text{tr}[H\rho_{\text{cg}}]| \leq 2\|H_{\text{int}}\| + k\Delta E, \quad (87)$$

which is Eq. (24) in the main text.  $\rho_{\text{cg}} = \sum_{E_1, E_2} \frac{p_{E_1 E_2}}{V_{E_1} V_{E_2}} P_{E_1} \otimes P_{E_2}$  is the coarse-grained state given

by the local energy coarse-grainings  $\mathcal{C} = \{P_{E_1} \otimes P_{E_2}\}$ .  $P_{E_i} = \sum_{E \in [E_i, E_i + \Delta E]} |E_i\rangle\langle E_i|$  are the coarse-grained projectors on local energies with resolution  $\Delta E$ , and  $H = H_1 + H_2 + H_{\text{int}}$ , where spectral decompositions of the local Hamiltonians are  $H_1 = \sum_{E_1} E_1 |E_1\rangle\langle E_1|$  and  $H_2 = \sum_{E_2} E_2 |E_2\rangle\langle E_2|$ .

We have

$$\begin{aligned} |\text{tr}[H\rho] - \text{tr}[H\rho_{\text{cg}}]| &= |\text{tr}[(H_1 + H_2)(\rho - \rho_{\text{cg}})] + \text{tr}[H_{\text{int}}(\rho - \rho_{\text{cg}})]| \\ &\leq \left| \text{tr} \left[ \left( \sum_{E_1, E_2} (E_1 + E_2) |E_1\rangle\langle E_1| \otimes |E_2\rangle\langle E_2| \right) (\rho - \rho_{\text{cg}}) \right] \right| + |\text{tr}[H_{\text{int}}(\rho - \rho_{\text{cg}})]| \\ &\leq \left| \sum_{E_1, E_2} (E_1 + E_2) \langle E_1, E_2 | (\rho - \rho_{\text{cg}}) | E_1, E_2 \rangle \right| + 2\|H_{\text{int}}\| \\ &\leq \left| \sum_{E'_1, E'_2} \sum_{E_1 \in [E'_1, E'_1 + \Delta E], E_2 \in [E'_2, E'_2 + \Delta E]} (E_1 + E_2) \langle E_1, E_2 | (\rho - \rho_{\text{cg}}) | E_1, E_2 \rangle \right| + 2\|H_{\text{int}}\| \quad (88) \\ &\leq \left| \sum_{E'_1, E'_2} (E'_1 + E'_2) \sum_{E_1 \in [E'_1, E'_1 + \Delta E], E_2 \in [E'_2, E'_2 + \Delta E]} \langle E_1, E_2 | (\rho - \rho_{\text{cg}}) | E_1, E_2 \rangle \right. \\ &\quad \left. + \sum_{E'_1, E'_2} \sum_{E_1 \in [E'_1, E'_1 + \Delta E], E_2 \in [E'_2, E'_2 + \Delta E]} (E_1 + E_2 - E'_1 - E'_2) \langle E_1, E_2 | \rho | E_1, E_2 \rangle \right| + 2\|H_{\text{int}}\| \\ &\leq 0 + \sum_{E_1, E_2} 2\Delta E p_{E_1 E_2} + 2\|H_{\text{int}}\| = 2\Delta E + 2\|H_{\text{int}}\|. \end{aligned}$$

In the above,  $E_1, E_2$  denote fine-grained energies, and  $E'_1, E'_2$  coarse-grained energies. Generalization for  $k$  par-

titions is  $|\text{tr}[H\rho] - \text{tr}[H\rho_{\text{cg}}]| \leq k\Delta E + 2\|H_{\text{int}}\|$ .  $\|H_{\text{int}}\|$  is also linearly proportional to  $k$ , therefore, both terms represent a finite size effect.