

THE CENTER AND INVARIANTS OF STANDARD FILIFORM LIE ALGEBRAS

VANDERLEI LOPES DE JESUS AND CSABA SCHNEIDER

ABSTRACT. We describe the centers of the universal enveloping algebras and the invariant rings of the standard filiform Lie algebras over fields of characteristic zero and also over large enough prime characteristic. We determine explicit generators for the quotient fields and also a compact form for the generators for the invariants rings. We prove several combinatorial results concerning the Hilbert series of these algebras.

1. INTRODUCTION

The problem of describing the center $Z(\mathfrak{g})$ of the universal enveloping algebra for nilpotent Lie algebras \mathfrak{g} has received quite a bit of attention. Dixmier [Dix58] determined $Z(\mathfrak{g})$ for nilpotent Lie algebras of dimension up to 5, and Ooms in [Oom09, Oom12] extended this work to algebras of dimension at most 7. Considering the descriptions of the centers $Z(\mathfrak{g})$ for these small-dimensional nilpotent Lie algebras, it becomes apparent that the most complicated structure is exhibited in the cases when \mathfrak{g} is standard filiform. So we decided that the study of the center $Z(\mathfrak{g})$ for standard filiform Lie algebras \mathfrak{g} is worth special attention. In this paper we consider both the cases of characteristic zero and prime characteristic.

As it turns out (and explained in Section 2), the center $Z(\mathfrak{g})$ of the standard filiform Lie algebra $\mathfrak{g} = \mathfrak{g}(n+2)$ (see Section 2 for the notation) is equal to the algebra of polynomial solutions $f \in \mathbb{F}[y_0, \dots, y_n]$ of the partial differential equation

$$\sum_{i \geq 0} y_i \frac{\partial f}{\partial y_{i+1}} = 0.$$

The operator $f \mapsto \sum y_i \partial f / \partial y_{i+1}$ is a locally nilpotent derivation, also referred to as a Weitzenböck derivation, on the polynomial algebra $\mathbb{F}[y_0, \dots, y_n]$ and was considered in [Bed08, Bed07, Bed09, Bed11b, Bed11a, BI15]. The same operator on the polynomial ring $\mathbb{F}[y_0, y_1, \dots]$ on infinitely many generators is referred to as the *down operator*, see [Fre13]. The crucial observation made by [Bed05, Bed10] is that the action of this operator on linear polynomials can be extended to an irreducible representation of the simple Lie algebra \mathfrak{sl}_2 and one can use the representation theory of \mathfrak{sl}_2 to obtain more information on the center $Z(\mathfrak{g})$. In fact, the elements of $Z(\mathfrak{g})$, considered as polynomials

Date: September 27, 2022.

in y_0, y_0, \dots, y_n , are known as *semi-invariants* of \mathfrak{sl}_2 acting on $\mathbb{F}[y_0, \dots, y_n]$ and Z_n is linked to the algebra of \mathfrak{sl}_2 -covariants of the binary form of degree n .

For $n \geq 1$, let Z_n denote $Z(\mathfrak{g}(n+2))$. If the characteristic of the field is zero, then Z_n can be viewed as a graded subalgebra of the polynomial algebra $\mathbb{F}[y_0, y_1, \dots, y_n]$. For $k \geq 0$, let $Z_{n,k}$ denote the degree- k homogeneous component of Z_n .

Theorem 1.1. *If \mathbb{F} has characteristic zero, then the algebra Z_n has Krull dimension n and the following are valid.*

- (1) *The fraction field $\text{Frac}(Z_n)$ can be written as*

$$\text{Frac}(Z_n) = \mathbb{F}(z_1, z_2, \dots, z_n) = \mathbb{F}(w_1, w_2, \dots, w_n)$$

where the elements z_1, z_2, \dots, z_n and w_1, w_2, \dots, w_n are defined in Section 5.

- (2) *$Z_n \subseteq \mathbb{F}[y_0^{-1}, z_1, z_2, \dots, z_n]$ and $Z_n \subseteq \mathbb{F}[y_0^{-1}, w_1, \dots, w_n]$.*

In Section 5 we define an operation \circ_d between two homogeneous elements of Z_n that are also h -eigenvectors for the generator h of the Cartan subalgebra of \mathfrak{sl}_2 . We show that Z_n is linearly spanned by elements of the form $y_0 \circ_{d_1} y_0 \circ_{d_2} \dots \circ_{d_k} y_0$ and one can choose minimal algebra generating sets out of such elements. Using computer calculation, we explicitly determine such generating sets for Z_n for $n \leq 6$ and for $n = 8$ in the Appendix A.

In Section 6, we consider the Hilbert series

$$H_n(t) = \sum_{k \geq 0} \dim Z_{n,k} t^k.$$

It is interesting that the dimensions $\dim Z_{n,k}$ occur also in the context of representation theory and combinatorics.

Theorem 1.2. *Assuming that \mathbb{F} has characteristic zero, that $n \geq 1$ and $k \geq 1$, the following are true for $\dim Z_{n,k}$:*

- (1) *$\dim Z_{n,k}$ coincides with the number of irreducible components of the \mathfrak{sl}_2 -module $S^k(U_n)$ where U_n is the $(n+1)$ -dimensional irreducible representation of \mathfrak{sl}_2 and S^k is the k -th symmetric power.*
- (2) *$\dim Z_{n,k}$ is equal to the number of partitions of $\lfloor kn/2 \rfloor$ into k blocks each of size n .*
- (3) *$\dim Z_{n,k} = \dim Z_{k,n}$.*

Theorem 1.2 follows from Theorem 6.1 (part (1)) and from Theorem 6.4 (parts (2) and (3)).

By the Mauer–Weitzenböck Theorem [Fre17, Theorem 6.1], Z_n is finitely generated and so $H_n(t)$ is a rational function. The combinatorial description of $\dim Z_{n,k}$ in Theorem 1.2 and the computations in [EZ22] make it possible to determine the Hilbert series $H_n(t)$ as rational expressions for $n \leq 16$. As in the situation considered by Almkvist [Alm80a, Alm80b], one also has the following closed formula for $H_n(t)$.

Theorem 1.3. *If \mathbb{F} has characteristic zero, then, for $n \geq 1$,*

$$(1) \quad H_n(t) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{1 + \exp(i\varphi)}{\prod_{k=0}^n (1 - t \exp(i(n-2k)\varphi))} d\varphi.$$

The integral formula in Theorem 1.3 appears in [Alm80a], since in that situation, the counting argument also boils down to counting the partitions in Theorem 1.2(2). We include a proof in Section 6 for easier reference.

In the final Section 7, we consider the case when the characteristic of the field is a prime. In this case, a well-known theorem of Zassenhaus states that $Z(\mathfrak{g})$ is a normal domain and our main result states that the generators z_2, \dots, z_n in Theorem 1.1 generate Z_n over the p -center $Z_p(\mathfrak{g})$ up to normal closure.

Theorem 1.4. *Suppose that \mathbb{F} has characteristic p and also that $p \geq n + 1$. Set $\mathfrak{g} = \mathfrak{g}(n + 2)$. Then $Z(\mathfrak{g})$ is the integral closure of $Z_p(\mathfrak{g})[z_2, \dots, z_n]$ in its field of fractions and it also coincides with the integral closure of $Z_p(\mathfrak{g})[z_2, \dots, z_n]$ in the localization $Z_p(\mathfrak{g})[y_0^{-1}, z_2, \dots, z_n]$. In particular, $Z(\mathfrak{g}) \subseteq Z_p(\mathfrak{g})[y_0^{-1}, z_2, \dots, z_n]$.*

The first author was financially supported by a PhD scholarship awarded by CNPq (Brazil). The second author acknowledges the financial support of the CNPq projects *Produtividade em Pesquisa* (project no.: 308212/2019-3) and *Universal* (project no.: 421624/2018-3 and 402934/2021-0). We thank Lucas Calixto for several useful comments on the earlier drafts, to Dmitry Shcheglov and to Csaba Noszaly for help with Theorem 1.3.

2. THE STANDARD FILIFORM LIE ALGEBRA

The *standard filiform Lie algebra* $\mathfrak{g}(n + 2)$ is the Lie algebra over a field \mathbb{F} with basis $\{x, y_0, y_1, \dots, y_n\}$ and nonzero brackets $[x, y_i] = y_{i-1}$ for $i \in \{1, \dots, n\}$ (in particular, $[x, y_0] = 0$ and y_0 is central). This is a nilpotent Lie algebra of dimension $n + 2$ and nilpotency class $n + 1$; such a Lie algebra is also referred to as a *Lie algebra of maximal class*. The center of $\mathfrak{g}(n + 2)$ is one-dimensional and is generated by y_0 .

The adjoint action of the Lie algebra $\mathfrak{g} = \mathfrak{g}(n + 2)$ on itself can be extended to the polynomial ring $\mathbb{F}[\mathfrak{g}] = \mathbb{F}[x, y_0, y_1, \dots, y_n]$ and we define the invariant ring $\mathbb{F}[\mathfrak{g}]^{\mathfrak{g}}$ as

$$\mathbb{F}[\mathfrak{g}]^{\mathfrak{g}} = \{f \in \mathbb{F}[\mathfrak{g}] \mid u(f) = 0 \text{ for all } u \in \mathfrak{g}\}.$$

The commutative algebra $\mathbb{F}[\mathfrak{g}]^{\mathfrak{g}}$ is referred to as the *algebra of polynomial invariants* of \mathfrak{g} .

Let $\mathbb{F}[\mathbf{y}]$ denote the polynomial algebra $\mathbb{F}[y_0, y_1, \dots]$ in infinitely many variables and let $\mathbb{F}[\mathbf{y}_n] = \mathbb{F}[y_0, \dots, y_n]$. Then the one-dimensional Lie algebra $\mathfrak{n} = \langle x \rangle$ acts on the vector space $V = \langle y_0, y_1, \dots \rangle$ by mapping y_i to y_{i-1} for $i \geq 1$ and y_0 to 0. Set $V_n = \langle y_0, \dots, y_n \rangle$ and note that V_n is an \mathfrak{n} -submodule of V . The Lie algebra $\mathfrak{g}(n + 2)$ can also be viewed as the semidirect product $V_n \rtimes \mathfrak{n}$.

The \mathfrak{n} -action on V can be extended to $\mathbb{F}[\mathbf{y}]$ by the Leibniz rule and $\mathbb{F}[\mathbf{y}_n]$ is \mathfrak{n} -invariant for all $n \geq 1$. We denote by $x(f)$ the image of $f \in \mathbb{F}[\mathbf{y}]$ under this action.

Simple computation shows that

$$(2) \quad x(f) = \sum_{i \geq 0} y_i \frac{\partial f}{\partial y_{i+1}}.$$

Thus, the algebra of x -invariants in $\mathbb{F}[\mathbf{y}]$ is denoted by $\mathbb{F}[\mathbf{y}]^x$; more precisely,

$$\mathbb{F}[\mathbf{y}]^x = \{f \in \mathbb{F}[\mathbf{y}] \mid x(f) = 0\}.$$

The operator $f \mapsto x(f)$ on $\mathbb{F}[\mathbf{y}]$ is often referred to as the *down operator*; see [Fre13]. The algebra of x -invariants can also be described as the set of solutions of the partial differential equation

$$(3) \quad \sum_{i \geq 0} y_i \frac{\partial f}{\partial y_{i+1}} = 0.$$

For a Lie algebra \mathfrak{g} , let $Z(\mathfrak{g})$ denote the center of the universal enveloping algebra $U(\mathfrak{g})$. Since y_0, \dots, y_n commute in $\mathfrak{g}(n+2)$, the Poincaré–Birkhoff–Witt Theorem gives an embedding of the polynomial algebra $\mathbb{F}[\mathbf{y}_n]$ into $U(\mathfrak{g}(n+2))$. Further, if \mathbb{F} has prime characteristic p and $p \geq n+1$, then $x^p \in Z(\mathfrak{g}(n+2))$ and $\mathbb{F}[x^p, y_0, \dots, y_n]$ is a subalgebra of $U(\mathfrak{g}(n+2))$. Note that the action of x on $\mathbb{F}[\mathbf{y}_n]$ is the same whether it is considered inside $U(\mathfrak{g}(n+2))$ or it is considered as a stand-alone polynomial algebra under the action defined in (2).

Lemma 2.1. *Let $\mathfrak{g} = \mathfrak{g}(n+2)$ be a standard filiform Lie algebra. If \mathbb{F} has characteristic zero, then*

$$Z(\mathfrak{g}) \subseteq \mathbb{F}[\mathbf{y}_n]$$

and

$$Z(\mathfrak{g}) = \mathbb{F}[\mathfrak{g}]^{\mathfrak{g}} = \mathbb{F}[\mathbf{y}_n]^x.$$

If the characteristic of \mathbb{F} is a prime p , then

$$Z(\mathfrak{g}) \subseteq \mathbb{F}[x^p, y_0, \dots, y_n]$$

and, if in addition $p \geq n+1$, then

$$Z(\mathfrak{g}) = \mathbb{F}[\mathfrak{g}]^{\mathfrak{g}} = \mathbb{F}[x^p, y_0, \dots, y_n]^x.$$

Proof. Easy induction shows, in $U(\mathfrak{g})$, that

$$[x^k, y_n] = x^k y_n - y_n x^k = k x^{k-1} y_{n-1} + h_k$$

where h_k is a linear combination of monomials whose degree in x is smaller than $k-1$. Suppose that $z \in Z(\mathfrak{g})$ and let $m = x^\alpha y_0^{\alpha_0} \cdots y_n^{\alpha_n}$ be the leading term of z in the lexicographic monomial order. Then the last displayed equation implies that the leading term of $[z, y_n]$ is

$$\alpha x^{\alpha-1} y_0^{\alpha_0} \cdots y_{n-1}^{\alpha_{n-1}+1} y_n^{\alpha_n}.$$

But, as z is central, $[z, y_n] = 0$, and so either $\alpha = 0$ or the characteristic of \mathbb{F} is p and $p \mid \alpha$. Thus the containments for $Z(\mathfrak{g})$ in $\mathbb{F}[\mathbf{y}_n]$ and, in the case of characteristic p , in $\mathbb{F}[x^p, y_0, \dots, y_n]$ is verified. One can similarly verify the same containment $\mathbb{F}[\mathfrak{g}]^{\mathfrak{g}} \subseteq \mathbb{F}[\mathbf{y}_n]$

and $\mathbb{F}[\mathfrak{g}]^{\mathfrak{g}} \subseteq \mathbb{F}[x^p, y_0, \dots, y_n]$. For the equations regarding $Z(\mathfrak{g})$, note that if \mathbb{F} has prime characteristic p and $p \geq n + 1$, then $x^p \in Z(\mathfrak{g})$. The rest of the lemma follows from the observation above that $\mathbb{F}[\mathfrak{y}_n]$ and $\mathbb{F}[x^p, y_0, \dots, y_n]$ are embedded into $U(\mathfrak{g})$ under the Poincaré–Birkhoff–Witt Theorem and the x -action on these subalgebras is the same as in (2). \square

From now on, we set $Z_n = Z(\mathfrak{g}_{n+2})$ considered as a subalgebra of $\mathbb{F}[\mathfrak{y}_n]$. Note that the x -action on $\mathbb{F}[\mathfrak{y}_n]$ preserves the grading, and hence Z_n is a graded subalgebra.

Example 2.2. Suppose that the characteristic of \mathbb{F} is zero. The ring Z_1 is the center of $U(\mathfrak{g}(3))$ which coincides with the invariant algebra of $\mathfrak{g}(3)$. Note that $\mathfrak{g}(3)$ is the three-dimensional Heisenberg Lie algebra. By Lemma 2.1,

$$Z_1 = \left\{ f \in \mathbb{F}[y_0, y_1] \mid y_0 \frac{\partial f}{\partial y_1} = 0 \right\} = \left\{ f \in \mathbb{F}[y_0, y_1] \mid \frac{\partial f}{\partial y_1} = 0 \right\} = \mathbb{F}[y_0].$$

3. THE ELEMENTS OF $\mathbb{F}[\mathfrak{y}_n]^x$

In this section, we prove some general facts concerning the elements of $\mathbb{F}[\mathfrak{y}_n]^x$ and of the fraction field $\mathbb{F}(\mathfrak{y}_n)^x$. We will use these facts to determine explicit generators in Section 3. In this section \mathbb{F} is a field of characteristic zero.

Lemma 3.1. *Suppose that $z \in \mathbb{F}[\mathfrak{y}_n]$ and write*

$$z = \sum_{i=0}^k y_n^i g_i$$

where $g_i \in \mathbb{F}[\mathfrak{y}_{n-1}]$. Then $z \in \mathbb{F}[\mathfrak{y}_n]^x$ if and only if

$$x(g_k) = 0 \quad \text{and} \quad x(g_i) = -(i+1)y_{n-1}g_{i+1} \quad \text{for all } i \leq k-1.$$

Proof. Compute that

$$\begin{aligned} x(z) &= x \left(\sum_{i=0}^k y_n^i g_i \right) = \sum_{i=1}^k i y_n^{i-1} y_{n-1} g_i + \sum_{i=0}^k y_n^i \cdot x(g_i) \\ &= y_n^k \cdot x(g_k) + \sum_{i=0}^{k-1} y_n^i ((i+1)y_{n-1}g_{i+1} + x(g_i)). \end{aligned}$$

Since $g_i, x(g_i) \in \mathbb{F}[\mathfrak{y}_{n-1}]$ for all $i \in \{1, \dots, k\}$, we obtain that the coefficient of y_n^k in $x(z)$ is $x(g_k)$ and, for $i \in \{0, \dots, k-1\}$, the coefficient of y_n^i is $(i+1)y_{n-1}g_{i+1} + x(g_i)$. Thus $x(z) = 0$ if and only if $x(g_k) = 0$ and $(i+1)y_{n-1}g_{i+1} + x(g_i) = 0$ for $i \in \{0, \dots, k-1\}$. \square

Corollary 3.2. *If $z \in \mathbb{F}[\mathfrak{y}_n]^x$ is written as in the previous lemma, then z is determined by the term $g_0 \in \mathbb{F}[\mathfrak{y}_{n-1}]$.*

In the following lemma, we consider monomials ordered lexicographically considering the exponents of y_n, \dots, y_1, y_0 in this order.

Lemma 3.3. *Suppose that $z \in \mathbb{F}[\mathbf{y}_n]^x$ and write*

$$z = \sum_{i=0}^k y_n^i g_i$$

where $g_i \in \mathbb{F}[\mathbf{y}_{n-1}]$ for all $i \in \{0, \dots, k\}$. Then

- (1) $g_k \in \mathbb{F}[\mathbf{y}_{n-1}]^x$;
- (2) the leading monomial of z does not contain y_1 .

Proof. Statement (1) follows from Lemma 3.1. Let us prove (2) by induction on n . If $n = 1$, then the statement is clear, since $Z_1 = \mathbb{F}[\mathbf{y}_1]^x = \mathbb{F}[y_0, y_1]^x = \mathbb{F}[y_0]$ (see Example 2.2). Suppose that the statement holds for some $n - 1 \geq 1$ and consider $z \in \mathbb{F}[\mathbf{y}_n]$. If z does not contain the variable y_n , then $z \in \mathbb{F}[\mathbf{y}_{n-1}]^x$ and we are done by the induction hypothesis. Otherwise, write $z = \sum_{i \leq k} y_n^i g_i$. Then the leading monomial of z is $y_n^k m$ where m is the leading monomial of g_k . By Statement (1), $g_k \in \mathbb{F}[\mathbf{y}_{n-1}]^x$, and so the induction hypothesis implies that m does not contain y_1 . Thus $y_n^k m$ does not contain y_1 also. \square

Theorem 3.4. *Suppose that $Z \subseteq \mathbb{F}[\mathbf{y}_n]$ is a graded subalgebra (with respect to the grading by degree) such that if $z \in Z$ then the leading monomial of z is not divisible by y_1 . Suppose that $z_1, \dots, z_n \in Z$ are homogeneous elements such that $z_1 = y_0$ and, for each $i \in \{2, \dots, n\}$,*

$$z_i = y_i y_0^{k_i} + z'_i \text{ with } k_i \geq 0 \text{ and } z'_i \in \mathbb{F}[\mathbf{y}_{i-1}].$$

Then the following are valid.

- (1) $Z \subseteq \mathbb{F}[y_0^{-1}, z_1, \dots, z_n]$.
- (2) The fraction field of Z is generated by z_1, \dots, z_n .
- (3) The elements z_1, \dots, z_n are algebraically independent over \mathbb{F} .
- (4) $\dim Z = n$ (Krull dimension).

Proof. (1) Suppose that $z \in Z$; we need to show that z can be written as a polynomial expression in y_0^{-1} and in the given z_i . We show this by induction on the leading monomial of z . The base case of the induction is when $z = 1$ and the statement in this case is obvious. Suppose now that z is a homogeneous element of Z and its leading monomial is

$$m = y_n^{\alpha_n} \cdots y_2^{\alpha_2} y_0^{\alpha_0}$$

with $m \neq 1$. Note, for $i \geq 2$, that the leading monomial of z_i is $y_i y_0^{k_i}$. Consider the element

$$z' = z - z_2^{\alpha_2} \cdots z_n^{\alpha_n} y_0^{\alpha_0 - \alpha_2 k_2 - \cdots - \alpha_n k_n}$$

Then the leading monomial of z' is smaller than m in the monomial ordering and $z' \in Z$. By the induction hypothesis, $z' \in \mathbb{F}[y_0^{-1}, z_1, \dots, z_n]$ which implies that $z \in \mathbb{F}[y_0^{-1}, z_1, \dots, z_n]$.

(2) follows from (1). To prove (3), take a polynomial $g \in \mathbb{F}[t_1, \dots, t_n]$ such that $g(z_1, \dots, z_n) = 0$. If $g \neq 0$, then, supposing that the leading term of g is $t_1^{\alpha_1} \cdots t_n^{\alpha_n}$,

we find that the leading term of $g(z_1, \dots, z_n)$ is $y_n^{\alpha_n} \dots y_2^{\alpha_2} y_0^k$ for some $k \geq 0$, which is nonzero. Thus $g = 0$ must hold and z_1, \dots, z_n are algebraically independent.

(4) follows from (3). \square

4. SOME FACTS CONCERNING THE REPRESENTATIONS OF \mathfrak{sl}_2

Suppose in this section that \mathbb{F} is a field of characteristic zero. It is well known that the Lie algebra $\mathfrak{sl}_2 = \langle e, f, h \rangle$, where $h = [e, f]$, acts on the polynomial ring $\mathbb{F}[x_1, x_2]$ by extending the action

$$\begin{aligned} e : x_1 &\mapsto 0, & x_2 &\mapsto x_1, \\ f : x_1 &\mapsto x_2, & x_2 &\mapsto 0, \\ h : x_1 &\mapsto x_1, & x_2 &\mapsto -x_2 \end{aligned}$$

using the Leibniz rule. This action on $\mathbb{F}[x_1, x_2]$ can be described, for $g \in \mathbb{F}[x_1, x_2]$, as

$$e(g) = x_1 \frac{\partial g}{\partial x_2} \quad \text{and} \quad f(g) = x_2 \frac{\partial g}{\partial x_1}.$$

Let U_n denote the space of homogeneous polynomials in $\mathbb{F}[x_1, x_2]$ of degree n with basis $x_1^n, x_1^{n-1}x_2, \dots, x_1x_2^{n-1}, x_2^n$. Then the action of e and f can be described as

$$e(x_1^i x_2^{n-i}) = (n-i)x_1^{i+1}x_2^{n-i-1} \quad \text{and} \quad f(x_1^i x_2^{n-i}) = ix_1^{i-1}x_2^{n-i+1}.$$

The basis $x_1^n, x_1^{n-1}x_2, \dots, x_2^n$ consists of eigenvectors for the operator h with eigenvalues $n, n-2, \dots, -n+2, -n$, respectively. The eigenvectors of h in a representation of \mathfrak{sl}_2 are often referred to as *weight vectors* and the corresponding eigenvalues as *weights*. The following result is well-known.

Theorem 4.1. *For $n \geq 0$, the space U_n is irreducible considered as an \mathfrak{sl}_2 -module. Furthermore, each finite-dimensional irreducible \mathfrak{sl}_2 -module is isomorphic to U_n for some $n \geq 0$.*

Note that the \mathfrak{sl}_2 -module U_n can also be recognised by the weights (that is, the h -eigenvalues) which are $n, n-2, \dots, -n+2, -n$. The reason why we are interested in the \mathfrak{sl}_2 -modules U_n is because the action of e on U_n is isomorphic to the action of x on $V_n = \langle y_0, \dots, y_n \rangle$ defined in Section 2 in the context of standard filiform Lie algebras.

Lemma 4.2. *Suppose that $n \geq 0$ and, for $i \in \{0, \dots, n\}$, let $w_i = (1/i!)x^{n-i}y^i$. Then $e(w_0) = 0$ and $e(w_i) = w_{i-1}$ for $i \in \{1, \dots, n\}$.*

Proof. This is easy: just calculate for $i \geq 0$ that

$$e(w_i) = \frac{1}{i!}e(x^{n-i}y^i) = i \cdot \frac{1}{i!}x^{n-i+1}y^{i-1} = \frac{1}{(i-1)!}x^{n-i+1}y^{i-1} = w_{i-1}.$$

\square

For a Lie algebra \mathfrak{g} and for a \mathfrak{g} -module U , let $U^{\mathfrak{g}}$ denote the space

$$U^{\mathfrak{g}} = \{u \in U \mid a(u) = 0 \text{ for all } a \in \mathfrak{g}\}$$

of \mathfrak{g} -invariants in U . If $\mathfrak{g} = \langle x \rangle$, then we write U^x for $U^{\mathfrak{g}}$ and U^x is the zero eigenspace of x acting on U .

Corollary 4.3. *The following are valid for a finite-dimensional \mathfrak{sl}_2 -module U written as a direct sum $U = W_1 \oplus \cdots \oplus W_d$ of simple \mathfrak{sl}_2 -submodules.*

- (1) *For $i \in \{1, \dots, d\}$, let $z_i \in W_i^e \setminus \{0\}$ (which is unique up scalar multiple). Then $\{z_1, \dots, z_d\}$ is a basis of U^e .*
- (2) *$\dim U^e = \dim U^f = d$.*
- (3) *$\dim U^e = \dim Y_0 + \dim Y_1$ where Y_0 and Y_1 are the eigenspaces in U of the operator h corresponding to the eigenvalues 0 and 1, respectively.*

Proof. By Weyl's Theorem (see [Jac79, Theorem 8, Section III.7]),

$$(4) \quad U = W_1 \oplus \cdots \oplus W_d$$

where the W_i are irreducible \mathfrak{sl}_2 -modules. Then each W_i is isomorphic to U_{k_i} where $k_i \geq 0$ (Theorem 4.1). By Lemma 4.2, the matrix of e on W_i is conjugate to a single Jordan block that corresponds to the eigenvalue zero. Thus the matrix of e on U is conjugate to a block-diagonal matrix with d Jordan blocks each of which corresponds to the eigenvalue zero. Noting that U^e is the zero eigenspace of e considered as an operator on U , we obtain assertions (1) and (2). For the assertion that $d = \dim Y_0 + \dim Y_1$, choose for each W_i a basis consisting of h -eigenvectors and note that each irreducible component W_i contributes with dimension one to either Y_0 (if k_i is even) or to Y_1 (if k_i is odd). \square

Lemma 4.4. *Suppose that U is a finite-dimensional \mathfrak{sl}_2 -module. Suppose that $z \in U^e$ is an h -eigenvector of eigenvalue w . Then z is contained in $\bigoplus_i (W_i)^e$ where the W_i are the irreducible components of U that are isomorphic to U_w . Furthermore, the \mathfrak{sl}_2 -submodule generated by z is irreducible and is isomorphic to U_w .*

Proof. The fact that $z \in \bigoplus_i (W_i)^e$ follows from the weight space decomposition (4) for U . One can show by induction that $f^i(z)$ is an h -eigenvector with eigenvalue $w - 2i$ for all $i \in \{0, \dots, n\}$ and these vectors form the basis of an \mathfrak{sl}_2 -submodule isomorphic to U_w . \square

Theorem 4.5. *Suppose that U_m and U_n are \mathfrak{sl}_2 -modules as defined above. Then the following decomposition of \mathfrak{sl}_2 -modules is valid:*

$$U_m \otimes U_n \cong \bigoplus_{i=0}^{\min\{m,n\}} U_{m+n-2i}.$$

Suppose that u_0, \dots, u_m and w_0, \dots, w_n are bases for U_m and U_n , respectively, such that $e(u_0) = e(w_0) = 0$ and $e(u_i) = u_{i-1}$ and $e(w_j) = w_{j-1}$ for $i \in \{1, \dots, m\}$ and

$j \in \{1, \dots, n\}$. Define, for $i = 0, 2, \dots, \min\{m, n\}$,

$$(5) \quad z_i = \sum_{d=0}^i (-1)^d u_d \otimes w_{i-d}.$$

Then $e(z_i) = 0$ and the irreducible component of $U_m \otimes U_n$ isomorphic to U_{m+n-2i} is generated by z_i .

Proof. The decomposition of $U_m \otimes U_n$ is well known as the Clebsch-Gordan formula for \mathfrak{sl}_2 [Kow14, Theorem 2.6.3]. The fact that $e(z_i) = 0$ is true since

$$\begin{aligned} e(z_i) &= e\left(\sum_{d=0}^i (-1)^d u_d \otimes w_{i-d}\right) = \sum_{d=0}^i (-1)^d e(u_d) \otimes w_{i-d} + \sum_{d=0}^i (-1)^d u_d \otimes e(w_{i-d}) \\ &= \sum_{d=1}^i (-1)^d u_{d-1} \otimes w_{i-d} + \sum_{d=0}^{i-1} (-1)^d u_d \otimes w_{i-d-1} = 0. \end{aligned}$$

Further, it also follows that z_i is an h -eigenvector of eigenvalue $n+m-2i$. By Lemma 4.4, z_i is contained in the direct sum of the \mathfrak{sl}_2 -submodules that are isomorphic to U_{n+m-2i} , but the Clebsch-Gordan decomposition implies that there is a unique such submodule and this must be generated by z_i . \square

The element z_i defined in (5) can be viewed as a Casimir element for the operator e on the tensor product $U_m \otimes U_n$.

Lemma 4.6. *Let $z \in U_n$ be an h -eigenvector of weight $n-2i$ with some $i \in \{0, \dots, n\}$. Then*

$$e(f(z)) = (n-i)(i+1)z.$$

Proof. We can assume without loss of generality that $U_n = \langle x_1^n, x_1^{n-1}x_2, \dots, x_1x_2^{n-1}, x_2^n \rangle$ and $z = x_1^{n-i}x_2^i$. Then

$$e(f(z)) = e(f(x_1^{n-i}x_2^i)) = (n-i)(i+1)x_1^{n-i}x_2^i.$$

\square

Corollary 4.7. *Let U_n be an $(n+1)$ -dimensional irreducible \mathfrak{sl}_2 -module and consider the polynomial algebra $\mathbb{F}[U_n]$ as an \mathfrak{sl}_2 -module. For $i \geq 0$, let $U_{n,i}$ denote the space of homogeneous polynomials of $\mathbb{F}[U_n]$ of degree i . Let $z_1 \in U_{n,k_1}$ and $z_2 \in U_{n,k_2}^e$ that are also h -eigenvectors of weight w_1 and w_2 , respectively. Suppose that $d \leq \min\{w_1, w_2\}$ and define for $i \in \{1, 2\}$ and for $k \in \{0, \dots, d\}$,*

$$z_{i,0} = z_i \quad \text{and} \quad z_{i,k} = \frac{1}{k(w_i - k + 1)} f(z_{i,k-1}) \quad \text{for } k \geq 1.$$

Set

$$z_1 \circ_d z_2 = \sum_{i=0}^d (-1)^i z_{1,i} z_{2,d-i}.$$

Then $z_1 \circ_d z_2 \in U_{n, k_1 + k_2}^e$ and it is an h -eigenvector of weight $w_1 + w_2 - 2d$. Furthermore, letting $y \in U_{n, 1}^e \setminus \{0\}$ be fixed and z_1, \dots, z_m be a basis of $U_{n, k-1}^e$ formed by h -eigenvectors, the elements $z_i \circ_d y$, with $i \in \{1, \dots, m\}$ and $d \geq 0$ generate $U_{n, k}^e$ and are h -eigenvectors.

Proof. Suppose that W_1 and W_2 are the irreducible components of U_{n, k_1} and U_{n, k_2} that contain z_1 and z_2 , respectively (they exist by Lemma 4.4). Then $W_1 \cong U_{w_1}$ and $W_2 \cong U_{w_2}$. The fact that $e(z_{i,0}) = 0$, for $i = 1, 2$, is clear. Suppose that $k \geq 1$ note that $z_{i, k-1}$ is an h -eigenvector of weight $w_i - 2(k-1)$. Thus Lemma 4.6 implies that

$$e(z_{i,k}) = \frac{1}{k(w_i - k + 1)} e(f(z_{i, k-1})) = \frac{k(w_i - k + 1)}{k(w_i - k + 1)} z_{i, k-1} = z_{i, k-1}.$$

The product space $W_1 W_2$ is an epimorphic image of the tensor product $W_1 \otimes W_2$ under the \mathfrak{sl}_2 -homomorphism $\psi : W_1 \otimes W_2 \rightarrow W_1 W_2$ induced by multiplication. Observe that $z_1 \circ_d z_2$ is the image of

$$\sum_{i=0}^d (-1)^i z_{1, i} \otimes z_{2, d-i}$$

under ψ , and so Theorem 4.5 implies that $e(z_1 \circ_d z_2) = 0$ and also that $(W_1 W_2)^e$ is generated by such elements. The final assertion follows from the fact that $U_{n, k} = U_{n, k-1} U_{n, 1}$ and that $U_{n, 1}$ is irreducible, and in particular $U_{n, 1}^e = \langle y \rangle$. \square

5. EXPLICIT GENERATORS OF Z_n

In this section we return to our investigation of the invariant algebra $Z_n = \mathbb{F}[\mathbf{y}_n]^x$ as seen in Sections 2–3. Assume throughout this section that \mathbb{F} is a field of characteristic zero. Note that x acts on the vector space $V_n = \langle y_0, y_1, \dots, y_n \rangle$ and Lemma 4.2 shows that its action is equivalent to the action of $e \in \mathfrak{sl}_2$. Thus there exists a $\hat{f}_n \in \text{End}(V_n)$ such that the map $x \mapsto e$, $\hat{f}_n \mapsto f$ can be extended to an isomorphism of Lie algebras $\langle x, \hat{f}_n \rangle \rightarrow \mathfrak{sl}_2$. In fact, \hat{f}_n is determined in [Bed10, Theorem 3.1] and in [Bed05, Theorem 3.1] as

$$(6) \quad \hat{f}_n(y_i) = (i+1)(n-i)y_{i+1} \quad \text{for all } i \in \{0, \dots, n\}.$$

Because of the isomorphism with \mathfrak{sl}_2 , $\hat{h} = [x, \hat{f}_n]$ is an operator on V_n with eigenvectors y_0, y_1, \dots, y_n corresponding to eigenvalues $n, n-2, \dots, -n+2, -n$. For $k \geq 0$, let $Z_{n, k}$ denote the degree- k homogeneous component of Z_n . The algebras $\mathbb{F}[\mathbf{y}_n]$ and $\mathbb{F}[U_n]$ in Corollary 4.7 are isomorphic as \mathfrak{sl}_2 -modules. Furthermore, under the obvious isomorphism between $\mathbb{F}[\mathbf{y}_n]$ and $\mathbb{F}[U_n]$, Z_n corresponds to $\mathbb{F}[U_n]^e$, while $Z_{n, k}$ corresponds to $U_{n, k}^e$. Thus the following result is an immediate consequence of Corollary 4.7.

Proposition 5.1. *Suppose that $z_1 \in Z_{n, k_1}$ and $z_2 \in Z_{n, k_2}$ are \hat{h} -eigenvectors with eigenvalues w_1 and w_2 respectively. Suppose that $d \leq \min\{w_1, w_2\}$ and define for $i \in \{1, 2\}$ and for $k \in \{0, \dots, d\}$,*

$$z_{i,0} = z_i \quad \text{and} \quad z_{i,k} = \frac{1}{k(w_i - k + 1)} \hat{f}_n(z_{i, k-1}) \quad \text{for } k \geq 1.$$

Define

$$(7) \quad z_1 \circ_d z_2 = \sum_{i=0}^d (-1)^i z_{1,i} z_{2,d-i}.$$

Then $z_1 \circ_d z_2 \in Z_{n,k_1+k_2}$ and this element is an \hat{h} -eigenvector of weight $w_1 + w_2 - 2d$. Furthermore, letting z_1, \dots, z_m be a basis of $Z_{n,k-1}$ formed by \hat{h} -eigenvectors, the elements $z_i \circ_d y_0$, with $i \in \{1, \dots, m\}$ and $d \geq 0$ generate $Z_{n,k}$ and are \hat{h} -eigenvectors.

The element $z \circ_d z$ appears as $\tau_d(z)$ in [Bed05, Bed10]. Note that using equation (7), we obtain that

$$y_0 \circ_d y_0 = \sum_{i=0}^d (-1)^i y_i y_{d-i}.$$

Applying Proposition 5.1 recursively for the homogeneous components $Z_{n,1}$, $Z_{n,2}$, etc, we obtain the following.

Corollary 5.2. *for $k \geq 1$, the homogeneous component $Z_{n,k}$ is generated as a vector space by elements of the form*

$$((y_0 \circ_{d_1} y_0) \circ_{d_2} y_0) \cdots \circ_{d_{k-1}} y_0$$

where the number of factors is k and the d_i are chosen in such a way that \circ_{d_i} be defined for each i . In particular, the ring Z_n admits a minimal generating set formed by polynomials of this form.

In some cases, it is possible to obtain explicit generators for the invariant ring $\mathbb{F}[\mathbf{y}_n]^x$ as shown in the following example.

Example 5.3. As noted in Example 2.2, $Z_1 = \mathbb{F}[y_0]$ and it is well-known that

$$Z_2 = \mathbb{F}[y_0, y_0 \circ_2 y_0] = \mathbb{F}[y_1, 2y_0y_2 - y_1^2]$$

(see [Dix58]). The generators for Z_3 were determined by Dixmier [Dix58] and we can write them as

$$Z_3 = \mathbb{F}[z_1, z_2, z_3, \zeta_4]$$

where

$$\begin{aligned} z_1 &= y_0; \\ z_2 &= 2y_0y_2 - y_1^2 = y_0 \circ_2 y_0; \\ z_3 &= 3y_0^2y_3 - 3y_0y_1y_2 + y_1^3 = -y_0 \circ_2 y_0 \circ_1 y_0; \\ \zeta_4 &= \frac{z_2^3 + z_3^2}{y_0^2} = -3y_1^2y_2^2 + 8y_0y_2^3 + 6y_1^3y_3 - 18y_0y_1y_2y_3 + 9y_0^2y_3^2 \\ &= (-3/2)y_0 \circ_2 y_0 \circ_1 y_0 \circ_3 y_0. \end{aligned}$$

The expressions in \circ_d are left normed; that is, $a \circ_{d_1} b \circ_{d_2} c = (a \circ_{d_1} b) \circ_{d_2} c$. According to [Oom12] The algebra $\mathbb{F}[\mathbf{y}_4]^x$ is generated by

$$\begin{aligned} z_1 &= y_0; \\ z_2 &= 2y_0y_2 - y_1^2 = y_0 \circ_2 y_0; \\ z_3 &= 3y_0^2y_3 - 3y_0y_1y_2 + y_1^3 = -y_0 \circ_2 y_0 \circ_1 y_0; \\ z_4 &= 2y_0y_4 - 2y_1y_3 + y_2^2 = y_0 \circ_4 y_0; \\ \zeta_5 &= \frac{z_2^3 + z_3^2 - 3y_0^2z_2z_4}{y_0^3} = 2y_2^3 - 6y_1y_2y_3 + 9y_0y_3^2 + 6y_1^2y_4 - 12y_0y_2y_4 \\ &= -2y_0 \circ_2 y_0 \circ_4 y_0. \end{aligned}$$

Since $\mathbb{F}[\mathbf{y}_n]^y$ is a noetherian ring, by the Mauer–Weitzenböck Theorem [Fre17, Theorem 6.1], one may want to describe its Noether normalization. The algebra $\mathbb{F}[\mathbf{y}_2]^x$ is freely generated by z_1 and z_2 and so there is nothing to do. On the other hand, $\mathbb{F}[\mathbf{y}_3]^x$ is an integral extension of $\mathbb{F}[z_1, z_2, \zeta_4]$, or of $\mathbb{F}[z_1, z_3, \zeta_4]$, but not of $\mathbb{F}[z_1, z_2, z_3]$. This is because z_3 satisfies the integral equation $t^2 + z_2^3 - y_0^2\zeta_4 = 0$. Similarly, we obtain that $\mathbb{F}[\mathbf{y}_4]$ is integral over $\mathbb{F}[z_1, z_2, z_4, \zeta_5]$ or over $\mathbb{F}[z_1, z_3, z_4, \zeta_5]$, but not over $\mathbb{F}[z_1, z_2, z_3, z_4]$.

Problem. Determine generators for the Noether normalization of $\mathbb{F}[\mathbf{y}_n]^x$ for $n \geq 5$.

The algebra $\mathbb{F}[\mathbf{y}_5]$ is known to be minimally generated by 23 generators that satisfy 168 relations; see the discussion after Example 27 in [Oom09]. Using the computational algebra system Magma [BCP97], we determined explicit expressions in terms of the \circ_d operation for Z_5 , Z_6 and Z_8 . See the Appendix A.

5.1. The first generator sequence. We use Proposition 4.7 to obtain an explicit generating set for the field of fractions of $\mathbb{F}[\mathbf{y}_n]^x$. Set $z_1 = y_0$, and, for $i \geq 2$, define

$$z_i = \begin{cases} y_0 \circ_i y_0 & \text{if } i \text{ is even} \\ y_0 \circ_1 (y_0 \circ_{i-1} y_0) & \text{if } i \text{ is odd.} \end{cases}$$

The explicit expressions for the first couple of values of the z_i are as follows:

$$\begin{aligned} z_1 &= y_0; \\ z_2 &= 2y_0y_2 - y_1^2; \\ z_3 &= 3y_0^2y_3 - 3y_0y_1y_2 + y_1^3; \\ z_4 &= 2y_0y_4 - 2y_1y_3 + y_2^2; \\ z_5 &= 5y_0^2y_5 - 5y_0y_1y_4 + y_0y_2y_3 + 2y_1^2y_3 - y_1y_2^2; \\ z_6 &= 2y_0y_6 - 2y_1y_5 + 2y_2y_4 - y_3^2; \\ z_7 &= 7y_0^2y_7 - 7y_0y_1y_6 + 3y_0y_2y_5 - y_0y_3y_4 + 2y_1^2y_5 - 2y_1y_2y_4 + y_1y_3^2. \end{aligned}$$

Recall that the lexicographic monomial order on $\mathbb{F}[\mathbf{y}_n]$ is taken ordering the variables in increasing order as y_n, y_{n-1}, \dots, y_0 .

Lemma 5.4. *The element $z_i \in \mathbb{F}[\mathbf{y}_n]^x$ for all $i \leq n$. Furthermore, the leading term of z_i in the lexicographic monomial order is $2y_i y_0$ if i is even, while the leading term is $i \cdot y_i y_0^2$ when i is odd.*

Proof. The fact that $z_i \in \mathbb{F}[\mathbf{y}_n]^x$ follows from Corollary 4.7. Let us show the statement concerning the leading term. For $i \geq 2$ even, we have that

$$z_i = \sum_{j=0}^i (-1)^j y_j y_{i-j}.$$

The leading term of z_i is the term that contains y_i which is $2y_0 y_i$ as claimed. Now for i odd, the definition of \circ_d in (7) and the definition of \hat{f} in (6) imply that

$$\begin{aligned} z_i &= \frac{1}{2n-2i+2} y_0 \hat{f}_n \left(\sum_{j=0}^{i-1} (-1)^j y_j y_{i-1-j} \right) - y_1 \sum_{j=0}^{i-1} (-1)^j y_j y_{i-1-j} \\ &= \frac{1}{2n-2i-2} y_0 \left(\sum_{j=0}^{i-1} (-1)^j (j+1)(n-j) y_{j+1} y_{i-1-j} + \right. \\ &\quad \left. \sum_{j=0}^{i-1} (-1)^j (i-j)(n-i+j+1) y_j y_{i-j} \right) \\ &\quad - y_1 \sum_{j=0}^{i-1} (-1)^j y_j y_{i-1-j}. \end{aligned}$$

The leading term of this expression is the only term containing y_i , which is $i y_0^2 y_i$ as claimed. \square

5.2. The second sequence. We define another sequence of elements in $\mathbb{F}[\mathbf{y}]^x$ which appears in [vW14]. (Recall we assume that the characteristic of \mathbb{F} is zero.) Set, $w_1 = y_0$ and for $n \geq 1$,

$$(8) \quad w_n = \frac{(-1)^n}{n!} y_1^n + \sum_{i=0}^{n-1} \frac{(-1)^i}{j!} y_0^{n-1-i} y_1^i y_{n-i}.$$

The first couple of values of the sequence w_i are as follows.

$$w_1 = y_0$$

$$w_2 = y_0 y_2 - \frac{1}{2} y_1^2$$

$$w_3 = y_0^2 y_3 - y_0 y_1 y_2 + \frac{1}{3} y_1^3$$

$$w_4 = y_0^3 y_4 - y_0^2 y_1 y_3 + \frac{1}{2} y_0 y_1^2 y_2 - \frac{1}{8} y_1^4$$

$$\begin{aligned}
w_5 &= y_0^4 y_5 - y_0^3 y_1 y_4 + \frac{1}{2} y_0^2 y_1^2 y_3 - \frac{1}{6} y_0 y_1^3 y_2 + \frac{1}{30} y_1^5 \\
w_6 &= y_0^5 y_6 - y_0^4 y_1 y_5 + \frac{1}{2} y_0^3 y_1^2 y_4 - \frac{1}{6} y_0^2 y_1^3 y_3 + \frac{1}{24} y_0 y_1^4 y_2 - \frac{1}{144} y_1^6 \\
w_7 &= y_0^6 y_7 - y_0^5 y_1 y_6 + \frac{1}{2} y_0^4 y_1^2 y_5 - \frac{1}{6} y_0^3 y_1^3 y_4 + \frac{1}{24} y_0^2 y_1^4 y_3 - \frac{1}{120} y_0 y_1^5 y_2 + \frac{1}{840} y_1^7 \\
w_8 &= y_0^7 y_8 - y_0^6 y_1 y_7 + \frac{1}{2} y_0^5 y_1^2 y_6 - \frac{1}{6} y_0^4 y_1^3 y_5 + \frac{1}{24} y_0^3 y_1^4 y_4 - \frac{1}{120} y_0^2 y_1^5 y_3 + \frac{1}{720} y_0 y_1^6 y_2 - \frac{1}{5760} y_1^8.
\end{aligned}$$

Lemma 5.5. *We have that $w_n \in \mathbb{F}[\mathbf{y}]^x$ for all $m \geq 1$. Furthermore, the leading term of w_n is $y_0^{i-1} y_n$.*

Proof. The claim that $w_n \in \mathbb{F}[\mathbf{y}]^x$ follows from [vW14, Lemma 3.1], while the claim concerning the leading term is easily verified by inspection of (8). \square

Theorem 5.6. *If the characteristic of \mathbb{F} is zero, then the following are valid.*

- (1) *The elements sets $\{z_1, z_2, \dots\}$ and $\{w_1, w_2, \dots\}$ are algebraically independent in $\mathbb{F}[\mathbf{y}]$.*
- (2) *The fraction field of the algebra Z_n is generated by z_1, \dots, z_n and also by w_1, \dots, w_n .*
- (3) *Z_n lies in $\mathbb{F}[y_0^{-1}, z_1, z_2, \dots, z_n]$ and also in $\mathbb{F}[y_0^{-1}, w_1, w_2, \dots, w_n]$.*
- (4) *The Krull dimension of Z_n is n .*

Proof. This follows from Theorem 3.4, Lemma 5.4 and from Lemma 5.5. \square

Theorem 5.6 implies Theorem 1.1.

6. THE HILBERT SERIES OF Z_n

In this section \mathbb{F} is a field of characteristic zero. Suppose, as in Section 2, that $V_n = \langle y_0, \dots, y_n \rangle$ is an \mathbb{F} -vector space of dimension $n+1$ and consider V_n as an x -module. The action of x on V_n is equivalent to the action of $e \in \mathfrak{sl}_2$ on the n -dimensional irreducible \mathfrak{sl}_2 -module U_n by Lemma 4.2. Thus the invariant ring $Z_n = \mathbb{F}[\mathbf{y}_n]^x$ is isomorphic to the algebra of polynomial invariants $\mathbb{F}[U_n]^e$ where $\mathbb{F}[U_n]$ is the polynomial algebra generated by U_n . Furthermore, this isomorphism is an isomorphism of graded algebras with respect to the grading by degree on $\mathbb{F}[\mathbf{y}_n]$ and $\mathbb{F}[U_n]$. For $n \geq 1$, and $d \geq 0$, let $\delta_{n,d}$ denote the dimension of the degree- d homogeneous component of Z_n (which is equal to the dimension of the degree- d homogeneous component of $\mathbb{F}[U_n]^e$). Set

$$H_n(t) = \sum_{d=0}^{\infty} \delta_{n,d} t^d.$$

Consider the basis of U_n formed by $u_i = x_1^i x_2^{n-i}$ for $i \in \{0, \dots, n\}$. Then

$$h(u_i) = (-n + 2i)u_i$$

and each monomial $m = u_0^{\alpha_0} \cdots u_n^{\alpha_n}$ in the u_i is an eigenvector of h with eigenvalue $w(m)$ (which can be calculated explicitly if needed). The eigenvalue $w(m)$ is also called the weight of m . The degree of such a monomial is $\sum_i \alpha_i$. Noting that $U_{n,d}$ is isomorphic, as an \mathfrak{sl}_2 -module, to the d -th symmetric power $S^d(U_n)$ of the irreducible \mathfrak{sl}_2 -module U_n , the following result follows at once from Corollary 4.3.

Theorem 6.1. $\delta_{n,d}$ is equal to the number of irreducible components of the symmetric power $S^d(U_n)$ considered as an \mathfrak{sl}_2 -module. Moreover, if either n or d is even then $\delta_{n,d}$ is equal to the number of monomials in the u_i with degree d and weight zero; otherwise $\delta_{n,d}$ is equal to the number of such monomials with degree d and weight one.

Example 6.2. Consider for example the case of $n = 1$. Then, for each $d \geq 0$,

$$\delta_{1,d} = 1$$

since, for even d , the only weight zero monomial of degree d is $u_0^{d/2} u_1^{d/2}$, while for odd d , the only weight one monomial of degree d is $u_0^{(d+1)/2} u_1^{(d-1)/2}$. This corresponds to the fact that Z_1 is generated by y_0 and the homogeneous component of degree d is generated by y_0^d (see Example 2.2). The Hilbert series is

$$H_1(t) = \sum_{d=0}^{\infty} t^d = \frac{1}{1-t}.$$

This shows also that Z_1 is generated by y_0 which is a generator of degree 1 and Z_1 is isomorphic to the polynomial algebra $\mathbb{F}[t]$ in one variable t .

Example 6.3. Suppose that $n = 2$. Then the weight zero monomials of degree d are $u_0^i u_1^{d-2i} u_2^i$ with $i \in \{0, \dots, \lfloor d/2 \rfloor\}$. Thus $\delta_{2,d} = \lfloor d/2 \rfloor + 1$ and the sequence $\delta_{2,d}$ is

$$1, 1, 2, 2, 3, 3, 4, 4, 5, 5, \dots$$

Easy computation (distinguishing between d odd and d even) shows that

$$\delta_{2,d} = \delta_{2,d-1} + \delta_{2,d-2} - \delta_{2,d-3}$$

for $d \geq 3$ with initial values $\delta_{2,0} = \delta_{2,1} = 1$ and $\delta_{2,2} = 2$. Hence the characteristic polynomial of $\delta_{2,d}$ considered as a recursive sequence is

$$t^3 - t^2 - t + 1 = (1-t)(1-t^2).$$

In fact, the Hilbert series is

$$H_2(t) = \sum_{d=0}^{\infty} \delta_{2,d} t^d = \frac{1}{(1-t)(1-t^2)}$$

reflecting the fact that $Z_2 = \mathbb{F}[z_1, z_2] = \mathbb{F}[y_0, 2y_0 y_2 - y_1^2]$ generated by a generator of degree one and a generator of degree two and these generators are algebraically independent.

For $k, d, n \in \mathbb{N}_0$, let $p(k, d, n)$ denote the number of partitions of k with at most d parts each of which has size n .

Theorem 6.4. *If $n, d \geq 0$, then $\delta_{n,d} = p(\lfloor dn/2 \rfloor, d, n) = p(\lfloor dn/2 \rfloor, n, d)$. Consequently, $\delta_{n,d} = \delta_{d,n}$*

Proof. Suppose that $m = u_0^{\alpha_0} u_1^{\alpha_1} \cdots u_n^{\alpha_n}$ is a monomial in the u_i of degree d ; that is $\sum_{i \geq 0} \alpha_i = d$. Define the function w' on the generators u_0, \dots, u_n as $w'(u_i) = i$ and extend it multiplicatively to the set of monomials in u_0, \dots, u_n . Then $w'(m) = (w(m) + dn)/2$ where $w(m)$ is the weight (that is, the h -eigenvalue) of the monomial m . If either n is even or d is even, then $\delta_{n,d}$ is equal to the number of monomials of degree d and weight zero, and if m has degree d , then $w(m) = 0$ if and only if $w'(m) = dn/2$. If both n and d are odd, then $\delta_{n,d}$ is equal to the number of monomials of degree d and weight minus one (which is equal to the number of monomials of degree d and weight one) and if the degree of m is d , then $w(m) = -1$ if and only if $w'(m) = (dn - 1)/2 = \lfloor dn/2 \rfloor$. So in both cases, $\delta_{n,d}$ is equal to the number of monomials m of degree d and $w'(m) = \lfloor dn/2 \rfloor$. That is, we need to count monomials $m = u_0^{\alpha_0} u_1^{\alpha_1} \cdots u_n^{\alpha_n}$ such that

$$\sum_{i=0}^n \alpha_i = d \quad \text{and} \quad \sum_{i=0}^n i \cdot \alpha_i = \lfloor dn/2 \rfloor.$$

Now the sequence $\alpha_0, \alpha_1, \dots, \alpha_n$ can be identified with the partition of $\lfloor dn/2 \rfloor$ in which there are α_i parts of size i for all $i \in \{0, \dots, n\}$. Omitting the parts of size zero, we obtain a correspondence between the set of such monomials m and partitions of $\lfloor dn/2 \rfloor$ with at most d parts each of which has size at most n .

Transposing the partitions in question, we obtain that $p(\lfloor dn/2 \rfloor, d, n) = p(\lfloor dn/2 \rfloor, n, d)$, and this implies the final statement of the theorem. \square

Example 6.5. Consider the case $n = 3$. Then $\delta_{3,d}$ is equal to the number of partitions of $\lfloor 3d/2 \rfloor$ with at most d parts of size at most 3. The first elements of this sequence are 1, 1, 2, 3, 5, 6, 8, 10, 13, 15, 18, 21, 25, 28, 32, 36, etc. According to [SI20, A001971], this sequence $a(d) = \delta_{3,d}$ satisfies the recurrence relation

$$a(d) = 2a(d-1) - a(d-2) + a(d-4) - 2a(d-5) + a(d-6).$$

Furthermore, its generating function is

$$H_3(t) = \frac{1 - t^6}{(1-t)(1-t^2)(1-t^3)(1-t^4)}.$$

It is known that $\mathbb{F}[\mathbf{y}_3]^x = Z_3$ is generated by y_0 , $2y_0y_2 - y_1^2$, $3y_0^2y_3 - 3y_0y_1y_2 + y_1^3$, and $-3y_1^2y_2^2 + 8y_0y_2^3 + 6y_1^3y_3 - 18y_0y_1y_2y_3 + 9y_0^2y_3^2$; that is, a generator of degree one, one of degree 2, one of degree three, and one of degree 4 (see Example 5.3).

Example 6.6. Consider the case $n = 4$. Then $\delta_{4,d}$ is equal to the number of partitions of $\lfloor 4d/2 \rfloor = 2d$ with at most d parts of size at most 4. The first elements of this sequence are 1, 1, 3, 5, 8, 12, 18, 24, 33, 43, 55, 69, 86, 104, 126, 150, etc. According to [SI20, A001973], this sequence $a(d) = \delta_{4,d}$ satisfies the recurrence relation

$$a(d) = 2a(d-1) - a(d-3) - a(d-4) + 2a(d-6) - a(d-7).$$

Furthermore, its generating function is

$$H_4(t) = \frac{1 + t^3}{(1 - t)(1 - t^2)^2(1 - t^3)}$$

It is known that $\mathbb{F}[\mathbf{y}_4]^x = Z_4$ is generated by z_1, z_2, z_3, z_4 (as defined in Section 5.1), and

$$\zeta_5 = 2y_2^3 - 6y_1y_2y_3 + 9y_0y_3^2 + 6y_1^2y_4 - 12y_0y_2y_4;$$

that is a generator of degree one, two of degree two and two of degree three (see Example 5.3).

Example 6.7. Consider the case $n = 5$. Then $\delta_{5,d}$ is equal to the number of partitions of $\lfloor 5d/2 \rfloor$ with at most d parts of size at most 5. The first elements of this sequence are 1, 1, 3, 6, 12, 20, 32, 49, 73, 102, 141, 190, 252, 325, 414, 521, 649, 795, 967, 1165, 1394, etc. According to [SI20, A001975], this sequence $a(d) = \delta_{5,d}$ satisfies the recurrence relation

$$\begin{aligned} a(d) &= 2a(d-1) - a(d-2) + a(d-4) - 2a(d-5) + 2a(d-6) - 2a(d-7) \\ &\quad + 2a(d-8) - 2a(d-9) + 2a(d-11) - 2a(d-12) + 2a(d-13) \\ &\quad - 2a(d-14) + 2a(d-15) - a(d-16) + a(d-18) - 2a(d-19) + a(d-20) \end{aligned}$$

Furthermore, its generating function is

$$H_5(t) = \frac{-(t^{14} - t^{13} + 2t^{12} + t^{11} + 2t^{10} + 3t^9 + t^8 + 5t^7 + t^6 + 3t^5 + 2t^4 + t^3 + 2t^2 - t + 1)}{(t^4 + 1)(t^2 + t + 1)(t^2 - t + 1)(t^2 + 1)^2(t + 1)^3(t - 1)^5}.$$

It is claimed in [Oom09, Oom12] that Z_5 is generated by 23 generators.

Theorem 1.3 stated in the Introduction gives a closed formula for the Hilbert series $H_n(t)$ for all n . The formula originates from [Alm80a] where it is derived in a different context with proof in [AF78]. We include a proof for completeness and easy reference.

The proof of Theorem 1.3. Suppose that $u_{-n}, u_{-n+2}, \dots, u_{n-2}, u_n$ are variables such that the weight of u_i is i . Suppose that I_n denotes the index set $\{-n, -n+2, \dots, n-2, n\}$. We denote by α a vector $(\alpha_{-n}, \alpha_{-n+2}, \dots, \alpha_n)$ and by A_d the set of α such that $0 \leq \alpha_i \leq d$ and $\sum_{j \in I_n} \alpha_j = d$. The weight of a monomial $\prod_{j \in I_n} u_j^{\alpha_j}$ is $\sum_{j \in I_n} \alpha_j \cdot j$ and its degree is $\sum_{j \in I_n} \alpha_j$. Recall, for $n \geq 0$ and $d \geq 0$, that $\delta_{n,d}$ is the number of monomials of degree d and weight zero plus the number of monomials of degree d and weight one in the u_i . We are required to show that the right-hand side of (1) can be written as

$$\sum_{d=0}^{\infty} \delta_{n,d} t^d.$$

Let us first work the denominator of the function inside the integral. More specifically, we compute that

$$\prod_{k=0}^n \frac{1}{1 - t \exp(i(n-2k)\varphi)} = \prod_{k \in I_n} \frac{1}{1 - t \exp(ik\varphi)} = \prod_{k \in I_n} \sum_{j \geq 0} \exp(ijk\varphi) t^j.$$

The product of the power series in the last expression is a power series $\sum_{d \geq 0} c_d t^d$ whose d -th coefficient is

$$c_d = \sum_{\alpha \in A_d} \exp\left(i\varphi \sum_{k \in I_n} k\alpha_k\right).$$

The integral in (1) can be split into two parts:

$$(9) \quad \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{1}{\prod_{k=0}^n (1 - t \exp(i(n-2k)\varphi))} d\varphi + \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{\exp(i\varphi)}{\prod_{k=0}^n (1 - t \exp(i(n-2k)\varphi))} d\varphi.$$

By the calculations above, the first summand in (9) can be written as

$$(10) \quad \frac{1}{2\pi} \left(\int_{-\pi}^{\pi} \sum_{d \geq 0} c_d d\varphi \right) t^d = \frac{1}{2\pi} \sum_{d \geq 0} \left(\sum_{\alpha \in A_d} \int_{-\pi}^{\pi} \exp\left(i\varphi \sum_{k \in I_n} k\alpha_k\right) d\varphi \right) t^d.$$

We have, for $k \in \mathbb{Z}$, that

$$\int_{-\pi}^{\pi} \exp(ik\varphi) d\varphi = \begin{cases} 0 & \text{if } k \neq 0 \\ 2\pi & \text{if } k = 0. \end{cases}$$

Hence, among the integrals in the coefficient of t^d in (10) only those are different from zero which are taken for $\alpha \in A_d$ with $\sum_{k \in I_n} k\alpha_k = 0$. Furthermore, for such an α , the value of the integral is 2π and hence, since there is a $1/(2\pi)$ factor, each such α contributes with one to the coefficient of t^d . Now each such choice of α corresponds to a monomial $u_{-n}^{\alpha_{-n}} \cdots u_n^{\alpha_n}$ with degree d and weight zero. Thus the first term in (9) can be written as $\sum_{d \geq 0} a_d t^d$ where a_d is the number of monomials in the u_i of degree d and weight zero.

Now consider the second summand in (9) which is equal

$$(11) \quad \frac{1}{2\pi} \left(\int_{-\pi}^{\pi} \exp(i\varphi) \sum_{d \geq 0} c_d d\varphi \right) t^d = \frac{1}{2\pi} \sum_{d \geq 0} \left(\sum_{\alpha \in A_d} \int_{-\pi}^{\pi} \exp\left(i\varphi \left(1 + \sum_{k \in I_n} k\alpha_k\right)\right) d\varphi \right) t^d.$$

For reasons similar to the ones in the previous paragraph, among the integrals in the coefficient of t^d the ones that are different from zero are taken for $\alpha \in A_d$ with $1 + \sum_{k \in I_n} k\alpha_k = 0$. Now each such choice of α corresponds to a monomial $u_{-n}^{a_{-n}} \cdots u_n^{a_n}$ with degree d and weight -1 . Thus the second term in (9) can be written as $\sum_{d \geq 0} b_d t^d$ where b_d is the number of monomials in the u_i of degree d and weight -1 which is the same as the number of monomials of degree d and weight one. \square

Let $V_{n,d}$ be the space of degree- d polynomials in the variables y_0, \dots, y_n considered as an \mathfrak{sl}_2 -module. That is, $V_{n,1}$ is considered as an irreducible \mathfrak{sl}_2 -module of dimension $n+1$ with weights $n, n-2, \dots, -n$ and $V_{n,d}$ is isomorphic to the symmetric power $S^d(V_{n,1})$. Then $V_{n,d}$ is spanned by monomials of the form $y_0^{\alpha_0} \cdots y_n^{\alpha_n}$ with $\alpha_0 + \cdots + \alpha_n = d$. As above, the weight of such a monomial is $n\alpha_0 + (n-2)\alpha_1 + \cdots - n\alpha_n$. Set

$$M_n(d, w) = \text{the number of monomials in } V_{n,d} \text{ of weight } w$$

and define the generating function $\mu_n(t, z)$ as

$$\mu_n(t, z) = \sum_{d \geq 0, w}^{\infty} M_n(d, w) t^d z^w.$$

Then standard computation with generating functions shows that the following theorem is valid.

Theorem 6.8. *For $n \geq 1$, we have*

$$(12) \quad \mu_n(t, z) = \prod_{k \in \{-n, -n+2, \dots, n-2, n\}} \frac{1}{1 - tz^k}.$$

Example 6.9. If $n = 1$, then

$$\mu_1(t, z) = 1 + t(z^{-1} + z) + t^2(z^{-2} + 1 + z^2) + t^3(z^{-3} + z^{-1} + z + z^3) + \cdots$$

In this case we have to count monomials of even degree and weight zero and monomials of odd degree and weight one. These monomials are counted by the terms

$$1 + tz + t^2 + t^3z + \cdots$$

showing that $\delta_{1,d} = 1$ for all $d \geq 0$. This agrees with the values of $\delta_{1,d}$ computed in Example 6.2.

Example 6.10. If $n = 2$, then

$$\begin{aligned} \mu_2(t, z) &= 1 + t(z^2 + z^{-2} + 1) + t^2(z^4 + z^{-4} + z^2 + z^{-2} + 2) \\ &\quad + t^3(z^6 + z^{-6} + z^4 + z^{-4} + 2z^2 + 2z^{-2} + 2) \\ &\quad + t^4(z^8 + z^{-8} + z^6 + z^{-6} + 2z^4 + 2z^{-4} + 2z^2 + 2z^{-2} + 3) \\ &\quad + t^5(z^{10} + z^{-10} + z^8 + z^{-8} + 2z^6 + 2z^{-6} + 2z^4 + 2z^{-4} + 3z^2 + 3z^{-2} + 3) + O(t^6) \end{aligned}$$

The part of this series expansion containing only the terms with z^0 is

$$1 + t + 2t^2 + 2t^3 + 3t^4 + 3t^5 + \cdots$$

and the coefficients agree with the values for $\delta_{2,d}$ computed in Example 6.3.

Example 6.11. Let us compute $H_3(t)$ using the method outlined by Ekhad and Zeilberger. By Theorem 6.8 we have that

$$\mu_3(t, z) = \frac{1}{(1 - z^{-3}t)(1 - z^{-1}t)(1 - zt)(1 - z^3t)} = z^4 \frac{1}{(z^3 - t)(z - t)(1 - zt)(1 - z^3t)}.$$

Using that the factors $z^3 - t$, $z - t$, $1 - zt$, $1 - z^3t$ are coprime polynomials in the variable z over the rational function field $\mathbb{Q}(t)$, we have that there are polynomials $Q_1, Q_2, Q_3, Q_4 \in \mathbb{Q}(t)[z]$ such that

$$\begin{aligned} \mu_3(t, z) &= z^4 \left(\frac{Q_1}{z^3 - t} + \frac{Q_2}{z + t} + \frac{Q_3}{1 - zt} + \frac{Q_4}{1 - z^3t} \right) \\ &= z \frac{Q_1}{1 - z^{-3}t} + z^3 \frac{Q_2}{1 - z^{-1}t} + z^4 \frac{Q_3}{1 - zt} + z^4 \frac{Q_4}{1 - z^3t}. \end{aligned}$$

The expressions Q_i in the last displayed equation can be determined in a computational algebra system, such as Magma [BCP97], using the Extended Euclidean Algorithm. Writing $\mu_3(t, z) = \sum_{k \in \mathbb{Z}} q_k z^k$ where $q_k \in \mathbb{Q}(t)$, remembering that $n = 3$ is odd, the Hilbert series $H_3(t)$ is determined by the coefficients q_1 and q_0 and they depend only on Q_1 and Q_2 . This way one obtains the same Hilbert series as in Example 6.5. The web page [EZ22] contains a report on computations that determined the rational expressions for $H_n(t)$ for $n \leq 16$ using partial fraction decomposition of $\mu_n(t, z)$ as given in equation (12); see also the note [EZ19] for details. This procedure was also implemented by the authors of the present paper and the implementation was used to verify the rational expressions for $H_n(t)$ in Examples 6.3, and 6.5–6.7.

7. PRIME CHARACTERISTIC

Throughout this final section, we let $n \geq 1$ fixed, \mathbb{F} denotes a field of characteristic p with $p \geq n + 1$, and $\mathfrak{g} = \mathfrak{g}(n + 2)$ is the $(n + 2)$ -dimensional standard filiform Lie algebra defined over \mathbb{F} . Note that $U(\mathfrak{g})$ can be viewed as $U_{\mathbb{Z}}(\mathfrak{g}) \otimes \mathbb{F}$ where $U_{\mathbb{Z}}(\mathfrak{g})$ is the universal enveloping algebra of the standard filiform Lie algebra $\mathfrak{g}_{\mathbb{Z}}(n + 2)$ defined over \mathbb{Z} . Since the elements z_1, \dots, z_n in Section 5.1 are central in $U_{\mathbb{Z}}(\mathfrak{g})$, they are also central in $U(\mathfrak{g}) = U_{\mathbb{Z}}(\mathfrak{g}) \otimes \mathbb{F}$. In characteristic p , however, there are other central elements that do not occur in characteristic zero. Let $Z_p(\mathfrak{g})$ denote the p -center of $U(\mathfrak{g})$; that is

$$Z_p(\mathfrak{g}) = \mathbb{F}[x^p, y_0, y_1^p, \dots, y_n^p].$$

The condition that $p \geq n + 1$ implies that $Z_p(\mathfrak{g})$ is isomorphic to the polynomial algebra $\mathbb{F}[\mathfrak{t}_{n+2}]$ and also that $Z_p(\mathfrak{g}) \subseteq Z(\mathfrak{g})$. Let $K(\mathfrak{g})$ and $K_p(\mathfrak{g})$ denote the fraction fields of $Z(\mathfrak{g})$ and $Z_p(\mathfrak{g})$ respectively. We also let $D(\mathfrak{g})$ denote the division algebra of fractions of $U(\mathfrak{g})$. Suppose that z_i is the sequence of elements defined in Section 5. The leading coefficient of z_i is either 2 or i by Lemma 5.4, which is nonzero by the assumption that $p \geq n + 1$.

In the rest of this section, the elements z_1, \dots, z_n are the same as defined in Section 5 and recall that $z_1 = y_0 \in Z_p(\mathfrak{g})$.

Theorem 7.1. *We have that $K(\mathfrak{g}) = K_p(\mathfrak{g})(z_2, \dots, z_n)$ and $\dim_{K(\mathfrak{g})} D(\mathfrak{g}) = p^2$.*

Proof. Consider the chain

$$K_p(\mathfrak{g}) \subset K(\mathfrak{g}) \subset D(\mathfrak{g})$$

of division algebras (where the first two terms are fields). We have that

$$\dim_{K_p(\mathfrak{g})} D(\mathfrak{g}) = p^{\dim \mathfrak{g} - \dim C(\mathfrak{g})} = p^{n+1} \quad \text{and} \quad \dim_{K(\mathfrak{g})} D(\mathfrak{g}) \geq p^2$$

(see [dJS22, Theorem 3.2]). Furthermore, since $a^p \in K_p(\mathfrak{g})$ for all $a \in K(\mathfrak{g})$, $K(\mathfrak{g})$ is a purely inseparable extension of $K_p(\mathfrak{g})$ of dimension p^k where $k \leq n-1$. Since $z_i \notin K_p(\mathfrak{g})(z_2, \dots, z_{i-1})$ but $z_i^p \in K_p(\mathfrak{g})$ for all $i \geq 2$ and $K_p(\mathfrak{g})(z_2, \dots, z_i)$ is a purely inseparable extension of $K_p(\mathfrak{g})(z_2, \dots, z_{i-1})$, it follows that

$$\dim_{K_p(\mathfrak{g})(z_2, \dots, z_{i-1})} K_p(\mathfrak{g})(z_2, \dots, z_{i-1}, z_i) = p.$$

This implies that $K(\mathfrak{g}) = K_p(z_2, \dots, z_n)$ and $\dim_{K(\mathfrak{g})} D(\mathfrak{g}) = p^2$. \square

Proposition 7.2. *Suppose that $R_n = Z_p(\mathfrak{g})[y_0^{-1}, z_2, \dots, z_n]$ and, for $i \in \{2, \dots, n\}$, set $\alpha_i = z_i^p$. Then $\alpha_i \in Z_p(\mathfrak{g})$ and following are valid.*

- (1) $R_n \cong Z_p(\mathfrak{g})[y_0^{-1}, t_2, \dots, t_n]/(t_2^p - \alpha_2, \dots, t_n^p - \alpha_n)$.
- (2) R_n is a regular ring.

Consequently, R_n is a normal domain (that is, integrally closed in its field of fractions).

Proof. The fact that $\alpha_i \in Z_p(\mathfrak{g})$ follows from the fact that α_i is an expression in the commuting variables y_0, y_1, \dots, y_n and that $y_0 \in Z_p(\mathfrak{g})$ and, for all $i \geq 1$, $y_i^p \in Z_p(\mathfrak{g})$.

We will prove the assertions (1)–(2) of the lemma simultaneously by induction on n . The basis of the induction is the case when $n = 1$ and the assertions in this case are valid since $R_1 = Z_p(\mathfrak{g})[y_0^{-1}] \cong \mathbb{F}[x^p, y_0, y_0^{-1}]$ (x^p and y_0 are algebraically independent) is a regular domain. Since a regular domain is also normal, we obtain that R_1 is a normal domain.

Suppose that assertions (1) and (2) hold for the algebra R_{n-1} that corresponds to the standard filiform Lie algebra $\mathfrak{g}(n+1)$. That is,

$$\begin{aligned} R_{n-1} &= \mathbb{F}[y_0^{-1}, y_0, y_1^p, \dots, y_n^p, z_2, \dots, z_{n-1}] \\ &\cong Z_p(\mathfrak{g}(n+1))[y_0^{-1}, z_2, \dots, z_{n-1}] \\ &\cong Z_p(\mathfrak{g}(n+1))[y_0^{-1}, t_2, \dots, t_{n-1}]/(t_2^p - \alpha_2, \dots, t_{n-1}^p - \alpha_{n-1}). \end{aligned}$$

Further, the induction hypothesis also states that R_{n-1} is regular and hence it is normal. Since, y_n^p is a free variable over R_{n-1} , $R_{n-1}[y_n^p]$ is normal. Also note that $\alpha_i \in R_{n-1}[y_n^p] \setminus (R_{n-1}[y_n^p])^p$ and $t_n^p - \alpha_n$ is prime in $R_{n-1}[y_n^p][t_n]$ (see [dJS22, Lemma 2.5]). Now since $t_n^p - \alpha_n$ is prime, we obtain that

$$\begin{aligned} R_n &= Z_p(\mathfrak{g})[y_0^{-1}, z_2, \dots, z_n] = R_{n-1}[y_n^p, z_n] \\ &\cong R_{n-1}[y_n^p][t_n]/(t_n^p - \alpha_n) \\ &\cong Z_p(\mathfrak{g})[y_0^{-1}, t_2, \dots, t_n]/(t_2^p - \alpha_2, \dots, t_n^p - \alpha_n). \end{aligned}$$

This proves claim (1).

Let us turn to claim (2). We identify $Z_p(\mathfrak{g})[y_0^{-1}, t_2, \dots, t_n]$ with the localization $\mathbb{F}[\mathbf{u}_{2n+1}]_{u_2}$ at the multiplicative set generated by u_2 of the polynomial ring $\mathbb{F}[\mathbf{u}_{2n+1}]$ in $2n+1$ variables where $x, y_0, y_1^p, \dots, y_n^p$ are identified with the variables u_1, u_2, \dots, u_{n+2}

and t_2, \dots, t_n are identified with the variables u_{n+3}, \dots, u_{2n+1} , respectively. In particular, y_0 corresponds to the variable u_2 and this is why the polynomial ring is localized at u_2 . Suppose, for $i \in \{2, \dots, n\}$ that f_i is the image of the polynomial $t_i^p - \alpha_i$ in $\mathbb{F}[\mathbf{u}_{2n-2}]_{u_2}$. The argument in the previous paragraph implies that

$$R_n \cong \mathbb{F}[\mathbf{u}_{2n+1}]_{u_2}/(f_2, \dots, f_n).$$

For $i = \{2, \dots, n\}$, let D_i denote the derivation $\partial/\partial u_{i+2}$ of $\mathbb{F}[\mathbf{u}_{2n+1}]_{u_2}$. Then we have from Lemma 5.4 that $D_i(f_j) = 0$ for all $j < i$ and $D_i(f_i) = c_i u_2^{k_i}$ where k_i and c_i are positive integers and $c_i < p$. Hence the Jacobian matrix $(D_i f_j)$ is upper triangular and $\det(D_i f_j) = c u_2^k$ where $c, k \in \mathbb{N}$ and $p \nmid c$.

Suppose that $\bar{P} \in \text{Spec}(R)$. Then $\bar{P} = P/(f_2, \dots, f_n)$ for some prime ideal $P \in \text{Spec}(\mathbb{F}[\mathbf{u}_{2n+1}]_{u_2})$ such that $u_2 \notin P$ and $(f_2, \dots, f_n) \subseteq P$. Thus $\det(D_i f_j) \notin P$, otherwise $u_2 \in P$ would be true. The Jacobian Criterion for Regularity [Mat89, Theorem 30.4] implies that $\mathbb{F}[\mathbf{u}_{2n+1}]_P/(f_2, \dots, f_n)_P$ is a regular ring; that is $(R_n)_{\bar{P}}$ is a regular ring. This shows that R_n is regular. Since a regular ring is normal, R_n is also a normal domain. \square

The proof of Theorem 1.4. Set $R = Z_p(\mathfrak{g})[z_2, \dots, z_n]$ and $R[y_0^{-1}] = Z_p(\mathfrak{g})[y_0^{-1}, z_2, \dots, z_n]$. Theorem 7.1 implies that

$$K(\mathfrak{g}) = K_p(\mathfrak{g})(z_2, \dots, z_n)$$

and so the fraction fields of R , $R[y_0^{-1}]$ and $Z(\mathfrak{g})$ coincide. Let us denote this field by K . For domains $A \subseteq B$, let A^B denote the integral closure of A in B . Since $R[y_0^{-1}]$ is integrally closed,

$$R^K \subseteq R[y_0^{-1}]^K = R[y_0^{-1}].$$

This means that the integral closure R^K must be contained in $R[y_0^{-1}]$ and hence $R^K = R^{R[y_0^{-1}]}$. For the other statement, recall that $Z(\mathfrak{g})$ is an integral extension of R in K and $Z(\mathfrak{g})$ is integrally closed in K (by Zassenhaus' Theorem [Zas54]). Thus

$$Z(\mathfrak{g}) \subseteq R^K \subseteq Z(\mathfrak{g})^K = Z(\mathfrak{g}).$$

It follows that equality must hold in each step of the previous chain which implies that $Z(\mathfrak{g}) = R^K$ as claimed. \square

REFERENCES

- [AF78] Gert Almkvist and Robert Fossum. Decomposition of exterior and symmetric powers of indecomposable $\mathbf{Z}/p\mathbf{Z}$ -modules in characteristic p and relations to invariants. In *Séminaire d'Algèbre Paul Dubreïl, 30ème année (Paris, 1976–1977)*, volume 641 of *Lecture Notes in Math.*, pages 1–111. Springer, Berlin, 1978.
- [Alm80a] Gert Almkvist. Invariants, mostly old ones. *Pacific J. Math.*, 86(1):1–13, 1980.
- [Alm80b] Gert Almkvist. Reciprocity theorems for representations in characteristic p . In *Séminaire d'Algèbre Paul Dubreïl et Marie-Paule Malliavin, 32ème année (Paris, 1979)*, volume 795 of *Lecture Notes in Math.*, pages 1–9. Springer, Berlin, 1980.

- [BCP97] Wieb Bosma, John Cannon, and Catherine Playoust. The Magma algebra system. I. The user language. *J. Symbolic Comput.*, 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- [Bed05] L. Bedratyuk. Casimir elements and kernel of Weitzenböck derivation. 2005. DOI: 10.48550/ARXIV.MATH/0512520; Preprint: arxiv.org/abs/math/0512520.
- [Bed07] Leonid Bedratyuk. On complete system of invariants for the binary form of degree 7. *J. Symbolic Comput.*, 42(10):935–947, 2007.
- [Bed08] L. P. Bedratyuk. The kernel of the generalized Weitzenböck derivation of a polynomial ring. *Mat. Stud.*, 29(2):115–120, 2008.
- [Bed09] Leonid Bedratyuk. A complete minimal system of covariants for the binary form of degree 7. *J. Symbolic Comput.*, 44(2):211–220, 2009.
- [Bed10] L. P. Bedratyuk. Kernels of derivations of polynomial rings and Casimir elements. *Ukrain. Mat. Zh.*, 62(4):435–452, 2010.
- [Bed11a] Leonid Bedratyuk. Bivariate Poincaré series for the algebra of covariants of a binary form. *ISRN Algebra*, pages Art. ID 312789, 11, 2011.
- [Bed11b] Leonid Bedratyuk. Weitzenböck derivations and classical invariant theory, II: The symbolic method. *Serdica Math. J.*, 37(2):87–106, 2011.
- [BI15] Leonid Bedratyuk and Nadia Ilash. The degree of the algebra of covariants of a binary form. *J. Commut. Algebra*, 7(4):459–472, 2015.
- [Dix58] Jacques Dixmier. Sur les représentations unitaires des groupes de Lie nilpotents. III. *Canadian J. Math.*, 10:321–348, 1958.
- [dJS22] Vanderlei Lopes de Jesus and Csaba Schneider. The center of the universal enveloping algebras of small-dimensional nilpotent Lie algebras in prime characteristic. To appear in *Beitr. Algebra Geom.* DOI: doi.org/10.1007/s13366-022-00631-5; Preprint: arxiv.org/abs/2111.13432, 2022.
- [EZ19] Shalosh B. Ekhad and Doron Zeilberger. In how many ways can I carry a total of n coins in my two pockets, and have the same amount in both pockets?, 2019. DOI: 10.48550/ARXIV.1901.08172; Preprint: arxiv.org/abs/1901.08172.
- [EZ22] Shalosh B. Ekhad and Doron Zeilberger. In how many ways can I carry a total of n coins in my two pockets, and have the same amount in both pockets? Exclusively published in the Personal Journal of Shalosh B. Ekhad and Doron Zeilberger and arxiv.org; URL: sites.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/change.html, accessed in August of 2022.
- [Fre13] Gene Freudenburg. Foundations of invariant theory for the down operator. *J. Symbolic Comput.*, 57:19–47, 2013.
- [Fre17] Gene Freudenburg. *Algebraic theory of locally nilpotent derivations*, volume 136 of *Encyclopedia of Mathematical Sciences*. Springer-Verlag, Berlin, second edition, 2017. Invariant Theory and Algebraic Transformation Groups, VII.
- [Jac79] Nathan Jacobson. *Lie algebras*. Dover Publications, Inc., New York, 1979. Republication of the 1962 original.
- [Kow14] Emmanuel Kowalski. *An introduction to the representation theory of groups*, volume 155 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2014.
- [Mat89] Hideyuki Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1989. Translated from the Japanese by M. Reid.
- [Oom09] Alfons I. Ooms. Computing invariants and semi-invariants by means of Frobenius Lie algebras. *J. Algebra*, 321(4):1293–1312, 2009.

- [Oom12] Alfons I. Ooms. The Poisson center and polynomial, maximal Poisson commutative subalgebras, especially for nilpotent Lie algebras of dimension at most seven. *J. Algebra*, 365:83–113, 2012.
- [SI20] Neil J. A. Sloane and The OEIS Foundation Inc. The on-line encyclopedia of integer sequences, 2020. URL: oeis.org.
- [vW14] Libor Šnobl and Pavel Winternitz. *Classification and identification of Lie algebras*, volume 33 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 2014.
- [Zas54] Hans Zassenhaus. The representations of Lie algebras of prime characteristic. *Proc. Glasgow Math. Assoc.*, 2:1–36, 1954.

APPENDIX A. THE GENERATING SETS OF Z_5 , Z_6 , Z_8

We computed a minimal generating set for Z_5 of this form consisting of the following 23 polynomials.

- degree 1: $z_1 = y_0$;
- degree 2: $z_2 = y_0 \circ_2 y_0$; $z_3 = y_0 \circ_4 y_0$;
- degree 3: $z_4 = y_0 \circ_2 y_0 \circ_1 y_0$; $z_5 = y_0 \circ_4 y_0 \circ_1 y_0$; $z_6 = y_0 \circ_4 y_0 \circ_2 y_0$;
- degree 4: $z_7 = y_0 \circ_4 y_0 \circ_2 y_0 \circ_1 y_0$; $z_8 = y_0 \circ_4 y_0 \circ_2 y_0 \circ_2 y_0$; $z_9 = y_0 \circ_4 y_0 \circ_1 y_0 \circ_5 y_0$;
- degree 5: $z_{10} = y_0 \circ_4 y_0 \circ_2 y_0 \circ_2 y_0 \circ_1 y_0$; $z_{11} = z_3^2 \circ_3 y_0$; $z_{12} = z_3^2 \circ_4 y_0$;
- degree 6: $z_{13} = z_3^2 \circ_4 y_0 \circ_1 y_0$; $z_{14} = z_3^2 \circ_3 y_0 \circ_3 y_0$;
- degree 7: $z_{15} = z_3^2 \circ_3 y_0 \circ_3 y_0 \circ_1 y_0$; $z_{16} = z_3^2 \circ_4 y_0 \circ_1 y_0 \circ_4 y_0$;
- degree 8: $z_{17} = z_3 z_{12} \circ_3 y_0$; $z_{18} = z_3 z_{11} \circ_5 y_0$;
- degree 9: $z_{19} = z_3 z_{12} \circ_3 y_0 \circ_2 y_0$;
- degree 11: $z_{20} = z_3 z_{17} \circ_4 y_0$;
- degree 12: $z_{21} = z_3^2 z_{16} \circ_5 y_0$;
- degree 13: $z_{22} = z_3 z_{12}^2 \circ_4 y_0$;
- degree 18: $z_{23} = z_3^2 z_{22} \circ_5 y_0$;

The minimal generating system for Z_6 consists of the following 26 polynomials:

- degree 1: $z_1 = y_0$;
- degree 2: $z_2 = y_0 \circ_2 y_0$; $z_3 = y_0 \circ_4 y_0$; $z_4 = y_0 \circ_6 y_0$;
- degree 3: $z_5 = y_0 \circ_2 y_0 \circ_1 y_0$; $z_6 = y_0 \circ_4 y_0 \circ_1 y_0$; $z_7 = y_0 \circ_4 y_0 \circ_2 y_0$; $z_8 = y_0 \circ_4 y_0 \circ_4 y_0$;
- degree 4: $z_9 = y_0 \circ_4 y_0 \circ_2 y_0 \circ_1 y_0$; $z_{10} = y_0 \circ_4 y_0 \circ_4 y_0 \circ_1 y_0$; $z_{11} = y_0 \circ_4 y_0 \circ_4 y_0 \circ_2 y_0$;
 $z_{12} = y_0 \circ_4 y_0 \circ_2 y_0 \circ_6 y_0$;
- degree 5: $z_{13} = y_0 \circ_4 y_0 \circ_4 y_0 \circ_2 y_0 \circ_1 y_0$; $z_{14} = y_0 \circ_4 y_0 \circ_4 y_0 \circ_2 y_0 \circ_3 y_0$;
 $z_{15} = y_0 \circ_4 y_0 \circ_4 y_0 \circ_2 y_0 \circ_4 y_0$;
- degree 6: $z_{16} = y_0 \circ_4 y_0 \circ_4 y_0 \circ_2 y_0 \circ_4 y_0 \circ_1 y_0$; $z_{17} = y_0 \circ_4 y_0 \circ_4 y_0 \circ_2 y_0 \circ_3 y_0 \circ_2 y_0$;
 $z_{18} = z_3 z_8 \circ_6 y_0$;

degree 7: $z_{19} = z_8^2 \circ_3 y_0; z_{20} = z_8^2 \circ_4 y_0;$

degree 8: $z_{21} = z_8^2 \circ_3 y_0 \circ_4 y_0;$

degree 9: $z_{22} = z_8^2 \circ_3 y_0 \circ_4 y_0 \circ_2 y_0;$

degree 10: $z_{23} = z_8^2 \circ_3 y_0 \circ_4 y_0 \circ_2 y_0 \circ_4 y_0; z_{24} = z_8^3 \circ_6 y_0;$

degree 12: $z_{25} = z_8 z_{21} \circ_4 y_0;$

degree 15: $z_{26} = z_8^2 z_{21} \circ_6 y_0;$

The generators of Z_8 are the following polynomials.

Degree 1: $z_1 = y_0;$

Degree 2 : $z_2 = y_0 \circ_2 y_0; z_3 = y_0 \circ_4 y_0; z_4 = y_0 \circ_6 y_0; z_5 = y_0 \circ_8 y_0;$

Degree 3 : $z_6 = y_0 \circ_2 y_0 \circ_1 y_0; z_7 = y_0 \circ_4 y_0 \circ_1 y_0; z_8 = y_0 \circ_6 y_0 \circ_1 y_0; z_9 = y_0 \circ_4 y_0 \circ_2 y_0;$

$z_{10} = y_0 \circ_6 y_0 \circ_2 y_0; z_{11} = y_0 \circ_6 y_0 \circ_3 y_0; z_{12} = y_0 \circ_6 y_0 \circ_4 y_0; z_{13} = y_0 \circ_4 y_0 \circ_8 y_0;$

Degree 4 : $z_{14} = y_0 \circ_4 y_0 \circ_2 y_0 \circ_1 y_0; z_{15} = y_0 \circ_6 y_0 \circ_2 y_0 \circ_1 y_0; z_{16} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_1 y_0;$

$z_{17} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_1 y_0; z_{18} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_2 y_0; z_{19} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_2 y_0;$

$z_{20} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0; z_{21} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0; z_{22} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0;$

$z_{23} = y_0 \circ_6 y_0 \circ_2 y_0 \circ_8 y_0;$

Degree 5 : $z_{24} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_2 y_0 \circ_1 y_0; z_{25} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0 \circ_1 y_0;$

$z_{26} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_1 y_0; z_{27} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_2 y_0;$

$z_{28} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0 \circ_2 y_0; z_{29} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0 \circ_3 y_0;$

$z_{30} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_3 y_0; z_{31} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0;$

$z_{32} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_4 y_0; z_{33} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0;$

$z_{34} = z_4^2 \circ_8 y_0;$

Degree 6 : $z_{35} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_1 y_0; z_{36} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0 \circ_1 y_0;$

$z_{37} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0; z_{38} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0 \circ_3 y_0;$

$z_{39} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_4 y_0 \circ_3 y_0; z_{40} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0;$

$z_{41} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_4 y_0 \circ_4 y_0; z_{42} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_3 y_0 \circ_6 y_0;$

$z_{43} = z_4 z_{12} \circ_8 y_0;$

Degree 7 : $z_{44} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0; z_{45} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0 \circ_3 y_0;$

$z_{46} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_4 y_0 \circ_4 y_0 \circ_3 y_0; z_{47} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0;$

$z_{48} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0 \circ_5 y_0; z_{49} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0 \circ_6 y_0;$

$z_{50} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0; z_{51} = z_{12}^2 \circ_8 y_0;$

Degree 8 : $z_{52} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0;$

$z_{53} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0 \circ_6 y_0 \circ_2 y_0;$

$z_{54} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0 \circ_5 y_0 \circ_4 y_0;$

$$z_{55} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0 \circ_4 y_0;$$

$$z_{56} = z_4 z_{33} \circ_6 y_0; z_{57} = y_0 \circ_6 y_0 \circ_3 y_0 \circ_5 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0 \circ_6 y_0; z_{58} = z_4 z_{32} \circ_8 y_0;$$

$$\text{Degree 9 : } z_{59} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0 \circ_5 y_0 \circ_4 y_0 \circ_4 y_0;$$

$$z_{60} = z_4 z_{42} \circ_6 y_0; z_{61} = z_{12} z_{33} \circ_6 y_0;$$

$$z_{62} = y_0 \circ_6 y_0 \circ_4 y_0 \circ_3 y_0 \circ_6 y_0 \circ_2 y_0 \circ_6 y_0 \circ_2 y_0 \circ_6 y_0; z_{63} = z_4 z_{41} \circ_8 y_0;$$

$$\text{Degree 10 : } z_{64} = z_4 z_{50} \circ_6 y_0; z_{65} = z_4 z_{49} \circ_6 y_0; z_{66} = z_4 z_{48} \circ_8 y_0;$$

$$\text{Degree 11 : } z_{67} = z_4 z_{56} \circ_6 y_0; z_{68} = z_4 z_{57} \circ_6 y_0;$$

$$\text{Degree 12 : } z_{69} = z_4 z_{60} \circ_6 y_0;$$

In all of these computations, we used the information given in [Fre13] concerning the maximum degree of the polynomials in a minimal generating set. The degree-15 polynomial in the generating set for Z_6 has 1370 nonzero terms, while the degree-12 polynomial in the generating set of Z_8 has 3651 nonzero terms.

(Lopes) DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE CIÊNCIAS EXATAS, UNIVERSIDADE FEDERAL DE MINAS GERAIS, AV. ANTÔNIO CARLOS 6627, BELO HORIZONTE, MG, BRAZIL. EMAIL: vanderleilopesbh@gmail.com

(Schneider) DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE CIÊNCIAS EXATAS, UNIVERSIDADE FEDERAL DE MINAS GERAIS, AV. ANTÔNIO CARLOS 6627, BELO HORIZONTE, MG, BRAZIL. EMAIL: csaba@mat.ufmg.br, URL: schcs.github.io/WP/