

Moduli of G -constellations and crepant resolutions II: the Craw-Ishii conjecture

Ryo Yamagishi

Abstract

For any given finite subgroup $G \subset SL_3(\mathbb{C})$, we show that every projective crepant resolution X of the quotient variety \mathbb{C}^3/G is isomorphic to the moduli space of θ -stable G -constellations for a generic stability condition θ , as conjectured by Craw and Ishii. We also show that generators of the Cox ring of X can be obtained from semi-invariants for representations of the McKay quiver of G . Moreover, we give an explicit description of the two symplectic resolutions of \mathbb{C}^4/T constructed by Lehn and Sorger for a binary tetrahedral group $T \subset Sp_4(\mathbb{C})$ as fine moduli spaces of T -constellations.

1 Introduction

Let G be a finite subgroup of $SL_n(\mathbb{C})$. The philosophy of the McKay correspondence is that the geometric properties of a crepant resolution of \mathbb{C}^n/G should reflect the representation-theoretic properties of G (cf. [Re2]). The natural way to see the explicit correspondence is to realize a crepant resolution of \mathbb{C}^n/G as the G -Hilbert scheme (see e.g. [Cr], [CCL]) or, more generally, a moduli space of G -constellations. In dimension two, it is well known that the unique crepant resolution is obtained as the G -Hilbert scheme. In dimension three, it was also proved by Craw and Ishii [CI] that every projective crepant resolution is obtained as the moduli space \mathcal{M}_θ of G -constellations for a suitable stability condition θ when G is abelian. In general \mathcal{M}_θ for any finite subgroup $G \subset SL_3(\mathbb{C})$ is known to be a crepant resolution for any generic stability condition θ by the result of Bridgeland, King and Reid [BKR], and it is conjectured that the result by Craw and Ishii still holds for non-abelian G [IINdC, Conjecture 1.4]. The main result of this article is that this conjecture is affirmative:

Theorem 1.1. (*=Theorem 4.1*) *Let $G \subset SL_3(\mathbb{C})$ be a finite subgroup. Then every projective crepant resolution of \mathbb{C}^3/G is isomorphic to the moduli space of θ -stable G -constellations for a generic stability condition θ .*

In this paper we also give an explicit construction of a torus-equivariant morphism from the spectrum of the Cox ring of a crepant resolution of \mathbb{C}^3/G to (a certain quotient of) a space of G -constellations, as a generalization of the construction in Part I [Y2] of the present paper. In [Y2], we gave a necessary and sufficient condition for a given (not-necessarily-projective) crepant resolution $X \rightarrow \mathbb{C}^n/G$ to admit a moduli description in

terms of a natural G -constellation family when G is abelian. The key idea to prove this result was to construct an equivariant morphism between certain toric varieties so that it descends to a birational morphism from X to a candidate of a fine moduli space. In this paper we generalize this construction to non-abelian G in dimension three so that we obtain an embedding $\mathcal{W} \hookrightarrow Y_{\mathcal{W}}$ of (a quotient of) a space of G -constellations into a toric variety such that it induces an embedding of \mathcal{M}_{θ} into a toric variety for any generic θ (Proposition 3.5). This embedding of \mathcal{M}_{θ} is so nice that it enables us to understand the behavior of \mathcal{M}_{θ} under variation of θ through the change of the ambient toric variety. We will demonstrate this embedding and investigate the geometric structure of the moduli space explicitly via concrete examples in Section 3.5. The embedding will also be used to prove Theorem 1.1 (see the proof of Lemma 4.3).

The embedding $\mathcal{W} \hookrightarrow Y_{\mathcal{W}}$ will be constructed by using the Cox ring $\text{Cox}(X)$ of $X = \mathcal{M}_{\theta}$, whose generators play a similar role to homogeneous coordinates of a projective space. On the other hand \mathcal{M}_{θ} is regarded as a moduli space of certain representations of the McKay quiver of G , and semi- (or relative) invariants for these representations also play a role of homogeneous coordinates of \mathcal{M}_{θ} . Then the equivariant morphism

$$\mathfrak{X} := \text{Spec Cox}(X) \rightarrow \mathcal{W}$$

is obtained by assigning to each semi-invariant a global section of a line bundle on X . Moreover, as an application of Theorem 1.1, we will show that generators of $\text{Cox}(X)$ can be computed from the semi-invariants for the McKay quiver. More precisely, there is a ring homomorphism φ from the ring $\mathbb{C}[\mathcal{W}]$ of semi-invariants to the (semi-)invariant ring $\mathbb{C}[\mathbb{C}^3]^{[G,G]}$ such that the ‘‘associated elements’’ to the images of homogeneous generators of $\mathbb{C}[\mathcal{W}]$ under φ give a generating system of $\text{Cox}(X)$ (Proposition 4.10).

The strategy for the proof of Theorem 1.1 is based on the one given by Craw and Ishii for abelian cases. To explain this, let us recall that each GIT-chamber $C \subset \Theta$ in the space of stability conditions gives a fine moduli space $X := \mathfrak{M}_C$ and its universal family \mathcal{U}_C . By the result of [BKR], the Fourier-Mukai transform by \mathcal{U}_C gives an equivalence between the bounded derived category of coherent sheaves on X and that of G -equivariant sheaves on \mathbb{C}^3 . This induces an isomorphism

$$\phi_C^* : \Theta \rightarrow F^1$$

from the space of stability conditions to the subspace F^1 inside the Grothendieck group $K(X)_{\mathbb{R}}$ generated by sheaves whose supports have codimension at least one. The proof of the theorem will be done by studying the behavior of $\phi_C^*(C)$ under crossing a wall of Type 0 inside Θ . Here a wall of Type 0 means a codimension-one face W of \bar{C} whose associated morphism $X \rightarrow \tilde{\mathfrak{M}}_W$ induced by variation of GIT is an isomorphism. The significant fact to prove Theorem 1.1 is that, similarly to abelian cases, crossing such a wall gives a wall-crossing in F^1 as well, up to tensoring a line bundle (Proposition 4.9). This enables us to find a wall \tilde{B} which realizes the small contraction $X \rightarrow X_B$ for any given flop $X \rightarrow X_B \leftarrow X'$ after crossing walls of Type 0. Using the technical lemma (Lemma 4.3), we can confirm that crossing the wall \tilde{B} gives the flop $X \dashrightarrow X'$. Since any two projective crepant resolutions are connected by a sequence of flops, this proves the theorem.

As an application of the construction of the equivariant map $\mathfrak{X} \rightarrow \mathcal{W}$, we will show that crepant resolutions of a certain four-dimensional quotient singularity \mathbb{C}^4/T can be realized as moduli spaces of T -constellations. More precisely, the group $T \subset Sp_4(\mathbb{C})$ is called the binary tetrahedral group, and \mathbb{C}^4/T is one of symplectic quotient singularities having symplectic (or equivalently, crepant) resolutions. The existence of the resolutions was shown by the results of Bellamy and Namikawa [B], [N], and later the resolutions were explicitly constructed by Lehn and Sorger [LS]. Although the abstract existence of an moduli interpretation of the resolutions is already known by the result of Bezrukavnikov and Kaledin [BK], we give an explicit description as moduli spaces with explicit stability conditions and also show that a component of the T -Hilbert scheme is a common blow-up of the two resolutions (Theorem 5.1). The fact that the two resolutions are dominated by a component of the T -Hilbert scheme was already observed in [LS]. In the proof of Theorem 5.1, we will describe how the parametrized T -constellations change under the birational contraction of the T -Hilbert scheme.

This paper is organized as follows. In Section 2 we review the construction of a moduli space of G -constellations and semi-invariants of quiver representations. In Section 3, we introduce the Cox ring of a crepant resolution and relate it with the semi-invariants for representations of the McKay quiver of G so that we obtain a nice embedding of a moduli space into a toric variety (Proposition 3.5). We also give concrete examples in Subsection 3.5. In Section 4 we give a proof of the main result (Theorem 4.1). In Section 5 we treat the symplectic quotient singularity \mathbb{C}^4/T for the binary tetrahedral group T . In particular we show that their symplectic resolutions are obtained as moduli spaces of T -constellations and that a certain component of the T -Hilbert scheme is isomorphic to a common blowing-up of these resolutions.

Conventions & Notations

In this paper all schemes are algebraic ones defined over \mathbb{C} . By a *point* of a scheme, we always mean a closed point. For an abelian group A , we denote by $A_{\mathbb{R}}$ its scalar extension $A \otimes_{\mathbb{Z}} \mathbb{R}$. Similarly, for an homomorphism f of abelian groups, $f_{\mathbb{R}}$ denotes the scalar extension. For a finite dimensional \mathbb{R} -vector space V and a subset $S \subset V$, we denote by $\text{Cone}(S)$ the cone generated by S .

Acknowledgements

The author is grateful to Alastair Craw and Akira Ishii for their useful comments to the draft of the present paper. This work was supported by the grant (NTU-110VV006) of the Taiwan Ministry of Education, by World Premier International Research Center Initiative (WPI), MEXT, Japan, and by JSPS KAKENHI Grant Number JP19K14504.

2 Moduli spaces of G -constellations

2.1 GIT-construction

In this section we review the construction of a moduli space of G -constellations for an arbitrary finite subgroup $G \subset SL(V)$ with $V = \mathbb{C}^n$ (cf. [CI, Section 2]). See e.g. [CMT]

and [Y2] for the special case where G is abelian.

We first recall that a G -constellation F is a G -equivariant $\mathbb{C}[V]$ -module which is isomorphic to the regular representation $R = \bigoplus_{g \in G} \mathbb{C}g$ as a $\mathbb{C}G$ -module where the G -action on $\mathbb{C}[V]$ comes from the inclusion $G \subset SL(V)$. Although G -constellations are defined as G -equivariant coherent sheaves on V in [CI], we identify them with their global sections since V is affine.

Let $\text{Irr}(G)$ be the set of isomorphism classes of irreducible representations of G . As is well known, R is isomorphic to $\bigoplus_{\rho \in \text{Irr}(G)} \rho^{\oplus \dim \rho}$ as a $\mathbb{C}G$ -module. We will construct a moduli space of G -constellations by taking a quotient of the affine scheme

$$\mathcal{N} = \{B \in \text{Hom}_{\mathbb{C}G}(V^* \otimes_{\mathbb{C}} R, R) \mid B \wedge B = 0\} \quad (2.1)$$

where V^* is the dual representation of V and $B \wedge B \in \text{Hom}_{\mathbb{C}[G]}(V^* \otimes_{\mathbb{C}} V^* \otimes_{\mathbb{C}} R, R)$ is defined by

$$(B \wedge B)(x \otimes y \otimes a) = B(x \otimes B(y \otimes a)) - B(y \otimes B(x \otimes a)).$$

We may regard \mathcal{N} as a space parametrizing all G -constellations, and two elements \mathcal{N} are isomorphic as G -equivariant $\mathbb{C}[V]$ -modules if and only if they lie in the same GL_R -orbit where $GL_R := \text{Aut}_{\mathbb{C}G}(R)$ is the G -equivariant automorphism group of R , which acts on \mathcal{N} by conjugation. By Shur's lemma, GL_R is written as a direct product $\prod_{\rho \in \text{Irr}(G)} GL_{\dim \rho}(\mathbb{C})$. Note that the diagonal subgroup \mathbb{C}^* acts trivially on \mathcal{N} and thus the action of GL_R descends to that of $PGL_R := GL_R/\mathbb{C}^*$.

Since the orbit space \mathcal{N}/PGL_R does not admit a reasonable structure as a scheme, we apply the construction of geometric invariant theory (GIT). Concretely, we consider a PGL_R -linearization of the trivial line bundle of \mathcal{N} and take the GIT-quotient with respect to it. Such a linearization is given by a character of PGL_R , and the group $\chi(PGL_R)$ of characters can be identified with the space of *stability conditions*

$$\bar{\Theta} := \{\theta \in \text{Hom}_{\mathbb{Z}}(R(G), \mathbb{Z}) \mid \theta(R) = 0\}$$

where $R(G) = \bigoplus_{\rho \in \text{Irr}(G)} \mathbb{Z}\rho$ is the representation ring of G . Note that $\bar{\Theta}$ has rank $s - 1$ with s being the number of conjugacy classes in G . The identification is given by

$$\begin{aligned} \chi(PGL_R) & \rightarrow \bar{\Theta} \\ \chi = (\bar{g}_\rho \mapsto \det(g_\rho)^{\theta_\rho}, \forall \rho \in \text{Irr}(G)) & \mapsto \theta_\chi = (\rho \mapsto \theta_\rho, \forall \rho \in \text{Irr}(G)). \end{aligned}$$

For a character $\chi \in \chi(PGL_R)$ we consider the χ -semistable locus

$$\mathcal{N}^{\chi\text{-ss}} = \{p \in \mathcal{N} \mid f(p) \neq 0 \text{ for some } f \in A_{\chi^k} \text{ and } k \in \mathbb{N}\}$$

where A_χ is the \mathbb{C} -vector space consisting of regular functions f on \mathcal{N} such that $g \cdot f = \chi(g)f$ for any $g \in PGL_R$. As a smaller open subset of $\mathcal{N}^{\chi\text{-ss}}$, the χ -stable locus $\mathcal{N}^{\chi\text{-s}}$ is defined as the set of points $p \in \mathcal{N}^{\chi\text{-ss}}$ such that the orbit $PGL_R \cdot p$ is closed in $\mathcal{N}^{\chi\text{-ss}}$ and that the stabilizer subgroup of p in PGL_R is finite. We say that a character χ is *generic* if $\mathcal{N}^{\chi\text{-ss}} = \mathcal{N}^{\chi\text{-s}}$.

For any character χ , we have a morphism

$$\pi_\chi : \mathcal{N}^{\chi\text{-ss}} \rightarrow \mathcal{N} //_\chi PGL_R := \text{Proj} \left(\bigoplus_{k=0}^{\infty} A_{\chi^k} \right)$$

which gives a one-to-one correspondence between the set of closed PGL_R -orbits in $\mathcal{N}^{\chi\text{-ss}}$ and the set of closed points in $\mathcal{N} //_\chi PGL_R$ (cf. [MFK]). We call $\mathcal{N} //_\chi PGL_R$ the *GIT-quotient* of \mathcal{N} by PGL_R with respect to χ . If χ is generic, then $\mathcal{N} //_\chi PGL_R$ is the orbit space of $\mathcal{N}^{\chi\text{-ss}}$, and in such a case $\mathcal{N} //_\chi PGL_R$ is called the *geometric quotient*.

The stability of a point of \mathcal{N} can be rephrased in representation-theoretic terms by the result of King:

Proposition 2.1. [Kin, Proposition 3.1] *For a point $p \in \mathcal{N}$ and a character $\chi \in \chi(PGL_R)$, p is in $\mathcal{N}^{\chi\text{-ss}}$ (resp. $\mathcal{N}^{\chi\text{-s}}$) if and only if the corresponding G -constellation F_p is θ_χ -semistable (resp. θ_χ -stable), that is, $\theta_\chi(M) \geq 0$ (resp. > 0) for any nonzero G -equivariant $\mathbb{C}[V]$ -submodule $M \subsetneq F_p$.*

We will use the scalar extension $\Theta := \bar{\Theta} \otimes_{\mathbb{Z}} \mathbb{R}$ as the space of stability conditions rather than $\bar{\Theta}$ itself in order to consider polyhedral cones inside Θ (or $\chi(PGL_R)_{\mathbb{R}}$). For $\theta \in \Theta$ and the corresponding character χ_θ , we denote $\mathcal{N} //_{\chi_\theta} PGL_R$ also by \mathcal{M}_θ . From King's result above, we see that Θ is divided into finitely many open chambers, on which the semistable locus is constant. We call such a chamber a *GIT-chamber* of Θ . Since G -constellations are regular representations as $\mathbb{C}G$ -modules, supporting hyperplanes of the GIT-chambers are of the form $\sum_{\rho \in \text{Irr}(G)} a_\rho \theta_\rho = 0$ with $a_\rho \in \{0, 1, \dots, \dim \rho\}$.

Typical examples of G -constellations are given by the coordinate rings of free G -orbits, that is, orbits $G \cdot x$ such that $\sharp(G \cdot x) = \sharp G$ in \mathbb{C}^n . Such G -constellations are simple $\mathbb{C}[V]$ -modules and hence stable for any $\theta \in \Theta$. As a natural generalization of free G -orbits, G -constellations obtained as quotients of $\mathbb{C}[V]$ by its ideal is called *G -clusters*. Since G -clusters are characterized as the G -constellations generated by the trivial representation, the moduli space of G -clusters, called the *G -Hilbert scheme*, is realized as \mathcal{M}_{θ_+} for $\theta_+ \in \Theta$ satisfying $\theta_+(\rho) > 0$ for any nontrivial $\rho \in \text{Irr}(G)$.

By [CI, Proposition 2.2], there exists an irreducible component of \mathcal{M}_0 which is isomorphic to \mathbb{C}^n/G such that its general points parametrize the coordinate rings of free G -orbits in \mathbb{C}^n . Accordingly, there is an irreducible component $\mathcal{V} \subset \mathcal{N}$ such that the restriction of the quotient map $\mathcal{N} \rightarrow \mathcal{M}_0$ to \mathcal{V} factors into

- the morphism $\mathcal{V} \rightarrow \mathbb{C}^n/G$ which sends a G -constellation to its support, and
- a closed immersion $\mathbb{C}^n/G \rightarrow \mathcal{M}_0$ onto an irreducible component of \mathcal{M}_0 .

In Subsection 3.3 we will give a more explicit description of \mathcal{V} (cf. Proposition 3.2). Note that the morphism $\mathfrak{M}_\theta := \mathcal{V} //_{\chi_\theta} PGL_R \rightarrow \mathbb{C}^n/G$ induced by the inclusion $\mathcal{V} \cap \mathcal{N}^{\chi_\theta\text{-ss}} \hookrightarrow \mathcal{V}$ is a birational morphism over \mathbb{C}^n/G for any $\theta \in \Theta$. The irreducible component \mathfrak{M}_θ of \mathcal{M}_θ is sometimes called the *coherent component*, and we will adopt this terminology in this article. For a GIT-chamber $C \subset \Theta$, the moduli space \mathfrak{M}_θ is constant for $\theta \in C$ and thus we also denote it by \mathfrak{M}_C .

The main result of [BKR] shows that the G -Hilbert scheme for $G \subset SL_3(\mathbb{C})$ is a crepant resolution of \mathbb{C}^3/G . More generally, the same proof of this result shows that the moduli space \mathcal{M}_θ of θ -semistable G -constellations for any generic $\theta \in \Theta$ is a crepant resolution of \mathbb{C}^3/G . In particular \mathcal{M}_θ is irreducible and hence $\mathcal{M}_\theta = \mathfrak{M}_\theta$ in this case.

As explained in [CI, §2.1], the moduli space \mathfrak{M}_C for a GIT-chamber C admits a universal family \mathcal{U}_C on $\mathfrak{M}_C \times \mathbb{C}^n$, and the associated tautological bundle $(p_{\mathfrak{M}_C})_*\mathcal{U}_C$ on \mathfrak{M}_C admits a decomposition

$$(p_{\mathfrak{M}_C})_*\mathcal{U}_C = \bigoplus_{\rho \in \text{Irr}(G)} \mathcal{R}_\rho \otimes \rho$$

into isotypic components where $p_{\mathfrak{M}_C} : \mathfrak{M}_C \times \mathbb{C}^n \rightarrow \mathfrak{M}_C$ is the projection. Note that each \mathcal{R}_ρ is a vector bundle on \mathfrak{M}_C of rank $\dim \rho$. Since tensoring a line bundle to $(p_{\mathfrak{M}_C})_*\mathcal{U}_C$ has no effect on the parametrized G -constellations, we assume that $\mathcal{R}_{\rho_0} \cong \mathcal{O}_{\mathfrak{M}_C}$ for the trivial representation ρ_0 whenever we talk about the universal family or the tautological bundle determined by C . Later we will identify G -constellations with representations of a quiver in Subsection 2.3. We will also use the associated tautological bundle \mathcal{R}_C on \mathfrak{M}_C as a moduli space of quiver representations rather than using the bundle $(p_{\mathfrak{M}_C})_*\mathcal{U}_C$.

For later use, we introduce the notion of an *orbit cone* for an affine scheme acted on by an algebraic group, following [BH, Definition 2.1]:

Definition 2.2. *Let H be an algebraic group acting on an affine algebraic scheme Z . For a point $z \in Z$, the **orbit cone** of z is the polyhedral cone $C_z \subset \chi(H)_{\mathbb{R}}$ generated by characters χ such that z is χ -semistable.*

Remark 2.1. The orbit cone of $z \in Z$ depends only on the H -orbit of z . If Z is the affine scheme \mathcal{N} acted on by $H = PGL_R$ as above, the orbit cone of a point of \mathcal{N} corresponding to a G -constellation F is identified with a cone in Θ consisting of stability conditions θ such that F is θ -semistable. Moreover, for a GIT-chamber $C \subset \Theta$, its closure \overline{C} is equal to the intersection of all the orbit cones for points of $\mathcal{N}^{C\text{-ss}}$.

2.2 Semi-invariants for quiver representations

Let us consider the subgroup $SL_R := \prod_{\rho \in \text{Irr}(G)} SL_{\dim \rho}(\mathbb{C})$ of GL_R . Since SL_R is reductive, there are finitely many generators of the invariant ring of the coordinate ring $\mathbb{C}[\mathcal{N}]$ of the affine scheme \mathcal{N} . Note that we have a decomposition of this ring into homogeneous components:

$$\mathbb{C}[\mathcal{N}]^{SL_R} = \bigoplus_{\chi \in \chi(PGL_R)} A_\chi.$$

As we will see in the next subsection, \mathcal{N} is identified with a space of quiver representations (with relations). In general the ring of semi-invariants for representations of a quiver (in characteristic zero) are generated by *determinantal semi-invariants* (see Theorem 2.2 below). In this subsection we explain how to obtain such semi-invariants.

Firstly, we briefly recall the concept of representations of a quiver. See e.g. [DeW2],[Kir] for more details. Let Q be a quiver, that is, an oriented graph consisting of a finite set I of vertices and a finite set A of arrows between two vertices. Here we allow Q to

have loops and oriented cycles. For any map $\alpha : I \rightarrow \mathbb{Z}_{\geq 0}$, we define a space $\text{Rep}(\alpha)$ of representations of Q with a dimension vector α as

$$\text{Rep}(\alpha) = \bigoplus_{a \in A} \text{Hom}_{\mathbb{C}}(\mathbb{C}^{\alpha(t_a)}, \mathbb{C}^{\alpha(h_a)})$$

where t_a and h_a denote the tail and the head of a respectively. For a representation $H = (H_a)_{a \in A} \in \text{Rep}(\alpha)$ and a path

$$p = a_1 \cdots a_k \quad (a_i \in A, h_{a_i} = t_{a_{i+1}}),$$

we set $H_p = H_{a_k} \circ \cdots \circ H_{a_1}$. For the trivial path e_v at a vertex v , we set $H_{e_v} = \text{id}_{\mathbb{C}^{\alpha(v)}}$. $\text{Rep}(\alpha)$ is acted on by $GL_{\alpha} := \prod_{v \in I} GL_{\alpha(v)}$ by conjugation.

Let $v_1, \dots, v_r, w_1, \dots, w_s$ be vertices of Q with possible repetition such that

$$\sum_{i=1}^r \alpha(v_i) = \sum_{j=1}^s \alpha(w_j),$$

and let $p_{i,j}$ be a \mathbb{C} -linear combination of paths from v_i to w_j . Then the function $f : \text{Rep}(\alpha) \rightarrow \mathbb{C}$ defined by

$$H \mapsto \det \begin{pmatrix} H_{p_{1,1}} & \cdots & H_{p_{1,s}} \\ \vdots & & \vdots \\ H_{p_{r,1}} & \cdots & H_{p_{r,s}} \end{pmatrix} \quad (2.2)$$

gives a semi-invariant satisfying

$$(g_v)_{v \in I} \cdot f = \left(\prod_{v \in I} \det g_v^{c_v} \right) f$$

for $(g_v)_{v \in I} \in GL_{\alpha}$ where

$$c_v = \alpha(v)(\#\{j \mid w_j = v\} - \#\{i \mid v_i = v\}).$$

A semi-invariant obtained in this way is called a *determinantal semi-invariant* of weight $(c_v)_{v \in I}$. Note that weights are identified with characters of GL_{α} .

Theorem 2.3. (*[DeW1],[DZ],[SV]*) *The ring $\mathbb{C}[\text{Rep}(\alpha)]^{SL_{\alpha}}$ of semi-invariants on $\text{Rep}(\alpha)$ with respect to the GL_{α} -action is generated by determinantal semi-invariants.*

This was first proved by Derksen and Weyman [DeW1] under the assumption that Q has no oriented cycles, and generalized later by Domokos and Zubkov [DZ] and by Schofield and van den Bergh [SV].

2.3 G -constellations as representations of the McKay quiver

In this subsection we realize the affine scheme \mathcal{N} in (2.1) as a space of quiver representations.

The *McKay quiver* Q_G of G is the quiver having $\text{Irr}(G)$ as the set of vertices and having $a_{\rho, \rho'} := \dim_{\mathbb{C}}(\text{Hom}_{\mathbb{C}G}(V^* \otimes_{\mathbb{C}} \rho, \rho'))$ arrows from ρ to ρ' for any $\rho, \rho' \in \text{Irr}(G)$. Decomposing R into irreducible summands, we see that $\text{Hom}_{\mathbb{C}G}(V^* \otimes_{\mathbb{C}} R, R)$ is isomorphic to the representation space $\text{Rep}(\alpha_G)$ of Q_G for the dimension vector $\alpha_G : \rho \mapsto \dim \rho$ (cf. [IN]). Thus, \mathcal{N} can be realized as a $GL_R (= GL_{\alpha_G})$ -invariant closed subscheme of $\text{Rep}(\alpha_G)$.

Via the identification of a G -constellation with a representation of Q_G , the moduli space \mathfrak{M}_C of G -constellations constructed in Subsection 2.1 is also regarded as a moduli space of representations of Q_G . Then the tautological bundle \mathcal{R}_C on \mathfrak{M}_C as the latter moduli space admits a decomposition $\mathcal{R}_C = \bigoplus_{\rho \in \text{Irr}(G)} \mathcal{R}_{\rho}$ into the isotypic components. Note that \mathcal{R}_C is equipped with a $\mathbb{C}[V]$ -action whose restriction to $\mathbb{C}[V]^G$ coincides with the $H^0(\mathcal{O}_{\mathfrak{M}_C})$ -action as a vector bundle on \mathfrak{M}_C . Hereafter, by *the tautological bundle on \mathfrak{M}_C* , we mean the bundle \mathcal{R}_C .

Example 2.1. (D_5 -singularities) We describe the space \mathcal{N} explicitly for the case of the D_5 -singularity. This singularity is given as a quotient of \mathbb{C}^2 by a binary dihedral group G of order 12:

$$G = \left\langle g_1 = \begin{pmatrix} \zeta & 0 \\ 0 & \zeta^{-1} \end{pmatrix}, g_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\rangle \subset SL_2(\mathbb{C})$$

where ζ is the sixth root of unity. $\text{Irr}(G)$ consists of four 1-dimensional representations $\rho_k : G \rightarrow \mathbb{C}^*$ ($k = 0, 1, 2, 3$) defined by

$$\rho_k(g_j) = \begin{cases} (-1)^k & \text{if } j = 1 \\ i^k & \text{if } j = 2 \end{cases},$$

and two 2-dimensional representations V_1, V_2 ; the first one is the inclusion $G \subset SL_2(\mathbb{C})$ presented above and the other is presented as

$$g_1 \mapsto \begin{pmatrix} \zeta^2 & 0 \\ 0 & \zeta^{-2} \end{pmatrix}, \quad g_2 \mapsto \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Let e_0, \dots, e_3 be a basis of ρ_0, \dots, ρ_3 respectively. Also, we let $\{v_{11}, v_{12}\}$ (resp. $\{v_{21}, v_{22}\}$) be a basis of V_1 (resp. V_2) such that it gives the matrix presentation above. We moreover prepare their copies $\{v'_{11}, v'_{12}\}$ and $\{v'_{21}, v'_{22}\}$ for the second direct summands of V_1 and V_2 in R respectively. When we emphasize that v_{11} and v_{12} are elements of V^* , we instead denote them by x and y respectively.

We fix isomorphisms of $\mathbb{C}G$ -modules as follows:

$$\begin{array}{cccc} V^* \otimes \rho_0 \cong V_1, & V^* \otimes \rho_1 \cong V_2, & V^* \otimes \rho_2 \cong V_1, & V^* \otimes \rho_3 \cong V_2 \\ x \otimes e_0 \mapsto v_{11} & x \otimes e_1 \mapsto v_{22} & x \otimes e_2 \mapsto v_{11} & x \otimes e_3 \mapsto v_{22} \\ y \otimes e_0 \mapsto v_{12} & y \otimes e_1 \mapsto iv_{21} & y \otimes e_2 \mapsto -v_{12} & y \otimes e_3 \mapsto -iv_{21} \\ \\ V^* \otimes V_1 \cong \rho_0 \oplus \rho_2 \oplus V_2, & V^* \otimes V_2 \cong \rho_1 \oplus \rho_3 \oplus V_1, \\ x \otimes v_{11} \mapsto & v_{21} & x \otimes v_{21} \mapsto & e_1 + e_3 \\ x \otimes v_{12} \mapsto & e_0 + e_2 & x \otimes v_{22} \mapsto & v_{12} \\ y \otimes v_{11} \mapsto & -e_0 + e_2 & y \otimes v_{21} \mapsto & -v_{11} \\ y \otimes v_{12} \mapsto & v_{22} & y \otimes v_{22} \mapsto & -ie_1 + ie_3. \end{array}$$

Thus the McKay quiver Q_G is presented as

$$\begin{array}{ccccccc}
& & \rho_0 & & \rho_1 & & \\
& & \uparrow & & \uparrow & & \\
& & A_0 & & A_1 & & \\
& & \downarrow & & \downarrow & & \\
& & B_0 & & B_1 & & \\
\rho_2 & \xleftarrow{B_2} & V_1 & \xleftarrow{D} & V_2 & \xleftarrow{A_3} & \rho_3 \\
& \xrightarrow{A_2} & & \xrightarrow{C} & & \xrightarrow{B_3} &
\end{array}$$

Using the isomorphisms above, matrices associated to the arrows in Q_G define an action of $\mathbb{C}[V]$ on the vector space

$$\mathbb{C}e_0 \oplus \cdots \oplus \mathbb{C}e_3 \oplus (\mathbb{C}v_{11} \oplus \mathbb{C}v_{12}) \oplus (\mathbb{C}v'_{11} \oplus \mathbb{C}v'_{12}) \oplus (\mathbb{C}v_{21} \oplus \mathbb{C}v_{22}) \oplus (\mathbb{C}v'_{21} \oplus \mathbb{C}v'_{22}) \cong R.$$

For instance, a matrix $\begin{pmatrix} a_1 \\ a_2 \end{pmatrix}$ ($a_i \in \mathbb{C}$) for A_1 lets V^* act on $\mathbb{C}e_1 = \rho_1$ as $x \cdot e_1 = a_1 v_{22} + a_2 v'_{22}$ and $y \cdot e_1 = a_1 i v_{21} + a_2 i v'_{21}$. Recall that the matrices must satisfy the condition $B \wedge B = 0$. By writing down the conditions $x \cdot (y \cdot v) - y \cdot (x \cdot v) = 0$ for each basis v of R , one sees that this condition is expressed as the following matrix relations:

$$B_0 A_0 = B_1 A_1 = B_2 A_2 = B_3 A_3 = A_0 B_0 - A_2 B_2 - DC = A_1 B_1 - A_3 B_3 - iCD = 0. \quad (2.3)$$

□

Remark 2.2. (1) Usually the quiver Q_G itself can be obtained easily from the character table of G , without giving explicit G -equivariant homomorphisms $V^* \otimes \rho \rightarrow \rho'$ as above. However, this computation will be necessary in order to construct embeddings of moduli spaces into toric varieties as we will see in the next section.

(2) The existence of an ideal I_G of the path algebra $\mathbb{C}Q_G$ such that

$$\mathcal{N} \cong \{H \in \text{Rep}(\alpha_G) \mid H_p = 0, p \in I_G\}$$

as in the above example is regarded as a consequence of the Morita equivalence between the smash product $G \rtimes \mathbb{C}[V]$ and a quotient of $\mathbb{C}Q_G$. For another method to calculate I_G , see [BTW, §3]. □

3 Embedding into toric varieties

In this section we explain how we embed moduli space \mathfrak{M}_θ into toric varieties such that the birational geometry of the embedded varieties is inherited from that of the ambient toric varieties.

3.1 The Cox ring of a resolution of a quotient singularity

In this subsection we treat the Cox ring, denoted by $\text{Cox}(X)$, of a crepant resolution X of a quotient singularity \mathbb{C}^n/G for $G \subset SL_n(\mathbb{C})$. $\text{Cox}(X)$ is the direct sum $\bigoplus_{L \in \text{Pic}(X)} H^0(X, L)$ as a $H^0(\mathcal{O}_{V/G})$ -module and admits a multiplicative structure inherited from the tensor product of global sections of line bundles (see [ADHL] for generalities

regarding Cox rings). Then $\text{Cox}(X)$ is a $\text{Pic}(X)$ -graded commutative ring and is known to be finitely generated by the fundamental result of Birker-Cascini-Hacon-McKernan [BCHM]. The most important feature of it is that X can be recovered from $\text{Cox}(X)$ as a GIT-quotient of $\text{Spec Cox}(X)$ by an algebraic torus with respect to a generic stability parameter if $X \rightarrow \mathbb{C}^n/G$ is projective (see the next subsection). Note that the isomorphism class of $\text{Cox}(X)$ is independent of the choice of X since crepant resolutions are mutually isomorphic in codimension one. The structure of $\text{Cox}(X)$ has been studied for special classes of G in [D], [DW], [G] and so on. See [Y1] for a general treatment. Moreover, algorithms finding generators of $\text{Cox}(X)$ are given in [DK] and [Y1].

Now we review how to describe the structure of $\text{Cox}(X)$ explicitly. We refer the reader to [Y1] for details. We first recall the notion of *age*. For an element $g \in G$, we choose a basis x_1, \dots, x_n of the dual space V^* of $V = \mathbb{C}^n$ which are eigenvectors for the action of g . Then we can write $g \cdot x_j = e^{\frac{2\pi i a_j}{r}}$ with an integer $0 \leq a_j < r$ where r is the order of G . We define the *age* of G as $\frac{1}{r} \sum_{j=1}^n a_j$, which is an integer since $g \in SL_n(\mathbb{C})$. We call an element $g \in G$ *junior* if its age is one. Note that the age is invariant under conjugation. We also define a discrete valuation $\nu_g : \mathbb{C}(V)^* \rightarrow \mathbb{Z}$ by setting $\nu_g(x_j) = a_j$.

It is shown in [IR] that there is a one-to-one correspondence between the set of conjugacy classes of junior elements in G and the set of irreducible exceptional divisors of the resolution $\pi : X \rightarrow V/G$, such that the divisorial valuation $\nu_E : \mathbb{C}(X)^* \rightarrow \mathbb{Z}$ along the π -exceptional divisor E corresponding to g coincides with the restriction of $\frac{1}{r}\nu_g$ to $\mathbb{C}(X)$ (via the birational map π). Let E_1, \dots, E_m be the set of π -exceptional irreducible divisors and let g_k ($k = 1, \dots, m$) be a representative of the junior conjugacy class corresponding to E_k . We also put $\nu_k = \nu_{g_k}$ and $r_k = \#\langle g_k \rangle$.

The Cox ring has an embedding into the Laurent polynomial ring $\mathbb{C}[V]^{[G,G]}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ with m variables over the invariant ring for the commutator group of G [Y1, Lemma 4.2]. Note that $\mathbb{C}[V]^{[G,G]}$ is naturally graded by the dual abelian group $Ab(G)^\vee$ of the abelianization $Ab(G) = G/[G, G]$ so that its degree-zero part is equal to $\mathbb{C}[V]^G$. For $Ab(G)^\vee$ -homogeneous generators f_1, \dots, f_ℓ of $\mathbb{C}[V]^{[G,G]}$, we associate to them the following $\ell + m$ elements

$$f_1 \prod_{k=1}^m t_k^{\nu_k(f_1)}, \dots, f_\ell \prod_{k=1}^m t_k^{\nu_k(f_\ell)}, t_1^{-r_1}, \dots, t_m^{-r_m} \quad (3.1)$$

of $\mathbb{C}[V]^{[G,G]}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$, which lie in $\text{Cox}(X)$. They do not give generators of $\text{Cox}(X)$ in general. In fact they give a generating system if and only if f_i 's satisfy the *valuation lifting condition* cf. [DG, Theorem 2.2]. We do not explain this condition here as we will not use it later.

Remark 3.1. The ring $\mathbb{C}[V]^{[G,G]}$ treated here should be understood as the Cox ring $\text{Cox}(V/G)$ of V/G ([AG, Theorem 3.1]), and the $Ab(G)^\vee$ -grading also corresponds to the $\text{Cl}(V/G)$ -grading on $\text{Cox}(V/G)$. Thus each f_i above defines a Weil divisor D_i on V/G , and the associated element $\tilde{f}_i := f_i \prod_{k=1}^m t_k^{\nu_k(f_i)}$ to f_i corresponds to a section defining the strict transform in X of D_i under the birational morphism $X \rightarrow V/G$. \square

The $\text{Pic}(X)$ -grading of $\mathbb{C}[V]^{[G,G]}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ can be read off from the \mathbb{Z}^m -grading given by the multi-degrees for the variables t_1, \dots, t_m . More precisely, the $\text{Pic}(X)$ -grading

is given so that the generator t_k^{-rk} lies in the homogeneous component corresponding to the divisor class of E_k . Note that the divisor classes of E_1, \dots, E_m are a \mathbb{Q} -basis (but usually not a \mathbb{Z} -basis) of $\text{Pic}(X)$.

3.2 Review of birational geometry of X

As mentioned in the previous subsection, any crepant resolution X of V/G has a finitely generated Cox ring. Such a variety is called a *Mori dream space*, and its birational contractions are controlled by polyhedral cones in $\text{Pic}(X)_{\mathbb{R}}$ (cf. [HK]). In this section we review this briefly.

Hereafter we assume $X \rightarrow V/G$ is projective. We first consider the polyhedral cone generated by the classes of divisors in $\text{Pic}(X)_{\mathbb{R}}$ whose base loci are of codimension greater than one. This cone is called the *movable cone* and denoted by $\text{Mov}(X)$. In our case, the movable cone is the union of the nef cones of all relative minimal models $X' \rightarrow V/G$ (cf. [Y1]). Here, the nef cone $\text{Nef}(X') \subset \text{Pic}(X')_{\mathbb{R}}$ is regarded as a cone inside $\text{Pic}(X)_{\mathbb{R}}$ via the birational map $X \dashrightarrow X'$ which is an isomorphism in codimension one. Note that, in dimension three, relative minimal models are nothing but projective crepant resolutions.

The nef cones $\text{Nef}(X') \subset \text{Pic}(X)_{\mathbb{R}}$ are full-dimensional and have disjoint interiors. If we take any L in the interior $\text{Amp}(X')$ of $\text{Nef}(X')$, the complete linear system of a sufficiently divisible power of L gives the birational map $X \dashrightarrow X'$. Set

$$T_X = \text{Hom}(\text{Pic}(X), \mathbb{C}^*) \cong (\mathbb{C}^*)^{\times m},$$

which naturally acts on $\mathfrak{X} := \text{Spec Cox}(X)$. By [HK, Theorem 2.3], this birational map is the same as the one $X \dashrightarrow \mathfrak{X} //_{\chi} T_X$ between the GIT-quotients with the character $\chi \in \chi(T_X)$ corresponding to L via the identification $\chi(T_X)_{\mathbb{R}} = \text{Pic}(X)_{\mathbb{R}}$. Moreover, the similar results hold for any *rational contraction*, that is, a rational map $f : X \dashrightarrow Z$ which is the composite of a sequence of flops $X \dashrightarrow X'$ and a proper morphism $f' : X' \rightarrow Z$ over V/G such that

- Z is a normal variety projective over V/G , and
- f' has connected fibers, i.e. $f'_* \mathcal{O}_{X'} = \mathcal{O}_Z$.

More precisely, for such f , there exists $L \in \text{Mov}(X)$ whose linear system recovers f , and moreover f equals the rational map $X \dashrightarrow \mathfrak{X} //_{\chi} T_X$ induced by variation of GIT for the character χ corresponding to L . Note that general points of a facet (i.e. codimension-one face) of $\text{Mov}(X)$ corresponds to a primitive divisorial rational contraction (i.e., a composite of a sequence of flops and a birational morphism contracting a single divisor which drops the Picard number by one). In contrast, small contractions are realized by taking divisors (or the corresponding characters) from the interior of $\text{Mov}(X)$. In particular every flop $X \dashrightarrow X'$ corresponds to the common facet $\text{Nef}(X) \cap \text{Nef}(X')$.

Remark 3.2. (1) We can also consider divisors (or corresponding characters) outside the movable cone, but they again give rational contractions (cf. [HK, Proposition 1.11]) and thus considering the movable cone is essential.

(2) Classifying the GIT-quotients of \mathfrak{X} (or equivalently, rational contractions of X) gives a fan structure on $\chi(T_X)_{\mathbb{R}}$ (or on $\text{Pic}(X)_{\mathbb{R}}$). This fan is called the *GIT-fan*. When X is toric, this is also called the *secondary fan*. See [CLS, Ch. 14] for a more detailed exposition of this fan.

Example 3.1. (Abelian group of type $\frac{1}{10}(1, 3, 6)$)

We exhibit how to compute the movable cone and nef cones for a crepant resolution of \mathbb{C}^n/G for abelian G through a particular example. Here we treat the case when $G \subset SL_3(\mathbb{C})$ is the cyclic group generated by

$$g = \begin{pmatrix} \zeta_{10} & 0 & 0 \\ 0 & \zeta_{10}^3 & 0 \\ 0 & 0 & \zeta_{10}^6 \end{pmatrix}$$

with ζ_{10} the primitive 10th root of unity. In this case g, g^2, g^4, g^5, g^7 are the junior elements, and the Cox ring of any crepant resolution X is generated by the following 8 elements in $\mathbb{C}[x, y, z][t_1^{\pm 1}, \dots, t_5^{\pm 1}]$:

$$xt_1t_2t_3^2t_4t_5^7, yt_1^3t_2^3t_3t_4t_5, zt_1^6t_2t_3^2t_5^2, t_1^{-10}, t_2^{-5}, t_3^{-5}, t_4^{-2}, t_5^{-10} \quad (3.2)$$

where x, y and z are the standard coordinates of \mathbb{C}^3 . Moreover, X is a toric variety corresponding to a fan Σ obtained as a triangulation of the cone $\sigma_0 = \text{Cone}(e_1, e_2, e_3) \subset \mathbb{R}^3$ such that the rays generated by $e_1, e_2, e_3, v_1, \dots, v_5$ is equal to the set of the rays of Σ where e_j is the standard basis of \mathbb{R}^3 and v_k 's are given as

$$v_1 = \begin{pmatrix} 1 \\ 3 \\ 6 \end{pmatrix}, v_2 = \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix}, v_3 = \begin{pmatrix} 2 \\ 1 \\ 2 \end{pmatrix}, v_4 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, v_5 = \begin{pmatrix} 7 \\ 1 \\ 2 \end{pmatrix}$$

so that v_k corresponds to E_k . See e.g. [Y2, §3] for details. One can check that there are six crepant resolutions, which are in fact all projective. These resolutions X_1, \dots, X_6 fit into the following diagram of flops:

$$\begin{array}{c} X_3 \\ \uparrow \\ \vdots \\ \downarrow \\ X_2 \leftarrow \text{---} \rightarrow X_1 \leftarrow \text{---} \rightarrow X_4 \leftarrow \text{---} \rightarrow X_5 \leftarrow \text{---} \rightarrow X_6. \end{array}$$

The cross sections of the fans Σ_1 and Σ_6 of X_1 and X_6 respectively are given in Figure 1. Among X_i 's, the G -Hilbert scheme corresponds to X_1 (cf. [CR]).

As is well known, each i -dimensional cone $\tau \in \Sigma_1$ represents a torus orbit O_τ in X_1 of codimension i (see e.g. [CLS, §3.2]). τ also defines a cone $C_\tau \subset \text{Pic}(X_1)_{\mathbb{R}}$ generated by line bundles having global sections which do not vanish on O_τ . Under the identification of $\text{Pic}(X_1)_{\mathbb{R}}$ with $\chi(T_X)_{\mathbb{R}}$, C_τ is the same as the orbit cone (see Definition 2.2) of some (and hence any) lift $\bar{x} \in \mathfrak{X}$ of some (and hence any) $x \in O_\tau$. C_τ is explicitly computed as the cone generated by the weights of the generators in (3.2) whose corresponding rays




Figure 1: The cross sections of the fans Σ_1 (left) and Σ_6 (right)

are not contained in τ . For example, C_τ for $\tau = \text{Cone}(e_3, v_1) \in \Sigma_1$ is generated by the weights of the generators in (3.2) with the third and fourth ones removed.

The movable cone is obtained as the intersection $\bigcap_{i=1}^5 C_{\tau_i}$ where $\tau_i = \mathbb{R}_{\geq 0} v_i$. Similarly, the nef cone of X_1 is the intersection $\bigcap_{\tau \in \Sigma_1} C_\tau$. In our case, however, we do not have to compute C_τ for all $\tau \in \Sigma_1$. $\text{Nef}(X_1)$ is given as the intersection of $\text{Mov}(X)$ and just one cone $\text{Cone}(v_2, v_4, v_5)$ since Σ_1 is clearly the only triangulation containing this three-dimensional cone.

The above computations of cones can be done, for instance by using the package “Polyhedra” of Macaulay2 [GS]. Then we see that $\text{Mov}(X_1)$ has 9 rays and 7 facets. The fact that $\text{Mov}(X_1)$ is non-simplicial comes from non-uniqueness of primitive rational contractions of exceptional divisors. For example, the divisor $E_3 \subset X_1$ is mapped to a curve by a primitive contraction, while it is mapped to a point by a primitive contraction inside X_6 . We also see that $\text{Nef}(X_1)$ is a simplicial cone having facets corresponding to 2 primitive divisorial contractions for E_1, E_3 and 3 flops along the curves $E_2 \cap E_4, E_2 \cap E_5$, and $E_4 \cap E_5$.

Later we will compare cones in $\text{Mov}(X_1)$ and orbit cones in Θ for moduli spaces of G -constellations (see Example 3.2). \square

3.3 Construction of embeddings

In this subsection we give a construction of nice embeddings of moduli spaces \mathfrak{M}_θ into toric varieties.

Let us consider the homomorphism $\iota_X^* : \mathbb{C}[y_1, \dots, y_{\ell+m}] \rightarrow \text{Cox}(X)$ sending y_i 's to homogeneous generators of $\text{Cox}(X)$ of the form in (3.1) so that we obtain an embedding ι_X of $\mathfrak{X} = \text{Spec}(\text{Cox}(X))$ into the affine space $\mathbb{A}^{\ell+m}$. We endow $\mathbb{C}[y_1, \dots, y_{\ell+m}]$ with a \mathbb{Z}^m -grading so that ι_X^* preserves the grading. Then we see that a character $\chi \in \chi(T_X)_{\mathbb{R}} = \text{Pic}(X)_{\mathbb{R}} \cong \mathbb{R}^m$ corresponding to an ample divisor on X satisfies $\mathfrak{X} //_{\chi} T_X = X$ by the GIT construction (see also the previous subsection). We can also consider the GIT-quotient $Y = \mathbb{A}^{\ell+m} //_{\chi} T_X$ for the same character, which is a normal toric variety such that $\text{Pic}(Y) \cong \chi(T_X) \cong \text{Pic}(X)$ [Co], and ι_X induces an embedding $\iota_X : X \hookrightarrow Y$

between the quotients.

Remark 3.3. If the generators of the Cox ring are minimal, then ι_X is a *neat embedding* in the sense of [Ro]. See [Ro, §2.3] for details. \square

This embedding has a nice property from the viewpoint of birational geometry. As explained in the previous subsection, every rational contraction $X \dashrightarrow X'$ is realized as the rational map $X \dashrightarrow \mathfrak{X} //_{\chi'} T_X$ for some χ' . The same character χ' gives a rational contraction $Y \dashrightarrow Y'$ as well, and $\iota_{\mathfrak{X}}$ again induces an embedding $X' \rightarrow Y'$. Note that GIT-chambers for $\mathbb{A}^{\ell+m}$ are finer than those for \mathfrak{X} in general, and in particular $Y \dashrightarrow Y'$ may not be an isomorphism even if $X \dashrightarrow X'$ is an isomorphism. In other words, the inclusion $\iota_X^*(\text{Nef}(Y)) \subset \text{Nef}(X)$ is strict in general. Note, however, that we have

$$\iota_X^*(\text{Mov}(Y)) = \text{Mov}(X)$$

since every exceptional divisor of $Y \rightarrow \mathbb{A}^{\ell+m} //_0 T_X$ restricts to an exceptional divisor of $X \rightarrow \mathbb{C}^n/G$ by the construction of Y .

Next we construct an embedding of a moduli space \mathfrak{M}_θ into a toric variety. Recall that \mathfrak{M}_θ is obtained as the GIT-quotient of the variety \mathcal{V} by the action of PGL_R with respect to the character $\chi_\theta \in \chi(PGL_R)$. Let $T_\Theta \subset PGL_R$ be the algebraic torus consisting of (the images of) the component-wise scalar matrices. Since the restriction map $\chi(PGL_R) \rightarrow \chi(T_\Theta)$ is an isomorphism, via this identification \mathfrak{M}_θ is isomorphic to $\mathcal{W} //_{\chi_\theta} T_\Theta$ where \mathcal{W} is the affine categorical quotient $\mathcal{V} // SL_R$. Note that the points of \mathcal{W} are in bijection with the closed SL_R -orbits in \mathcal{V} .

In the proof of [CI, Proposition 2.2], it is shown that $\mathcal{M}_0 = \mathcal{N} // PGL_R$ admits an irreducible component which is isomorphic to \mathbb{C}^n/G (see also Subsection 2.1) by constructing a morphism $V \rightarrow \mathcal{M}_0$ induced from a G -constellation family \mathcal{F} over $V = \mathbb{C}^n$. We will explicitly describe $V \rightarrow \mathcal{M}_0$ by lifting it to a morphism $V \rightarrow \mathcal{V}$. To do this, let

$$\mu : \mathbb{C}[V] \rightarrow \mathbb{C}[G] \otimes_{\mathbb{C}} \mathbb{C}[V]$$

be the ring homomorphism corresponding to the action of $G \subset SL(V)$. We let G act on $\mathbb{C}[G]$ as $(g \cdot f)(h) = f(g^{-1}h)$ for any $g, h \in G$ and $f \in \mathbb{C}[G]$. In terms of algebras, the $\mathbb{C}[V]$ -action on each fiber $\mathcal{F}_p := \mathbb{C}[G]$ ($p \in V$) of the family \mathcal{F} is given by the morphism

$$\mathbb{C}[V] \otimes \mathbb{C}[G] \rightarrow \mathbb{C}[G]; \quad x \otimes f \mapsto (\mu(x)f \bmod \mathfrak{m}_p)$$

where $\mathfrak{m}_p \subset \mathbb{C}[V]$ is the maximal ideal for p . This is indeed G -equivariant since we have

$$(g \cdot (x \cdot f))(h) = (x \cdot f)(g^{-1}h) = (\mu(x)f \bmod \mathfrak{m}_p)(g^{-1}h) = x(g^{-1}h \cdot p)f(g^{-1}h)$$

while

$$\begin{aligned} ((g \cdot x) \cdot (g \cdot f))(h) &= (\mu(g \cdot x)(g \cdot f) \bmod \mathfrak{m}_p)(h) = ((g \cdot x)(h \cdot p))((g \cdot f)(h)) \\ &= x(g^{-1}h \cdot p)f(g^{-1}h) \end{aligned}$$

for any $g, h \in G$, $x \in \mathbb{C}[V]$ and $f \in \mathcal{F}_p = \mathbb{C}[G]$.

For each $\rho \in \text{Irr}(G)$, we fix its \mathbb{C} -basis $v_1, \dots, v_{\dim \rho}$. The matrix presentation of the representation ρ with respect to this basis gives a matrix $(f_{i,j})_{i,j}$ with entries in $\mathbb{C}[G]$, that is, $g \cdot v_j = \sum_{i=1}^{\dim \rho} f_{i,j}(g)v_i$ for all $g \in G$. Let $v_1^*, \dots, v_{\dim \rho}^*$ be the dual basis to $v_1, \dots, v_{\dim \rho}$. Then the assignment $f_{i,j} \mapsto v_i^*$ gives G -equivariant isomorphisms from the vector subspaces $V_{\rho,j} := \bigoplus_{i=1}^{\dim \rho} \mathbb{C}f_{i,j} \subset \mathbb{C}[G]$ to the dual (or contragredient) representations ρ^* of ρ . This can be checked by comparing the coefficients of v_i 's in

$$g^{-1}h \cdot v_j = \sum_{i=1}^{\dim \rho} f_{i,j}(g^{-1}h)v_i = \sum_{i=1}^{\dim \rho} (g \cdot f_{i,j})(h)v_i$$

and

$$g^{-1} \cdot (h \cdot v_j) = g^{-1} \cdot \left(\sum_{i=1}^{\dim \rho} f_{i,j}(h)v_i \right) = \sum_{i=1}^{\dim \rho} f_{i,j}(h) \left(\sum_{k=1}^{\dim \rho} f_{k,i}(g^{-1})v_k \right)$$

with $g, h \in G$. Since $\mathbb{C}[G]$ is a direct sum of $\{V_{\rho,j}\}_{\rho,j}$ by the character theory for finite groups, we obtain an explicit G -equivariant isomorphism $\mathbb{C}[G] \rightarrow \bigoplus_{\rho \in \text{Irr}(G)} \rho^{\oplus \dim \rho}$. We will describe the family \mathcal{F} as quiver representations with the matrix presentation with respect to the basis of $\mathbb{C}[G]$ obtained in this way. To this end, we also fix a basis of the $a_{\rho,\rho'}$ -dimensional vector space $\text{Hom}_{\mathbb{C}G}(V^* \otimes_{\mathbb{C}} \rho, \rho')$ for each pair $\rho, \rho' \in \text{Irr}(G)$.

For each arrow $a \in A$ from ρ to ρ' in Q_G , we have an embedding $\rho' \rightarrow V^* \otimes \rho$. Regarding the fixed basis for ρ and ρ' , we obtain a $(\dim \rho' \times \dim \rho)$ -matrix \tilde{H}_a with entries in V^* . Then we obtain a set of matrices $\{H_a^p := (\tilde{H}_a \bmod \mathfrak{m}_p)\}_{a \in A}$, which can be regarded as a set of representations of Q_G parametrized by V (see Subsection 2.3).

Lemma 3.1. *The representation of Q_G corresponding to the G -constellation \mathcal{F}_p is isomorphic to $\{H_a^p\}_{a \in A}$ for any $p \in V$.*

Proof. We take any arrow $a \in A$ from ρ to ρ' and set $m = \dim \rho$ and $n = \dim \rho'$. Let $(f_{i,j})_{1 \leq i,j \leq m}$ (resp. $(g_{i,j})_{1 \leq i,j \leq n}$) be the matrix presentation of ρ^* (resp. ρ'^*) with respect to the fixed basis. We write the $n \times m$ -matrix \tilde{H}_a as $(h_{i,j})_{i,j}$ with $h_{i,j} \in V^*$. Regarded as a representation of Q_G , H_a^p defines an action of V^* on the vector space $V_{\rho^*,j} = \bigoplus_{i=1}^m \mathbb{C}f_{i,j} \subset \mathbb{C}[G]$, which is isomorphic to ρ as a $\mathbb{C}G$ -module, so that

$$\begin{aligned} \sum_{i=1}^m h_{1,i} \cdot f_{i,k} &= \sum_{j=1}^n h_{j,k}(p)g_{1,j} \\ \sum_{i=1}^m h_{2,i} \cdot f_{i,k} &= \sum_{j=1}^n h_{j,k}(p)g_{2,j} \\ &\vdots \\ \sum_{i=1}^m h_{n,i} \cdot f_{i,k} &= \sum_{j=1}^n h_{j,k}(p)g_{n,j} \end{aligned} \tag{3.3}$$

for each $k = 1, \dots, m$. We show that this action coincides with the one for \mathcal{F}_p . By the definition of the $\mathbb{C}[V]$ -module structure on $\mathcal{F}_p = \mathbb{C}[G]$, the LHS of the j -th row of (3.3), as an element of F_p , is the function

$$g \mapsto h_{j,1}(g \cdot p)f_{1,k}(g) + h_{j,2}(g \cdot p)f_{2,k}(g) + \dots + h_{j,m}(g \cdot p)f_{m,k}(g)$$

on G for each j . Note that the value of this function at the identity $1 \in G$ is $h_{j,k}(p)$ and this is also equal to the value of the RHS of the j -th row of (3.3) (as a function on G) at $1 \in G$. By the choice of \tilde{H}_a , the LHSs and the RHSs of (3.3) give G -equivariantly isomorphic basis of ρ' . Noticing that $f(g) = g^{-1} \cdot f(1)$ for any $f \in \mathbb{C}[G]$ and $g \in G$, this implies that the two functions attain the same values for all $g \in G$, and hence the claim. \square

We will give concrete examples of the representations $\{H_a^p\}_{a \in A}$ in Subsection 3.5.

The generic representation $\{\tilde{H}_a\}_a$ gives a morphism $V \rightarrow \mathcal{N}$, which factors through $V \rightarrow \mathcal{V}$ since V is mapped onto $V/G \subset \mathcal{M}_0$. Let f be a semi-invariant function on \mathcal{N} and hence on \mathcal{V} . Then the (scheme-theoretic) zero locus of f on \mathcal{V} descends to a Weil divisor D_f on V/G . Thus, by sending f to the section of the divisorial sheaf $\mathcal{O}_{V/G}(D_f)$ defines a ring homomorphism

$$\varphi : \mathbb{C}[\mathcal{N}]^{SL_R} \rightarrow \text{Cox}(V/G) \cong \mathbb{C}[V]^{[G,G]}$$

(cf. Remark 3.1). For a determinantal semi-invariant $f : H \mapsto \det(H_{p_{i,j}})_{i,j}$ defined by paths $p_{i,j}$ as in (2.3), $\varphi(f)$ is given as $\det \mathcal{H} \in \mathbb{C}[V]$ where $\mathcal{H} = (\tilde{H}_{p_{i,j}})_{i,j}$ is defined similarly to \tilde{H}_a . Note that if f has weight $(c_\rho)_{\rho \in \text{Irr}(G)}$, then $\varphi(f)$ is acted on by G by the character

$$\det \left(\bigoplus_{\rho \in \text{Irr}(G)} \rho^{\oplus c_\rho} \right) : G \rightarrow \mathbb{C}^*.$$

In particular this naturally gives a homomorphism $\bar{\Theta} \rightarrow \text{Ab}(G)^\vee$.

Let us consider the $\bar{\Theta}$ -graded algebra $\bigoplus_{\theta \in \bar{\Theta}} \mathbb{C}[V]^{[G,G]} t_\theta$ where t_θ is a formal variable indicating the graded component for $\theta \in \bar{\Theta}$. Using φ , we can give a description of the coordinate ring $\mathbb{C}[\mathcal{V}]^{SL_R}$ of $\mathcal{W} = \mathcal{V} // SL_R$ similarly to the description of $\text{Cox}(X)$ in (3.1).

Proposition 3.2. *The algebra $\mathbb{C}[\mathcal{W}] = \mathbb{C}[\mathcal{V}]^{SL_R}$ is isomorphic to the graded subring $\mathbf{S}_{\mathcal{W}}$ of $\bigoplus_{\theta \in \bar{\Theta}} \mathbb{C}[V]^{[G,G]} t_\theta$ generated by $\varphi(h)t_\theta$ for semi-invariants $h \in \mathbb{C}[\mathcal{N}]^{SL_R}$ of weight θ .*

Proof. The T_Θ -equivariant morphism

$$V/[G, G] \times T_\Theta \rightarrow \mathcal{N} // SL_R$$

corresponding to the homomorphism $\mathbb{C}[\mathcal{N}]^{SL_R} \rightarrow \bigoplus_{\theta \in \bar{\Theta}} \mathbb{C}[V]^{[G,G]} t_\theta$ is induced from the morphism $V \rightarrow \mathcal{V} \subset \mathcal{N}$ (and thus descends to the closed immersion $V/G \rightarrow \mathcal{M}_0$). Then the claim follows since the spectrum of the integral domain $\mathbf{S}_{\mathcal{W}}$ is isomorphic to the scheme-theoretic image of the equivariant morphism above, which is equal to $\mathcal{V} // SL_R = \mathcal{W}$. \square

From now on we assume $n = 3$, i.e. $V = \mathbb{C}^3$. We fix any GIT-chamber $C \subset \Theta$ and let $X := \mathfrak{M}_C$ be the crepant resolution of V/G . Similar to the construction of φ , we define a ring homomorphism

$$\varphi_C : \mathbb{C}[\mathcal{N}]^{SL_R} \rightarrow \text{Cox}(X)$$

whose composition with

$$\text{Cox}(X) \hookrightarrow \mathbb{C}[V]^{[G,G]}[t_1^{\pm 1}, \dots, t_m^{\pm 1}] \xrightarrow{t_k \mapsto 1} \mathbb{C}[V]^{[G,G]}$$

is equal to φ . This also induces a homomorphism $\psi_C : \bar{\Theta} \rightarrow \text{Pic}(X)$ such that the degree of $\varphi_C(f)$ for a homogeneous element of weight $\theta : \rho \mapsto c_\rho$ is equal to $\psi_C(\theta)$.

Now we are ready to define an embedding of \mathcal{W} into a toric variety. We first choose $Ab(G)^\vee$ -homogeneous generators $f_1, \dots, f_\ell \in \mathbb{C}[V]^{[G,G]}$ such that the associated elements in (3.1) generate $\text{Cox}(X)$. Let $\iota_{\mathfrak{X}} : \mathfrak{X} \rightarrow \mathbb{A}^{\ell+m}$ be the associated embedding constructed in the beginning of this subsection. Set $N_X = \mathbb{Z}^\ell \oplus \text{Pic}(X)^\vee$ and let $\sigma_{\mathbb{A}} \subset (N_X)_{\mathbb{R}} \cong \mathbb{R}^{\ell+m}$ be the cone defining the affine toric variety $\mathbb{A}^{\ell+m}$ with the T_X -action. We also set $N_{\mathcal{W}} = \mathbb{Z}^\ell \oplus \bar{\Theta}^\vee$ and define $\sigma_{\mathcal{W}} \subset (N_{\mathcal{W}})_{\mathbb{R}} \cong \mathbb{R}^{\ell+s-1}$ as the cone generated by the set of cones $\{(\text{id}_{\mathbb{Z}^\ell} \oplus \psi_C^*)_{\mathbb{R}}(\sigma_{\mathbb{A}})\}_X$ where $X = \mathfrak{M}_C$ runs for all GIT-chambers $C \subset \bar{\Theta}$ and the choice of f_i 's is consistent regardless of the choice of X . Note that ψ_C and its pullback $\psi_C^* : \text{Pic}(X)^\vee \rightarrow \bar{\Theta}^\vee$ depend on X .

Let $\bar{Y}_{\mathcal{W}}$ be the affine normal toric variety defined by $\sigma_{\mathcal{W}}$. We show that the normalization $\bar{\mathcal{W}}$ of \mathcal{W} has an embedding into $\bar{Y}_{\mathcal{W}}$. By definition, the coordinate ring of $\bar{Y}_{\mathcal{W}}$ is the semigroup algebra $\mathbb{C}[S_{\mathcal{W}}]$ for the semigroup $S_{\mathcal{W}} = M_{\mathcal{W}} \cap \sigma_{\mathcal{W}}^\vee$ where $M_{\mathcal{W}} = N_{\mathcal{W}}^\vee$ and $\sigma_{\mathcal{W}}^\vee$ is the dual cone

$$\{f \in (M_{\mathcal{W}})_{\mathbb{R}} \mid f(v) \geq 0 \text{ for all } v \in \sigma_{\mathcal{W}}\}$$

to $\sigma_{\mathcal{W}}$. Since f_i 's generate $\mathbb{C}[V]^{[G,G]}$, we can choose homogeneous generators $\{h_j\}_j$ of $\mathbb{C}[\mathcal{W}] = \mathbb{C}[\mathcal{V}]^{S_{LR}}$ such that each $\varphi(h_j)$ is of the form $f_1^{a_1} \cdots f_\ell^{a_\ell}$. This presentation may not be unique in general, but we fix it for each $\varphi(h_j)$. Then \mathcal{W} admits a closed immersion into the spectrum $Y_{\mathcal{W}}$ of the semigroup algebra $\mathbb{C}[S'_{\mathcal{W}}]$ for the semigroup $S'_{\mathcal{W}} \subset M_{\mathcal{W}}$ generated by $\{(a_1, \dots, a_\ell, \theta_j)\}_j$ where $\theta_j \in \bar{\Theta}$ is the weight of h_j . This also induces a closed immersion of $\bar{\mathcal{W}}$ into the spectrum of the integral closure of $\mathbb{C}[S'_{\mathcal{W}}]$.

Lemma 3.3. *The semigroup algebra $\mathbb{C}[S_{\mathcal{W}}]$ coincides with the integral closure of $\mathbb{C}[S'_{\mathcal{W}}]$.*

Proof. For each $f \in S_{\mathcal{W}}$, it defines a rational function \bar{f} on $\bar{\mathcal{W}}$ via the above embedding of $\bar{\mathcal{W}}$. It suffices to show that \bar{f} is regular on $\bar{\mathcal{W}}$. Since $\bar{\mathcal{W}}$ is normal it suffices to show that \bar{f} is regular outside a locus of codimension greater than one. By the construction of $\sigma_{\mathcal{W}}$, \bar{f} is regular at general points of any divisor $D \subset \bar{\mathcal{W}}$ which descends to a divisor on a resolution $\bar{\mathcal{W}}//_{\theta} T_{\bar{\Theta}}$ of V/G for some generic stability condition θ . Then the claim follows by the lemma below. \square

Lemma 3.4. *The complement of the open subset $\bigcup_{\theta:\text{generic}} \mathcal{W}^{\theta\text{-ss}}$ in \mathcal{W} is of codimension at least two.*

Proof. Assume that the complement contains an irreducible divisor $D \subset \mathcal{W}$ in order to deduce a contradiction. Let $C \subset \bar{\Theta}$ be the cone generated by the weights of semi-invariants of \mathcal{V} which do not vanish at general points of D . By the assumption, C has positive codimension and, equivalently, general points of D has a positive-dimensional stabilizer subgroup in $T_{\bar{\Theta}}$. Then the image of D under the quotient map $\mathcal{W}^{\theta\text{-ss}} \rightarrow \mathcal{W}//_{\theta} T_{\bar{\Theta}}$

for θ in the relative interior of C has dimension greater than or equal to $n = 3$. This is contrary to the irreducibility of \mathcal{W} since the T_Θ -orbits of general points of \mathcal{W} , which correspond to free G -orbits (see Subsection (2.1)), are closed and hence $\mathcal{W} //_\theta T_\Theta$ has dimension three for any (non-generic) θ . \square

Thus, we obtain the following commutative diagram of equivariant maps:

$$\begin{array}{ccccc} \mathbb{A}^{n+m} & \longrightarrow & \bar{Y}_{\mathcal{W}} & \longrightarrow & Y_{\mathcal{W}} \\ \iota_X \uparrow & & \uparrow & & \uparrow \\ \mathfrak{X} & \longrightarrow & \bar{\mathcal{W}} & \longrightarrow & \mathcal{W} \end{array} \quad (3.4)$$

where vertical arrows are closed immersions. Note that $Y_{\mathcal{W}}$ is a possibly non-normal toric variety whose normalization is $\bar{Y}_{\mathcal{W}}$ by Lemma 3.3. For any $\theta \in C$, we also obtain the following diagram of maps between quotients induced from (3.4):

$$\begin{array}{ccccc} Y & \longrightarrow & \bar{Y}_{\mathcal{W}} //_{\chi_\theta} T_\Theta & \longrightarrow & Y_{\mathcal{W}} //_{\chi_\theta} T_\Theta \\ \iota_X \uparrow & & \uparrow & & \uparrow \\ X & \longrightarrow & \bar{\mathcal{W}} //_{\chi_\theta} T_\Theta & \longrightarrow & \mathcal{W} //_{\chi_\theta} T_\Theta \end{array} \quad (3.5)$$

where vertical arrows are again closed immersions and horizontal arrows are all isomorphisms. The results obtained so far are summarized as follows:

Proposition 3.5. *The irreducible component \mathcal{W} of $\mathcal{N} // SL_R$ such that $\mathcal{W} // T_\Theta \cong \mathbb{C}^3 / G$ admits a T_Θ -equivariant closed immersion $\iota_{\mathcal{W}} : \mathcal{W} \rightarrow Y_{\mathcal{W}}$ into a possibly non-normal toric variety whose torus contains T_Θ , satisfying the following property:*

- for any generic $\theta \in \Theta$, the induced morphism $\mathcal{W} //_{\chi_\theta} T_\Theta \rightarrow Y_{\mathcal{W}} //_{\chi_\theta} T_\Theta$ between the GIT-quotients is identified with a closed immersion $\iota_X : X \rightarrow Y$ of the resolution $X := \mathfrak{M}_\theta$ into a normal toric variety such that the restriction map $\iota_X^* : \text{Pic}(Y)_{\mathbb{R}} \rightarrow \text{Pic}(X)_{\mathbb{R}}$ is an isomorphism and $\text{Mov}(X) = \iota_X^*(\text{Mov}(Y))$.

In the next subsection we will explain how the (semi)stable locus of $\bar{Y}_{\mathcal{W}}$ for a given stability condition is determined in terms of the fan $\sigma_{\mathcal{W}}$.

Remark 3.4. (1) Let $v_1, \dots, v_m \in \sigma_{\mathcal{W}}$ be the primitive generators of the rays corresponding to the exceptional divisors E_k of $X = \mathfrak{M}_C \rightarrow V/G$. Recall that the order of zeros (or poles) of a rational function $f \in M_{\mathcal{W}}$ along the corresponding divisor to v_k is given by $f(v_k)$, by toric geometry. Thus, the image of a homogeneous semi-invariant $h := f_1^{a_1} \cdots f_\ell^{a_\ell} t_\theta$ ($a_i \in \mathbb{Z}$) under φ_C is given by

$$\tilde{f}_1^{a_1} \cdots \tilde{f}_\ell^{a_\ell} \prod_{k=1}^m t_k^{-\mathbf{a}(v_k)r_k} \quad (3.6)$$

(see Remark 3.1) where $\mathbf{a} = (a_1, \dots, a_\ell, \theta) \in \mathbb{Z}^\ell \oplus \bar{\Theta}$ is identified with an element of $M_{\mathcal{W}}$. In particular, $\psi_C(\theta) \in \text{Pic}(X)$ is given by the multi-degree of t_k 's in (3.6). Note that ψ_C only depends on v_k 's and therefore $\psi_{C'}$ is equal to ψ_C as long as we take a GIT-chamber

C' inside the intersection of the orbit cones of G -constellations corresponding to general points of the divisors E_1, \dots, E_m .

(2) The homomorphism $\varphi_C : \mathbb{C}[\mathcal{N}]^{SL_R} \rightarrow \text{Cox}(X)$ is never surjective unless G is trivial since each $t_k^{-r_k} \in \text{Cox}(X)$ is clearly outside of the image. However, we will show that, for homogeneous generators $\{h_i\}$ of $\mathbb{C}[\mathcal{N}]^{SL_R}$, the associated elements to the images $\varphi(h_i)$ generate $\text{Cox}(X)$ (Proposition 4.10). \square

3.4 GIT-quotients of a toric variety by a subtorus-action

As we saw in the previous subsection, every crepant resolution of the form \mathfrak{M}_θ is embedded in a quotient of the toric variety $\bar{Y}_\mathcal{W}$ (or $Y_\mathcal{W}$) by the algebraic torus T_Θ . In this subsection we show how to compute the θ -semistable locus of $\bar{Y}_\mathcal{W}$ for $\theta \in \Theta$ in terms of the fan of $\bar{Y}_\mathcal{W}$. This also determines the semistable locus of $\bar{\mathcal{W}}$ since we have

$$\bar{Y}_\mathcal{W}^{\theta\text{-ss}} \cap \bar{\mathcal{W}} = \bar{\mathcal{W}}^{\theta\text{-ss}}$$

by the construction of $\bar{Y}_\mathcal{W}$. If θ is generic, $\bar{\mathcal{W}}^{\theta\text{-ss}}$ is also identified with $\mathcal{W}^{\theta\text{-ss}}$ since the geometric quotient of $\mathcal{W}^{\theta\text{-ss}}$ is smooth.

To consider quotients of $\bar{Y}_\mathcal{W}$, first note that T_Θ is regarded as a subtorus of the big torus $T_\mathcal{W} := \text{Hom}_{\mathbb{Z}}(N_\mathcal{W}, \mathbb{C}^*) \subset Y_\mathcal{W}$. Quotients of a toric variety by an action of a subtorus of the big torus have been studied in [KSZ],[AH],[Hu] and so on. When we consider GIT-quotients (or more generally good quotients), the quotient map naturally induces a one-to-one correspondence between the sets of the maximal cones for the toric variety and its quotient (see e.g. [AH, Proposition 3.2]). To state this more precisely for our situation, recall that the toric variety $\bar{Y}_\mathcal{W}$ is defined by the cone $\sigma_\mathcal{W}$ in $(N_\mathcal{W})_{\mathbb{R}}$. For any $\theta \in \Theta$, its semistable locus $\bar{Y}_\mathcal{W}^{\theta\text{-ss}}$ is $T_\mathcal{W}$ -invariant since functions on $\bar{\mathcal{W}}$ defining the semistable locus can be taken as homogeneous ones with respect to $T_\mathcal{W}$. Thus, $\bar{Y}_\mathcal{W}^{\theta\text{-ss}}$ is also a toric variety defined by a subfan $\bar{\Sigma}_\theta$ of $\Sigma_\mathcal{W}$ where $\Sigma_\mathcal{W}$ is the fan consisting of all faces of $\sigma_\mathcal{W}$. Then the GIT-quotient $Y_\theta := \bar{Y}_\mathcal{W} //_\theta T_\Theta$ is a toric variety defined by a fan $\bar{\Sigma}_\theta$, and the quotient map $\bar{Y}_\mathcal{W}^{\theta\text{-ss}} \rightarrow Y_\theta$ is a toric morphism induced by the projection

$$p : (N_\mathcal{W})_{\mathbb{R}} \rightarrow (N_\mathcal{W}/\bar{\Theta}^\vee)_{\mathbb{R}} = \mathbb{R}^\ell$$

satisfying the following condition (*):

- For any cone $\sigma \in \bar{\Sigma}_\theta$, the cone $p^{-1}(\sigma) \cap |\Sigma_\theta|$ belongs to Σ_θ where $|\Sigma_\theta| \subset \mathbb{R}^\ell$ is the support of the fan Σ_θ . Moreover, if θ is generic, p induces an isomorphism of Σ_θ and $\bar{\Sigma}_\theta$.

Now we give an explicit method to compute $\bar{\Sigma}_\theta$ and hence $\bar{Y}_\mathcal{W}^{\theta\text{-ss}}$. To this end, it is sufficient to determine a general point of $p^{-1}(v) \cap |\Sigma_\theta|$ for each $v \in \mathbb{R}^\ell$ by the condition (*). Let $q : (N_\mathcal{W})_{\mathbb{R}} \cong \mathbb{R}^\ell \times \Theta^\vee \rightarrow \Theta^\vee$ be the second projection to the dual vector space of $\Theta \cong \mathbb{R}^{s-1}$ (while p is regarded as the first projection). Then we set $P_v = q(p^{-1}(v) \cap |\Sigma_\mathcal{W}|) \subset \Theta^\vee$, which is a rational convex polyhedron since $\sigma_\mathcal{W}$ is rational and strongly convex. Note that θ is a linear function on P_v . The following result is regarded as a generalization of [CMT, Theorem 7.2] to the case where G is non-abelian, see Example 3.5.1.

Proposition 3.6. *For any $\theta \in \Theta$, a face F of $\sigma_{\mathcal{W}}$ belongs to Σ_{θ} if and only if $q(F) \cap P_{p(\tilde{v})}$ equals the set of points of $P_{p(\tilde{v})}$ for which the function $\theta : P_{p(\tilde{v})} \rightarrow \mathbb{R}$ is minimized for some (and hence any) point \tilde{v} of the relative interior $\text{relint}(F)$ of F .*

Although this result can be proved e.g. by using a similar argument to the the proof of [CMT, Theorem 7.2], we give another proof using the Hilbert-Mumford numerical criterion in order to understand the result in terms of one-parameter subgroups. The numerical criterion states that a point $x \in \bar{Y}_{\mathcal{W}}$ is θ -semistable if and only if we have $\lambda \cdot \theta \geq 0$ for any 1-parameter subgroup (1-PS) $\lambda : \mathbb{C}^* \rightarrow T_{\Theta}$ such that $\lim_{t \rightarrow 0} (\lambda(t) \cdot x)$ exists, where $\lambda \cdot \theta$ means the natural paring between a 1-PS and a character of T_{Θ} .

To apply this criterion to toric varieties, let us recall that faces F of $\sigma_{\mathcal{W}}$ and $T_{\mathcal{W}}$ -orbits of \mathcal{W} are in one-to-one correspondence which assigns to F the orbit $O(F)$ of $x_F := \lim_{t \rightarrow 0} \lambda^{\tilde{v}}(t)$ for the 1-PS $\lambda^{\tilde{v}} : \mathbb{C}^* \rightarrow T_{\mathcal{W}}$ corresponding to $\tilde{v} \in \text{relint}(F)$. Moreover, the closure of $O(F)$ is equal to $\bigcup_{F' \subset F \in \Sigma_{\mathcal{W}}} O(F')$, and we have $\lim_{t \rightarrow 0} (\lambda^u(t) \cdot x_F) \in O(F')$ for any $u \in \text{relint}(F')$ where λ^u is the 1-PS corresponding to u . See [CLS, Proposition 3.2.2] and its proof for these standard facts. We also observe that the limit $\lim_{t \rightarrow 0} (\lambda^u(t) \cdot x_F)$ for $u \in N_{\mathcal{W}}$ exists if and only if there exist (sufficiently small) $h > 0$ and a face F' of $\sigma_{\mathcal{W}}$ containing F such that $u_1 + hu$ lies in F' for some (and hence any) $u_1 \in \text{relint}(F)$.

Proof of Proposition 3.6.

According to the observation above, for a 1-PS λ^w of T_{Θ} corresponding to a vector $w \in \bar{\Theta}^{\vee}$ and any point x of the orbit $O(F)$, the limit $\lim_{t \rightarrow 0} (\lambda^w(t) \cdot x)$ exists if and only if there exist $h > 0$ and a face F' containing F such that $u_1 + hu \in F'$ for some $u_1 \in \text{relint}(F)$.

Let us assume that $q(F) \cap P_{p(\tilde{v})}$ does not minimize θ . Then there is another face F' of $\sigma_{\mathcal{W}}$ such that $q(F') \cap P_{p(\tilde{v})}$ minimizes θ , and $(w_1 - q(\tilde{v})) \cdot \theta < 0$ for any $w_1 \in \text{relint}(q(F') \cap P_{p(\tilde{v})})$. Since $q(\tilde{v}) + 1 \cdot (w_1 - q(\tilde{v})) = w_1$ is inside $P_{p(\tilde{v})}$, the convexity of $P_{p(\tilde{v})}$ implies that the limit for the 1-PS corresponding to w_1 exists and thus F is not in Σ_{θ} by the numerical criterion.

Conversely, if $q(F) \cap P_{p(\tilde{v})}$ minimizes θ , then $q(F) \cap P_{p(\tilde{v})}$ is a face of $P_{p(\tilde{v})}$ having $\{\theta = (\text{the minimal value})\} \subset \Theta$ as the supporting hyperplane. Therefore, $(w - q(\tilde{v})) \cdot \theta \geq 0$ for any $w \in P_{p(\tilde{v})}$ and F is in Σ_{θ} by the numerical criterion again. \square

Remark 3.5. Whenever we are given an affine toric variety and a subtorus of the big torus, the similar statement to the above proposition applies to compute the semistable locus of the given toric variety with respect to a character of the subtorus.

3.5 Examples

In this subsection we demonstrate the construction of the toric embedding and the homomorphism φ_C in the previous subsection via concrete examples.

3.5.1 Abelian cases

We first consider the case where $G \subset SL_3(\mathbb{C})$ is a finite abelian subgroup of order r . In this case all irreducible representations of G are 1-dimensional, which particularly implies

that SL_R is trivial and hence $\mathcal{V} = \mathcal{W}$. Then there is a natural choice of generators of the Cox ring of a crepant resolution X and generators of the coordinate ring of \mathcal{V} as follows.

Once we fix a basis of $V = \mathbb{C}^3$ such that G is a diagonal group, we obtain generators of $\mathbb{C}[V]^{[G,G]} = \mathbb{C}[V]$ as the standard basis x_1, x_2, x_3 of V^* and also standard generators of $\text{Cox}(X)$ (cf. [Y2, Proposition 3.5]). As explained in [CMT] (see also [Y2]), the coordinate ring of \mathcal{V} is generated by $3r$ variables $\{x_{i,j}\}_{0 \leq i \leq r-1, 1 \leq j \leq 3}$ with certain relations where r is the order of G . In terms of the quiver representation, the variable $x_{i,j}$ is a homogeneous semi-invariant corresponding to an arrow ρ_i to $\rho_i \otimes \chi_j$ of the McKay quiver of G where $\text{Irr}(G) = \{\rho_0, \dots, \rho_{r-1}\}$ and we regard the character $\chi_j : G \rightarrow \mathbb{C}^*$ defined by $g \cdot x_j = \chi_j(g)x_j$ as a 1-dimensional representation of G . Then \mathcal{V} is already a possibly non-normal toric variety [CMT, Theorem 3.10], and the ambient toric variety $Y_{\mathcal{V}}$ constructed from these generators is nothing but \mathcal{V} itself.

For each orbit $O \subset \mathcal{V}$ by the big torus, it has the *distinguished point* $v \in O$ satisfying $x_{i,j}(v) = 1$ for all i, j such that $x_{i,j}(O) \neq \{0\}$. For any θ , the fan Σ_{θ} of the θ -semistable locus of \mathcal{V} is computed using Proposition 3.6, and one obtains the distinguished point corresponding to a cone $\sigma \in \Sigma_{\theta}$ by setting

$$x_{i,j}(v) = \begin{cases} 1 & \text{if } \sigma \subset H_{i,j} \\ 0 & \text{if } \sigma \not\subset H_{i,j} \end{cases}$$

where $H_{i,j} \subset (N_{\mathcal{V}})_{\mathbb{R}}$ is the hyperplane defined by $x_{i,j} \in M_{\mathcal{V}}$. This is essentially the same procedure as the one given in [CMT, Theorem 7.2], which enables us to obtain the distinguished point corresponding to the cone $\sigma_w \in \Sigma_{\theta}$ such that $p(\sigma_w)$ contains a given $w \in \mathbb{R}^3$ in its relative interior. Note that one can also consider distinguished points of $\bar{Y}_{\mathcal{W}}$ for non-abelian G but these points usually do not lie inside $\bar{\mathcal{W}}$.

In [Y2], an explicit description of the homomorphism $\varphi_C : \mathbb{C}[\mathcal{V}] \rightarrow \text{Cox}(X)$ is given in terms of the notion of a *G-nat family* introduced by Logvineko [Lo]. To put it shortly, a *G-nat family* on a crepant resolution $X \rightarrow \mathbb{C}^3/G$ is a family of G -constellations which extends to X the family over the open subset of \mathbb{C}^3/G parametrizing the free G -orbits. To give such a family is equivalent to choosing rays of $\sigma_{\mathcal{V}}$ such that their images under $p : (N_{\mathcal{V}})_{\mathbb{R}} \rightarrow \mathbb{R}^3$ are the rays of the toric fan of X . The main result of [Y2] for $n = 3$ states that X is realized as a fine moduli space of G -constellations (not-necessarily of the form \mathfrak{M}_{θ}) if and only if X admits a *G-nat family* \mathcal{F} such that the maximal cones of the ‘‘associated’’ subfan $\Sigma_{X,\mathcal{F}}$ of $\Sigma_{\mathcal{V}}$ has the expected dimension 3 (see [Y2, §4] for details). Note that the results in [Y2] work for abelian $G \subset SL_n(\mathbb{C})$ with arbitrary n .

Remark 3.6. If we add redundant homogeneous generators (like $x_j^2 + x_{j'}$ if $\chi_j^2 = \chi_{j'}$) to the generators x_1, \dots, x_n of $\mathbb{C}[V]$, we obtain a strictly bigger toric variety $\bar{Y}_{\mathcal{V}}$ than $\bar{\mathcal{V}}$. \square

Example 3.2. (Continuation of Example 3.1)

We consider again the abelian G treated in Example 3.1. In order to describe $\bar{\mathcal{V}}$ concretely, we fix coordinates $\theta_0, \dots, \theta_9$ of $\theta \in \Theta$ so that $\theta_i = \theta(\rho^{\otimes i}) \in \mathbb{R}$ where $\rho : G \rightarrow \mathbb{C}^*$ is the one-dimensional representation defined by $\rho(g) = \zeta_{10}$. We identify Θ with \mathbb{R}^9 via $\theta \mapsto (\theta_1, \dots, \theta_9)$ from now on. Note that $\sum_{i=0}^9 \theta_i = 0$. Then we see that the

cone $\sigma_{\mathcal{V}}$ admits 65 rays, whose images under $p : \mathbb{R}^3 \times \mathbb{R}^9 \rightarrow \mathbb{R}^3$ are equal to the rays $\mathbb{R}_{\geq 0}e_j$ ($j = 1, 2, 3$) and $\mathbb{R}_{\geq 0}v_k$ ($k = 1, \dots, 5$), as expected.

In order to determine which rays of $\sigma_{\mathcal{V}}$ correspond to exceptional divisors of X_1 , recall that the G -Hilbert scheme is realized as $\mathcal{V} //_{\theta_+} T_{\Theta}$ for any $\theta_+ \in \Theta$ satisfying $(\theta_+)_i > 0$ for all $i > 0$. Note that GIT-chamber C_1 for the G -Hilbert scheme contains all such θ_+ . Using Proposition 3.6, we see that the rays generated by the following vectors in $\mathbb{R}^3 \times \mathbb{R}^9$ are the rays in Σ_{θ_+} :

$$\begin{aligned} w_1 &= (1, 3, 6, -1, -2, -3, -4, -5, -6, -7, -8, -9) \\ w_2 &= (1, 2, 3, -1, -2, -3, -4, -5, -1, -2, -3, -4) \\ w_3 &= (2, 1, 2, -2, -4, -1, -3, -5, -2, -4, -6, -3) \\ w_4 &= (1, 1, 0, -1, 0, -1, 0, -1, 0, -1, 0, -1) \\ w_5 &= (7, 1, 2, -7, -4, -1, -8, -5, -2, -9, -6, -3). \end{aligned}$$

One can also determine w_k 's by using the fact that the universal family for the G -Hilbert scheme is the *maximal shift family*, see [Lo, §3.5] for details.

Now we can explicitly describe φ_{C_1} as explained in Remark 3.4(1). If we embed $\text{Pic}(X_1)$ into \mathbb{Z}^5 as we have been doing so far (cf. (3.2)), then the map $(\psi_{C_1})_{\mathbb{R}} : \mathbb{R}^9 \rightarrow \mathbb{R}^5$ is given by the matrix

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 \\ 1 & 2 & 3 & 4 & 5 & 1 & 2 & 3 & 4 \\ 2 & 4 & 1 & 3 & 5 & 2 & 4 & 6 & 3 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 7 & 4 & 1 & 8 & 5 & 2 & 9 & 6 & 3 \end{pmatrix}.$$

One can check that we have $(\psi_{C_1})_{\mathbb{R}}(\overline{C}_1) = \text{Nef}(X_1)$.

Now we can explicitly confirm that every crepant resolution of \mathbb{C}^3/G is realized as \mathfrak{M}_{θ} for some generic θ , which is guaranteed by the main result of [CI]. As mentioned in Remark 3.4(1), the map ψ_C for a GIT-chamber C coincides with ψ_{C_1} as long as C is contained in the intersection $C_+ := \bigcap_{k=1}^5 C_{w_k}$ of the orbit cones

$$C_{w_k} = \text{Cone}(\{\text{wt}(x_{i,j}) \in \Theta \mid w_k \subset \{x_{i,j} = 0\}\}).$$

One sees that we have $(\psi_{C_1})_{\mathbb{R}}(C_+) = \bigcup_{i=1}^5 \text{Nef}(X_i)$, which implies that the interior of

$$\overline{C}_i := (\psi_{C_1})_{\mathbb{R}}^{-1}(\text{Nef}(X_i)) \cap C_+ \quad (i = 1, \dots, 5)$$

is a GIT-chamber C_i whose associated moduli space equals X_i . The resolution X_6 is also achieved as a moduli space by crossing a boundary of C_+ . More precisely, one can check that the hyperplane $\{\theta_0 + \theta_1 + \theta_3 = 0\}$ forms a facet of both C_+ and C_3 and that the adjacent chamber C'_3 to C_3 regarding this facet corresponds to the ray of $\sigma_{\mathcal{V}}$ generated by

$$w'_3 = (2, 1, 2, -2, 1, -1, 2, 0, 3, 1, -1, 2).$$

Then the counterpart C'_+ of C_+ obtained by replacing C_3 with C'_3 satisfies

$$(\psi_{C'_3})_{\mathbb{R}}(C'_+) \supset \text{Nef}(X_6).$$

Therefore, we obtain a GIT-chamber $C_6 \subset C'_+$ such that $\mathfrak{M}_{C_6} \cong X_6$. \square

3.5.2 D_5 -singularity, continuation of Example 2.1

We next consider the case of D_5 -singularity again (see Example 2.1). Note that the construction of embeddings in Subsection 3.3 is valid in dimension two since the moduli spaces for any generic θ are crepant (or minimal) resolutions of \mathbb{C}^2/G . We fix a basis of the regular representation of G determined by the matrix presentations in Example 2.1 (see Subsection 3.3).

If we use x, y as the standard coordinates of $V = \mathbb{C}^2$ again, the generic matrices for the G -constellation family over \mathbb{C}^2 are given as

$$A_0 = \begin{pmatrix} x \\ y \end{pmatrix}, B_0 = \begin{pmatrix} -y & x \end{pmatrix}, A_1 = \begin{pmatrix} -iy \\ x \end{pmatrix}, B_1 = \begin{pmatrix} x & iy \end{pmatrix}, A_2 = \begin{pmatrix} x \\ -y \end{pmatrix}, B_2 = \begin{pmatrix} y & x \end{pmatrix},$$

$$A_3 = \begin{pmatrix} iy \\ x \end{pmatrix}, B_3 = \begin{pmatrix} x & -iy \end{pmatrix}, C = \begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix}, D = \begin{pmatrix} -y & 0 \\ 0 & x \end{pmatrix}$$

for the choice of basis of the vector spaces $\text{Hom}_{\mathbb{C}G}(V^* \otimes_{\mathbb{C}} \rho, \rho')$ in Example 2.1. Here we regard the arrows of Q_G as the corresponding matrices by abuse of notation. Note that these matrices satisfy the condition (2.3). We take $Ab(G)^\vee$ -homogeneous elements

$$f_1 = x^3 + iy^3, \quad f_2 = xy, \quad f_3 = x^3 - iy^3$$

so that $\mathbb{C}[V]^{[G,G]} = \mathbb{C}[f_1, f_2, f_3]$. Note that $g \in G$ acts on f_k via the character ρ_k . As is shown in [D, §6] and [Y1, Example 1], these f_k 's give rise to generators of the Cox ring of the minimal resolution X of V/G . We will consider the embedding of \mathcal{W} into a toric variety with respect to these f_k 's.

Let us consider the semi-invariants obtained as the determinants of the following 42 square (block) matrices:

$$C, D, B_2A_0, B_0A_2, B_3A_1, B_1A_3, (A_0 \ A_2), \begin{pmatrix} B_0 \\ B_2 \end{pmatrix}, (A_1 \ A_3), \begin{pmatrix} B_1 \\ B_3 \end{pmatrix}, B_1CA_0, B_3CA_0,$$

$$B_1CA_2, B_3CA_2, B_0DA_1, B_2DA_1, B_0DA_3, B_0DA_3, (CA_0 \ A_1), (CA_0 \ A_3),$$

$$(CA_2 \ A_1), (CA_2 \ A_3), (A_0 \ DA_1), (A_0 \ DA_3), (A_2 \ DA_1), (A_2 \ DA_3),$$

$$\begin{pmatrix} B_0 \\ B_1C \end{pmatrix}, \begin{pmatrix} B_0 \\ B_3C \end{pmatrix}, \begin{pmatrix} B_2 \\ B_1C \end{pmatrix}, \begin{pmatrix} B_2 \\ B_3C \end{pmatrix}, \begin{pmatrix} B_0D \\ B_1 \end{pmatrix}, \begin{pmatrix} B_0D \\ B_3 \end{pmatrix}, \begin{pmatrix} B_2D \\ B_1 \end{pmatrix}, \begin{pmatrix} B_2D \\ B_3 \end{pmatrix}, \begin{pmatrix} A_0 \ D \\ 0 \ B_1 \end{pmatrix},$$

$$\begin{pmatrix} A_0 \ D \\ 0 \ B_3 \end{pmatrix}, \begin{pmatrix} A_2 \ D \\ 0 \ B_1 \end{pmatrix}, \begin{pmatrix} A_2 \ D \\ 0 \ B_3 \end{pmatrix}, \begin{pmatrix} B_0 \ 0 \\ C \ A_1 \end{pmatrix}, \begin{pmatrix} B_0 \ 0 \\ C \ A_1 \end{pmatrix}, \begin{pmatrix} B_2 \ 0 \\ C \ A_3 \end{pmatrix}, \begin{pmatrix} B_2 \ 0 \\ C \ A_3 \end{pmatrix}.$$

We will see that these semi-invariants give a generating system of $\mathbb{C}[\mathcal{W}] = \mathbb{C}[\mathcal{V}]^{SL_R}$. Although it might be possible to check this directly (with help of a computer) by using algorithms such as the one in [DeK, 4.1], we adopt another approach.

In order to construct $Y_{\mathcal{W}}$ explicitly, we use coordinates $\theta_0, \dots, \theta_5$ of $\theta \in \Theta$ such that $\theta_i = \theta(\rho_i)$ for $i = 0, 1, 2, 3$ and $\theta_i = \theta(V_{i-3})$ for $i = 4, 5$. We then identify Θ with \mathbb{R}^5 via $\theta \mapsto (\theta_1, \dots, \theta_5)$. Note that we have $\theta_0 + \theta_1 + \theta_2 + \theta_3 + 2\theta_4 + 2\theta_5 = 0$ in this case. Each of the 42 semi-invariants above is mapped to one of f_i 's under φ up to constant multiplication, and we construct an embedding $\mathcal{W} \hookrightarrow Y_{\mathcal{W}}$ with respect to f_i 's. For example, the associated vectors in $M_{\mathcal{W}} \subset \mathbb{Z}^{3+5}$ to the semi-invariants for the matrices C and $\begin{pmatrix} B_2 & 0 \\ C & A_3 \end{pmatrix}$ are given as $(0, 1, 0, 0, 0, 0, -1, 1)$ and $(1, 0, 0, 0, 1, -1, -1, 1)$ respectively.

Let $\sigma' \subset \mathbb{R}^{3+5}$ be the dual cone to the vectors in $M_{\mathcal{W}}$ associated to the above semi-invariants. We will show that σ' equals the cone $\sigma_{\mathcal{W}} \subset \mathbb{R}^8$ defining $\bar{Y}_{\mathcal{W}}$. Let \mathcal{W}' be the spectrum of the subalgebra of $\mathbb{C}[\mathcal{W}]$ generated by the 42 semi-invariants above. Then we obtain an embedding of \mathcal{W}' into a toric variety $Y'_{\mathcal{W}}$ whose normalization is defined by σ' , similar to the construction of $\mathcal{W} \hookrightarrow Y_{\mathcal{W}}$. In fact one can check that $Y'_{\mathcal{W}}$ is normal by comparing $\sigma'^{\vee} \cap M_{\mathcal{W}}$ and the semi-group generated by the elements of $M_{\mathcal{W}}$ associated to the 42 semi-invariants. This computation can be done e.g. by using the package “Normaliz” of Macaulay2 [GS].

We see, by calculation, that σ' admits 165 rays. The primitive generators in \mathbb{Z}^8 of these rays include the first three coordinate vectors $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$, and the other 162 vectors are divided into five types by looking at their first three coordinates:

$$v_1 = (1, 2, 1), \quad v_2 = (1, 1, 1), \quad v_3 = (3, 2, 3), \quad v_4 = (3, 2, 5), \quad \text{or } v_5 = (5, 2, 3). \quad (3.7)$$

These five types correspond to the prime exceptional divisors E_1, \dots, E_5 of the minimal resolution X defined by the conjugacy classes of $g_1, g_1^2, g_1^3, g_2, g_1g_2$ respectively (cf. [Y1, Example 1]). More precisely, the rays for E_1, \dots, E_5 in the fan of the ambient toric variety Y of X (with respect to f_i 's) are generated by the vectors in (3.7). Note that σ' particularly satisfies the following condition:

$$\text{the polyhedral cone } p^{-1}(\mathbb{R}_{\geq 0}v_k) \cap \sigma' \text{ is generated by rays of } \sigma' \text{ for each } k. \quad (3.8)$$

This will be important when one computes a generating system of semi-invariants efficiently (see Remark 3.8).

The fact that the rays of σ' are projected to those of the fan of X implies that $Y'_{\mathcal{W}}//_{\theta}T_{\Theta}$ is isomorphic to the toric variety Y for any generic $\theta \in \Theta$. Therefore, we have $\sigma_{\mathcal{W}} = \sigma'$, and the natural toric morphism $\bar{Y}_{\mathcal{W}} \rightarrow Y'_{\mathcal{W}}$ is an isomorphism. This also implies that the 42 semi-invariants generate $\mathbb{C}[\mathcal{W}]$.

Since that the G -Hilbert scheme is given by a stability condition θ_+ satisfying $(\theta_+)_i > 0$ for all $i > 0$, we see that generators of the rays of $\sigma_{\mathcal{W}}$ corresponding the exceptional divisors E_1, \dots, E_5 of the G -Hilbert scheme are given by


$$\begin{aligned} w_1 &= (1, 2, 1, -1, -2, -1, -2, -2), & w_2 &= (1, 1, 1, -1, -1, -1, -2, -2), \\ w_3 &= (3, 2, 3, -3, -2, -3, -4, -6), & w_4 &= (3, 2, 5, -3, -2, -5, -4, -6), \\ w_5 &= (5, 2, 3, -5, -2, -3, -4, -6) \end{aligned}$$

respectively. From these data, one can also explicitly compute the map $(\psi_C)_{\mathbb{R}} : \mathbb{R}^5 \rightarrow \mathbb{R}^5$ associated to the GIT-chamber C containing θ_+ .

Remark 3.7. In [Le, §2], the G -Hilbert schemes X for binary dihedral groups are studied, and in particular the author introduced distinguished G -clusters which give rise to open affine covering of X . For example, the distinguished G -cluster F of *type B* is defined by an homogeneous ideal of $\mathbb{C}[x, y]$ and is presented as the following \mathbb{C} -vector subspace:

$$\mathbb{C} \oplus (\mathbb{C}x \oplus \mathbb{C}y) \oplus \mathbb{C}xy \oplus (\mathbb{C}x^2 \oplus \mathbb{C}y^2) \oplus \mathbb{C}x^3 \oplus \mathbb{C}y^3 \oplus (\mathbb{C}x^4 \oplus \mathbb{C}y^4) \oplus (\mathbb{C}x^5 \oplus \mathbb{C}y^5).$$

Its structure as a $\mathbb{C}[V]$ -module is also presented as in the following diagram:



where each arrow $\rho \rightarrow \rho'$ represents a nontrivial action of V^* on ρ into ρ' similarly to the McKay quiver. One sees that, among the 42 semi-invariants, the ones h_1, \dots, h_9 for the matrices

$$B_1CA_0, B_2A_0, B_3CA_0, (CA_0 \ A_1), (CA_0 \ A_3), (A_0 \ DA_1), (A_0 \ DA_3), \begin{pmatrix} A_0 & D \\ 0 & B_1 \end{pmatrix}, \begin{pmatrix} A_0 & D \\ 0 & B_3 \end{pmatrix}$$

do not vanish at F , and thus the face of $\sigma_{\mathcal{W}}$ corresponding to F is given as the intersection $\sigma_{\mathcal{W}} \cap \{w_{h_1} = \dots = w_{h_9} = 0\}$ where $w_{h_i} \in M_{\mathcal{W}}$ is the associated element to h_i . This is equal to $\text{Cone}(w_1, w_2)$, and thus F corresponds to the the intersection point of the curves E_1 and E_2 . \square

3.5.3 Trihedral group of order 21

Here we deal with the following three-dimensional example:


$$G = \left\langle g_1 = \begin{pmatrix} \zeta_7 & 0 & 0 \\ 0 & \zeta_7^2 & 0 \\ 0 & 0 & \zeta_7^4 \end{pmatrix}, g_2 = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \right\rangle \subset SL_3(\mathbb{C})$$

where ζ_7 is the seventh root of unity. Then $\text{Irr}(G)$ consists of three 1-dimensional representations $\rho_k : G \rightarrow \mathbb{C}^*$ ($k = 0, 1, 2$) defined by $\rho_k(g_1) = 1$, $\rho_k(g_2) = \omega^k$ where ω is the third root of unity, and two 3-dimensional representations V_1, V_2 , the first one is the inclusion $G \subset SL_3(\mathbb{C})$ presented above and the other is presented as

$$g_1 \mapsto \begin{pmatrix} \zeta_7^3 & 0 & 0 \\ 0 & \zeta_7^6 & 0 \\ 0 & 0 & \zeta_7^5 \end{pmatrix}, g_2 \mapsto \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

We again fix the basis obtained in this way for the subsequent computations.

The McKay quiver Q_G for this case is as follows:



where C_i stands for two arrows C_1 and C_2 .

With the standard coordinates x, y, z of $V = \mathbb{C}^3$, the generic matrices are given as

$$\begin{aligned} A_0 &= \begin{pmatrix} x \\ y \\ z \end{pmatrix}, B_0 = (z \ x \ y), A_1 = \begin{pmatrix} x \\ \omega y \\ \omega^2 z \end{pmatrix}, B_1 = (\omega z \ x \ \omega^2 y), \\ A_2 &= \begin{pmatrix} x \\ \omega^2 y \\ \omega z \end{pmatrix}, B_2 = (\omega^2 z \ x \ \omega y), C_1 = \begin{pmatrix} 0 & x & 0 \\ 0 & 0 & y \\ z & 0 & 0 \end{pmatrix}, C_2 = \begin{pmatrix} y & 0 & 0 \\ 0 & z & 0 \\ 0 & 0 & x \end{pmatrix}, \\ D &= \begin{pmatrix} 0 & y & 0 \\ 0 & 0 & z \\ x & 0 & 0 \end{pmatrix}, L_1 = \begin{pmatrix} 0 & 0 & z \\ x & 0 & 0 \\ 0 & y & 0 \end{pmatrix}, L_2 = \begin{pmatrix} 0 & z & 0 \\ 0 & 0 & x \\ y & 0 & 0 \end{pmatrix} \end{aligned}$$

for some choice of basis $\text{Hom}_{\mathbb{C}G}(\rho \otimes_{\mathbb{C}} V^*, \rho')$ with $\rho, \rho' \in \text{Irr}(G)$.

In this case, it seems hard to obtain generators of semi-invariants directly using the algorithm in [DeK, Ch. 4] since the input data are large. Instead we give a candidate of a system of generators and then confirm the correctness as we did in the previous example.

As computed in [DG, §5.1], there are $Ab(G)^\vee$ -homogeneous polynomials $f_1, \dots, f_{13} \in \mathbb{C}[V]^{[G,G]}$ such that the associated elements

$$\tilde{f}_1, \dots, \tilde{f}_{13}, t_1^{-7}, t_2^{-3}, t_3^{-3} \in \mathbb{C}[V]^{[G,G]}[t_1^{\pm 1}, t_2^{\pm 1}, t_3^{\pm 1}]$$

generate the Cox ring of a crepant resolution X of V/G , where we set the variables t_1, t_2, t_3 so that they correspond to junior elements g_1, g_2, g_2^2 respectively (see Subsection 3.1).

Similarly to the previous examples, we realize the cone $\sigma_{\mathcal{W}}$ inside \mathbb{R}^{13+5-1} by ordering the elements of $\text{Irr}(G)$ as $\rho_0, \rho_1, \rho_2, V_1, V_2$. Regarding the embedding $\iota_X : X \hookrightarrow Y$ with respect to f_1, \dots, f_{13} , the generators of the rays in \mathbb{R}^{13} corresponding to the exceptional divisors E_1, E_2, E_3 are given as

$$\begin{aligned} \mathbf{v}_1 &= (1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1), \\ \mathbf{v}_2 &= (0, 0, 2, 1, 0, 2, 1, 3, 2, 1, 3, 2, 4), \\ \mathbf{v}_3 &= (0, 0, 1, 2, 0, 1, 2, 3, 1, 2, 3, 4, 2) \end{aligned}$$

respectively. Then we define $\sigma' \subset \mathbb{R}^{17}$ as the cone generated by the coordinate vectors $\mathbf{e}_1, \dots, \mathbf{e}_{13}$ and the following 16 vectors:

$$\begin{aligned} \mathbf{v}_{1,1} &= (\mathbf{v}_1, -1, -1, -1, -2), \quad \mathbf{v}_{1,2} = (\mathbf{v}_1, 0, -1, -1, -2), \quad \mathbf{v}_{1,3} = (\mathbf{v}_1, -1, 0, -1, -2), \\ \mathbf{v}_{1,4} &= (\mathbf{v}_1, 0, 0, -1, -2), \quad \mathbf{v}_{1,5} = (\mathbf{v}_1, 0, 0, -1, 0), \quad \mathbf{v}_{1,6} = (\mathbf{v}_1, 0, 0, 0, 1), \\ \mathbf{v}_{1,7} &= (\mathbf{v}_1, 0, 0, 2, 1), \quad \mathbf{v}_{1,8} = (\mathbf{v}_1, 0, 1, 2, 1), \quad \mathbf{v}_{1,9} = (\mathbf{v}_1, 1, 0, 2, 1), \quad \mathbf{v}_{1,10} = (\mathbf{v}_1, 1, 1, 2, 1), \\ \mathbf{v}_{2,1} &= (\mathbf{v}_2, -1, -2, -3, -3), \quad \mathbf{v}_{2,2} = (\mathbf{v}_2, -1, 1, 0, 0), \quad \mathbf{v}_{2,3} = (\mathbf{v}_2, 2, 1, 3, 3), \\ \mathbf{v}_{3,1} &= (\mathbf{v}_3, -2, -1, -3, -3), \quad \mathbf{v}_{3,2} = (\mathbf{v}_3, 1, -1, 0, 0), \quad \mathbf{v}_{3,3} = (\mathbf{v}_3, 1, 2, 3, 3). \end{aligned}$$

By computing σ' explicitly (with help of Macaulay2 [GS]), one sees that it has 400 facets and accordingly we obtain 400 primitive elements of $M_{\mathcal{W}}$ defining these facets. In

fact we can take determinantal semi-invariants corresponding to these primitive elements. One can also check that the toric variety $Y'_{\mathcal{W}}$ constructed from these elements is already normal again. Since the rays of σ' are projected to those of the fan of X , we have that $\sigma' = \sigma_{\mathcal{W}}$ again and that the associated 400 semi-invariants generate $\mathbb{C}[\mathcal{W}]$.

Remark 3.8. In general, a candidate σ' of $\sigma_{\mathcal{W}}$ can be given by computing determinantal semi-invariants h (from ones with $\varphi(h)$ having smaller degrees) until the dual cone

$$\sigma'' := \bigcap_h \{v \in \mathbb{R}^\ell \oplus \Theta^\vee \mid w_h(v) \geq 0\}$$

satisfies the condition (3.8). This may be achieved by computing relatively small number of semi-invariants (compared to the number of generators of $\mathbb{C}[\mathcal{W}]$). Then the candidate σ' is defined as the cone generated by the rays of σ'' which are projected to the rays of the fan of X .

In the above case, such σ'' is obtained from determinantal semi-invariants coming from 3×3 -matrices only, while the 400 generators of $\mathbb{C}[\mathcal{W}]$ include ones coming from 6×6 -matrices. One may also verify that $\sigma_{\mathcal{W}} = \sigma'$ by checking the existence of G -constellations corresponding to rays of σ' , as will be done below. \square

Once we know the cone $\sigma_{\mathcal{W}}$, one can investigate the structures of G -constellations presented by faces of $\sigma_{\mathcal{W}}$. As an demonstration, we take the ray generated by $\mathbf{v}_{1,1}$. If we take the generators $\mathbf{v}_{1,1}, \dots, \mathbf{v}_{1,10}$ of $\sigma'_{\mathcal{W}}$ whose former coordinates is \mathbf{v}_1 , the orbit cone $C_{\mathbf{v}_{1,j}} \subset \Theta$ of the corresponding divisor to $\mathbf{v}_{1,j}$ is given as the cone

$$\{\theta \in \Theta \mid p(\mathbf{v}_{1,j}) \cdot \theta = \min_{j=1, \dots, 10} p(\mathbf{v}_{1,j}) \cdot \theta\}$$

by Proposition 3.6. Note that $\{C_{\mathbf{v}_{1,j}}\}_{j=1, \dots, 10}$ give a subdivision of Θ into polyhedral cones. The orbit cone $C_{\mathbf{v}_{1,1}} \subset \Theta$ is defined by the following 5 inequalities:

$$\theta_1 \geq 0, \theta_2 \geq 0, \theta_1 + \theta_2 + 2\theta_4 \geq 0, \theta_1 + \theta_2 + \theta_3 + 3\theta_4 \geq 0, \theta_1 + \theta_2 + 3\theta_3 + 3\theta_4 \geq 0.$$

Notice that the coefficients of these inequalities are obtained as the direction vectors $\overrightarrow{\mathbf{v}_{1,1}\mathbf{v}_{1,j}}$ for j such that the line segment $\overline{\mathbf{v}_{1,1}\mathbf{v}_{1,j}}$ forms an edge of the convex hull $\text{Conv}(\mathbf{v}_{1,1}, \dots, \mathbf{v}_{1,10})$. One can infer the structure of the G -constellation F of a general point of E_1 from these inequalities. In this case the structure of F is presented as the following diagram for some decomposition $F = \rho_0 \oplus \rho_1 \oplus \rho_2 \oplus V_1^{\oplus 3} \oplus V_2^{\oplus 3}$:


$$\begin{array}{ccccccc} \rho_0 & \longrightarrow & V_1 & \longrightarrow & V_2 & & V_2 & \longrightarrow & \rho_1 \\ & & \downarrow & & \downarrow & \nearrow & & \searrow & \\ & & V_1 & \longrightarrow & V_1 & \longrightarrow & V_2 & \longrightarrow & \rho_2 \end{array}$$

As another example, the orbit cone $C_{\mathbf{v}_{1,6}}$ for $\mathbf{v}_{1,6} = (\mathbf{v}_1, 0, 0, 0, 1)$ is defined by

$$\theta_3 \geq 0, \theta_1 + 2\theta_3 \geq 0, \theta_2 + 2\theta_3 \geq 0, \theta_1 + \theta_2 + 2\theta_3 \geq 0,$$

$$\theta_3 + \theta_4 \leq 0, \theta_1 + \theta_3 + 3\theta_4 \leq 0, \theta_2 + \theta_3 + 3\theta_4 \leq 0, \theta_1 + \theta_2 + \theta_3 + 3\theta_4 \leq 0$$

and the structure of the G -constellation F for $\mathbf{v}_{1,6}$ is presented as



As a more explicit description, F can be realized as a quotient of the ideal of $\mathbb{C}[V]$ generated by xy, yz and zx . Note that $\mathbb{C}xy \oplus \mathbb{C}yz \oplus \mathbb{C}zx$ is isomorphic to V_2 as a $\mathbb{C}G$ -module.

Finally, we compute the homomorphism ψ_C for certain chambers C and see that every projective crepant resolution of V/G is obtained as a moduli space of G -constellations. Let $C_{i,j} \subset \Theta$ be the orbit cones corresponding to the rays of $\sigma_{\mathcal{W}}$ generated by $\mathbf{v}_{i,j}$. There are 42 choices of a triple (i, j, k) such that $\overline{C}_{1,i} \cap \overline{C}_{2,j} \cap \overline{C}_{3,k}$ has the full dimension 4. Among them, $C_+ := \overline{C}_{1,1} \cap \overline{C}_{2,1} \cap \overline{C}_{3,1}$ is the one containing the GIT-chamber C_{++} for the G -Hilbert scheme. C_{++} is in fact defined by the inequalities $\theta_i > 0$ ($i = 1, 2, 3, 4$).

Similarly to the abelian case in Example 3.2, one can compute $\psi_{C_{++}}$ and obtain $\psi_{C_{++}}(C_+) = \text{Mov}(X)$. Thus, every projective crepant resolution of V/G is realized as \mathfrak{M}_θ for some generic $\theta \in C_+$. In this case the movable cone $\text{Mov}(X)$ is a three-dimensional simplicial cone divided into four nef cones (cf. [DG, Proposition 5.2]). The nef cone for the G -Hilbert scheme is given as $\psi_{C_{++}}(\overline{C}_{++})$ and we see that it is the central one which can pass to the three remaining (corner) ones. Note that not all of $\overline{C}_{1,i} \cap \overline{C}_{2,j} \cap \overline{C}_{3,k}$ of dimension 4 realize all the resolutions. For example, $\psi_{C'}(C)$ is strictly smaller than $\text{Mov}(X)$ for the cone $C = \overline{C}_{1,2} \cap \overline{C}_{2,1} \cap \overline{C}_{3,1}$ adjacent to C_+ and any GIT-chamber $C' \subset C$.

4 Proof of the main result

In this section we give a proof of the following theorem:

Theorem 4.1. (*=Theorem 1.1*) *Let $G \subset SL_3(\mathbb{C})$ be a finite subgroup and $X \rightarrow \mathbb{C}^3/G$ a projective crepant resolution. Then there is a generic stability condition $\theta \in \Theta$ such that $X \cong \mathfrak{M}_\theta$.*

To prove Theorem 4.1, we will use a similar argument to the one in [CI, §8]. To this end, we should know how the or the tautological bundle \mathcal{R}_C of \mathfrak{M}_C changes when we cross a wall in Θ . Here we say that the intersection $W = \overline{C} \cap \overline{C}'$ for GIT-chambers $C, C' \subset \Theta$ is a *wall* (of C) if W is a codimension-one face of C (and hence of C').

For any given GIT-chamber $C \subset \Theta$, we choose any general point θ_0 of a wall $W \subset C$. Then we have $\mathcal{W}^{C\text{-ss}} \subset \mathcal{W}^{\theta_0\text{-ss}}$ and this inclusion induces a morphism $\alpha : \mathfrak{M}_C \rightarrow \overline{\mathfrak{M}}_{\theta_0}$ over $\mathfrak{M}_0 = \mathbb{C}^3/G$ where $\overline{}$ denotes the normalization. Similarly to the case of \mathfrak{M}_C , we

denote the moduli space $\bar{\mathfrak{M}}_{\theta_0}$ by $\bar{\mathfrak{M}}_W$. We also denote the crepant resolution \mathfrak{M}_C by X as in previous sections.

Recall that each line bundle $L \in \text{Pic}(X)$ defines a character $\chi_L \in \chi(T_X)$ and its associated GIT-quotient $X_L := \mathfrak{X} //_{\chi_L^k} T_X$ with a sufficiently divisible $k > 0$. As observed in Subsection 3.2, X_L is the same as the image of the morphism obtained by the complete linear system of (some power of) L . Note that $X_0 = \mathbb{C}^3/G$ and that X_L makes sense for $L \in \text{Pic}(X)_{\mathbb{R}}$. One can determine $\alpha : \mathfrak{M}_C \rightarrow \bar{\mathfrak{M}}_W$ by looking at the homomorphism $\psi_C : \Theta \rightarrow \text{Pic}(X)_{\mathbb{R}}$ constructed in Subsection 3.3:

Lemma 4.2. (cf. [CI, Lemma 3.3]) *For any point $\theta \in \bar{C}$, the normalized GIT-quotient $\bar{\mathfrak{M}}_{\theta}$ is isomorphic to $X_{\psi_C(\theta)}$.*

Proof. Recall that $X = \mathfrak{M}_C$ is regarded as a GIT-quotient of $\mathfrak{X} = \text{Spec Cox}(X)$ via the restriction $\mathfrak{X}^{\psi_C(C)\text{-ss}} \rightarrow \mathcal{W}^{C\text{-ss}}$ of the equivariant map $\mathfrak{X} \rightarrow \mathcal{W}$ obtained from φ_C (see Subsection 3.3). Then the line bundle $L := \psi_C(\theta)$ is just the pullback of the ample line bundle $\mathcal{O}_{\bar{\mathfrak{M}}_{\theta}}(1)$ (induced by the GIT-construction) under the morphism $X \rightarrow \bar{\mathfrak{M}}_{\theta}$. Then the claim follows since $X \rightarrow \bar{\mathfrak{M}}_{\theta}$ is identified with the morphism coming from the linear system of L . \square

Lemma 4.3. *Let $F \subset \Theta$ be a face of an orbit cone $\tilde{C}_x \subset \Theta$ of the G -constellation corresponding to a point $x \in \mathfrak{M}_C$. If the relative interior $\text{relint}(F)$ of F intersects with \bar{C} and if $\psi_C(F) \subset \text{Pic}(X)_{\mathbb{R}}$ is not of full-dimension, then $\psi_C(F)$ is on the boundary of the orbit cone $C_x \subset \text{Pic}(X)_{\mathbb{R}}$ of a point of \mathfrak{X} which is a lift of $x \in X$.*

Proof. We first choose homogeneous generators h_1, \dots, h_{ℓ} of $\mathbb{C}[\mathcal{W}] = \mathbb{C}[\mathcal{V}]^{SL_R}$ so that we have an associated embedding $\mathcal{W} \hookrightarrow Y_{\mathcal{W}}$ and its quotient $X \hookrightarrow Y$ as in Proposition 3.5. Let $\bar{x} \in \mathcal{W}$ be any lift of x . By reordering h_i 's, we may assume that

(1) there exists $\ell_1 < \ell$ such that h_1, \dots, h_{ℓ_1} do not vanish at \bar{x} and their weights generate the cone F and that

(2) for any h_k with $k > \ell_1$, either h_k vanishes at \bar{x} or $\text{wt}(h_k)$ is not in F .

The locally closed subset of \mathcal{W} consisting of points having F as their orbit cones is given by

$$\mathcal{W}_F := \mathcal{W} \cap \{h_1 h_2 \cdots h_{\ell_1} \neq 0\} \cap \{h_{\ell_1+1} = \cdots = h_{\ell} = 0\}.$$

Note that torus-orbits of the toric varieties Y , $\mathbb{A}^{\ell+m}$, and $Y_{\mathcal{W}}$ restrict to stratification of X , \mathfrak{X} , and \mathcal{W} respectively, and that \mathcal{W}_F is such a stratum of \mathcal{W} .

By construction, the equivariant map $\varphi_C^* : \mathfrak{X} \rightarrow \mathcal{W}$ associated to C is the restriction of a toric morphism $\mathbb{A}^{\ell+m} \rightarrow Y_{\mathcal{W}}$ which is obtained by sending the rays $\tilde{\tau}_k$ of the fan of $\mathbb{A}^{\ell+m}$ corresponding to the exceptional divisors E_k to the rays $\tau_{C,k}$ of $\sigma_{\mathcal{W}}$ corresponding to E_k in \mathfrak{M}_C . For $\theta_0 \in \text{relint}(F) \cap \bar{C}$, the restriction $\mathfrak{X}^{\psi_C(\theta_0)\text{-ss}} \rightarrow \mathcal{W}^{\theta_0\text{-ss}}$ of φ_C^* descends to an isomorphism $X_{\psi_C(\theta_0)} \rightarrow \bar{\mathfrak{M}}_{\theta_0}$ by Lemma 4.2. Note that \mathcal{W}_F is a closed subset of $\mathcal{W}^{\theta_0\text{-ss}}$ which intersects with closure of the T_{Θ} -orbit of \bar{x} . In particular, the categorical quotient of \mathcal{W}_F by T_{Θ} is again a closed subset in $\bar{\mathfrak{M}}_{\theta_0}$.

We ‘‘perturb’’ the map φ_C^* to obtain another equivariant map $\varphi'^* : \mathfrak{X} \rightarrow \mathcal{W}$ such that its restriction to the semistable loci still descends to an isomorphism $X_{\psi_C(\theta_0)} \rightarrow \bar{\mathfrak{M}}_{\theta_0}$ and that every point of $\mathcal{W}_F \subset \mathcal{W}^{\theta_0\text{-ss}}$ is inside a T_{Θ} -orbit of a point in the image of φ'^* .

Such φ'^* is explicitly constructed as follows. For each $k = 1, \dots, m$, let σ_k be the face of $\sigma_{\mathcal{W}}$ generated by $\tau_{C,k}$ and the rays τ of $\sigma_{\mathcal{W}}$ (if any) such that τ is projected to τ_k under $p : (N_{\mathcal{W}})_{\mathbb{R}} \rightarrow (N_{\mathcal{W}}/\Theta^{\vee})_{\mathbb{R}}$ (see Subsection 3.4) and that the orbit cone of τ contains F as a face. Note that σ_k is equal to $\tau_{C,k}$ if $\tau_{C,k} \not\subset \sigma_F$ and otherwise σ_k is the intersection $\sigma_F \cap p^{-1}(\tau_k)$ where σ_F is the face of $\sigma_{\mathcal{W}}$ corresponding to the stratum \mathcal{W}_F . Then φ'^* is obtained as the restriction of the toric morphism $\mathbb{A}^{\ell+m} \rightarrow Y_{\mathcal{W}}$ obtained by sending each ray $\tilde{\tau}_k$ to a ray generated by a general point of σ_k . Note that φ'^* indeed induces an isomorphism $X_{\psi_C(\theta_0)} \rightarrow \tilde{\mathfrak{M}}_{\theta_0}$ since the projected cone $p(\sigma)$ of the image σ of each cone of the fan of $\mathfrak{X}^{\psi_C(\theta_0)\text{-ss}}$ under the induced homomorphism $\mathbb{R}^{\ell+m} \rightarrow \mathbb{R}^{\ell+s-1}$ by φ_C^* remains the same even if we replace φ_C^* by φ'^* .

The map φ'^* gives an associated homomorphism $\psi' : \Theta \rightarrow \text{Pic}(X)_{\mathbb{R}}$ similarly to ψ_C . More concretely, for a homogeneous semi-invariant $f \in \mathbb{C}[\mathcal{W}]$ of weight $c \in \Theta$, $\psi'(c)$ is the class of the sum of the strict transform of the image in V/G of $\{f = 0\}$ and $\sum_k v_k(f)E_k$ where v_k is the primitive element in σ_k defining φ'^* (cf. Remarks 3.1 and 3.4(1)).

One sees that $\psi'|_F = c\psi_C|_F$ for some constant $c \in \mathbb{Q}$ and in particular $\psi'(F)$ is not full-dimensional by the assumption. Let $\varphi' : \mathbb{C}[\mathcal{W}] \rightarrow \mathbb{C}[\mathfrak{X}]$ be the ring homomorphism associated to φ'^* . The equivariant map φ'^* restricts to a map from the nonempty (affine) locally closed subset

$$\mathfrak{X}_F := (\varphi'^*)^{-1}(\mathcal{W}_F) = \mathfrak{X} \cap \{\varphi'(h_1 h_2 \cdots h_{\ell_1}) \neq 0\} \cap \{\varphi'(h_{\ell_1+1}) = \cdots = \varphi_C(h_{\ell}) = 0\}$$

of \mathfrak{X} to \mathcal{W}_F , which descends to an isomorphism between the (categorical) quotients by T_X and T_{Θ} .

We assume that $\psi'(F)$ does not lie on the boundary of C_x to deduce a contradiction. Then the quotients of \mathfrak{X}_F and \mathcal{W}_F are isomorphic to the stratum S_x of X containing x . If \mathfrak{X}_F itself is a stratum of \mathfrak{X} , then it must be the set \mathfrak{X}_x of points of \mathfrak{X} whose orbit cones are equal to C_x . The fact that $\psi'(F)$ is not of full-dimension implies that every point of \mathfrak{X}_x has a positive-dimensional stabilizer in T_X , which is a contradiction since C_x is full-dimensional. If there exists a stratum $\mathfrak{X}_{x'} \subsetneq \mathfrak{X}_F$ not equal to \mathfrak{X}_x , then this is again a contradiction since the orbit cone of such $\mathfrak{X}_{x'}$ is a proper face of C_x and in particular $\mathfrak{X}_{x'}$ cannot be mapped to S_x under the quotient map. Note that proper faces of an orbit cone of \mathfrak{X} always correspond to nontrivial contractions, in contrast to faces of orbit cones in Θ such as walls of *Type 0* introduced below. \square

If a wall $W = \overline{C} \cap \overline{C}'$ is not contained in the pullback of a hyperplane of $\text{Pic}(X)_{\mathbb{R}}$ under ψ_C , then the morphism $\mathfrak{M}_C \rightarrow \tilde{\mathfrak{M}}_W$ is an isomorphism by Lemma 4.2, and we call such a wall of *Type 0* following [CI]. Conversely, for any wall W whose image under ψ_C is not of full dimension, the morphism $\mathfrak{M}_C \rightarrow \tilde{\mathfrak{M}}_W$ is a non-trivial contraction of either a divisor or a curve by Lemma 4.3. For a wall W of Type 0, the two moduli spaces \mathfrak{M}_C and $\mathfrak{M}_{C'}$ are isomorphic but the tautological bundle changes. In fact $\mathcal{R}_{C'}$ is a modification of \mathcal{R}_C by tensoring a line bundle to a subbundle of \mathcal{R}_C similar to the abelian cases [CI, Corollary 4.2], as we will see below.

In order to study what happens when we cross a wall W of Type 0, we first see that the *unstable locus*

$$D_W = \{x \in X \mid \text{any lift } \bar{x} \in \mathcal{W} \text{ of } x \text{ is not } W\text{-stable}\}$$

is a reduced compact divisor. Note that D_W has a natural scheme structure as the set of zeros of sections of line bundles. Later we will also see that D_W is connected (cf. Proposition 4.9).

Lemma 4.4. *For any given GIT-chamber $C \subset \Theta$ and its wall W of Type 0, the unstable locus D_W is a reduced compact divisor.*

Proof. We first show that D_W is reduced. Note that D_W is a geometric quotient of (an open subset of) a closed subscheme \tilde{D}_W of \mathcal{W} . By the definition of the stability, \tilde{D}_W is the zero locus of the semi-invariants which vanish at the (closed) points of \tilde{D}_W . Since semi-invariants are nothing but regular functions on the affine variety \mathcal{W} , the subscheme \tilde{D}_W is reduced and so is D_W .

Next we show that D_W is a divisor. We fix an embedding $\mathcal{W} \hookrightarrow Y_{\mathcal{W}}$ again, and let Σ_{θ} be the subfan of $\sigma_{\mathcal{W}}$ defining $Y_{\mathcal{W}}^{\theta\text{-ss}}$ for $\theta \in \Theta$ (see Subsection 3.4). We also consider the ambient toric variety Y of X and let σ_x be the cone of the fan Σ_X of Y corresponding to a point $x \in X \subset Y$.

For any point $x \in D_W$, there exists a unique cone $\tilde{\sigma}_x \in \Sigma_{\theta}$ for $\theta \in C$ such that $p(\tilde{\sigma}_x) = \sigma_x$. Since $\mathfrak{M}_C \rightarrow \bar{\mathfrak{M}}_W$ is an isomorphism, we have a cone $\bar{\sigma}_x \in \Sigma_{\theta_0}$ strictly containing $\tilde{\sigma}_x$ (as a facet) such that it satisfies $p(\bar{\sigma}_x) = \sigma_x$ where θ_0 is a general point of W . Recall that the maximal cones of the fan of $\mathcal{W}^{W\text{-ss}}$ are projected under p to the maximal cones of Σ_X (see Subsection 3.4). Since the image of any ray of $\sigma_{\mathcal{W}}$ under p is again a ray of Σ_X , there must be a ray τ of $\tilde{\sigma}_x$ and a face $\bar{\tau} \subset \bar{\sigma}_x$ such that $\bar{\tau} \supsetneq \tau$ and $p(\tau) = p(\bar{\tau})$. This implies that x is contained in the unstable divisor of X corresponding to $p(\bar{\tau})$. Thus, D_W is of pure codimension one.

For the compactness of D_W , one may apply exactly the same argument in [CI, Proposition 4.4]. \square

Now we describe how G -constellations over unstable points change when we cross the wall W . For convenience, we will consider corresponding representations of Q_G rather than G -constellations in the rest of this section.

Since the regular representation R is primitive as an element of $R(G)$, one can take nonzero $\mathbb{C}G$ -modules M_1 and M_2 which are primitive as elements of $R(G)$ such that $\theta_0(M_1) = 0$ for any θ_0 of W and that $R = aM_1 + M_2$ in $R(G)$ for some $a \in \mathbb{N}$. We may assume that M_1 is not contained in M_2 (as a direct summand) and vice versa. We also assume $\theta(M_1) > 0$ for $\theta \in C$ since the other case $\theta(M_1) < 0$ can be treated similarly by switching the roles of C and C' . Let F be the representation of Q_G corresponding to any point $x \in D_W \subset X = \mathfrak{M}_C$. Then there exists a unique maximal $\mathbb{C}[V]$ -submodule $S \subsetneq F$ such that $\theta_0(S) = 0$ for any θ_0 of W , and let Q be the quotient of F by S . Then Q is indecomposable since it is θ_0 -semistable for general $\theta_0 \in W$. Note that S and Q as $\mathbb{C}G$ -modules may a priori vary on different connected components of D_W if $a > 0$, but the primitiveness of the quotient Q is independent of the choice of x .

We take a 1-PS $\lambda : \mathbb{C}^* \rightarrow GL_R$ defined by Q (for a fixed connected component of D_W), that is, λ is positive on C and zero on W as a function on $\Theta \cong \chi(GL_R)_{\mathbb{R}}$. Then λ induces a filtration of F by its submodules (cf. [Kin, §3]) such that Q is one of its associated graded component. This works for a family of representations of Q_G and thus

we obtain an exact sequence

$$0 \rightarrow \mathcal{S} \rightarrow \mathcal{R}_C|_{D_W} \rightarrow \mathcal{Q} \rightarrow 0 \quad (4.1)$$

of flat families \mathcal{S} and \mathcal{Q} of $\mathbb{C}[V]$ -modules corresponding to S and Q respectively. Note that this sequence splits as vector bundles on D_W , whereas it is a nontrivial extension as $\mathbb{C}[V]$ -modules. The ranks of \mathcal{S} and \mathcal{Q} as vector bundles on D_W a priori vary on different connected components of D_W but it will be shown that D_W is connected in Proposition 4.9. Let \mathcal{K} be the kernel of the natural surjection $\mathcal{R}_C \rightarrow \mathcal{Q}$. Then \mathcal{K} is a vector bundle on X (with a $\mathbb{C}[V]$ -action) called the *elementary transformation* of \mathcal{R}_C (associated to $\mathcal{R}_C \rightarrow \mathcal{Q}$) in the sense of [M]. Applying the inverse of the elementary transformation to \mathcal{K} , we obtain an exact sequence

$$0 \rightarrow \mathcal{Q} \otimes \mathcal{O}_X(-D_W) \rightarrow \mathcal{K}|_{D_W} \rightarrow \mathcal{S} \rightarrow 0 \quad (4.2)$$

of flat families of $\mathbb{C}[V]$ -modules. Note that \mathcal{K} parametrizes the same representations of Q_G as \mathcal{R}_C outside D_W while sub and quotient modules are switched over D_W .

Lemma 4.5. *The family \mathcal{K} coincides with the tautological bundle $\mathcal{R}_{C'}$ for the GIT-chamber C' up to tensoring a line bundle on X .*

Proof. First recall that the bundle $\mathcal{R}_{C'} = \bigoplus_{\rho \in \text{Irr}(G)} \mathcal{R}'_{\rho}$ is normalized by tensoring a line bundle so that the component \mathcal{R}'_{ρ_0} becomes the trivial line bundle. In order to see that \mathcal{K} parametrizes θ' -(semi)stable representations of Q_G for $\theta' \in C'$, we only have to show that the fibers of $\mathcal{K}|_{D_W}$ are indecomposable as $\mathbb{C}[V]$ -modules. To do this, it suffices to show that the fibers of $\mathcal{K}|_{D_W}$ are indecomposable over a Zariski open dense subset of D_W since \mathcal{K} and $\mathcal{R}_{C'}$ are vector bundles on a smooth variety.

Let $x \in D_W$ be a general point contained in a single irreducible component $D \subset D_W$ and let F be the corresponding representation of Q_G in \mathfrak{M}_C . We first consider the case when the fiber S of \mathcal{S} at x is indecomposable. Since having a nontrivial direct summand (of a fixed $\mathbb{C}[V]$ -module structure) is a closed condition, every fiber of $\mathcal{S}|_D$ is indecomposable by the genericity of x . The same argument as in the proof of [CI, Proposition 4.1] shows that the sequence (4.2) is a nontrivial extension of $\mathbb{C}[V]$ -modules and therefore the fibers of $\mathcal{K}|_D$ are indecomposable.

Next we consider the other case when there exists a nontrivial irreducible direct summand S_1 of S . Then one can take a 1-PS λ of GL_R such that $\lim_{t \rightarrow 0} (\lambda(t) \cdot F)$ admits exactly two irreducible direct summands one of which is S_1 . Then, similarly to (4.1), we have an exact sequence

$$0 \rightarrow \mathcal{S}_1 \rightarrow \mathcal{R}_C|_D \rightarrow \mathcal{Q}_1 \rightarrow 0$$

of families corresponding to the sequence $0 \rightarrow S_1 \rightarrow F \rightarrow Q_1 \rightarrow 0$. Applying the elementary transformation to $\mathcal{R}_C \rightarrow \mathcal{Q}_1$, we obtain a family \mathcal{K}' with the fibers of $\mathcal{K}'|_D$ indecomposable similarly as in the first case. By repeating this procedure for direct summands of S , we will obtain the family \mathcal{K} with indecomposable fibers and thus the claim follows. \square

Remark 4.1. In abelian cases every irreducible representation of G has multiplicity one in R and thus the modules S and Q are clearly indecomposable. When G is non-abelian and S is non-primitive in $R(G)$, the module S may be decomposable. Even if S is indecomposable, the θ_0 -polystable representation \bar{F} of Q_G (i.e. the direct sum of θ_0 -stable modules) corresponding to the image of x under $\mathfrak{M}_C \rightarrow \mathfrak{M}_W$ is not isomorphic to the direct sum $S \oplus Q$ in general. S may degenerate in \bar{F} to decompose into a direct sum $\bigoplus_i S_i$ satisfying $\theta_0(S_i) = 0$. \square

Next we describe the behavior of the Grothendieck group $K(X)$ of locally free sheaves on X under the wall-crossing. As we will see below, it is basically the same as the abelian cases and this enables us to proceed the proof of the main result in a similar way to [CI]. To this end we briefly review the necessary notions. See [CI, §2.4, §5, and §7.1] and also [BKR, §9.2] for details.

The result of [BKR] shows that the Fourier-Mukai transform by the universal family \mathcal{U}_C gives an equivalence $\Phi_C : D(X) \cong D^G(\mathbb{C}^3)$ whose inverse $\Phi_C^{-1} : D^G(\mathbb{C}^3) \rightarrow D(X)$ is given by $\mathcal{O}_{\mathbb{C}^3} \otimes \rho \mapsto \mathcal{R}_\rho^\vee$. Φ_C induces an isomorphism between $K(X)$ and the Grothendieck group $K^G(\mathbb{C}^3)$ of $D^G(\mathbb{C}^3)$. Let $K_0(X)$ be the subgroup of $K(X)$ generated by the classes of complexes whose supports are contained in the fiber $\pi^{-1}(0)$ for $\pi : X \rightarrow \mathbb{C}^3/G$. Then $K_0(X)$ is identified with the dual of $K(X)$ via the perfect pairing $\chi(-, -) : K(X) \times K_0(X) \rightarrow \mathbb{Z}$ defined by

$$(\alpha, \beta) \mapsto \sum_i (-1)^i \dim \text{Ext}^i(\alpha, \beta),$$

which is skew-symmetric on $K_0(X)$. Via the equivalence Φ_C we also have a perfect pairing $\chi^G(-, -)$ between $K^G(\mathbb{C}^3)$ and the Grothendieck group $K_0^G(\mathbb{C}^3)$ of G -equivariant sheaves on \mathbb{C}^3 supported at the origin. With respect to this pairing, the classes of $\{\mathcal{O}_{\mathbb{C}^3} \otimes \rho\}_{\rho \in \text{Irr}(G)}$ and $\{\mathcal{O}_0 \otimes \rho\}_{\rho \in \text{Irr}(G)}$ give mutually dual basis of $K(\mathbb{C}^3)$ and $K_0(\mathbb{C}^3)$ respectively where \mathcal{O}_0 is the skyscraper sheaf at the origin of \mathbb{C}^3 .

The equivalence Φ_C also induces an isomorphism $\phi_C : K_0(X) \rightarrow K_0^G(\mathbb{C}^3)$ and its pullback $\phi_C^* : K^G(\mathbb{C}^3) \rightarrow K(X)$. The vector space Θ is regarded as a subspace of $K^G(\mathbb{C}^3)_{\mathbb{R}}$, and the restriction of ϕ_C^* to Θ gives an isomorphism $\Theta \rightarrow F^1$ where we have a filtration

$$K(X)_{\mathbb{R}} = F^0 \supset F^1 \supset F^2 \supset F^3 = 0$$

with F^i being the subspace generated by sheaves whose supports are of codimension at least i . The homomorphism $\phi_C^*|_{\Theta} : \Theta \rightarrow F^1$ is explicitly given as

$$\theta = (\theta_\rho)_\rho \mapsto \sum_{\rho \in \text{Irr}(G)} \theta_\rho [\mathcal{R}_\rho^\vee]$$

where $[\mathcal{R}_\rho^\vee]$ denotes the class of the dual sheaf \mathcal{R}_ρ^\vee in F^1 . Then we obtain a homomorphism $L_C : \Theta \rightarrow \text{Pic}(X)_{\mathbb{R}}$ such that the following diagram of vector spaces is commutative:

$$\begin{array}{ccc} \Theta & \xrightarrow{\phi_C^*|_{\Theta}} & F^1 \\ L_C \downarrow & & \text{pr} \downarrow \\ \text{Pic}(X)_{\mathbb{R}} & \xleftarrow{\det^{-1}} & F^1/F^2 \end{array} \quad (4.3)$$

where pr is the projection and the lower horizontal arrow denotes the isomorphism $F^1/F^2 \cong \text{Pic}(X)_{\mathbb{R}}$ given by sending the class $[\mathcal{F}] \in F^1$ to $\det(\mathcal{F})^{-1}$.

Lemma 4.6. L_C is equal to the homomorphism $\psi_C : \Theta \rightarrow \text{Pic}(X)_{\mathbb{R}}$ defined in Subsection 3.3.

Proof. By definition L_C is given as

$$\theta = (\theta_{\rho})_{\rho} \mapsto \det \left(\bigoplus_{\rho \in \text{Irr}(G)} \mathcal{R}_{\rho}^{\oplus \theta_{\rho}} \right).$$

For a determinantal semi-invariant f having $\theta = (\theta_{\rho})_{\rho}$ as its weight, $\varphi_C(f)$ is regarded as a section of a line bundle $L_C(\theta)$ by noticing that each arrow $a_{\rho, \rho'}$ in Q_G is regarded as a section of $\mathcal{R}_{\rho}^{\vee} \otimes \mathcal{R}_{\rho'}$. Thus, $L_C(\theta)$ is nothing but the line bundle $\psi_C(\theta)$. Since Θ is spanned by the weights of determinantal semi-invariants by Theorem 2.2, L_C and ψ_C are the same on the whole Θ . \square

When we cross a wall from a GIT-chamber C to an adjacent one C' , the corresponding chambers $\phi_C^*(C)$ and $\phi_{C'}^*(C')$ in F^1 may not be adjacent in general. For example, in dimension two, the automorphism $\phi_{C'}^* \circ (\phi_C^*)^{-1}$ of $F^1 \cong \text{Pic}(X)$ is a reflection along the hyperplane containing $\phi_C^*(W)$ for every wall $W = \overline{C} \cap \overline{C}'$. In the proof of the main result of [CI], namely the abelian case of Theorem 4.1, the fact that, for a wall-crossing of Type 0, the chambers $\phi_C^*(C)$ and $\phi_{C'}^*(C')$ in F^1 are adjacent up to tensoring a line bundle plays a crucial role. We will show that the same property holds for non-abelian groups as well (cf. Proposition 4.9).

Lemma 4.7. *Let $W = \overline{C} \cap \overline{C}'$ be a wall of Type 0 and consider the exact sequence on the unstable locus D_W as in (4.1). Then, for each connected component $D \subset D_W$, exactly one of the following two situations occurs:*

- (1) *Either \mathcal{S} or \mathcal{Q} parametrizes mutually isomorphic $\mathbb{C}[V]$ -modules on D .*
- (2) *D is a product of two curves C_1, C_2 where each C_i is a tree of rational curves of Dynkin type, that is, C_i is isomorphic to the (reduced) exceptional locus of the minimal resolution of an ADE surface singularity.*

Proof. Let $R_1 \subset R$ (resp. $R_2 \subset R$) be the $\mathbb{C}G$ -module obtained by forgetting the structure as $\mathbb{C}[V]$ -modules on the G -equivariant modules parametrized by \mathcal{S}_x (resp. \mathcal{Q}_x) of $x \in D$. Since the morphism $\alpha : \mathfrak{M}_C \rightarrow \mathfrak{M}_W$ is an isomorphism, we may regard the divisor D as a moduli space parametrizing pairs of isomorphism types of structures as $\mathbb{C}[V]$ -modules on R_1 and R_2 . We construct moduli spaces $\mathfrak{M}_{\mathcal{S}}$ (resp. $\mathfrak{M}_{\mathcal{Q}}$) parametrizing $\mathbb{C}[V]$ -modules appearing as fibers of $\mathcal{S}|_D$ (resp. $\mathcal{Q}|_D$), as GIT-quotients of certain affine schemes $\mathcal{V}_{\mathcal{S}}, \mathcal{V}_{\mathcal{Q}}$ by the GL_{R_i} -actions ($i = 1, 2$) in a similar way to the case of \mathfrak{M}_{θ} (see Section 2). The stability condition for $\mathfrak{M}_{\mathcal{S}}$ (resp. $\mathfrak{M}_{\mathcal{Q}}$) is chosen so that its associated semi-stable locus in $\mathcal{V}_{\mathcal{S}}$ (resp. $\mathcal{V}_{\mathcal{Q}}$) contains the points corresponding to fibers of $\mathcal{S}|_D$ (resp. $\mathcal{Q}|_D$) and is minimal satisfying this property. Then we have a natural equivariant morphism $\mathcal{V}_{\mathcal{S}} \times \mathcal{V}_{\mathcal{Q}} \rightarrow \mathcal{V}$ which descends to an isomorphism $\mathfrak{M}_{\mathcal{S}} \times \mathfrak{M}_{\mathcal{Q}} \rightarrow \bar{D} \subset \mathfrak{M}_W$ where \bar{D} is the image of D under the morphism $\mathfrak{M}_C \rightarrow \mathfrak{M}_W$.

Since D_W is a surface by Lemma 4.4, we only have to show that \mathfrak{M}_S and \mathfrak{M}_Q are curves of Dynkin type when they are one-dimensional. Indeed, this particularly implies that each irreducible component of \bar{D} is smooth and thus $D \rightarrow \bar{D}$ is an isomorphism. Note that $\mathfrak{M}_C \rightarrow \mathfrak{M}_W$ has connected fibers since each fiber of the quotient map $\mathcal{V}^{W\text{-ss}} \rightarrow \mathfrak{M}_W$ is the closure in $\mathcal{V}^{W\text{-ss}}$ of a GL_R -orbit.

Let $E'_1, \dots, E'_{m'}$ be the irreducible components of D and $C_{E'_k} \subset \Theta$ their orbit cones. We also let $H \subset \Theta$ be the hyperplane containing W . Then, by moving from the relative interior of $\bigcap_k C_{E'_k} \cap H$ to its smaller faces, we obtain $\theta \in \Theta$ and its associated birational map $\alpha_\theta : \mathfrak{M}_C \dashrightarrow \mathfrak{M}_\theta$ such that α_θ is defined on D and that α_θ only collapses the first factor \mathfrak{M}_S of D and in particular the image $\alpha_\theta(D)$ is the curve \mathfrak{M}_Q . This is indeed possible since, if we encounter a wall corresponding to a contraction of the second factor \mathfrak{M}_Q during the process, then we just have to cross it. Note that moving from a face to its codimension-one face gives an isomorphism or a contraction of relative Picard number one between the corresponding (normalized) moduli spaces, and in particular each process does not contract the two factors at the same time.

Therefore, the curve \mathfrak{M}_S is identified with the fiber, under α_θ , over a general point of the one-dimensional singular locus of $\bar{\mathfrak{M}}_\theta$. Then it must be a tree of rational curves of Dynkin type since $\bar{\mathfrak{M}}_\theta$ has only canonical singularities and α_θ is crepant (cf. [Re1, Corollary 1.14]). The same argument works for \mathfrak{M}_Q as well. \square

For given $\mathcal{E} \in K_0(X)$, we consider the automorphism $\mathcal{T}_\mathcal{E}$ of $K(X)$ defined by

$$\mathcal{T}_\mathcal{E}(\xi) = \xi - \chi(\xi, \mathcal{E}) \cdot \mathcal{E}.$$

Note that $\mathcal{T}_\mathcal{E}$ has positive determinant and fixes the hyperplane

$$\mathcal{E}^\perp := \{\xi \in K(X) \mid \chi(\xi, \mathcal{E}) = 0\}$$

pointwise. We denote by $\mathcal{T}'_\mathcal{E}$ the inverse of $\mathcal{T}_\mathcal{E}$. We will show that the automorphism $\phi_{C'}^* \circ (\phi_C^*)^{-1}$ of F^1 coincides with $\mathcal{T}_\mathcal{E}$ or $\mathcal{T}'_\mathcal{E}$ for some $\mathcal{E} \in K_0(X)$ up to an automorphism obtained by tensoring a line bundle of X (Proposition 4.9).

For the families \mathcal{S} and \mathcal{Q} in (4.1) associated to the wall W , let \mathcal{S}_ρ and \mathcal{Q}_ρ be their ρ -isotypic components respectively.

Lemma 4.8. *For any $\rho, \sigma \in \text{Irr}(G)$, we have $H^i(D_W, \mathcal{S}_\rho^\vee \otimes \mathcal{Q}_\sigma) = 0$ for all i .*

Proof. We first note that the G -equivariant bundle $\bigoplus_{\rho \in \text{Irr}(G)} \mathcal{R}_\rho^\vee \otimes \rho^*$ on $X \times \mathbb{C}^3$ can be regarded as the universal family of the moduli space $\mathfrak{M}_{\theta'}$ for $\theta' \in \Theta$ defined by $\theta'(\rho) = -\theta(\rho^*)$ with $\theta \in C$ by [CI, Lemma 2.6(ii)]. Similarly, \mathcal{K}^\vee is regarded as a tautological bundle for a moduli space obtained from $\mathfrak{M}_{\theta'}$ by crossing a wall of Type 0. Then we see that the dual bundles are related as

$$0 \rightarrow \mathcal{R}_C^\vee \rightarrow \mathcal{K}^\vee \rightarrow \mathcal{Q}^\vee \otimes \mathcal{O}_X(D_W) \rightarrow 0 \quad (4.4)$$

noticing that \mathcal{R}_C and \mathcal{K} are related by the exact sequences

$$0 \rightarrow \mathcal{K} \rightarrow \mathcal{R}_C \rightarrow \mathcal{Q} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow \mathcal{R}_C(-D_W) \rightarrow \mathcal{K} \rightarrow \mathcal{S} \rightarrow 0. \quad (4.5)$$

From (4.4) we obtain a long exact sequence

$$\begin{aligned} \cdots \rightarrow \mathrm{Ext}_{\mathcal{O}_X}^i(\mathcal{R}_C^\vee, \mathcal{R}_C^\vee) &\rightarrow \mathrm{Ext}_{\mathcal{O}_X}^i(\mathcal{R}_C^\vee, \mathcal{K}^\vee) \rightarrow \mathrm{Ext}_{\mathcal{O}_X}^i(\mathcal{R}_C^\vee, \mathcal{Q}^\vee \otimes \mathcal{O}_X(D_W)) \\ &\rightarrow \mathrm{Ext}_{\mathcal{O}_X}^{i+1}(\mathcal{R}_C^\vee, \mathcal{R}_C^\vee) \rightarrow \cdots \end{aligned} \quad (4.6)$$

since \mathcal{R}_C^\vee is locally free. Applying the equivalence Φ_C , we have

$$\mathrm{Ext}_{\mathcal{O}_X}^i(\mathcal{R}_C^\vee, \mathcal{R}_C^\vee) = \mathrm{Ext}_{\mathcal{O}_{\mathbb{C}^3}}^i(\mathcal{O}_{\mathbb{C}^3} \otimes_{\mathbb{C}} R, \mathcal{O}_{\mathbb{C}^3} \otimes_{\mathbb{C}} R)^G = \begin{cases} \mathbb{C}[V] & \text{if } i = 0 \\ 0 & \text{if } i \neq 0 \end{cases} \quad (4.7)$$

(cf. [CI, Lemma 5.4]). Since the kernel of the map from the second term to the third in (4.6) is isomorphic to

$$H^i(D_W, \mathcal{R}_C|_{D_W} \otimes \mathcal{S}^\vee) = H^i(D_W, (\mathcal{S} \oplus \mathcal{Q}) \otimes \mathcal{S}^\vee),$$

we obtain $H^i(D_W, \mathcal{S}^\vee \otimes \mathcal{Q}) = 0$ for $i > 0$ by (4.7). Since elements of $\mathbb{C}[V]$ of positive degree are zero on $\mathcal{R}_C^\vee|_{D_W}$, we also see that $H^0(D_W, \mathcal{S}^\vee \otimes \mathcal{Q}) = 0$. Decomposing $\mathcal{S}^\vee \otimes \mathcal{Q}$ into isotypic components, we obtain the claim. \square

Proposition 4.9. *Let $W = \overline{C} \cap \overline{C}'$ be a wall of Type 0 with the associated sequence (4.1) on the unstable locus D_W . Then D_W is connected. Moreover, if we let R_1 and R_2 be the underlying $\mathbb{C}G$ -modules of fibers of \mathcal{S} and \mathcal{Q} respectively, then there exists $\mathcal{E} \in K_0(X)$ such that*

- (i) *the image $\phi_C(\mathcal{E}) \in K_0(\mathbb{C}^3) \cong R(G)$ is equal to either R_1 or R_2 , and that*
- (ii) *the automorphism of $K(X)$ induced by the autoequivalence $\Phi_{C'}^{-1} \circ \Phi_C$ of $D(X)$ is equal to one of the four automorphisms: (a) $\mathcal{T}_\mathcal{E}$, (b) $\mathcal{T}'_\mathcal{E}$, (c) $\xi \mapsto \mathcal{T}_\mathcal{E}(\xi) \otimes \mathcal{O}(-D_W)$, and (d) $\xi \mapsto \mathcal{T}'_\mathcal{E}(\xi \otimes \mathcal{O}(D_W))$.*

In particular, $\phi_{C'}^(C')$ is adjacent to $\phi_C^*(C) \otimes L$ for some L in $\mathrm{Pic}^c(X)$, the subgroup of $\mathrm{Pic}(X)$ generated by the classes of compact divisors.*

Proof. Let $D_W = \bigcup_i D_i$ be a decomposition into connected components and let $R_{1,i}$ (resp. $R_{2,i}$) be the $\mathbb{C}G$ -modules for fibers of $\mathcal{S}|_{D_i}$ (resp. $\mathcal{Q}|_{D_i}$). Note that all $R_{1,i}$ (resp. $R_{2,i}$) are proportional in $R(G)$ since D_i 's are defined by the same wall W .

We first show that we can take \mathcal{E} for each D_i such that $\phi_C^*(\mathcal{E})$ is proportional to $R_{1,i}$ or $R_{2,i}$. We consider the two cases (1), (2) in Lemma 4.7 separately. For Case (1), we first assume that $\mathcal{Q}|_{D_i}$ is a trivial family of G -equivariant $\mathbb{C}[V]$ -modules. Since every fiber of \mathcal{Q} is indecomposable, there exists a line bundle L on D_i such that $\mathcal{Q}_\rho|_{D_i} \cong L^{\oplus \mathrm{rank}(\mathcal{Q}_\rho|_{D_i})}$ for all ρ . Note that $R_{2,i} = \bigoplus_{\rho \in \mathrm{Irr}(G)} \rho^{\oplus \mathrm{rank}(\mathcal{Q}_\rho|_{D_i})}$ in $R(G)$.

We set $\mathcal{E} := [L^{-1} \otimes \omega_{D_i}] \in K_0(X)$. The cohomology sheaves of $\Phi_C(L^{-1} \otimes \omega_{D_i})$, which

are G -equivariant $\mathcal{O}_{\mathbb{C}^3}$ -modules supported at the origin, are computed as

$$\begin{aligned}
\Phi_C^j(L^{-1} \otimes \omega_{D_i}) &\cong \bigoplus_{\rho \in \text{Irr}(G)} H^j(X, \mathcal{R}_\rho \otimes L^{-1} \otimes \omega_{D_i}) \otimes_{\mathbb{C}} \rho \\
&\cong \bigoplus_{\rho \in \text{Irr}(G)} H^{2-j}(D_i, \mathcal{R}_\rho^\vee \otimes L)^\vee \otimes_{\mathbb{C}} \rho \\
&\cong \bigoplus_{\rho \in \text{Irr}(G)} H^{2-j}(D_i, ((L^{-1})^{\oplus \text{rank}(\mathcal{Q}_\rho|_{D_i})} \oplus \mathcal{S}_\rho^\vee) \otimes L)^\vee \otimes_{\mathbb{C}} \rho \\
&\cong \bigoplus_{\rho \in \text{Irr}(G)} H^{2-j}\left(D_i, \mathcal{O}_{D_i}^{\oplus \text{rank}(\mathcal{Q}_\rho|_{D_i})}\right)^\vee \otimes_{\mathbb{C}} \rho
\end{aligned}$$

where the last isomorphism holds by Lemma 4.8. Therefore, the class $\phi_C^*(\mathcal{E})$ is equal to $\chi(\mathcal{O}_{D_i}) \cdot R_{2,i} \in R(G)$.

When $\mathcal{S}|_{D_i}$ is a trivial family, one can similarly show that $\phi_C^*([L'])$ is equal to $\chi(\mathcal{O}_{D_i}) \cdot R_{1,i} \in R(G)$ for the line bundle L' on D_i such that $\mathcal{S}_\rho|_{D_i} \cong L' \oplus^{\text{rank}(\mathcal{S}_\rho|_{D_i})}$.

Next we consider the case (2) in Lemma 4.7. In this case each D_i is the product of curves C_1 and C_2 which are regarded as moduli spaces parametrizing fibers of $\mathcal{S}|_{D_i}$ and $\mathcal{Q}|_{D_i}$ respectively. Since each C_j is a union of \mathbb{P}^1 's, the bundle $\mathcal{S}|_{D_i}$ (resp. $\mathcal{Q}|_{D_i}$) on D_i is a direct sum of line bundles such that the restrictions of these line bundles to fibers of the projection $D_i \rightarrow C_1$ (resp. $D_i \rightarrow C_2$) are mutually isomorphic. We take any line bundles L_1 and L_2 on D_i appearing as direct summands of $\mathcal{S}_\rho|_{D_i}$ and $\mathcal{Q}_{\rho'}|_{D_i}$ respectively for any ρ, ρ' . By Lemma 4.8, we have $H^j(D_i, L_1^{-1} \otimes L_2) = 0$ for all j . Then either

$$H^j(C_1, (L_1^{-1} \otimes L_2)|_{C_1}) = 0 \ (\forall j) \quad \text{or} \quad H^j(C_2, (L_1^{-1} \otimes L_2)|_{C_2}) = 0 \ (\forall j)$$

holds where C_1 and C_2 are regarded as fibers of $D_i \rightarrow C_2$ and $D_i \rightarrow C_1$ respectively. From this we see that one has either $H^j(C_2, (L_1^{-1} \otimes L_2')|_{C_2}) = 0 \ (\forall j)$ for all line bundles L_2' appearing as direct summands of $\mathcal{Q}|_{D_i}$ or $H^j(C_1, (L_1^{-1} \otimes L_2)|_{C_1}) = 0 \ (\forall j)$ for all line bundles L_1' appearing as direct summands of $\mathcal{S}|_{D_i}$. In the former case one has $\phi_C([L_2^{-1}]) = R_{2,i} \in R(G)$, and in the latter case $\phi_C([L_1] \otimes \omega_{D_i}) = R_{1,i}$. This can be checked by noticing that the Euler characteristics $\chi(L_2'|_{C_2})$ (resp. $\chi(L_1'|_{C_1})$) are the same for all the direct summands L_2' of $\mathcal{Q}|_{D_i}$ in the former case (resp. L_1' of $\mathcal{S}|_{D_i}$ in the latter case). Note that the Euler characteristic of a line bundle L on C_i is equal to $1 + \sum_k \text{deg}(L|_{C_{i,k}})$ where $C_{i,k}$ are the irreducible components of C_i .

We have shown that the image in $R(G)$ of a line bundle on D_i is proportional to $R_{1,i}$ (and to $R_{2,i}$) modulo $\mathbb{Z}R$ in any case. Since D_i 's are linearly independent in $\text{Pic}(X)_{\mathbb{R}} \cong F^1/F^2$, the classes of the line bundles on D_i 's in $K(X)$ are also linearly independent (cf. the proof of [CI, Proposition 5.5(ii)]). Therefore, D_W must be connected. The similar argument in the proof of [CI, Proposition 5.5(iii)] also shows that we have $H^1(\mathcal{O}_{D_W}) = H^2(\mathcal{O}_{D_W}) = 0$ in the case (1) in Lemma 4.7.

To summarize, the following two cases may occur:

- (I) we have $\chi(\mathcal{R}_\rho^\vee, \mathcal{E}) = \text{rank}(\mathcal{Q}_\rho)$ for the class $\mathcal{E} = [L^{-1} \otimes \omega_{D_W}] \in K_0(X)$ with an arbitrary line bundle L on D_W which is a direct summand of \mathcal{Q} , and
- (II) we have $\chi(\mathcal{R}_\rho^\vee, \mathcal{E}) = \text{rank}(\mathcal{S}_\rho)$ for the class $\mathcal{E} = [L] \in K_0(X)$ with an arbitrary line

bundle L on D_W which is a direct summand of \mathcal{S} .

In the former case (I), one has

$$\mathcal{T}_\mathcal{E}([\mathcal{R}_\rho^\vee]) = [\mathcal{R}_\rho^\vee] - \text{rank}(\mathcal{Q}_\rho)[L^{-1} \otimes \omega_{D_W}] = [\mathcal{R}_\rho^\vee] - [\mathcal{Q}_\rho^\vee(D)]$$

in $K(X)$. By the sequence (4.4) and the fact that $\{\mathcal{R}_\rho^\vee\}_\rho$ forms a basis of $K(X)$, we can conclude that $\mathcal{T}_\mathcal{E}$ coincides with the automorphism induced by $\Phi_{C'}^{-1} \circ \Phi_C$ or its composition with tensoring $\mathcal{O}_X(-D_W)$. Which of the cases (a) and (c) happens depends on whether $\mathcal{Q}_{\rho_0} = 0$ or $\neq 0$. The latter case (II) can be treated similarly by switching the roles of C and C' , so that we fall into the cases (b) and (d). This completes the proof. \square

Remark 4.2. The above proposition shows that the wall-crossing gives a combination of a spherical twist and tensoring a line bundle at least at the level of the Grothendieck group. In the rigid case (1) in Lemma 4.7, one can use the same argument as in the proof of [CI, Proposition 7.3] to prove the similar statement at the level of the derived category. In the case (2), however, this argument does not work since the line bundles in \mathcal{S} or \mathcal{Q} are not mutually isomorphic (while they have the same Euler characteristics) in contrast to the rigid case. By analogy with the abelian case [CI, Corollary 4.6], it is expected that the case (2) does not occur. \square

Proof of Theorem 4.1

Since all projective crepant resolutions of \mathbb{C}^3/G are connected with each other by a sequence of flops, we only have to show that every flop of $X = \mathfrak{M}_C$ can be realized by (not necessarily single) wall-crossings. Let $B \subset \overline{\text{Amp}}(X)$ be a facet corresponding to a flop $X \rightarrow X_B \leftarrow X'$. If we can reach a GIT-chamber C'' through walls of Type 0 such that C'' admits a wall $\tilde{B} \subset \overline{C}''$ satisfying $\psi_{C''}(\tilde{B}) \subset B$, then crossing \tilde{B} induces the desired flop. Indeed, otherwise \tilde{B} would be contained in a facet of the orbit cone of a general point of a divisor since the homomorphism $\psi_{C''}$ must change when we cross \tilde{B} . In such a case, however, Lemma 4.3 shows that \tilde{B} would induce a divisorial contraction, which is a contradiction.

As shown in [CI, §8], each compact exceptional divisor $E \subset X$ acts on F^1 by

$$\xi \mapsto \xi \otimes \mathcal{O}_X(E) = \xi + [\xi \otimes \mathcal{O}_E]$$

with $[\xi \otimes \mathcal{O}_E] \in F^2$. Moreover, for the set of all irreducible compact exceptional divisors S_1, \dots, S_d and for any $\xi \in \text{pr}^{-1}(\text{Amp}(X) \cup \text{relint}(B))$, the set $[\xi \otimes \mathcal{O}_{S_1}], \dots, [\xi \otimes \mathcal{O}_{S_d}]$ forms a basis of F^2 where $\text{Amp}(X)$ is identified with a cone in F^1/F^2 (see (4.3)).

The existence of the desired chamber C'' can be shown similarly to the proof of [CI, Proposition 8.2]. More precisely, one can show from Proposition 4.9 that we have

$$\text{pr}^{-1}(\text{Amp}(X) \cup \text{relint}(B)) \subset \bigcup_{L \in \text{Pic}^c(X)} \bigcup_{\mathfrak{M}_C \cong X} L \otimes \phi_C^*(\overline{C}) \quad (4.8)$$

noticing that GIT-chambers in Θ are finite. Thus we obtain a chamber C'' and its wall \tilde{B} such that $\text{pr}(\phi_C^*(\tilde{B})) \subset B$. \square

Theorem 4.1 together with (4.8) implies that $\{\psi_C^*(\bar{C})\}_{C \in \Theta: \text{chamber}}$ covers the movable cone of X . Thus, every rational contraction $X \dashrightarrow X'$ (see Subsection 3.2) is realized as the birational map $\mathfrak{M}_C \dashrightarrow \mathfrak{M}_\theta$ induced by variation of GIT for some θ . This fact has the following application:

Proposition 4.10. *Let h_1, \dots, h_ℓ be homogeneous generators of the normalization of $\mathbb{C}[\mathcal{W}]$. Then the associated $\ell + m$ elements $\varphi(h_1), \dots, \varphi(h_\ell) \in \mathbb{C}[V]^{[G, G]}$ (see (3.1)) generate the Cox ring $\text{Cox}(X) \subset \mathbb{C}[V]^{[G, G]}[t_1^{\pm 1}, \dots, t_m^{\pm 1}]$ of any crepant resolution $X \rightarrow \mathbb{C}^3/G$.*

Proof. Let $f_1, \dots, f_\ell \in \mathbb{C}[V]^{[G, G]}$ be homogeneous elements such that the associated elements in (3.1) generate $\text{Cox}(X)$. We consider the ambient toric varieties Y of X and $\bar{Y}_{\mathcal{W}}$ of $\bar{\mathcal{W}}$ associated to f_i 's (see Subsection 3.3). Note that each f_i corresponds to a non-exceptional torus-invariant divisor $D_i \subset Y$ and that the image of the corresponding section $f_{D_i} \in H^0(Y, \mathcal{O}(-D_i))$ (i.e. $D_i = \{f_{D_i} = 0\}$) in $\text{Cox}(X)$ is equal to \tilde{f}_i (cf. Remark 3.1). Regarding the weight of f_{D_i} as a character of T_X , we consider its associated GIT-quotient Y_i of $\mathbb{A}^{\ell+m}$. In terms of a fan, Y_i is characterized as a birational model $Y \dashrightarrow Y_i$ with $\dim \text{Pic}(Y_i)_{\mathbb{R}} = 1$ such that the toric fan of Y_i contains the cone generated by the rays corresponding to the torus-invariant divisors of Y except D_i .

For each i , let $X_i \subset Y_i$ be the induced GIT-quotient of $\text{Spec}(\text{Cox}(X))$. As observed above, there exists $\theta \in \Theta$ such that $X_i \cong \mathfrak{M}_\theta$. Since $\bar{Y}_{\mathcal{W}}$ is normal, there is a homogeneous semi-invariant f in $\mathbb{C}[\bar{\mathcal{W}}]$ such that its (scheme-theoretic) zero locus in $\mathfrak{M}_\theta \cong X_i$ is equal to the strict transform of $D_i \cap X$ under $X \dashrightarrow X_i$. If we take a GIT-chamber C so that \bar{C} contains θ , this implies $\varphi_C(\tilde{f})$ is equal to \tilde{f}_i (up to constant multiplication). This particularly implies that each \tilde{f}_i can be obtained as the associated element of $\varphi(h)$ for some homogeneous semi-invariant h . Thus, the claim follows. \square

5 The symplectic resolutions for the binary tetrahedral group

In this section we treat crepant resolutions of a symplectic quotient singularity defined by the binary tetrahedral group $T \subset Sp_4(\mathbb{C})$ (cf. [LS]). It is generated by two elements

$$g_1 = \begin{pmatrix} i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \\ 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \end{pmatrix}, \quad g_2 = -\frac{1}{2} \begin{pmatrix} (1+i)\omega & (-1+i)\omega & 0 & 0 \\ (1+i)\omega & (1-i)\omega & 0 & 0 \\ 0 & 0 & (1+i)\omega^2 & (-1+i)\omega^2 \\ 0 & 0 & (1+i)\omega^2 & (1-i)\omega \end{pmatrix}$$

where ω is the third root of unity. T is isomorphic to a subgroup of $SL_2(\mathbb{C})$ corresponding to E_6 -singularity via

$$g_1 \mapsto \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad g_2 \mapsto -\frac{1}{2} \begin{pmatrix} 1+i & -1+i \\ 1+i & 1-i \end{pmatrix}.$$


We denote this representation by V_0 . Also, $\text{Irr}(G)$ has three 1-dimensional representations $\rho_k : G \rightarrow \mathbb{C}^*$ ($k = 0, 1, 2$) defined by $\rho_k(g_1) = 1$, $\rho_k(g_2) = \omega^k$. Note that the

inclusion $T \subset Sp_4(\mathbb{C})$ corresponds to the direct sum $V_1 \oplus V_2$ where $V_k = V_0 \otimes \rho_k$. Moreover, $\text{Irr}(G)$ contains a unique 3-dimensional representation W given by

$$g_1 \mapsto \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad g_2 \mapsto \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix},$$

so that $\text{Irr}(G) = \{\rho_0, \rho_1, \rho_2, V_0, V_1, V_2, W\}$. In the subsequent computations, we use the basis of the regular representation of G coming from the above presentations similarly as before.

The McKay quiver Q_T is as follows:



where the arrow A_{ki} (resp. B_{ki}) stands for two arrows A_{k1}, A_{k2} (resp. B_{k1}, B_{k2}).

We denote the standard coordinates of $V = \mathbb{C}^4$ by x, y, z, w . Then the generic matrices are given as

$$\begin{aligned} a_{01} &= \begin{pmatrix} x \\ y \end{pmatrix}, b_{10} = (-w \ z), a_{12} = \begin{pmatrix} x \\ y \end{pmatrix}, b_{21} = (-w \ z), a_{20} = \begin{pmatrix} x \\ y \end{pmatrix}, b_{02} = (-w \ z), \\ a_{02} &= \begin{pmatrix} z \\ w \end{pmatrix}, b_{20} = (-y \ x), a_{10} = \begin{pmatrix} z \\ w \end{pmatrix}, b_{01} = (-y \ x), a_{21} = \begin{pmatrix} z \\ w \end{pmatrix}, b_{12} = (-y \ x), \\ A_{01} &= \begin{pmatrix} \omega^2 x & y & -i\omega y \\ -\omega^2 y & x & i\omega x \end{pmatrix}, A_{02} = \begin{pmatrix} i\omega^2 z & i\omega w & w \\ -i\omega^2 w & i\omega z & -z \end{pmatrix}, B_{01} = \begin{pmatrix} y & x \\ -\omega x & \omega y \\ -i\omega^2 x & -i\omega^2 y \end{pmatrix}, B_{02} = \begin{pmatrix} w & z \\ -\omega^2 z & \omega^2 w \\ -i\omega z & -i\omega w \end{pmatrix}, \\ A_{11} &= \begin{pmatrix} x & y & -iy \\ -y & x & ix \end{pmatrix}, A_{12} = \begin{pmatrix} \omega^2 z & w & -i\omega w \\ -\omega^2 w & z & i\omega z \end{pmatrix}, B_{11} = \begin{pmatrix} y & x \\ -\omega^2 x & \omega^2 y \\ -i\omega x & -i\omega y \end{pmatrix}, B_{12} = \begin{pmatrix} w & z \\ -z & w \\ -iz & -iw \end{pmatrix}, \\ A_{21} &= \begin{pmatrix} i\omega^2 x & i\omega y & y \\ -i\omega^2 y & i\omega x & -x \end{pmatrix}, A_{22} = \begin{pmatrix} z & w & -iw \\ -w & z & iz \end{pmatrix}, B_{21} = \begin{pmatrix} y & x \\ -x & y \\ -ix & -iy \end{pmatrix}, B_{22} = \begin{pmatrix} w & z \\ -\omega z & \omega w \\ -i\omega^2 z & -i\omega^2 w \end{pmatrix} \end{aligned}$$

for some choices of basis of the vector spaces $\text{Hom}_{\mathbb{C}G}(V^* \otimes_{\mathbb{C}} \rho, \rho')$.

The commutator group $[T, T] = \langle g_1, g_2 g_1 g_2^{-1} \rangle$ is isomorphic to the quaternion group of order eight, and we can choose 18 generators

$$\bar{w}_{01}, \dots, \bar{w}_{07}, \bar{w}_{11}, \dots, \bar{w}_{15}, \bar{w}_{21}, \dots, \bar{w}_{25}, \bar{w}_3$$

of $\mathbb{C}[V]^{[T, T]}$ such that the associated 20 elements $w_{01}, \dots, w_3, t_1^{-3}, t_2^{-3}$ in $\mathbb{C}[V]^{[T, T]}[t_1^{\pm 1}, t_2^{\pm 1}]$ generate the Cox ring of a crepant (or equivalently symplectic) resolution X of V/T

[Y2, Example 3], [G], where the variables t_1, t_2 correspond to junior elements g_2, g_2^2 respectively.

To state the main result for this section, let us recall that V/T admits two projective symplectic resolutions which are isomorphic as varieties (but not as varieties over V/G). The nef cones in $\text{Pic}(X)_{\mathbb{R}} = \mathbb{R}E_1 \oplus \mathbb{R}E_2$ of these two resolutions are given as $\text{Cone}(-E_1 - E_2, -2E_1 - E_2)$ and $\text{Cone}(-E_1 - E_2, -E_1 - 2E_2)$, and we denote the corresponding resolutions by X_1 and X_2 respectively.

Theorem 5.1. *The symplectic resolution X_1 (resp. X_2) of V/T is isomorphic to the coherent component of the moduli space $\mathfrak{M}_{\theta_{X_1}}$ of T -constellations for certain generic θ_{X_1} (resp. θ_{X_2}). Moreover, the coherent component of the T -Hilbert scheme is isomorphic to the common blow-up of X_1 and X_2 along closed subschemes which are isomorphic to \mathbb{P}^2 .*

In the rest of this section we give a proof of this theorem. We will first show that X_1 is isomorphic to $\mathfrak{M}_{\theta_{X_1}}$ for some θ_{X_1} . The claim for X_2 can be shown similarly using the symmetry of T which we will see below.

Let us consider the action of $\mathbb{C}^* \times \mathbb{C}^*$ on $V = V_1 \oplus V_2$ given by the component-wise dilation. Then $\mathbb{C}^* \times \mathbb{C}^*$ naturally acts on the set of ideals of $\mathbb{C}[V]$ and we can find 13 T -clusters $\mathbb{C}[V]/I$ which are $\mathbb{C}^* \times \mathbb{C}^*$ -invariant as mentioned in [LS]. We describe some of them as T -invariant \mathbb{C} -vector subspaces of $\mathbb{C}[V]$ using the formal variables z_1, z_2 corresponding to the weight for the $\mathbb{C}^* \times \mathbb{C}^*$ -actions as in [LS]:

$$\begin{aligned}
p_1 : & \rho_0 \oplus V_2 z_2 \oplus W z_2^2 \oplus (V_1 \oplus V_2) z_2^3 \oplus (\rho_1 \oplus W) z_2^4 \oplus (V_0 \oplus V_1) z_2^5 \oplus W z_2^6 \oplus V_0 z_2^7 \oplus \rho_2 z_2^8 \\
p_2 : & \rho_0 \oplus V_1 z_1 \oplus V_2 z_2 \oplus W z_1 z_2 \oplus W z_2^2 \oplus V_0 z_1 z_2^2 \oplus (V_1 \oplus V_2) z_2^3 \oplus \rho_2 z_1 z_2^3 \oplus (\rho_1 \oplus W) z_2^4 \oplus V_0 z_2^5 \\
p_{3,1} : & F \oplus (V_2 \quad z_2^3 \oplus \rho_1 \quad z_2^4) \oplus (V_1 z_1 z_2^2 \oplus \rho_2 z_1^2 z_2^2) \\
p_{3,3} : & F \oplus (V_2 \quad z_2^3 \oplus \rho_1 \quad z_2^4) \oplus (V_1 \quad z_2^3 \oplus \rho_2 z_1 z_2^3) \\
p_{4,2} : & F \oplus (V_2 z_1 z_2^2 \oplus \rho_1 z_1^2 z_2^2) \oplus (V_1 z_1^3 \quad \oplus \rho_2 z_1^4 \quad) \\
p_{4,3} : & F \oplus (V_2 z_1 z_2^2 \oplus \rho_1 z_1^2 z_2^2) \oplus (V_1 \quad z_2^3 \oplus \rho_2 z_1 z_2^3) \\
p_{5,1} : & F \oplus (V_2 z_1^3 \quad \oplus \rho_1 z_1^3 z_2) \oplus (V_1 z_1 z_2^2 \oplus \rho_2 z_1^2 z_2^2) \\
p_{5,2} : & F \oplus (V_2 z_1^3 \quad \oplus \rho_1 z_1^3 z_2) \oplus (V_1 z_1^3 \quad \oplus \rho_2 z_1^4 \quad) \\
p_6 : & \rho_0 \oplus V_1 z_1 \oplus V_2 z_2 \oplus W z_1^2 \oplus W z_1 z_2 \oplus (V_1 \oplus V_2) z_1^3 \oplus V_0 z_1^2 z_2 \oplus (\rho_2 \oplus W) z_1^4 \oplus \rho_1 z_1^3 z_2 \oplus V_0 z_1^5 \\
p_7 : & \rho_0 \oplus V_1 z_1 \oplus W z_1^2 \oplus (V_1 \oplus V_2) z_1^3 \oplus (\rho_2 \oplus W) z_1^4 \oplus (V_0 \oplus V_2) z_1^5 \oplus W z_1^6 \oplus V_0 z_1^7 \oplus \rho_1 z_1^8
\end{aligned}$$

where

$$F = \rho_0 \oplus V_1 z_1 \oplus V_2 z_2 \oplus W z_1^2 \oplus W z_1 z_2 \oplus W z_2^2 \oplus V_0 z_1^2 z_2 \oplus V_0 z_1 z_2^2.$$

The corresponding ideals are also presented similarly. For example, the ideal for p_1 is generated by $V_1 z_1, \rho_0 z_2^4, \rho_0 z_2^6, V_2 z_2^9$.

One can give an explicit basis of each direct summand of p_i and $p_{i,j}$ as polynomials using the generic matrices above. For example, a basis of $W z_1 z_2$ in F is given as the three entries of $B_{12} a_{01}$ (or $B_{21} a_{02}$). Note that p_1 (resp. p_2) plays a dual role to p_7 (resp. p_6) with respect to the involutions $\rho_1 \leftrightarrow \rho_2, V_1 \leftrightarrow V_2$ and $z_1 \leftrightarrow z_2$. We have three other $\mathbb{C}^* \times \mathbb{C}^*$ -invariant T -clusters but in fact they do not lie in the coherent component of the T -Hilbert scheme.

We also consider $\mathbb{C}^* \times \mathbb{C}^*$ -invariant T -constellations p_3, p_4, p_5 which are not T -clusters:

$$\begin{aligned}
p_3 : & F \oplus (V_2 \quad z_2^3 \oplus \rho_1 \quad z_2^4) \oplus (V_1 z_1^2 \quad z_2^{-1} \oplus \rho_2 z_1^3 z_2^{-1}) \\
p_4 : & F \oplus (V_2 z_1 z_2^2 \oplus \rho_1 z_1^2 z_2^2) \oplus (V_1 \quad z_2 \quad \oplus \rho_2 z_1 z_2) \\
p_5 : & F \oplus (V_2 z_1^3 \quad \oplus \rho_1 z_1^3 z_2) \oplus (V_1 z_1^{-1} z_2^2 \quad \oplus \rho_2 \quad z_2^2)
\end{aligned}$$

where each p_i is indecomposable as a G -equivariant $\mathbb{C}[V]$ -module, admits a quotient module which is isomorphic to $V_1 \oplus \rho_2$, and the complement of this quotient is generated by ρ_0 . Note that $p_{i,j}$'s ($i = 3, 4, 5$) have submodules which are isomorphic to $V_1 \oplus \rho_2$, in contrast to these p_i 's.

For each i , we consider the orbit cone $C_{p_i} = \{\theta \in \Theta \mid p_i \text{ is } \chi_\theta\text{-semistable}\}$, which is equal to

$$\{\theta = (\theta_\rho)_\rho \in \Theta \mid \sum_{\rho} \theta_\rho \cdot c_\rho \geq 0 \text{ for any submodule } p_i \supset M \cong \bigoplus_{\rho} \rho^{c_\rho}\}. \quad (5.1)$$

by King's result (Proposition 2.1). Each C_{p_i} defines a neighborhood $\tilde{U}_i \subset \mathcal{W} = \mathcal{V} // SL_R$ of p_i consisting of points $p \in \mathcal{W}$ such that $C_p = C_{p_i}$. From the structure of p_i presented above, one can compute C_{p_i} and check that the intersection $\bigcap_{i=1}^7 C_{p_i}$ is full-dimensional. In particular we can take a stability condition θ_{X_1} from its interior.

By [G, Theorem 4.18], the χ -(semi)stable locus of \mathfrak{X} is a union of seven affine open subsets U_1, \dots, U_7 :

$$U_1 = \{w_{12}t_1^{-3} \neq 0\}, U_2 = \{w_{12}w_{23} \neq 0\}, U_3 = \{w_{12}w_3 \neq 0\}, U_4 = \{w_{11}w_3 \neq 0\}, \\ U_5 = \{w_{13}w_3 \neq 0\}, U_6 = \{w_{13}w_{22} \neq 0\}, U_7 = \{w_{22}t_2^{-3} \neq 0\}.$$

We show that the quotient of \tilde{U}_i by the torus T_Θ is isomorphic to U_i and that p_i corresponds to the origin of $U_i \cong \mathbb{C}^4$ via this isomorphism. For this, we define a ring map $\varphi_{X_1} : \mathbb{C}[\mathcal{W}] \rightarrow \mathbb{C}[V]^{[T,T]}[t_1^{\pm 1}, t_2^{\pm 1}]$ by sending a semi-invariant h of weight (c_0, \dots, c_6) to $\varphi(h)t_1^{c_1+c_4+c_5+c_6}t_2^{c_2+c_3+c_5+c_6}$ where the ordering for $\text{Irr}(G)$ is given by $\rho_0, \rho_1, \rho_2, V_0, V_1, V_2, W$.

We will construct an isomorphism $U_i \cong \tilde{U}_i // T_\Theta$ for $i = 1$. The other cases can be done in a similar way. We choose determinantal semi-invariants $\{h_i\}_i$ such that

- (1) their weights in Θ generate a full-dimensional cone (contained in C_{p_1}),
- (2) they do not vanish at p_1 , and
- (3) each $\varphi(h_i)$ is equal to $\bar{w}_{12} = z^4 + (-4\omega - 2)z^2w^2 + w^4$ up to scalar multiplication.

For instance, $\{h_i\}_i$ can be chosen as the determinants of the following six matrices:

$$A_{02}B_{12}, (b_{02} \quad b_{02}A_{02}B_{02}), (b_{21} \quad b_{21}A_{22}B_{22}), \begin{pmatrix} a_{10} \\ A_{02}B_{02}a_{10} \end{pmatrix}, \begin{pmatrix} a_{02} \\ A_{22}B_{22}a_{02} \end{pmatrix}, (b_{21}A_{22} \quad A_{02}).$$

Remark 5.1. One can find h_i 's as semi-invariants whose weights are part of generators of the cone C_{p_1} . \square

Since each $\varphi(h_i)$ is of the form $c\bar{w}_{12}$ ($c \in \mathbb{C}^*$), φ_{X_1} induces a homomorphism

$$(\mathbb{C}[\mathcal{W}]_{h_1 \dots h_6})^{T_\Theta} \rightarrow (\mathbb{C}[V]^{[T,T]}[t_1^{\pm 1}, t_2^{\pm 1}]_{w_{12}})^{T_X}$$

between the torus-invariant parts of the localizations. The latter ring contains $H^0(\mathcal{O}_{U_1}) = \mathbb{C}[w_{02}, w_{05}, \frac{w_{23}}{w_{12}^2 t_1^{-3}}, \frac{w_{24}}{w_{12}^2 t_1^{-3}}] \subset \mathbb{C}(V)^T$ [G, Theorem 4.18], and we can show that the above homomorphism is an isomorphism onto $H^0(\mathcal{O}_{U_1})$ by constructing its inverse by

$$w_{02} \mapsto \det(b_{10}A_{12}B_{12}A_{12}B_{22}a_{02}), w_{05} \mapsto \det(b_{10}A_{12}B_{22}a_{02}), \\ \frac{w_{23}}{w_{12}^2 t_1^{-3}} \mapsto \frac{\det(b_{12} \quad b_{02}A_{02}B_{12})}{h_2 h_4}, \frac{w_{24}}{w_{12}^2 t_1^{-3}} \mapsto \frac{\det(b_{12}A_{02}B_{11} \quad b_{02}A_{02}B_{12})}{h_2 h_4}$$

where

$$\begin{aligned}\bar{w}_{02} &= z^5 w - z w^5, \quad \bar{w}_{05} = z^4 + (4\omega + 2)z^2 w^2 + w^4, \\ \bar{w}_{23} &= xz^3 + (2\omega + 1)yz^2 w + (2\omega + 1)xz w^2 + yw^3, \quad \bar{w}_{24} = yz^5 + 5xz^4 w - 5yz w^4 - xw^5.\end{aligned}$$

One can check that the elements of the both sides of this assignment indeed give the same rational functions on V/T . This implies that the open subset $\{h_1 \cdots h_6 \neq 0\} \subset \mathcal{W}$ descends to an open subset of the moduli space $\mathfrak{M}_{\theta_{X_1}}$ which is isomorphic to U_1 . Note that here we use the assumption that the weights of h_i 's generate a full-dimensional cone.

In this way we obtain open subsets of $\mathfrak{M}_{\theta_{X_1}}$ which are isomorphic to U_i for all i . Thus we can regard X_1 as an open subset of $\mathfrak{M}_{\theta_{X_1}}$. Since X_1 is already proper over V/T , we obtain $X_1 \cong \mathfrak{M}_{\theta_{X_1}}$.

Finally, we show that the coherent component of the T -Hilbert scheme is isomorphic to the blowup \tilde{X} of X_1 along a closed subscheme $P \cong \mathbb{P}^2$. The center $P \subset X_1$ is defined as the zero locus of all the coordinates w_{01}, \dots, w_{25} but $w_{11}, w_{12}, w_{13}, w_3$ [G, Proposition 4.11], and is covered with U_3, U_4 and U_5 . Inside

$$U_3 = \text{Spec } \mathbb{C} \left[\frac{w_{11}}{w_{12}}, \frac{w_{13}}{w_{12}}, \frac{w_{12}w_{23}}{w_3}, \frac{w_{12}w_{21}}{w_3} \right],$$

for instance, P is the zero locus of the last two coordinates, and the blowup along $P \cap U_3$ has the following affine open covering

$$U_{3,1} = \text{Spec } \mathbb{C} \left[\frac{w_{11}}{w_{12}}, \frac{w_{13}}{w_{12}}, \frac{w_{23}}{w_{21}}, \frac{w_{12}w_{21}}{w_3} \right], \quad U_{3,3} = \text{Spec } \mathbb{C} \left[\frac{w_{11}}{w_{12}}, \frac{w_{13}}{w_{12}}, \frac{w_{12}w_{23}}{w_3}, \frac{w_{21}}{w_{23}} \right].$$

Similar to the above argument, we can construct an isomorphism between a neighborhood of $p_{3,1}$ (resp. $p_{3,3}$) and $U_{3,1}$ (resp. $U_{3,3}$). Applying the same construction to U_4 and U_5 as well, we finally obtain the claim. \square

References

- [AH] A. A'Campo-Neuen and J. Hausen, Quotients of toric varieties by the action of a subtorus, *Tohoku Math. J. (2)* **51** (1999), no. 1, 1–12.
- [ADHL] I. Arzhantsev, U. Derenthal, J. Hausen and A. Laface, *Cox rings*, Cambridge Studies in Advanced Mathematics, 144, Cambridge University Press, Cambridge, 2015.
- [AG] I. V. Arzhantsev and S. A. Gařfullin, *Sb. Math.* **201** (2010), no. 1-2, 1–21; translated from *Mat. Sb.* **201** (2010), no. 1, 3–24.
- [B] G. Bellamy, On singular Calogero-Moser spaces, *Bull. Lond. Math. Soc.* **41** (2009), no. 2, 315–326.
- [BH] F. BERTHOLD and J. Hausen, GIT equivalence beyond the ample cone, *Michigan Math. J.* **54** (2006), no. 3, 483–515.

- [BK] R. V. Bezrukavnikov and D. B. Kaledin, Proc. Steklov Inst. Math. **2004**, no. 3(246), 13–33; translated from Tr. Mat. Inst. Steklova **246** (2004), Algebr. Geom. Metody, Svyazi i Prilozh., 20–42.
- [BCHM] C. Birkar et al., Existence of minimal models for varieties of log general type, J. Amer. Math. Soc. **23** (2010), no. 2, 405–468.
- [BTW] R. Bocklandt, T. Schedler and M. Wemyss, Superpotentials and higher order derivations, J. Pure Appl. Algebra **214** (2010), no. 9, 1501–1522.
- [BKR] T. Bridgeland, A. King and M. Reid, The McKay correspondence as an equivalence of derived categories, J. Amer. Math. Soc. **14** (2001), no. 3, 535–554.
- [CCL] S. Cautis, A. Craw and T. Logvinenko, Derived Reid’s recipe for abelian subgroups of $SL_3(\mathbb{C})$, J. Reine Angew. Math. **727** (2017), 1–48.
- [Co] D. A. Cox, The homogeneous coordinate ring of a toric variety, J. Algebraic Geom. **4** (1995), no. 1, 17–50.
- [CLS] D. A. Cox, J. B. Little and H. K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, 124, American Mathematical Society, Providence, RI, 2011.
- [Cr] A. Craw, An explicit construction of the McKay correspondence for A -Hilb \mathbb{C}^3 , J. Algebra **285** (2005), no. 2, 682–705.
- [CI] A. Craw and A. Ishii, Flops of G -Hilb and equivalences of derived categories by variation of GIT quotient, Duke Math. J. **124** (2004), no. 2, 259–307.
- [CMT] A. Craw, D. Maclagan and R. R. Thomas, Moduli of McKay quiver representations. I. The coherent component, Proc. Lond. Math. Soc. (3) **95** (2007), no. 1, 179–198.
- [CR] A. Craw and M. Reid, How to calculate A -Hilb \mathbb{C}^3 , in *Geometry of toric varieties*, 129–154, Sémin. Congr., 6, Soc. Math. France, Paris.
- [DeK] H. Derksen and G. Kemper, *Computational invariant theory*, second enlarged edition, Encyclopaedia of Mathematical Sciences, 130, Springer, Heidelberg, 2015.
- [DK] M. Donten-Bury and S. Keicher, Computing resolutions of quotient singularities, J. Algebra **472** (2017), 546–572.
- [DeW1] H. Derksen and J. Weyman, Semi-invariants of quivers and saturation for Littlewood-Richardson coefficients, J. Amer. Math. Soc. **13** (2000), no. 3, 467–479.
- [DeW2] H. Derksen and J. Weyman, *An introduction to quiver representations*, Graduate Studies in Mathematics, 184, American Mathematical Society, Providence, RI, 2017.
- [DZ] M. Domokos and A. N. Zubkov, Semi-invariants of quivers as determinants, Transform. Groups **6** (2001), no. 1, 9–24.

- [D] M. Donten-Bury, Cox rings of minimal resolutions of surface quotient singularities, *Glasg. Math. J.* **58** (2016), no. 2, 325–355.
- [DG] M. Donten-Bury and M. Grab, Crepant resolutions of 3-dimensional quotient singularities via Cox rings, *Exp. Math.* **28** (2019), no. 2, 161–180.
- [DW] M. Donten-Bury and J. A. Wiśniewski, On 81 symplectic resolutions of a 4-dimensional quotient by a group of order 32, *Kyoto J. Math.* **57** (2017), no. 2, 395–434.
- [G] M. Grab, Cox rings and symplectic quotient singularities with torus action, arXiv:1807.11438v2.
- [GS] D. Grayson and M. Stillman, Macaulay 2: A software system for research in algebraic geometry; available at <http://www.math.uiuc.edu/Macaulay2>.
- [Ha] M. Haiman, Combinatorics, symmetric functions, and Hilbert schemes, in *Current developments in mathematics, 2002*, 39–111, Int. Press, Somerville, MA.
- [Hu] Y. Hu, Combinatorics and quotients of toric varieties, *Discrete Comput. Geom.* **28** (2002), no. 2, 151–174.
- [HK] Y. Hu and S. Keel, Mori dream spaces and GIT, *Michigan Math. J.* **48** (2000), 331–348.
- [IINdC] A. Ishii, Y. Ito and Á. Nolla de Celis, On G/N -Hilb of N -Hilb, *Kyoto J. Math.* **53** (2013), no. 1, 91–130.
- [IN] Y. Ito and H. Nakajima, McKay correspondence and Hilbert schemes in dimension three, *Topology* **39** (2000), no. 6, 1155–1191.
- [IR] Y. Ito and M. Reid, The McKay correspondence for finite subgroups of $SL(3, \mathbf{C})$, in *Higher-dimensional complex varieties (Trento, 1994)*, 221–240, de Gruyter, Berlin.
- [Kin] A. D. King, Moduli of representations of finite-dimensional algebras, *Quart. J. Math. Oxford Ser. (2)* **45** (1994), no. 180, 515–530.
- [Kir] A. Kirillov, Jr., *Quiver representations and quiver varieties*, Graduate Studies in Mathematics, 174, American Mathematical Society, Providence, RI, 2016.
- [KSZ] M. M. Kapranov, B. Sturmfels and A. V. Zelevinsky, Quotients of toric varieties, *Math. Ann.* **290** (1991), no. 4, 643–655.
- [LS] M. Lehn and C. Sorger, A symplectic resolution for the binary tetrahedral group, in *Geometric methods in representation theory. II*, 429–435, Sémin. Congr., 24-II, Soc. Math. France, Paris.
- [Le] B. Leng. The McKay correspondence and orbifold Riemann-Roch. PhD thesis, University of Warwick, 2002.

- [Lo] Logvinenko, Timothy Natural G -constellation families. *Doc. Math.* **13** (2008), 803–823.
- [M] M. Maruyama, On a family of algebraic vector bundles, in *Number theory, algebraic geometry and commutative algebra, in honor of Yasuo Akizuki*, 95–146, Kinokuniya, Tokyo.
- [MFK] D. Mumford, J. Fogarty and F. Kirwan, *Geometric invariant theory*, third edition, *Ergebnisse der Mathematik und ihrer Grenzgebiete (2)*, **34**, Springer-Verlag, Berlin, 1994.
- [N] Namikawa, Poisson deformations of affine symplectic varieties, *Duke Math. J.* **156** (2011), no. 1, 51–85.
- [Re1] M. Reid, Canonical 3-folds, in *Journées de Géométrie Algébrique d’Angers, Juillet 1979/Algebraic Geometry, Angers, 1979*, 273–310, Sijthoff & Noordhoff, Alphen aan den Rijn.
- [Re2] M. Reid. McKay correspondence. *Proc. of Algebraic Geom. Symposium (Kinosaki, November 1996)*, pages 14–41, 1997.
- [Ro] M. Rossi, Embedding non-projective Mori dream space, *Geom. Dedicata* **207** (2020), 355–393.
- [SV] A. Schofield and M. van den Bergh, Semi-invariants of quivers for arbitrary dimension vectors, *Indag. Math. (N.S.)* **12** (2001), no. 1, 125–138.
- [T] M. Thaddeus, Geometric invariant theory and flips, *J. Amer. Math. Soc.* **9** (1996), no. 3, 691–723.
- [W] K. Watanabe, Certain invariant subrings are Gorenstein. I, II, *Osaka Math. J.* **11** (1974), 1–8; *ibid.* **11** (1974), 379–388.
- [Y1] R. Yamagishi, On smoothness of minimal models of quotient singularities by finite subgroups of $SL_n(\mathbb{C})$, *Glasg. Math. J.* **60** (2018), no. 3, 603–634.
- [Y2] R. Yamagishi, Moduli of G -constellations and crepant resolutions I, arXiv:2209.11900, to appear in *Advanced Studies in Pure Mathematics*.

National Center for Theoretical Sciences (NCTS), National Taiwan University, Taipei,
Taiwan

Email address: yamagishi@ncts.ntu.edu.tw