

Universal Densities Exist for Every Finite Reference Measure

Łukasz Dębowski*

Abstract

As it is known, universal codes, which estimate the entropy rate consistently, exist for any stationary ergodic source over a finite alphabet but not over a countably infinite one. We cast the problem of universal coding into the problem of universal densities with respect to a given reference measure on a countably generated measurable space, examples being the counting measure or the Lebesgue measure. We show that universal densities, which estimate the differential entropy rate consistently, exist if the reference measure is finite, which disproves that the assumption of a finite alphabet is necessary in general. To exhibit a universal density, we combine the prediction by partial matching (PPM) code with the non-parametric differential (NPD) entropy rate estimator, putting a prior both over all Markov orders and all quantization levels. The proof of universality applies Barron's asymptotic equipartition for densities and continuity of f -divergences for filtrations. As an application, we demonstrate that any universal density induces a strongly consistent Cesàro mean estimator of the conditional density given an infinite past, which solves the problem of universal prediction with the $0-1$ loss for a countable alphabet, by the way. We also show that there exists a strongly consistent entropy rate estimator with respect to the Lebesgue measure in the class of stationary ergodic Gaussian processes.

Keywords: stationary ergodic processes; entropy rate; universal coding; prediction by partial matching; quantization; density estimation; universal prediction

MSC 2000: 94A29, 62M20

*Ł. Dębowski is with the Institute of Computer Science, Polish Academy of Sciences, ul. Jana Kazimierza 5, 01-248 Warszawa, Poland (e-mail: ldebowski@ipipan.waw.pl).

1 Introduction

Consider the family of stationary ergodic measures over a given alphabet. It is known that universal measures, i.e., those consistently estimating the entropy rate in the almost sure sense and in expectation, respectively, exist for any finite alphabet. A simple example thereof is the PPM (prediction by partial matching) measure, also called the R -measure, constructed gradually by Cleary and Witten and by Ryabko [10, 27, 26].

Universal measures are important for many reasons. They matter not only in data compression but also in non-parametric statistical inference. The Shannon-Fano code taken with respect to a universal measure is an instance of a universal code. Other important instances of universal codes were discovered in [30, 24, 21]. Universal measures induce universal predictors with the 0 – 1 loss under mild conditions [15], see also [23]. Universal measures are an important building block also in estimation of the Markov order and the hidden Markov order [22, 31] as well as bounding mutual information [14].

It is known, alas, that universal measures or codes do not exist for a countably infinite alphabet [20, 18, 29]. It may seem that the assumption of a finite alphabet is necessary in general. In this paper, we disprove this hypothesis by casting the problem of universal measures into universal densities, i.e., Radon-Nikodym derivatives with respect to a given reference measure. In doing so, we draw ideas from the Shannon-McMillan-Breiman (SMB) theorem for densities by Barron [5]. In this setting, we show that universal densities exist if the reference measure is finite, such as the Lebesgue measure on a finite interval. Moreover, inspired by work [18], we show that if a universal density exists then it induces a strongly consistent Cesàro mean estimator of the conditional density given an infinite past, in the total variation, as it follows by the Pinsker inequality. Any such conditional density estimator solves also the problem of universal prediction with the 0 – 1 loss for a countable alphabet, see [15].

We prove the existence of universal densities for a finite reference measure in a constructive way. To produce a universal density, we combine the construction of the known PPM measure [10, 27, 26] with the NPD (non-parametric differential) entropy rate estimator, a recent simple idea of quantization by Feutrill and Roughan [17]. Simply speaking, we put a prior over all quantization levels, which is analogous to putting a prior over all Markov orders in the PPM measure [27, 26]. We note that the constructed universal density is defined as an infinite series. It is finite only almost everywhere. The proof of universality itself rests on the SMB theorem for densities by Barron [5] and the monotone convergence of f -divergences for filtrations, see e.g. [13].

Thus, it is not finiteness of the alphabet but rather finiteness of the reference measure that allows for universal densities. Non-existence of universal codes for a countably infinite alphabet is tightly connected to non-existence of a uniform probability measure over the set of natural numbers.

The organization of the paper is as follows. In Section 2, we amply recall

preliminaries to keep the paper relatively self-contained. Section 3 contains the main results. The first one is the construction of the universal density with respect to a finite measure (Section 3.1). Drawing the inspiration from [17] and [27, 26], we call it simply the total NPD density. Having an instance of a universal density, we construct a consistent Cesàro mean estimator of the conditional density given an infinite past, drawing the idea from work [18] (Section 3.2). Using techniques detailed in work [15], the consistent estimator of the conditional density can be turned into a universal predictor in the countable alphabet setting (Section 3.3). As some further examples, we present applications of the total NPD density to the subclasses of processes over a countably infinite alphabet (Section 3.4) and on the real line (Section 3.5). We easily name some sufficient conditions that allow for consistent estimation of the entropy rate with respect to infinite reference measures in these domains, see also [29]. In particular, as we show, there exists a strongly consistent entropy rate estimator with respect to the Lebesgue measure in the class of stationary ergodic Gaussian processes.

2 Preliminaries

In this section we establish our setting and report the received knowledge. We discuss the introductory material such as the entropy rate, the SMB theorem, also called the asymptotic equipartition, the definition of universal measures, and the universal PPM measure for a finite alphabet.

2.1 General setting

Let $(\mathbb{X}, \mathcal{X}, \mu)$ be a countably generated measurable space with a σ -finite measure μ on it. Measure μ will be called the reference measure. Simple familiar examples are the counting measure $\mu(A) = \gamma(A) := \text{card } A$ for a countable alphabet \mathbb{X} or the Lebesgue measure $\mu([a, b]) = \lambda([a, b]) := b - a$ for $\mathbb{X} = \mathbb{R}$. Consider the product space $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$ and put random variables $X_k : \mathbb{X}^{\mathbb{Z}} \ni (x_i)_{i \in \mathbb{Z}} \mapsto x_k \in \mathbb{X}$. We write the tuples of points as $x_{j:k} := (x_j, x_{j+1}, \dots, x_k)$. For a probability measure R on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$, we denote its finite-dimensional restrictions $R_n(A) := R(X_{1:n} \in A)$ and if $R_n \ll \mu^n$ then we write the densities

$$R_\mu(x_{1:n}) := \frac{dR_n}{d\mu^n}(x_{1:n}). \quad (1)$$

The space of stationary ergodic measures on $(\mathbb{X}^{\mathbb{Z}}, \mathcal{X}^{\mathbb{Z}})$ with respect to the shift operation will be denoted as \mathbb{E} . A measure $P \in \mathbb{E}$ where $P_n \ll \mu^n$ is called a memoryless source if $P_\mu(x_{1:n}) = \prod_{i=1}^n P_\mu(x_i)$.

Fix a measure $P \in \mathbb{E}$ where $P_n \ll \mu^n$. The expectation operator \mathbf{E} and the quantifier “almost surely” (a.s.) will be taken throughout with respect

to P . Throughout the paper, symbol \log denotes the logarithm to a fixed underspecified base. We define the block entropy

$$h_\mu(n) := \mathbf{E} [-\log P_\mu(X_{1:n})] = - \int P_\mu(x_{1:n}) \log P_\mu(x_{1:n}) d\mu^n(x_{1:n}). \quad (2)$$

A short notice, if the reference measure μ is the counting measure then $h_\mu(n) \geq 0$ since $P_\mu(x_{1:n}) \leq 1$. In contrast, if μ is a probability measure then $h_\mu(n) \leq 0$ since $-h_\mu(n)$ is the Kullback-Leibler divergence between P_n and μ^n .

In any case, by stationarity, the block entropy is subadditive, $h_\mu(n+m) \leq h_\mu(n) + h_\mu(m)$. In particular, $nh_\mu(n+1) - (n+1)h_\mu(n) \leq nh_\mu(1) - h_\mu(n) \leq 0$. Hence sequence $h_\mu(n)/n$ is decreasing and there exists the entropy rate

$$h_\mu := \lim_{n \rightarrow \infty} \frac{h_\mu(n)}{n} = \inf_{n \geq 1} \frac{h_\mu(n)}{n}, \quad (3)$$

which, by the way, constitutes a simpler proof of the known Fekete lemma for subadditive sequences [16] than reported for instance in [13, Theorem 5.11].

Moreover, by the Shannon-McMillan-Breiman (SMB) theorem, we have the asymptotic equipartition property, namely,

$$\lim_{n \rightarrow \infty} [-\log P_\mu(X_{1:n})] / n = h_\mu \text{ a.s.}, \quad (4)$$

as consecutively noticed by Shannon [28] for memoryless sources and generalized later in [7, 9, 5, 2] to other settings. Consider the class of stationary ergodic measures with a finite entropy rate,

$$\mathbb{E}(\mu) := \{P \in \mathbb{E} : P_n \ll \mu^n \text{ and } |h_\mu| < \infty\}. \quad (5)$$

Let us also denote the conditional density

$$P_\mu(x_{n+1:n+m} | x_{1:n}) := \frac{P_\mu(x_{1:n+m})}{P_\mu(x_{1:n})}. \quad (6)$$

As shown by Barron [5, Theorem 1], we have the asymptotic equipartition (4) for an arbitrary $P \in \mathbb{E}(\mu)$. This fact is a consequence of the Breiman ergodic theorem [7], [1, Theorem 12(c)] since, as shown by Barron [5, Proof of Theorem 1], for $P \in \mathbb{E}(\mu)$ there exists the limit of conditional densities

$$P_\mu(X_0 | X_{-\infty:-1}) := \lim_{n \rightarrow \infty} P_\mu(X_0 | X_{-n:-1}) \text{ a.s.} \quad (7)$$

and $\mathbf{E} \sup_{n \in \mathbb{N}} |\log P_\mu(X_0 | X_{-n:-1})| < \infty$, whereas

$$h_\mu = \mathbf{E} [-\log P_\mu(X_0 | X_{-\infty:-1})]. \quad (8)$$

Obviously, the asymptotic equipartition is an interesting property from a statistical perspective since it suggests that the entropy rate can be estimated, which is known to be the case for a finite counting reference measure. However,

the problem of directly applying statement (4) is that in order to estimate the entropy rate, we need to know the unknown probability measure.

To approach the problem of entropy estimation in a reasonable way, let us consider another probability measure R where $R_n \ll \mu^n$, which need not be stationary or ergodic. As it is a part of an older information-theoretic folklore, for any $m > 0$ we have an inequality, called sometimes the Barron inequality,

$$P \left(\log \frac{dP_n}{dR_n}(X_{1:n}) \leq -\log m \right) \leq \frac{1}{m}, \quad (9)$$

shown for example in [6, Theorem 3.1] or applied implicitly much earlier in [8]. Since $dP_n/dR_n(x_{1:n}) = P_\mu(x_{1:n})/R_\mu(x_{1:n})$, as a result, by an easy application of the Borel-Cantelli lemma, we obtain

$$\liminf_{n \rightarrow \infty} [-\log R_\mu(X_{1:n})] / n \geq h_\mu \text{ a.s.} \quad (10)$$

Since the Kullback-Leibler divergence is non-negative in general,

$$D(P_n || R_n) := \mathbf{E} \left[\log \frac{dP_n}{dR_n}(X_{1:n}) \right] = \mathbf{E} \left[\log \frac{P_\mu(X_{1:n})}{R_\mu(X_{1:n})} \right] \geq 0, \quad (11)$$

we also have a similar result in expectation:

$$\liminf_{n \rightarrow \infty} \mathbf{E} [-\log R_\mu(X_{1:n})] / n \geq h_\mu \text{ a.s.} \quad (12)$$

Consequently, we will consider two definitions of a universal measure. By definition, such a measure can be used for estimating the entropy rate of an unknown stationary ergodic process.

Definition 1. *A probability measure R where $R_n \ll \mu^n$ is called strongly universal with respect to a reference measure μ if for every measure $P \in \mathbb{E}(\mu)$ we have*

$$\lim_{n \rightarrow \infty} [-\log R_\mu(X_{1:n})] / n = h_\mu \text{ a.s.} \quad (13)$$

Definition 2. *A probability measure R where $R_n \ll \mu^n$ is called universal in expectation with respect to a reference measure μ if for every measure $P \in \mathbb{E}(\mu)$ we have*

$$\lim_{n \rightarrow \infty} \mathbf{E} [-\log R_\mu(X_{1:n})] / n = h_\mu. \quad (14)$$

If the reference measure is the counting measure over a countable alphabet \mathbb{X} , i.e., $\mu(A) = \gamma(A) := \text{card } A$ for $A \subset \mathbb{X}$, then we speak of measures that are universal with respect to alphabet \mathbb{X} , respectively. In this case, we drop the subscript γ : $P_\gamma(x) \rightarrow P(x)$, $R_\gamma(x) \rightarrow R(x)$, and $h_\gamma \rightarrow h$.

2.2 Finite alphabet

For a finite alphabet, universal measures are known to exist. We recall an example of a universal measure for this case, called the PPM (prediction by partial matching). The PPM measure was discovered gradually. Cleary and Witten [10] considered roughly Markov approximations PPM_k defined in (15) and coined name PPM, which we prefer as more distinctive. Ryabko [27, 26] considered the infinite series PPM defined in (17), called it the R -measure, and proved that it is universal. To be precise, Ryabko used the Krichevsky-Trofimov smoothing $(+1/2)$ rather than the Laplace smoothing $(+1)$ applied in (15). This difference is minor and does not affect universality.

The definition that we use is as follows.

Definition 3 (PPM density). *Let the alphabet be $\mathbb{X} = \{a_1, \dots, a_D\}$. Adapting the definitions from works [10, 27, 26, 12], the PPM density of order $k \geq 0$ is defined as*

$$\text{PPM}_k^D(x_{1:n}) := \begin{cases} D^{-k-1} \prod_{i=k+2}^n \frac{N(x_{i-k:i}|x_{1:i-1}) + 1}{N(x_{i-k:i-1}|x_{1:i-2}) + D}, & k \leq n-2, \\ D^{-n}, & k \geq n-1, \end{cases} \quad (15)$$

where the frequency of a substring $w_{1:k}$ in a string $x_{1:n}$ is

$$N(w_{1:k}|x_{1:n}) := \sum_{i=1}^{n-k+1} \mathbf{1}\{x_{i:i+k-1} = w_{1:k}\}. \quad (16)$$

Subsequently, we define the (total) PPM density as

$$\text{PPM}^D(x_{1:n}) := \sum_{k=0}^{\infty} w_k \text{PPM}_k^D(x_{1:n}), \quad w_k := \frac{1}{k+1} - \frac{1}{k+2}. \quad (17)$$

Infinite series (17) is finite and computable in the sense of computability theory since we have

$$\text{PPM}_k^D(x_{1:n}) = D^{-n} \text{ for } k \geq L(x_{1:n}), \quad (18)$$

where $L(x_{1:n})$ is the maximal length of a repetition in $x_{1:n}$,

$$L(x_{1:n}) := \max \{k \geq 0 : x_{i+1:i+k} = x_{j+1:j+k} \text{ for some } 0 \leq i < j \leq n-k\}. \quad (19)$$

Theorem 1. *There exists a measure R such that $R(x_{1:n}) = \text{PPM}^D(x_{1:n})$.*

Remark: This measure R will be denoted PPM^D .

Proof. By the Kolmogorov process theorem, it suffices to show that

$$\sum_{x_{n+1} \in \mathbb{X}} \text{PPM}^D(x_{1:n+1}) = \text{PPM}^D(x_{1:n}). \quad (20)$$

But this follows by the monotone convergence from

$$\sum_{x_{n+1} \in \mathbb{X}} \text{PPM}_k^D(x_{1:n+1}) = \text{PPM}_k^D(x_{1:n}), \quad (21)$$

which in turn follows by the definition of $\text{PPM}_k^D(x_{1:n})$. \square

The strong universality of the total PPM measure follows by the Stirling approximation and by the Birkhoff ergodic theorem. Hence, by $\text{PPM}^D(x_{1:n}) \geq [\frac{1}{n} - \frac{1}{n+1}] \text{PPM}_{n-1}^D(x_{1:n}) \geq D^{-n}/(n+1)^2$ and by the dominated convergence follows also the universality in expectation.

Theorem 2 ([27, 26, 12]). *Measure PPM^D is universal, both strongly and in expectation, with respect to alphabet $\mathbb{X} = \{a_1, \dots, a_D\}$.*

3 Main results

In this section, we will exhibit more novel results. In particular, we will show that universal measures exist if the reference measure is an arbitrary finite measure. Our constructive example of a universal measure is called the NPD measure and bears to the quantization idea by Feutrill and Roughan [17], who carried out the construction half-way. We also show that if a universal measure exists then we may construct a strongly consistent Cesàro mean estimator for the limiting conditional density given an infinite past, see [18]. Consequently, we show that any strongly consistent estimator of this conditional density yields a universal predictor for a countable alphabet, compare [15]. Moreover, as some supplementary examples, we will scale down the NPD measure to two particular cases where consistent estimation of the entropy rate is not possible in general, namely, the countably infinite alphabet and the real line. This general infeasibility will become intuitive by virtue of the additional assumptions that we are bound to make, see also [29]. Once these conditions are satisfied, the corrected NPD estimator is strongly consistent.

3.1 Finite reference measure

Let the reference measure be an arbitrary finite measure. We will demonstrate universality of the following constructive object.

Definition 4 (NPD density). *Let $(\mathbb{X}, \mathcal{X}, \mu)$ be a countably generated finite measure space. Let $\mathcal{X}_l \uparrow \mathcal{X}$, where $l = 0, 1, 2, \dots$, be a filtration where σ -fields \mathcal{X}_l are finite with $\mathcal{X}_0 = \{\mathbb{X}, \emptyset\}$. Such a filtration exists since \mathcal{X} is countably*

generated. Let χ_l be the finite partitions that generate σ -fields \mathcal{X}_l respectively. We treat sets χ_l as finite alphabets of symbols $A \in \chi_l$. We put $D_l := \text{card } \chi_l$ and we introduce quantizations of points $x \in \mathbb{X}$ as symbols $x^l := A$ for $x \in A \in \chi_l$. We define the NPD density of order $l \geq 0$ as

$$\text{NPD}_\mu^l(x_{1:n}) := \frac{\text{PPM}^{D_l}(x_{1:n}^l)}{\prod_{i=1}^n \mu(x_i^l)}. \quad (22)$$

Subsequently, we define the (total) NPD density as

$$\text{NPD}_\mu(x_{1:n}) := \sum_{l=0}^{\infty} w_l \text{NPD}_\mu^l(x_{1:n}), \quad w_l := \frac{1}{l+1} - \frac{1}{l+2}. \quad (23)$$

In particular, we observe that $\text{NPD}_\mu(x_{1:n}) \geq \frac{1}{2} \text{NPD}_0^\mu(x_{1:n}) = \frac{1}{2}$.

Actually, for the universality of the NPD density, it does not matter which universal measure for a finite alphabet we use in definition (22). Densities (22) for the Lebesgue measure and partitions of the real line into intervals of an equal length were considered by Feutrill and Roughan [17] under the name of the non-parametric differential (NPD) entropy rate estimator. The total density (23) was not considered in [17]. There is some analogy between the PPM series (17) and the NPD series (23). Weights w_l in series (23) weigh different quantization levels, whereas weights w_k in series (17) weigh different Markov approximations.

The total NPD density is measurable and finite μ -almost everywhere, as it follows by the monotone convergence. Just an explicit proof for a sanity check.

Theorem 3. *We have $\text{NPD}_\mu(x_{1:n}) < \infty$ for μ^n -almost all $x_{1:n}$.*

Proof. For each $l \geq 0$, we have

$$\int \text{NPD}_\mu^l(x_{1:n}) d\mu^n(x_{1:n}) = \sum_{x_{1:n}^l} \text{PPM}^{D_l}(x_{1:n}^l) = 1. \quad (24)$$

Since $\text{NPD}_\mu^l \geq 0$, hence by the monotone convergence, we obtain

$$\int \text{NPD}_\mu(x_{1:n}) d\mu^n(x_{1:n}) = \sum_{l=0}^{\infty} w_l \int \text{NPD}_\mu^l(x_{1:n}) d\mu^n(x_{1:n}) = 1. \quad (25)$$

Since the integral is finite, the integrand is finite almost surely. \square

However, the series defining the total NPD density can be divergent for particular tuples $x_{1:n}$. In general, we have

$$\text{NPD}_\mu^l(x_{1:n}) = \frac{D_l^{-n}}{\prod_{i=1}^n \mu(x_i^l)} \text{ for } l \geq M(x_{1:n}), \quad (26)$$

where $M(x_{1:n})$ is the minimal quantization level that puts points x_1, \dots, x_n into different bins,

$$M(x_{1:n}) := \min \{l \geq 0 : L(x_{1:n}^l) = 0\}. \quad (27)$$

A particularly regular case arises for uniformly dyadic partitions:

$$D_l = 2^l \text{ and } \mu(x_i^l) = 2^{-l}. \quad (28)$$

Such partitioning is feasible if the reference measure μ is a non-atomic probability measure, such as the normal distribution $N(m, \sigma^2)$ to be discussed in Section 3.5. Then $\text{NPD}_\mu^l(x_{1:n}) = 1$ for $l \geq M(x_{1:n})$ and series $\text{NPD}_\mu(x_{1:n})$ is finite if $M(x_{1:n})$ is finite, whereas statistic $M(X_{1:n})$ is finite $\mu^{\mathbb{Z}}$ -almost surely.

Universality was stated in Definitions 1 and 2 as a property of measures rather than their densities. First, let us see the following statement.

Theorem 4. *There exists a measure R such that $R_\mu(x_{1:n}) = \text{NPD}_\mu(x_{1:n})$.*

Remark: This measure R will be denoted NPD.

Proof. We may construct measures

$$R_n(A) := \int_{x_{1:n} \in A} \text{NPD}_\mu(x_{1:n}) d\mu^n(x_{1:n}). \quad (29)$$

To show that measures R_n induce measure R on infinite sequences, by the Kolmogorov process theorem, it suffices to show that $R_{n+1}(A \times \mathbb{X}) = R_n(A)$. In turn, using the Fubini theorem this is implied by condition

$$\int_{x_{n+1} \in \mathbb{X}} \text{NPD}_\mu(x_{1:n+1}) d\mu(x_{n+1}) = \text{NPD}_\mu(x_{1:n}), \quad (30)$$

The above follows by the monotone convergence from

$$\int_{x_{n+1} \in \mathbb{X}} \text{NPD}_\mu^l(x_{1:n+1}) d\mu(x_{n+1}) = \text{NPD}_\mu^l(x_{1:n}), \quad (31)$$

which is true since PPM^{D_l} is a measure. \square

Let us note that the NPD measure depends implicitly not only on the filtration $\mathcal{X}_l \uparrow \mathcal{X}$ but also on the measure μ . In order to prove strong universality of the total NPD measure, we will apply the following lemma concerning convergence of f -divergences for filtrations:

Theorem 5 ([13, Chapter 3, Problem 4]). *For an interval A , let $f : A \rightarrow [0, \infty]$ be a non-negative, continuous, and convex measurable function, let $\nu \ll \rho$ be two finite measures on a measurable space, and let $\mathcal{G}_n \uparrow \mathcal{G}$ be a filtration. We have*

$$\lim_{n \rightarrow \infty} \int f \left(\frac{d\nu|_{\mathcal{G}_n}}{d\rho|_{\mathcal{G}_n}} \right) d\rho = \int f \left(\frac{d\nu|_{\mathcal{G}}}{d\rho|_{\mathcal{G}}} \right) d\rho, \quad (32)$$

where the sequence on the left hand side is increasing.

Remark: Theorem 5 follows by a synergy of the Fatou lemma and the Jensen inequality. One yields the lower bound, whereas another yields the upper bound.

Now we will derive the main result of this section.

Theorem 6. *Measure NPD is strongly universal with respect to the finite measure μ .*

Proof. It suffices to show that for $P \in \mathbb{E}(\mu)$, we have

$$\limsup_{n \rightarrow \infty} \frac{[-\log \text{NPD}_\mu(X_{1:n})]}{n} \leq h_\mu \text{ a.s.} \quad (33)$$

The strong universality follows hence by the converse bound (10).

So as to demonstrate (33), we first observe that by the strong universality of the total PPM measures, we have

$$\lim_{n \rightarrow \infty} \frac{[-\log \text{PPM}^{D_l}(X_{1:n}^l)]}{n} = \inf_{n \geq 1} \frac{\mathbf{E} [-\log P_n(X_{1:n}^l)]}{n} \text{ a.s.} \quad (34)$$

On the other hand, by the Birkhoff ergodic theorem, we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n [-\log \mu(X_i^l)] = C_l := \mathbf{E} [-\log \mu(X_i^l)] \text{ a.s.} \quad (35)$$

Moreover, each cross entropy C_l is finite since each C_l is a sum over finitely many finite elements—by $P_1 \ll \mu$. Denote quantities

$$h_\mu^l(n) := \mathbf{E} \left[-\log \frac{P_n(X_{1:n}^l)}{\prod_{i=1}^n \mu(X_i^l)} \right] = \mathbf{E} [-\log P_n(X_{1:n}^l)] - n C_l. \quad (36)$$

Since cross entropies C_l are finite, equations (34) and (35) imply

$$\lim_{n \rightarrow \infty} \frac{[-\log \text{NPD}_\mu^l(X_{1:n})]}{n} = h_\mu^l := \inf_{n \geq 1} \frac{h_\mu^l(n)}{n} \text{ a.s.} \quad (37)$$

Since $\text{NPD}_\mu(x_{1:n}) \geq w_l \text{NPD}_\mu^l(x_{1:n})$ then for any $l \geq 0$, we obtain

$$\limsup_{n \rightarrow \infty} \frac{[-\log \text{NPD}_\mu(X_{1:n})]}{n} \leq \limsup_{n \rightarrow \infty} \frac{[-\log w_l \text{NPD}_\mu^l(X_{1:n})]}{n} = h_\mu^l \text{ a.s.} \quad (38)$$

It remains to show that $\inf_{l \geq 0} h_\mu^l = h_\mu$. For this goal we observe that

$$\frac{P_n(x_{1:n}^l)}{\prod_{i=1}^n \mu(x_i^l)} = \frac{dP_n|_{\mathcal{X}_l^n}}{d\mu^n|_{\mathcal{X}_l^n}}(x_{1:n}). \quad (39)$$

Hence we have

$$h_\mu^l = \inf_{n \geq 1} \frac{h_\mu^l(n)}{n} = \inf_{n \geq 1} \frac{1}{n} \int \eta \left(\frac{dP_n|_{\mathcal{X}_l^n}}{d\mu^n|_{\mathcal{X}_l^n}} \right) d\mu^n, \quad (40)$$

where $\eta(x) := -x \log x$. We switch the order of infimums,

$$\inf_{l \geq 0} h_\mu^l = \inf_{n \geq 1} \frac{1}{n} \inf_{l \geq 0} \int \eta \left(\frac{dP_n|_{\mathcal{X}_l^n}}{d\mu^n|_{\mathcal{X}_l^n}} \right) d\mu^n \quad (41)$$

and we apply Theorem 5 to function $f(x) = \log 2 - \eta(x)$. Hence

$$\inf_{l \geq 0} h_\mu^l = \inf_{n \geq 1} \frac{1}{n} \int \eta \left(\frac{dP_n}{d\mu^n} \right) d\mu^n = \inf_{n \geq 1} \frac{h_\mu(n)}{n} = h_\mu. \quad (42)$$

The proof is complete. \square

The proof of universality in expectation applies a bit different idea than in the finite alphabet case. The role of the dominated convergence is played by a synergy of the reversed Fatou lemma and the source coding inequality (12).

Theorem 7. *Measure NPD is universal in expectation with respect to the finite measure μ .*

Proof. Since $\text{NPD}_\mu(x_{1:n}) \geq \frac{1}{2}$, we have $-\log \text{NPD}_\mu(X_{1:n}) \leq \log 2$. Hence by the reversed Fatou lemma and Theorem 6, we obtain

$$\limsup_{n \rightarrow \infty} \frac{\mathbf{E} [-\log \text{NPD}_\mu(X_{1:n})]}{n} \leq \mathbf{E} \limsup_{n \rightarrow \infty} \frac{[-\log \text{NPD}_\mu(X_{1:n})]}{n} = h_\mu. \quad (43)$$

Contrasting this with inequality (12) yields the claim. \square

3.2 Conditional density estimation

In this section we will show that a universal density, if it exists, induces a strongly consistent Cesàro mean estimator of the conditional density. For this goal, we consider a general reference measure μ as in Section 2.1.

A stochastic process $(R_\mu^{(n)})_{n \geq 1}$ is called a conditional density estimator if each random variable $R_\mu^{(n)}(x)$ is a (marginal) probability density with respect to measure μ and each function $R_\mu^{(n)}$ is a measurable function of random variables $X_{-n:-1}$. We also denote the respective random measure $R^{(n)}(A) := \int_A R_\mu^{(n)}(x) d\mu(x)$. We will seek for a conditional density estimator that converges in some strong sense to the conditional density

$$P_\mu^{(\infty)}(x) := P_\mu(x|X_{-\infty:-1}) = P_\mu^{(n)}(x) := \lim_{n \rightarrow \infty} P_\mu(x|X_{-n:-1}) \quad (44)$$

for any $P \in \mathbb{E}(\mu)$. We denote the respective random measure $P^{(\infty)}(A) := \int_A P_\mu^{(\infty)}(x) d\mu(x)$. Obviously $P^{(\infty)} = P_1$ if P is a memoryless source.

Inspired by the construction of [18], we will consider the following object:

Definition 5. *Consider a probability measure R where $R_n \ll \mu^n$. The Cesàro mean measure \bar{R} is defined via conditional densities*

$$\bar{R}_\mu(x_n|x_{1:n-1}) := \frac{1}{n} \sum_{i=0}^{n-1} R_\mu(x_n|x_{n-i:n-1}). \quad (45)$$

Let us observe that we may introduce a conditional density estimator

$$\bar{R}_\mu^{(n)}(x) := \bar{R}_\mu(x|X_{-n:-1}). \quad (46)$$

We will call it the Cesàro mean density estimator. We denote the respective random measure $\bar{R}^{(n)}(A) := \int_A \bar{R}_\mu^{(n)}(x) d\mu(x)$. In work [18], a similar conditional density estimator was considered, albeit with a reflected time arrow. The Cesàro mean measure \bar{R} reminds also of linear interpolation models used for statistical language modeling in the 1990's [19].

For the Cesàro mean density estimator, we have the following result which generalizes [18, Theorem 1] for memoryless sources.

Theorem 8. *Consider a measure $P \in \mathbb{E}(\mu)$, a probability measure R where $R_n \ll \mu^n$, and the Cesàro mean density estimator $(\bar{R}_\mu^{(n)})_{n \geq 1}$. We have inequality*

$$\mathbf{E} [-\log R(X_{1:n})] / n - h_\mu \geq \mathbf{E} D(P^{(\infty)} | \bar{R}^{(n)}). \quad (47)$$

Proof. We essentially apply the proof idea of [18, Theorem 1], which is a restriction of the present claim to μ being the counting measure and P being a memoryless source. In the reasoning rewritten in a more transparent notation we apply the stationarity and the Jensen inequality,

$$\begin{aligned} \mathbf{E} [-\log R(X_{1:n})] / n - h_\mu &= \mathbf{E} \left[-\frac{1}{n} \sum_{i=1}^n \log R_\mu(X_i | X_{1:i-1}) \right] - h_\mu \\ &= \mathbf{E} \left[-\frac{1}{n} \sum_{i=0}^{n-1} \log R_\mu(X_0 | X_{-i:-1}) \right] - h_\mu \\ &\geq \mathbf{E} \left[-\log \frac{1}{n} \sum_{i=0}^{n-1} R_\mu(X_0 | X_{-i:-1}) \right] - h_\mu \\ &= \mathbf{E} D(P^{(\infty)} | \bar{R}^{(n)}), \end{aligned} \quad (48)$$

where the last transition is due to equality $h_\mu = \mathbf{E} [-\log P_\mu^{(\infty)}(X_0)]$ shown by Barron [5, Proof of Theorem 1]. \square

We recall the total variation distance of probability measures P_1 and R_1 on a measurable space $(\mathbb{X}, \mathcal{X})$, defined as

$$\delta(P_1, R_1) := \sup_{A \in \mathcal{X}} |P_1(A) - R_1(A)|. \quad (49)$$

If $P_1, R_1 \ll \mu$ then $\delta(P_1, R_1) = \frac{1}{2} \int |P_\mu(x) - R_\mu(x)| d\mu(x)$. We also recall the Pinsker inequality [11], which reads

$$\delta(P_1, R_1) \leq \sqrt{\frac{D(P_1 | R_1)}{2 \log e}}. \quad (50)$$

In consequence, if a universal density exists and the entropy rate is finite then the Cesàro mean density estimator is strongly consistent in the total variation.

Theorem 9. *Suppose that measure R is universal in expectation with respect to a reference measure μ . Then for every measure $P \in \mathbb{E}(\mu)$, we have*

$$\lim_{n \rightarrow \infty} \delta(P^{(\infty)}, \bar{R}^{(n)}) = \lim_{n \rightarrow \infty} \delta(P^{(n)}, \bar{R}^{(n)}) = 0 \text{ a.s.} \quad (51)$$

Proof. If R is universal in expectation then for every measure $P \in \mathbb{E}(\mu)$, we have

$$0 = \lim_{n \rightarrow \infty} \mathbf{E} [-\log R(X_{1:n})] / n - h_\mu \geq \lim_{n \rightarrow \infty} \mathbf{E} D(P^{(\infty)} || \bar{R}^{(n)}) \geq 0. \quad (52)$$

Now by the Pinsker inequality (50) and by the dominated convergence, we obtain

$$0 = \lim_{n \rightarrow \infty} \mathbf{E} [\delta(P^{(\infty)}, \bar{R}^{(n)})]^2 = \mathbf{E} \left[\lim_{n \rightarrow \infty} \delta(P^{(\infty)}, \bar{R}^{(n)}) \right]^2, \quad (53)$$

which implies the almost sure convergence for $\delta(P^{(\infty)}, \bar{R}^{(n)})$. Since

$$h_\mu = \lim_{n \rightarrow \infty} \mathbf{E} [-\log P_\mu^{(n)}(X_0)] \quad (54)$$

for a stationary P then we also have

$$\lim_{n \rightarrow \infty} \mathbf{E} D(P^{(n)} || \bar{R}^{(n)}) = \lim_{n \rightarrow \infty} \mathbf{E} D(P^{(\infty)} || \bar{R}^{(n)}) = 0. \quad (55)$$

Thus, analogously we derive the almost sure convergence for $\delta(P^{(n)}, \bar{R}^{(n)})$. \square

3.3 Universal prediction

Theorem 9 generalizes the celebrated Ornstein theorem, originally stated for binary stationary ergodic processes [25]. Ornstein's theorem plays an important role in the theory of universal prediction of binary processes with the 0–1 loss [23]. Analogously, we can apply Theorem 9 to develop a theory of universal prediction for processes over an arbitrary countable alphabet, strengthening the recent result of [15] by the way.

The exact development, recalling the basic facts from [15], is as follows. For a countable alphabet \mathbb{X} , we will consider densities with respect to the counting measure γ , denoted without subscript γ according to our earlier convention. A predictor is an arbitrary function $f : \mathbb{X}^* \rightarrow \mathbb{X}$.

Definition 6. *The predictor induced by a probability measure P is defined as a maximizer*

$$f_P(x_{1:n-1}) \in \arg \max_{x_n \in \mathbb{X}} P(x_n | x_{1:n-1}). \quad (56)$$

In the problem of universal prediction we seek for a predictor that minimizes the relative frequency of prediction mistakes. That is, we apply the 0 – 1 loss rather than the logarithmic loss encountered in the problem of universal coding. By the Azuma-Hoeffding inequality [3], for any probability measure P , the rate of mistakes is equal to the rate of their conditional probabilities, namely,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n [\mathbf{1}\{X_i \neq f(X_{1:i-1})\} - P(X_i \neq f(X_{1:i-1})|X_{1:i-1})] = 0 \text{ a.s.}, \quad (57)$$

see [15, Theorem 3.5] for the derivation. Therefore, as shown in [1], [15, Theorem 3.5] using the Breiman ergodic theorem [7], [1, Theorem 12(c)], for any measure $P \in \mathbb{E}$ and any predictor f , we have

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \mathbf{1}\{X_i \neq f(X_{1:i-1})\} \geq u \text{ a.s.} \quad (58)$$

with the equality for $f = f_P$, where we define the unpredictability rate

$$u := \mathbf{E} \left[1 - \max_{x \in \mathbb{X}} P(x|X_{-\infty:-1}) \right]. \quad (59)$$

Thus by an analogy to universal measures, we propose universal predictors.

Definition 7. *A predictor f is called almost surely universal with respect to a reference measure μ if (58) holds with the equality for any measure $P \in \mathbb{E}(\mu)$.*

In work [15, Theorems 3.12 and 3.18], it was shown that for a finite alphabet \mathbb{X} and an almost surely universal measure R , the induced predictor f_R is almost surely universal if we have a uniform bound for conditional probabilities of form

$$-\log R(x_{n+1}|x_1^n) \leq \epsilon_n \sqrt{n/\log n}, \quad \lim_{n \rightarrow \infty} \epsilon_n = 0. \quad (60)$$

Subsequently, we will show that condition (60) can be dropped if R is universal in expectation and we consider predictor $f_{\bar{R}}$, induced by the Cesàro mean measure \bar{R} , rather than predictor f_R , induced by measure R .

Theorem 10. *Consider a countable alphabet \mathbb{X} . Suppose that measure R is universal in expectation with respect to a reference measure μ . The Cesàro mean predictor $f_{\bar{R}}$ is almost surely universal with respect to measure μ .*

Proof. Consider a measure $P \in \mathbb{E}(\mu)$. By Theorem 9, we have a generalization of the Ornstein theorem [25], namely,

$$\lim_{n \rightarrow \infty} 2\delta(P^{(n)}, \bar{R}^{(n)}) = \lim_{n \rightarrow \infty} \sum_{x \in \mathbb{X}} |P(x|X_{-n:-1}) - \bar{R}(x|X_{-n:-1})| = 0 \text{ a.s.} \quad (61)$$

since $P^{(n)}, \bar{R}^{(n)} \ll \gamma$. As it follows from a simple application of the Breiman ergodic theorem [7], [1, Theorem 12(c)], statement (61) implies a generalization of the Bailey theorem [4], namely,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n \sum_{x \in \mathbb{X}} |P(x|X_{1:i-1}) - \bar{R}(x|X_{1:i-1})| = 0 \text{ a.s.} \quad (62)$$

Since we have a so called prediction inequality

$$\begin{aligned} & P(X_i = f_P(X_{1:i-1})|X_{1:i-1}) - P(X_i = f_{\bar{R}}(X_{1:i-1})|X_{1:i-1}) \\ & \leq \sum_{x \in \mathbb{X}} |P(x|X_{1:i-1}) - \bar{R}(x|X_{1:i-1})| \end{aligned} \quad (63)$$

noticed in work [15, Proposition 3.11] then by corollary (57) of the Azuma-Hoeffding inequality, we obtain

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=1}^n [\mathbf{1}\{X_i \neq f_{\bar{R}}(X_{1:i-1})\} - \mathbf{1}\{X_i \neq f_P(X_{1:i-1})\}] = 0 \text{ a.s.} \quad (64)$$

That is, the Cesàro mean predictor $f_{\bar{R}}$ is strongly universal. \square

3.4 Countably infinite alphabet

Subsequently, let us consider a countably infinite alphabet \mathbb{X} . For probability measures P and R , we denote the Shannon entropy and the Kullback-Leibler divergence taken with respect to their marginal densities:

$$H(P_1) := - \sum_{x \in \mathbb{X}} P(x) \log P(x), \quad (65)$$

$$D(P_1||R_1) := \sum_{x \in \mathbb{X}} P(x) \log \frac{P(x)}{R(x)}. \quad (66)$$

We recall that there are no consistent conditional density estimators for a countably infinite alphabet.

Theorem 11 ([18, Theorem 2]). *Let the alphabet \mathbb{X} be countably infinite. Let $(R_\mu^{(n)})_{n \geq 1}$ be an arbitrary conditional density estimator. Then there is a memoryless source P such that $H(P_1) < \infty$ and $D(P_1||R^{(n)}) = \infty$ a.s. for all n .*

In view of Theorems 8 and 11, for a countably infinite alphabet \mathbb{X} , for each measure R there exists a memoryless source P with $H(P_1) < \infty$ such that

$$\lim_{n \rightarrow \infty} \mathbf{E} [-\log R(X_{1:n})] / n = \infty. \quad (67)$$

Since each memoryless source P is stationary ergodic with entropy rate $H(P_1)$ we obtain the known result.

Theorem 12 ([20], [18, Theorem 3]). *There is no measure universal in expectation with respect to a countably infinite alphabet.*

For a countably infinite alphabet \mathbb{X} , consider now a reference measure μ to be contrasted with the counting measure γ . We suppose that $\mu \ll \gamma$ and $\mu \gg \gamma$. Entropies of P with respect to μ are written as $h_\mu(n)$ or h_μ , whereas entropies of P with respect to γ are written as $h(n)$ or h . We have

$$P(x_{1:n}) = P_\mu(x_{1:n}) \frac{d\mu^n}{d\gamma^n}(x_{1:n}) = P_\mu(x_{1:n}) \prod_{i=1}^n \mu(x_i). \quad (68)$$

Let us write the marginal cross entropy

$$H(P_1||R_1) := H(P_1) + D(P_1||R_1) = - \sum_{x \in \mathbb{X}} P(x) \log R(x). \quad (69)$$

If $H(P_1||\mu) < \infty$ then from equation (68), applying stationarity, we obtain

$$h(n) = h_\mu(n) - n \sum_{x \in \mathbb{X}} P(x) \log \mu(x) = h_\mu(n) + nH(P_1||\mu). \quad (70)$$

Hence $h = h_\mu + H(P_1||\mu) \geq 0$. If μ is a probability measure then $h_\mu \leq 0$. Consequently, we have $P \in \mathbb{E}(\mu)$ if there holds $P \in \mathbb{E}$, $\mu(x) > 0$ for all $x \in \mathbb{X}$, and $H(P_1||\mu) < \infty$.

In particular, we may estimate the entropy rate in the following way, compare it with more complicated characterizations in [29]:

Theorem 13. *Consider a countably infinite alphabet \mathbb{X} and a probability measure μ such that $\mu(x) > 0$ for all $x \in \mathbb{X}$. Let $P \in \mathbb{E}$ with $H(P_1||\mu) < \infty$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left[-\log \text{NPD}_\mu(X_{1:n}) - \sum_{i=1}^n \log \mu(X_i) \right] = h \text{ a.s.} \quad (71)$$

Proof. Quantity $-n^{-1} \log \text{NPD}_\mu(X_{1:n})$ is a strongly consistent estimator of h_μ , whereas $-n^{-1} \sum_{i=1}^n \log \mu(X_i)$ is a strongly consistent estimator of $H(P_1||\mu)$ by the Birkhoff ergodic theorem. Since $H(P_1||\mu) < \infty$ then the sum of these two estimators converges to h . \square

Estimator (71) differs from the PPM measure since $\mu(X_i) \leq \mu(X_i^l)$. To observe some simplifications of estimator (71), let us suppose that the alphabet is the set of natural numbers, $\mathbb{X} = \mathbb{N}$, and we take partitions

$$\chi_l := \{\{1\}, \{2\}, \dots, \{l-1\}, \{l, l+1, \dots\}\}, \quad l \in \mathbb{N}, \quad (72)$$

so that $D_l = l$. Then for each quantization level l , we have the bijection $x^l \leftrightarrow x \wedge l$, where $x \wedge a := \min\{x, a\}$. Let us denote the optimal quantization level Q_n and the optimal Markov order R_n for sample $X_{1:n}$ as some elements

$$(Q_n, R_n) \in \arg \max_{(q,r) \in \mathbb{N}_0^2} \frac{\text{PPM}_r^q(X_{1:n}^q)}{\mu^n(X_{1:n}^q)}, \quad (73)$$

where $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. Since we have $\text{PPM}_r^{q+1}(x_{1:n}) < \text{PPM}_r^q(x_{1:n})$ in general and $\text{PPM}_r^q(x_{1:n}) = q^{-n}$ for $r \geq n - 1$, we may fix pair (Q_n, R_n) so that

$$Q_n \leq \max X_{1:n}, \quad R_n \leq n - 1. \quad (74)$$

Moreover, using the known idea for the PPM measure, we may bound

$$0 \leq -\log \text{NPD}_\mu(X_{1:n}) + \log \frac{\text{PPM}_{R_n}^{Q_n}(X_{1:n}^{Q_n})}{\mu^n(X_{1:n}^{Q_n})} \leq -\log w_{Q_n} - \log w_{R_n}. \quad (75)$$

Consequently, if the maximal observation X_i does not grow too fast, namely, when we have

$$\lim_{n \rightarrow \infty} \frac{\log \max X_{1:n}}{n} = 0 \text{ a.s.} \quad (76)$$

then we may rewrite statement (71) as proposition

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left[-\log \text{PPM}_{R_n}^{Q_n}(X_{1:n} \wedge Q_n) + C_n(Q_n) \right] = h \text{ a.s.}, \quad (77)$$

where we define

$$C_n(q) := -\sum_{i=1}^n \log \frac{\mu(X_i)}{\mu(X_i^q)} = -\sum_{i=1}^n \mathbf{1}\{X_i \geq q\} \log \mu(X_i | X_i \geq q) \geq 0. \quad (78)$$

Term $C_n(Q_n)$ is a non-negative correction of the finite-alphabet PPM entropy estimator taken for the optimal quantization level and the optimal Markov order. It is not obvious that $\lim_{n \rightarrow \infty} C_n(Q_n)/n = 0$ holds almost surely but in some cases it seems a reasonable hypothesis. For instance, let us suppose that

$$P(X_i \geq Q_i) \geq P(X_i \geq Q_n) \text{ for } n \geq i, \quad (79)$$

$$\lim_{n \rightarrow \infty} P(X_i \geq Q_i) = 0, \quad (80)$$

$$\mathbf{E} \log^2 \mu(X_i) < \infty. \quad (81)$$

Then by the Cauchy-Schwarz inequality and by the Toeplitz lemma, we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\mathbf{E} C_n(Q_n)}{n} &\leq \lim_{n \rightarrow \infty} \mathbf{E} \sum_{i=1}^n \frac{\mathbf{1}\{X_i \geq Q_n\}}{\sqrt{n}} \frac{[-\log \mu(X_i)]}{\sqrt{n}} \\ &\leq \lim_{n \rightarrow \infty} \mathbf{E} \sqrt{\sum_{i=1}^n \frac{\mathbf{1}\{X_i \geq Q_n\}}{n}} \sqrt{\sum_{i=1}^n \frac{\log^2 \mu(X_i)}{n}} \\ &\leq \lim_{n \rightarrow \infty} \sqrt{\mathbf{E} \sum_{i=1}^n \frac{\mathbf{1}\{X_i \geq Q_n\}}{n}} \sqrt{\mathbf{E} \sum_{i=1}^n \frac{\log^2 \mu(X_i)}{n}} \end{aligned}$$

$$\begin{aligned}
&\leq \lim_{n \rightarrow \infty} \sqrt{\frac{1}{n} \sum_{i=1}^n P(X_i \geq Q_i)} \sqrt{\mathbf{E} \log^2 \mu(X_i)} \\
&= 0 \cdot \sqrt{\mathbf{E} \log^2 \mu(X_i)} = 0.
\end{aligned} \tag{82}$$

Hence $\lim_{n \rightarrow \infty} C_n(Q_n)/n = 0$ holds in probability in this case.

3.5 Real line and Gaussian processes

Consider alphabet $\mathbb{X} = \mathbb{R}$ and the reference measure μ being the normal distribution $N(m, \sigma^2)$ to be contrasted with the Lebesgue measure λ . Entropies of P with respect to μ are written as $h_\mu(n)$ or h_μ , whereas entropies of P with respect to λ are written as $h_\lambda(n)$ or h_λ . We have

$$\frac{d\mu}{d\lambda}(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-m)^2}{2\sigma^2}\right). \tag{83}$$

Let us write the rescaled second moment of the marginal distribution

$$M_2 := \int_{-\infty}^{\infty} \frac{(x-m)^2}{2\sigma^2} P_\lambda(x) dx = \frac{\text{Var } X_i + (\mathbf{E} X_i - m)^2}{2\sigma^2}, \tag{84}$$

denoting variance $\text{Var } X_i := \mathbf{E} (X_i - \mathbf{E} X_i)^2$.

If $M_2 < \infty$, we obtain like in Section 3.4 that

$$\begin{aligned}
h_\lambda(n) &= h_\mu(n) - n \int_{-\infty}^{\infty} P_\lambda(x) \log \frac{d\mu}{d\lambda}(x) dx. \\
&= h_\mu(n) + n \left[M_2 \log e + \log \sigma\sqrt{2\pi} \right].
\end{aligned} \tag{85}$$

Hence $h_\lambda = h_\mu + M_2 \log e + \log \sigma\sqrt{2\pi}$. Since μ is a probability measure then $h_\mu \leq 0$. Consequently, we have $P \in \mathbb{E}(\mu)$ if there holds $P \in \mathbb{E}$, $P_n \ll \lambda^n$, $|\mathbf{E} X_i| < \infty$, $\text{Var } X_i < \infty$, and $|h_\lambda| < \infty$.

In particular, we may estimate the entropy rate in the following way:

Theorem 14. *Consider $\mathbb{X} = \mathbb{R}$. Let μ be the normal distribution $N(m, \sigma^2)$, whereas λ be the Lebesgue measure. Suppose that $P \in \mathbb{E}$ with $P_n \ll \lambda^n$, $|\mathbf{E} X_i| < \infty$, $\text{Var } X_i < \infty$, and $|h_\lambda| < \infty$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \left[-\log \text{NPD}_\mu(X_{1:n}) + \left[\sum_{i=1}^n \frac{(X_i - m)^2}{2\sigma^2} \right] \log e \right] + \log \sigma\sqrt{2\pi} = h_\lambda \text{ a.s.} \tag{86}$$

Proof. Quantity $-n^{-1} \log \text{NPD}_\mu(X_{1:n})$ is a strongly consistent estimator of the entropy rate h_μ , whereas $-n^{-1} \sum_{i=1}^n (X_i - m)^2 / 2\sigma^2$ is a strongly consistent estimator of the rescaled second moment M_2 . Since $M_2 < \infty$ then the linear combination of these estimators tends to h_λ . \square

Thus, the corrected NPD estimator (86) is a strongly consistent estimator of the entropy rate for all non-deterministic stationary ergodic Gaussian processes since $|\mathbf{E} X_i| < \infty$, $\text{Var} X_i < \infty$, and $|h_\lambda| < \infty$ holds in this case. This idea, although it was not carried out to its end, motivated the work of Feutrill and Roughan [17].

We may suppose that the corrected NPD estimator (86) improves if we take m and σ close to the expectation and variance of X_i with respect to P . Obviously, we can prove easily:

Theorem 15. *Consider $\mathbb{X} = \mathbb{R}$. Let μ be the normal distribution $N(m, \sigma^2)$, whereas λ be the Lebesgue measure. Suppose that $P \in \mathbb{E}$ with $P_n \ll \lambda^n$, $\mathbf{E} X_i = m$, $\text{Var} X_i = \sigma^2$, and $|h_\lambda| < \infty$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} [-\log \text{NPD}_\mu(X_{1:n})] + \log \sigma \sqrt{2\pi e} = h_\lambda \text{ a.s.} \quad (87)$$

The problem with the above estimator is that we need to know the exact expectation and variance of X_i . Can we estimate them from sample $X_{1:n}$ and plug the result into the NPD estimator? Consider random measures $\mu_n \sim N(m_n, \sigma_n^2)$, where

$$m_n := \frac{1}{n} \sum_{i=1}^n X_i, \quad \sigma_n := \sqrt{\frac{1}{n} \sum_{i=1}^n (X_i - m_n)^2}. \quad (88)$$

We may ask whether it holds still

$$\lim_{n \rightarrow \infty} \frac{1}{n} [-\log \text{NPD}_{\mu_n}(X_{1:n})] + \log \sigma_n \sqrt{2\pi e} = h_\lambda \text{ a.s.} \quad (89)$$

However, this question is not stated precisely enough. Let us note that measure NPD depends implicitly on the quantization path $\mathcal{X}_l \uparrow \mathcal{X}$ and measure μ . When we variate the reference measure $\mu \rightarrow \mu_n$, it is not clear whether we also have to variate the σ -fields $\mathcal{X}_l \rightarrow \mathcal{X}_{l,n}$. In any case, the proof technique of Theorem 6 works no longer and we have no simple guarantee of consistency.

Funding

This work was supported by the National Science Centre Poland grant 2018/31/B/HS1/04018.

References

- [1] P. H. Algoet. The strong law of large numbers for sequential decisions under uncertainty. *IEEE Trans. Inform. Theory*, 40(3):609–633, 1994.

- [2] P. H. Algoet and T. M. Cover. A sandwich proof of the Shannon-McMillan-Breiman theorem. *Ann. Probab.*, 16:899–909, 1988.
- [3] K. Azuma. Weighted sums of certain dependent random variables. *Tohoku Math. J. Sec. Ser.*, 19(3):357–367, 1967.
- [4] D. H. Bailey. *Sequential schemes for classifying and predicting ergodic processes*. PhD thesis, Stanford University, 1976.
- [5] A. R. Barron. The strong ergodic theorem for densities: Generalized Shannon-McMillan-Breiman theorem. *Ann. Probab.*, 13:1292–1303, 1985.
- [6] A. R. Barron. *Logically Smooth Density Estimation*. PhD thesis, Stanford University, 1985.
- [7] L. Breiman. The individual ergodic theorem of information theory. *Ann. Math. Statist.*, 28:809–811, 1957.
- [8] G. J. Chaitin. A theory of program size formally identical to information theory. *J. ACM*, 22:329–340, 1975.
- [9] K. L. Chung. A note on the ergodic theorem of information theory. *Ann. Math. Statist.*, 32:612–614, 1961.
- [10] J. G. Cleary and I. H. Witten. Data compression using adaptive coding and partial string matching. *IEEE Trans. Comm.*, 32:396–402, 1984.
- [11] I. Csiszár and J. Körner. *Information Theory: Coding Theorems for Discrete Memoryless Systems*. Cambridge University Press, 2011.
- [12] Ł. Dębowski. Is natural language a perigraphic process? The theorem about facts and words revisited. *Entropy*, 20(2):85, 2018.
- [13] Ł. Dębowski. *Information Theory Meets Power Laws: Stochastic Processes and Language Models*. Wiley & Sons, 2021.
- [14] Ł. Dębowski. A refutation of finite-state language models through Zipf’s law for factual knowledge. *Entropy*, 23:1148, 2021.
- [15] Ł. Dębowski and T. Steifer. Universal coding and prediction on ergodic random points. *Bull. Symb. Logic*, 28(2):387–412, 2022.
- [16] M. Fekete. Über die Verteilung der Wurzeln bei gewissen algebraischen Gleichungen mit ganzzahligen Koeffizienten. *Math. Z.*, 17:228–249, 1923.
- [17] A. Feutrill and M. Roughan. NPD entropy: A non-parametric differential entropy rate estimator. <https://arxiv.org/abs/2105.11580>, 2021.
- [18] L. Györfi, I. Páli, and E. C. van der Meulen. There is no universal source code for infinite alphabet. *IEEE Trans. Inform. Theory*, 40:267–271, 1994.

- [19] F. Jelinek. *Statistical Methods for Speech Recognition*. The MIT Press, 1997.
- [20] J. Kieffer. A unified approach to weak universal source coding. *IEEE Trans. Inform. Theory*, 24:674–682, 1978.
- [21] J. C. Kieffer and E. Yang. Grammar-based codes: A new class of universal lossless source codes. *IEEE Trans. Inform. Theory*, 46:737–754, 2000.
- [22] N. Merhav, M. Gutman, and J. Ziv. On the estimation of the order of a Markov chain and universal data compression. *IEEE Trans. Inform. Theory*, 35(5):1014–1019, 1989.
- [23] G. Morvai and B. Weiss. On universal algorithms for classifying and predicting stationary processes. *Probab. Surveys*, 18:77–131, 2021.
- [24] D. Neuhoff and P. C. Shields. Simplistic universal coding. *IEEE Trans. Inform. Theory*, IT-44:778–781, 1998.
- [25] D. S. Ornstein. Guessing the next output of a stationary process. *Israel J. Math.*, 30(3):292–296, 1978.
- [26] B. Ryabko. Compression-based methods for nonparametric density estimation, on-line prediction, regression and classification for time series. In *2008 IEEE Information Theory Workshop, Porto*, pages 271–275. Institute of Electrical and Electronics Engineers, 2008.
- [27] B. Y. Ryabko. Prediction of random sequences and universal coding. *Probl. Inform. Transm.*, 24(2):87–96, 1988.
- [28] C. Shannon. A mathematical theory of communication. *Bell Syst. Tech. J.*, 30:379–423,623–656, 1948.
- [29] J. F. Silva and P. Piantanida. Universal weak variable-length source coding on countably infinite alphabets. *IEEE Trans. Inform. Theory*, 66(1):649–668, 2020.
- [30] J. Ziv and A. Lempel. A universal algorithm for sequential data compression. *IEEE Trans. Inform. Theory*, 23:337–343, 1977.
- [31] J. Ziv and N. Merhav. Estimating the number of states of a finite-state source. *IEEE Trans. Inform. Theory*, 38(1):61–65, 1992.