

Convex polygons and separation of convex sets*

Eduardo Rivera-Campo¹ and Jorge Urrutia²

¹Departamento de Matemáticas, Universidad Autónoma
Metropolitana - Iztapalapa, erc@xanum.uam.mx

²Instituto de Matemáticas, Universidad Nacional Autónoma de
México, urrutia@matem.uam.mx

Abstract

We prove that for any collection F of $n \geq 2$ pairwise disjoint compact convex sets in the plane there is a pair of sets A and B in F such that any line that separates A from B separates either A or B from a subcollection of F with at least $n/18$ sets.

Keywords.- Convex polygon. Plane Compact Convex Set. Separating line.

1 Introduction

H. Tverberg [5] proved that for each positive integer k , there is a minimum integer $f(k)$ such that for every collection F of $f(k)$ or more plane compact convex sets with pairwise disjoint interiors, there is a line that separates one set in F from a subcollection of F with at least k sets. R. Hope and M. Katchalski [3] showed that $3k + 1 \leq f(k) \leq 12k - 1$.

Later E. Rivera-Campo and J. Töröcsik [4] proved that in any collection F of $n \geq 5$ pairwise disjoint compact convex sets in the plane, there is a pair of sets A and B such that any line that separates A from B separates either A or B from at least $\frac{n+28}{30}$ sets in F . In this paper we give a higher lower bound of $\frac{n}{18}$ sets in F for any collection F of $n \geq 2$ sets.

*Partially supported by Conacyt, México.

2 Preliminary results

H. Edelsbrunner *et al* [1] proved the following theorem, see L. Fejes-Toth [2] for a related result.

Theorem 1. *Any collection of $n \geq 3$ compact, convex and pairwise disjoint sets in the plane may be covered with non-overlapping convex polygons with a total of not more than $6n - 9$ sides. Furthermore no more than $3n - 6$ distinct slopes are required.*

We adapt part of the proof given in [1] to obtain the following result.

Lemma 1. *Let $\mathcal{P} = \{P_1, P_2, \dots, P_n\}$ be a collection of $n \geq 3$ pairwise disjoint convex polygons in the plane. There exists a collection $\mathcal{R} = \{R_1, R_2, \dots, R_n\}$ of pairwise non-overlapping convex polygons such that:*

1. *For $i = 1, 2, \dots, n$, P_i is contained in R_i .*
2. *For $i = 1, 2, \dots, n$, each side of R_i supports a side of P_i .*
3. *The total number of sides among polygons R_1, R_2, \dots, R_n is at most $9n - 9$.*

Proof. A side s of a polygon $P_i \in \mathcal{P}$ is *reducible with respect to \mathcal{P}* if the triangle t_s (not containing P_i), bounded by s and the lines supporting the sides of P_i incident to s , does not intersect the interior of any another polygon P_j . Equivalently, a side s of a polygon $P_i \in \mathcal{P}$ is reducible with respect to \mathcal{P} if it vanishes before reaching the interior of a polygon P_j when it is translated in the direction orthogonal to s and away from P_i (see Fig. 1).




Figure 1: Polygon P with a reducible side s .

We modify \mathcal{P} by growing, one at a time, polygons in \mathcal{P} as follows: if some polygon $P_i \in \mathcal{P}$ has a reducible side s with respect to \mathcal{P} substitute P_i in \mathcal{P} with polygon $P_i \cup t_s$. Repeat this until no polygon in \mathcal{P} has a reducible side. Denote by $\mathcal{R} = \{R_1, R_2, \dots, R_n\}$ the family of polygons thus obtained.

Its is clear that \mathcal{R} satisfies conditions 1) and 2), we claim it also satisfies condition 3. To prove this consider a third family $\mathcal{Q} = \{Q_1, Q_2, \dots, Q_n\}$ of convex polygons obtained from \mathcal{R} by further growing polygons R_1, R_2, \dots, R_n in the following way.

A side s of a polygon R_i in \mathcal{R} is *free* with respect to \mathcal{R} if the open polygonal region, not containing R_i , which is determined by s and the lines supporting the two sides of R_i adjacent to s contains no points of polygons in \mathcal{R} .

Let C be a circle containing all polygons in \mathcal{R} . Starting with the non free sides of polygons $R_i \in \mathcal{R}$, translate one at a time a side s of a polygon $R_i \in \mathcal{R}$ (also in the direction orthogonal to s and away from R_i) as far as possible without intersecting the interior of another (possibly already grown) polygon R_j or the exterior of C (see Fig. 2).




Figure 2: Side s of R_i reaches a polygon R_j or reaches circle C .

Stop when no side can be translated as indicated and let $\mathcal{Q} = \{Q_1, Q_2, \dots, Q_n\}$ where, for $i = 1, 2, \dots, n$, Q_i is the polygon obtained from R_i in this manner. We claim that if C is chosen large enough, then each polygon in \mathcal{Q}

contains at most 3 free sides and all non free sides of a polygon Q_i in \mathcal{Q} are in contact with a polygon Q_j with $i \neq j$.

By the choice of \mathcal{R} no side of a polygon R_j vanishes, therefore for $i = 1, 2, \dots, n$ the number of sides of R_i is equal to the number of sides of Q_i .

We define a plane graph G with one vertex v_i placed in the interior of each polygon $Q_i \in \mathcal{Q}$ and a vertex v_{n+1} placed outside circle C . The edge set of G consists of 6 types of edges as follows:

a) If a side of a polygon Q_i and a side of a polygon Q_j have a segment l in common choose an arbitrary point p in l and draw an edge $v_i v_j$ through p as in Fig. 3 (left).

b) If a vertex p of a polygon Q_i lies in the interior of a side of a polygon Q_j and p is not a vertex of a polygon in \mathcal{Q} other than Q_i , then draw an edge $v_i v_j$ through p as in Fig. 3 (center).

c) If two or more polygons $Q_{i_1}, Q_{i_2}, \dots, Q_{i_r}$ share a vertex p which lies in the interior of a side of a polygon Q_j and p is not a point of any polygon Q_k other than $Q_{i_1}, Q_{i_2}, \dots, Q_{i_r}$, then there is a neighbourhood N of p containing no points of polygons Q_k with $k \neq j, i_1, i_2, \dots, i_r$. In this case draw a cycle with edges $v_j v_{i_1}, v_{i_1} v_{i_2}, v_{i_2} v_{i_3}, \dots, v_{i_{r-1}} v_{i_r}, v_{i_r} v_j$ as in Fig. 3 (right).




Figure 3: Edge type a) (left), edge type b) (center) and edges type c) (right).

d) If three or more polygons $Q_{i_1}, Q_{i_2}, \dots, Q_{i_r}$ share a vertex p and p is not a point of any polygon Q_k other than $Q_{i_1}, Q_{i_2}, \dots, Q_{i_r}$, then there is a neighbourhood N of p containing no points of any polygon Q_k with $k \neq i_1, i_2, \dots, i_r$. In this case draw a cycle with edges $v_{i_1} v_{i_2}, v_{i_2} v_{i_3}, \dots, v_{i_{r-1}} v_{i_r}, v_{i_r} v_{i_1}$ as in Fig. 4 (left).

e) If two polygons Q_i and Q_j share a vertex p and p is not a point of any polygon in \mathcal{R} other than Q_i and Q_j , then draw an edge $v_i v_j$ through p as in Fig. 4 (right).

f) For each free side s of a polygon Q_i add an edge $v_i v_{n+1}$ drawn through an interior of point p in s .




Figure 4: Edges type d) (left) and edge type e) (right).

Notice that edges of type f) are the only possible multiple edges of G . Therefore G is a plane multigraph with $n + 1$ vertices and $m \leq (3(n + 1) - 6) + (x_2 + 2x_3) = (3n - 3) + (x_2 + 2x_3)$ edges, where for $i = 2, 3$, x_i denotes the number of polygons in \mathcal{Q} with i free sides.




Figure 5: Graph G with $n = 18$, $x_1 = 6$, $x_2 = 2$, $m_e = 1$ and $m_f = 10$.

We say that an edge $e = v_i v_j$ of G intersects a side s of a polygon in \mathcal{Q}

if e is drawn through either an interior point of s or through a vertex of s . Each non free side among polygons in \mathcal{Q} is intersected by at least one edge of G of type a), b), c), d) or e) and each free side is intersected by an edge of type f). We claim that if an edge $v_i v_j$ of G is of type e) then two of the 4 sides intersected by $v_i v_j$ are intersected by at least another edge of G .

Proof of Claim. Let $v_i v_j$ be an edge of G of type e). Then either two sides s, t of Q_i (of Q_j) or one side s of Q_i and one side t of Q_j are such that the lines supporting s and t separate Q_i and Q_j (see Fig. 6).



Figure 6: Lines supporting s and t separate Q_i and Q_j .

Without loss of generality we assume the line supporting a side s of Q_i separates Q_i and Q_j . Let b be the other side of Q_i incident in the common vertex p of Q_i and Q_j . Then side b must contain a point of a polygon Q_k other than Q_i and Q_j , otherwise b could be translated away from Q_i in the direction orthogonal to b which is not possible by the properties of \mathcal{Q} (see Fig. 7). This means that side b is intersected by an edge of G other than edge $v_i v_j$.




Figure 7: Side b could be translated away from Q_i .

Analogously a side $c \neq b$, incident with t , is intersected by an edge of G other than $v_i v_j$. \square

Let T denote the total number of sides among polygons in \mathcal{Q} . Also let m , m_e and m_f denote the number of edges of G , the number of edges of G of

type e) and the number of edges of G of type f), respectively. Since polygons in \mathcal{Q} contain at most 3 free edges, $m_f = x_1 + 2x_2 + 3x_3$, where for $i = 1, 2, 3$ x_i is the number of polygons in \mathcal{Q} containing i free edges.

In order to bound the number of sides among polygons in \mathcal{Q} we add the number of sides intersected by edges of G and subtract the number of sides intersected by at least two edges of G (2 sides for each edge of type e)). Considering that edges of types a), b), c), and d), intersect at most 3 sides; edges of type e) intersect 4 sides and edges of type f) intersect one side, we have:

$$\begin{aligned}
 T &\leq (3(m - (m_e + m_f)) + 4m_e + m_f) - 2m_e \\
 &= 3m - m_e - 2m_f \\
 &\leq 3m - 2m_f \\
 &\leq 3((3n - 3) + (x_2 + 2x_3)) - 2(x_1 + 2x_2 + 3x_3) \\
 &= 9n - 9 - 2x_1 - x_2 \\
 &\leq 9n - 9.
 \end{aligned}$$

□

Consider the following family of plane geometric graphs G_t , $t = 1, 2, \dots$ given by Edelsbrunner *et al* [1] where it is used to show that the bounds in the number of sides and slopes on Theorem 1 are tight.

G_1 is the graph with 7 vertices shown on Fig. 8 (left); for $t \geq 1$, G_{t+1} is the graph obtained from G_t as in Fig. 8 (right). For $k \geq 1$ the graph G_k is a plane geometric graph with $n = 3k$ internal faces. The three inner-most faces are quadrilaterals while all other faces are hexagons.




Figure 8: Left: Graph G_1 . Right: Graph G_{t+1} obtained from graph G_t (placed upside down inside the dotted triangle) .

Place an hexagon inside each of the three inner-most faces of G_k as in Fig. 9 (left), an octagon inside each outer-most internal face of G_k as in Fig. 9 (right) and a nonagon inside each other internal face of G_k as in Fig. 10.




Figure 9: Hexagon inside inner-most face of G_k . Octagon inside outer-most internal face of G_k .




Figure 10: Nonagon inside internal face of G_k

Since each face of G_k has an even number of sides, the placement of the polygons described above may be done in such a way that for each internal edge e of G_k a side s_e of the polygon R_i lying on one side of e is parallel (and closed enough) to e while a vertex v_e of the polygon R_j lying on the other side of e lies on e as in Fig. 11.

Let $\mathcal{R} = \{R_1, R_2, \dots, R_n\}$ be the set of polygons described above. The total number of sides among polygons in \mathcal{R} is

$$m = 3(6) + 3(8) + (n - 6)(9) = 9n - 12$$

Each side among polygons in \mathcal{R} is relevant with respect to \mathcal{R} , therefore our bound on the number of sides in Lemma 1 is almost tight.




Figure 11: Polygons on both sides of an internal edge of G_k .

3 Main Results

In this section we present our main results.

Lemma 2. *Let $\mathcal{P} = \{P_1, P_2, \dots, P_n\}$ be a collection of $n \geq 3$ pairwise disjoint convex polygons in the plane. There is a side s of a polygon P_i such that the line supporting s separates P_i from a subcollection \mathcal{F} of \mathcal{P} with at least $\frac{n}{18}$ polygons.*

Proof. Let $\mathcal{R} = \{R_1, R_2, \dots, R_n\}$ be a collection of pairwise disjoint convex polygons satisfying conditions 1), 2) and 3) in Lemma 1 and let $\mathcal{L} = \{l_1, l_2, \dots, l_m\}$ be the (multi)set of lines supporting the sides of each polygon in \mathcal{R} . We include c copies of a line l if l supports sides of c polygons in \mathcal{R} . Therefore we can associate a unique polygon $R_{i(k)}$ to each line $l_k \in \mathcal{L}$ such that a side of $R_{i(k)}$ is supported by l_k . For $k = 1, 2, \dots, m$ let H_k^- be the closed halfplane determined by l_k that does not contain $R_{i(k)}$.

Define a bipartite graph F with a vertex v_j for each polygon $R_j \in \mathcal{R}$ and a vertex w_k for each line $l_k \in \mathcal{L}$. Graph F has an edge $v_j w_k$ if polygon R_j is contained in H_k^- .

For each pair of polygons R_i, R_j there is at least one side s of one of them such that the line supporting s separates R_i from R_j . Therefore graph F has at least one edge for each pair $\{i, j\}$ with $1 \leq i < j \leq n$. This implies that there is a vertex w_k whose degree in F is at least $\binom{n}{2}/m \geq \binom{n}{2}/(9n - 9) = n/18$.

This means that the line l_k separates polygon $R_{i(k)}$ from at least $n/18$ polygons in \mathcal{R} . By Property 2) in Lemma 1, line l_k supports a side of polygon $P_{i(k)}$. □

Theorem 2. *In any collection \mathcal{C} of $n \geq 2$ pairwise disjoint compact convex sets in the plane, there is a pair of sets A and B such that any line that separates A from B separates either A or B from a subcollection \mathcal{F} of \mathcal{C} with at least $\frac{n}{18}$ sets.*

Proof. Let $n \geq 2$ be an integer and $\mathcal{C} = \{C_1, C_2, \dots, C_n\}$ be a collection of pairwise disjoint compact convex sets in the plane and let T be a triangle containing all sets in \mathcal{C} in its interior.

For any line l let t_l^- and t_l^+ be the number of sets in \mathcal{C} lying to the left and to the right of l , respectively; for a horizontal line l , t_l^- and t_l^+ are the number of sets in \mathcal{C} lying above and below l , respectively. For $1 \leq i < j \leq n$ let l_{ij} be a line separating C_i from C_j for which $\max\{t_{l_{ij}}^-, t_{l_{ij}}^+\}$ is as small as possible.

Each line l_{ij} determines two closed halfplanes H_{ij} and H_{ji} containing C_i and C_j , respectively. For $i = 1, 2, \dots, n$, let $P_i = T \cap (\bigcap_{j \neq i}^n H_{ij})$. Then For $i = 1, 2, \dots, n$, P_i is a convex polygon that contains set C_i such that each side is supported by a line l_{ij} or by a side of T .

Let $\mathcal{P} = \{P_1, P_2, \dots, P_n\}$. By Lemma 2, there is a side s of a polygon P_i such that the line $l(s)$ supporting s separates P_i from a subcollection of \mathcal{P} with at least $\frac{n}{18}$ polygons.

Since no side of T separates polygons in \mathcal{P} , $l(s)$ is of the form l_{ij} . By the choice of l_{ij} each line that separates sets C_i from C_j separates either C_i or C_j from at least $\frac{n}{18}$ sets in \mathcal{C} . □

References

- [1] Herbert Edelsbrunner, Arch D. Robison and Xiao Jun Shen. Covering convex sets with non-overlapping polygons, *Discrete Mathematics* 81: 153 – 164, 1990.
- [2] László Fejes Tóth. Illumination of convex discs, *Acta Mathematica Academiae Scientiarum Hungaricae* 29(3 - 4): 355 – 360, 1997.

- [3] Rafael Hope and Meir Katchalski. Separating plane convex sets, *Mathematica Scandinavica* 66 (1): 44–46, 1990.
- [4] Eduardo Rivera-Campo and Jeno Töröcsik. On separation of plane convex sets, *European Journal of Combinatorics* 14: 113 – 116, 1993.
- [5] Helge Tverberg. A separation property of plane convex sets, *Mathematica Scandinavica* 45: 255–260, 1979.