

MACLANE-VAQUIÉ CHAINS AND VALUATION-TRANSCENDENTAL EXTENSIONS

SNEHA MAVI AND ANUJ BISHNOI*

ABSTRACT. In this paper, for a valued field (K, v) of arbitrary rank and an extension w of v to $K[X]$, we give a connection between complete sets of ABKPs for w and MacLane-Vaquié chains of w .

1. INTRODUCTION

In 1936, MacLane [3], proved that an extension w of a discrete rank one valuation v of K to $K[X]$ can be obtained as a chain of augmentations

$$w_0 \xrightarrow{\phi_1, \gamma_1} w_1 \xrightarrow{\phi_2, \gamma_2} \cdots \longrightarrow w_{n-1} \xrightarrow{\phi_n, \gamma_n} w_n \longrightarrow \cdots$$

for some suitable key polynomials $\phi_i \in K[X]$ with respect to the intermediate valuations, and values γ_i belonging to some totally ordered abelian group containing the value group of v as an ordered subgroup. Later, Vaquié generalized MacLane's theory to arbitrary valued fields (see [10]). Recently, Nart gave a survey of generalized MacLane-Vaquié theory in [5] and [6]. Starting with a valuation w_0 which admit key polynomials of degree one, Nart also introduced **MacLane-Vaquié chains**, consisting of a mixture of ordinary and limit augmentations (see Definition 2.22). The main result, Theorem 4.3 of [6] says that, all extensions w of v to $K[X]$ fall exactly in one of the following categories:

- (i) It is the last valuation of a complete finite MacLane-Vaquié chain, i.e., after a finite number r of augmentation steps, we get $w_r = w$.
- (ii) After a finite number r of augmentation steps, it is the stable limit of a continuous family of augmentations of w_r defined by key polynomials of constant degree.
- (iii) It is the stable limit of a complete infinite MacLane-Vaquié chain.

In this paper, we study MacLane-Vaquié chains of the first type and prove that a precise complete finite MacLane-Vaquié chain can be obtained using a given complete set $\{Q_i\}_{i \in \Delta}$ of ABKPs for w such that Δ has a last element. Conversely, if w is the last valuation of some complete finite MacLane-Vaquié

Date: 09,19,2022.

2020 Mathematics Subject Classification. 12F20, 12J10, 13A18.

Key words and phrases. Abstract key polynomials, key polynomials, MacLane-Vaquié chains.

*Corresponding author, E-mail address: abishnoi@maths.du.ac.in.

chain, then it is possible to derive, from the data of this chain, a complete set $\{Q_i\}_{i \in \Delta}$ of ABKPs for w such that Δ has a last element.

To state the main results of the paper, we first recall some notation, definitions and preliminary results.

2. BACKGROUND AND NOTATION

Throughout (K, v) denote a valued field of arbitrary rank with value group Γ_v , residue field k_v and by \bar{v} we denote an extension of v to a fixed algebraic closure \bar{K} of K .

An extension w of v to the simple transcendental extension $K(X)$ of K is said to be **value-transcendental** if the quotient Γ_w/Γ_v is a torsion-free group and the corresponding residue field extension $k_w|k_v$ is algebraic. The extension w of v to $K(X)$ is called **residually transcendental** (abbreviated as r. t.) if Γ_w/Γ_v is a torsion group and $k_w|k_v$ is transcendental. We call w **valuation-transcendental** if it is either value-transcendental or is residually transcendental.

We fix a common extension \bar{w} of w and \bar{v} to $\bar{K}(X)$. By [2, Lemma 3.3], \bar{w} is valuation-transcendental if and only if w is valuation-transcendental.

Definition 2.1. For any valuation w' on $K(X)$ taking values in a subgroup of Γ_w , we say that $w' \leq w$ if and only if

$$w'(f) \leq w(f), \quad \forall f \in K[X].$$

If $w' < w$, we denote by $\Phi(w', w)$ the set of all monic polynomials $g \in K[X]$ of minimal degree such that $w'(g) < w(g)$.

Definition 2.2. A pair (α, δ) in $\bar{K} \times \Gamma_{\bar{w}}$ is called a **pair of definition** for w if $\bar{w} = \bar{w}_{\alpha, \delta}$, where $\bar{w}_{\alpha, \delta}$ is a valuation on $\bar{K}[X]$ defined by

$$\bar{w}_{\alpha, \delta} \left(\sum_{i \geq 0} c_i (X - \alpha)^i \right) := \min_{i \geq 0} \{ \bar{v}(c_i) + i\delta \}, \quad c_i \in \bar{K}.$$

Definition 2.3. For any polynomial f in $K[X]$, we denote

$$\delta(f) = \max\{ \bar{w}(X - \alpha) \mid f(\alpha) = 0 \}.$$

This value $\delta(f)$ is independent of the choice of \bar{w} [8, Proposition 3.1].

Definition 2.4 (Abstract key polynomials). A monic polynomial Q in $K[X]$ is said to be an **abstract key polynomial** (abbreviated as ABKP) for w if for each polynomial f in $K[X]$ with $\deg f < \deg Q$ we have $\delta(f) < \delta(Q)$.

It is immediate from the definition that all monic linear polynomials are ABKPs for w . Also an ABKP for w is an irreducible polynomial (see [7, Proposition 2.4]).

Definition 2.5. For a polynomial Q in $K[X]$ the Q -**truncation** of w is a map $w_Q : K[X] \rightarrow \Gamma_w$ defined by

$$w_Q(f) := \min_{0 \leq i \leq n} \{w(f_i Q^i)\},$$

where $f = \sum_{i=0}^n f_i Q^i$, $\deg f_i < \deg Q$, is the Q -expansion of f .

The Q -truncation w_Q of w need not be a valuation [7, Example 2.5]. However, if Q is an ABKP for w , then w_Q is a valuation on $K(X)$ (see [7, Proposition 2.6]). Also any ABKP, Q for w , is also an ABKP for the truncation valuation w_Q .

Lemma 2.6 (Lemma 2.11, [7]). *If Q is an ABKP for w , then every element $F \in \Phi(w_Q, w)$ is also an ABKP for w and $\delta(Q) < \delta(F)$.*

Definition 2.7. A family $\Lambda = \{Q_i\}_{i \in \Delta}$ of ABKPs for w is said to be a **complete set of ABKPs** for w if the following conditions are satisfied:

- (i) $\delta(Q_i) \neq \delta(Q_j)$ for every $i \neq j \in \Delta$.
- (ii) Λ is well-ordered with respect to the ordering given by $Q_i < Q_j$ if and only if $\delta(Q_i) < \delta(Q_j)$ for every $i < j \in \Delta$.
- (iii) For any $f \in K[X]$, there exists some $Q_i \in \Lambda$ such that $\deg Q_i \leq \deg f$ and $w_{Q_i}(f) = w(f)$.

Theorem 2.8 (Theorem 1.1, [7]). *Every valuation w on $K(X)$ admits a complete set of ABKPs for w .*

Remark 2.9. As shown in [4, Remark 4.6] and [7, proof of Theorem 1.1], there is a complete set $\Lambda = \{Q_i\}_{i \in \Delta}$ of ABKPs for w having the following properties.

- (i) $\Delta = \bigcup_{j \in I} \Delta_j$ with $I = \{0, 1, \dots, N\}$ or $\mathbb{N} \cup \{0\}$, and for each $j \in I$ we have $\Delta_j = \{j\} \cup \vartheta_j$, where ϑ_j is an ordered set without last element or is empty.
- (ii) Q_0 is a monic polynomial of degree one.
- (iii) For all $j \in I \setminus \{0\}$ we have $j - 1 < i < j$, for all $i \in \vartheta_{j-1}$.
- (iv) All polynomials Q_i with $i \in \Delta_j$ have the same degree and have degree strictly less than the degree of the polynomials $Q_{i'}$ for every $i' \in \Delta_{j+1}$.
- (v) For each $i < i' \in \Delta$ we have $w(Q_i) < w(Q_{i'})$ and $\delta(Q_i) < \delta(Q_{i'})$.

All complete set of ABKPs used in this paper will be assumed to satisfy these properties.

Even though the set $\{Q_i\}_{i \in \Delta}$ of ABKPs for w is not unique, the cardinality of I and the degree of an abstract key polynomial Q_i for each $i \in I$ are uniquely determined by w .

The ordered set Δ has a last element if and only if the set $I = \{0, 1, \dots, N\}$ is finite and $\Delta_N = \{N\}$, i.e., $\vartheta_N = \emptyset$.

Definition 2.10 (Limit ABKPs). Let $\Lambda = \{Q_i\}_{i \in \Delta}$ be a complete set of ABKPs for w . For an element $i \in I \setminus \{0\}$, we say that Q_i is a **limit ABKP** for w if i has no immediate predecessor in Δ ; that is, $\vartheta_{i-1} \neq \emptyset$.

We now recall the notion of key polynomials which was first introduced by MacLane in 1936 and generalized by Vaquié in 2007 (see [3] and [10]).

Definition 2.11 (Key polynomials). For a valuation w on $K(X)$ and polynomials f, g in $K[X]$, we say that

- (i) f and g are w -equivalent and write $f \sim_w g$ if $w(f - g) > w(f) = w(g)$.
- (ii) g is w -divisible by f (denoted by $f \mid_w g$) if there exists some polynomial $h \in K[X]$ such that $g \sim_w fh$.
- (iii) f is w -irreducible if for any $h, q \in K[X]$, whenever $f \mid_w hq$, then either $f \mid_w h$ or $f \mid_w q$.
- (iv) f is w -minimal if for every nonzero polynomial $h \in K[X]$, whenever $f \mid_w h$, then $\deg h \geq \deg f$.

Any monic polynomial f satisfying (iii) and (iv) is called a (MacLane-Vaquié) **key polynomial** for w .

In view of Proposition 2.10 of [1], any ABKP, Q for w is a key polynomial for w_Q of minimal degree. Let $\text{KP}(w)$ denote the set of all key polynomials for w . For any $\phi \in \text{KP}(w)$, we denote by $[\phi]_w$ the set of all key polynomials which are w -equivalent to ϕ .

The existence of key polynomials is characterized as follows.

Theorem 2.12 (Theorem 4.4, [5]). *A valuation w on $K(X)$ has $\text{KP}(w) \neq \emptyset$ if and only if it is valuation-transcendental.*

Definition 2.13 (Ordinary augmentation). Let ϕ be a key polynomial for a valuation w' on $K(X)$ and let $\gamma > w'(\phi)$ be an element of a totally ordered abelian group Γ containing $\Gamma_{w'}$ as an ordered subgroup. The map $w : K[X] \rightarrow \Gamma \cup \{\infty\}$ defined by

$$w(f) := \min\{w'(f_i) + i\gamma\},$$

where $f = \sum_{i \geq 0} f_i \phi^i$, $\deg f_i < \deg \phi$, is the ϕ -expansion of $f \in K[X]$, gives a valuation on $K(X)$ (see [3, Theorem 4.2]) called the **ordinary augmentation** of w and is denoted by $w = [w'; \phi, \gamma]$.

Clearly, $\phi \in \Phi(w', w)$. On the other hand, the polynomial ϕ is a key polynomial of minimal degree for the augmented valuation w (see [5, Corollary 7.3]).

If ϕ is a key polynomial for w of minimal degree, then we define

$$\deg(w) := \deg \phi.$$

Theorem 2.14 (Theorem 1.15, [10]). *Let w be a valuation on $K(X)$ and $w' < w$. Then any $\phi \in \Phi(w', w)$ is a key polynomial for w' and*

$$w' < [w'; \phi, w(\phi)] \leq w.$$

For any non-zero polynomial $f \in K[X]$, the equality $w'(f) = w(f)$ holds if and only if $\phi \nmid_{w'} f$.

Corollary 2.15 (Corollary 2.5, [6]). *Let $w' < w$ be as above. Then*

- (i) $\Phi(w', w) = [\phi]_{w'}$ for all $\phi \in \Phi(w', w)$.
- (ii) If $w' < \mu \leq w$ is a chain of valuations, then $\Phi(w', w) = \Phi(w', \mu)$. In particular, for all $f \in K[X]$ we have

$$w'(f) = w(f) \iff w'(f) = \mu(f).$$

Let (K, v) be a valued field and w' be an extension of v to $K(X)$. We now recall the definition of a continuous family of augmentations of w' ([10, 5]).

Definition 2.16. Let w' be a valuation-transcendental valuation on $K(X)$. A **continuous family of augmentations** of w' is a family of ordinary augmentations of w'

$$\mathcal{W} = (\rho_i = [w'; \chi_i, \gamma_i])_{i \in \mathbf{A}},$$

indexed by a totally ordered set \mathbf{A} , satisfying the following conditions:

- (i) The set \mathbf{A} has no last element.
- (ii) All key polynomials $\chi_i \in \text{KP}(w')$ have the same degree.
- (iii) For all $i < j$ in \mathbf{A} , χ_j is a key polynomial for ρ_i and satisfies:

$$\chi_j \not\sim_{\rho_i} \chi_i \text{ and } \rho_j = [\rho_i; \chi_j, \gamma_j].$$

The common degree $m = \deg \chi_i$, for all i , is called the **stable degree** of the family \mathcal{W} and is denoted by $\deg(\mathcal{W})$.

A polynomial $f \in K[X]$ is said to be **\mathcal{W} -stable** with respect to the family $\mathcal{W} = (\rho_i)_{i \in \mathbf{A}}$ if

$$\rho_i(f) = \rho_{i_0}(f), \text{ for every } i \geq i_0$$

for some index $i_0 \in \mathbf{A}$. This stable value is denoted by $\rho_{\mathcal{W}}(f)$. By Corollary 2.15 (ii), a polynomial $f \in K[X]$ is **\mathcal{W} -unstable** if and only if

$$\rho_i(f) < \rho_j(f), \quad \forall i < j \in \mathbf{A}.$$

The minimal degree of an \mathcal{W} -unstable polynomial is denoted by m_{∞} . If all polynomials are \mathcal{W} -stable, then we set $m_{\infty} = \infty$.

Remark 2.17. The following properties hold for any continuous family $\mathcal{W} = (\rho_i)_{i \in \mathbf{A}}$ of augmentations (see [6, p. 9]):

- (i) For all $i \in \mathbf{A}$, χ_i is a key polynomial for ρ_i of minimal degree.
- (ii) $\Phi(\rho_i, \rho_j) = [\chi_j]_{\rho_i} \quad \forall i < j \in \mathbf{A}$.
- (iii) All the value groups Γ_{ρ_i} coincide and the common value group is denoted by $\Gamma_{\mathcal{W}}$.

Definition 2.18 (MacLane-Vaquié limit key polynomials). Let \mathcal{W} be a continuous family of augmentations of a valuation w' . A monic \mathcal{W} -unstable polynomial of minimal degree is called a **MacLane-Vaquié limit key polynomial** (abbreviated as MLV) for \mathcal{W} .

We denote by $\text{KP}_{\infty}(\mathcal{W})$ the set of all MLV limit key polynomials. Since the product of stable polynomials is stable, so all MLV limit key polynomials are irreducible in $K[X]$.

Definition 2.19. We say that \mathcal{W} is an **essential continuous family** of augmentations if $\deg(\mathcal{W}) = m < m_\infty < \infty$.

Let \mathcal{W} be an essential continuous family of augmentations of a valuation w' . Then \mathcal{W} admit MLV limit key polynomials. If Q is any MLV limit key polynomial, then any polynomial f in $K[X]$ with $\deg f < \deg Q$ is \mathcal{W} -stable.

Definition 2.20 (Limit augmentation). Let Q be any MLV limit key polynomial for an essential continuous family of augmentations $\mathcal{W} = (\rho_i)_{i \in \mathbf{A}}$ of w' and $\gamma > \rho_i(Q)$, for all $i \in \mathbf{A}$, be an element of a totally ordered abelian group Γ containing $\Gamma_{\mathcal{W}}$ as an ordered subgroup. Then, the map $w : K[X] \rightarrow \Gamma \cup \{\infty\}$ defined by

$$w(f) = \min_{i \geq 0} \{\rho_{\mathcal{W}}(f_i) + i\gamma\},$$

where $f = \sum_{i \geq 0} f_i Q^i$, $\deg f_i < \deg Q$, is the Q -expansion of $f \in K[X]$, gives a valuation on $K(X)$, called the **limit augmentation** of \mathcal{W} and we denote it by $w = [\mathcal{W}; Q, \gamma]$.

Note that $w(Q) = \gamma$ and $\rho_i < w$ for all $i \in \mathbf{A}$. On the other hand, Q is a key polynomial for w of minimal degree [5, Corollary 7.13].

We now recall the definition of MacLane-Vaquié chains given by Nart in [6]. For this, we first consider a finite, or countably infinite, chain of mixed augmentations

$$(2.1) \quad w_0 \xrightarrow{\phi_1, \gamma_1} w_1 \xrightarrow{\phi_2, \gamma_2} \dots \longrightarrow w_n \xrightarrow{\phi_{n+1}, \gamma_{n+1}} w_{n+1} \longrightarrow \dots$$

in which every valuation is an augmentation of the previous one and is of one of the following type:

- Ordinary augmentation: $w_{n+1} = [w_n; \phi_{n+1}, \gamma_{n+1}]$, for some $\phi_{n+1} \in \text{KP}(w_n)$.
- Limit augmentation: $w_{n+1} = [\mathcal{W}_n; \phi_{n+1}, \gamma_{n+1}]$, for some $\phi_{n+1} \in \text{KP}_\infty(\mathcal{W}_n)$, where $\mathcal{W}_n = (\rho_i)_{i \in \mathbf{A}_n}$ is an essential continuous family of augmentations of w_n .

Let $\phi_0 \in \text{KP}(w_0)$ be a key polynomial of minimal degree and let $\gamma_0 = w_0(\phi_0)$. Then, in view of Theorem 2.14, Proposition 6.3 of [5], Proposition 2.1, 3.5 of [6] and Corollary 2.15, we have the following properties of a chain (2.1) of augmentations.

- Remark 2.21.* (i) $\gamma_n = w_n(\phi_n) < \gamma_{n+1}$.
(ii) For all $n \geq 0$, the polynomial ϕ_n is a key polynomial for w_n of minimal degree and therefore

$$\deg(w_n) (= \deg \phi_n) \text{ divides } \deg(\Phi(w_n, w_{n+1})).$$

(iii)

$$\Phi(w_n, w_{n+1}) = \begin{cases} [\phi_{n+1}]_{w_n}, & \text{if } w_n \rightarrow w_{n+1} \text{ is an ordinary augmentation} \\ \Phi(w_n, \rho_i), & \text{for all } i \in \mathbf{A}_n, \text{ if } w_n \rightarrow w_{n+1} \text{ is a limit augmentation.} \end{cases}$$

(iv)

$$\deg(\Phi(w_n, w_{n+1})) = \begin{cases} \deg \phi_{n+1}, & \text{if } w_n \rightarrow w_{n+1} \text{ is an ordinary augmentation} \\ \deg(\mathcal{W}_n), & \text{if } w_n \rightarrow w_{n+1} \text{ is a limit augmentation.} \end{cases}$$

Definition 2.22 (MacLane-Vaquié chains). A finite, or countably infinite chain of mixed augmentations as in (2.1) is called a **MacLane-Vaquié chain** (abbreviated as MLV), if every augmentation step satisfies:

- (i) if $w_n \rightarrow w_{n+1}$ is an ordinary augmentation, then $\deg(w_n) < \deg(\Phi(w_n, w_{n+1}))$.
- (ii) if $w_n \rightarrow w_{n+1}$ is a limit augmentation, then $\deg(w_n) = \deg(\Phi(w_n, w_{n+1}))$ and $\phi_n \notin \Phi(w_n, w_{n+1})$.

An MLV chain is said to be **complete** if w_0 is a depth zero valuation; that is, $w_0 = \bar{w}_{a,\delta}$ for some $a \in K$ and $\delta \in \Gamma$ (see Definition 2.2).

3. STATEMENTS OF MAIN RESULTS

In the following result, we use a complete set of ABKPs for a valuation w to construct a complete finite MLV chain whose last valuation is w .

Theorem 3.1. *Let (K, v) be a valued field and let w be an extension of v to $K(X)$. If $\{Q_i\}_{i \in \Delta}$ is a complete set of ABKPs for w such that N is the last element of Δ , then*

$$(3.1) \quad w_0 \xrightarrow{Q_1, \gamma_1} w_1 \xrightarrow{Q_2, \gamma_2} \cdots \longrightarrow w_{N-1} \xrightarrow{Q_N, \gamma_N} w_N = w,$$

is a complete finite MLV chain of w such that

- (i) if $\vartheta_j = \emptyset$, then $w_j \rightarrow w_{j+1}$ is an ordinary augmentation.
- (ii) if $\vartheta_j \neq \emptyset$, then $w_j \rightarrow w_{j+1}$ is a limit augmentation.

In both cases, $w_{j+1} = w_{Q_{j+1}}$ and $\gamma_{j+1} = w(Q_{j+1})$.

The converse of the above result also holds.

Theorem 3.2. *Let (K, v) be a valued field and let w be an extension of v to $K(X)$. If*

$$(3.2) \quad w_0 \xrightarrow{\phi_1, \gamma_1} w_1 \xrightarrow{\phi_2, \gamma_2} \cdots \longrightarrow w_{N-1} \xrightarrow{\phi_N, \gamma_N} w_N = w,$$

is a complete finite MLV chain of w , then $\Lambda = \{Q_i\}_{i \in \Delta}$ forms a complete set of ABKPs for w , where

- (i) $\Delta = \bigcup_{j \in I} \Delta_j$, with $I = \{0, 1, \dots, N\}$, and $\Delta_j = \{j\} \cup \mathbf{A}_j$ for all $j \in I$. Moreover, $Q_j = \phi_j$ for all $j \in I$.
- (ii) if $w_j \rightarrow w_{j+1}$ is an ordinary augmentation, then $\vartheta_j = \emptyset$.
- (iii) if $w_j \rightarrow w_{j+1}$ is a limit augmentation, with respect to an essential continuous family $\mathcal{W}_j = (\rho_i)_{i \in \mathbf{A}_j}$ of augmentations of w_j , then $\vartheta_j = \mathbf{A}_j$. Moreover, $Q_i = \chi_i$ for all $i \in \mathbf{A}_j$, where χ_i is the key polynomial for w_j such that $\rho_i = [w_j; \chi_i, w(\chi_i)]$.

It is known that if $\{Q_i\}_{i \in \Delta}$ is a complete set of ABKPs for w , then w is a valuation-transcendental extension of v to $K(X)$ if and only if Δ has a last element, say, N , and then $w = w_{Q_N}$ (see [4, Theorem 5.6]). Therefore, as an immediate consequence of Theorems 3.1 and 3.2, we have the following result.

Corollary 3.3. *Let (K, v) and $(K(X), w)$ be as above. Then the following are equivalent:*

- (i) *The extension w is valuation-transcendental.*
- (ii) *There exist a complete set $\{Q_i\}_{i \in \Delta}$ of ABKPs for w such that Δ has a last element.*
- (iii) *The extension w is the last valuation of a complete finite MLV chain.*

Definition 3.4. A valuation w on $K(X)$ is said to be **inductive** if it is the last valuation of a complete finite MLV chain, all whose augmentations are ordinary.

The following result is a consequence of Theorem 3.1 and Theorem 3.2.

Corollary 3.5. *Let (K, v) be a valued field and w an extension of v to $K(X)$. Then w is inductive if and only if it admits a finite complete set of ABKPs.*

4. PROOF OF MAIN RESULTS

Let (K, v) be a valued field and $(\overline{K}, \overline{v})$ be as before. Let w be an extension of v to $K(X)$ and \overline{w} be a common extension of w and \overline{v} to $\overline{K}(X)$. With notations and definitions as in the previous section, we first give some preliminary results which will be used to prove the main results.

In the following result we recall some basic properties of ABKPs for w (see Proposition 3.8, Corollary 3.13 and Theorem 6.1 of [9]).

Proposition 4.1. *For ABKPs, Q and Q' for w the following holds:*

- (i) *If $\delta(Q) < \delta(Q')$, then $w_Q(Q') < w(Q')$.*
- (ii) *If $\deg Q = \deg Q'$, then*

$$w(Q) < w(Q') \iff w_Q(Q') < w(Q') \iff \delta(Q) < \delta(Q').$$

Hence $Q' \in \Phi(w_Q, w)$ in this case.

- (iii) *Suppose that $\delta(Q) < \delta(Q')$. For any polynomial $f \in K[X]$, we have*

$$w_Q(f) \leq w_{Q'}(f) \leq w(f).$$

Moreover, if $w_{Q'}(f) < w(f)$, then $w_Q(f) < w_{Q'}(f)$.

- (iv) *If $Q' \in \Phi(w_Q, w)$, then Q and Q' are key polynomials for w_Q . Moreover, $w_{Q'} = [w_Q; Q', w(Q')]$.*

The following result gives a comparison between key polynomials and ABKPs.

Theorem 4.2 (Theorem 2.17, [1]). *Suppose that $w' < w$ and Q is a key polynomial for w' . Then Q is an ABKP polynomial for w if and only if it satisfies one of the following two conditions:*

- (i) $Q \in \Phi(w', w)$,
- (ii) $Q \notin \Phi(w', w)$ and $\deg Q = \deg w'$.

In the first case $w_Q = [w'; Q, w(Q)]$. In the second case $w_Q = w'$.

Proof of Theorem 3.1. Let $\Lambda = \{Q_i\}_{i \in \Delta}$ be a complete set of ABKPs for w with N the last element of Δ . If $N = 0$, then $w = w_{Q_0}$ is a depth zero valuation and the result holds trivially.

Assume now that $N \geq 1$. Then by Remark 2.9 (i), $\Delta = \bigcup_{j=0}^N \Delta_j$, where $\Delta_j = \{j\} \cup \vartheta_j$ and ϑ_j is either empty or an ordered set without a last element. Since for each $i \in \Delta$, Q_i is an ABKP for w , so w_{Q_i} is a valuation on $K(X)$ and we denote it by w_i . Since $Q_i \notin \Phi(w_i, w)$, Theorem 4.2 shows that Q_i is a key polynomial of minimal degree for w_i . Hence, $\deg(w_i) = \deg Q_i$ for all $i \in \Delta$.

Suppose first that $\vartheta_j = \emptyset$ for some $0 \leq j \leq N - 1$. We first show that $Q_{j+1} \in \Phi(w_j, w)$. Since $\delta(Q_j) < \delta(Q_{j+1})$, Proposition 4.1 (i) shows that $w_j(Q_{j+1}) < w(Q_{j+1})$. Therefore, it is enough to show that for any polynomial $f \in K[X]$ with $\deg f < \deg Q_{j+1}$, we have $w_j(f) = w(f)$. For such a polynomial f , by definition of the complete set Λ of ABKPs, there exists some $Q_i \in \Lambda$ such that $\deg Q_i \leq \deg f$ and

$$w_i(f) = w(f).$$

Since $\deg Q_i < \deg Q_{j+1}$, we have $i < j + 1$. As $\vartheta_j = \emptyset$, we necessarily have $i \leq j$. By Proposition 4.1 (iii) and the above equality, we have

$$w_i(f) \leq w_j(f) \leq w(f) = w_i(f),$$

so that $w_j(f) = w(f)$. This proves that $Q_{j+1} \in \Phi(w_j, w)$.

By Proposition 4.1 (iv), Q_j and Q_{j+1} are key polynomials for w_j and

$$w_{j+1} = [w_j; Q_{j+1}, w(Q_{j+1})]$$

is an ordinary augmentation of w_j . The condition of a an MLV chain is fulfilled because $\deg(w_j) = \deg Q_j < \deg Q_{j+1} = \deg(\Phi(w_j, w_{j+1}))$.

Assume now that $\vartheta_j \neq \emptyset$ for some $0 \leq j < N$. By Proposition 4.1 (i), for each $i < i'$ in ϑ_j the ABKPs $Q_i, Q_{i'}$ satisfies

$$w_j(Q_i) < w(Q_i) = w_i(Q_i), \quad w_i(Q_{i'}) < w(Q_{i'}) = w_{i'}(Q_{i'}).$$

By a completely analogous argument as above, we deduce that $Q_i \in \Phi(w_j, w)$, $Q_{i'} \in \Phi(w_i, w)$ and

$$w_i = [w_j; Q_i, w(Q_i)], \quad w_{i'} = [w_i, Q_{i'}, w(Q_{i'})]$$

are ordinary augmentations. By Corollary 2.15, we have $\Phi(w_i, w) = \Phi(w_i, w_{i'})$ and as $Q_i \notin \Phi(w_i, w)$, so

$$(4.1) \quad w_i(Q_i) = w_{i'}(Q_i), \quad \forall i' > i \text{ in } \vartheta_j,$$

which in view of Theorem 2.14, implies that $Q_{i'} \not\sim_{w_i} Q_i$. Hence $\mathcal{W}_j := (w_i)_{i \in \vartheta_j}$ is a continuous family of augmentations such that for each $i \in \vartheta_j$, w_i is an ordinary augmentation of w_j with respect to $w(Q_i)$ and from (4.1), Q_i is \mathcal{W}_j -stable with stability degree m_j . As $\vartheta_j \neq \emptyset$, so by Definition 2.10, Q_{j+1} is a limit ABKP for w . Since $\delta(Q_i) < \delta(Q_{i'})$, for every $i < i' \in \vartheta_j$, and $w_{i'}(Q_{j+1}) < w(Q_{j+1})$ for every $i' \in \vartheta_j$, (because $i' < j+1 \in \Delta$), so in view of Proposition 4.1 (iii), we have that

$$w_i(Q_{j+1}) < w_{i'}(Q_{j+1}) \text{ for every } i < i' \in \vartheta_j.$$

We now show that Q_{j+1} has minimal degree with this property. For any polynomial $f \in K[X]$ with $\deg f < \deg Q_{j+1}$, there exists some ABKP $Q_{i_0} \in \Lambda$ such that $\deg Q_{i_0} \leq \deg f$ and $w_{i_0}(f) = w(f)$. Clearly, $i_0 < j+1$. Therefore, for any $i, i' \in \vartheta_j$ with $i_0 < i < i'$, Proposition 4.1 (iii) shows that

$$w_{i_0}(f) \leq w_i(f) \leq w_{i'}(f) \leq w(f) = w_{i_0}(f),$$

so that f is \mathcal{W}_j -stable. Therefore, Q_{j+1} is a an MLV limit key polynomial for \mathcal{W}_j . Now, since $\deg(\mathcal{W}_j) = \deg Q_j < \deg Q_{j+1}$, we see that \mathcal{W}_j is an essential continuous family of augmentations of w_j .

Let γ_{j+1} denote the value $w(Q_{j+1})$. Clearly, $\gamma_{j+1} > w_i(Q_{j+1})$ for all $i \in \vartheta_j$. Let

$$\rho_{j+1} = [\mathcal{W}_j; Q_{j+1}, \gamma_{j+1}]$$

be the limit augmentation determined by these data. Since Q_{j+1} is a key polynomial of minimal degree for ρ_{j+1} , Theorem 4.2 shows that ρ_{j+1} is the truncation

$$\rho_{j+1} = w_{Q_{j+1}} = w_{j+1}.$$

Thus, the step $w_j \rightarrow w_{j+1}$ is a limit augmentation. The condition of a an MLV chain is fulfilled because $Q_j \notin \Phi(w_j, w) = \Phi(w_j, w_{j+1})$, the last equality follows from Corollary 2.15.

Clearly, $w_N = w$, for if there exists some polynomial $f \in K[X]$ such that $w_N(f) < w(f)$, then as Λ is a complete set, so $w_i(f) = w(f)$ for some $0 \leq i < N$. But this will imply that $w(f) = w_i(f) \leq w_N(f) < w(f)$.

Thus from the above arguments it follows that

$$w_0 \xrightarrow{Q_1, \gamma_1} w_1 \xrightarrow{Q_2, \gamma_2} \cdots \longrightarrow w_{N-1} \xrightarrow{Q_N, \gamma_N} w_N = w,$$

where $w_i = w_{Q_i}$, $\gamma_i = w(Q_i)$ for $0 \leq i \leq N$, is a an MLV chain whose last valuation is $w_N = w$, such that:

- if $\vartheta_j = \emptyset$, then $w_j \rightarrow w_{j+1}$ is an ordinary augmentation.
- if $\vartheta_j \neq \emptyset$, then $w_j \rightarrow w_{j+1}$ is a limit augmentation of an essential continuous family $\mathcal{W}_j = (w_i)_{i \in \vartheta_j}$ of augmentations of w_j .

Finally the chain is complete because $w_0 = w_{Q_0}$, where $Q_0 = X$, defined by the pair $(0, w_0(X))$ and is a depth zero valuation. \square

Proof of Theorem 3.2. Let

$$w_0 \xrightarrow{\phi_1, \gamma_1} w_1 \xrightarrow{\phi_2, \gamma_2} \cdots \longrightarrow w_{N-1} \xrightarrow{\phi_N, \gamma_N} w_N = w,$$

be a complete finite MLV chain of w . If $N = 0$, then $w_0 = w$ is a depth zero valuation and the result holds trivially.

Assume now that $N \geq 1$. Then each ϕ_j is a key polynomial for w_j of minimal degree, i.e., $\deg \phi_j = \deg(w_j)$, and $w_j(\phi_j) = w(\phi_j)$. Therefore $\phi_j \notin \Phi(w_j, w)$ which in view of Theorem 4.2, implies that ϕ_j is an ABKP for w and

$$(4.2) \quad w_j = w_{\phi_j} \text{ for } 0 \leq j \leq N.$$

Suppose first that $w_j \rightarrow w_{j+1}$ is an ordinary augmentation. Then by definition of MLV chain of w , we have $\phi_{j+1} \in \Phi(w_j, w_{j+1}) = \Phi(w_j, w)$, the last equality by Corollary 2.15. Now, keeping in mind (4.2), this implies that $\phi_{j+1} \in \Phi(w_{\phi_j}, w)$ and hence from Lemma 2.6, it follows that

$$\delta(\phi_j) < \delta(\phi_{j+1}).$$

Assume now that $w_j \rightarrow w_{j+1}$ is a limit augmentation. Then ϕ_{j+1} is an MLV limit key polynomial for an essential continuous family (say) \mathcal{W}_j of augmentations of w_j . Let $\mathcal{W}_j = (\rho_i)_{i \in \mathbf{A}_j}$, where \mathbf{A}_j is some totally ordered set without a last element and for each $i \in \mathbf{A}_j$, $\rho_i = [w_j, \chi_i, \gamma_i]$ is an ordinary augmentation of w_j with stability degree, (say) $m_j = \deg \phi_j = \deg \chi_i$. Also, for all $i < i' \in \mathbf{A}_j$, $\chi_{i'}$ is a key polynomial for ρ_i such that

$$\chi_{i'} \not\prec_{\rho_i} \chi_i, \text{ i.e., } \chi_{i'} \uparrow_{\rho_i} \chi_i \text{ and } \rho_{i'} = [\rho_i; \chi_{i'}, \gamma_{i'}].$$

Since $\rho_i < w$ and $\chi_{i'} \uparrow_{\rho_i} \chi_i$, so by Theorem 2.14, $\rho_i(\chi_{i'}) = w(\chi_{i'})$, which together with Corollary 2.15, gives

$$(4.3) \quad \chi_i \notin \Phi(\rho_i, w) = \Phi(\rho_i, \rho_{i'}) = [\chi_{i'}]_{\rho_i}.$$

Now by Remark 2.17 (i), for each $i \in \mathbf{A}_j$, χ_i is a key polynomial for ρ_i of minimal degree, i.e., $\deg \chi_i = \deg(\rho_i)$, therefore keeping in mind that $\rho_i < w$, equation (4.3) in view of Theorem 4.2, implies that each $Q_i := \chi_i$ is an ABKP for w and

$$(4.4) \quad \rho_i = w_{Q_i}, \text{ for all } i \in \mathbf{A}_j.$$

Hence for each $i < i' \in \mathbf{A}_j$, Q_i and $Q_{i'}$ are ABKPs for w such that

$$w(Q_i) = w_{Q_i}(Q_i) < w_{Q_{i'}}(Q_{i'}) = w(Q_{i'}) \text{ and } \deg Q_i = \deg Q_{i'},$$

which in view of Proposition 4.1 (ii), implies that $w_{Q_i}(Q_{i'}) < w(Q_{i'})$. Therefore, by Lemma 2.6, we get that

$$Q_{i'} \in \Phi(w_{Q_i}, w) \text{ and } \delta(Q_i) < \delta(Q_{i'}) \text{ for every } i < i' \in \mathbf{A}_j.$$

As $\deg \phi_j = \deg Q_i$, for every $i \in \mathbf{A}_j$ and $w(\phi_j) < w(Q_i)$, so again by Proposition 4.1 (ii), we have that $Q_i \in \Phi(w_{\phi_j}, w)$ and then

$$\delta(\phi_j) < \delta(Q_i) \forall i \in \mathbf{A}_j,$$

follows from Lemma 2.6. Since \mathcal{W}_j is essential, so $\deg \phi_j = \deg Q_i < \deg \phi_{j+1}$, for every $i \in \mathbf{A}_j$, this together with the fact that ϕ_{j+1} is an ABKP

for w , implies that

$$\delta(\phi_j) < \delta(\phi_{j+1}), \delta(Q_i) < \delta(\phi_{j+1}) \text{ and } \phi_{j+1} \notin \Phi(w_{\phi_j}, w).$$

For every $0 \leq j \leq N$, let $\Delta_j = \{j\} \cup \mathbf{A}_j$, and $\Delta = \bigcup_{j=0}^N \Delta_j$. We now set $Q_j := \phi_j$ for all $j \in I = \{0, 1, \dots, N\}$, and show that $\Lambda = \{Q_i\}_{i \in \Delta}$ is a complete set of ABKPs for w . Clearly, as shown above for every $i < i' \in \Delta$, we have $\delta(Q_i) < \delta(Q_{i'})$. Therefore, the set Λ is well-ordered with respect to the ordering given by: $Q_i < Q_{i'}$ if and only if $\delta(Q_i) < \delta(Q_{i'})$ for every $i < i' \in \Delta$. Let $f \in K[X]$ be any polynomial. Then on using the fact that $w_N = w$ and (4.2), we get $w_{Q_N} = w$; that is $w_{Q_N}(f) = w(f)$. If $\deg Q_N \leq \deg f$, then we are done.

So suppose that $\deg Q_j \leq \deg f < \deg Q_{j+1}$, for some $j \in I \setminus \{N\}$. If $w_j \rightarrow w_{j+1}$ is an ordinary augmentation, then by Remark 2.21 (iii) and (iv), we have that $f \notin \Phi(w_{Q_j}, w_{Q_{j+1}})$, which together with Corollary 2.15 implies that $f \notin \Phi(w_{Q_j}, w)$ and consequently, $w_{Q_j}(f) = w(f)$. Now if $w_j \rightarrow w_{j+1}$ is a limit augmentation, then as Q_{j+1} is a \mathcal{W}_j -unstable polynomial of minimal degree, so we must have that $\rho_i(f) = \rho_{i'}(f)$ for all $i > i' \in \mathbf{A}_j$, for some index $i' \in \mathbf{A}_j$, which in view of (4.4) implies that $w_{Q_i}(f) = w_{Q_{i'}}(f)$, i.e., $f \notin \Phi(w_{Q_{i'}}, w_{Q_i}) = \Phi(w_{Q_{i'}}, w)$ (see Corollary 2.15). Hence $w_{Q_{i'}}(f) = w(f)$, for $i' \in \mathbf{A}_j$. Thus $\{Q_i\}_{i \in \Delta}$ is a complete set of ABKPs for w such that

- if $w_j \rightarrow w_{j+1}$ is an ordinary augmentation, then $\vartheta_j = \emptyset$.
- if $w_j \rightarrow w_{j+1}$ is a limit augmentation, with respect to an essential continuous family $\mathcal{W}_j = (\rho_i)_{i \in \mathbf{A}_j}$ of augmentations of w_j , then $\vartheta_j = \mathbf{A}_j$, $Q_i = \chi_i$ for all $i \in \mathbf{A}_j$, and $Q_{j+1} = \phi_{j+1}$ is a limit ABKP for w .

□

ACKNOWLEDGEMENT

We would like to thank the anonymous referee for a careful reading and providing useful suggestions which led to an improvement in the presentation of the paper. The research of first author is supported by CSIR (Grant No. 09/045(1747)/2019-EMR-I).

REFERENCES

- [1] M. Alberich-Carramināna, A. F. F. Boix, J. Fernández, J. Guàrdia, E. Nart and J. Roé, Of limit key polynomials. *Illinois J. Math.* **65** (2021), no. 1, 201–229.
- [2] F. -V. Kuhlmann, Value groups, residue fields, and bad places of rational function fields. *Trans. Amer. Math. Soc.* **356** (2004), no. 11, 4559–4600.
- [3] S. MacLane, A construction for absolute values in polynomial rings. *Trans. Amer. Math. Soc.* **40** (1936), no. 3, 363–395.
- [4] W. Mahboub, A. Mansour and M. Spivakovsky, On common extensions of valued fields. *J. Algebra* **584** (2021), 1–18.
- [5] E. Nart, Key polynomials over valued fields. *Publ. Mat.* **64** (2020), No. 1, 195–232.
- [6] E. Nart, MacLane-Vaquíé chains of valuations on a polynomial ring. *Pac. J. Math.* **311** (2021), 165–195.
- [7] J. Novacoski and M. Spivakovsky, Key polynomials and pseudo-convergent sequences. *J. Algebra* **495** (2018), 199–219.

- [8] J. Novacoski, Key polynomials and minimal pairs. *J. Algebra* **523** (2019), 1-14.
- [9] J. Novacoski, On MacLane-Vaquié key polynomials. *J. Pure Appl. Algebra* **225** (2021), no. 8, 106644.
- [10] M. Vaquié, Extension d'une valuation. *Trans. Amer. Math. Soc.* **359** (2007), no. 7, 3439-3481

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI-110007, INDIA.
Email address: `mavisneha@gmail.com`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF DELHI, DELHI-110007, INDIA.
Email address: `abishnoi@maths.du.ac.in`