

Integer group determinants for C_2^4

Yuka Yamaguchi and Naoya Yamaguchi

October 3, 2022

Abstract

We determine all possible values of the integer group determinant of C_2^4 , where C_2 is the cyclic group of order 2.

1 Introduction

For a finite group G , assigning a variable x_g for each $g \in G$, the group determinant of G is defined as $\det(x_{gh^{-1}})_{g,h \in G}$. When the variables x_g are all integers, the group determinant is called an integer group determinant of G . Let $S(G)$ denote the set of all possible values of the integer group determinant of G :

$$S(G) := \left\{ \det(x_{gh^{-1}})_{g,h \in G} \mid x_g \in \mathbb{Z} \right\}.$$

Let C_2 be the cyclic group of order 2. The complete descriptions of $S(C_2^2)$ and $S(C_2^3)$ are obtained in [1, Theorem 5.3] and [4, Theorem 3.1], respectively:

- $S(C_2^2) = \{4m + 1, 2^4(2m + 1), 2^6m \mid m \in \mathbb{Z}\};$
- $S(C_2^3) = \{8m + 1, 2^8(4m + 1), 2^{12}m \mid m \in \mathbb{Z}\}.$

In this paper, we determine $S(C_2^4)$.

Theorem 1.1. *Let $A := \{(8k - 3)(8l + 3) \mid k, l \in \mathbb{Z}\} \subsetneq \{8m - 1 \mid m \in \mathbb{Z}\}$. Then we have*

$$S(C_2^4) = \{16m + 1, 2^{16}(4m + 1), 2^{24}(4m + 1), 2^{24}(8m + 3), 2^{24}m', 2^{26}m \mid m \in \mathbb{Z}, m' \in A\}.$$

For every group G of order at most 15, $S(G)$ was determined (see [3], [4]). For the groups of order 16, the complete descriptions of $S(G)$ were obtained for the dihedral group [1, Theorem 5.3] and the cyclic group [6]. There are fourteen groups of order 16 up to isomorphism [2], [7]. Theorem 1.1 determines $S(G)$ for one of the unsolved twelve groups.

2 Relations with group determinants of subgroups

For any $g = (\varepsilon_0, \varepsilon_1, \dots, \varepsilon_{n-1}) \in C_2^n$ with $\varepsilon_i \in \{0, 1\}$, we denote the variable x_g by x_j , where $j := \varepsilon_{n-1} \cdot 2^{n-1} + \varepsilon_{n-2} \cdot 2^{n-2} + \dots + \varepsilon_0 \cdot 2^0$, and let

$$D_n(x_0, x_1, \dots, x_{2^n-1}) := \det (x_{gh^{-1}})_{g, h \in C_2^n}.$$

From the $H = C_2^{n-1}$ and $K = C_2$ case of [5, Theorem 1.1], we have the following corollary.

Corollary 2.1. *For any positive integer n ,*

$$D_{n+1}(x_0, \dots, x_{2^{n+1}-1}) = D_n(x_0 + x_{2^n}, \dots, x_{2^n-1} + x_{2^{n+1}-1}) D_n(x_0 - x_{2^n}, \dots, x_{2^n-1} - x_{2^{n+1}-1}).$$

Throughout this paper, we assume that $a_0, a_1, \dots, a_{15} \in \mathbb{Z}$ and, for any $0 \leq i \leq 3$, put

$$\begin{aligned} b_i &:= (a_i + a_{i+8}) + (a_{i+4} + a_{i+12}), & c_i &:= (a_i + a_{i+8}) - (a_{i+4} + a_{i+12}), \\ d_i &:= (a_i - a_{i+8}) + (a_{i+4} - a_{i+12}), & e_i &:= (a_i - a_{i+8}) - (a_{i+4} - a_{i+12}). \end{aligned}$$

The following relations will be frequently used in this paper:

$$\begin{aligned} D_4(a_0, a_1, \dots, a_{15}) &= D_3(a_0 + a_8, \dots, a_7 + a_{15}) D_3(a_0 - a_8, \dots, a_7 - a_{15}), \\ &= D_2(b_0, b_1, b_2, b_3) D_2(c_0, c_1, c_2, c_3) D_2(d_0, d_1, d_2, d_3) D_2(e_0, e_1, e_2, e_3). \end{aligned}$$

Remark 2.2. *The following hold:*

- (1) $b_i \equiv c_i \equiv d_i \equiv e_i \pmod{2}$ for $0 \leq i \leq 3$;
- (2) $b_i + c_i + d_i + e_i \equiv 0 \pmod{4}$ for $0 \leq i \leq 3$.

Noting that $\det(\alpha_{st})_{s,t} \equiv \det(\beta_{st})_{s,t} \pmod{2}$ holds if $\alpha_{st} \equiv \beta_{st} \pmod{2}$ for any s and t , we have the following lemma.

Lemma 2.3. *The following hold:*

- (1) $D_4(a_0, a_1, \dots, a_{15}) \equiv D_3(a_0 + a_8, \dots, a_7 + a_{15}) \equiv D_3(a_0 - a_8, \dots, a_7 - a_{15}) \pmod{2}$;
- (2) $D_4(a_0, a_1, \dots, a_{15}) \equiv D_2(b_0, b_1, b_2, b_3) \equiv D_2(c_0, c_1, c_2, c_3) \equiv D_2(d_0, d_1, d_2, d_3) \equiv D_2(e_0, e_1, e_2, e_3) \pmod{2}$.

Let \mathbb{Z}_{odd} be the set of all odd numbers.

Lemma 2.4. *We have $D_4(a_0, a_1, \dots, a_{15}) \in 2\mathbb{Z} \iff b_0 + b_2 \equiv b_1 + b_3 \pmod{2}$.*

Proof. From Lemma 2.3 (2) and

$$\begin{aligned} D_2(b_0, b_1, b_2, b_3) &= D_1(b_0 + b_2, b_1 + b_3) D_1(b_0 - b_2, b_1 - b_3) \\ &= \{(b_0 + b_2)^2 - (b_1 + b_3)^2\} \{(b_0 - b_2)^2 - (b_1 - b_3)^2\} \\ &\in \begin{cases} 2\mathbb{Z}, & b_0 + b_2 \equiv b_1 + b_3 \pmod{2}, \\ \mathbb{Z}_{\text{odd}}, & b_0 + b_2 \not\equiv b_1 + b_3 \pmod{2}, \end{cases} \end{aligned}$$

the lemma is proved. □

3 Group determinant of C_2^2

In this section, we prove five lemmas which give properties of the group determinant of C_2^2 . These lemmas are used in the next section.

By direct calculation, we have the following lemma.

Lemma 3.1. *The identity $D_2(x_0, x_1, x_2, x_3) = \sum_{i=0}^3 x_i^4 - 2 \sum_{0 \leq i < j \leq 3} x_i^2 x_j^2 + 8x_0 x_1 x_2 x_3$ holds. Therefore, $D_2(x_0, x_1, x_2, x_3)$ is a symmetric polynomial in x_0, x_1, x_2, x_3 .*

Lemma 3.2. *For any $k, l, m, n \in \mathbb{Z}$, the following hold:*

- (1) $D_2(2k, 2l, 2m, 2n + 1) \equiv 8(k + l + m) + 1 \pmod{16}$;
- (2) $D_2(2k, 2l + 1, 2m + 1, 2n + 1) \equiv 8k - 3 \pmod{16}$.

Proof. We have

$$\begin{aligned}
& D_2(2k, 2l, 2m, 2n + 1) \\
&= D_1(2k + 2m, 2l + 2n + 1) D_1(2k - 2m, 2l - 2n - 1) \\
&= \{4(k + m)^2 - 4(l + n)^2 - 4(l + n) - 1\} \{4(k - m)^2 - 4(l - n)^2 + 4(l - n) - 1\} \\
&\equiv -4(k + m)^2 + 4(l + n)^2 + 4(l + n) - 4(k - m)^2 + 4(l - n)^2 - 4(l - n) + 1 \\
&\equiv -8k^2 - 8m^2 + 8l^2 + 8n(n + 1) + 1 \\
&\equiv 8(k + l + m) + 1 \pmod{16}, \\
& D_2(2k, 2l + 1, 2m + 1, 2n + 1) \\
&= D_1(2k + 2m + 1, 2l + 2n + 2) D_1(2k - 2m - 1, 2l - 2n) \\
&= \{4(k + m)^2 + 4(k + m) + 1 - 4(l + n + 1)^2\} \{4(k - m)^2 - 4(k - m) + 1 - 4(l - n)^2\} \\
&\equiv 4(k + m)^2 + 4(k + m) - 4(l + n + 1)^2 + 4(k - m)^2 - 4(k - m) - 4(l - n)^2 + 1 \\
&\equiv 8k^2 + 8m(m + 1) - 8l(l + 1) - 8n(n + 1) - 3 \\
&\equiv 8k - 3 \pmod{16}.
\end{aligned}$$

□

Lemma 3.3. *For any $k, l, m, n \in \mathbb{Z}$, let $\alpha := D_2(2k, 2l, 2m, 2n)$. The following hold:*

- (1) *If $k + m \not\equiv l + n$, $km \equiv ln \pmod{2}$, then $\alpha \in \{2^4(8a + 1) \mid a \in \mathbb{Z}\}$;*
- (2) *If $k + m \not\equiv l + n$, $km \not\equiv ln \pmod{2}$, then $\alpha \in \{2^4(8a - 3) \mid a \in \mathbb{Z}\}$;*
- (3) *If k, l, m, n are even and $k + m \not\equiv l + n \pmod{4}$, then $\alpha \in \{2^8(4a + 1) \mid a \in \mathbb{Z}\}$;*
- (4) *If k, l, m, n are odd and $k + m \not\equiv l + n \pmod{4}$, then $\alpha \in \{2^8(4a - 1) \mid a \in \mathbb{Z}\}$;*
- (5) *If $k \equiv l \equiv m \equiv n \pmod{2}$ and $k + m \equiv l + n \pmod{4}$, then $\alpha \in 2^{12}\mathbb{Z}$;*
- (6) *If exactly two of k, l, m, n are even numbers, then $\alpha \in 2^{10}\mathbb{Z}$.*

Proof. First, we prove (1) and (2). We have $(k+m)^2 - (l+n)^2 = (k-m)^2 - (l-n)^2 + 4(km - ln)$. Therefore, if $k + m \not\equiv l + n \pmod{2}$, then

$$\begin{aligned} 2^{-4}\alpha &= \{(k+m)^2 - (l+n)^2\} \{(k-m)^2 - (l-n)^2\} \\ &\equiv \begin{cases} 1 \pmod{8}, & km \equiv ln \pmod{2}, \\ -3 \pmod{8}, & km \not\equiv ln \pmod{2}. \end{cases} \end{aligned}$$

Second, we prove (3). If k, l, m, n are even and $k + m \not\equiv l + n \pmod{4}$, then there exist $k', l', m', n' \in \mathbb{Z}$ satisfying $k = 2k', l = 2l', m = 2m', n = 2n'$ and $k' + m' \not\equiv l' + n' \pmod{2}$. Therefore,

$$2^{-8}\alpha = \{(k' + m')^2 - (l' + n')^2\} \{(k' - m')^2 - (l' - n')^2\} \equiv 1 \pmod{4}.$$

Third, we prove (4). If k, l, m, n are odd and $k + m \not\equiv l + n \pmod{4}$, then there exist $k', l', m', n' \in \mathbb{Z}$ satisfying $k = 2k' + 1, l = 2l' + 1, m = 2m' + 1, n = 2n' + 1$ and $k' + m' \not\equiv l' + n' \pmod{2}$. Therefore,

$$2^{-8}\alpha = \{(k' + m' + 1)^2 - (l' + n' + 1)^2\} \{(k' - m')^2 - (l' - n')^2\} \equiv -1 \pmod{4}.$$

Fourth, we prove (5). If $k \equiv l \equiv m \equiv n \pmod{2}$ and $k + m \equiv l + n \pmod{4}$, then $(k+m)^2 - (l+n)^2 \equiv (k-m)^2 - (l-n)^2 \equiv 0 \pmod{16}$. Therefore,

$$\alpha = 2^4 \{(k+m)^2 - (l+n)^2\} \{(k-m)^2 - (l-n)^2\} \in 2^{12}\mathbb{Z}.$$

Finally, we prove (6). If $(k, l, m, n) \equiv (0, 0, 1, 1) \pmod{2}$, then $(k+m)^2 - (l+n)^2 \equiv (k-m)^2 - (l-n)^2 \equiv 0 \pmod{8}$. Therefore,

$$\alpha = 2^4 \{(k+m)^2 - (l+n)^2\} \{(k-m)^2 - (l-n)^2\} \in 2^{10}\mathbb{Z}.$$

Lemma 3.1 completes the proof of (6). □

Lemma 3.4. *For any $k, l, m, n \in \mathbb{Z}$, let $\alpha := D_2(2k, 2l + 1, 2m, 2n + 1)$. The following hold:*

- (1) *If $k \equiv m \equiv 0 \pmod{2}$ and (I) hold, then $\alpha \in \{2^6(8a - 1)(4b - 1) \mid a, b \in \mathbb{Z}\}$;*
- (2) *If $k \equiv m \equiv 0 \pmod{2}$ and (II) hold, then $\alpha \in \{2^6(8a + 3)(4b + 1) \mid a, b \in \mathbb{Z}\}$;*
- (3) *If $k \equiv m \equiv 1 \pmod{2}$ and (I) hold, then $\alpha \in \{2^6(8a + 3)(4b - 1) \mid a, b \in \mathbb{Z}\}$;*
- (4) *If $k \equiv m \equiv 1 \pmod{2}$ and (II) hold, then $\alpha \in \{2^6(8a - 1)(4b + 1) \mid a, b \in \mathbb{Z}\}$,*

where

(I) $k + m \equiv 1 - l - n \equiv 0 \pmod{4}$ or $k - m \equiv 2 - l + n \equiv 0 \pmod{4}$;

(II) $k + m \equiv 1 - l - n \equiv 2 \pmod{4}$ or $k - m \equiv 2 - l + n \equiv 2 \pmod{4}$.

Proof. Note that

$$\begin{aligned}\alpha &= D_1(2k + 2m, 2l + 2n + 2)D_1(2k - 2m, 2l - 2n) \\ &= 2^4 \{(k + m)^2 - (l + n + 1)^2\} \{(k - m)^2 - (l - n)^2\}.\end{aligned}$$

If $k \equiv m \equiv 0 \pmod{2}$, then

$$(k + m)^2 - (l + n + 1)^2 \equiv \begin{cases} -4 \pmod{16}, & \text{if } k + m \equiv 1 - l - n \equiv 0 \pmod{4}, \\ 4 \pmod{16}, & \text{if } k + m \equiv 1 - l - n \equiv 2 \pmod{4}, \\ -1 \pmod{8}, & \text{if } k - m \equiv 2 - l + n \equiv 0 \pmod{4}, \\ 3 \pmod{8}, & \text{if } k - m \equiv 2 - l + n \equiv 2 \pmod{4}, \end{cases}$$

$$(k - m)^2 - (l - n)^2 \equiv \begin{cases} -1 \pmod{8}, & \text{if } k + m \equiv 1 - l - n \equiv 0 \pmod{4}, \\ 3 \pmod{8}, & \text{if } k + m \equiv 1 - l - n \equiv 2 \pmod{4}, \\ -4 \pmod{16}, & \text{if } k - m \equiv 2 - l + n \equiv 0 \pmod{4}, \\ 4 \pmod{16}, & \text{if } k - m \equiv 2 - l + n \equiv 2 \pmod{4}. \end{cases}$$

If $k \equiv m \equiv 1 \pmod{2}$, then

$$(k + m)^2 - (l + n + 1)^2 \equiv \begin{cases} -4 \pmod{16}, & \text{if } k + m \equiv 1 - l - n \equiv 0 \pmod{4}, \\ 4 \pmod{16}, & \text{if } k + m \equiv 1 - l - n \equiv 2 \pmod{4}, \\ 3 \pmod{8}, & \text{if } k - m \equiv 2 - l + n \equiv 0 \pmod{4}, \\ -1 \pmod{8}, & \text{if } k - m \equiv 2 - l + n \equiv 2 \pmod{4}, \end{cases}$$

$$(k - m)^2 - (l - n)^2 \equiv \begin{cases} 3 \pmod{8}, & \text{if } k + m \equiv 1 - l - n \equiv 0 \pmod{4}, \\ -1 \pmod{8}, & \text{if } k + m \equiv 1 - l - n \equiv 2 \pmod{4}, \\ -4 \pmod{16}, & \text{if } k - m \equiv 2 - l + n \equiv 0 \pmod{4}, \\ 4 \pmod{16}, & \text{if } k - m \equiv 2 - l + n \equiv 2 \pmod{4}. \end{cases}$$

From the above, the lemma is proved. □

Note that for any integers a, b and c , where b is odd, it holds that

$$ab \equiv c \pmod{8} \iff a \equiv bc \pmod{8}$$

since $b^2 \equiv 1 \pmod{8}$. From this, we have the following.

Remark 3.5. For any $k, l, m, n \in \mathbb{Z}$, the following hold:

- (1) $(2k + 2l + 1)(2m + 2n + 1) \equiv 1 \pmod{8} \iff k - m \equiv -l + n \pmod{4}$;
- (2) $(2k + 2l + 1)(2m + 2n + 1) \equiv -1 \pmod{8} \iff k + m \equiv -l - n - 1 \pmod{4}$;
- (3) $(2k + 2l + 1)(2m + 2n + 1) \equiv 3 \pmod{8} \iff k + m \equiv 1 - l - n \pmod{4}$;

$$(4) \quad (2k + 2l + 1)(2m + 2n + 1) \equiv -3 \pmod{8} \iff k - m \equiv 2 - l + n \pmod{4}.$$

Lemma 3.6. *For any $k, l, m, n \in \mathbb{Z}$, the following hold:*

$$(1) \quad D_2(2k, 2l, 2m, 2n) \in \begin{cases} 2^4\mathbb{Z}_{\text{odd}}, & k + m \not\equiv l + n \pmod{2}, \\ 2^8\mathbb{Z}_{\text{odd}} \cup 2^{10}\mathbb{Z}, & k + m \equiv l + n \pmod{2}; \end{cases}$$

$$(2) \quad D_2(2k + 1, 2l + 1, 2m + 1, 2n + 1) \in \begin{cases} 2^4\mathbb{Z}_{\text{odd}}, & k + m \not\equiv l + n \pmod{2}, \\ 2^9\mathbb{Z}, & k + m \equiv l + n \pmod{2}; \end{cases}$$

$$(3) \quad D_2(2k, 2l + 1, 2m, 2n + 1) \in \begin{cases} 2^7\mathbb{Z}, & k \not\equiv m \pmod{2}, \\ 2^8\mathbb{Z}, & k \equiv m \pmod{2} \text{ and } (2k + 2l + 1)(2m + 2n + 1) \equiv \pm 1 \pmod{8}, \\ 2^6\mathbb{Z}_{\text{odd}}, & k \equiv m \pmod{2} \text{ and } (2k + 2l + 1)(2m + 2n + 1) \equiv \pm 3 \pmod{8}. \end{cases}$$

Proof. From Lemma 3.3, it follows that (1) holds. Also, we obtain (2) from

$$\begin{aligned} D_2(2k + 1, 2l + 1, 2m + 1, 2n + 1) &= D_1(2k + 2m + 2, 2l + 2n + 2)D_1(2k - 2m, 2l - 2n) \\ &= 2^4 \{(k + m + 1)^2 - (l + n + 1)^2\} \{(k - m)^2 - (l - n)^2\}. \end{aligned}$$

Below, we prove (3). Let $\alpha := D_2(2k, 2l + 1, 2m, 2n + 1)$. Note that

$$\begin{aligned} \alpha &= D_1(2k + 2m, 2l + 2n + 2)D_1(2k - 2m, 2l - 2n) \\ &= 2^4 \{(k + m)^2 - (l + n + 1)^2\} \{(k - m)^2 - (l - n)^2\}. \end{aligned}$$

If $k \not\equiv m \pmod{2}$, then $\alpha \in 2^7\mathbb{Z}$. If $k \equiv m \pmod{2}$ and $(2k + 2l + 1)(2m + 2n + 1) \equiv 1 \pmod{8}$, then $(k - m)^2 - (l - n)^2 \in 2^4\mathbb{Z}$ from Remark 3.5 (1). Thus, $\alpha \in 2^8\mathbb{Z}$. If $k \equiv m \pmod{2}$ and $(2k + 2l + 1)(2m + 2n + 1) \equiv -1 \pmod{8}$, then $(k + m)^2 - (l + n + 1)^2 \in 2^4\mathbb{Z}$ from Remark 3.5 (2). Thus, $\alpha \in 2^8\mathbb{Z}$. The cases of $k \equiv m \pmod{2}$ and $(2k + 2l + 1)(2m + 2n + 1) \equiv \pm 3 \pmod{8}$ are proved from Lemma 3.4 and Remark 3.5 (3) and (4). \square

4 Impossible values

In this section, we consider impossible values.

Lemma 4.1. *We have $S(C_2^4) \cap \mathbb{Z}_{\text{odd}} \subset \{16m + 1 \mid m \in \mathbb{Z}\}$.*

Proof. Let $D_4(a_0, \dots, a_{15}) = D_2(b_0, b_1, b_2, b_3)D_2(c_0, c_1, c_2, c_3)D_2(d_0, d_1, d_2, d_3)D_2(e_0, e_1, e_2, e_3)$ be an odd number. Then, $b_0 + b_2 \not\equiv b_1 + b_3 \pmod{2}$ holds from Lemma 2.4. We divide the proof into the following cases:

- (i) Exactly three of b_0, b_1, b_2, b_3 are even;
- (ii) Exactly one of b_0, b_1, b_2, b_3 is even.

First, we consider the case of (i). If $(b_0, b_1, b_2, b_3) \equiv (0, 0, 0, 1) \pmod{2}$, then there exist $k_i, l_i, m_i \in \mathbb{Z}$ satisfying $(b_0, b_1, b_2) = (2k_0, 2l_0, 2m_0)$, $(c_0, c_1, c_2) = (2k_1, 2l_1, 2m_1)$, $(d_0, d_1, d_2) = (2k_2, 2l_2, 2m_2)$, $(e_0, e_1, e_2) = (2k_3, 2l_3, 2m_3)$ and $\sum_{i=0}^3 k_i \equiv \sum_{i=0}^3 l_i \equiv \sum_{i=0}^3 m_i \equiv 0 \pmod{2}$ from Remark 2.2. Therefore, from Lemma 3.2 (1), we have

$$D_4(a_0, a_1, \dots, a_{15}) \equiv \prod_{i=0}^3 \{8(k_i + l_i + m_i) + 1\} \equiv 8 \sum_{i=0}^3 (k_i + l_i + m_i) + 1 \equiv 1 \pmod{16}.$$

Lemma 3.1 completes the proof for the case (i). Next, we consider the case of (ii). If $(b_0, b_1, b_2, b_3) \equiv (0, 1, 1, 1) \pmod{2}$, then there exist $k'_i \in \mathbb{Z}$ satisfying $b_0 = 2k'_0$, $c_0 = 2k'_1$, $d_0 = 2k'_2$, $e_0 = 2k'_3$ and $\sum_{i=0}^3 k'_i \equiv 0 \pmod{2}$ from Remark 2.2. Therefore, from Lemma 3.2 (2), we have

$$D_4(a_0, a_1, \dots, a_{15}) \equiv \prod_{i=0}^3 (8k'_i - 3) \equiv 8 \sum_{i=0}^3 k'_i + 1 \equiv 1 \pmod{16}.$$

Lemma 3.1 completes the proof for the case (ii). \square

Lemma 4.2. *We have $S(C_2^4) \cap 2\mathbb{Z} \subset 2^{16}\mathbb{Z}_{\text{odd}} \cup 2^{24}\mathbb{Z}_{\text{odd}} \cup 2^{26}\mathbb{Z}$.*

Proof. Let $D_4(a_0, \dots, a_{15}) = D_2(b_0, b_1, b_2, b_3)D_2(c_0, c_1, c_2, c_3)D_2(d_0, d_1, d_2, d_3)D_2(e_0, e_1, e_2, e_3)$ be an even number. Then, $b_0 + b_2 \equiv b_1 + b_3 \pmod{2}$ holds from Lemma 2.4. We prove the following:

- (i) If b_0, b_1, b_2, b_3 are even, then $D_4(a_0, \dots, a_{15}) \in 2^{16}\mathbb{Z}_{\text{odd}} \cup 2^{24}\mathbb{Z}_{\text{odd}} \cup 2^{26}\mathbb{Z}$;
- (ii) If b_0, b_1, b_2, b_3 are odd, then $D_4(a_0, \dots, a_{15}) \in 2^{16}\mathbb{Z}_{\text{odd}} \cup 2^{26}\mathbb{Z}$;
- (iii) If exactly two of b_0, b_1, b_2, b_3 are even, then $D_4(a_0, \dots, a_{15}) \in 2^{24}\mathbb{Z}_{\text{odd}} \cup 2^{26}\mathbb{Z}$.

First, we prove (i). If b_0, b_1, b_2, b_3 are even, then there exist $k_i, l_i, m_i, n_i \in \mathbb{Z}$ satisfying

$$\begin{aligned} (b_0, b_1, b_2, b_3) &= (2k_0, 2l_0, 2m_0, 2n_0), & (c_0, c_1, c_2, c_3) &= (2k_1, 2l_1, 2m_1, 2n_1), \\ (d_0, d_1, d_2, d_3) &= (2k_2, 2l_2, 2m_2, 2n_2), & (e_0, e_1, e_2, e_3) &= (2k_3, 2l_3, 2m_3, 2n_3) \end{aligned}$$

and $\sum_{i=0}^3 k_i \equiv \sum_{i=0}^3 l_i \equiv \sum_{i=0}^3 m_i \equiv \sum_{i=0}^3 n_i \equiv 0 \pmod{2}$ from Remark 2.2. Let N denote the cardinal number of the set $\{i \mid 0 \leq i \leq 3, k_i + m_i \equiv l_i + n_i \pmod{2}\}$. Then, $N \in \{0, 2, 4\}$ holds from $\sum_{i=0}^3 (k_i + m_i - l_i - n_i) \equiv 0 \pmod{2}$. Therefore, from Lemma 3.6 (1), we obtain (i). Also, in the same way, we can prove (ii) by using Lemma 3.6 (2). Finally, we prove (iii). If $(b_0, b_1, b_2, b_3) \equiv (0, 1, 0, 1) \pmod{2}$, then there exist $k'_i, m'_i \in \mathbb{Z}$ satisfying $(b_0, b_2) = (2k'_0, 2m'_0)$, $(c_0, c_2) = (2k'_1, 2m'_1)$, $(d_0, d_2) = (2k'_2, 2m'_2)$, $(e_0, e_2) = (2k'_3, 2m'_3)$ and $\sum_{i=0}^3 k'_i \equiv \sum_{i=0}^3 m'_i \equiv 0 \pmod{2}$ from Remark 2.2. Let N' denote the cardinal number of the set $\{i \mid 0 \leq i \leq 3, k'_i \equiv m'_i \pmod{2}\}$. Then, $N' \in \{0, 2, 4\}$ holds from $\sum_{i=0}^3 (k'_i - m'_i) \equiv 0 \pmod{2}$. Therefore, from Lemma 3.6 (3), we have $D_4(a_0, \dots, a_{15}) \in 2^{24}\mathbb{Z}_{\text{odd}} \cup 2^{26}\mathbb{Z}$. Lemma 3.1 completes the proof of (iii). \square

Lemma 4.3. *We have $S(C_2^4) \cap 2^{16}\mathbb{Z}_{\text{odd}} \subset \{2^{16}(4m+1) \mid m \in \mathbb{Z}\}$.*

Proof. Note that we have $S(C_2^3) \cap 2\mathbb{Z} = \{2^8(4m+1), 2^{12}m \mid m \in \mathbb{Z}\}$ from the description in the introduction. Let $D_4(a_0, \dots, a_{15}) = D_3(a_0 + a_8, \dots, a_7 + a_{15})D_3(a_0 - a_8, \dots, a_7 - a_{15}) \in 2^{16}\mathbb{Z}_{\text{odd}}$. Then, from Lemma 2.3 (1), there exist $k, l \in \mathbb{Z}$ satisfying $D_3(a_0 + a_8, \dots, a_7 + a_{15}) = 2^8(4k+1)$ and $D_3(a_0 - a_8, \dots, a_7 - a_{15}) = 2^8(4l+1)$. Therefore, we have $D_4(a_0, \dots, a_{15}) \in \{2^{16}(4m+1) \mid m \in \mathbb{Z}\}$. \square

Let $A := \{(8k-3)(8l+3) \mid k, l \in \mathbb{Z}\}$.

Lemma 4.4. *If $m \equiv -1 \pmod{8}$ and $m \notin A$, then $2^{24}m \notin S(C_2^4)$.*

To prove Lemma 4.4, we use the following two lemmas.

Lemma 4.5. *If $m \equiv -1 \pmod{8}$ and $m \notin A$, then $D_4(a_0, a_1, \dots, a_{15}) \neq 2^{24}m$ for any $a_0, a_1, \dots, a_{15} \in \mathbb{Z}$ with $b_0 \equiv b_1 \equiv b_2 \equiv b_3 \equiv 0 \pmod{2}$.*

Proof. We prove by contradiction. Assume that there exist $a_0, a_1, \dots, a_{15} \in \mathbb{Z}$ with $b_0 \equiv b_1 \equiv b_2 \equiv b_3 \equiv 0 \pmod{2}$ satisfying $D_4(a_0, a_1, \dots, a_{15}) = 2^{24}m$. Then there exist $k_i, l_i, m_i, n_i \in \mathbb{Z}$ satisfying

$$\begin{aligned} D_2^{(0)} &:= D_2(2k_0, 2l_0, 2m_0, 2n_0) = D_2(b_0, b_1, b_2, b_3), \\ D_2^{(1)} &:= D_2(2k_1, 2l_1, 2m_1, 2n_1) = D_2(c_0, c_1, c_2, c_3), \\ D_2^{(2)} &:= D_2(2k_2, 2l_2, 2m_2, 2n_2) = D_2(d_0, d_1, d_2, d_3), \\ D_2^{(3)} &:= D_2(2k_3, 2l_3, 2m_3, 2n_3) = D_2(e_0, e_1, e_2, e_3) \end{aligned}$$

and $\sum_{i=0}^3 k_i \equiv \sum_{i=0}^3 l_i \equiv \sum_{i=0}^3 m_i \equiv \sum_{i=0}^3 n_i \equiv 0 \pmod{2}$ from Remark 2.2. Note that $\sum_{i=0}^3 (k_i + m_i - l_i - n_i) \equiv 0 \pmod{2}$. Since $D_2^{(0)}D_2^{(1)}D_2^{(2)}D_2^{(3)} \in 2^{24}\mathbb{Z}_{\text{odd}}$, one of the following holds: (i) Three of $D_2^{(i)}$ are of type (1) or (2) in Lemma 3.3 and the other one is of type (5) in Lemma 3.3; (ii) Two of $D_2^{(i)}$ are of type (1) or (2) in Lemma 3.3 and the others are of type (3) or (4) in Lemma 3.3. In the case of (i), we have $\sum_{i=0}^3 (k_i + m_i - l_i - n_i) \not\equiv 0 \pmod{2}$. This is a contradiction. In the case of (ii), since $m \equiv -1 \pmod{8}$ and $m \notin A$, we have

$$D_2^{(\alpha)}, D_2^{(\beta)} \in \{2^4(8a+1) \mid a \in \mathbb{Z}\}, D_2^{(\gamma)} \in \{2^8(4a+1) \mid a \in \mathbb{Z}\}, D_2^{(\delta)} \in \{2^8(4a-1) \mid a \in \mathbb{Z}\},$$

where $\{\alpha, \beta, \gamma, \delta\} = \{1, 2, 3, 4\}$. This implies that

$$\begin{aligned} (k_\alpha, l_\alpha, m_\alpha, n_\alpha) &\equiv (0, 0, 0, 1), (0, 0, 1, 0), (0, 1, 0, 0) \text{ or } (1, 0, 0, 0) \pmod{2}, \\ (k_\beta, l_\beta, m_\beta, n_\beta) &\equiv (0, 0, 0, 1), (0, 0, 1, 0), (0, 1, 0, 0) \text{ or } (1, 0, 0, 0) \pmod{2}, \\ (k_\gamma, l_\gamma, m_\gamma, n_\gamma) &\equiv (0, 0, 0, 0) \pmod{2}, \\ (k_\delta, l_\delta, m_\delta, n_\delta) &\equiv (1, 1, 1, 1) \pmod{2}. \end{aligned}$$

Therefore, at least two of $\sum_{i=0}^3 k_i, \sum_{i=0}^3 l_i, \sum_{i=0}^3 m_i, \sum_{i=0}^3 n_i$ are odd. This is a contradiction. \square

Lemma 4.6. *If $m \equiv -1 \pmod{8}$ and $m \notin A$, then $D_4(a_0, a_1, \dots, a_{15}) \neq 2^{24}m$ for any $a_0, a_1, \dots, a_{15} \in \mathbb{Z}$, where exactly two of b_0, b_1, b_2, b_3 are even.*

Proof. We prove by contradiction. Assume that there exist $a_0, a_1, \dots, a_{15} \in \mathbb{Z}$, where exactly two of b_0, b_1, b_2, b_3 are even, satisfying $D_4(a_0, a_1, \dots, a_{15}) = 2^{24}m$. If $(b_0, b_1, b_2, b_3) \equiv (0, 1, 0, 1) \pmod{2}$, then there exist $k_i, l_i, m_i, n_i \in \mathbb{Z}$ satisfying

$$\begin{aligned} D_2^{(0)} &:= D_2(2k_0, 2l_0 + 1, 2m_0, 2n_0 + 1) = D_2(b_0, b_1, b_2, b_3), \\ D_2^{(1)} &:= D_2(2k_1, 2l_1 + 1, 2m_1, 2n_1 + 1) = D_2(c_0, c_1, c_2, c_3), \\ D_2^{(2)} &:= D_2(2k_2, 2l_2 + 1, 2m_2, 2n_2 + 1) = D_2(d_0, d_1, d_2, d_3), \\ D_2^{(3)} &:= D_2(2k_3, 2l_3 + 1, 2m_3, 2n_3 + 1) = D_2(e_0, e_1, e_2, e_3) \end{aligned}$$

and $\sum_{i=0}^3 k_i \equiv \sum_{i=0}^3 l_i \equiv \sum_{i=0}^3 m_i \equiv \sum_{i=0}^3 n_i \equiv 0 \pmod{2}$ from Remark 2.2. Since $D_2^{(0)} D_2^{(1)} D_2^{(2)} D_2^{(3)} \in 2^{24}\mathbb{Z}_{\text{odd}}$, it follows from Lemma 3.6 (3) that $k_i \equiv m_i \pmod{2}$ and $(2k_i + 2l_i + 1)(2m_i + 2n_i + 1) \equiv \pm 3 \pmod{8}$ for any $0 \leq i \leq 3$. Then, from Remark 3.5 (3) and (4), we find that every $D_2^{(i)}$ is of type (1) or (4) in Lemma 3.4 since $m \notin A$. In addition, since $m \equiv -1 \pmod{8}$, exactly one or three of $D_2^{(i)}$ are of type (1) in Lemma 3.4. Then we have $\sum_{i=0}^3 k_i \equiv 1 \pmod{2}$. This is a contradiction. Lemma 3.1 completes the proof. \square

Proof of Lemma 4.4. From Lemma 2.4, it is sufficient to prove that $D_4(a_0, \dots, a_{15}) \neq 2^{24}m$ for any $a_0, \dots, a_{15} \in \mathbb{Z}$ with $b_0 + b_2 \equiv b_1 + b_3 \pmod{2}$. If b_0, b_1, b_2, b_3 are odd, then as mentioned in the proof of Lemma 4.2, it holds that $D_4(a_0, \dots, a_{15}) \in 2^{16}\mathbb{Z}_{\text{odd}} \cup 2^{26}\mathbb{Z}$. Thus, $D_4(a_0, \dots, a_{15}) \neq 2^{24}m$. If all or exactly two of b_0, b_1, b_2, b_3 are even, then we have $D_4(a_0, \dots, a_{15}) \neq 2^{24}m$ from Lemmas 4.5 and 4.6. \square

5 Possible values

In this section, we determine all possible values. Lemmas 4.1–4.4 imply that $S(C_2^4)$ does not include every integer that is not mentioned in the following lemma.

Lemma 5.1. *For any $m, n \in \mathbb{Z}$, the following hold:*

- (1) $16m + 1 \in S(C_2^4)$;
- (2) $2^{16}(4m + 1) \in S(C_2^4)$;
- (3) $2^{24}(4m + 1) \in S(C_2^4)$;
- (4) $2^{24}(4m + 1)(8n + 3) \in S(C_2^4)$;
- (5) $2^{26}m \in S(C_2^4)$.

Proof. We obtain (1) from

$$\begin{aligned}
D_4(m+1, m, m, \dots, m) &= D_3(2m+1, 2m, 2m, \dots, 2m)D_3(1, 0, 0, \dots, 0) \\
&= D_2(4m+1, 4m, 4m, 4m)D_2(1, 0, 0, 0)^3 \\
&= D_1(8m+1, 8m)D_1(1, 0)^7 \\
&= (8m+1)^2 - (8m)^2 \\
&= 16m+1.
\end{aligned}$$

We obtain (2) from

$$\begin{aligned}
D_4(k+2, k, k, \dots, k) &= D_3(2k+2, 2k, 2k, \dots, 2k)D_3(2, 0, 0, \dots, 0) \\
&= D_2(4k+2, 4k, 4k, 4k)D_2(2, 0, 0, 0)^3 \\
&= D_1(8k+2, 8k)D_1(2, 0)^7 \\
&= \{(8k+2)^2 - (8k)^2\} \cdot 4^7 \\
&= 2^{16}(8k+1)
\end{aligned}$$

and

$$\begin{aligned}
&D_4(1-k, k, 1-k, k, -k, k, 1-k, k-1, 1-k, k, -k, k-1, -k, k, -k, k) \\
&= D_3(2-2k, 2k, 1-2k, 2k-1, -2k, 2k, 1-2k, 2k-1)D_3(0, 0, 1, 1, 0, 0, 1, -1) \\
&= D_2(2-4k, 4k, 2-4k, 4k-2)D_2(2, 0, 0, 0)D_2(0, 0, 2, 0)D_2(0, 0, 0, 2) \\
&= D_1(4-8k, 8k-2)D_1(0, 2)^2D_1(2, 0)^3D_1(-2, 0)D_1(0, -2) \\
&= \{(4-8k)^2 - (8k-2)^2\} (-4)^3 \cdot 4^4 \\
&= 2^{16}(8k-3).
\end{aligned}$$

We obtain (3) from

$$\begin{aligned}
D_4(m+3, m+1, m, m, \dots, m) &= D_3(2m+3, 2m+1, 2m, 2m, \dots, 2m)D_3(3, 1, 0, 0, \dots, 0) \\
&= D_2(4m+3, 4m+1, 4m, 4m)D_2(3, 1, 0, 0)^3 \\
&= D_1(8m+3, 8m+1)D_1(3, 1)^7 \\
&= \{(8m+3)^2 - (8m+1)^2\} \cdot 8^7 \\
&= 2^{24}(4m+1).
\end{aligned}$$

We obtain (4) by substituting

$$\begin{aligned}
a_0 &= m-n+2, & a_1 &= -(m+n+1), & a_2 &= m+n+1, & a_3 &= n-m, \\
a_4 &= m-n-1, & a_5 &= -(m+n), & a_6 &= m+n+1, & a_7 &= n-m, \\
a_8 &= m-n-1, & a_9 &= -(m+n), & a_{10} &= m+n+1, & a_{11} &= n-m, \\
a_{12} &= m-n-1, & a_{13} &= -(m+n), & a_{14} &= m+n+1, & a_{15} &= n-m.
\end{aligned}$$

In fact,

$$\begin{aligned}
D_4(a_0, \dots, a_{15}) &= D_2(b_0, b_1, b_2, b_3)D_2(c_0, c_1, c_2, c_3)D_2(d_0, d_1, d_2, d_3)D_2(e_0, e_1, e_2, e_3) \\
&= D_2(4m - 4n - 1, -4m - 4n - 1, 4m + 4n + 4, 4n - 4m)D_2(3, -1, 0, 0)^3 \\
&= D_1(8m + 3, -8m - 1)D_1(-8n - 5, -8n - 1)D_1(3, -1)^6 \\
&= \{(8m + 3)^2 - (8m + 1)^2\} \{(8n + 5)^2 - (8n + 1)^2\} \cdot 8^6 \\
&= 2^{24}(4m + 1)(8n + 3).
\end{aligned}$$

We obtain (5) from

$$\begin{aligned}
&D_4(k + 1, 1 - k, k + 1, -k, k - 1, -k, k + 1, -k, k, -k, k + 1, -k, k - 2, 1 - k, k + 1, -k) \\
&= D_3(2k + 1, 1 - 2k, 2k + 2, -2k, 2k - 3, 1 - 2k, 2k + 2, -2k)D_3(1, 1, 0, 0, 1, -1, 0, 0) \\
&= D_2(4k - 2, 2 - 4k, 4k + 4, -4k)D_2(4, 0, 0, 0)D_2(2, 0, 0, 0)D_2(0, 2, 0, 0) \\
&= D_1(8k + 2, 2 - 8k)D_1(-6, 2)D_1(4, 0)^2D_1(2, 0)^2D_1(0, 2)^2 \\
&= \{(8k + 2)^2 - (2 - 8k)^2\} \cdot 32 \cdot 16^2 \cdot 4^2 \cdot (-4)^2 \\
&= 2^{26}(2k)
\end{aligned}$$

and

$$\begin{aligned}
&D_4(k - 1, k + 1, k + 1, k + 2, k + 1, k + 2, k + 2, k, k, k, k, k, k, k, k, k) \\
&= D_3(2k - 1, 2k + 1, 2k + 1, 2k + 2, 2k + 1, 2k + 2, 2k + 2, 2k)D_3(-1, 1, 1, 2, 1, 2, 2, 0) \\
&= D_2(4k, 4k + 3, 4k + 3, 4k + 2)D_2(-2, -1, -1, 2)^2D_2(0, 3, 3, 2) \\
&= D_1(8k + 3, 8k + 5)D_1(-3, 1)^4D_1(-1, -3)^2D_1(3, 5) \\
&= \{(8k + 3)^2 - (8k + 5)^2\} \cdot 8^4 \cdot (-8)^2 \cdot (-16) \\
&= 2^{26}(2k + 1).
\end{aligned}$$

□

Remark that Lemma 5.1 (4) implies that $2^{24}(8m + 3), 2^{24}m' \in S(C_2^4)$ for any $m \in \mathbb{Z}$ and $m' \in A$. From Lemmas 4.1–4.4 and 5.1, Theorem 1.1 is proved.

References

- [1] Ton Boerkoel and Christopher Pinner. Minimal group determinants and the Lind-Lehmer problem for dihedral groups. *Acta Arith.*, 186(4):377–395, 2018.
- [2] Otto Hölder. Die Gruppen der Ordnungen p^3 , pq^2 , pqr , p^4 . *Math. Ann.*, 43(2-3):301–412, 1893.
- [3] Bishnu Paudel and Chris Pinner. Integer circulant determinants of order 15. *Integers*, 22:Paper No. A4, 21, 2022.
- [4] Christopher Pinner and Christopher Smyth. Integer group determinants for small groups. *Ramanujan J.*, 51(2):421–453, 2020.
- [5] Naoya Yamaguchi and Yuka Yamaguchi. Remark on laquer’s theorem for circulant determinants. *International Journal of Group Theory*, 2022.
- [6] Yuka Yamaguchi and Naoya Yamaguchi. Integer circulant determinants of order 16. *The Ramanujan Journal*, 2022.
- [7] J. W. A. Young. On the Determination of Groups Whose Order is a Power of a Prime. *Amer. J. Math.*, 15(2):124–178, 1893.

Yuka Yamaguchi
Faculty of Education
University of Miyazaki
1-1 Gakuen Kibanadai-nishi
Miyazaki 889-2192
JAPAN
y-yamaguchi@cc.miyazaki-u.ac.jp

Naoya Yamaguchi
Faculty of Education
University of Miyazaki
1-1 Gakuen Kibanadai-nishi
Miyazaki 889-2192
JAPAN
n-yamaguchi@cc.miyazaki-u.ac.jp