

DIRAC COHOMOLOGY FOR THE BGG CATEGORY \mathcal{O}

SPYRIDON AFENTOULIDIS-ALMPANIS

ABSTRACT. We study Dirac cohomology $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ for modules belonging to category \mathcal{O} of a finite dimensional complex semisimple Lie algebra. We prove Vogan's conjecture, a nonvanishing result for $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ while we show that in the case of a Hermitian symmetric pair $(\mathfrak{g}, \mathfrak{k})$ and an irreducible unitary module $M \in \mathcal{O}$, Dirac cohomology coincides with the nilpotent Lie algebra cohomology with coefficients in M . In the last part, we show that the higher Dirac cohomology and index introduced by Pandžić and Somberg satisfy nice homological properties for $M \in \mathcal{O}$.

CONTENTS

1. Introduction	1
2. Spin module	6
3. Dirac cohomology and BGG category \mathcal{O}	8
4. Application to simple Verma modules	13
5. Dirac and Lie algebra cohomology for \mathcal{O}	14
6. Higher Dirac cohomology and index for the BGG category \mathcal{O}	17
References	25

1. INTRODUCTION

Algebraic Dirac operators were introduced by Vogan in a series of lectures at MIT in the late 90's. More precisely, if \mathfrak{g} is a complex semisimple Lie algebra and \mathfrak{k} a maximal compact subalgebra of \mathfrak{g} , to each \mathfrak{g} -module (X, π) , he associated an operator $D_{\mathfrak{g}, \mathfrak{k}}(X) : X \otimes S \rightarrow X \otimes S$ given by

$$(1) \quad D_{\mathfrak{g}, \mathfrak{k}}(X) = \sqrt{2} \sum_j \pi(Z_j) \otimes \gamma(Z_j)$$

where $\{Z_j\}$ is an orthonormal basis of $\mathfrak{p} := \mathfrak{k}^\perp$, the orthogonal complement of \mathfrak{k} with respect to the Killing form $\langle \cdot, \cdot \rangle$ of \mathfrak{g} , S is a spin module for the Clifford algebra $C(\mathfrak{p})$ of \mathfrak{p} and γ stands for the Clifford multiplication of

2020 Mathematics Subject Classification. Primary 17B10; Secondary 17B20, 17B56.

Key words and phrases. Dirac cohomology, BGG category \mathcal{O} , Verma modules.

The author was supported by the GAČR grant EXPRO 19-28628X.

$\mathbf{C}(\mathfrak{p})$ on S . The operator $D_{\mathfrak{g},\mathfrak{k}}(X)$ is well defined in the sense that it does not depend on the choice of the basis $\{Z_j\}$. The tensor product $X \otimes S$ has a \mathfrak{k} -action via the diagonal embedding

$$(\cdot)_{\Delta} : \mathfrak{k} \longrightarrow U(\mathfrak{g}) \otimes \mathbf{C}(\mathfrak{p})$$

of \mathfrak{k} into $U(\mathfrak{g}) \otimes \mathbf{C}(\mathfrak{p})$, given by the embedding of \mathfrak{k} in $U(\mathfrak{g})$ and the embedding (9) of \mathfrak{k} in $\mathbf{C}(\mathfrak{p})$, while, with respect to this action, $D_{\mathfrak{g},\mathfrak{k}}(X)$ is \mathfrak{k} -equivariant, so that $\ker D_{\mathfrak{g},\mathfrak{k}}(X)$ and $\text{im } D_{\mathfrak{g},\mathfrak{k}}(X)$ are \mathfrak{k} -modules.

Vogan defined the Dirac cohomology of a \mathfrak{g} -module to be the \mathfrak{k} -module

$$(2) \quad H_D^{\mathfrak{g},\mathfrak{k}}(X) = \frac{\ker D_{\mathfrak{g},\mathfrak{k}}(X)}{\text{im } D_{\mathfrak{g},\mathfrak{k}}(X) \cap \ker D_{\mathfrak{g},\mathfrak{k}}(X)}.$$

Assuming that X is an irreducible (\mathfrak{g}, K) -module, so that $H_D^{\mathfrak{g},\mathfrak{k}}(X)$ is a module for the spin 2-cover \tilde{K} of K , Vogan conjectured that if $H_D^{\mathfrak{g},\mathfrak{k}}(X)$ contains a \tilde{K} -module with highest weight β , then V has infinitesimal character $\beta + \rho_{\mathfrak{k}}$. In other words, the infinitesimal character of a module can be recovered from its Dirac cohomology. Vogan's conjecture was first proved by Huang and Pandžić in [HP02]. Crucial for the proof of Vogan's conjecture (as well for many other results and applications) is the fact that the square $D_{\mathfrak{g},\mathfrak{k}}(X)^2$ differs from the Casimir by a scalar. Namely,

$$(3) \quad D_{\mathfrak{g},\mathfrak{k}}(X)^2 = \Omega_{\mathfrak{g}} \otimes 1 - (\Omega_{\mathfrak{k}})_{\Delta} + \|\rho\|^2 - \|\rho_{\mathfrak{k}}\|^2$$

where $\Omega_{\mathfrak{g}}$ (respectively $\Omega_{\mathfrak{k}}$) denotes the Casimir element in the enveloping algebra $U(\mathfrak{g})$ (respectively $U(\mathfrak{k})$) of \mathfrak{g} (respectively \mathfrak{k}), ρ (respectively $\rho_{\mathfrak{k}}$) is half the sum of the positive roots for some fixed positive system Δ^+ (respectively $\Delta_{\mathfrak{k}}^+$) of the root system Δ (respectively $\Delta_{\mathfrak{k}}$) of \mathfrak{g} (respectively \mathfrak{k}) and $\|\cdot\|$ is the norm induced by the Killing form of \mathfrak{g} .

In the case of equal rank \mathfrak{g} and \mathfrak{k} , another interesting invariant of X can be defined using $D_{\mathfrak{g},\mathfrak{k}}(X)$. More precisely, in this case the spin module S splits into $S = S^+ \oplus S^-$ and $D_{\mathfrak{g},\mathfrak{k}}(X)$ interchanges $X \otimes S^{\pm}$ so that there are operators

$$D_{\mathfrak{g},\mathfrak{k}}(X)^{\pm} : X \otimes S^{\pm} \rightarrow X \otimes S^{\mp}.$$

Then, we can define "half"-Dirac cohomology spaces

$$H_D^{\mathfrak{g},\mathfrak{k}}(X)^{\pm} := \frac{\ker D_{\mathfrak{g},\mathfrak{k}}(X)^{\pm}}{\text{im } D_{\mathfrak{g},\mathfrak{k}}(X)^{\mp} \cap \ker D_{\mathfrak{g},\mathfrak{k}}(X)^{\pm}}$$

and the virtual \mathfrak{k} -module

$$I_D(X) := H_D^{\mathfrak{g},\mathfrak{k}}(X)^+ - H_D^{\mathfrak{g},\mathfrak{k}}(X)^-$$

known as the Dirac index of X . In [MPV17], Mehdi, Pandžić and Vogan proved a translation principle for $I_D(X)$ and provided interesting applications.

In [Goe99] and [Kos99], Goette and independently Kostant generalized the above construction of Dirac operators to more general pairs $(\mathfrak{g}, \mathfrak{h})$ where \mathfrak{h} is a subalgebra of \mathfrak{g} such that

- (a) \mathfrak{h} is reductive
- (b) the restriction of the Killing form of \mathfrak{g} to \mathfrak{h} is nondegenerate.

Then, \mathfrak{g} decomposes into an orthogonal direct sum

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q},$$

where $\mathfrak{q} := \mathfrak{h}^\perp$ with respect to the Killing form of \mathfrak{g} , and we consider again a spin module S for the Clifford algebra $\mathbf{C}(\mathfrak{q})$ of \mathfrak{q} . For a \mathfrak{g} -module X , they defined the corresponding Dirac operator $D_{\mathfrak{g},\mathfrak{h}}(X) : X \otimes S \rightarrow X \otimes S$, known as cubic Dirac operator, to be given by

$$(4) \quad D_{\mathfrak{g},\mathfrak{h}}(X) = \sqrt{2} \left\{ \sum_j \pi(Z_j) \otimes \gamma(Z_j) - 1 \otimes \gamma(c) \right\},$$

where $\{Z_j\}$ is an orthonormal basis of \mathfrak{q} and $\gamma(c)$ is given by

$$(5) \quad \gamma(c) = \frac{1}{6} \sum_{i,j,k} B(\tilde{Z}_i, [\tilde{Z}_j, \tilde{Z}_k]) \gamma(Z_i) \gamma(Z_j) \gamma(Z_k).$$

We note that in the case where \mathfrak{h} is a symmetric Lie algebra, $\gamma(c) = 0$. The operator $D_{\mathfrak{g},\mathfrak{h}}(X)$ turns out to have the same "good" properties with $D_{\mathfrak{g},\mathfrak{k}}(X)$. Namely, $D_{\mathfrak{g},\mathfrak{h}}(X)$ does not depend on the choice of the basis $\{Z_j\}$ while it is \mathfrak{h} -equivariant with respect to the action of \mathfrak{h} on $X \otimes S$ given by a diagonal embedding of \mathfrak{h} in $U(\mathfrak{g}) \otimes \mathbf{C}(\mathfrak{q})$. The modified formula, with the extra cubic term $1 \otimes \gamma(c)$, is used instead of formula (1) so that we have the convenient formula for $D_{\mathfrak{g},\mathfrak{h}}(X)^2$:

$$(6) \quad D_{\mathfrak{g},\mathfrak{h}}(X)^2 = \Omega_{\mathfrak{g}} \otimes 1 - (\Omega_{\mathfrak{h}})_{\Delta} + \|\rho\|^2 - \|\rho_{\mathfrak{h}}\|^2,$$

where $\rho_{\mathfrak{h}}$ is half the sum of the positive roots of the positive system $\Delta_{\mathfrak{h}}^+ := \Delta_{\mathfrak{h}} \cap \Delta^+$ for $\Delta_{\mathfrak{h}} := \Delta(\mathfrak{h}, \mathfrak{t})$. For a more detailed discussion about the case where the cubic term $\gamma(c)$ is omitted, see [Afe21a] and [Afe21b]. Moreover, as we did in the case of the symmetric pair $(\mathfrak{g}, \mathfrak{k})$, we can still define the Dirac cohomology $H_D^{\mathfrak{g},\mathfrak{h}}(X)$ of a \mathfrak{g} -module X to be the \mathfrak{h} -module

$$H_D^{\mathfrak{g},\mathfrak{h}}(X) = \frac{\ker D_{\mathfrak{g},\mathfrak{h}}(X)}{\text{im } D_{\mathfrak{g},\mathfrak{h}}(X) \cap \ker D_{\mathfrak{g},\mathfrak{h}}(X)}.$$

In the case where X is a finite-dimensional \mathfrak{g} -module, $H_D^{\mathfrak{g},\mathfrak{h}}(X)$ coincides with $\ker D_{\mathfrak{g},\mathfrak{h}}(X)$. If, moreover, \mathfrak{h} contains a Cartan subalgebra \mathfrak{t} of \mathfrak{g} , Kostant gave a complete decomposition of $\ker D_{\mathfrak{g},\mathfrak{h}}(X)$ into simple \mathfrak{h} -modules:

$$(7) \quad \ker(D_{\mathfrak{g},\mathfrak{h}}(X)) = \bigoplus_{w \in W^1} F_{w(\lambda+\rho)-\rho_{\mathfrak{h}}},$$

where $\lambda \in \mathfrak{t}^*$ is the highest weight of X , F_μ stands for the simple highest weight \mathfrak{h} -module with extremal weight $\mu \in \mathfrak{t}^*$, and

$$W^1 := \{w \in W \mid \Delta_{\mathfrak{h}}^+ \subseteq w\Delta^+\},$$

with W being the Weyl group of \mathfrak{g} . In fact, a similar formula was proved in the unequal rank situation, in [HKP09] for the pair $(\mathfrak{g}, \mathfrak{k})$ and in [MZ14] for pairs $(\mathfrak{g}, \mathfrak{h})$. In particular, in the unequal rank case, the decomposition of $H_D^{\mathfrak{g}, \mathfrak{h}}(X)$ into irreducibles is no longer multiplicity free.

Apart from finite dimensional modules, Dirac cohomology has been studied for various families of modules, including highest weight modules in [HX12], $A_{\mathfrak{q}}(\lambda)$ modules in [HKP09], generalized Enright-Varadarajan modules in [MP11], unipotent representations in [BP11] and [BP15]. In all the above cases, some finiteness assumption has been made on the restriction of the corresponding \mathfrak{g} -module to \mathfrak{k} (or more generally to \mathfrak{h}). For instance, in the case of a (\mathfrak{g}, K) -module X , we automatically assume that $X|_{\mathfrak{k}}$, the restriction of X to \mathfrak{k} , admits an algebraic direct sum decomposition into simple finite dimensional \mathfrak{k} -modules.

The aim of this paper is to study Dirac cohomology for modules M belonging to Bernstein-Gelfand-Gelfand category \mathcal{O} . Category \mathcal{O} , introduced by J. Bernstein, I. Gelfand and S. Gelfand in [BGG76], consists of the finitely generated, locally n -finite weight modules of \mathfrak{g} . Examples of modules belonging to \mathcal{O} include finite dimensional modules, Verma modules, (twisted) dual Verma modules, etc. This category has nice homological properties and it seems to be the "correct" module category to study questions raised by Verma [Ver68] concerning composition series and embeddings of Verma modules, and Jantzen [Jan79] concerning his so-called translation functors.

In [HX12], Huang and Xiao computed Dirac cohomology in terms of Kazhdan-Lusztig polynomials in the case of modules belonging to the parabolic category $\mathcal{O}^{\mathfrak{p}}$; here, \mathfrak{p} is a standard parabolic subalgebra of \mathfrak{g} with Levi decomposition $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{u}$, and $\mathfrak{h} := \mathfrak{l}$. The parabolic category $\mathcal{O}^{\mathfrak{p}}$ is the full subcategory of \mathcal{O} consisting of the \mathfrak{g} -modules $M \in \mathcal{O}$ such that the restriction $M|_{\mathfrak{l}}$ of M to the Levi factor \mathfrak{l} decomposes into an algebraic direct sum of finite dimensional \mathfrak{l} -submodules. Hence, there is again a finiteness condition satisfied for $M|_{\mathfrak{l}}$.

The main difference in our case is that M belongs to the larger category \mathcal{O} , so that $M|_{\mathfrak{h}}$ need not satisfy any finiteness condition and \mathfrak{h} need not be a Levi factor of some parabolic. For instance, the case of a Verma module $M(\lambda)$ could be illustrative. Whenever \mathfrak{h} strictly contains a Cartan subalgebra \mathfrak{t} of \mathfrak{g} , any \mathfrak{h} -submodule of $M(\lambda)|_{\mathfrak{h}}$ turns out to be infinite dimensional. Indeed, for such an \mathfrak{h} , there will be some $A \subset \Delta^+$ such that \mathfrak{h} decomposes

into

$$\mathfrak{h} = \mathfrak{t} \oplus \mathfrak{n}_{\mathfrak{h}}^+ \oplus \mathfrak{n}_{\mathfrak{h}}^-,$$

where

$$\mathfrak{n}_{\mathfrak{h}}^{\pm} = \bigoplus_{\alpha \in A} \mathfrak{g}_{\pm\alpha},$$

and

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{n}_{\mathfrak{q}}^+ \oplus \mathfrak{n}_{\mathfrak{q}}^-$$

with

$$\mathfrak{n}_{\mathfrak{q}}^{\pm} := \bigoplus_{\alpha \in \Delta \setminus A} \mathfrak{g}_{\pm\alpha}.$$

Then, as $\mathfrak{n}_{\mathfrak{h}}^-$ -modules,

$$M(\lambda)|_{\mathfrak{n}_{\mathfrak{h}}^-} = U(\mathfrak{g}) \otimes_{U(\mathfrak{t} \oplus \mathfrak{n}_{\mathfrak{h}}^+ \oplus \mathfrak{n}_{\mathfrak{q}}^+)} \mathbb{C}\lambda \cong U(\mathfrak{n}_{\mathfrak{h}}^- \oplus \mathfrak{n}_{\mathfrak{q}}^-) \cong U(\mathfrak{n}_{\mathfrak{h}}^-) \otimes_{\mathbb{C}} U(\mathfrak{n}_{\mathfrak{q}}^-).$$

In other words, $M(\lambda)|_{\mathfrak{n}_{\mathfrak{h}}^-}$ is a free $U(\mathfrak{n}_{\mathfrak{h}}^-)$ -module so that every $U(\mathfrak{n}_{\mathfrak{h}}^-)$ -submodule is infinite dimensional. This should be the case for every $U(\mathfrak{h})$ -submodule of $M(\lambda)|_{\mathfrak{h}}$. Another problem that may arise is the fact that the restriction $M|_{\mathfrak{h}}$ of $M \in \mathcal{O}$ is not finitely generated in general. For example, one can consider the case where M is a Verma module $M(\lambda)$ and \mathfrak{h} coincides with the Cartan subalgebra \mathfrak{t} of \mathfrak{g} . For a thorough discussion about restrictions of Verma modules, e.g., [Kob12, KØSS15, PS16].

In this paper, we adapt the methods used in the study of Dirac cohomology of (\mathfrak{g}, K) -modules in order to study Dirac cohomology $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ in the case where M belongs to \mathcal{O} . More precisely, after collecting the basic notions about Clifford algebras and spinors in Section 2, in Section 3 we show that, even if the decomposition of $M \otimes S$ into \mathfrak{h} -submodules can be tricky, $M \otimes S$ has a generalized eigenspace decomposition for the operator $D_{\mathfrak{g}, \mathfrak{h}}(M)^2$ (Corollary 3.3). As a consequence, Dirac cohomology $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ is finitely generated and Vogan's conjecture holds (Theorem 3.4). Moreover, we show that Dirac cohomology does not vanish whenever $M \neq 0$ (Theorem 3.5). As a direct application, in Section 4, we show that the Dirac cohomology of a simple Verma module is again a simple Verma module for \mathfrak{h} (Theorem 4.1).

Formula (7) resembles Kostant's formula for nilpotent Lie algebra cohomology with coefficients in a finite dimensional simple module [Kos61]. Indeed, in [HPR06], Huang, Pandžić and Renard related nilpotent Lie algebra cohomology and Dirac cohomology in the case of Hermitian symmetric pairs $(\mathfrak{g}, \mathfrak{k})$ and irreducible unitary (\mathfrak{g}, K) -modules. In Section 5, we show that a similar result holds when X is replaced by an irreducible unitary module in \mathcal{O} (Theorem 5.2).

In [PS16], for equal rank \mathfrak{g} and \mathfrak{h} , Pandžić and Somberg studied Dirac cohomology for (\mathfrak{g}, K) -modules having generalized infinitesimal character and satisfying some finiteness condition for \mathfrak{h} . In contrast to the case of a \mathfrak{g} -module X having infinitesimal character where $D_{\mathfrak{g},\mathfrak{h}}(X)^2$, by (6), acts on each \mathfrak{h} -isotypic component of $X \otimes S$ by a scalar, when X has a generalized infinitesimal character, this is not true in general. As a consequence, $\ker D_{\mathfrak{g},\mathfrak{h}}(X)^2$ need not exhaust the generalized kernel of $D_{\mathfrak{g},\mathfrak{h}}(X)$ and $H_D^{\mathfrak{g},\mathfrak{h}}(X)$ may have strange behaviour in terms of Homological Algebra. In order to overcome this problem, they introduced a notion of higher Dirac cohomology and proved that this new \mathfrak{h} -module has nice homological properties. In the last section, we adapt their methods in order to prove similar results for $H_D^{\mathfrak{g},\mathfrak{h}}(M)$ when M is a module in \mathcal{O} .

2. SPIN MODULE

In this section, we recall the basic notions of Clifford algebras and spinors. More details can be found in [GW09, Chapter 6] and [HP06, Chapter 2]. Let \mathfrak{g} be a complex semisimple Lie algebra, $\langle \cdot, \cdot \rangle$ its Killing form, θ a Cartan involution of \mathfrak{g} and \mathfrak{h} a Lie subalgebra of \mathfrak{g} such that:

$$(8) \quad \begin{cases} \mathfrak{h} \text{ is reductive} \\ \mathfrak{h} \text{ is } \theta\text{-stable} \\ \langle \cdot, \cdot \rangle|_{\mathfrak{h} \times \mathfrak{h}} \text{ is nondegenerate.} \end{cases}$$

Then, \mathfrak{g} decomposes into a direct sum

$$\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{q},$$

where $\mathfrak{q} := \mathfrak{h}^\perp$ with respect to $\langle \cdot, \cdot \rangle$. Consider the Clifford algebra $\mathbf{C}(\mathfrak{q})$ of \mathfrak{q} , i.e. the quotient of the tensor algebra $T(\mathfrak{q})$ of \mathfrak{q} with the ideal generated by all elements

$$X \otimes Y + Y \otimes X - \langle X, Y \rangle, \quad X, Y \in \mathfrak{q}.$$

For a and b in \mathfrak{q} , let $R_{a,b}$ be the element of

$$\mathfrak{so}(\mathfrak{q}) := \{T \in \mathfrak{gl}(\mathfrak{q}) \mid \langle Tu, v \rangle + \langle u, Tv \rangle = 0, \forall u, v \in \mathfrak{q}\}.$$

defined by

$$R_{a,b}(v) := \langle b, v \rangle a - \langle a, v \rangle b, \quad v \in \mathfrak{q}.$$

Then $\{R_{a,b} \mid a, b \in \mathfrak{q}\}$ spans $\mathfrak{so}(\mathfrak{q})$ and there is a Lie algebra monomorphism $\varphi : \mathfrak{so}(\mathfrak{q}) \rightarrow \mathbf{C}(\mathfrak{q})$ given by

$$\varphi : R_{a,b} \mapsto \frac{1}{2}[\gamma(a), \gamma(b)].$$

Let \mathfrak{q}^- be a maximal isotropic subspace of \mathfrak{q} and set $S := \bigwedge \mathfrak{q}^-$. Then, $\mathbf{C}(\mathfrak{q})$ acts on S by Clifford multiplication while S equipped with the \mathfrak{h} -action defined by

$$(9) \quad \mathfrak{h} \xrightarrow{\text{ad}} \mathfrak{so}(\mathfrak{q}) \xrightarrow{\varphi} \mathbf{C}(\mathfrak{q}) \longrightarrow \text{End}(S)$$

is a Lie algebra module called spin module of \mathfrak{h} . We note that the spin module S does not depend on the choice of \mathfrak{q}^- .

Suppose that there is a common Cartan subalgebra \mathfrak{t} of \mathfrak{g} and \mathfrak{h} . Then, the root system $\Delta := \Delta(\mathfrak{g}, \mathfrak{t})$ splits into $\Delta = \Delta_{\mathfrak{h}} \sqcup \Delta_{\mathfrak{q}}$ so that

$$\mathfrak{h} = \mathfrak{t} \oplus \bigoplus_{\alpha \in \Delta_{\mathfrak{h}}} \mathfrak{g}_{\alpha} \text{ and } \mathfrak{q} = \bigoplus_{\beta \in \Delta_{\mathfrak{q}}} \mathfrak{g}_{\beta}.$$

Choose a positive system $\Delta^+ \subset \Delta$ for Δ and set $\Delta_{\mathfrak{h}}^+ := \Delta^+ \cap \Delta_{\mathfrak{h}}$ to be a positive system for the root system $\Delta_{\mathfrak{h}}$ and $\Delta_{\mathfrak{q}}^+ := \Delta^+ \cap \Delta_{\mathfrak{q}}$ while ρ and $\rho_{\mathfrak{h}}$ stand for the half-sum of positive roots in Δ and $\Delta_{\mathfrak{h}}$ respectively. Then,

$$(10) \quad \mathfrak{n}_{\mathfrak{q}}^{\pm} := \bigoplus_{\beta \in \Delta_{\mathfrak{q}}^{\pm}} \mathfrak{g}_{\pm\beta}.$$

are maximal dual isotropic subspace of \mathfrak{q} and S can be chosen to be

$$S := \bigwedge \mathfrak{n}_{\mathfrak{q}}^-.$$

Then, as \mathfrak{h} -module, S is a weight module, i.e. it can be decomposed into a direct sum of its weight subspaces. More precisely, let β_1, \dots, β_l be the positive roots lying in \mathfrak{q} and $e_{\pm 1}, \dots, e_{\pm k}$ the corresponding root vectors of $\pm\beta_1, \dots, \pm\beta_l$ respectively, such that

$$\langle e_i, e_j \rangle = \begin{cases} 1 & \text{if } i + j = 0 \\ 0 & \text{otherwise.} \end{cases}$$

If $I = (i_1, \dots, i_s)$ with $1 \leq s \leq k$, set

$$u_I := e_{-i_1} \wedge \dots \wedge e_{-i_s} \in S.$$

Then u_I is a weight vector of S , the weight being

$$(11) \quad \rho - \rho_{\mathfrak{h}} - \sum_{i \in I} \beta_i = \frac{1}{2} \left\{ \sum_{i \notin I} \beta_i - \sum_{i \in I} \beta_i \right\},$$

where, by abuse of notation, we write $i \in I$ if $i \in \{i_1, \dots, i_s\}$. One easily checks that as \mathfrak{t} -modules

$$S \cong \bigwedge \mathfrak{n}_{\mathfrak{q}}^- \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{h}}}.$$

Moreover, from the \mathbb{Z}_2 -splitting of the vector space $\bigwedge \mathfrak{n}_{\mathfrak{q}}^-$, the vector space S admits a \mathbb{Z}_2 -splitting

$$(12) \quad S = S^+ \oplus S^-.$$

Under the assumption of equal rank \mathfrak{g} and \mathfrak{h} , this splitting is preserved by the action of \mathfrak{h} .

3. DIRAC COHOMOLOGY AND BGG CATEGORY \mathcal{O}

Let \mathfrak{g} be a complex semisimple Lie algebra and \mathfrak{h} a Lie subalgebra of \mathfrak{g} satisfying (8). Moreover, assume that \mathfrak{h} contains a Cartan subalgebra \mathfrak{t} of \mathfrak{g} . In this section, we show that for a module $M \in \mathcal{O}$, $M \otimes S$ admits a direct sum decomposition in terms of generalized infinitesimal character subspaces. As a corollary, $M \otimes S$ has a generalized eigenspace decomposition for the operator $D_{\mathfrak{g},\mathfrak{h}}(M)^2$ while Vogan's conjecture holds for $H_D^{\mathfrak{g},\mathfrak{h}}(M)$. At the end of the section, we show that whenever $M \neq 0$, $H_D^{\mathfrak{g},\mathfrak{h}}(M)$ is nonzero.

Let M be a module belonging to \mathcal{O} . Then, M can be decomposed into a direct sum

$$(13) \quad M = \bigoplus_{\lambda} M^{\chi_{\lambda}},$$

where $\chi_{\lambda} : Z(\mathfrak{g}) \rightarrow \mathbb{C}$ is the infinitesimal character determined by $\lambda \in \mathfrak{t}^*$ using the Harish-Chandra map, and

$$M^{\chi_{\lambda}} := \{v \in M \mid \forall z \in Z(\mathfrak{g}), \exists n := n(z, v) > 0 \text{ s.t. } (z - \chi_{\lambda}(z))^n v = 0\}$$

is the generalized infinitesimal character subspace of λ . In other words, category \mathcal{O} decomposes into a direct sum

$$\mathcal{O} = \bigoplus_{\lambda} \mathcal{O}_{\chi_{\lambda}}$$

with $\mathcal{O}_{\chi_{\lambda}}$ being the full subcategory of \mathcal{O} consisting of modules $M \in \mathcal{O}$ such that $M^{\chi_{\lambda}} = M$ [Hum08]. For the study of $D_{\mathfrak{g},\mathfrak{h}}(M)$, it would be convenient to have a similar decomposition into generalized infinitesimal character subspaces for the \mathfrak{h} -module $M \otimes S$.

Lemma 3.1. *Let $M \in \mathcal{O}$. Then $M \otimes S$ can be decomposed into a direct sum*

$$M \otimes S = \bigoplus_{\nu \in \mathfrak{t}^*} (M \otimes S)^{\xi_{\nu}}$$

where $(M \otimes S)^{\xi_{\nu}}$ stands for the generalized infinitesimal character subspace corresponding to $\xi_{\nu} \in \max\text{spec } Z(\mathfrak{h})$. Moreover, each $(M \otimes S)^{\xi_{\nu}}$ is finitely generated.

Proof of Lemma 3.1. Since the actions of $Z(\mathfrak{h})$ and \mathfrak{t} on $M \otimes S$ commute, it suffices to prove the statement for any weight subspace $(M \otimes S)_{\mu}$, $\mu \in \mathfrak{t}^*$, of $M \otimes S$. Since M belongs to the category \mathcal{O} , $(M \otimes S)_{\mu}$ is finite-dimensional and the result follows from a standard Linear Algebra argument.

For the second statement, if $(M \otimes S)^{\xi_\nu} = 0$, there is nothing to prove. If not, let v_1 be a maximal weight vector of $(M \otimes S)^{\xi_\nu}$ and set

$$X_1 := U(\mathfrak{h})v_1 \subseteq (M \otimes S)^{\xi_\nu}.$$

If $(M \otimes S)^{\xi_\nu} = X_1$, there is nothing more to prove. If not, let \bar{v}_2 be a maximal weight vector of $(M \otimes S)^{\xi_\nu}/X_1$ and set $\bar{X}_2 := U(\mathfrak{h})\bar{v}_2$. Let X_2 be the preimage of \bar{X}_2 in $(M \otimes S)^{\xi_\nu}$. Inductively, if $(M \otimes S)^{\xi_\nu}/X_{i-1}$ is nontrivial, we define \bar{v}_i to be a maximal weight of $(M \otimes S)^{\xi_\nu}/X_{i-1}$, \bar{X}_i the cyclic submodule generated by \bar{v}_i and X_i the preimage of \bar{X}_i in $(M \otimes S)^{\xi_\nu}$. Therefore, we have created a filtration

$$(14) \quad 0 = X_0 \subsetneq X_1 \subsetneq \dots \subseteq (M \otimes S)^{\xi_\nu}.$$

It suffices to show that (14) is finite. Suppose that $(M \otimes S)^{\xi_\nu}/X_{i-1}$ is not trivial and let $\bar{v}_i = v_i + X_{i-1}$ be a nonzero maximal weight vector of it, say of weight ν_i . Then, $Z(\mathfrak{h})$ acts on it by the character $\xi_{\nu_i + \rho_{\mathfrak{h}}}$ and so $\xi_{\nu_i + \rho_{\mathfrak{h}}}$ must coincide with ξ_ν so that ν and $\nu_i + \rho_{\mathfrak{h}}$ belong to the same W -orbit. Since there are only finitely many weights ν_i of $M \otimes S$ satisfying this condition and for such a weight the corresponding weight subspace $(M \otimes S)_{\nu_i}$ is finite dimensional, we deduce that filtration (14) is of finite length and so $(M \otimes S)^{\xi_\nu}$ must be finitely generated. \square

Thanks to (6), we have the following lemma.

Lemma 3.2. *Let $M \in \mathcal{O}_{\chi_\lambda}$. Then $(M \otimes S)^{\xi_\nu}$ is a subspace of the generalized eigenspace of $D_{\mathfrak{g}, \mathfrak{h}}(M)^2$ with eigenvalue*

$$c_\nu := \chi_\lambda(\Omega_{\mathfrak{g}}) + \|\rho\|^2 - \xi_\nu(\Omega_{\mathfrak{h}}) - \|\rho_{\mathfrak{h}}\|^2 = \|\lambda + \rho\|^2 - \|\mu + \rho_{\mathfrak{h}}\|^2.$$

Proof of Lemma 3.2. Let $x \in (M \otimes S)^{\xi_\nu}$. Then, there are $n_1, n_2 \geq 1$ such that

$$(\Omega_{\mathfrak{g}} \otimes 1 - \chi_\lambda(\Omega_{\mathfrak{g}}))^{n_1} x = 0$$

and

$$(\Omega_{\mathfrak{h}_\Delta} - \xi_\nu(\Omega_{\mathfrak{h}}))^{n_2} x = 0.$$

Set

$$c_\nu := \chi_\lambda(\Omega_{\mathfrak{g}}) + \|\rho\|^2 - \xi_\nu(\Omega_{\mathfrak{h}}) - \|\rho_{\mathfrak{h}}\|^2.$$

We will show that

$$(D_{\mathfrak{g}, \mathfrak{h}}(M)^2 - c_\nu)^{n_1 + n_2 - 1} x = 0.$$

Recall that according to (6),

$$D_{\mathfrak{g}, \mathfrak{h}}(X)^2 = \Omega_{\mathfrak{g}} \otimes 1 - (\Omega_{\mathfrak{h}})_\Delta + \|\rho\|^2 - \|\rho_{\mathfrak{h}}\|^2$$

while $\Omega_{\mathfrak{g}} \otimes 1$ and $\Omega_{\mathfrak{h}_\Delta}$ commute. Therefore

$$\begin{aligned} (D_{\mathfrak{g},\mathfrak{h}}(M)^2 - c_\nu)^{n_1+n_2-1} &= \left((\Omega_{\mathfrak{g}} \otimes 1 - \chi_\lambda(\Omega_{\mathfrak{g}})) - (\Omega_{\mathfrak{h}_\Delta} - \xi_\nu(\Omega_{\mathfrak{h}})) \right)^{n_1+n_2-1} \\ &= \sum_{k=0}^{n_1+n_2-1} (-1)^k \binom{n_1+n_2-1}{k} (\Omega_{\mathfrak{g}} \otimes 1 - \chi_\lambda(\Omega_{\mathfrak{g}}))^k \\ &\quad (\Omega_{\mathfrak{h}_\Delta} - \xi_\nu(\Omega_{\mathfrak{h}}))^{n_1+n_2-1-k} \end{aligned}$$

and one can easily check that the exponent of $(\Omega_{\mathfrak{g}} \otimes 1 - \chi_\lambda(\Omega_{\mathfrak{g}}))$ is greater than n_1 or the exponent of $(\Omega_{\mathfrak{h}_\Delta} - \xi_\nu(\Omega_{\mathfrak{h}}))$ is greater than n_2 . Hence, $(\Omega_{\mathfrak{g}} \otimes 1 - \chi_\lambda(\Omega_{\mathfrak{g}}))^k x = 0$ or $(\Omega_{\mathfrak{h}_\Delta} - \xi_\nu(\Omega_{\mathfrak{h}}))^{n_1+n_2-1-k} x = 0$, and

$$(D_{\mathfrak{g},\mathfrak{h}}(M)^2 - c_\nu)^{n_1+n_2-1} x = 0.$$

□

Let $(M \otimes S)(c)$ stand for the generalized eigenspace of $D_{\mathfrak{g},\mathfrak{h}}(M)^2$ with eigenvalue $c \in \mathbb{R}$. Combining, now, Lemmas 3.1 and 3.2, we obtain the following corollary.

Corollary 3.3. *Let $M \in \mathcal{O}$. Then $M \otimes S$ can be decomposed into a direct sum*

$$M \otimes S = \bigoplus_{c \in \mathbb{R}} (M \otimes S)(c)$$

of generalized eigenspaces $(M \otimes S)(c)$ of $D_{\mathfrak{g},\mathfrak{h}}(M)^2$. Moreover, every generalized eigenspace $(M \otimes S)(c)$ is finitely generated.

Proof. Without loss of generality, we assume that $M \in \mathcal{O}_{\chi_\lambda}$. According to Lemma 3.1, $(M \otimes S)$ decomposes into a direct sum of generalized infinitesimal character subspaces

$$M \otimes S = \bigoplus_{\nu \in \mathfrak{t}^*} (M \otimes S)^{\xi_\nu}$$

while from Lemma 3.2, $D_{\mathfrak{g},\mathfrak{h}}(M)^2$ acts on each $(M \otimes S)^{\xi_\nu}$ by the scalar

$$c_\nu := \chi_\lambda(\Omega_{\mathfrak{g}}) + \|\rho\|^2 - \xi_\nu(\Omega_{\mathfrak{h}}) - \|\rho_{\mathfrak{h}}\|^2.$$

Hence, $M \otimes S$ decomposes into a direct sum of generalized eigenspaces $(M \otimes S)(c)$ for $D_{\mathfrak{g},\mathfrak{h}}(M)^2$ while

$$(M \otimes S)(c) = \bigoplus_{\substack{\nu \text{ s.t.} \\ c=c_\nu}} (M \otimes S)^{\xi_\nu}.$$

For the second statement, since there are finitely many weights ν of $M \otimes S$ satisfying the condition $c_\nu = c$, the above direct sum must be finite.

According to Lemma 3.1, every $(M \otimes S)^{\xi_\nu}$ is finitely generated and so this is the case for $(M \otimes S)(c)$. \square

Corollary 3.3 results in the following theorem.

Theorem 3.4 (Vogan's conjecture). *Let $M \in \mathcal{O}_{\chi_\lambda}$. Then $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ is finitely generated so that it belongs to the BGG category $\mathcal{O}(\mathfrak{h})$ for \mathfrak{h} . Moreover, $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ has generalized infinitesimal character ξ_λ .*

Proof. For the first statement, we note that $\ker D_{\mathfrak{g}, \mathfrak{h}}(M)$ is a submodule of the generalized eigenspace $(M \otimes S)(0)$ which according to Corollary 3.3 is finitely generated. Since the universal enveloping algebra $U(\mathfrak{h})$ of \mathfrak{h} is Noetherian [Kna96, Proposition 3.27], $\ker D_{\mathfrak{g}, \mathfrak{h}}(M)$ must be finitely generated so that $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ is finitely generated too.

The proof of the second statement follows from [HP06, Kos03]. More precisely, Huang and Pandžić in the case of (\mathfrak{g}, K) -modules and Kostant in the more general case of (\mathfrak{g}, H) -modules proved Vogan's conjecture [Vog97] by showing that there is an algebra morphism $\zeta : Z(\mathfrak{g}) \rightarrow Z(\mathfrak{h}_\Delta)$ such that for every $z \in Z(\mathfrak{g})$ there is $a \in (U(\mathfrak{g}) \otimes \mathbb{C}(\mathfrak{q}))^{\mathfrak{h}}$ such that

$$z - \zeta(z) = D_{\mathfrak{g}, \mathfrak{h}}(M)a + aD_{\mathfrak{g}, \mathfrak{h}}(M).$$

Then, $aD_{\mathfrak{g}, \mathfrak{h}}(M)$ acts trivially on $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$, while, for $x \in \ker D_{\mathfrak{g}, \mathfrak{h}}(M)$, since $D_{\mathfrak{g}, \mathfrak{h}}(M)^2$ is central in $(U(\mathfrak{g}) \otimes \mathbb{C})^{\mathfrak{h}}$, we have

$$D_{\mathfrak{g}, \mathfrak{h}}(M)(D_{\mathfrak{g}, \mathfrak{h}}(M)ax) = D_{\mathfrak{g}, \mathfrak{h}}(M)^2ax = aD_{\mathfrak{g}, \mathfrak{h}}(M)^2x = 0.$$

Hence $D_{\mathfrak{g}, \mathfrak{h}}(M)ax \in \ker D_{\mathfrak{g}, \mathfrak{h}}(M) \cap \text{im } D_{\mathfrak{g}, \mathfrak{h}}(M)$ so that $D_{\mathfrak{g}, \mathfrak{h}}(M)a$ acts trivially on $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$. Consequently, $z \otimes 1 - \zeta(z)$ acts trivially on $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$. In other words, $\zeta(z)$ acts by the scalar $\chi_\lambda(z)$ on $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ for every $z \in Z(\mathfrak{g})$. \square

In the last part of this section, we prove the following nonvanishing result for $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$.

Theorem 3.5. *Let M be a nonzero \mathfrak{g} -module in \mathcal{O} . Then*

$$H_D^{\mathfrak{g}, \mathfrak{h}}(M) \neq 0.$$

Proof. Let M be a nonzero \mathfrak{g} -module in \mathcal{O} . Let $\lambda \in \mathfrak{t}^*$ be the maximal weight of M and v_λ^+ a nonzero maximal weight vector of the corresponding weight subspace M_λ . It suffices to show that the vector $v_\lambda^+ \otimes \mathbf{1}$ belongs to $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$.

First, we show that $v_\lambda^+ \otimes \mathbf{1}$ is annihilated by $D_{\mathfrak{g}, \mathfrak{h}}(M)$. Let $\{e_\alpha\}_{\alpha \in \Delta_{\mathfrak{q}}}$ be a set of root vectors belonging to \mathfrak{q} and such that

$$\langle e_\alpha, e_{-\alpha} \rangle = 1 \text{ for every } \alpha \in \Delta_{\mathfrak{q}}.$$

For every $\alpha \in \Delta_{\mathfrak{q}}^+$, set

$$Z_{\alpha}^{(1)} := \frac{\sqrt{2}}{2}(e_{\alpha} + e_{-\alpha})$$

and

$$Z_{\alpha}^{(2)} := \frac{\sqrt{2}}{2i}(e_{\alpha} - e_{-\alpha}).$$

Then $\{Z_{\alpha}^{(i)} \mid \alpha \in \Delta_{\mathfrak{q}}^+, i = 1, 2\}$ is an orthonormal basis of \mathfrak{q} . Replacing this basis in (4) gives

$$D_{\mathfrak{g},\mathfrak{h}}(M) = \sqrt{2}D_+ + \sqrt{2}D_- - 1 \otimes \gamma(c)$$

where

$$D_{\pm} := \sum_{\alpha \in \Delta_{\mathfrak{q}}^+} \pi(e_{\pm\alpha}) \otimes \gamma(e_{\mp\alpha})$$

and $\gamma(c)$ is the cubic term (5). Then, for every $\alpha \in \Delta_{\mathfrak{q}}^+$, $\pi(e_{\alpha})$ annihilates v_{λ}^+ while $\gamma(e_{\alpha})$ annihilates $\mathbf{1}$ so that

$$D_{\pm}(v_{\lambda} \otimes \mathbf{1}) = 0.$$

On the other hand, the cubic term $\gamma(c)$ annihilates $\mathbf{1}$. More precisely, if V_0 stands for the 1-dimensional simple \mathfrak{g} -module of highest weight 0, replacing V_0 into (7) gives

$$\ker \gamma(c) \cong \ker D_{\mathfrak{g},\mathfrak{h}}(V_0) \cong \bigoplus_{w \in W^1} F_{w\rho - \rho_{\mathfrak{h}}},$$

where

$$W^1 := \{w \in W \mid \Delta_{\mathfrak{h}}^+ \subseteq w\Delta^+\}$$

and F_{μ} stands for the simple \mathfrak{h} -module of highest weight $\mu \in \mathfrak{t}^*$. For $w = 1$, $F_{\rho - \rho_{\mathfrak{h}}}$ is contained in $\ker \gamma(c)$ while the weight subspace $S_{\rho - \rho_{\mathfrak{h}}}$ of S is 1-dimensional and generated by $\mathbf{1}$. Therefore, $\mathbf{1} \in \ker \gamma(c)$ and we conclude that $v_{\lambda}^+ \otimes \mathbf{1} \in \ker D_{\mathfrak{g},\mathfrak{h}}(M)$.

Now we show that $v_{\lambda}^+ \otimes \mathbf{1} \notin \text{im } D_{\mathfrak{g},\mathfrak{h}}(M)$. Indeed, assume that the weight vector $v_{\lambda}^+ \otimes \mathbf{1} \in (M \otimes S)_{\lambda + \rho - \rho_{\mathfrak{h}}}$ belongs to $\text{im } D_{\mathfrak{g},\mathfrak{h}}(M)$. Then, there must be a weight vector $u \in (M \otimes S)_{\lambda + \rho - \rho_{\mathfrak{h}}}$ such that

$$D_{\mathfrak{g},\mathfrak{h}}(M)u = v_{\lambda}^+ \otimes \mathbf{1}.$$

But λ and $\rho - \rho_{\mathfrak{h}}$ are maximal weights of M and S respectively, so that u is a linear combination of tensor products $v_i \otimes \mathbf{1}$ with each v_i being a nonzero maximal weight vector of M_{λ} . Replacing v_{λ}^+ by v_i in the above argument, we deduce that $D_{\mathfrak{g},\mathfrak{h}}(M)$ annihilates every element $v_i \otimes \mathbf{1}$. Therefore,

$$0 = D_{\mathfrak{g},\mathfrak{h}}(M)u \neq v_{\lambda}^+ \otimes \mathbf{1}$$

and $v_\lambda^+ \otimes \mathbf{1} \notin \text{im } D_{\mathfrak{g}, \mathfrak{h}}(M)$. Hence $v_\lambda^+ \otimes \mathbf{1} \in H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ and $H_D^{\mathfrak{g}, \mathfrak{h}}(M)$ is nonzero. \square

4. APPLICATION TO SIMPLE VERMA MODULES

Let \mathfrak{g} be a complex semisimple Lie algebra and \mathfrak{h} a Lie subalgebra of \mathfrak{g} satisfying (8). Moreover, assume that \mathfrak{h} contains a Cartan subalgebra \mathfrak{t} of \mathfrak{g} . In this section, we provide an application for simple Verma modules using results from Section 3. More precisely, we show that the Dirac cohomology of a simple Verma \mathfrak{g} -module turns out to be a simple Verma \mathfrak{h} -module.

Theorem 4.1. *Let $M(\lambda)$ be the simple Verma module of highest weight $\lambda \in \mathfrak{t}^*$. Then*

$$H_D(M(\lambda)) \cong M_{\mathfrak{h}}(\lambda + \rho - \rho_{\mathfrak{h}}),$$

where $M_{\mathfrak{h}}(\lambda + \rho - \rho_{\mathfrak{h}})$ stands for the simple Verma \mathfrak{h} -module of highest weight $\lambda + \rho - \rho_{\mathfrak{h}}$.

Proof. Let $M(\lambda)$, $\lambda \in \mathfrak{t}^*$, be the Verma module with highest weight $\lambda \in \mathfrak{t}^*$, and suppose that $M(\lambda)$ is simple. Then λ is ρ -antidominant [Hum08, Section 4.8], i.e.

$$2 \frac{\langle \lambda + \rho, \alpha \rangle}{\|\alpha\|^2} \notin \mathbb{Z}_{>0} \text{ for every } \alpha \in \Delta^+.$$

According to Theorem 3.4, $H_D^{\mathfrak{g}, \mathfrak{h}}(M(\lambda))$ belongs to $\mathcal{O}(\mathfrak{h})$ so that it is of finite length, while, from Theorem 3.5, $H_D^{\mathfrak{g}, \mathfrak{h}}(M(\lambda))$ is nonzero. Let the simple \mathfrak{h} -module $L_{\mathfrak{h}}(\mu)$ of highest weight $\mu \in \mathfrak{t}^*$ be a composition factor of $H_D^{\mathfrak{g}, \mathfrak{h}}(M(\lambda))$. According to Theorem 3.4, $\mu + \rho_{\mathfrak{h}}$ must belong to the W -orbit of $\lambda + \rho$, i.e. $\mu = w(\lambda + \rho) - \rho_{\mathfrak{h}}$ for some $w \in W$. On the other hand, since λ is ρ -antidominant, $\lambda + \rho \leq w(\lambda + \rho)$ [Hum08, Proposition 3.5] so that $\mu = w(\lambda + \rho) - \rho_{\mathfrak{h}} \geq \lambda + \rho - \rho_{\mathfrak{h}}$. But $\lambda + \rho - \rho_{\mathfrak{h}}$ is the maximal weight of $M(\lambda) \otimes S$ and one concludes that $\mu = \lambda + \rho - \rho_{\mathfrak{h}}$. Therefore, every composition factor of $H_D^{\mathfrak{g}, \mathfrak{h}}(M(\lambda))$ is isomorphic to $L_{\mathfrak{h}}(\lambda + \rho - \rho_{\mathfrak{h}})$.

On the other hand, the weight subspace $(M \otimes S)_{\lambda + \rho - \rho_{\mathfrak{h}}}$ of $M \otimes S$ is generated by a nonzero highest weight vector v_λ^+ of $M(\lambda)$ tensorized with the weight vector $\mathbf{1}$ of S , and is 1-dimensional. Therefore, $L_{\mathfrak{h}}(\lambda + \rho - \rho_{\mathfrak{h}})$ appears only once in $H_D^{\mathfrak{g}, \mathfrak{h}}(M(\lambda))$ so that $H_D^{\mathfrak{g}, \mathfrak{h}}(M(\lambda)) \cong L_{\mathfrak{h}}(\lambda + \rho - \rho_{\mathfrak{h}})$. The equality $L_{\mathfrak{h}}(\lambda + \rho - \rho_{\mathfrak{h}}) = M_{\mathfrak{h}}(\lambda + \rho - \rho_{\mathfrak{h}})$ follows from the fact that the element $\lambda + \rho - \rho_{\mathfrak{h}}$ turns out to be $\rho_{\mathfrak{h}}$ -antidominant so that $M_{\mathfrak{h}}(\lambda + \rho - \rho_{\mathfrak{h}})$ is simple [Hum08, Section 4.8]. \square

5. DIRAC AND LIE ALGEBRA COHOMOLOGY FOR \mathcal{O}

In this section, we adapt the method used in [HPR06], in order to relate Dirac cohomology of unitary \mathfrak{g} -modules $M \in \mathcal{O}$ and nilpotent Lie algebra cohomology with coefficients in M .

Fix \mathfrak{g}_0 to be a θ -stable real form of \mathfrak{g} so that $\mathfrak{g}_0 = \mathfrak{k}_0 \oplus_{\theta} \mathfrak{p}_0$ is the corresponding Cartan decomposition.

Definition 5.1. A \mathfrak{g} -module $(\pi, M) \in \mathcal{O}$ is said to be unitary if it admits a Hermitian inner product $\langle \cdot, \cdot \rangle$ such that

$$\langle \pi(X)u, v \rangle + \langle u, \pi(X)v \rangle = 0$$

for every $u, v \in M$ and every $X \in \mathfrak{g}_0$.

In other words, $\pi(X)^{\text{adj}} = -\pi(X)$ for every $X \in \mathfrak{g}_0$. One can easily check that, if $\tilde{\cdot}$ stands for the conjugation with respect to \mathfrak{g}_0 , then

$$(15) \quad \pi(X)^{\text{adj}} = -\pi(\tilde{X}) \text{ for every } X \in \mathfrak{g}.$$

For more details concerning the classification of unitary highest weight modules, see [EHW83] for the case when \mathfrak{g}_0 is a compact real form of \mathfrak{g} , and [GL90] for the case when \mathfrak{g}_0 is a noncompact real form of \mathfrak{g} .

Assume now that \mathfrak{g} and $\mathfrak{k} := (\mathfrak{k}_0)^{\mathbb{C}}$ form a Hermitian symmetric pair. Then, $\text{rank}_{\mathbb{C}}(\mathfrak{g}) = \text{rank}_{\mathbb{C}}(\mathfrak{k})$ and $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}^+ \oplus \mathfrak{p}^-$, with \mathfrak{p}^+ and \mathfrak{p}^- being maximal isotropic dual subspaces of $\mathfrak{p} := (\mathfrak{p}_0)^{\mathbb{C}}$ with respect to the restriction of the Killing form on \mathfrak{p} . Moreover \mathfrak{p} is abelian and $\mathfrak{k} \oplus \mathfrak{p}^+$ is a parabolic subalgebra of \mathfrak{g} while we make the assumption that $\mathfrak{k} \oplus \mathfrak{p}^+ \supseteq \mathfrak{b}$.

Theorem 5.2. *Let $\mathfrak{g}, \mathfrak{k}$ be a Hermitian symmetric pair and M a unitary module belonging to \mathcal{O} . Then*

$$H_D^{\mathfrak{g}, \mathfrak{k}}(M) \cong \ker D_{\mathfrak{g}, \mathfrak{k}}(M) \cong H(\mathfrak{p}^+, M) \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}} \cong H(\mathfrak{p}^-, M) \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}},$$

where $H(\mathfrak{p}^+, M)$ (respectively $H(\mathfrak{p}^-, M)$) stands for the nilpotent Lie algebra cohomology (respectively homology) with coefficients in M .

In what follows, in order to prove Theorem 5.2, we provide a Hodge decomposition for the operator $D_{\mathfrak{g}, \mathfrak{k}}(M)$. Let $\{Z_i\}$ be an orthonormal basis of \mathfrak{p}_0 and set

$$(16) \quad e_j := \frac{Z_{2j-1} + iZ_{2j}}{\sqrt{2}}, \quad e_{-j} := \frac{Z_{2j-1} - iZ_{2j}}{\sqrt{2}}.$$

Then $\{e_j\}$ and $\{e_{-j}\}$ are dual bases of the maximal isotropic dual subspaces $\mathfrak{p}^+ := \text{span}\{e_j\}_j$ and $\mathfrak{p}^- := \text{span}\{e_{-j}\}_j$ of \mathfrak{p} , respectively. Define operators

$$C^+ := \sum_j \pi(e_j) \otimes \gamma(e_{-j})$$

$$C^- := \sum_j \pi(e_{-j}) \otimes \gamma(e_j)$$

so that $C^+ + C^- = D_{\mathfrak{g}, \mathfrak{k}}(M)$.

Since \mathfrak{p}^+ and \mathfrak{p}^- are maximal isotropic dual subspaces of \mathfrak{p} ,

$$S \cong \wedge \mathfrak{p}^- \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}}$$

and

$$(17) \quad \begin{aligned} M \otimes S &\cong M \otimes \wedge \mathfrak{p}^- \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}} \cong M \otimes \wedge (\mathfrak{p}^+)^* \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}} \\ &\cong M \otimes (\wedge \mathfrak{p}^+)^* \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}} \cong \text{Hom}(\wedge \mathfrak{p}^+, M) \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}} \end{aligned}$$

while $\text{Hom}(\wedge \mathfrak{p}^+, M)$ provides the Chevalley-Eilenberg resolution which calculates the nilpotent Lie algebra cohomology with coefficients in M . The differential

$$d^k : \text{Hom}(\wedge^k \mathfrak{p}^+, M) \rightarrow \text{Hom}(\wedge^{k+1} \mathfrak{p}^+, M)$$

of this resolution is given by

$$\begin{aligned} d^k \omega(X_0 \wedge \dots \wedge X_k) &:= \sum_{i=0}^k (-1)^i X_i (\omega(X_0 \wedge \dots \wedge \widehat{X}_i \wedge \dots \wedge X_k)) \\ &+ \sum_{i < j} (-1)^{i+j} \omega([X_i, X_j] \wedge X_0 \wedge \dots \wedge \widehat{X}_i \wedge \dots \wedge \widehat{X}_j \wedge \dots \wedge X_k), \end{aligned}$$

where $X_i \in \mathfrak{p}^+$, $\omega \in \text{Hom}(\wedge^k \mathfrak{p}^+, M)$ and $\widehat{}$ indicates that we omit the corresponding element.

On the other hand, $M \otimes \wedge \mathfrak{p}^-$ provides the resolution for the nilpotent Lie algebra homology with the differential

$$\partial^k : M \otimes \wedge^k \mathfrak{p}^- \rightarrow M \otimes \wedge^{k-1} \mathfrak{p}^-$$

given by:

$$\begin{aligned} \partial^k (v \otimes Y_1 \wedge \dots \wedge Y_{k+1}) &:= \sum_{i=1}^{k+1} (-1)^{i+1} Y_i v \otimes Y_1 \wedge \dots \wedge \widehat{Y}_i \wedge \dots \wedge Y_{k+1} \\ &+ \sum_{i < j} (-1)^{i+j} v \otimes [Y_i, Y_j] \wedge Y_1 \wedge \dots \wedge \widehat{Y}_i \wedge \dots \wedge \widehat{Y}_j \wedge \dots \wedge Y_{k+1}, \end{aligned}$$

Lemma 5.3. *Under the identification (17), $C^+ = d$ and $C^- = \partial$ so that*

$$D_{\mathfrak{g}, \mathfrak{k}}(M) = d + \partial.$$

Proof. See [HP06, Section 9.1.4]. \square

Lemma 5.4. *Let M be a unitary \mathfrak{g} -module. Then, there is a Hermitian inner product $\langle \cdot, \cdot \rangle$ on $M \otimes S$ such that*

$$(C^\pm)^{\text{adj}} = -C^\mp.$$

Hence $D_{\mathfrak{g}, \mathfrak{k}}(M)$ is skew-adjoint.

Proof. According to [Afe21a, HP06], $S = \bigwedge \mathfrak{p}^-$ admits a Hermitian inner product $\langle \cdot, \cdot \rangle_S$ and

$$\gamma(X)^{\text{adj}} = \gamma(\tilde{X}) \text{ for every } X \in \mathfrak{p},$$

where $\tilde{\cdot}$ stands for the conjugation with respect to \mathfrak{g}_0 . On the other hand, by hypothesis and (15), there is a Hermitian inner product $\langle \cdot, \cdot \rangle_M$ on M such that

$$\pi(X)^{\text{adj}} = -\pi(\tilde{X}) \text{ for every } X \in \mathfrak{p}.$$

On $M \otimes S$ define the natural inner product $\langle \cdot, \cdot \rangle$ coming from tensorizing $\langle \cdot, \cdot \rangle_S$ and $\langle \cdot, \cdot \rangle_M$, i.e.

$$\langle v_1 \otimes u_1, v_2 \otimes u_2 \rangle := \langle v_1, v_2 \rangle_M \langle u_1, u_2 \rangle_S$$

for every $v_1 \otimes u_1, v_2 \otimes u_2 \in M \otimes S$. Then, if e_j and e_{-j} are as in (16), we have

$$\begin{aligned} (\pi(e_j) \otimes \gamma(e_{-j}))^{\text{adj}} &= \pi(e_j)^{\text{adj}} \otimes \gamma(e_{-j})^{\text{adj}} \\ &= -\pi(e_{-j}) \otimes \gamma(e_j) \end{aligned}$$

so that $(C^+)^{\text{adj}} = -C^-$, $(C^-)^{\text{adj}} = -C^+$ and

$$D_{\mathfrak{g}, \mathfrak{k}}(M)^{\text{adj}} = -D_{\mathfrak{g}, \mathfrak{k}}(M).$$

\square

Lemma 5.5. *Let $(\mathfrak{g}, \mathfrak{k})$ be a Hermitian symmetric pair and M a unitary module in \mathcal{O} . Then*

$$M \otimes S = \ker D_{\mathfrak{g}, \mathfrak{k}}(M) \oplus \text{im } D_{\mathfrak{g}, \mathfrak{k}}(M).$$

Proof. According to Corollary 3.3,

$$M \otimes S = (M \otimes S)(0) \oplus \bigoplus_{c \neq 0} (M \otimes S)(c),$$

where $(M \otimes S)(c)$, $c \in \mathbb{R}$, stands for the generalized eigenspace of $D_{\mathfrak{g}, \mathfrak{k}}(M)^2$ of eigenvalue c . It suffices to show that

$$\ker D_{\mathfrak{g}, \mathfrak{k}}(M) = (M \otimes S)(0)$$

$$\text{and } \text{im } D_{\mathfrak{g}, \mathfrak{k}}(M) = \bigoplus_{c \neq 0} (M \otimes S)(c).$$

For the first equality, since $D_{\mathfrak{g},\mathfrak{k}}(M)$ is skew-adjoint, $\ker D_{\mathfrak{g},\mathfrak{k}}(M)^2 = \ker D_{\mathfrak{g},\mathfrak{k}}(M)$. Therefore, one easily checks that $\ker D_{\mathfrak{g},\mathfrak{k}}(M)^k = \ker D_{\mathfrak{g},\mathfrak{k}}(M)$ for every $k > 0$ so that $\ker D_{\mathfrak{g},\mathfrak{k}}(M) = (M \otimes S)(0)$.

For the second equality, using again the fact that $D_{\mathfrak{g},\mathfrak{k}}(M)$ is skew-adjoint and the first equality, we obtain

$$\begin{aligned} \operatorname{im} D_{\mathfrak{g},\mathfrak{k}}(M) &= \ker D_{\mathfrak{g},\mathfrak{k}}(M)^\perp \\ &= (M \otimes S)(0)^\perp \\ &= \bigoplus_{c \neq 0} (M \otimes S)(c) \end{aligned}$$

□

Proposition 5.6. *Let $(\mathfrak{g}, \mathfrak{k})$ be a Hermitian symmetric pair and M a unitary module in \mathcal{O} . Then*

- (i) $\ker C^+ = \operatorname{im} C^+ \oplus \ker D_{\mathfrak{g},\mathfrak{k}}(M)$
- (ii) $\ker C^- = \operatorname{im} C^- \oplus \ker D_{\mathfrak{g},\mathfrak{k}}(M)$.

The above direct sums are orthogonal with respect to $\langle \cdot, \cdot \rangle$.

Proof. The proof is the same as in the case of unitary (\mathfrak{g}, K) -modules [HP06, Lemma 9.3.3 & Proposition 9.3.4]. □

Consequently, we obtain

$$H_D^{\mathfrak{g},\mathfrak{k}}(M) \cong \ker D_{\mathfrak{g},\mathfrak{k}}(M) \cong H(\mathfrak{p}^+, M) \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}} \cong H(\mathfrak{p}^-, M) \otimes \mathbb{C}_{\rho - \rho_{\mathfrak{k}}}$$

which is exactly the statement of Theorem 5.2. Here, the first equivalence is due to Lemma 5.5 while the second and the third ones due to Lemma 5.3 and Proposition 5.6.

6. HIGHER DIRAC COHOMOLOGY AND INDEX FOR THE BGG CATEGORY \mathcal{O}

Let \mathfrak{g} be a complex semisimple Lie algebra and \mathfrak{h} a Lie subalgebra of \mathfrak{g} satisfying (8). Let \mathfrak{h} contain a Cartan subalgebra \mathfrak{t} of \mathfrak{g} , so that the spin module S has a splitting into $S = S^+ \oplus S^-$. Let X_1, X_2, X_3 and X be \mathfrak{g} -modules such that the restrictions $(X_1)|_{\mathfrak{h}}, (X_2)|_{\mathfrak{h}}, (X_3)|_{\mathfrak{h}}$ and $X|_{\mathfrak{h}}$ to \mathfrak{h} decompose into algebraic finite-multiplicity direct sums of finite dimensional simple \mathfrak{h} -modules. Assume that X_1, X_2, X_3 fit into a short exact sequence

$$0 \longrightarrow X_1 \longrightarrow X_2 \longrightarrow X_3 \longrightarrow 0.$$

In the case where each X_1, X_2, X_3 , and X has infinitesimal character (or it is a direct sum of modules having infinitesimal characters), we can construct \mathfrak{h} -morphisms so that the following diagram of Dirac cohomology spaces is exact [HP06, Subsection 9.6.2]:

$$(18) \quad \begin{array}{ccccc} H_D^{\mathfrak{g},\mathfrak{h}}(X_1)^+ & \longrightarrow & H_D^{\mathfrak{g},\mathfrak{h}}(X_2)^+ & \longrightarrow & H_D^{\mathfrak{g},\mathfrak{h}}(X_3)^+ \\ \uparrow & & & & \downarrow \\ H_D^{\mathfrak{g},\mathfrak{h}}(X_3)^- & \longleftarrow & H_D^{\mathfrak{g},\mathfrak{h}}(X_2)^- & \longleftarrow & H_D^{\mathfrak{g},\mathfrak{h}}(X_1)^- \end{array}$$

while for the Dirac index $I_D(X)$ of X , we have [MPV17]:

$$(19) \quad I_D(X) = X \otimes S^+ - X \otimes S^-.$$

Nevertheless, when X_1 , X_2 , X_3 and X , instead of infinitesimal characters, have generalized infinitesimal characters, the above statements may fail [PS16, Section 2].

In order to overcome this problem, Pandžić and Somberg defined the following notion of higher Dirac cohomology [PS16]:

Definition 6.1. For \mathfrak{g} , \mathfrak{h} and X as above, we call k -th higher Dirac cohomology of X the \mathfrak{h} -module

$$H_{\text{top}}^k(X) := \frac{\ker D_{\mathfrak{g},\mathfrak{h}}(X)^{2k+1}}{\ker D_{\mathfrak{g},\mathfrak{h}}(X)^{2k+1} \cap \text{im } D_{\mathfrak{g},\mathfrak{h}}(X) + \ker D_{\mathfrak{g},\mathfrak{h}}(X)^{2k}}$$

and higher Dirac cohomology of X the \mathfrak{h} -module

$$H_{\text{top}}(X) := \bigoplus_{k \geq 0} H_{\text{top}}^k(X).$$

Using the above definition of higher Dirac cohomology, they introduced a notion of higher Dirac index:

Definition 6.2. For \mathfrak{g} , \mathfrak{h} and X as above, the virtual \mathfrak{h} -module

$$I_{\text{top}}(X) := H_{\text{top}}(X)^+ - H_{\text{top}}(X)^-$$

is called the higher Dirac index of X .

Remark 6.3. Note that in the case where X has infinitesimal character, $H_D^{\mathfrak{g},\mathfrak{h}}(X)$ (respectively $I_D(X)$) and $H_{\text{top}}(X)$ (respectively $I_{\text{top}}(X)$) coincide.

Then, they proved that if, instead of the standard notions of Dirac cohomology $H_D^{\mathfrak{g},\mathfrak{h}}(X)$ and index $I_D(X)$, we use the higher Dirac cohomology $H_{\text{top}}(X)$ and index $I_{\text{top}}(X)$, the statements for (18) and (19) remain true whenever X_1 , X_2 , X_3 and X have generalized infinitesimal characters. The proofs of their results are based on the fact that the generalized kernel $(X \otimes S)(0)$ of $D_{\mathfrak{g},\mathfrak{h}}(X)$ is finite dimensional and can be decomposed into Jordan blocks for $D_{\mathfrak{g},\mathfrak{h}}(X)$.

Remark 6.4. For X_1, X_2 and X_3 having generalized infinitesimal characters, the maps used in the corresponding diagram (18) depend on the chosen Jordan decomposition and, so, are noncanonical.

In this section, we prove similar results in the case where, instead of \mathfrak{g} -modules X_1, X_2, X_3 and X satisfying the previous finiteness condition for their restrictions to \mathfrak{h} , we have \mathfrak{g} -modules M_1, M_2, M_3 and M belonging to \mathcal{O} . Of course, we can again have a notion of higher Dirac cohomology $H_{\text{top}}(M)$ and index $I_{\text{top}}(M)$. Nevertheless, the generalized kernel $(M \otimes S)(0)$ of $D_{\mathfrak{g},\mathfrak{h}}(M)$ need not be finite dimensional and may not have a Jordan decomposition for $D_{\mathfrak{g},\mathfrak{h}}(M)$. In order to overcome this problem, we decompose $(M \otimes S)(0)$ into a direct sum of weight subspaces

$$(M \otimes S)(0) = \bigoplus_{\mu \in \mathfrak{t}^*} [(M \otimes S)(0)]_{\mu},$$

where each weight subspace $[(M \otimes S)(0)]_{\mu}$ is finite dimensional. On the other hand, since $\mathfrak{t} \subseteq \mathfrak{h}$ and $D_{\mathfrak{g},\mathfrak{h}}(M)$ is \mathfrak{h} -equivariant, $D_{\mathfrak{g},\mathfrak{h}}(M)$ is \mathfrak{t} -equivariant so that it preserves each weight subspace. Now, each weight subspace $[(M \otimes S)(0)]_{\mu}$ has a Jordan block decomposition for $D_{\mathfrak{g},\mathfrak{h}}(M)$ and we can apply the methods used in [PS16] and construct maps so that

$$(20) \quad \begin{array}{ccccc} H_{\text{top}}(M_1)_{\mu}^{+} & \longrightarrow & H_{\text{top}}(M_2)_{\mu}^{+} & \longrightarrow & H_{\text{top}}(M_3)_{\mu}^{+} \\ \uparrow & & & & \downarrow \\ H_{\text{top}}(M_3)_{\mu}^{-} & \longleftarrow & H_{\text{top}}(M_2)_{\mu}^{-} & \longleftarrow & H_{\text{top}}(M_1)_{\mu}^{-} \end{array}$$

forms an exact circle of weight subspaces, i.e. of \mathfrak{t} -modules, while for the weight subspaces $I_{\text{top}}(M)_{\mu}$ and $(M \otimes S^{\pm})_{\mu}$ we have

$$(21) \quad I_{\text{top}}(M)_{\mu} = (M \otimes S^{+})_{\mu} - (M \otimes S^{-})_{\mu}.$$

We start with the result concerning the higher Dirac index which is the analogue of [PS16, Theorem 1.11].

Proposition 6.5. *Let $M \in \mathcal{O}$. Then*

$$I_{\text{top}}(M) = M \otimes S^{+} - M \otimes S^{-}$$

Proof of Proposition 6.5. Let $(M \otimes S)(c)$ be the generalized eigenspace of $D_{\mathfrak{g},\mathfrak{h}}(M)^2$ with eigenvalue $c \in \mathbb{R}$. Since $D_{\mathfrak{g},\mathfrak{h}}(M)^2$ is even in the Clifford factor, $(M \otimes S)(c)$ decomposes into

$$(M \otimes S)(c) := (M \otimes S)(c)^{+} \oplus (M \otimes S)(c)^{-}.$$

Moreover, since $D_{\mathfrak{g},\mathfrak{h}}(M)$ is \mathfrak{t} -equivariant, one obtains a more refined decomposition in terms of weight subspaces:

$$[(M \otimes S)(c)]_{\mu} := [(M \otimes S)(c)^+]_{\mu} \oplus [(M \otimes S)(c)^-]_{\mu}.$$

If $c \neq 0$, $D_{\mathfrak{g},\mathfrak{h}}(M)$ maps injectively $[(M \otimes S)(c)^{\pm}]_{\mu}$ into $[(M \otimes S)(c)^{\mp}]_{\mu}$. More precisely, if $x \in [(M \otimes S)(c)^{\pm}]_{\mu}$ and $D_{\mathfrak{g},\mathfrak{h}}(M)x = 0$, then $D_{\mathfrak{g},\mathfrak{h}}(M)^2x = 0$ and so x must be 0. Since $[(M \otimes S)(c)^{\pm}]_{\mu}$ is finite dimensional, one deduces that the restriction

$$D_{\mathfrak{g},\mathfrak{h}}(M) : [(M \otimes S)(c)^{\pm}]_{\mu} \rightarrow [(M \otimes S)(c)^{\mp}]_{\mu}$$

is an isomorphism. Hence, $(M \otimes S)(c)^+ \cong (M \otimes S)(c)^-$ for every $c \neq 0$ so that

$$M \otimes S^+ - M \otimes S^- = (M \otimes S)(0)^+ - (M \otimes S)(0)^-.$$

The following lemma completes the proof. \square

Lemma 6.6. *Let $M \in \mathcal{O}$. Then*

$$(M \otimes S)(0)^+ - (M \otimes S)(0)^- = H_{\text{top}}(M)^+ - H_{\text{top}}(M)^-.$$

Proof. Since we are working in the Grothendieck's group of \mathfrak{h} , it suffices to prove the statement for the corresponding formal characters. In other words, it suffices to show that, for every $\mu \in \mathfrak{t}^*$,

$$\dim[(M \otimes S)(0)_{\mu}^+] - \dim[(M \otimes S)(0)_{\mu}^-] = \dim H_{\text{top}}(M)_{\mu}^+ - \dim H_{\text{top}}(M)_{\mu}^-.$$

For every $\mu \in \mathfrak{t}^*$, $[(M \otimes S)(0)]_{\mu}$ is finite dimensional and can be decomposed into Jordan blocks for $D_{\mathfrak{g},\mathfrak{h}}(M)$:

$$[(M \otimes S)(0)]_{\mu} = J_1^{(\mu)} \oplus \dots \oplus J_{n_{\mu}}^{(\mu)}.$$

For every $j \in \{1, \dots, n_{\mu}\}$, the Jordan block $J_j^{(\mu)}$ can be decomposed into a direct sum of 1-dimensional subspaces

$$W_1^{(\mu,j)} \oplus W_2^{(\mu,j)} \oplus \dots \oplus W_{m_{\mu,j}}^{(\mu,j)}$$

such that

$$D_{\mathfrak{g},\mathfrak{h}}(M)W_1^{(\mu,j)} = 0$$

and

$$D_{\mathfrak{g},\mathfrak{h}}(M) : W_i^{(\mu,j)} \xrightarrow{\cong} W_{i-1}^{(\mu,j)} \text{ for } i = 2, \dots, m_{\mu,j}.$$

We say that the Jordan block $J_j^{(\mu)}$ is of size $m_{\mu,j}$. Moreover, we can choose each $W_i^{(\mu,j)}$ to be contained either in $(M \otimes S)(0)^+$ or in $(M \otimes S)(0)^-$. For every $k > 0$, set

$$N_{\mu,k} := \bigoplus_{j=1}^{n_{\mu}} W_k^{(\mu,j)} \text{ and } N_{\mu,k}^{\pm} = N_{\mu,k} \cap (M \otimes S)^{\pm}.$$

$$\begin{array}{ccccccc}
& & & N_{\mu,3} & & N_{\mu,2} & & N_{\mu,1} & & \\
& & & \vdots & & \vdots & & \vdots & & \\
J_1^{(\mu)} : & \dots & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_3^{(\mu,1)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_2^{(\mu,1)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_1^{(\mu,1)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & 0 \\
J_2^{(\mu)} : & \dots & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_3^{(\mu,2)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_2^{(\mu,2)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_1^{(\mu,2)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & 0 \\
J_3^{(\mu)} : & \dots & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_3^{(\mu,3)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_2^{(\mu,3)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & \boxed{W_1^{(\mu,3)}} & \xrightarrow{D_{\mathfrak{g},\mathfrak{h}}(M)} & 0 \\
& & & \vdots & & \vdots & & \vdots & &
\end{array}$$

FIGURE 1. The above decomposition of $[(M \otimes S)(c)]_\mu$

Note that $N_{\mu,k} = N_{\mu,k}^+ \oplus N_{\mu,k}^-$ while by convention $W_k^{(\mu,j)} = 0$ whenever $k > m_{\mu,j}$. Then

$$(\ker D_{\mathfrak{g},\mathfrak{h}}(M)_\mu^{2k+2})^\pm = N_{\mu,1}^\pm \oplus \dots \oplus N_{\mu,2k+2}^\pm$$

and

$$\operatorname{im} D_{\mathfrak{g},\mathfrak{h}}(M)|_{N_{\mu,2k+2}^\mp} = \bigoplus_{\substack{j=1 \text{ s.t.} \\ m_{\mu,j} \geq 2k+2}}^{n_\mu} (W_{2k+1}^{(\mu,j)})^\pm \cong N_{\mu,2k+2}^\mp$$

as vector spaces. Therefore

$$\begin{aligned}
\frac{N_{\mu,2k+1}^\pm}{\operatorname{im} D_{\mathfrak{g},\mathfrak{h}}(M)|_{N_{\mu,2k+2}^\mp}} &\cong \bigoplus_{\substack{j=1 \text{ s.t.} \\ m_{\mu,j} < 2k+2}}^{n_\mu} (W_{2k+1}^{(\mu,j)})^\pm = \bigoplus_{\substack{j=1 \text{ s.t.} \\ m_{\mu,j} = 2k+1}}^{n_\mu} (W_{2k+1}^{(\mu,j)})^\pm \\
&\cong \frac{(\ker D_{\mathfrak{g},\mathfrak{h}}(M)_\mu^{2k+1})^\pm}{(\ker D_{\mathfrak{g},\mathfrak{h}}(M)_\mu^{2k+1} \cap \operatorname{im} D_{\mathfrak{g},\mathfrak{h}}(M)_\mu + \ker D_{\mathfrak{g},\mathfrak{h}}(M)_\mu^{2k})^\pm} \cong H_{\text{top}}^k(M)_\mu^\pm
\end{aligned}$$

so that

$$\begin{aligned}
\dim H_{\text{top}}^k(M)_\mu^\pm &= \dim N_{\mu,2k+1}^\pm - \dim \operatorname{im} D_{\mathfrak{g},\mathfrak{h}}(M)|_{N_{\mu,2k+2}^\mp} \\
&= \dim N_{\mu,2k+1}^\pm - \dim N_{\mu,2k+2}^\mp.
\end{aligned}$$

Then

$$\begin{aligned}
\dim H_{\text{top}}(M)_\mu^+ - \dim H_{\text{top}}(M)_\mu^- &= \sum_{k \geq 0} \dim H_{\text{top}}^k(M)_\mu^+ - \sum_{k \geq 0} \dim H_{\text{top}}^k(M)_\mu^- \\
&= \sum_{k \geq 0} (\dim N_{\mu, 2k+1}^+ - \dim N_{\mu, 2k+2}^-) \\
&\quad - \sum_{k \geq 0} (\dim N_{\mu, 2k+1}^- - \dim N_{\mu, 2k+2}^+) \\
&= \sum_{k \geq 0} (\dim N_{\mu, 2k+1}^+ + \dim N_{\mu, 2k+2}^+) \\
&\quad - \sum_{k \geq 0} (\dim N_{\mu, 2k+1}^- + \dim N_{\mu, 2k+2}^-) \\
&= \sum_{k \geq 1} \dim N_{\mu, k}^+ - \sum_{k \geq 1} \dim N_{\mu, k}^- \\
&= (M \otimes S)(0)_\mu^+ - (M \otimes S)(0)_\mu^-.
\end{aligned}$$

□

The following theorem concerning the exact circle (18) is the analogue of [PS16, Theorem 1.13]. Nevertheless, the maps that we obtain are \mathfrak{t} -equivariant.

Theorem 6.7. *Let $M_1, M_2, M_3 \in \mathcal{O}$ and*

$$0 \longrightarrow M_1 \longrightarrow M_2 \longrightarrow M_3 \longrightarrow 0$$

be an exact sequence. Then,

$$(22) \quad \begin{array}{ccccc}
H_{\text{top}}(M_1)^+ & \longrightarrow & H_{\text{top}}(M_2)^+ & \longrightarrow & H_{\text{top}}(M_3)^+ \\
\uparrow & & & & \downarrow \\
H_{\text{top}}(M_3)^- & \longleftarrow & H_{\text{top}}(M_2)^- & \longleftarrow & H_{\text{top}}(M_1)^-
\end{array}$$

forms an exact circle of \mathfrak{t} -modules.

Proof. We adapt the proof of [PS16, Theorem 1.13]; instead of \mathfrak{h} -isotypic components we consider finite dimensional weight subspaces.

Since tensorizing with a finite dimensional module is an exact functor, we obtain the exact sequence of \mathfrak{h} -modules:

$$(23) \quad 0 \longrightarrow M_1 \otimes S \xrightarrow{i} M_2 \otimes S \xrightarrow{p} M_3 \otimes S \longrightarrow 0.$$

We note that the maps i and p of the above exact sequence commute with the action of $U(\mathfrak{g}) \otimes \mathbf{C}(\mathfrak{q})$. As a consequence, they commute with the

corresponding Dirac operators. Namely

$$\begin{aligned} i \circ D_{\mathfrak{g},\mathfrak{h}}(M_1) &= D_{\mathfrak{g},\mathfrak{h}}(M_2) \circ i \\ \text{and } p \circ D_{\mathfrak{g},\mathfrak{h}}(M_2) &= D_{\mathfrak{g},\mathfrak{h}}(M_3) \circ p. \end{aligned}$$

Each \mathfrak{h} -module $M_i \otimes S$, $i = 1, 2, 3$, admits a direct sum decomposition in terms of generalized eigenspaces for $D_{\mathfrak{g},\mathfrak{h}}(M_i)^2$ and in terms of weight subspaces while these decompositions are compatible. Therefore, (23) breaks up into a direct sum of exact sequences

$$0 \longrightarrow (M_1 \otimes S)(c)_\mu \xrightarrow{i} (M_2 \otimes S)(c)_\mu \xrightarrow{p} (M_3 \otimes S)(c)_\mu \longrightarrow 0,$$

where $\mu \in \mathfrak{t}^*$ and $(M_i \otimes S)(c)_\mu$ is the corresponding weight subspace of the generalized eigenspace with eigenvalue c . We are interested in the case of the generalized kernel of $D_{\mathfrak{g},\mathfrak{h}}(M_i)$, i.e. when $c = 0$:

$$(24) \quad 0 \longrightarrow (M_1 \otimes S)(0)_\mu \xrightarrow{i} (M_2 \otimes S)(0)_\mu \xrightarrow{p} (M_3 \otimes S)(0)_\mu \longrightarrow 0.$$

Since $(M_3 \otimes S)(0)_\mu$ is finite dimensional, it can be decomposed into a direct sum of Jordan blocks $({}^3J_1^\mu, \dots, {}^3J_{n_\mu}^\mu$ for $D_{\mathfrak{g},\mathfrak{h}}(M_3)$. As we saw in the proof of Lemma 6.6, every such block $({}^3J_j^\mu$ can be decomposed into a direct sum

$$({}^3J_j^\mu = ({}^3W_1^{(\mu,j)} \oplus \dots \oplus ({}^3W_{m_{\mu,j}}^{(\mu,j)}$$

such that

$$D_{\mathfrak{g},\mathfrak{h}}(M_3)({}^3W_1^{(\mu,j)}) = 0$$

and

$$D_{\mathfrak{g},\mathfrak{h}}(M_3) : ({}^3W_i^{(\mu,j)}) \xrightarrow{\cong} ({}^3W_{i-1}^{(\mu,j)}) \text{ for } i = 2, \dots, m_{\mu,j}.$$

For every top summand $({}^3W_{m_{\mu,j}}^{(\mu,j)})$, let $({}^2W_{l_{\mu,j}}^{(\mu,j)})$ be a preimage in $(M_2 \otimes S)(0)_\mu$. Then, $({}^2W_{l_{\mu,j}}^{(\mu,j)})$ must be a top summand of a Jordan block of $(M_2 \otimes S)(0)_\mu$. Indeed, if we suppose that there exists $({}^2W_{l_{\mu,j}+1}^{(\mu,j)}) \neq 0$ such that

$$D_{\mathfrak{g},\mathfrak{h}}(M_2) : ({}^2W_{l_{\mu,j}+1}^{(\mu,j)}) \xrightarrow{\cong} ({}^2W_{l_{\mu,j}}^{(\mu,j)}),$$

then

$$({}^3W_{m_{\mu,j}}^{(\mu,j)}) = p \circ D_{\mathfrak{g},\mathfrak{h}}(M_2)({}^2W_{l_{\mu,j}+1}^{(\mu,j)}) = D_{\mathfrak{g},\mathfrak{h}}(M_3) \circ p({}^2W_{l_{\mu,j}+1}^{(\mu,j)}) = 0$$

which is a contradiction. Hence, $({}^2W_{l_{\mu,j}}^{(\mu,j)})$ is the top summand of a Jordan block, say $({}^2J_j^\mu)$, in $(M_2 \otimes S)(0)_\mu$. Then

$$({}^1J_j^\mu := \ker ({}^2J_j^\mu \rightarrow ({}^3J_j^\mu)$$




FIGURE 2. Decomposition of (24) into compatible Jordan blocks

is a Jordan block of length $k_{\mu,j} := l_{\mu,j} - m_{\mu,j}$ in $(M_1 \otimes S)_\mu$, say of top summand $(1)W_{k_{\mu,j}}^{(\mu,j)}$, and

$$0 \longrightarrow (1)J_j^\mu \xrightarrow{i} (2)J_j^\mu \xrightarrow{p} (3)J_j^\mu \longrightarrow 0$$

forms a short exact sequence of vector spaces.

Then, either $k_{\mu,j}$, $l_{\mu,j}$ and $m_{\mu,j}$ are all even so that the corresponding Jordan blocks do not contribute to the higher Dirac cohomology, or exactly one of them is even while the other two are odd. Assume that $k_{\mu,j}$ is even. Then,

$$\begin{aligned}
 H_{\text{top}}((1)J_j^\mu) &= 0 \\
 H_{\text{top}}((2)J_j^\mu) &= (2)W_{l_{\mu,j}+1}^{(\mu,j)} \\
 \text{and } H_{\text{top}}((3)J_j^\mu) &= (3)W_{k_{\mu,j}+1}^{(\mu,j)}
 \end{aligned}$$

so that

$$\begin{array}{ccc}
 & H_{\text{top}}((1)J_j^\mu) & \\
 & \nearrow 0 & \searrow 0 \\
 H_{\text{top}}((3)J_j^\mu) & \xleftarrow{1} & H_{\text{top}}((2)J_j^\mu)
 \end{array}$$

forms an exact circle. In the same way, we treat the other two cases. Therefore, applying the same procedure for the other Jordan blocks of $(M_3 \otimes$

$S)(0)_\mu$, we obtain an exact circle

$$\begin{array}{ccc} & H_{\text{top}}(M_1)_\mu & \\ & \nearrow & \searrow \\ H_{\text{top}}(M_3)_\mu & \longleftarrow & H_{\text{top}}(M_2)_\mu \end{array}$$

Taking direct sums over $\mu \in \mathfrak{t}^*$, we obtain an exact circle of \mathfrak{t} -modules

$$\begin{array}{ccc} & H_{\text{top}}(M_1) & \\ & \nearrow & \searrow \\ H_{\text{top}}(M_3) & \longleftarrow & H_{\text{top}}(M_2) \end{array}$$

while considering the natural \mathbb{Z}_2 -gradings, we obtain the exact circle (22). \square

We close this section with the analogue of [PS16, Theorem 1.10].

Theorem 6.8. *Let $M \in \mathcal{O}_{\chi_\lambda}$. Then $H_{\text{top}}(M) \in \mathcal{O}(\mathfrak{h})_{\xi_\lambda}$.*

Proof. The proof is the same as in [PS16, Theorem 1.10]. \square

REFERENCES

- [Afe21a] S. Afentoulidis-Almpanis, *Noncubic Dirac operators for finite-dimensional modules*, Phd Thesis, Advisor: S. Mehdi, Université de Lorraine (2021).
- [Afe21b] S. Afentoulidis-Almpanis, *Noncubic Dirac operators for finite-dimensional modules*, J. Lie Theory 31 (2021), no. 4, pp. 1113-1140.
- [BGG76] I. N. Bernstein, I. M. Gel'fand, S. I. Gel'fand, *A certain category of \mathfrak{g} -modules*, (in Russian), Funkcional. Anal. i Priložen 10 (1976), no. 2, pp. 1-8.
- [BP11] D. Barbasch, P. Pandžić, *Dirac cohomology and unipotent representations of complex groups*, Noncommutative geometry and global analysis, 1-22, Contemp. Math., 546, Amer. Math. Soc., Providence, RI, 2011.
- [BP15] D. Barbasch, P. Pandžić, *Dirac cohomology of unipotent representations of $Sp(2n, \mathbb{R})$ and $U(p, q)$* , J. Lie Theory 25 (2015), no. 1, pp. 185-213.
- [EHW83] T. Enright, R. Howe, N. Wallach, *A classification of unitary highest weight modules*, Representation theory of reductive groups (Park City, Utah, 1982), 97-143, Progr. Math., 40, Birkhäuser Boston, Boston, MA, 1983.
- [GL90] J. García-Escudero, M. Lorente, *Classification of unitary highest weight representations for noncompact real forms*, J. Math. Phys. 31 (1990), no. 4, pp. 781-790.

- [Goe99] S. Goette, *Equivariant η -invariants on homogeneous spaces*, Math. Z. 232 (1999), pp. 1-42.
- [GW09] R. Goodman, N.R. Wallach, *Symmetry, Representations and Invariants*, Graduate texts in mathematics 255, Springer-Verlag, New York, (2009).
- [HKP09] J.-S. Huang, Yi-F. Kang, P. Pandžić, *Dirac cohomology of some Harish-Chandra modules*, Transformation Groups 14 (2009), pp. 163-173.
- [HP02] J.-S. Huang, P. Pandžić, *Dirac cohomology, unitary representations and a proof of a conjecture of Vogan*, J. Amer. Math. Soc. 15 (2002), pp. 185-202.
- [HP06] J.-S. Huang, P. Pandžić, *Dirac operators in representation theory*, Mathematics: Theory and Applications. Birkhäuser Boston, (2006), xii+199 pp.
- [HPR06] J.-S. Huang, P. Pandžić, D. Renard, *Dirac operators and Lie algebra cohomology*, Represent. Theory 10 (2006), pp. 299-313.
- [Hum08] J. E. Humphreys, *Representations of semisimple Lie algebras in the BGG category \mathcal{O}* , Graduate Studies in Mathematics 94, American Mathematical Society, Providence, (2008), xvi+289 pp.
- [HX12] J.-S. Huang, W. Xiao, *Dirac cohomology of highest weight modules*, Selecta Math. (N.S.) 18 (2012), no. 4, pp. 803-824.
- [Jan79] J. C. Jantzen, *Moduln mit einem höchsten Gewicht*, (in German), Lecture Notes in Mathematics 750, Springer, Berlin, (1979), ii+195 pp.
- [Kna96] A. W. Knaapp, *Lie Groups Beyond an Introduction*, 2nd edition, Progress in Mathematics 140, Birkhäuser, Boston, (1996).
- [Kob12] T. Kobayashi, *Restrictions of generalized Verma modules to symmetric pairs*, Transform. Groups 17 (2012), no. 2, pp. 523-546.
- [Kos61] B. Kostant, *Lie algebra cohomology and the generalized Borel-Weil theorem*, Ann. of Math. (2) 74 (1961), pp. 329-387.
- [Kos99] B. Kostant, *A cubic Dirac operator and the emergence of Euler number multiplets of representations for equal rank subgroups*, Duke Mathematical Journal 100 (1999), pp. 447-501.
- [Kos03] B. Kostant, *Dirac cohomology for the cubic Dirac operator*, Studies in memory of Issai Schur, Vol. 210. Progr. Math. Birkhäuser Boston, 2003, pp. 69-93.
- [KØSS15] T. Kobayashi, B. Ørsted, P. Somberg, V. Souček, *Branching laws for Verma modules and applications in parabolic geometry I*, Adv. Math. 285 (2015), pp. 1796-1852.
- [MP11] S. Mehdi, R. Parthasarathy, *Cubic Dirac cohomology for generalized Enright-Varadarajan modules*, J. Lie Theory 21.4 (2011), pp. 861-884.
- [MPV17] S. Mehdi, P. Pandžić, D. Vogan, *Translation principle for Dirac index*, Amer. J. Math. 139 (2017), no. 6, pp. 1465-1491.

- [MZ14] S. Mehdi, R. Zierau, *The Dirac cohomology of a finite dimensional representation*, Proceedings of the American Mathematical Society 142 (2014), pp. 1507-1512.
- [PS16] P. Pandžić, P. Somberg, *Higher Dirac cohomology of modules with generalized infinitesimal character*, Transform. Groups 21.3 (2016), pp. 803-819.
- [Ver68] D.-N. Verma, *Structure of certain induced representations of complex semisimple Lie algebras*, Bull. Amer. Math. Soc. 74 (1968), pp. 160-166.
- [Vog97] D. Vogan, *Dirac operators and unitary representations*, Three talks at the MIT Lie Groups Seminar, (1997).

MATHEMATICAL INSTITUTE OF CHARLES UNIVERSITY, SOKOLOVSKA 49/83, 18675
PRAHA 8, CZECH REPUBLIC

Email address: `almpanis@karlin.mff.cuni.cz`