

# THE GOLOMB TOPOLOGY OF POLYNOMIAL RINGS, II

DARIO SPIRITO

**ABSTRACT.** We study the interplay of the Golomb topology and the algebraic structure in polynomial rings  $K[X]$  over a field  $K$ . In particular, we focus on infinite fields  $K$  of positive characteristic such that the set of irreducible polynomials of  $K[X]$  is dense in the Golomb space  $G(K[X])$ . We show that, in this case, the characteristic of  $K$  is a topological invariant, and that any self-homeomorphism of  $G(K[X])$  is the composition of multiplication by a unit and a ring automorphism of  $K[X]$ .

## 1. INTRODUCTION

Let  $R$  be an integral domain, i.e., a commutative unitary ring without zero-divisors. The *Golomb space*  $G(R)$  on  $R$  is the topological space having  $R^\bullet := R \setminus \{0\}$  as its base space and whose topology (the *Golomb topology*) is generated by the cosets  $a + I$ , where  $a \in R$  and  $I$  is an ideal such that  $a^\bullet$  and  $I$  are coprime, i.e.,  $\langle a, I \rangle = R$ . This construction was originally considered on the set  $\mathbb{N}$  of natural numbers by Brown [4] and Golomb [7, 8], as part of a series of coset topologies [9], and subsequently extended to arbitrary rings by [5] (following ideas introduced in [1]) with particular focus on what happens when  $R$  is a Dedekind domain with infinitely many maximal ideals. In this case,  $G(R)$  is a Hausdorff space that is not regular, and is a connected space that is disconnected at each of its points.

An interesting question is how much the topological structure of  $G(R)$  reflects the algebraic structure of  $R$ : for example, it is an open question whether the fact that  $G(R)$  and  $G(S)$  are homeomorphic implies that  $R$  and  $S$  are isomorphic (as rings). Relatedly, one can ask if there are self-homeomorphisms of  $G(R)$  besides the one arising from the algebraic structure, i.e., multiplication by units and automorphisms of  $R$  (and their compositions).

These problems were studied in [10] for  $R = \mathbb{Z}$ , showing that the only self-homeomorphisms are the trivial ones (the identity and the multiplication by  $-1$ ) [10, Theorem 7.7] and that  $G(\mathbb{Z}) \simeq G(R)$  cannot happen when  $R \neq \mathbb{Z}$  is contained in the algebraic closure of  $\mathbb{Q}$  [10, Theorem

---

*Date:* September 27, 2022.

*2020 Mathematics Subject Classification.* 54G99; 13F20; 13F05; 12E99.

*Key words and phrases.* Golomb topology; polynomial rings; Dedekind domains.

7.8]; a variant of its method showed that  $\mathbb{N}$ , with the Golomb topology, is rigid, i.e., it does not have any nontrivial self-homeomorphism [2]. As a second case, [11] studied the space  $G(R)$  when  $R = K[X]$  is a polynomial ring over a field  $K$ , showing that several algebraic properties of  $K$  (for example, having positive or zero characteristic, being algebraically or separably closed) imply different properties on the Golomb space. In particular, it was shown that if  $K, K'$  are fields of positive characteristic that are algebraic over their base field, then  $G(K[X]) \simeq G(K'[X])$  imply  $K \simeq K'$  when one of them is algebraically closed [11, Theorem 5.11 and Corollary 7.2] and when they have the same characteristic [11, Theorem 7.5].

In this paper, we improve these results in two ways. In Section 2, we show that, if the set of irreducible polynomials of  $K[X]$  is dense in  $G(K[X])$ , then the characteristic of  $K$  can be detected from the Golomb topology; that is, if  $K, K'$  satisfy these hypothesis and  $G(K[X]) \simeq G(K'[X])$ , then the characteristic of  $K$  and  $K'$  are equal (Theorem 2.6). As a consequence, we show that if  $G(K[X]) \simeq G(K'[X])$  and  $K$  is algebraic over  $\mathbb{F}_p$ , then  $K$  and  $K'$  must be isomorphic (Theorem 2.7). In Section 3, we concentrate on self-automorphisms of the Golomb space  $G(K[X])$  and show that (under the above density hypothesis, and the condition that the characteristic of  $K$  is positive) all such self-automorphisms are algebraic in nature, being compositions of a multiplication by a unit and a ring automorphism of  $K[X]$  (Theorem 3.9). The key to both results is a variant of the proofs of [1, Lemma 5.10] and [11, Theorem 6.7], which allows to prove that, under appropriate hypothesis, any homeomorphism of Golomb spaces respects powers, i.e.,  $h(a^n) = h(a)^n$  for every  $a \in G(R)$ ,  $n \in \mathbb{N}$ .

Throughout the paper,  $K$  is a field, and if  $R$  is an integral domain then  $G(R)$  denotes the Golomb space as defined above. Given a set  $X \subseteq R$ , we set  $X^\bullet := X \setminus \{0\}$ . By a ‘‘polynomial ring’’, we always mean a polynomial ring in one variable. We denote by  $U(R)$  the set of the units of  $R$ , so that  $U(K[X]) = K^\bullet$ .

If  $h : G(R) \rightarrow G(S)$  is a homeomorphism, then  $h$  sends units to units [5, Theorem 13] and prime ideals to prime ideals (i.e., if  $P$  is a prime ideal of  $R$ , then  $h(P^\bullet) \cup \{0\}$  is a prime ideal of  $S$ ; equivalently,  $h(P^\bullet) = Q^\bullet$  for some prime ideal  $Q$  of  $S$ ) [11, Theorem 3.6].

If  $x$  is a prime element of  $R$ , we set  $\text{pow}(x) := \{ux^n \mid u \in U(R), n \in \mathbb{N}\}$ . If  $h : G(R) \rightarrow G(S)$  is a homeomorphism, then  $h(\text{pow}(x)) = \text{pow}(h(x))$  (see [11, Section 2]).

If  $P$  is a prime ideal of  $R$ , the  $P$ -topology on  $R \setminus P$  is just the  $P$ -adic topology, i.e., the topology having the sets  $a + P^n$ , for  $n \in \mathbb{N}$ , as a local basis for  $a$ . The  $P$ -topology can be recovered from the Golomb topology, and thus, for any prime ideal  $P$  of  $R$ , a homeomorphism  $h : G(R) \rightarrow G(S)$  restricts to a homeomorphism between  $R \setminus P$

endowed with the  $P$ -topology and  $S \setminus Q$  endowed with the  $Q$ -topology (where  $h(P^\bullet) = Q^\bullet$ ) [10, Theorem 4.3].

We say that  $R$  is *Dirichlet* if the set of irreducible elements of  $R$  is dense, with respect to the Golomb topology. If  $R = K[X]$  is Dirichlet, where  $K$  is a field, we say for brevity that  $K$  itself is Dirichlet. This happens, for example, when  $K$  is pseudo-algebraically closed and admits separable irreducible polynomials of arbitrary large degree [3, Theorem A]; in particular, it happens when  $K$  is an algebraic extension of a finite field that is not algebraically closed [6, Corollary 11.2.4].

## 2. DETECTING THE CHARACTERISTIC

In this section, we show how to detect the characteristic of a polynomial ring from its Golomb topology.

Given an element  $t \in R^\bullet$ , we denote by  $\sigma_t$  the multiplication by  $t$ , i.e.,

$$\begin{aligned} \sigma_t: G(R) &\longrightarrow G(R), \\ y &\longmapsto ty. \end{aligned}$$

It is easy to see that the map  $\sigma_t$  is always continuous, and that it is a homeomorphism if  $t$  is a unit of  $R$ .

If  $R$  is a polynomial ring over a Dirichlet field, and  $h : G(R) \longrightarrow G(R)$  is a self-homeomorphism such that  $h(P^\bullet) = P^\bullet$  for every prime ideal  $P$ , then  $h$  must be equal to  $\sigma_u$  for some unit  $u$  [11, Proposition 7.4]; this result can be improved in the following way.

**Proposition 2.1.** *Let  $R, S$  be polynomial rings over Dirichlet fields. If  $h : G(R) \longrightarrow G(S)$  is a homeomorphism such that  $h(1) = 1$ , then  $h(uz) = h(u)h(z)$  for every  $u \in U(R)$  and  $z \in R^\bullet$ .*

*Proof.* Let  $u$  be a unit of  $R$ . Consider the map  $h' := h^{-1} \circ \sigma_{h(u)} \circ h$ , that is,

$$h'(z) = h^{-1}(h(u)h(z))$$

for every  $z \in R^\bullet$ . Since  $u$  is a unit, so is  $h(u)$ , and thus  $\sigma_{h(u)}$  is a self-homeomorphism of  $G(S)$ ; therefore,  $h'$  is a homeomorphism of  $G(R)$ .

Let now  $P$  be a prime ideal of  $R$ ; then,  $h(P^\bullet) = Q^\bullet$  for some prime ideal  $Q$  of  $S$ . For every  $z \in P$ , we have  $h(u)h(z) \in Q$ , and thus  $h'(z) = h^{-1}(h(u)h(z)) \in P$ . Hence,  $h'(P) \subseteq P$ , i.e.,  $h$  fixes every prime ideal of  $R$ ; by [11, Proposition 7.4],  $h' = \sigma_t$  for some unit  $t$  of  $R$ . However, since  $h(1) = 1$ ,

$$h'(1) = h^{-1}(h(u)h(1)) = h^{-1}(h(u)) = u,$$

and thus it must be  $t = u$ , i.e.,  $h'(z) = uz$  for every  $z \in R^\bullet$ ; it follows that,  $h(uz) = h(u)h(z)$  for every  $z \in R^\bullet$ .  $\square$

We now want to study the relationship between a homeomorphism of Golomb topologies and powers of elements. For an element  $z \in R^\bullet$ , we denote by  $z^\mathbb{N}$  the set of powers of  $z$ , i.e.,  $z^\mathbb{N} := \{z^n \mid n \in \mathbb{N}\}$ .

**Lemma 2.2.** *Let  $R, S$  be polynomial rings over Dirichlet fields, and let  $h : G(R) \rightarrow G(S)$  be a homeomorphism such that  $h(1) = 1$ . The set*

$$\mathcal{X}_h := \{z \in R \mid z \text{ is irreducible and } h(z^{\mathbb{N}}) = h(z)^{\mathbb{N}}\}$$

*is dense in  $G(R)$ .*

*Proof.* Let  $Q$  be a prime ideal of  $S$ , and let  $P$  be the prime ideal of  $R$  such that  $h(P^\bullet) = Q^\bullet$ . Then,  $h$  restricts to a homeomorphism between the  $P$ -topology on  $R \setminus P$  and the  $Q$ -topology on  $S \setminus Q$ ; it follows that for every  $m \in \mathbb{N}$  there is an  $n$  such that  $h(1 + P^n) \subseteq 1 + Q^m$ . Let  $z$  be any irreducible element in  $1 + P^n$  (which exists since  $R$  is Dirichlet); then, for every  $t \in \mathbb{N}$ , we have  $z^t \in 1 + P^n$  too, and thus also  $h(z^t) \in 1 + Q^m$ . Moreover,  $z^t \in \text{pow}(z)$ , and thus there are a unit  $u$  of  $S$  and an integer  $\alpha(t)$  such that  $h(z^t) = uh(z)^{\alpha(t)}$ . Since  $U(S) \rightarrow S/Q$  is injective, it follows that  $u = 1$  for every  $t$ . Hence,  $h(z^{\mathbb{N}}) = h(z)^{\mathbb{N}}$ .

Let now  $\Omega = c + I$  be a subbasic open set of  $G(R)$ , and let  $d + J$  be a subbasic open set contained in  $h(\Omega)$ . Let  $Q$  be a prime ideal of  $S$  that is coprime with  $J$ ; then, the prime  $P$  such that  $h(P^\bullet) = Q^\bullet$  is coprime with  $I$ . Since  $R$  is Dirichlet,  $(c + I) \cap (1 + P^n)$  contains irreducible elements for every  $n$ ; in particular, if  $n$  is large enough, every irreducible polynomial in  $1 + P^n$  is in  $\mathcal{X}_h$ , and thus  $\mathcal{X}_h \cap (c + I)$  is nonempty. Thus  $\mathcal{X}_h$  is dense.  $\square$

The following proof follows the one of [11, Theorem 6.8], which in turn was based on the one of [1, Lemma 5.10].

**Lemma 2.3.** *Let  $R, S$  be polynomial rings over Dirichlet fields, and let  $h : G(R) \rightarrow G(S)$  be a homeomorphism such that  $h(1) = 1$ . For every  $a \in R$ , we have  $h(a^{\mathbb{N}}) = h(a)^{\mathbb{N}}$ .*

*Proof.* If  $a$  is a unit, the statement follows from Proposition 2.1. Take any  $a \in R$ ,  $a \notin U(R)$ , and let  $b := h(a)$ . We first claim that  $h(a^{\mathbb{N}}) \subseteq b^{\mathbb{N}}$ . Take any  $n \in \mathbb{N}$  and let  $f : G(R) \rightarrow G(R)$  be defined as  $f(y) = y^n$ , and let  $\phi := h \circ f \circ h^{-1}$ . Then,  $\phi : G(S) \rightarrow G(S)$  is continuous (since  $f$  is continuous) and  $\phi(P^\bullet) \subseteq P^\bullet$  for every prime ideal  $P$  of  $R$ . Let

$$c := \phi(b) = h(f(a)) = h(a^n),$$

and suppose that  $c \notin b^{\mathbb{N}}$ . Take an integer  $k$  such that  $\deg b^k > \deg h(a^n) = \deg c$ ; then, since  $b$  and  $b^k - 1$  are coprime, so are  $c$  and  $b^k - 1$  (since  $c$  and  $b$  belong to the same prime ideals, by [10, Theorem 3.6]), and thus  $\Omega := c + (b^k - 1)S$  is an open set. Since  $\phi$  is open, there is a neighborhood  $b + dS$  of  $b$  such that  $\phi(b + dS) \subseteq \Omega$ .

By Lemma 2.2, we can find an irreducible polynomial  $z \in h^{-1}(b + d(b^k - 1)S)$  such that  $h(z^{\mathbb{N}}) = h(z)^{\mathbb{N}}$ ; setting  $y := h(z)$ , we have

$$\phi(y) = (h \circ f \circ h^{-1})(h(z)) = h(z^n) = h(z)^l$$

for some integer  $l$ . Thus, we have

$$\phi(y) \in \phi(b + dS) \subseteq \Omega = c + (b^k - 1)S$$

and

$$\phi(y) = h(z)^l \in (b + d(b^k - 1)S)^l \subseteq b^l + (b^k - 1)S;$$

hence,  $c \equiv b^l \pmod{(b^k - 1)S}$ . If  $l = ik + j$ , we have  $b^j \equiv b^l \pmod{(b^k - 1)S}$ ; hence, we have  $c \equiv b^j \pmod{(b^k - 1)S}$  for some  $0 \leq j < k$ . However, by hypothesis,  $c \neq b^j$ , and the degree of both  $c$  and  $b^j$  is less than the degree of  $b^k - 1$ ; this is a contradiction, and thus we must have  $h(a^{\mathbb{N}}) \subseteq b^{\mathbb{N}}$ .

The opposite inclusion is obtained using the homeomorphism  $h^{-1}$ . Thus,  $h(a^{\mathbb{N}}) = b^{\mathbb{N}} = h(a)^{\mathbb{N}}$ , as claimed.  $\square$

**Lemma 2.4.** *Let  $R$  be an integral domain. If  $U(R)$  is infinite, then its exponent is infinite.*

*Proof.* Suppose that the exponent of  $U(R)$  is finite, say equal to  $n$ . Since  $R$  is a domain, there can be at most  $n$  units of order divisible by  $n$  (the roots of  $X^n - 1 = 0$ ); however, this is impossible since  $U(R)$  is infinite. Hence, the exponent is infinite.  $\square$

**Proposition 2.5.** *Let  $R, S$  be polynomial rings over infinite Dirichlet fields, and let  $h : G(R) \rightarrow G(S)$  be a homeomorphism such that  $h(1) = 1$ . For every  $a \in R$  and every  $n \in \mathbb{N}$ , we have  $h(a^n) = h(a)^n$ .*

*Proof.* If  $a$  is a unit the claim follows from Proposition 2.1. Take  $a \in R \setminus U(R)$  and  $n \in \mathbb{N}$ ; by Lemma 2.3, we have  $h(a^n) = h(a)^s$  for some  $s \in \mathbb{N}$ . By hypothesis,  $R$  has infinitely many units, and thus by Lemma 2.4 there is a unit  $u$  whose order is larger than  $n$  and  $s$ ; then, using Proposition 2.1 we have

$$h((ua)^n) = (h(ua))^t = (h(u)h(a))^t = h(u)^t h(a)^t$$

for some  $t$  and

$$h((ua)^n) = h(u^n a^n) = h(u^n)h(a^n) = h(u)^n h(a)^s.$$

The equality

$$h(u)^t h(a)^t = h(u)^n h(a)^s$$

can only hold if  $n = t = s$ ; in particular,  $n = s$  and  $h(a^n) = h(a)^n$ , as claimed.  $\square$

**Theorem 2.6.** *Let  $K, K'$  be infinite Dirichlet fields. If  $G(K[X])$  and  $G(K'[X])$  are homeomorphic, then they have the same characteristic.*

*Proof.* If one of  $K$  and  $K'$  has characteristic 0, the claim follows from [11, Corollary 4.2]. Suppose thus that  $\text{char } K = p$  and  $\text{char } K' = q$ , with  $p, q > 0$ .

Let  $h : G(K[X]) \rightarrow G(K'[X])$  be a homeomorphism such that  $h(1) = 1$ , and let  $a$  be an irreducible element. Let  $f$  be a factor of  $a - 1$ . By [11, Proposition 5.5], the sequence  $a^{p^n}$  converges to 1 in the  $(f)$ -topology, and thus also  $h(a^{p^n})$  converges to 1 in the  $h((f))$ -topology. By Proposition 2.5,  $h(a^{p^n}) = h(a)^{p^n}$ . By [11, Proposition 5.5], it follows

that  $v_q(p^n) \rightarrow \infty$ , where  $v_q$  is the  $q$ -adic valuation; thus, it must be  $q = p$ . The claim is proved.  $\square$

We note that the proof of the previous theorem actually needs only *one* irreducible element  $a$  such that  $h(a^n) = h(a)^n$  for every  $n$ .

**Theorem 2.7.** *Let  $K, K'$  be fields. If  $K$  is algebraic over  $\mathbb{F}_p$  and  $G(K[X]) \simeq G(K'[X])$ , then  $K \simeq K'$ .*

*Proof.* If  $K$  is finite then  $|U(K[X])| = |K| - 1$ . Since the set of units is preserved by homeomorphisms of the Golomb topology,  $K$  and  $K'$  must have the same cardinality, and thus they are isomorphic. Suppose  $K, K'$  are infinite.

By [11, Corollary 4.2],  $K'$  has positive characteristic, and by [11, Corollary 7.2]  $K'$  must be algebraic over its base field  $\mathbb{F}_q$ .

If  $K$  is algebraically closed, then the set  $G_1(K[X]) := \{z \in K[X] \mid z \text{ is contained in a unique prime ideal}\}$  is not dense in  $G(K[X])$  [11, Proposition 5.2(a)]; conversely, if  $K$  is not algebraically closed then  $G_1(K[X])$  is dense since it contains the irreducible polynomials. The same holds for  $K'$ . Since a homeomorphism of Golomb topologies sends  $G_1(K[X])$  to  $G_1(K'[X])$  [10, Theorem 3.6(b)], it follows that if  $K$  is algebraically closed then so is  $K'$ , and in this case  $p = q$  by [11, Theorem 5.11]; hence  $K \simeq K'$ .

If  $K$  and  $K'$  are not algebraically closed, they are pseudo-algebraically closed fields containing separable irreducible polynomials of arbitrarily large degree [3, Theorem A]; by Theorem 2.6, it follows that  $p = q$ . By [11, Theorem 7.5], it follows that  $K \simeq K'$ .  $\square$

### 3. SELF-HOMEOMORPHISMS AND AUTOMORPHISMS

In this section, we concentrate on the study of self-homeomorphisms of the Golomb space  $G(K[X])$ . We shall work under the following assumptions:

#### Hypothesis 3.1.

- $K$  is an infinite field of characteristic  $p > 0$ ;
- $R := K[X]$ ;
- $R$  is Dirichlet, i.e., the set of irreducible polynomials is dense in  $G(R)$ ;
- $h : G(R) \rightarrow G(R)$  is a self-homeomorphism such that  $h(1) = 1$ .

In particular, under these hypothesis, we have  $h(a^n) = h(a)^n$  for every  $a \in R$  and  $n \in \mathbb{N}$ , by Proposition 2.5.

Given a set  $X \subseteq R$ , we define the  $p$ -radical of  $X$  as

$$\text{rad}_p(X) := \{f \in R \mid f^{p^n} \in X \text{ for every large } n\}.$$

This construction is invariant under  $h$ , in the following sense.

**Lemma 3.2.** *Assume Hypothesis 3.1. For every  $X \subseteq R^\bullet$ , we have  $h(\text{rad}_p(X)) = \text{rad}_p(h(X))$ .*

*Proof.* If  $f \in \text{rad}_p(X)$ , then  $f^{p^n} \in X$  for every  $n \geq N$ , and thus  $h(f^{p^n}) \in h(X)$ . By Proposition 2.5, it follows that  $h(f)^{p^n} \in h(X)$  for  $n \geq N$ , and thus  $h(f) \in \text{rad}_p(h(X))$ . Conversely, if  $g = h(f) \in \text{rad}_p(h(X))$ , then  $g^n = h(f)^{p^n} = h(f^{p^n})$  belongs to  $h(X)$  for large  $n$ , and thus applying  $h^{-1}$  we have  $f^{p^n} \in X$  for large  $n$ . Thus  $f \in \text{rad}_p(X)$  and  $g \in h(\text{rad}_p(X))$ .  $\square$

**Lemma 3.3.** *Let  $P$  a prime ideal of a ring  $R$  of characteristic  $p$ . For every  $m \in \mathbb{N}^+$ , we have  $\text{rad}_p(1 + P^m) = 1 + P$ .*

*Proof.* If  $t \in 1 + P$ , then  $t = 1 + x$  with  $x \in P$ ; if  $p^n \geq m$ , then  $t^{p^n} = (1 + x)^{p^n} = 1 + x^{p^n} \in 1 + P^m$ , and thus  $t \in \text{rad}_p(1 + P^m)$ . Conversely, if  $t \in \text{rad}_p(1 + P^m)$  then  $t^{p^n} \in 1 + P^m$  for some  $m$ ; in particular,  $t^{p^n} \in 1 + P$ , and thus  $t^{p^n} - 1 = (t - 1)^{p^n} \in P$ . Hence  $t - 1 \in P$ , i.e.,  $t \in 1 + P$ .  $\square$

**Proposition 3.4.** *Assume Hypothesis 3.1, and let  $P, Q$  be prime ideals of  $R$  with  $h(P^\bullet) = Q^\bullet$ . Then,  $h(1 + P) = 1 + Q$ .*

*Proof.* Since  $h(1) = 1$ ,  $h(1 + P)$  is open in the  $Q$ -topology, and thus  $1 + Q^m \subseteq h(1 + P)$  for some  $m$ . Therefore,

$$1 + Q = \text{rad}_p(1 + Q^m) \subseteq \text{rad}_p(h(1 + P)) = h(\text{rad}_p(1 + P)) = h(1 + P)$$

using Lemmas 3.2 and 3.3. Applying the same reasoning to  $h^{-1}$  gives  $1 + P \subseteq h^{-1}(1 + Q)$ , i.e.,  $h(1 + P) \subseteq 1 + Q$ . Hence,  $h(1 + P) = 1 + Q$ , as claimed.  $\square$

**Corollary 3.5.** *Assume Hypothesis 3.1, and let  $P, Q$  be prime ideals of  $R$  with  $h(P^\bullet) = Q^\bullet$ . If  $u \in K^\bullet$ , then  $h(u + P) = h(u) + Q$ . In particular, if  $f \equiv u \pmod{P}$  for some unit  $u$ , then  $h(f) \equiv h(u) \pmod{Q}$ .*

*Proof.* By Propositions 2.1, we have  $u + P = u(1 + P)$ ; by Proposition 3.4, it follows that

$$h(u + P) = h(u)h(1 + P) = h(u)(1 + Q) = h(u) + Q,$$

as claimed. The ‘‘in particular’’ statement follows from the fact that  $f \equiv u \pmod{P}$  is equivalent to  $f \in u + P$ .  $\square$

**Corollary 3.6.** *Assume Hypothesis 3.1. If  $f$  is a linear polynomial, so is  $h(f)$ .*

*Proof.* Since  $f$  is linear,  $(f)$  is prime and every polynomial  $g \notin (f)$  is equivalent to a unit modulo  $(f)$ . By Corollary 3.5, it follows that every  $g \notin (h(f))$  is equivalent to a unit modulo  $(h(f))$ ; hence,  $h(f)$  must be linear too.  $\square$

In particular, by Corollary 3.6,  $h(X)$  is a linear polynomial; it follows that there is an automorphism  $\sigma$  of  $R$  sending  $h(X)$  to  $h$ . Since  $\sigma$  restricts to a self-homeomorphism of  $G(R)$ , passing from  $h$  to  $H := \sigma \circ h$  we obtain a self-homeomorphism of  $G(R)$  that fixes both 1 and  $X$ . Thus, it is not restrictive to assume also that  $h(X) = X$ , as we do in the remaining part of the section.

**Lemma 3.7.** *Assume Hypothesis 3.1 and suppose  $h(X) = X$ . For every  $u \in K^\bullet$ , we have  $h(X + u) = X + h(u)$ .*

*Proof.* We have

$$\begin{cases} X + u \equiv u \pmod{(X)} \\ X \equiv -u \pmod{(X + u)}. \end{cases}$$

Applying  $h$  to both equivalences, using  $h(X) = X$  and Corollary 3.5, we have

$$\begin{cases} h(X + u) \equiv h(u) \pmod{(X)} \\ X \equiv -h(u) \pmod{(h(X + u))}. \end{cases}$$

The first equation implies that  $X$  divides  $h(X + u) - h(u)$ ; since  $h(X + u)$  is linear, it follows that  $h(X + u) = vX + h(u)$  for some  $v \in K^\bullet$ . The second equation implies that  $vX + h(u)$  divides  $X + h(u)$ ; the only possibility is  $v = 1$ , i.e.,  $h(X + u) = X + h(u)$ . The claim is proved.  $\square$

**Lemma 3.8.** *Assume Hypothesis 3.1 and suppose  $h(X) = X$ . Then, the map*

$$H: K \longrightarrow K, \\ a \longmapsto \begin{cases} h(a) & \text{if } a \neq 0 \\ 0 & \text{if } a = 0 \end{cases}$$

*is an automorphism of  $K$ .*

*Proof.* Let  $a, b \in K$ . If  $a = 0$  or  $b = 0$  then clearly  $H(ab) = H(a)H(b)$  and  $H(a + b) = H(a) + H(b)$ .

Suppose  $a \neq 0 \neq b$ . Then,  $H = h$  on these values, and  $h(ab) = h(a)h(b)$  by Proposition 2.1. Furthermore,

$$X + a + b \equiv b \pmod{(X + a)};$$

applying  $h$  and using Lemma 3.7, we have

$$X + h(a + b) \equiv h(b) \pmod{(X + h(a))},$$

that is,  $X + h(a)$  divides  $X + h(a + b) - h(b)$ . Thus, it must be  $X + h(a) = X + h(a + b) - h(b)$  and  $h(a) = h(a + b) - h(b)$ , that is,  $h(a) + h(b) = h(a + b)$ . It follows that  $H$  is an automorphism of  $K$ , as claimed.  $\square$

**Theorem 3.9.** *Let  $K$  is an infinite Dirichlet field of positive characteristic, and let  $h$  be a self-homeomorphism of  $G(K[X])$ . Then, there are a unit  $u \in K^\bullet$  and an automorphism  $\sigma$  of  $K[X]$  such that  $h(f) = u\sigma(f)$  for every  $f \in K[X]^\bullet$ .*



*Proof.* By [5, Theorem 13],  $h(1) = u$  is a unit of  $R$ ; since the multiplication  $\sigma_u$  is a self-homeomorphism of  $G(R)$ , the map  $h_1 := \sigma_u^{-1} \circ h$  is a self-homeomorphism such that  $h_1(1) = 1$ .

By Corollary 3.6,  $h_1(X)$  is a linear polynomial, and thus there is an automorphism  $\sigma_1$  of  $R$  such that  $h_1(X) = \sigma_1(X)$ . Thus,  $h_2 := \sigma_1^{-1} \circ h_1$  is again a self-homeomorphism of  $G(R)$ , and furthermore both 1 and  $X$  are fixed points of  $h_2$ .

By Lemma 3.8, the restriction of  $h_2$  to  $K$  is an automorphism  $H$ ; this map can be extended to an automorphism  $\sigma_2$  of  $R$  by setting  $\sigma_2(\sum_i a_i X^i) = \sum_i H(a_i) X^i$ . Thus,  $h_3 := \sigma_2^{-1} \circ h_2$  is a self-homeomorphism of  $G(R)$  such that  $h_3(u) = u$  for every  $u \in K^\bullet$  and  $h_3(X) = X$ . By Lemma 3.7, it follows that  $h_3(X + u) = X + u$  for every  $u \in K^\bullet$ .

Let now  $f \in R$ , and take any  $t \in K$ . Then,

$$\begin{cases} f \equiv f(t) \pmod{X - t} \\ h_3(f) \equiv h_3(f)(t) \pmod{X - t}. \end{cases}$$

Since  $h_3((X - t)) = (X - t)$ , by Corollary 3.5 the first equivalence also implies  $h_3(f) \equiv h_3(f(t)) \pmod{X - t}$ . Hence,  $h_3(f)(t) = h_3(f(t)) = f(t)$ . Since this happens for every  $t \in K$ , and  $K$  is infinite, it follows that  $f = h_3(f)$ , that is,  $h_3$  is the identity on  $G(R)$ .

Going back to the definition,

$$h_3 = \sigma_2^{-1} \circ h_2 = \sigma_2^{-1} \circ \sigma_1^{-1} \circ h_1 = \sigma_2^{-1} \circ \sigma_1^{-1} \circ \sigma_u^{-1} \circ h,$$

i.e.,  $h = \sigma_u \circ \sigma_1 \circ \sigma_2$ . Since  $\sigma_u(x) = ux$  for every  $x$ , setting  $\sigma := \sigma_1 \circ \sigma_2$  (which is still an automorphism of  $R$ ), we obtain  $h(f) = u\sigma(f)$  for every  $f \in G(R)$ , as claimed.  $\square$

## REFERENCES

- [1] Taras Banach, Jerzy Mioduszewski, and Sławomir Turek. On continuous self-maps and homeomorphisms of the Golomb space. *Comment. Math. Univ. Carolin.*, 59(4):423–442, 2018.
- [2] Taras Banach, Dario Spirito, and Sławomir Turek. The Golomb space is topologically rigid. *Comment. Math. Univ. Carolin.*, 62(3):347–360, 2021.
- [3] Lior Bary-Soroker. Dirichlet’s theorem for polynomial rings. *Proc. Amer. Math. Soc.*, 137(1):73–83, 2009.
- [4] Morton Brown. A countable connected Hausdorff space. In L.W. Cohen, editor, *The April meeting in New York*, volume 4, pages 330–371. Bull. Amer. Math. Soc., 1953. Abstract 423.
- [5] Pete L. Clark, Noah Lebowitz-Lockard, and Paul Pollack. A note on Golomb topologies. *Quaest. Math.*, 42(1):73–86, 2019.
- [6] Michael D. Fried and Moshe Jarden. *Field arithmetic*, volume 11 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 2005.
- [7] Solomon W. Golomb. A connected topology for the integers. *Amer. Math. Monthly*, 66:663–665, 1959.

- [8] Solomon W. Golomb. Arithmetica topologica. In *General Topology and its Relations to Modern Analysis and Algebra (Proc. Sympos., Prague, 1961)*, pages 179–186. Academic Press, New York; Publ. House Czech. Acad. Sci., Prague, 1962.
- [9] John Knopfmacher and Stefan Porubsky. Topologies related to arithmetical properties of integral domains. *Exposition. Math.*, 15(2):131–148, 1997.
- [10] Dario Spirito. The Golomb topology on a Dedekind domain and the group of units of its quotients. *Topology Appl.*, 273:107101, 20, 2020.
- [11] Dario Spirito. The Golomb topology of polynomial rings. *Quaest. Math.*, 44(4):447–468, 2021.

DIPARTIMENTO DI SCIENZE MATEMATICHE, INFORMATICHE E FISICHE, UNIVERSITÀ DEGLI STUDI DI UDINE, UDINE, ITALY  
*Email address:* `dario.spirito@uniud.it`